MYB93 regulates responses to environmental sulphur in *Arabidopsis* and

tomato.

Xulyu Cao[1,2], Helen B. Wilkinson[2], Bethany Hutton[2], Nancy McMulkin[2], Neil S.

Graham[3], Ross Etherington[2], Alice Oliver[2], Clare M. Clayton[2], Harjeet Kaur[2], Juliet

C. Coates[2*].

[1] College of Resources and Environment, Academy of Agricultural Sciences, Key

Laboratory of Efficient Utilization of Soil and Fertilizer resources, Southwest

University, Chongqing, 400716, China.

[2] School of Biosciences, University of Birmingham, Birmingham B15 2TT, UK.

[3] Plant and Crop Sciences Division, School of Biosciences, Sutton Bonington

Campus, University of Nottingham, LE12 5RD, UK.

[*]Corresponding author. ORCID 0000-0002-2381-0298

Abstract

Sulphur (S) is an important nutrient that has wide-ranging effects on plant health and

metabolism. Several classes of transcription factor respond to S deprivation,

including R2R3-MYBs. In *Arabidopsis*, the *AtMYB93* transcription factor-encoding

gene is upregulated by S deprivation. *At*MYB93 has a non-redundant function in

lateral root development and redundant functions in suberin biosynthesis alongside

related MYB transcription factors, but *At*MYB93's role in S signalling, and how it

relates to lateral root development, is unknown.

We show that the transcriptome of *Atmyb93* mutant roots implicates *At*MYB93 in

responses to S, including changes in S transport and metabolism, and flavonoid- and

carbohydrate metabolism. Elemental analysis demonstrates that the *Atmyb93*

mutant has elevated shoot S levels while tomato *Sl*MYB93-overexpressing plants

have reduced shoot S. We uncover a stimulatory effect of S deprivation on

adventitious root development. However, *Atmyb93* mutants do not show significant

changes in sensitivity to S with respect to lateral- or adventitious root development,

most likely due to some functional redundancy. Moreover, *AtMYB93* promoter

activity is not spatially regulated by S deprivation. Taken together, our data suggest

that *AtMYB93* has a role in mediating root responses to S in alongside other root

transcription factors.

Keywords: MYB transcription factor, Sulphur, Root development, transcriptome.

Introduction

Plants require a range of nutrients to grow and develop optimally. The soil provides macronutrients and micronutrients, in relatively large and small amounts respectively. The key macronutrients that plants obtain from soil are nitrogen (N), phosphorus (P), potassium (K), calcium (Ca), magnesium (Mg) and sulphur (S). S is often considered a 'neglected' macronutrient: S fertilisation can improve plant yield, abiotic- and biotic stress tolerance (Ali et al. 2021; Narayan et al. 2023; Sarda et al. 2013; Zenda et al. 2021) although S deprivation may improve resistance to some pathogens (Criollo-Arteaga et al. 2021). In terms of human nutrition, plants are the major source of methionine, an essential S-containing amino acid (Kopriva et al. 2019).

Problems with S deficiency in crops are increasing globally due to the decreasing amount of rainfall-derived S in combination with the continued development of high-yielding crop varieties (Sharma et al. 2024) (Zenda et al. 2021) (Sarda et al. 2013) (Grant et al. 2012). Crop requirements for S vary, with Brassicas (especially oilseed rape) requiring high S, cotton and sugarcane requiring medium S and cereals having relatively low S requirements (Zenda et al. 2021). However, naturally high S *Brassica* crops used for animal feed can promote risk of goitre and kale anemia (Smith 1980; Smith et al. 1974; Paxman and Hill 1974). Moreover, S levels can affect crop quality, for example high S can cause increased acrylamide formation in cooked potatoes (Elmore et al. 2007; Elmore et al. 2010). In marine ecosystems, sulfide stress in seagrass beds, caused by microbial anaerobic digestion of sulfate in sediments, increases with elevated temperature (Zhang et al. 2024). Thus, S levels in plants may require careful management on a case-by-case basis.

S deficiency in plants has wide-ranging effects including reduced chlorophyll, changes in sugar- and amino acid metabolism, changes in phenylpropanoid metabolism (particularly flavonoids) and changes in antioxidant levels (e.g. (Lunde et al. 2008; Wawrzynska et al. 2022; Bielecka et al. 2014; Chandra and Pandey 2014; Forieri et al. 2017; Henriquez-Valencia et al. 2018; Canales et al. 2020; Robe et al. 2020; Nikiforova et al. 2003; Wang et al. 2022). Moreover, various S-containing

compounds affect plant hormone biosynthesis directly or indirectly (Wawrzynska and Sirko 2024).

Several classes of transcription factor (TF) have been identified that may co-ordinate responses to S stress. The ETHYLENE INSENSITIVE LIKE (EIL) TF family member SULFUR LIMITATION 1 (SLIM1/EIL3) is a key regulator of metabolism under S deficiency in *Arabidopsis* and tomato roots (Maruyama-Nakashita et al. 2006; Canales et al. 2020) partnering with EIL1 (Dietzen et al. 2020). Bioinformatic analysis suggested NUCLEAR FACTOR Y-A2 (NFY-A2) and the circadian regulator REVEILLE2 (RVE2) as 'hub' genes co-ordinating expression of S-response genes (Henriquez-Valencia et al. 2018). S availability affects expression of *Arabidopsis* R2R3-MYB family TFs that regulate biosynthesis of glucosinolates, S-containing secondary metabolites with defence functions, likely via SLIM1 (Frerigmann and Gigolashvili 2014a, b; Li et al. 2013). Moreover, S stress increases expression of a suite of *Arabidopsis* root TFs including several R2R3-MYBs; in particular, *AtMYB93* shows a sustained increase upon S deprivation and S resupply (Bielecka et al. 2014).

AtMYB93 is part of a flowering plant-specific clade (the S24 clade) of plant R2R3-MYB transcription factors (Du et al. 2015; Gibbs et al. 2014). In *Arabidopsis*, the S24 clade consists of *AtMYB93*, *AtMYB92* and *AtMYB53* (Du et al. 2015; Gibbs et al. 2014). All three *Arabidopsis* S24 *R2R3-MYB*s show root-enriched expression with *AtMYB93* being expressed exclusively in the root (Gibbs et al. 2014). The function of the three *Arabidopsis* S24 *MYB* genes is not fully redundant. *AtMYB93*, but not *AtMYB92*, is a negative regulator of LR development and only *AtMYB93* expression is induced by auxin (Gibbs et al. 2014). As mentioned above, *AtMYB93* is significantly upregulated upon S deprivation with persistent elevated expression upon S resupply in 7-day old seedlings (Bielecka et al. 2014). *AtMYB53* is upregulated by S deprivation to a lesser extent than *AtMYB93* and is not persistently upregulated upon S resupply (Bielecka et al. 2014). *AtMYB92* shows a limited S deprivation response: upregulation in seedlings when seeds are germinated directly on S-deficient medium but not when S-replete seedlings are transferred to S-deficient medium after 8 days of growth (Nikiforova et al. 2003). Interestingly, S deprivation inhibits LR development in young *Arabidopsis* seedlings (Dan et al. 2007; Dong et al. 2019; Joshi et al. 2019) although the converse occurs in older plants (Kutz et al. 2002).

AtMYB93 is expressed only in the root endodermis (Gibbs et al. 2014), a single-celled layer separating the outer epidermis and cortex from the inner pericycle and vasculature. The endodermis forms a selective barrier to passage of solutes, nutrients and water between the soil and the root vasculature (and hence the shoot) (Andersen et al. 2015; Geldner 2013). *AtMYB93* promoter activity is restricted to a very small number of endodermal cells overlying lateral root primordia (LRP) during early lateral root (LR) development (Gibbs et al. 2014; Shukla et al. 2021). S24 *MYB* genes and the most closely related *Arabidopsis* genes in clades S10, S11 and S42 (Du et al. 2015) are involved in synthesis of the lipophilic biopolymer suberin that surrounds mature root endodermal cells and is present in fruit and in seeds (Andersen et al. 2015; Vishwanath et al. 2015; Barberon et al. 2016). *MYB* genes act, potentially in a hierarchical fashion (Xu et al. 2022; Xu et al. 2023), to enhance suberin production. Analysis in tomato, apple and *Arabidopsis* identified a conserved cross-species gene expression 'signature' for suberin biosynthesis (Gou et al. 2017; Lashbrooke et al. 2016). Apple *MdMYB93* promotes biosynthesis and cell export of suberin in the fruit (Legay et al. 2016). *AtMYB39* (*SUBERMAN*), *AtMYB41*, *AtMYB53*, *AtMYB92* and *AtMYB93* are all expressed in the root endodermis and can promote suberin production (Cohen et al. 2020; Hu 2018; Kosma et al. 2014; Shukla et al. 2021; To et al. 2020; Wang et al. 2020). Lipid-, wax-, fatty acid and suberin biosynthesis genes are differentially expressed in an *Atmyb93/Atmyb92/Atmyb53* triple mutant and an *At*MYB53-overexpressing line (Klein 2019). *At*MYB41, *At*MYB93, *At*MYB53, *At*MYB92 function redundantly to promote suberin biosynthesis in the root endodermis with *At*MYB41 having the dominant role (Shukla et al. 2021). *SlMYB92* promotes suberin biosynthesis in the tomato exodermis (Canto-Pastor et al. 2024), while three *MYB*s, *OsMYB93a*, *OsMYB41* and *OsMYB76* also regulate root suberin biosynthesis in a monocot, rice (Huang et al. 2024). Expression of *AtMYB93*, *AtMYB53*, *AtMYB9* and two other MYB transcription factors is upregulated in the *kcs1-5* mutant that has impaired synthesis of very long chain fatty acids (VLCFAs) (Uemura et al. 2023), which are components of suberin (Serra and Geldner 2022).

Interestingly, an *Arabidopsis* mutant lacking all four of *At*MYB41, *At*MYB93, *At*MYB53 and *At*MYB92 has reduced suberin but has no change in lateral root development (Shukla et al. 2021), suggesting the functions of *At*MYB93 in suberin

formation and root development are separable. The function of *At*MYB93 and its

relatives in plant responses to S has not been investigated.

Given that *AtMYB93* has a non-redundant root function, highly specific root

expression and a unique pattern of regulation by hormones and environmental

signals compared to other closely related MYB genes, we hypothesise that *AtMYB93*

has functions in addition to suberin biosynthesis during *Arabidopsis* root

development.

Results

Transcriptome analysis of plants lacking *AtMYB93* reveals a potential function

in responding to S.

To ascertain how *AtMYB93* carries out its function, we conducted transcriptome

analysis comparing gene expression in roots of wild type (Col-0) and *Atmyb93*

seedlings using RNAseq

(Fig. 1; Online Resource 2). A total of 255 genes were differentially expressed

(p<0.001) between Col-0 and *Atmyb93*, of which 44 were downregulated (including

AtMYB93 itself) and 211 were upregulated (Online Resource 2).

When analysed for Gene Ontology (GO) term enrichment, genes downregulated in

the *Atmyb93* mutant compared to wild type showed enrichment of Biological Process

(BP) terms associated with (i) S transport and metabolism, (ii) carbohydrate and

sugar metabolism, (iii) phenylpropanoid and flavonoid metabolism and (iv) responses

to stimuli including light, auxin and karrikin (Fig. 1a). Genes upregulated in *Atmyb93*

compared to wild type were enriched for GO-BP terms associated with (i)

carbohydrate (including trehalose) metabolism, transport and responses and (ii)

responses to stimuli including carbohydrates and hypoxia/oxygen as well as other

abiotic stimuli (Fig. 1b). These changes are reminiscent of both S deprivation

responses in other studies, as outlined in the introduction to this paper, and of the

transcriptome of *slim1* and *eil1* mutant plants [25].

We collated the differentially expressed genes with known functions in S transport

and metabolism (Fig. 1c) and compared their differential expression in *Atmyb93* with

their responses to S deprivation in whole root tissue and root endodermal tissue

(Iyer-Pascuzzi et al. 2011; Winter et al. 2007). Four of the six genes with S-related

functions downregulated in *Atmyb93* showed upregulation under S deprivation, while

three of the four S-related genes upregulated in *Atmyb93* showed downregulation under S deprivation (Fig. 1c). Two genes, At1g22150 SULTR1;3 and At1g61740 sulfite exporter, show different regulation in the endodermis compared to the whole root (Fig. 1c).

There is no overlap between *Atmyb93* differentially expressed genes (Online Resource 2) and the largely suberin-related genes differentially expressed in the *Atmyb53/92/93* triple mutant (Klein 2019). Collectively, these data suggest that *At*MYB93, directly or indirectly, regulates genes related to S-deprivation responses in *Arabidopsis*.

MYB93 homologues in dicots show both root-specific expression and upregulation by S deprivation.

Our previous data demonstrated that *At*MYB93 shows the most root-specific expression of *Arabidopsis* S24 clade of R2R3-MYB genes (Gibbs et al. 2014). To determine whether expression of *MYB93*-related genes outside *Arabidopsis* is root-specific, we analysed tissue-specific expression of three tomato *MYB93* homologues. The closest tomato *MYB93* homologue is *Sl04g074170* while *Sl04g056310* has previously been annotated as *SlMYB53L* (Canales et al. 2020) and *Sl05g051550* has previously been annotated as *SlMYB92* (Canales et al. 2020; Canto-Pastor et al. 2024). Note that *Sl04g074170* is different to the '*SlMYB93*' identified in Kajala et al (2021), *Sl11g011050*, which is closest in protein sequence to *At*MYB48 (Kajala et al. 2021). All three *SlMYB93* homologues show root enrichment (Fig. 2a). We made transgenic *Arabidopsis* plants heterologously overexpressing *SlMYB93* (Fig. 2c) and showed that *SlMYB93* can reduce lateral root density and root length (Fig. 2c,d) similarly to *AtMYB93* overexpression (Gibbs et al. 2014). To extend our analysis beyond *Arabidopsis* and tomato, we analysed publicly available transcriptome data and demonstrated that root-specific or root-enriched expression of putative *MYB93* homologues is present in a wide range of, but not all, dicot species (Fig. 3). In poplar and cassava, which each have 2 closely related *MYB93* homologues, only one homologue shows root enrichment or specificity, whereas both soybean homologues show root specificity (Fig. 3). Dicot *MYB92/53*-like genes also show root enrichment (Fig. 3). Some dicot *AtMYB93*-like genes from outside the *MYB93* clade show root enrichment (*AtMYB48* and *Sl11g011050*) (Fig. 3). Some monocot *AtMYB93*-related genes show root enrichment to a lesser extent:

all rice homologues (*OsMYB93a* (Os08g37970), *OsMYB93b* (Os06g17780) and

OsMYB41(Os02g51799) (Huang et al. 2024)) are root-enriched as is one barley

homologue and one maize homologue but not the genes in *Brachypodium* (Fig. 3).

Overall, root specificity/enrichment is most pronounced in dicot *AtMYB93*

homologues (compared to *MYB92* and *MYB53*).

We also analysed, where available, the cell type specificity of *MYB93* homologues

within the root and their response to S deprivation. Both *AtMYB93* and *Sl04g074170*

show endodermal-specific and meristematic expression (Fig. 3; (Gibbs et al. 2014;

Kajala et al. 2021; Waese et al. 2017)). In contrast, *Sl11g011050* is expressed in

both exodermis and endodermis (Fig. 3; (Gibbs et al. 2014; Kajala et al. 2021;

Waese et al. 2017)). *Sl04g056310* (*SlMYB53L*) is present in endodermis, exodermis

and xylem (Fig. 3; (Gibbs et al. 2014; Kajala et al. 2021; Waese et al. 2017)) and

Sl05g051550 (*SlMYB92*) is expressed in the exodermis (Fig. 3; (Canto-Pastor et al.

2024; Kajala et al. 2021; Waese et al. 2017)). All three closest tomato *MYB93*

homologues are upregulated by S deprivation (Fig. 3; (Canales et al. 2020)) with

SlMYB93 and *SlMYB53L* showing the strongest response, similarly to *Arabidopsis*

(Fig. 3; (Bielecka et al. 2014)). Outside the MYB93 clade, *AtMYB107* and

Sl11g011050 also show some upregulation by S deprivation (Fig. 3; (Canales et al.

2020)). However, no change in rice *MYB93*-like genes is seen upon S deprivation

(Fig. 3; (Wang et al. 2022)). Taken together, these data suggest a conserved role for

dicot *MYB93*-like genes in regulating root responses to S deprivation, with the

closest *MYB93* homologues specifically functioning in the endodermis.

Changes in *MYB93* level alter plant shoot element composition.

To ask whether *MYB93*'s potential link to root S-deprivation responses is associated

with changes in the shoot, we first measured the elemental composition of key

macronutrients (Mg, P, S, K, Ca) and micronutrients (Mn, Fe, Zn, B, Cu, Mo) in

Arabidopsis wild type and *Atmyb93* mutant shoot tissue. Under normal growth

conditions, the *Atmyb93* mutant showed generally elevated macronutrient and

micronutrient levels but significant differences in S, Mg and B (Fig. 4a; Online

Resource 2). Next, we investigated elemental composition of transgenic tomato lines

with increased *SlMYB93* expression, *SlMYB93-OX* (Fig. 4b,c). The trends observed

were in accordance with the *Arabidopsis* result, with two out of three independent

SlMYB93-OX lines showing reduced shoot elements (Fig. 4C; Online Resource 2).

One line showed a significant reduction in Mg and B and another showed a

significant reduction in S, B and Cu (Fig. 4C; Online Resource 2). These data

suggest that altering *MYB93* function in the root can lead to elemental changes

particularly to S, Mg and B in the shoot.

***Atmyb93* mutants respond to S deprivation.**

To further investigate the link between *At*MYB93 and plant responses to S, we

compared root development responses of wild type and *Atmyb93* mutant plants

under S deprivation.

A reduction in lateral root number under S starvation has previously been reported

(Dong et al. 2019). A reduction in lateral root density and an increase in root length

was seen in seedlings transferred to -S for 3 days after 5 days' growth on S-replete

medium that was slightly, but not significantly, less pronounced in the *Atmyb93*

mutant (Fig. 5a,b). We also observed that S deprivation increases the number of

adventitious roots (AR) produced at the root-shoot junction (collet) in both wild type

and *Atmyb93* plants transferred to medium lacking S for 3 days after 5 days' growth

on S-replete medium, with no significant differences in AR number between wild type

and *Atmyb93* being seen (Fig. 5c). We also tested the sensitivity of *Atmyb93* to O-

acetyl serine (OAS), a compound that is part of the S assimilation pathway and

increases in response to S deprivation (Hirai et al. 2003; Hubberten et al. 2012a;

Hubberten et al. 2012b). We did not observe significant effects of OAS on either the

primary root length or lateral root density of wild type or *Atmyb93* mutant plants (Fig.

5d,e).

Collectively, these results show that S deprivation can increase adventitious root

formation at the root-hypocotyl junction and suggest that, while *Atmyb93* may show

slight insensitivity to S deprivation, factors in addition to *AtMYB93* also contribute to

root S deprivation responses.

S stress does not affect the spatial localisation of *AtMYB93* promoter activity.

As S deprivation increases *AtMYB93* gene expression (Bielecka et al. 2014) and as

AtMYB93 transcriptional activity is restricted to a very few cells in the root, we

investigated whether endodermal localization of *AtMYB93* changes under S

deprivation using transgenic *Arabidopsis* expressing *pAtMYB93::GUS* (Gibbs et al.

2014). No change in the localisation and extent of *AtMYB93* promoter activity was seen under S stress (Fig. 6).

Discussion

The *Atmyb93* mutant transcriptome highlights root responses to S.

Despite the known (albeit redundant) role for *At*MYB93 in endodermal suberin production (Klein 2019; Shukla et al. 2021) the *Atmyb93* mutant root transcriptome does not show changes in suberin metabolism or transport genes. This is in agreement with an experiment that identified genes differentially expressed between wild type and *Atmyb93* root segments containing gravity-induced 46h lateral root primordia, which identified cell wall metabolism as an enriched biological process but no suberin metabolism genes (Uemura et al. 2023). However, there is almost no gene overlap between the dataset from root segments at this very specific, synchronised developmental time point (Uemura et al. 2023) and our whole-root dataset (Online Resource 2).

Instead, genes differentially expressed in *Atmyb93* mutant whole roots compared to wild type are enriched in GO terms for S transport and metabolism, carbohydrate metabolism, flavonoid- and phenylpropanoid metabolism and responses to oxygen and other stimuli. This profile is similar to S deprivation response transcriptomes (e.g. (Bielecka et al. 2014; Canales et al. 2020; Chandra and Pandey 2014; Forieri et al. 2017; Henriquez-Valencia et al. 2018; Lunde et al. 2008; Robe et al. 2020; Wawrzynska et al. 2022)). The trend of down- and upregulated genes suggests that the *Atmyb93* mutant is in a 'S-replete' state as the key S deprivation response genes *APR2* and *SDI2* are downregulated in *Atmyb93* but upregulated upon S stress (Fig. 1c; (Maruyama-Nakashita et al. 2003; Aarabi et al. 2016)). The two sulphate transporters downregulated in the *Atmyb93* mutant are the vascular transporters *SULTR1;3* (phloem) and *SULTR3;5* (xylem, pericycle and LRP), which mediate long-distance transport from source to sink and from root to shoot, respectively (Kataoka et al. 2004; Yoshimoto et al. 2003). The genomic region of At1g61740 sulfite exporter, which is upregulated in *Atmyb93*, was also identified as binding to *At*MYB93 in a DNA-affinity purification-sequencing (DAP-seq) experiment (O'Malley et al. 2016). Interestingly, root to shoot movement of sulphate under drought stress affects leaf ABA biosynthesis and stomatal closure via S-incorporation into the amino acid cysteine (Batool et al. 2018; Cao et al. 2014). As *AtMYB93* is expressed only in

a very few cells of the root (Gibbs et al. 2014) it may be mediating localised changes in or responses to S levels in the root that affect responses to the environment.

***AtMYB93*, S stress and root development.**

*At*MYB93 is a known negative regulator of lateral root development (Gibbs et al. 2014) that is upregulated by S stress ((Bielecka et al. 2014); this work). Previous research has shown that S stress can reduce LR development (Dan et al. 2007; Dong et al. 2019; Joshi et al. 2019) and enhance primary root elongation (Zhao et al. 2014). Although we hypothesised that these changes in root development upon S stress would be mediated by *Atmyb93*, we did not see clear differences in S deprivation root phenotypes in wild type or *Atmyb93* plants in our experiments. Previous experiments showing LR changes in wild type plants either measured lateral root number (rather than density) and lateral root length (Joshi et al. 2019), or changes were measured only in the context of S resupply (Dong et al. 2019), or in much older (15 day) seedings (Dan et al. 2007). This suggests that S regulation of LR development may have context-dependent nuances, only some of which involve *At*MYB93, or involve *At*MYB93 working partly in concert with other related R2R3 MYBs. Moreover, S deprivation does not extend the localisation of *pMYB93::GUS* expression beyond those endodermal cells overlying lateral root primordia, similarly to what is seen with ABA (Gibbs et al. 2014), suggesting that the extent of *AtMYB93* promoter activity is tightly restricted even under stress conditions.

S stress promotes flavonoid biosynthesis and increases anthocyanin levels ((Dong et al. 2019; Bielecka et al. 2014; Nikiforova et al. 2003)). GO terms for phenylpropanoid- and flavonoid biosynthesis are enriched in the genes downregulated in the *Atmyb93* mutant compared to wild type (Fig. 1a). The key flavonoid regulator *AtMYB12* (Mehrtens et al. 2005) is downregulated In the *Atmyb93* mutant, along with genes in the flavonoid biosynthesis pathway (*CYP75B1* (a flavonoid 3' hydroxylase), *UGT78D2*, *UGT84A1*, *UGT84A2*, *UGT90A1*; Online Resource 2). Moreover, *AtMYBL2*, an inhibitor of anthocyanin biosynthesis (Dubos et al. 2008; Matsui et al. 2008), is upregulated in the *Atmyb93* mutant (Online Resource 2). These data suggest that *Atmyb93* mutant roots would contain reduced flavonoids including anthocyanins. These gene expression changes could at least in part explain the increased root phenotypes of the *Atmyb93* mutants as flavonoids inhibit lateral root development (Brown et al. 2001; Chapman and Muday 2021) while

*At*MYB12 is known to regulates cell division orientation and cell differentiation in the

root (Wybouw et al. 2023).

Phenylpropanoids (which include flavonoids and lignin precursors) are required for

both lignin- and suberin production and maintenance in the endodermis (Andersen et

al. 2021). This could explain the role of *AtMYB93* in suberin deposition. Moreover,

the transcription factor DEWAX, a negative regulator of very long chain fatty acid

(VLCFA) biosynthesis (Go et al. 2014), is upregulated in the *Atmyb93* mutant (Online

Resource 2) suggesting that *At*MYB93 could affect VLCFA levels surrounding the

root cells in which it is expressed. However, neither the quadruple

AtMYB41/53/92/93 mutant with highly reduced suberin nor the *Atmyb92* mutant (with

reduced suberin (Klein 2019)) has a lateral root phenotype (Gibbs et al. 2014;

Shukla et al. 2021), arguing against a clear role for suberin itself in LR development.

AtMYB12 is downregulated in the *Atmyb39*/*suberman* mutant (Cohen et al. 2020),

but the lateral root phenotype of this mutant is not known.

Interestingly, in our experiments S deprivation increases adventitious root formation

at the root-hypocotyl junction. To our knowledge, this is the first time that S

deprivation has been shown to affect adventitious rooting at the collet in *Arabidopsis*.

S stress reduces crown root formation in rice (Grewal et al. 2018) most likely via a

non-*MYB93* mechanism as *OsMYB93*s are not upregulated by S stress (Fig. 3).

Flavonoids are known to inhibit AR formation in the hypocotyl in the presence of

auxin (Correa Lda et al. 2012) although a mutant lacking flavonoids makes fewer

ARs than wild type (Correa Lda et al. 2012). Whether the *Atmyb93* adventitious root

sensitivity is linked to flavonoids is unknown.

***AtMYB93*, shoot element composition, root architecture and root barriers.**

The *Atmyb93* mutant shows a general trend of elevated shoot elements, significantly

S, Mg and B, while two tomato *SlMYB93*-overexpressing lines show the opposite

trend. The third *SlMYB93*-overexpressing line broadly resembles wild type plants,

perhaps due to position-specific insertion effects of the transgene (Schnell et al.

2015).

The role of suberin on S uptake and assimilation is not fully understood. S

deprivation increases and extends suberisation in the root leading to fewer passage

cells (Barberon et al. 2016; Ogden et al. 2018) and loss of endodermal suberin

enhances the S deficiency phenotype of an S transporter mutant (Barberon et al.

2016) emphasising the requirement of an in-tact endodermal barrier for S uptake and retention. However, ionomic analysis shows shoot sulphate ion composition either not changing or being elevated in suberin-deficient transgenic plants made by expressing a suberin-degrading enzyme in the endodermis (*ELTP::CDEF1* and *CASP1::CDEF1* respectively, (Barberon et al. 2016)). Furthermore, another ionomic analysis shows no change in S in *ELTP::CDEF1* plants, a slight S reduction in the *Atmyb41/Atmyb53/Atmyb92/Atmyb93* quadruple mutant plants and no change/a slight reduction in S in plants overexpressing *AtMYB41* in the endodermis (*ELTP::MYB41* and *CDEF1::MYB41* respectively, (Shukla et al. 2021)). In both studies, the profiles of Mg and B are more in-line with our *Atmyb93* mutant data, being generally elevated with reduced suberin and reduced with elevated suberin ((Barberon et al. 2016; Shukla et al. 2021); Fig. 4). Plants with elevated suberin due to *AtMYB39* overexpression show reduced shoot elemental S (and several other elements), similarly to our *SlMYB93* overexpression data, but the *Atmyb39* mutant does not show elevated S (Cohen et al. 2020). Conversely, the enhanced suberin mutant *esb1* has elevated shoot sulphate levels (Baxter et al. 2009). Together, these data imply that changes in suberin are not the primary driver of the changes in shoot S seen in the *Atmyb93* mutant: these changes are more readily explained by alterations in root architecture and/or changes in S assimilation and transport caused by the perturbed *Atmyb93* transcriptome. Although *SULTR3;5* is downregulated in the *Atmyb39* mutant as in *Atmyb93*, *AtSULTR3;5* and another S transporter *AtSULTR1;2* are both downregulated in an *At*MYB39-overexpressing line and there is no other S-related overlap between the *Atmyb93* and *Atmyb39* mutant transcriptomes (Cohen et al. 2020). Importantly, the fatty acid elongation required for suberin biosynthesis requires acetyl coenzyme A (acetyl coA), a S-containing compound (Woolfson et al. 2022) suggesting a possible link between suberin biosynthesis and S metabolism.

We suggest that *AtMYB93* is indirectly involved, via phenylpropanoid metabolism, in increasing the amount and extent of suberisation specifically in its highly localised regions of expression during S deprivation, whilst also additionally responding to S levels to change S transport and metabolism.

Conclusion and future directions.

We have shown that MYB93 regulates responses to S levels. The *Atmyb93*

transcriptome shows a S stress signature, and MYB93 homologues in two dicot

plants (*Arabidopsis* and tomato) are upregulated by S deprivation. *Arabidopsis* and

tomato with perturbed *MYB93* levels show changes in shoot S content. S deprivation

changes root architecture in *Arabidopsis* but this is not mediated solely by *AtMYB93*.

*At*MYB93's effect on lateral root development is likely via flavonoids rather than

suberin as there is no evidence for direct suberin regulation by *At*MYB93: future work

will investigate this link further.

Materials and Methods

Plant materials and growth conditions.

Arabidopsis ecotype Col-0 wild type, *Atmyb93* mutant (Gibbs et al. 2014), *Atmpk3-1*

mutant (SALK_151594, NASC ID N869692; (Merkouropoulos et al. 2008)) and

pAtMYB93::GUS transgenic lines (Gibbs et al. 2014) were grown in the glasshouse

in Levington M3 compost/vermiculite mix at 22˚C under 16h light. For growth on

plates, Col-0, *Atmyb93*, *Atmpk3* seeds were sterilised for 10 minutes in 10%

Parozone™ Bleach (Jeyes, Hemel Hempstead, UK) followed by 3 rinses in sterile

distilled water and resuspension in 200µl distilled water. Seeds were vernalised at

4˚C for 2 days in the dark and plated in rows to grow vertically at the top of half-

strength Murashige and Skoog (MS) medium (M0404, Sigma-Aldrich, St Louis,

Missouri, USA) pH5.6-5.8 with 1% agar.

For RNAseq analysis, the medium used was Sigma-Aldrich MS M0404 and

seedlings were grown for 7 days. For root assays of Col-0, *Atmyb93* and *Atmpk3*

(comparing genotypes, or OAS treatment) the medium used was half-strength MS

(Sigma-Aldrich M0404), with seedlings grown for 8 days.

For S deprivation root assay experiments, the medium used was half-strength MS

either containing (+S) or not containing (-S) S (MS basal salts MSP01 and MSP44;

Caisson Labs, Smithfield, Utah USA) with 1% agar, with seedlings grown for 8 days.

For some experiments, seedlings were initially germinated and grown on +S for 5

days and then transferred to -S for 3 days, using sterile forceps in a sterile laminar

flow cabinet. All plates were sealed with micropore tape (3M, St Paul, Minnesota,

USA).

For *Arabidopsis* fresh weight and shoot element analysis, Col-0 and *Atmyb93* were

grown in Magenta pots (Sigma) containing 4 evenly spaced seedlings per pot. Pots

were filled to half capacity (125ml) with 0.5 strength MS made using the Merck

classic protocol (KGaA 2018) with 1% sucrose and 0.7% agar (Sigma-Aldrich CAS

9002-18-0). For medium without S, all sulphate ions were replaced with the chloride

equivalent. Foil was wrapped round the part of the pot with agar to mimic soil

darkness and facilitate root penetration of the agar. Pots were sealed with micropore

tape (3M).

Solanum lycopersicum cultivar Micro-Tom was grown in an incubator (24°C) or

glasshouse (22°C) in Levington M3 compost/vermiculite mix under 16h light. Seeds

were sown directly onto compost unless transformation was to be carried out (see

tomato transformation section); soil-grown plants were used for elemental analysis,

RT-PCR analysis and cloning (roots were washed before making RNA).

RNA extraction and cDNA generation

14 Plant tissue was ground in liquid nitrogen using RNAse-free ceramic pestles and

15 mortars and RNA extraction was performed using an ISOLATE II Plant RNA kit

16 (Bioline, Meridian Biosciences, Memphis, TN, USA).

Transcriptome sequencing and analysis

RNA from ~30mg of tissue per sample of whole 7-day old roots of wild type and

Atmyb93 (triplicate samples) was extracted and quality checked by both calculating

the OD260/280 and OD 260/230 ratios of a 1μl sample using a Nanodrop analyser

and by agarose gel electrophoresis to visualise ribosomal RNA integrity (Online

Resource 1). Samples were sent to Novogene for further quality control (Agilent

2100 analysis), library preparation and sequencing. Briefly, sample mRNA was

enriched using oligo(dT) coupled beads and RNA was fragmented using buffer

containing 0.1mM $ZnCl_2$. cDNA libraries were synthesised from mRNA fragments

using random hexamers and reverse transcriptase followed by second strand cDNA

synthesis using dNTPs, RNaseH and *E. coli* DNA polymerase I. The library was

finished by carrying out end-repair, A-tailing, adaptor ligation, size selection and

amplification by PCR. Library insert size and concentration was checked before

Illumina sequencing. Raw data were transformed by base calling into sequenced

reads (including sequence information and sequence quality information), which

were exported in FASTQ format. For quality control, Phred scores were used to

estimate the error rate of sequenced reads. Raw reads were filtered to remove

adapters and low-quality reads. Cleaned FASTQ data were aligned to the

Arabidopsis reference genome using HISAT2 (Kim et al. 2019). Clean, mapped

fragments representing transcript abundance were used to estimate the expression

level of each gene in each sample via fragments per kilobase of transcript sequence

per million fragments paired sequenced (FKPM) values using HTseq software in

union mode (Anders et al. 2015). Normalised FKPM values were subject to

Pearson's correlation analysis to estimate similarity between biological repeats.

Differentially expressed genes (DEGs) were identified using DEseq2 software with

the DEseq normalisation method with a false discovery rate (FDR) < 0.05. GO

enrichment analysis was carried out separately on down- and up-regulated DEGs,

using the GO enrichment tool in PlantRegMap (Tian et al. 2020) with a cutoff of

p<0.01. Enriched GO terms were visualised using ReviGo (Supek et al. 2011).

Phylogeny and *in silico* expression analysis of MYB93-related genes.

AtMYB93-related protein sequences were selected as follows: (i) the *Arabidopsis*

genes known to be involved in suberin formation (refs) that are (ii) also most closely

related at the protein sequence level to *At*MYB93 (Clades S24, S10, S11, S42; Du et

al 2015), (iii) sequences from representative dicots and monocots identified as being

most similar to *At*MYB93 by BLASTP (Altschul et al), mostly from species for which

published expression data was also available. Chosen species were *Brassica napus*,

Brassica rapa, *Glycine max* (soybean), *Populus trichocarpa* (poplar), *Manihot*

esculenta (cassava), *Solanum lycopersicum* (tomato), *Malus domestica* (apple),

Oryza sativa (rice), *Zea mays* (maize), *Brachypodium distachyon*, *Hordeum vulgare*

(barley). Sequences were obtained from NCBI, Phytozome and BrassicaDB

(http://brassicadb.cn/#/). Sequences were aligned using MUSCLE

(https://www.ebi.ac.uk/jdispatcher/msa) and a simple Neighbour-Joining tree was

generated, which was visualized using FigTree

(http://tree.bio.ed.ac.uk/software/figtree/).

Expression data was extracted from a combination of the ePlant (Waese et al. 2017)

and the *Arabidopsis* eFP browser RootII dataset (Iyer-Pascuzzi et al. 2011; Winter et

al. 2007) and papers where expression of MYB93-related genes or whole-plant

responses to S deprivation are recorded (Bielecka et al. 2014; Canales et al. 2020;

Du et al. 2012; Jain et al. 2007; Kajala et al. 2021; Libault et al. 2009; Sekhon et al.

2011; Wang et al. 2022; Wilkins et al. 2009; Wilson et al. 2017). Data was visualised semi-quantitatively using colour intensity to provide an overall summary of trends.

RT-PCR and semi quantitative RT-PCR.

For cloning and RT-PCR in tomato, cDNA synthesis was performed from RNA using the SuperScript™ III first-strand synthesis system (Invitrogen, ThermoFisher Scientific, Waltham, MA, USA). For expression analysis of *SlMYB93* homologues, PCR was carried out on the cDNA using Taq polymerase using primers in Online Resource 2.

Generation of an *SlMYB93* overexpression construct and transformation into *Arabidopsis* and tomato.

Full-length *SlMYB93* (*Solyc04g074170*) cDNA was cloned from tomato roots cv. Ailsa Craig using a proofreading DNA polymerase (Phusion™, New England Biolabs, Ipswich, MA, USA) and relevant primers (Online Resource 2) into a modified pBI121 vector (Clontech), mpBI12135S, with modifications to the left and right border (Yongsheng Liu personal communication; (Cao et al. 2012)). mpBI121 contains additional restriction sites compared to pBI121, including *Kpn*I, *Sal*I and *Xho*I, and the *GUS* reporter gene has been removed, so that only the NOS terminator remains at the end of the multiple cloning site. mpBI121 contains the *nptII* kanamycin resistance gene for selection in plants. *SlMYB93* in mpBI121 was transformed into electrocompetent *Agrobacterium* strain GV3101 by electroporation at 2.5kV, 25μF capacitance and 400 Ω resistance.The *p35S::SlMYB93* construct was transformed into *Arabidopsis* via floral dip (Clough and Bent 1998).

For transformation of *p35S::SlMYB93* into tomato, *Solanum lycopersicum* cv. Micro Tom seeds were imbibed in water for 1h at 37°C and sterilised by soaking in 20% Parozone™ (Jeyes, Hemel Hempstead, UK) bleach for 20 min followed by 3 washes with 30ml autoclaved water and drying on autoclaved filter paper in a laminar flow hood. Seeds were then rinsed in sterile water at least 3 further times in fresh tubes to completely remove all bleach residues and air-dried. Seeds were placed on half-strength MS basal salts (M5524, Sigma-Aldrich, St Louis, Missouri, USA) containing 2% sucrose in Magenta pots (Sigma-Aldrich, St Louis, Missouri, USA) and cold-treated (4°C) in the dark for at 2-3 days before moving to a growth room at 22°C in

the dark for 2 days. If most seeds germinated, pots were transferred to a 16h light

/8h dark cycle for 4-5 days. From these plants, tomato transformation was carried

out as in (Cao 2022). Briefly, cotyledons were wounded and transformed with

Agrobacterium containing *p35S::SlMYB93* before rounds of kanamycin selection to

generate transformed callus producing shoots, followed by transfer to rooting

medium and further transfer of rooted, transformed plants to soil.

Shoot element analysis of *Arabidopsis* ICP-MS

Arabidopsis plants were grown in Magenta pots as detailed previously and harvested

using tweezers at 21 days, before flowering had begun. Roots were removed with a

razor blade and pooled fresh shoot tissue was heated at 60°C for 48h in an oven to

remove all water. At least 100mg of dry shoot tissue for each condition was analysed

by ICP-MS as in (Thomas et al. 2016). Significant differences were identified by

Mann Whitney U-tests comparing *Atmyb93* mutants to wild type.

Shoot element analysis of tomato by ICP-OES

For tomato shoot element analysis, ~20g fresh weight of young leaves (the first 10-

12 leaves behind the growing tip) were harvested from ~6-week old plants of each

genotype, which were dried at 50°C in an oven for 48h. At least 0.2g dry weight of

each type of leaf tissue was sent for analysis by inductively coupled plasma optical

emission spectroscopy (ICP-OES) at Yara Analytical Services, York, UK.

Briefly, samples were dried, milled, and sieved prior to analysis. Samples were

digested in nitric acid using a microwave digester (CEM, Buckingham, UK). The

digested samples were analysed on an Agilent ICP 5900 ICP-OES instrument

(Agilent Technologies LDA UK Ltd, Stockport, UK) calibrated against standards of

known concentrations. Significant differences between wild type and *SlMYB93-*

overexpressing lines were identified by Mann Whitney U- tests.

***Arabidopsis* root assay quantification.**

Arabidopsis seedlings growing vertically on +S or -S medium were photographed.

Emerged lateral roots and adventitious roots (roots emerging from the collet) were

counted by eye from plates and root length was measured from photographs using

the freehand drawing tool in ImageJ (https://imagej.net/ij/). Lateral root density was

calculated for each seedling by dividing lateral root number by primary root length.

All root data was visualised using SuperPlots ((Lord et al. 2020);

https://huygens.science.uva.nl/SuperPlotsOfData/) Statistical significance was

calculated via a Kruskal-Wallis test with a post-hoc Dunn's test, or by a Mann-

Whitney U-test for pairwise comparisons.

***AtMYB93* promoter activity.**

Seedlings expressing *pAtMYB93::GUS* (Gibbs et al. 2014) were grown vertically on

0.5MS (KGaA 2018) plates with or without S and with 1% agar (Sigma-Aldrich CAS

9002-18-0) for 5-8 days. GUS assays were carried out as in Gibbs et al (2014).

Images were acquired by light microscopy using a GXCAM-HiChrome-S camera (GT

vision, Stansfield, UK).

References

Aarabi F, Kusajima M, Tohge T, Konishi T, Gigolashvili T, Takamune M, Sasazaki Y, Watanabe M, Nakashita H, Fernie AR, Saito K, Takahashi H, Hubberten HM, Hoefgen R, Maruyama-Nakashita A (2016) Sulfur deficiency-induced repressor proteins optimize glucosinolate biosynthesis in plants. Sci Adv 2 (10):e1601087. doi:10.1126/sciadv.1601087

Ali MM, Shafique MW, Gull S, Naveed WA, Javed T, Ahmed FY, Mauro RP (2021) Alleviation of Heat Stress in Tomato by Exogenous Application of Sulfur. Horticulturae 7 (2). doi:https://doi.org/10.3390/horticulturae7020021

Anders S, Pyl PT, Huber W (2015) HTSeq--a Python framework to work with high-throughput sequencing data. Bioinformatics 31 (2):166-169. doi:10.1093/bioinformatics/btu638

Andersen TG, Barberon M, Geldner N (2015) Suberization - the second life of an endodermal cell. Curr Opin Plant Biol 28:9-15. doi:10.1016/j.pbi.2015.08.004

Andersen TG, Molina D, Kilian J, Franke RB, Ragni L, Geldner N (2021) Tissue-Autonomous Phenylpropanoid Production Is Essential for Establishment of Root Barriers. Curr Biol 31 (5):965-977 e965. doi:10.1016/j.cub.2020.11.070

Barberon M, Vermeer JE, De Bellis D, Wang P, Naseer S, Andersen TG, Humbel BM, Nawrath C, Takano J, Salt DE, Geldner N (2016) Adaptation of Root Function by Nutrient-Induced Plasticity of Endodermal Differentiation. Cell 164 (3):447-459. doi:10.1016/j.cell.2015.12.021

Batool S, Uslu VV, Rajab H, Ahmad N, Waadt R, Geiger D, Malagoli M, Xiang CB, Hedrich R, Rennenberg H, Herschbach C, Hell R, Wirtz M (2018) Sulfate is Incorporated into Cysteine to Trigger ABA Production and Stomatal Closure. Plant Cell 30 (12):2973-2987. doi:10.1105/tpc.18.00612

Baxter I, Hosmani PS, Rus A, Lahner B, Borevitz JO, Muthukumar B, Mickelbart MV, Schreiber L, Franke RB, Salt DE (2009) Root suberin forms an extracellular barrier that affects water relations and mineral nutrition in Arabidopsis. PLoS Genet 5 (5):e1000492. doi:10.1371/journal.pgen.1000492

Bielecka M, Watanabe M, Morcuende R, Scheible WR, Hawkesford MJ, Hesse H, Hoefgen R (2014) Transcriptome and metabolome analysis of plant sulfate starvation and resupply provides novel information on transcriptional regulation of metabolism associated with sulfur, nitrogen and phosphorus nutritional responses in Arabidopsis. Front Plant Sci 5:805. doi:10.3389/fpls.2014.00805

Brown DE, Rashotte AM, Murphy AS, Normanly J, Tague BW, Peer WA, Taiz L, Muday GK (2001) Flavonoids act as negative regulators of auxin transport in vivo in arabidopsis. Plant Physiol 126 (2):524-535. doi:10.1104/pp.126.2.524

Canales J, Uribe F, Henriquez-Valencia C, Lovazzano C, Medina J, Vidal EA (2020) Transcriptomic analysis at organ and time scale reveals gene regulatory networks controlling the sulfate starvation response of Solanum lycopersicum. BMC Plant Biol 20 (1):385. doi:10.1186/s12870-020-02590-2

Canto-Pastor A, Kajala K, Shaar-Moshe L, Manzano C, Timilsena P, De Bellis D, Gray S, Holbein J, Yang H, Mohammad S, Nirmal N, Suresh K, Ursache R, Mason GA, Gouran M, West DA, Borowsky AT, Shackel KA, Sinha N, Bailey-Serres J, Geldner N, Li S, Franke RB, Brady SM (2024) A suberized exodermis is required for tomato drought tolerance. Nat Plants 10 (1):118-130. doi:10.1038/s41477-023-01567-x

Cao MJ, Wang Z, Zhao Q, Mao JL, Speiser A, Wirtz M, Hell R, Zhu JK, Xiang CB (2014) Sulfate availability affects ABA levels and germination response to ABA and salt stress in Arabidopsis thaliana. Plant J 77 (4):604-615. doi:10.1111/tpj.12407

Cao X (2022) Understanding gene and protein function of MYB93 in tomato and Arabidopsis. University of Birmingham, Birmingham, UK

Cao Y, Tang X, Giovannoni J, Xiao F, Liu Y (2012) Functional characterization of a tomato COBRA-like gene functioning in fruit development and ripening. BMC Plant Biol 12:211. doi:10.1186/1471-2229-12-211

Chandra N, Pandey N (2014) Influence of Sulfur Induced Stress on Oxidative Status and Antioxidative Machinery in Leaves of Allium cepa L. Int Sch Res Notices 2014:568081. doi:10.1155/2014/568081

Chapman JM, Muday GK (2021) Flavonols modulate lateral root emergence by scavenging reactive oxygen species in Arabidopsis thaliana. J Biol Chem 296:100222. doi:10.1074/jbc.RA120.014543

Clough SJ, Bent AF (1998) Floral dip: a simplified method for Agrobacterium-mediated transformation of Arabidopsis thaliana. Plant J 16 (6):735-743. doi:10.1046/j.1365-313x.1998.00343.x

Cohen H, Fedyuk V, Wang C, Wu S, Aharoni A (2020) SUBERMAN regulates developmental suberization of the Arabidopsis root endodermis. Plant J 102 (3):431-447. doi:10.1111/tpj.14711

Correa Lda R, Troleis J, Mastroberti AA, Mariath JE, Fett-Neto AG (2012) Distinct modes of adventitious rooting in Arabidopsis thaliana. Plant Biol (Stuttg) 14 (1):100-109. doi:10.1111/j.1438-8677.2011.00468.x

Criollo-Arteaga S, Moya-Jimenez S, Jimenez-Meza M, Gonzalez-Vera V, Gordon-Nunez J, Llerena-Llerena S, Ramirez-Villacis DX, van 't Hof P, Leon-Reyes A (2021) Sulfur Deprivation Modulates Salicylic Acid Responses via Nonexpressor of Pathogenesis-Related Gene 1 in Arabidopsis thaliana. Plants (Basel) 10 (6). doi:10.3390/plants10061065

Dan H, Yang G, Zheng ZL (2007) A negative regulatory role for auxin in sulphate deficiency response in Arabidopsis thaliana. Plant Mol Biol 63 (2):221-235. doi:10.1007/s11103-006-9084-0

Dietzen C, Koprivova A, Whitcomb SJ, Langen G, Jobe TO, Hoefgen R, Kopriva S (2020) The Transcription Factor EIL1 Participates in the Regulation of Sulfur-Deficiency Response. Plant Physiol 184 (4):2120-2136. doi:10.1104/pp.20.01192

Dong W, Wang Y, Takahashi H (2019) CLE-CLAVATA1 Signaling Pathway Modulates Lateral Root Development under Sulfur Deficiency. Plants (Basel) 8 (4). doi:10.3390/plants8040103

Du H, Liang Z, Zhao S, Nan MG, Tran LS, Lu K, Huang YB, Li JN (2015) The Evolutionary History of R2R3-MYB Proteins Across 50 Eukaryotes: New Insights Into Subfamily Classification and Expansion. Sci Rep 5:11037. doi:10.1038/srep11037

Du H, Yang SS, Liang Z, Feng BR, Liu L, Huang YB, Tang YX (2012) Genome-wide analysis of the MYB transcription factor superfamily in soybean. BMC Plant Biol 12:106. doi:10.1186/1471-2229-12-106

Dubos C, Le Gourrierec J, Baudry A, Huep G, Lanet E, Debeaujon I, Routaboul JM, Alboresi A, Weisshaar B, Lepiniec L (2008) MYBL2 is a new regulator of flavonoid biosynthesis in Arabidopsis thaliana. Plant J 55 (6):940-953. doi:10.1111/j.1365-313X.2008.03564.x

Elmore JS, Dodson AT, Muttucumaru N, Halford NG, Parry MAJ, Mottram DS (2010) Effects of sulphur nutrition during potato cultivation on the formation of acrylamide and aroma compounds during cooking. Food Chemistry 122 (3):753-760

Elmore JS, Mottram DS, Muttucumaru N, Dodson AT, Parry MAJ, Halford NG (2007) Changes in Free Amino Acids and Sugars in Potatoes Due to Sulfate Fertilization and the Effect on Acrylamide Formation. J Agric Food Chem 55:5363-5366

Forieri I, Sticht C, Reichelt M, Gretz N, Hawkesford MJ, Malagoli M, Wirtz M, Hell R (2017) System analysis of metabolism and the transcriptome in Arabidopsis thaliana roots reveals differential co-regulation upon iron, sulfur and potassium deficiency. Plant Cell Environ 40 (1):95-107. doi:10.1111/pce.12842

Frerigmann H, Gigolashvili T (2014a) MYB34, MYB51, and MYB122 distinctly regulate indolic glucosinolate biosynthesis in Arabidopsis thaliana. Mol Plant 7 (5):814-828. doi:10.1093/mp/ssu004

Frerigmann H, Gigolashvili T (2014b) Update on the role of R2R3-MYBs in the regulation of glucosinolates upon sulfur deficiency. Front Plant Sci 5:626. doi:10.3389/fpls.2014.00626

Geldner N (2013) The endodermis. Annu Rev Plant Biol 64:531-558. doi:10.1146/annurev-arplant-050312-120050

Gibbs DJ, Voss U, Harding SA, Fannon J, Moody LA, Yamada E, Swarup K, Nibau C, Bassel GW, Choudhary A, Lavenus J, Bradshaw SJ, Stekel DJ, Bennett MJ, Coates JC (2014) AtMYB93 is a novel negative regulator of lateral root development in Arabidopsis. New Phytol 203 (4):1194-1207. doi:10.1111/nph.12879

Go YS, Kim H, Kim HJ, Suh MC (2014) Arabidopsis Cuticular Wax Biosynthesis Is Negatively Regulated by the DEWAX Gene Encoding an AP2/ERF-Type Transcription Factor. Plant Cell 26 (4):1666-1680. doi:10.1105/tpc.114.123307

Gou M, Hou G, Yang H, Zhang X, Cai Y, Kai G, Liu CJ (2017) The MYB107 Transcription Factor Positively Regulates Suberin Biosynthesis. Plant Physiol 173 (2):1045-1058. doi:10.1104/pp.16.01614

Grant CA, Mahli SS, Karamanos RE (2012) Sulfur management for rapeseed. Field Crops Research 128:119-128

Grewal RK, Saraf S, Deb A, Kundu S (2018) Differentially Expressed MicroRNAs Link Cellular Physiology to Phenotypic Changes in Rice Under Stress Conditions. Plant Cell Physiol 59 (10):2143-2154. doi:10.1093/pcp/pcy136

Henriquez-Valencia C, Arenas MA, Medina J, Canales J (2018) Integrative Transcriptomic Analysis Uncovers Novel Gene Modules That Underlie the Sulfate Response in Arabidopsis thaliana. Front Plant Sci 9:470. doi:10.3389/fpls.2018.00470

Hirai MY, Fujiwara T, Awazuhara M, Kimura T, Noji M, Saito K (2003) Global expression profiling of sulfur-starved Arabidopsis by DNA macroarray reveals the role of O-acetyl-l-serine as a general regulator of gene expression in response to sulfur nutrition. Plant J 33 (4):651-663. doi:10.1046/j.1365-313x.2003.01658.x

Hu H (2018) The role of transcription factor MYB53 from Arabidopsis thaliana in the regulated production of suberin. Carleton University,

Huang X, Li Y, Chang Z, Yan W, Xu C, Zhang B, He Z, Wang C, Zheng M, Li Z, Xia J, Li G, Tang X, Wu J (2024) Regulation by distinct MYB transcription factors defines the roles of OsCYP86A9 in anther development and root suberin deposition. Plant J. doi:10.1111/tpj.16722

Hubberten HM, Drozd A, Tran BV, Hesse H, Hoefgen R (2012a) Local and systemic regulation of sulfur homeostasis in roots of Arabidopsis thaliana. Plant J 72 (4):625-635. doi:10.1111/j.1365-313X.2012.05105.x

Hubberten HM, Klie S, Caldana C, Degenkolbe T, Willmitzer L, Hoefgen R (2012b) Additional role of O-acetylserine as a sulfur status-independent regulator during plant growth. Plant J 70 (4):666-677. doi:10.1111/j.1365-313X.2012.04905.x

Iyer-Pascuzzi AS, Jackson T, Cui H, Petricka JJ, Busch W, Tsukagoshi H, Benfey PN (2011) Cell identity regulators link development and stress responses in the Arabidopsis root. Dev Cell 21 (4):770-782. doi:10.1016/j.devcel.2011.09.009

Jain M, Nijhawan A, Arora R, Agarwal P, Ray S, Sharma P, Kapoor S, Tyagi AK, Khurana JP (2007) F-box proteins in rice. Genome-wide analysis, classification, temporal and spatial gene expression during panicle and seed development, and regulation by light and abiotic stress. Plant Physiol 143 (4):1467-1483. doi:10.1104/pp.106.091900

Joshi NC, Meyer AJ, Bangash SAK, Zheng ZL, Leustek T (2019) Arabidopsis gamma-glutamylcyclotransferase affects glutathione content and root system architecture during sulfur starvation. New Phytol 221 (3):1387-1397. doi:10.1111/nph.15466

Kajala K, Gouran M, Shaar-Moshe L, Mason GA, Rodriguez-Medina J, Kawa D, Pauluzzi G, Reynoso M, Canto-Pastor A, Manzano C, Lau V, Artur MAS, West DA, Gray SB, Borowsky AT, Moore BP, Yao AI, Morimoto KW, Bajic M, Formentin E, Nirmal NA, Rodriguez A, Pasha A, Deal RB, Kliebenstein DJ, Hvidsten TR, Provart NJ, Sinha NR, Runcie DE, Bailey-Serres J, Brady SM (2021) Innovation, conservation, and repurposing of gene function in root cell type development. Cell 184 (12):3333-3348 e3319. doi:10.1016/j.cell.2021.04.024

Kataoka T, Hayashi N, Yamaya T, Takahashi H (2004) Root-to-shoot transport of sulfate in Arabidopsis. Evidence for the role of SULTR3;5 as a component of low-affinity sulfate transport system in the root vasculature. Plant Physiol 136 (4):4198-4204. doi:10.1104/pp.104.045625

KGaA M (2018) Classical Plant Media Formulations. https://www.sigmaaldrich.com/GB/en/technical-documents/technical-article/cell-culture-and-cell-culture-analysis/plant-tissue-culture/classic-plant-media.

Kim D, Paggi JM, Park C, Bennett C, Salzberg SL (2019) Graph-based genome alignment and genotyping with HISAT2 and HISAT-genotype. Nat Biotechnol 37 (8):907-915. doi:10.1038/s41587-019-0201-4

Klein D (2019) The downstream targets of MYB-type transcription factors involved in suberin biosynthesis. Carleton University, Ottawa, Canada

Kopriva S, Malagoli M, Takahashi H (2019) Sulfur nutrition: impacts on plant development, metabolism, and stress responses. J Exp Bot 70 (16):4069-4073. doi:10.1093/jxb/erz319

Kosma DK, Murmu J, Razeq FM, Santos P, Bourgault R, Molina I, Rowland O (2014) AtMYB41 activates ectopic suberin synthesis and assembly in multiple plant species and cell types. Plant J 80 (2):216-229. doi:10.1111/tpj.12624

Kutz A, Muller A, Hennig P, Kaiser WM, Piotrowski M, Weiler EW (2002) A role for nitrilase 3 in the regulation of root morphology in sulphur-starving Arabidopsis thaliana. Plant J 30 (1):95-106. doi:10.1046/j.1365-313x.2002.01271.x

Lashbrooke J, Cohen H, Levy-Samocha D, Tzfadia O, Panizel I, Zeisler V, Massalha H, Stern A, Trainotti L, Schreiber L, Costa F, Aharoni A (2016) MYB107 and MYB9 Homologs Regulate Suberin Deposition in Angiosperms. Plant Cell 28 (9):2097-2116. doi:10.1105/tpc.16.00490

Legay S, Guerriero G, Andre C, Guignard C, Cocco E, Charton S, Boutry M, Rowland O, Hausman JF (2016) MdMyb93 is a regulator of suberin deposition in russeted apple fruit skins. New Phytol 212 (4):977-991. doi:10.1111/nph.14170

Li Y, Sawada Y, Hirai A, Sato M, Kuwahara A, Yan X, Hirai MY (2013) Novel insights into the function of Arabidopsis R2R3-MYB transcription factors regulating aliphatic glucosinolate biosynthesis. Plant Cell Physiol 54 (8):1335-1344. doi:10.1093/pcp/pct085

Libault M, Joshi T, Takahashi K, Hurley-Sommer A, Puricelli K, Blake S, Finger RE, Taylor CG, Xu D, Nguyen HT, Stacey G (2009) Large-scale analysis of putative soybean regulatory gene expression identifies a Myb gene involved in soybean nodule development. Plant Physiol 151 (3):1207-1220. doi:10.1104/pp.109.144030

Lord SJ, Velle KB, Mullins RD, Fritz-Laylin LK (2020) SuperPlots: Communicating reproducibility and variability in cell biology. J Cell Biol 219 (6). doi:10.1083/jcb.202001064

Lunde C, Zygadlo A, Simonsen HT, Nielsen PL, Blennow A, Haldrup A (2008) Sulfur starvation in rice: the effect on photosynthesis, carbohydrate metabolism, and oxidative stress protective pathways. Physiol Plant 134 (3):508-521. doi:10.1111/j.1399-3054.2008.01159.x

Maruyama-Nakashita A, Inoue E, Watanabe-Takahashi A, Yamaya T, Takahashi H (2003) Transcriptome profiling of sulfur-responsive genes in Arabidopsis reveals global effects of sulfur nutrition on multiple metabolic pathways. Plant Physiol 132 (2):597-605. doi:10.1104/pp.102.019802

Maruyama-Nakashita A, Nakamura Y, Tohge T, Saito K, Takahashi H (2006) Arabidopsis SLIM1 is a central transcriptional regulator of plant sulfur response and metabolism. Plant Cell 18 (11):3235-3251. doi:10.1105/tpc.106.046458

Matsui K, Umemura Y, Ohme-Takagi M (2008) AtMYBL2, a protein with a single MYB domain, acts as a negative regulator of anthocyanin biosynthesis in Arabidopsis. Plant J 55 (6):954-967. doi:10.1111/j.1365-313X.2008.03565.x

Mehrtens F, Kranz H, Bednarek P, Weisshaar B (2005) The Arabidopsis transcription factor MYB12 is a flavonol-specific regulator of phenylpropanoid biosynthesis. Plant Physiol 138 (2):1083-1096. doi:10.1104/pp.104.058032

Merkouropoulos G, Andreasson E, Hess D, Boller T, Peck SC (2008) An Arabidopsis protein phosphorylated in response to microbial elicitation, AtPHOS32, is a substrate of MAP kinases 3 and 6. J Biol Chem 283 (16):10493-10499. doi:10.1074/jbc.M800735200

Narayan OP, Kumar P, Yadav B, Dua M, Johri AK (2023) Sulfur nutrition and its role in plant growth and development. Plant Signal Behav 18 (1):2030082. doi:10.1080/15592324.2022.2030082

Nikiforova V, Freitag J, Kempa S, Adamik M, Hesse H, Hoefgen R (2003) Transcriptome analysis of sulfur depletion in Arabidopsis thaliana: interlacing of biosynthetic pathways provides response specificity. Plant J 33 (4):633-650. doi:10.1046/j.1365-313x.2003.01657.x

O'Malley RC, Huang SC, Song L, Lewsey MG, Bartlett A, Nery JR, Galli M, Gallavotti A, Ecker JR (2016) Cistrome and Epicistrome Features Shape the Regulatory DNA Landscape. Cell 166 (6):1598. doi:10.1016/j.cell.2016.08.063

Ogden M, Hoefgen R, Roessner U, Persson S, Khan GA (2018) Feeding the Walls: How Does Nutrient Availability Regulate Cell Wall Composition? Int J Mol Sci 19 (9). doi:10.3390/ijms19092691

Paxman PJ, Hill R (1974) The goitrogenicity of kale and its relation to thiocyanate content. J Sci Food Agric 25 (3):329-337. doi:10.1002/jsfa.2740250314

Robe K, Gao F, Bonillo P, Tissot N, Gaymard F, Fourcroy P, Izquierdo E, Dubos C (2020) Sulphur availability modulates Arabidopsis thaliana responses to iron deficiency. PLoS One 15 (8):e0237998. doi:10.1371/journal.pone.0237998

Sarda X, Diquelou S, Abdallah M, Nesi N, Cantat O, Le Gouee P, Avice JC, Ourry A (2013) Assessment of sulphur deficiency in commercial oilseed rape crops from plant analysis. The Journal of Agricultural Science 152 (4):616-633

Schnell J, Steele M, Bean J, Neuspiel M, Girard C, Dormann N, Pearson C, Savoie A, Bourbonniere L, Macdonald P (2015) A comparative analysis of insertional effects in genetically engineered plants: considerations for pre-market assessments. Transgenic Res 24 (1):1-17. doi:10.1007/s11248-014-9843-7

Sekhon RS, Lin H, Childs KL, Hansey CN, Buell CR, de Leon N, Kaeppler SM (2011) Genome-wide atlas of transcription during maize development. Plant J 66 (4):553-563. doi:10.1111/j.1365-313X.2011.04527.x

Serra O, Geldner N (2022) The making of suberin. New Phytol 235 (3):848-866. doi:10.1111/nph.18202

Sharma RK, Cox MS, Oglesby C, Dhillon JS (2024) Revisiting the role of sulfur in crop production: A narrative review. Journal of Agriculture and Food Research 15. doi:10.1016/j.jafr.2024.101013

Shukla V, Han JP, Cleard F, Lefebvre-Legendre L, Gully K, Flis P, Berhin A, Andersen TG, Salt DE, Nawrath C, Barberon M (2021) Suberin plasticity to developmental and exogenous cues is regulated by a set of MYB transcription factors. Proc Natl Acad Sci U S A 118 (39). doi:10.1073/pnas.2101730118

Smith RH (1980) Kale poisoning: the brassica anaemia factor. Vet Rec 107 (1):12-15. doi:10.1136/vr.107.1.12

Smith RH, Earl CR, Matheson NA (1974) The probable role of S-methylcysteine sulphoxide in kale poisoning in ruminants. Biochemical Society Transactions 2:101-104

Supek F, Bosnjak M, Skunca N, Smuc T (2011) REVIGO summarizes and visualizes long lists of gene ontology terms. PLoS One 6 (7):e21800. doi:10.1371/journal.pone.0021800

Thomas CL, Alcock TD, Graham NS, Hayden R, Matterson S, Wilson L, Young SD, Dupuy LX, White PJ, Hammond JP, Danku JM, Salt DE, Sweeney A, Bancroft I, Broadley MR (2016) Root morphology and seed and leaf ionomic traits in a Brassica napus L. diversity panel show wide phenotypic variation and are characteristic of crop habit. BMC Plant Biol 16 (1):214. doi:10.1186/s12870-016-0902-5

Tian F, Yang DC, Meng YQ, Jin J, Gao G (2020) PlantRegMap: charting functional regulatory maps in plants. Nucleic Acids Res 48 (D1):D1104-D1113. doi:10.1093/nar/gkz1020

To A, Joubes J, Thueux J, Kazaz S, Lepiniec L, Baud S (2020) AtMYB92 enhances fatty acid synthesis and suberin deposition in leaves of Nicotiana benthamiana. Plant J 103 (2):660-676. doi:10.1111/tpj.14759

Uemura Y, Kimura S, Ohta T, Suzuki T, Mase K, Kato H, Sakaoka S, Uefune M, Komine Y, Hotta K, Shimizu M, Morikami A, Tsukagoshi H (2023) A very long chain fatty acid responsive transcription factor, MYB93, regulates lateral root development in Arabidopsis. Plant J 115 (5):1408-1427. doi:10.1111/tpj.16330

Vishwanath SJ, Delude C, Domergue F, Rowland O (2015) Suberin: biosynthesis, regulation, and polymer assembly of a protective extracellular barrier. Plant Cell Rep 34 (4):573-586. doi:10.1007/s00299-014-1727-z

Waese J, Fan J, Pasha A, Yu H, Fucile G, Shi R, Cumming M, Kelley LA, Sternberg MJ, Krishnakumar V, Ferlanti E, Miller J, Town C, Stuerzlinger W, Provart NJ (2017) ePlant: Visualizing and Exploring Multiple Levels of Data for Hypothesis Generation in Plant Biology. Plant Cell 29 (8):1806-1821. doi:10.1105/tpc.17.00073

Wang C, Wang H, Li P, Li H, Xu C, Cohen H, Aharoni A, Wu S (2020) Developmental programs interact with abscisic acid to coordinate root suberization in Arabidopsis. Plant J 104 (1):241-251. doi:10.1111/tpj.14920

Wang RY, Liu LH, Zhao FJ, Huang XY (2022) Local and Systemic Response to Heterogeneous Sulfate Resupply after Sulfur Deficiency in Rice. Int J Mol Sci 23 (11). doi:10.3390/ijms23116203

Wawrzynska A, Piotrowska J, Apodiakou A, Bruckner F, Hoefgen R, Sirko A (2022) The SLIM1 transcription factor affects sugar signaling during sulfur deficiency in Arabidopsis. J Exp Bot 73 (22):7362-7379. doi:10.1093/jxb/erac371

Wawrzynska A, Sirko A (2024) Sulfate Availability and Hormonal Signaling in the Coordination of Plant Growth and Development. Int J Mol Sci 25 (7). doi:10.3390/ijms25073978

Wilkins O, Nahal H, Foong J, Provart NJ, Campbell MM (2009) Expansion and diversification of the Populus R2R3-MYB family of transcription factors. Plant Physiol 149 (2):981-993. doi:10.1104/pp.108.132795

Wilson MC, Mutka AM, Hummel AW, Berry J, Chauhan RD, Vijayaraghavan A, Taylor NJ, Voytas DF, Chitwood DH, Bart RS (2017) Gene expression atlas for the food security crop cassava. New Phytol 213 (4):1632-1641. doi:10.1111/nph.14443

Winter D, Vinegar B, Nahal H, Ammar R, Wilson GV, Provart NJ (2007) An "Electronic Fluorescent Pictograph" browser for exploring and analyzing large-scale biological data sets. PLoS One 2 (8):e718. doi:10.1371/journal.pone.0000718

Woolfson KN, Esfandiari M, Bernards MA (2022) Suberin Biosynthesis, Assembly, and Regulation. Plants (Basel) 11 (4). doi:10.3390/plants11040555

Wybouw B, Arents HE, Yang B, Nolf J, Smet W, Vandorpe M, Minne M, Luo X, De Clercq I, Van Damme D, Glanc M, De Rybel B (2023) The transcription factor AtMYB12 is part of a feedback loop regulating cell division orientation in the root meristem vasculature. J Exp Bot 74 (6):1940-1956. doi:10.1093/jxb/erad020

Xu H, Liu P, Wang C, Wu S, Dong C, Lin Q, Sun W, Huang B, Xu M, Tauqeer A, Wu S (2022) Transcriptional networks regulating suberin and lignin in endodermis link development and ABA response. Plant Physiol 190 (2):1165-1181. doi:10.1093/plphys/kiac298

Xu X, Guerriero G, Domergue F, Beine-Golovchuk O, Cocco E, Berni R, Sergeant K, Hausman JF, Legay S (2023) Characterization of MdMYB68, a suberin master regulator in russeted apples. Front Plant Sci 14:1143961. doi:10.3389/fpls.2023.1143961

Yoshimoto N, Inoue E, Saito K, Yamaya T, Takahashi H (2003) Phloem-localizing sulfate transporter, Sultr1;3, mediates re-distribution of sulfur from source to sink organs in Arabidopsis. Plant Physiol 131 (4):1511-1517. doi:10.1104/pp.014712

Zenda T, Liu S, Dong A, Duan H (2021) Revisiting Sulphur—The Once Neglected Nutrient: It's Roles in Plant Growth, Metabolism, Stress Tolerance and Crop Production. Agriculture 11 (7). doi:https://doi.org/10.3390/agriculture11070626

Zhang Y, Yue S, Gao Y, Zhao P, Liu M, Qiao Y, Xu S, Gu R, Zhang X, Zhou Y (2024) Insights into response of seagrass (Zostera marina) to sulfide exposure at morphological, physiochemical and molecular levels in context of coastal eutrophication and warming. Plant Cell Environ. doi:10.1111/pce.15048

Zhao Q, Wu Y, Gao L, Ma J, Li CY, Xiang CB (2014) Sulfur nutrient availability regulates root elongation by affecting root indole-3-acetic acid levels and the stem cell niche. J Integr Plant Biol 56 (12):1151-1163. doi:10.1111/jipb.12217

Funding and acknowledgements

XC was funded by China Scholarship Council studentship 201606690032. CC and HW were funded by UK Biotechnology and Biological Sciences Research Council (BBSRC) doctoral training grant BB/M01116X/1. NSG was funded by BBSRC SARIC grant, BB/N004302/1.

We thank Jasmine Carlson, Maxwell Ware, Ahmed Hussain, Lisa King, Elizabeth Chapman, Jessica Finch and Adam Elgey for related preliminary work on aspects of *At*MYB93 function and abiotic stress that informed the direction of this paper.

Competing interests

The authors declare no competing interests.

Author Contributions

XC and JCC designed research. All authors performed research and analysed data.

XC, HW, AO, RE and JCC visualised data. JCC supervised the project. XC and JCC

wrote the manuscript. All authors reviewed the manuscript.

Data availability

The datasets generated and analysed during this study are all included in the

publication and supplemental material.

Figures.

Fig. 1 Differentially expressed genes in the *Atmyb93* mutant are enriched for

functions relating to S-, phenylpropanoid/flavonoid- and carbohydrate

metabolism, and responses to a range of environmental stimuli. S-related

differentially expressed genes in *Atmyb93* largely also show differential

expression in response to S deprivation.

a) and b) Gene Ontology (GO) Biological Process (BP) term enrichment for genes

(a) significantly downregulated (q value < 0.05) in the *Atmyb93* mutant defined by

biological process and (b) significantly upregulated (q value < 0.05) in the *Atmyb93*

mutant. In each panel, GO term enrichment was calculated using PlantRegMap

(p<0.01) and represented with ReviGO, based on the semantic distances between

GO terms. LogSize represents the $\log_{10}$ (number of annotations for GO Term ID in

selected species in the EBI GOA database).

c) Genes with S-related functions significantly differentially expressed in the

Atmyb93 mutant, showing the gene, gene name, function and $\log_2$ fold-change

(down, blue and up, yellow) in *Atmyb93* or under S stress in the root. Data in the

right-hand column highlights genes that also have a fold-change change >1 (down,

blue or up, yellow) under S stress in either endodermis (endo, 3h S deprivation) or a whole root time series (time of largest increase shown) published by (Iyer-Pascuzzi et al. 2011) and visualized at the eFP browser (Winter et al. 2007). Two genes, At1g22150 *SULTR1;3* and At1g61740 sulfite exporter, show different regulation in the endodermis compared to the whole root

Fig. 2 Tomato S24-related *MYB93* homologues are root-enriched and *Sl*MYB93

functions in *Arabidopsis* similarly to *At*MYB93.

a) Root-enriched expression of tomato *MYB93* homologues demonstrated by RT-

PCR. R, root; S, Stem; L, leaf; F, flower; TF, 4 days post anthesis (DPA) tomato fruit;

IMG, immature green; MG, mature green; Br, breaker; RR, red ripen; SD, seed.

SlMYB93, Solyc04g074170; *SlMYB53L*, Solyc04g056310; *SlMYB92*

Solyc05g051550. The *ACTIN2* gene is shown as a control.

b) RT-PCR showing heterologous expression of *SlMYB93* in *Arabidopsis* using a

p35S::SlMYB93 construct (3 independent transgenic lines) compared to a wild type

control (WT, Col-0).

c) Lateral root density of 3 independent transgenic *Arabidopsis* lines overexpressing

SlMYB93 (*Sl*MYB93-OX).

d) Primary root length of 3 independent transgenic *Arabidopsis* lines overexpressing

SlMYB93. 8-day old seedlings from 3 combined biological repeats are shown with

data points (small coloured points) for each repeat coloured differently. Larger

coloured circles represent the means of the biological repeats and black bars

represent the overall mean and standard deviation of the means.

In c) and d) 8-day old seedlings from 3 combined biological repeats are shown with

data points (small coloured points) for each repeat coloured differently. Larger

coloured circles represent the means of the biological repeats and black bars

represent the overall mean and standard deviation of the means. The number of

seedlings per treatment ranges from 35-75; these are the same seedlings in both

panels

Fig. 3 Expression of flowering plant *MYB93* homologues: root-enrichment and

dicot-specific response to S deprivation.

Phylogeny of *AtMYB93*-related genes: the clade in which the *Arabidopsis* S24 genes

fall is coloured in pink, a moncot *AtMYB93*-related clade is purple and a clade

containing the additional suberin-producing *Arabidopsis* MYBs is in black.

Arabidopsis genes are shown in green and tomato genes are shown in blue. Semi-

quantitative expression analysis from publicly available/published data is shown: R,

root; S, shoot, F, fruit, silique or seed; End, endodermis; Exo, exodermis; Sul, S

deprivation response. Pink, tissue expression; yellow, upregulation in response to S

deprivation.

Key to species: At, *Arabidopsis thaliana*; Bn, *Brassica napus;* Br, *Brassica rapa*; Gm,

Glycine max (soybean); Potri, *Populus trichocarpa* (poplar); Manes, *Manihot*

esculenta (cassava); Solyc *Solanum lycopersicum* (tomato); Md, *Malus domestica*

(apple); Os, *Oryza sativa* (rice); Zm, *Zea mays* (maize); Bradi, *Brachypodium*

distachyon; HORVU, *Hordeum vulgare* (barley). According to the notation of Huang

et al (2024), Os2g51799 corresponds to *OsMYB41*, Os08g37970 corresponds to

OsMYB93a and Os06g17780 corresponds to *OsMYB93b*

Fig. 4 *Arabidopsis Atmyb93* mutants and tomato overexpressing *SlMYB93*
show altered shoot element composition.

a) Comparison of shoot element composition in Col-0 and *Atmyb93* 21-day old
plants. Heatmap shows the percentage change of means from 3 combined biological
repeats. Asterisks represent significant differences (p=0.04 for each) from Mann
Whitney tests on the raw data.

b) RT-PCR detection of *SlMYB93* in the roots of wild type tomato plants and two
independent *SlMYB93*-overexpressing lines.

c) Comparison of shoot element composition in wild type and *SlMYB93*-
overexpressing tomato plants. Heatmap shows the percentage change of means
from 3 combined biological repeats. Asterisks represent significant differences (all
p=0.04) from Mann Whitney tests on the raw data

104

Fig. 5 S deprivation affects root development but *Atmyb93* mutants do not show strong root or shoot phenotypes upon S deprivation or OAS treatment.

a), b) Lateral root density (a) and primary root length (b) in wild type (Col-0) and *Atmyb93* seedlings grown with S for 3 days then transferred to –S medium (or +S medium control) for 5 days. 2 combined biological repeats are shown with data points (small coloured points) for each repeat coloured differently. Larger coloured circles represent the means of the biological repeats and black bars represent the overall mean and standard deviation of the means.

c) Formation of adventitious roots at the collet in wild type (Col-0) and *Atmyb93* 8-day old seedlings grown for 3 days with S before transfer to –S medium (or a +S medium control). 5 combined biological repeats are shown with data points (small coloured points) and distributions (shaded areas) for each repeat coloured differently. Larger coloured circles represent the means of the biological repeats and black bars represent the standard deviation of the means.

d), e) Primary root length (d) and lateral root density (e) of wild type (Col-0) and *Atmyb93* seedlings grown for 8 days on 0.1mM O-acetyl serine (OAS). 2 combined biological repeats are shown with data points (small coloured points) for each repeat

coloured differently. Larger coloured circles represent the means of the biological repeats and black bars represent the overall mean and standard deviation of the means.

In all panels, letters indicate differences $p<0.05$ in post-hoc Dunn's test carried out after a Kruskal-Wallis test. For adventitious roots, seedling numbers range from 9 to 20 per treatment in a biological repeat. For primary/lateral root experiments ±S, seedling numbers range from 30 to 98 per treatment depending on the biological repeat. For OAS experiments, seedling numbers range from 18 to 20 per treatment in one biological repeat and 56 to 59 in the other

Fig. 6 Spatial localization of *AtMYB93* promoter activity is not changed by S

stress.

*At*MYB93 promoter activity visualised in wild type plants expressing *pAtMYB93::GUS*

(Gibbs et al 2014) under normal sulphur levels (+S) and S deprivation (-S) in 7-day

old roots. 2 examples of each treatment are shown. Cells expressing

pAtMYB93::GUS are highlighted with arrows. Developing Lateral Root Primordia are

highlighted with arrowheads. Scale bars represent 50µm