Differential activation of p53-Lamin A/C and p16-RB mediated senescence pathways in

trophoblast from pregnancies complicated by type A2 Gestational Diabetes Mellitus

Leena Kadam[1]*, Kaylee Chan[1], Kylia Ahuna[1], Nicole Marshall[1] Leslie Myatt[1]

[1]Department of Obstetrics & Gynecology, Oregon Health & Science University, Portland, OR,
USA
*Corresponding author: kadam@ohsu.edu
7   3303 S Bond Avenue,
8   CHH1, OHSU, Portland, OR 97239
9

10 **FUNDING SOURCES**: This research was funded by the National Institutes of Health grant

11 1R01HD095610

14 **Keywords:** Trophoblast, senescence, placenta, GDM

Abstract:

Gestational diabetes mellitus (GDM) increases maternal risks such as hypertension and future

type 2 diabetes while also contributing to fetal complications such as large-for-gestational-age

infants and stillbirth. The placenta which is crucial for fetal development, exhibits structural and

functional changes in GDM, but the impact of these alterations on placental trophoblast function

remains unclear. During their differentiation villous cytotrophoblast display several characteristics

of senescent cells however the role of senescence pathways in placental function remains

unexplored in GDM. Here we investigate whether placental senescence pathways are altered in

GDM, utilizing term villous tissue and primary trophoblasts to assess molecular changes, and

determined fetal sex-based differences. Our data suggest that both p21 and p16 mediated

senescence pathways are activated during trophoblast differentiation and are dysregulated in

GDM placenta in a sexually dimorphic manner. We also provide evidence for increased activation

of p53-Lamin A/C and p16-RB pathways in trophoblast from GDM placentas. Reduced expression

of p21 and its downstream effects on GCM1 expression and βhCG secretion outline how altered

physiological senescence can affect trophoblast differentiation and function. This is a seminal

study highlighting how placental senescence pathways are altered in pregnancies complicated

by GDM.

Introduction:

Gestational diabetes mellitus (GDM) is a metabolic disorder of pregnancy which manifests as impaired glucose tolerance with onset or first recognition during pregnancy [1]. It is considered as one of the great obstetrical syndromes with an increasing incidence, currently being ~14.7% worldwide and ~7% of pregnancies in the USA [2,3]. Women with GDM are more susceptible to hypertensive disorders, dysfunctional labor and thromboembolic events during pregnancy and are also at higher risk of developing type 2 diabetes and cardiovascular disease in later life [4-7]. GDM can lead to fetal complications such as large-for-gestational-age (LGA) infants (which can lead to cesarean delivery, birth trauma, shoulder dystocia and asphyxia), fetal growth restriction (FGR) or stillbirth [4,5,8]. The offspring of women with GDM are also at a higher risk for development of obesity and diabetes in later life [4,5,8-10]. It has been observed that women carrying male fetuses are at a higher risk of developing GDM and associated co-morbidities highlighting that fetal sex might influence the development and severity of the disease [11,12]. The placenta, being the maternal-fetal interface, ensures optimal fetal nutrition, growth and development. The maternal metabolic conditions of hyperglycemia and hyperinsulinemia in GDM appear to affect placental development as GDM placentas are heavier and show histopathological defects such as immature villi, villous fibrinoid necrosis, increased angiogenesis and chorangiosis [13,14]. Extensive reduction of microvilli on thin hyper-vacuolized syncytiotrophoblast, a thick cytotrophoblast layer and a thick trophoblast basal membrane have also been reported in GDM placentas [15,16]. However, the effect of these changes on placental trophoblast function at the cellular level is not completely understood.

Cellular senescence is an irreversible cycle arrest wherein the cells stop proliferating but continue to be metabolically active. It is a physiological process integral to normal tissue homeostasis and functions with crucial roles in tumor suppression, wound healing and embryonic development [17-19]. Disruption in physiological senescence pathways is associated with declining tissue function

in diseases such as hypertension and Type 2 Diabetes Mellitus in non-pregnant individuals

[20,21]. Furthermore, stressors such as increased oxidative stress, inflammation, metabolic

imbalance and insulin resistance - characteristic of pregnancies with GDM, are also known to

increase senescence in non-pregnant individuals but their impact on placental senescence

remains relatively unexplored in GDM tissues [22-30]. Interestingly, placental syncytiotrophoblast

display several characteristics of senescent cells in being multinucleate, terminally differentiated

cells with irreversible replication arrest. In recent years, elevated senescence has been reported

in placentae from pathologies such as FGR, stillbirth, pre-eclampsia and GDM [31-33]. However,

the role of senescence in placental function has not been evaluated.

In the present study, we aimed to determine if placental senescence pathways are altered in GDM

potentially affecting its function. We utilized villous tissue as well as primary trophoblast isolated

at term from placentas of normoglycemic and GDM pregnancies to evaluate changes in

senescence pathway mediators. As it is known that pregnancies with male fetuses are more

susceptible to developing GDM, we also evaluated fetal sex-based effects on trophoblast

senescence.

METHODS

Patient recruitment and collection of placental tissue:

This study was performed in line with the principles of the Declaration of Helsinki. Approval was

granted by the Institutional Review Board of Oregon Health & Science University (IRB 16328).

Informed consent was obtained from all patients before tissue collection. Placentas (n=10 per

group with n=5 per sex) were collected from normoglycemic women (NW) and women with type

A2 GDM (GDM). GDM was defined using IADPSG criteria which recommends 75-grams two

hours OGTT with at least one abnormal result: Fasting plasma glucose (FPG) at ≥ 92 mg/dl or 1-

hour ≥ 180 mg/dl or 2-hour ≥ 153mg/dl to be classified as GDM [34]. Women with type A2 GDM

were those with greater than 30% of FPG ≥90-95 mg/dl or 1 hr Postprandial ≥130-140 mg/dl and

needed medication to control their blood glucose levels. To keep the samples consistent, we

focused on women with type A2GDM and collected samples from women who received insulin

as medication to control their diabetes. Patients from both NW and GDM groups satisfied the

following inclusion/exclusion criteria.

Inclusion criteria included a singleton pregnancy, an age range of 18-45 and delivery by cesarean

section at term with no labor. Exclusion criteria included concurrent disease (including, but not

limited to IUGR, hypertension, pre-eclampsia, eating disorders, infection, inflammatory disorders),

use of tobacco, drugs, or medications other than to treat GDM, excessive weight gain or loss prior

to pregnancy (>20 lbs), or bariatric surgery in the last year and labor defined by regular uterine

contractions (every 3-4 minutes, verified by tocodynamometry) resulting in cervical dilatation

and/or effacement.

For all patients, maternal age, gestational age, fetal and placental weights were recorded, and

placental tissue was processed within 30 minutes of delivery. Placental villous tissue was

immediately sampled from 5 random sites of the placenta and processed for trophoblast cell

isolation as outlined below. Small villous pieces (~5mm^3) from all 5 sites were also flash frozen

and stored at -80^{0}C. Subsequently these villous tissue pieces from all 5 sites were powdered

under liquid nitrogen and equal amounts from all sites were combined to be used for downstream

analysis.

Primary cell isolation and culture:

The chorionic plate and decidua were removed from each of the randomly collected placental

pieces leaving only villous tissue, which was thoroughly rinsed in PBS to remove excess blood.

Primary cytotrophoblast were isolated from villous tissue using a protocol adapted from Eis *et al.*

[35] using trypsin/DNAse digestion followed by density gradient purification. Isolated

cytotrophoblast cells were then frozen in freezing media (10% DMSO in FBS) and stored in liquid

nitrogen until usage. For culture, cytotrophoblast cells were rapidly thawed in a 37°C water bath

and immediately diluted in Iscove's modified Dulbecco's medium (25mM glucose, 4mM glutamine

and 1mM pyruvate) supplemented with 10% FBS and 1% penicillin/streptomycin (complete

media). Cells were centrifuged at 1000g for 10 min and re-suspended in fresh complete media.

Trophoblast cells were then plated in either 6 well-plates for collecting proteins and RNA or in 8

well chamber slides for staining. The cytotrophoblast (CT) when cultured differentiate into

syncytiotrophoblast (ST) over a period of 72 hours. This differentiation is confirmed visually by

observing cells under a light microscope. All studies were performed at two time points - 24 hrs

denoted as CT stage and 72hrs – denoted as ST stage. Media was collected at both time points

to measure secretion of βhCG as an additional way to confirm cytotrophoblast differentiation.

Protein extraction and Western blotting:

<u>Extraction from tissue:</u> Approximately 25mg of powdered tissue was mixed with 250μL of RIPA

buffer supplemented with protease/phosphatase inhibitors (ThermoFisher Scientific, Cat.

#A32959) and incubated on ice for 30 minutes. Post incubation, the tubes were centrifuged at

1000 rpm for 5 minutes and supernatant/lysate was transferred to a new tube. Total protein was

quantified using the Pierce BCA Protein Assay Kit as per manufacturers protocol. (ThermoFisher

Scientific, Cat. #23225).

<u>Extraction from cells</u>: Proteins were extracted from cultured trophoblast cells at 24 and 72hrs.

Briefly, cells were washed in cold PBS with protease and phosphatase inhibitors (Halt^{TM} Protease

Inhibitor Cocktail) followed by lysing in 150μL RIPA buffer supplemented with

protease/phosphatase inhibitors (ThermoFisher Scientific, Cat. #A32959) for 20 minutes on ice.

The lysate is then transferred to a 1.5ml tube and centrifuged at 1000g for 10 min at 4°C to remove

cell debris. Total protein was quantified using the Pierce BCA Protein Assay Kit.

Approximately 30μg of protein was separated on 12% sodium dodecyl sulphate-polyacrylamide

gel electrophoresis (SDS-PAGE) hand-cast gels for approximately 30 min at 30V followed by 2

hr at 100V and transferred for 1 hr at 100V onto nitrocellulose membranes using Mini-PROTEAN

tetra cell electrophoresis chamber (BioRad, Cat. # 1658004). Membranes were blocked in 5%

(w/v) nonfat milk in TBS + 0.1% Tween 20 (TBST) for 1 hr and incubated with primary antibodies

(outlined in supplementary table 1) overnight at 4 °C. On the next day, membranes were washed

three times in TBST for 5 min each and incubated with HRP-conjugated secondary antibodies.

Membranes were washed and the antibody binding was detected using Supersignal West Pico

Plus ECL Substrate (ThermoFisher Scientific, Cat. #34578) for 1 min and imaged using

ChemiDoc Imaging System (Bio-Rad Laboratories, USA) and Image Lab V.5.1 software (Bio-Rad

Laboratories, USA). Densities of immunoreactive bands were measured as arbitrary units by

ImageJ software. Protein levels were normalized to a housekeeping protein β-actin (1: 20,000;

Abcam, UK).

Senescence associated β- galactosidase staining assay:

Senescence-Associated β-Galactosidase (SA-βgal) is a lysosomal enzyme that cleaves β-D-

galactose residues in β-D-galactosides. Detectable SA-βgal activity is the most extensively used

marker for senescent or aging cells whether in culture or in mammalian tissues [36,37]. Senescent

cells in trophoblasts isolated from LN and GDM placentae, CT were cultured for 24hrs in 8

chambered slides and then processed for SA-βgal staining using the CellEvent™ Senescence

Green Detection Kit (Thermo Fisher) as per manufacturers protocol. Briefly, the cells were first

fixed with 200uL of 1X fixative solution for 10 mins at RT after which the cells were washed with

PBS containing 1% Bovine serum albumin. 200uL of prewarmed working solution was then

applied to the cells. The slide chambers were sealed and incubated in dark for 2 hours in a CO_2

free incubator at 37^0C. After incubation, the working solution was drained, and cells were washed

3 times with 200 µL of PBS followed by staining with DAPI nuclear stain for 10 minutes. The cells

were washed again once with PBS, the chamber walls separated, and the slides were

coverslipped using SlowFade Diamond Antifade Mountant (Thermofisher Scientific, Cat.

#S36972). After allowing to set, coverslips were sealed in place using clear nail polish. Images

were captured using a Zeiss LSM 880 confocal microscope and processed using ImageJ

Software. From each chamber, 3 randomly chosen areas were imaged. For each area, the total

number of cells and cells positive for SA-βgal staining were counted to determine the % of

senescent cells per sample.

Enzyme linked Immunosorbent (ELISA) assay:

The media collected from cultured trophoblast at 24 and 72hrs was assayed for levels of human

chorionic gonadotropin β (βhCG) hormone using the Duo Set ELISA development kits (R&D

Systems, USA) as per the manufacturers protocol. The optical density of the final-colored reaction

product was measured at 450 nm using multispectral UV/VIS plate reader (Bio-Tek, VT). Standard

curves were used to calculate protein content in the samples. The level of proteins detected was

normalized to cellular protein.

Statistical analysis: The data was analyzed for statistical difference using the GraphPad Prism

7.0 software. All analysis consisted of n=5 samples per sex per clinical group. Outliers were

detected using the ROUT method and wherever detected maximum 1 outlier was excluded from

analysis. The values between CT and respective ST were analyzed using paired t-test whereas

differences between groups were analyzed using ANOVA followed by post hoc Tukey analysis.

RESULTS:

I. Clinical characteristics:

The clinical characteristics of patients for which placental tissue and isolated trophoblast were

utilized in the study are outlined in Table 1.0. There was no difference in maternal age and

gestational age between the normoglycemic (NW) and type A2 GDM (GDM) group. However,

patients from the GDM group had significantly higher BMI. As obesity increases the incidence of

GDM by 3 times, we expected to see this in our GDM cohort [8,38,39]. All neonates had good

outcomes and there was no difference in placental or fetal weight at birth between the NW and

GDM groups.

II. GDM placentas have elevated levels of senescence marker p16:

Placental villous tissue from NW and GDM women was used to determine the expression of

senescence markers p21$^{WAF1/CIP1}$ (p21) and p16^{INK4A}(p16). We observed that expression of p21

was similar between the NW and GDM placentas whereas expression of p16 was significantly

higher in GDM than NW (Fig 1A).

	Normoglycemic (NW, n=10)		Gestational Diabetes (GDM, n=10)	
	Male (n=5)	Female (n=5)	Male (n=5)	Female (n=5)
Maternal age (yrs)	33.8 $\pm$ 3.5		34.7 $\pm$ 4.5	
	35 $\pm$ 2.2	32.6 $\pm$ 4.5	36 $\pm$ 3.7	33.4 $\pm$ 5.3
Maternal BMI (wt[kg]/height[m^2])	21.29 $\pm$ 1.8		34.77 $\pm$ 4.8*	
	22.5 $\pm$ 1.4	20.0 $\pm$ 1.2	35.2 $\pm$ 3.0*, †	34.2 $\pm$ 6.5*, †
Gestational age (wks)	38.62 $\pm$ 0.90		38.08 $\pm$ 1.15	
	38.84 $\pm$ 0.93	38.4 $\pm$ 0.92	38.32 $\pm$ 0.93	37.84 $\pm$ 1.36
Fetal birthweight (g)	3459 $\pm$ 319		3212 $\pm$ 258	
	3556 $\pm$ 214	3361 $\pm$ 399	3326 $\pm$ 205	3098 $\pm$ 275
Placental weight (g)	556.4 $\pm$ 107.7		638.0 $\pm$ 150.5	
	609.7 $\pm$ 79.1	503.1 $\pm$ 112.9	690.8 $\pm$ 174.3	572.0 $\pm$ 97.3
Fetal:Placental weight ratio	6.3 $\pm$ 1.1		5.5 $\pm$ 1.3	
	5.8 $\pm$ 0.6	6.8 $\pm$ 1.3	5.5 $\pm$ 1.6	5.6 $\pm$ 1.0

Table 1: Clinical characteristics of study participants. Data presented as mean ± SD. Significant differences between NW and GDM groups were determined using the student's *t* test. * *p* < 0.05 for NW vs. GDM, † p<0.05 for same sex comparison between the NW and GDM groups.

The data was then stratified based on fetal sex creating 4 subgroups: NW-male, NW-female,

GDM-male and GDM-female. This analysis showed that expression of p21 was higher in the

GDM-female compared to the other 3 subgroups, whereas the expression of p16 in the GDM-

male subgroup was significantly higher compared to the other 3 subgroups. (Fig 1B). As outlined

earlier, the women from the GDM group had significantly higher BMI compared to NW. We

therefore also assessed expression of p21 and p16 in placentas from obese women with no type

A2GDM to delineate if the changes seen in GDM placentas were due to maternal obesity.

Interestingly obese placentas showed a pattern similar to the NW group (Supplementary figure 1)

implying the changes in GDM placentas were specific to maternal diabetes. For the remaining

experiments, we therefore continued using the NW group as our control.

III. Trophoblast isolated from GDM placentae have higher numbers of senescent cells:

As trophoblast are the major cell type that display characteristics of senescent cells in the

placenta, we focused on them [40]. Cytotrophoblast isolated from NW and GDM placentas were

Figure 1: Protein expression of senescence markers p21 and p16 in villous tissue. A. Expression in NW and GDM villous tissue **B.** Expression in NW and GDM tissues stratified by fetal sex. Data presented Mean ± SEM with N=10 per clinical group with each containing n=5 males and n=5 females. * p<0.05, **p<0.01, *** p<0.001.

cultured for 24 hours and the number of SA-β-gal positive cells was assessed at CT stage

(Supplementary figure 2 shows representative images). We observed that CT from GDM

placentas had a higher percentage of SA-β-gal positive cells compared to NW (Fig 2A). Stratifying

this data by fetal sex showed that NW-male and GDM-male subgroups had comparable number

of senescent cells, but the GDM-female subgroup had a higher number of senescent cells

compared to the other 3 subgroups (Fig 2B).

Figure 2: Percentage of senescent (SA-β-gal positive) cytotrophoblast after 24 hours of culture. A. Percentage of SA-β-Gal positive cells in CT isolated and cultured from NW and GDM placenta **B.** Percentage of SA-β-Gal positive cells stratified by fetal sex. N=6 per clinical group with each containing n=3 males and n=3 females. * p<0.05, **p<0.01 between groups

IV. Differentiating cytotrophoblast from GDM placentas have distinct expression of senescence markers:

We assessed the expression of p16 and p21 in cultured trophoblast at CT stage and post differentiation at ST stage. We observed that the levels of both p21 and p16 were lower in ST when compared to their respective CT in both NW and GDM groups - a trend previously identified in cells undergoing senescence [41] (Fig 3A, C). Comparison between the groups showed that CT from GDM group had significantly lower expression of p21 and higher expression of p16 (Fig 3A, C) compared to CT from NW group implying that trophoblast from both groups were undergoing senescence during trophoblast differentiation, but via activation of different pathways.

Figure 3: Protein expression of senescence markers p21 and p16 in CT and differentiated ST. Expression of p21 in NW and GDM CT and ST in **A**. sex combined manner, **B**. fetal sex stratified manner. Expression of p16 in NW and GDM CT and ST in **C**. sex combined manner, **D**. fetal sex stratified manner. Data presented Mean ± SEM with N=10 per clinical group with each containing n=5 males and n=5 females. * $p<0.05$, **$p<0.01$, *** $p<0.001$ between groups and # $p<0.05$ for comparison between CT vs ST in the same group.

Fetal sex-stratified analysis showed that the expression of p21 significantly declined in ST

compared to their respective CT in all 4 subgroups (NW-male, NW-female and GDM-male, GDM-

female). CT from the NW-male subgroup had higher p21 levels compared to NW-female as well

as GDM-male and GDM-female subgroups (Fig 3B). Expression of p16 followed a similar trend

wherein its levels also significantly declined in ST compared to their respective CT in all

subgroups. NW-female CT also had significantly lower expression of p16 compared to CT from

NW-male and GDM-male subgroups (Fig 3D).

V. The p53-p21 and p16-pRB pathways are differentially activated in GDM trophoblast:

To understand the mechanism behind differential senescence pathway activation, we assessed

the expression of proteins, Tumor protein 53 (p53) and Retinoblastoma (RB) which act upstream

and downstream of p16 and p21 respectively [42,43]. Phosphorylation of p53 is needed for its

activation, whereas RB needs to be hypo-phosphorylated for mediating cell cycle arrest. We

therefore compared the ratio of phosphorylated forms to total protein levels to determine their

levels of activation.

We observed a higher proportion of activated p53 in GDM compared to NW at both the CT and

ST stage (Fig 4A) implying higher activity of p53 in GDM trophoblast. Levels of phosphorylated 3

(normalized to Actin) were higher in ST from GDM compared to ST from NW group respectively

(Supplementary Fig 3A). Similarly, total protein levels of p53 (normalized to Actin) were also

significantly higher in ST from GDM compared to NW, whereas their levels in CT from both groups

remained comparable between (Supplementary Fig 3C). Fetal sex-stratified analysis showed that

the proportion of activated p53 declined in ST compared to their respective CT in NW-male and

GDM-male subgroups but the proportions in NW-female and GDM-female subgroups were

comparable (Fig 4B). Levels of phosphorylated p53 (normalized to Actin) were higher in ST from

GDM-female compared to ST from the other 3 subgroups (Supplementary Fig 3B). Total protein

levels of p53 were higher in ST from GDM-male and female groups compared to their respective

CT, but this was not observed in NW-male or NW-female subgroups (Supplementary Fig 3D).

The proportion of phosphorylated RB increased in ST from both groups compared to their

respective CT (Fig 4C) however, GDM ST had significantly lower levels compared to NW ST,

suggesting hypo-phosphorylation of RB in GDM trophoblast. The levels of phosphorylated RB

(normalized to Actin) were lower in CT of GDM group compared to NW, but the levels in ST were

comparable (Supplementary Fig 4A). The levels of total RB protein (normalized to Actin) showed

a similar trend, with ST from both groups having lower levels compared to their respective CT

(Supplementary Fig 4C). Fetal sex-stratified analysis showed that both male and female ST from

NW group have a significantly higher proportion of phosphorylated RB compared to their

Figure 4: Protein expression of senescence pathway mediators p53 and RB in CT and differentiated ST. Ratio of expression of phosphorylated p53/total p53 in NW and GDM CT and ST in **A**. sex combined manner, **B.** fetal sex stratified manner. Ratio of expression of phosphorylated RB/total RB in NW and GDM CT and ST in **C**. sex combined manner, **D.** fetal sex stratified manner. Data presented Mean ± SEM with N=10 per clinical group with each containing n=5 males and n=5 females. * p<0.05, **p<0.01, *** p<0.001 between groups and # p<0.05 for comparison between CT vs ST in the same group.

respective CT and ST from NW-male subgroup had significantly higher proportion compared to

ST in the other 3 subgroups (Fig 4D). These results suggest that during trophoblast differentiation,

NW increased phosphorylation of Rb potentially to reduce its activity which was not attained in

GDM. Levels of phosphorylated RB (normalized to Actin) were comparable in CT and ST in the

subgroups except in NW-female wherein ST had significantly lower levels compared to their CT

(Supplementary Fig 4B). For total RB protein (normalized to Actin), CT and ST from GDM-male

subgroup had comparable levels whereas the remaining 3 subgroups had significantly lower

levels in ST compared to their respective CT (Sup Fig 4D).

VI. Expression of Lamin A/C is high in GDM trophoblast

Lamins are a type of intermediate filament protein that line the inner surface of the nuclear

envelope and contribute to the size, shape, and stability of the nucleus which is critical in

multinucleated cells like the syncytiotrophoblast [44]. However, elevated levels of Lamin A/C were

shown to mediate p53 induced senescence via destabilizing nuclear structure [45,46]. We

observed significantly higher expression of Lamin A/C in both CT and ST from GDM placentas

compared to CT and ST from NW placentas respectively (Fig 5A)). Fetal sex stratified analysis

showed that CT and ST in both NW-male and NW-female subgroups had comparable expression.

Expression in CT from GDM-female subgroup was significantly higher compared to CT from NW-

Figure 5: Protein expression of Lamin A/C in CT and differentiated ST. Expression of Lamin A/C in CT and ST **A**. sex combined manner, **B.** fetal sex stratified manner. Data presented Mean ± SEM with N=10 per clinical group with each containing n=5 males and n=5 females. * p<0.05, **p<0.01, *** p<0.001 between groups.

male and NW-female subgroup whereas ST from GDM-male subgroup had significantly higher

expression compared to ST from the NW-male and NW-female subgroup (Fig 5B).

VII. Expression of both GCM1 and βhCG is reduced in differentiating trophoblast from GDM

placentas

A recent report suggested that p21 is involved in regulation of Glial cell missing -1 (GCM1)

implying a physiological link between activation of senescence and trophoblast differentiation [47].

We observed that levels of GCM1 were significantly lower in ST from both groups compared to

their respective CT suggesting induction of trophoblast differentiation in both groups. However,

both CT and ST from GDM placentas had significantly lower levels of GCM1 compared to CT and

Figure 6: Protein expression of GCM1 and βhCG secretion in CT and differentiated ST. Expression of GCM1 CT and ST from NW and GDM **A**. sex combined manner, **B.** fetal sex stratified manner. Levels of βhCG secreted in media by CT and ST from NW and GDM **C**. sex combined manner, D**.** fetal sex stratified manner Data presented Mean ± SEM with N=10 per clinical group with each containing n=5 males and n=5 females. * p<0.05, **p<0.01, *** p<0.001 between groups and # p<0.05 for comparison between CT vs ST in the same group.

ST from NW placentas respectively (Fig 6A) suggesting reduced induction of differentiation in

these cells. Sex stratified analysis reflected the pattern of downregulation in GCM1 expression in

ST compared to CT in all subgroups however CT from the GDM-male subgroup had significantly

lower expression of GCM1 compared to the other 3 subgroups (Fig 6B). The expression in CT

and ST from NW-male and female groups was comparable, implying induction of GCM1 in

maternal GDM was sexually dimorphic.

As GCM1 is known to regulate human chorionic gonadotropin-β (βhCG) – also involved in

trophoblast differentiation, we next evaluated βhCG secretion from these cells. We observed that

secretion of βhCG significantly increased as CT differentiated to ST in trophoblast from NW and

GDM placentas, in line with previous literature and again implying induction of differentiation in

CT from both groups (Fig 6C). However, the βhCG levels secreted GDM ST by them were

significantly lower than ST from NW placentas implying reduced trophoblast differentiation. Fetal

sex stratified analysis showed that βhCG secretion significantly increased in ST compared to their

respective CT in all four subgroups. Interestingly, ST from the NW-female subgroup had

significantly higher levels compared to ST from NW-male but the levels secreted by ST from GDM-

males and GDM-female subgroups were comparable suggesting a sexual dimorphic pattern in

normoglycemic placentas which was absent in GDM. The **β**hCG levels secreted by ST from the

NW-female subgroup were also significantly higher than GDM-male and female subgroups (Fig

6D).

DISCUSSION:

The current literature exploring senescence in human placenta focuses mainly on expression of

senescence associated markers with several studies reporting upregulation in post mature

placentas, and in placentas from pregnancies complicated by PE, FGR and stillbirth [31,32,48].

However, the functional role of senescence in placental tissue remains unexplored. Here, using

the primary trophoblast cell model we demonstrate that while senescence pathways are induced

during physiological trophoblast differentiation, their induction is reduced in the maternal condition

of GDM and affects differentiation which in turn might affect placental function. We also assessed

sexual dimorphism in these pathways which has not been previously addressed.

Cellular senescence is often synonymously used with ageing as accumulation of senescent cells

in tissues has been observed in aging and age-related pathologies [49]. However, senescence

also plays crucial physiological roles in maintaining tissue homeostasis, wound healing and as an

anti-cancer mechanism in tissues [50]. Cellular senescence has also emerged as a response to

different stressors including genotoxic stress, nutrient imbalance and mitochondrial dysfunction.

The latter two being extensively reported in GDM placentas [17,51]. Programmed cellular

senescence was shown to be crucial for embryonic development and was also proposed to play

a role in timing of parturition [31]. Studies exploring aging and/or senescence in placental tissues

have reported an increase in p21, p16 and p53 markers in pregnancies complicated by PE, IUGR,

stillbirth and miscarriage [31,32,48,52]. Increase in senescence markers was also reported to

coincide with increasing oxidative stress in these pathological states as well as post mature

placentas [32]. Most previous studies were conducted using villous tissues from these

pregnancies or trophoblast cell lines of cancerous origins that might have already bypassed cell

cycle-related processes which may be dysregulated in senescence *in-vivo* [31,32,48,40,53].

Biron-Shental *et al* reported an increase in the senescence markers – SA-β-gal staining,

senescence-associated heterochromatin foci and reduced telomere lengths in GDM placentas

[33]. Garcia-Martin *et al* also reported reduced telomere lengths in GDM placentas, which was

reversed when mothers received interventions such as metformin or insulin but focused only on

placentas from male fetuses [54]. While these studies support our findings, they lack the ability to

distinguish between physiological and pathological senescence as they rely on histological

markers or protein levels in bulk placental tissue and do not consider sexual dimorphism. Higuchi

et al performed a longitudinal study across placental tissue from 1st, 2nd and 3rd trimester and

reported higher senescence in CT from early to mid-pregnancy in association with syncytial

fusion, while senescence was observed in ST in late pregnancy. The authors suggested this

change in senescence across gestation may play a role in maintaining placental function [53],

however the physiological role of senescence in placental function remains unevaluated in GDM.

In our analysis with villous tissue, we also observed altered expression of senescence regulators

p21 and p16 in GDM placentas. P21 and p16 are cyclin-dependent kinase (CDK) inhibitors that

prevent cell cycle progression by interacting respectively with the transcription factors p53 and

RB that regulate expression of genes involved in nutrient metabolism, DNA replication, cell growth

and apoptosis (Fig. 7) [55]. We observed an increase in expression of p16 in GDM placentas

compared to NW controls implying an increase in p16 mediated senescence in GDM. We also

observed sexual dimorphism in expression, wherein male placentas from the GDM group had

significantly higher levels of p16 compared to female placentas. These results suggest that male

and female placentas differ in activation of senescence pathways in response to maternal GDM

conditions. As trophoblast display characteristics of senescent cells, we focused on them and

utilized cytotrophoblast isolated from pregnancies complicated by NW and GDM to study

expression of senescence pathway mediators as they differentiated to syncytiotrophoblast [40].

This also enabled us to understand differences between physiological senescence in NW and

how it is altered in GDM pathology. At 24 hours post culture, we observed SA-β-gal positive cells

in both NW and GDM CT with GDM CT having a significantly higher number suggesting premature

activation of senescence pathways. We also observed that CT from both groups activated both

p16 and p21 mediated senescence as part of their differentiation process and their levels declined

as CT differentiated to ST (Fig. 1), which is synonymous with observations made in non-placental

senescent cells wherein levels of p21 and p16 decline once the senescence program is initiated

[41]. P21 is proposed to be activated as part of the developmental senescence program as p21

mutant mice showed defects in embryonic senescence among other developmental defects

which would explain the robust expression of p21 in CT from both sexes in NW and GDM

placentas [19,56]. Taken together, these results further solidify the role for physiological

senescence in trophoblast differentiation. Studies from other models have also suggested that

p21 is activated in the initial stages of senescence whereas p16 is activated in the late stages

and an increase in expression of p16 has been linked to a senescence-mediated decline in

progenitor stem cell populations and function [57,58]. We observed lower p21 expression and

higher p16 levels in GDM CT compared to CT from NW placentas. P16 also showed a sexually

dimorphic pattern wherein female CT from both NW and GDM groups had lower expression

compared to their respective male CT. Reduced levels of p21 and higher levels of p16 in CT from

GDM trophoblast with higher number of SA-β-gal positive cells could indicate disruption in

physiological senescence in these placentas.

We also observed an increase in activated p53 in which typically acts upstream of p21 in both

GDM CT and ST. Cellular stressors including metabolic stress, oxidative stress and mitochondrial

stress can all activate p53 as a way to combat that stress and can lead to cell death [59,60]. The

non-significant increase in p53 observed here could be in response to cellular stress in GDM.

However, increased activation of p53 did not coincide with a concomitant increase in p21 as

expected, which might be because p21 can be regulated in a p53 independent manner [61,62].

However, p53 was recently shown to induce p16 expression via Lamin A/C stabilization [45].

Indeed, we observed increase in Lamin A/C levels in both CT and their differentiated ST from

GDM placentas suggest that even though there was a nominal activation of p53, this was enough

to induce p16. While there were no specific sexually dimorphic trends in expression of p53 and

Lamin A/C, it is worth highlighting that both CT and ST from GDM females displayed higher

expression of both proteins. We also assessed phosphorylated and total levels of RB protein

which acts downstream of p16. P16 enforces senescence by inhibiting CDK4/6, leading to RB

hypo-phosphorylation which results in its binding to E2F's forming a repressive complex that binds

and prevents expression of genes needed for cell cycle progression [63,64] (Fig 7). In our analysis

we observed that CT from both NW and GDM groups had comparable proportion of

phosphorylated/total RB which increased in their respective ST. This increase in

phosphorylated/total RB could be explained by reduced levels of total RB protein implying rapid

Figure 7: Overview of changes in senescence pathways as cytotrophoblast from NW and GDM placentas differentiate into syncytiotrophoblast.

degradation of RB once the cell cycle arrest pathways have been activated. Interestingly, we

observed that ST from the GDM group were more hypo-phosphorylated compared to ST from the

NW group which aligns with increased cell cycle arrest and senescence in GDM ST [43].

Additionally, p21 is also known to regulate RB phosphorylation and formation of the repressive

RB-E2F complexes. P21 mediated inhibition of CDK's also contributes to reduced

phosphorylation of RB and progress to cell cycle arrest [65]. The decline in p21 levels as CT from

NW placentas differentiate to ST coincides with an increase in RB phosphorylation status in these

cells highlighting the role of the p21-RB axis in physiological senescence. However, low levels of

p21 in GDM placental CT with a further decline in ST suggest that this axis is also dysregulated

in GDM trophoblast. To determine if this dysregulation in senescence affects placental function,

we analyzed levels of trophoblast differentiation factors GCM and βhCG. Expression of GCM1

was significantly downregulated in both CT and ST from the GDM group compared to the CT and

ST from NW group respectively, suggesting reduced differentiation. As CT from both groups

differentiated to ST, they increased their secretion of βhCG confirming differentiation however,

the βhCG levels secreted by GDM ST were significantly lower compared to ST from the NW group

again confirming dysregulated differentiation in GDM trophoblast. GCM1 may induce p21

expression to arrest the cell cycle, and in turn p21 enhances the function of GCM1 to begin the

fusion program [47]. The reduced expression of both p21 and GCM1, and the downstream target

βhCG suggest that the dysregulation in p21 mediated physiological senescence in GDM

placentas affects trophoblast differentiation. This dysregulated differentiation could explain the

presence of immature villi observed in GDM placentas [13]. Taken together, our results suggest

that GDM cytotrophoblast may be undergoing premature senescence, driven by p53 mediated

Lamin A/C accumulation and p16/RB activation (Fig 7). Our results also suggest that loss in p21

in GDM trophoblast alters physiological senescence and is capable of disrupting trophoblast

differentiation by altering GCM1 and βhCG levels. Disruption in trophoblast differentiation can

alter placental function in numerous ways such as altered secretion of hormones and signaling

mediators, altered expression of nutrient transporters, etc. This is the first study to outline

disruption of placental senescence pathways in GDM and their potential impact on trophoblast

differentiation. Several stressors including oxidative stress, mitochondrial dysfunction, and

metabolic stress have all been shown to induce senescence pathways in non-pregnant individuals

tissues [66,67,27,22-26]. GDM placentas must adapt to the altered maternal metabolic

environment and have been reported to have increased oxidative stress as well as mitochondrial

dysfunction [68-70]. It is plausible that a combination of these conditions could alter physiological

senescence in GDM placentas (mediated via p21) and induce premature elevated senescence

(mediated by p16). We also report sexual dimorphism in placenta senescence pathways further

highlighting how fetal sex plays a role in shaping placental function and its response to maternal

stress. Our study warrants further research into the potential stimulus for the reported

dysregulated placental senescence in GDM and the sexual dimorphism therein.

AUTHOR CONTRIBUTIONS: LK conceptualized, designed and performed the experiments. LK

was also responsible for data generation and analysis. NM was involved with identifying patients

and KC and KA were involved with sample collection and running experiments. LK wrote the

manuscript with contribution from LM.

CONFLICT OF INTEREST: The authors declare no conflict of interests.

REFERENCES:

1. AmericanDiabetesAssociation (2018). 2. Classification and Diagnosis of Diabetes: Standards
of Medical Care in Diabetes-2018. *Diabetes Care, 41*(Suppl 1), S13-S27,
doi:10.2337/dc18-S002.
2. Saeedi, M., Cao, Y., Fadl, H., Gustafson, H., & Simmons, D. (2021). Increasing prevalence of
gestational diabetes mellitus when implementing the IADPSG criteria: A systematic review
and meta-analysis. *Diabetes Res Clin Pract, 172*, 108642,
doi:10.1016/j.diabres.2020.108642.
3. Hur, K. Y. (2019). New Diagnostic Criteria for Gestational Diabetes Mellitus and Pregnancy
Outcomes in Korea. *Diabetes Metab J, 43*(6), 763-765, doi:10.4093/dmj.2019.0247.
4. Yogev, Y., & Catalano, P. M. (2009). Pregnancy and obesity. *Obstet Gynecol Clin North Am,
36*(2), 285-300, viii, doi:10.1016/j.ogc.2009.03.003.
5. Reece, E. A. (2008). Obesity, diabetes, and links to congenital defects: a review of the evidence
and recommendations for intervention. *J Matern Fetal Neonatal Med, 21*(3), 173-180,
doi:10.1080/14767050801929885.
6. Vasudevan, C. R., M. McGuire, W. (2011). Fetal and perinatal consequences of maternal
obesity. *Arch Dis Child Fetal Neonatal Ed, 96*(5), F378-382, doi:adc.2009.170928 [pii]
10.1136/adc.2009.170928.
7. England, L. J., Dietz, P. M., Njoroge, T., Callaghan, W. M., Bruce, C., Buus, R. M., et al. (2009).
Preventing type 2 diabetes: public health implications for women with a history of
gestational diabetes mellitus. *Am J Obstet Gynecol, 200*(4), 365 e361-368,
doi:10.1016/j.ajog.2008.06.031.
8. Chu, S. Y., Kim, S. Y., Lau, J., Schmid, C. H., Dietz, P. M., Callaghan, W. M., et al. (2007).
Maternal obesity and risk of stillbirth: a metaanalysis. *Am J Obstet Gynecol, 197*(3), 223-
228, doi:10.1016/j.ajog.2007.03.027.
9. Frias, A. E., Morgan, T. K., Evans, A. E., Rasanen, J., Oh, K. Y., Thornburg, K. L., et al. (2011).
Maternal high-fat diet disturbs uteroplacental hemodynamics and increases the frequency
of stillbirth in a nonhuman primate model of excess nutrition. *Endocrinology, 152*(6), 2456-
2464, doi:en.2010-1332 [pii] 10.1210/en.2010-1332.
10. Rogers, L. K. V., M. (2011). Maternal inflammation, growth retardation, and preterm birth:
insights into adult cardiovascular disease. *Life Sci, 89*(13-14), 417-421, doi:S0024-
3205(11)00363-8 [pii] 10.1016/j.lfs.2011.07.017.
11. Retnakaran, R., Kramer, C. K., Ye, C., Kew, S., Hanley, A. J., Connelly, P. W., et al. (2015).
Fetal sex and maternal risk of gestational diabetes mellitus: the impact of having a boy.
Diabetes Care, 38(5), 844-851, doi:10.2337/dc14-2551.
12. Retnakaran, R., & Shah, B. R. (2015). Fetal Sex and the Natural History of Maternal Risk of
Diabetes During and After Pregnancy. *J Clin Endocrinol Metab, 100*(7), 2574-2580,
doi:10.1210/jc.2015-1763.
13. Jarmuzek, P., Wielgos, M., & Bomba-Opon, D. (2015). Placental pathologic changes in
gestational diabetes mellitus. *Neuro Endocrinol Lett, 36*(2), 101-105.
14. Tabacu, M. C., Istrate-Ofiţeru, A. M., Manolea, M. M., Dijmărescu, A. L., Rotaru, L. T.,
Boldeanu, M. V., et al. (2022). Maternal obesity and placental pathology in correlation with

adverse pregnancy outcome. *Rom J Morphol Embryol, 63*(1), 99-104, doi:10.47162/rjme.63.1.09.

15. Meng, Q., Shao, L., Luo, X., Mu, Y., Xu, W., Gao, C., et al. (2015). Ultrastructure of placenta of gravidas with gestational diabetes mellitus. *Obstetrics and gynecology international, 2015*.

16. Abdelghany, A., Eissa, T., & Idris, S. (2018). Study of the ultrastructure of the placenta in gestational Diabetes mellitus. *Int J Anat Var, 11*, 4-10.

17. Gorgoulis, V., Adams, P. D., Alimonti, A., Bennett, D. C., Bischof, O., Bishop, C., et al. (2019). Cellular Senescence: Defining a Path Forward. *Cell, 179*(4), 813-827, doi:10.1016/j.cell.2019.10.005.

18. Farfán-Labonne, B., Leff-Gelman, P., Pellón-Díaz, G., & Camacho-Arroyo, I. (2023). Cellular senescence in normal and adverse pregnancy. *Reproductive Biology, 23*(1), 100734, doi:10.1016/j.repbio.2023.100734.

19. Muñoz-Espín, D., Cañamero, M., Maraver, A., Gómez-López, G., Contreras, J., Murillo-Cuesta, S., et al. (2013). Programmed cell senescence during mammalian embryonic development. *Cell, 155*(5), 1104-1118, doi:10.1016/j.cell.2013.10.019.

20. Iwasaki, K., Abarca, C., & Aguayo-Mazzucato, C. (2023). Regulation of Cellular Senescence in Type 2 Diabetes Mellitus: From Mechanisms to Clinical Applications. *Diabetes Metab J, 47*(4), 441-453, doi:10.4093/dmj.2022.0416.

21. McCarthy, C. G., Wenceslau, C. F., Webb, R. C., & Joe, B. (2019). Novel Contributors and Mechanisms of Cellular Senescence in Hypertension-Associated Premature Vascular Aging. *Am J Hypertens, 32*(8), 709-719, doi:10.1093/ajh/hpz052.

22. Passos, J. F., Saretzki, G., Ahmed, S., Nelson, G., Richter, T., Peters, H., et al. (2007). Mitochondrial Dysfunction Accounts for the Stochastic Heterogeneity in Telomere-Dependent Senescence. *PLoS Biology, 5*(5), e110, doi:10.1371/journal.pbio.0050110.

23. Passos, J. F., Nelson, G., Wang, C., Richter, T., Simillion, C., Proctor, C. J., et al. (2010). Feedback between p21 and reactive oxygen production is necessary for cell senescence. *Mol Syst Biol, 6*, 347, doi:10.1038/msb.2010.5.

24. Huang, W., Xie, W., Zhong, H., Cai, S., Huang, Q., Liu, Y., et al. (2022). Cytosolic p53 Inhibits Parkin-Mediated Mitophagy and Promotes Acute Liver Injury Induced by Heat Stroke. *Front Immunol, 13*, 859231, doi:10.3389/fimmu.2022.859231.

25. Ahmad, T., Sundar, I. K., Lerner, C. A., Gerloff, J., Tormos, A. M., Yao, H., et al. (2015). Impaired mitophagy leads to cigarette smoke stress-induced cellular senescence: implications for chronic obstructive pulmonary disease. *Faseb j, 29*(7), 2912-2929, doi:10.1096/fj.14-268276.

26. Correia-Melo, C., Marques, F. D. M., Anderson, R., Hewitt, G., Hewitt, R., Cole, J., et al. (2016). Mitochondria are required for pro-ageing features of the senescent phenotype. *The EMBO Journal, 35*(7), 724-742-742, doi:https://doi.org/10.15252/embj.201592862.

27. Maeda, M., Hayashi, T., Mizuno, N., Hattori, Y., & Kuzuya, M. (2015). Intermittent High Glucose Implements Stress-Induced Senescence in Human Vascular Endothelial Cells: Role of Superoxide Production by NADPH Oxidase. *PLOS ONE, 10*(4), e0123169, doi:10.1371/journal.pone.0123169.

28. Chow, H. M., Shi, M., Cheng, A., Gao, Y., Chen, G., Song, X., et al. (2019). Age-related hyperinsulinemia leads to insulin resistance in neurons and cell-cycle-induced senescence. *Nat Neurosci, 22*(11), 1806-1819, doi:10.1038/s41593-019-0505-1.

29. Saucedo, R., Ortega-Camarillo, C., Ferreira-Hermosillo, A., Díaz-Velázquez, M. F., Meixueiro-Calderón, C., & Valencia-Ortega, J. (2023). Role of Oxidative Stress and Inflammation in Gestational Diabetes Mellitus. *Antioxidants (Basel), 12*(10), doi:10.3390/antiox12101812.

30. Choudhury, A. A., & Devi Rajeswari, V. (2021). Gestational diabetes mellitus - A metabolic and reproductive disorder. *Biomedicine & Pharmacotherapy, 143*, 112183, doi:https://doi.org/10.1016/j.biopha.2021.112183.

31. Kajdy, A., Modzelewski, J., Cymbaluk-Płoska, A., Kwiatkowska, E., Bednarek-Jędrzejek, M., Borowski, D., et al. (2021). Molecular Pathways of Cellular Senescence and Placental Aging in Late Fetal Growth Restriction and Stillbirth. *International Journal of Molecular Sciences, 22*(8), 4186, doi:10.3390/ijms22084186.

32. Scaife, P. J., Simpson, A., Kurlak, L. O., Briggs, L. V., Gardner, D. S., Pipkin, F. B., et al. (2021). Increased Placental Cell Senescence and Oxidative Stress in Women with Pre-Eclampsia and Normotensive Post-Term Pregnancies. *International Journal of Molecular Sciences, 22*(14), 7295, doi:10.3390/ijms22147295.

33. Biron-Shental, T., Sukenik-Halevy, R., Naboani, H., Liberman, M., Kats, R., & Amiel, A. (2015). Telomeres are shorter in placentas from pregnancies with uncontrolled diabetes. *Placenta, 36*(2), 199-203, doi:10.1016/j.placenta.2014.11.011.

34. Li-Zhen, L., Yun, X., Xiao-Dong, Z., Shu-Bin, H., Zi-Lian, W., Adrian Sandra, D., et al. (2019). Evaluation of guidelines on the screening and diagnosis of gestational diabetes mellitus: systematic review. *BMJ Open, 9*(5), e023014, doi:10.1136/bmjopen-2018-023014.

35. Eis, A. L., Brockman, D. E., Pollock, J. S., & Myatt, L. (1995). Immunohistochemical localization of endothelial nitric oxide synthase in human villous and extravillous trophoblast populations and expression during syncytiotrophoblast formation in vitro. *Placenta, 16*(2), 113-126, doi:10.1016/0143-4004(95)90000-4.

36. Gerland, L. M., Peyrol, S., Lallemand, C., Branche, R., Magaud, J. P., & Ffrench, M. (2003). Association of increased autophagic inclusions labeled for beta-galactosidase with fibroblastic aging. *Exp Gerontol, 38*(8), 887-895, doi:10.1016/s0531-5565(03)00132-3.

37. Dimri, G. P., Lee, X., Basile, G., Acosta, M., Scott, G., Roskelley, C., et al. (1995). A biomarker that identifies senescent human cells in culture and in aging skin in vivo. *Proc Natl Acad Sci U S A, 92*(20), 9363-9367, doi:10.1073/pnas.92.20.9363.

38. Kim, S. Y., England, L., Wilson, H. G., Bish, C., Satten, G. A., & Dietz, P. (2010). Percentage of gestational diabetes mellitus attributable to overweight and obesity. *American journal of public health, 100*(6), 1047-1052, doi:10.2105/AJPH.2009.172890.

39. Torloni, M. R., Betrán, A. P., Horta, B. L., Nakamura, M. U., Atallah, A. N., Moron, A. F., et al. (2009). Prepregnancy BMI and the risk of gestational diabetes: a systematic review of the literature with meta-analysis. *Obes Rev, 10*(2), 194-203, doi:10.1111/j.1467-789X.2008.00541.x.

40. Chuprin, A., Gal, H., Biron-Shental, T., Biran, A., Amiel, A., Rozenblatt, S., et al. (2013). Cell fusion induced by ERVWE1 or measles virus causes cellular senescence. *Genes Dev, 27*(21), 2356-2366, doi:10.1101/gad.227512.113.

41. Stein, G. H., Drullinger, L. F., Soulard, A., & Dulić, V. (1999). Differential roles for cyclin-dependent kinase inhibitors p21 and p16 in the mechanisms of senescence and differentiation in human fibroblasts. *Mol Cell Biol, 19*(3), 2109-2117, doi:10.1128/mcb.19.3.2109.

42. el-Deiry, W. S., Harper, J. W., O'Connor, P. M., Velculescu, V. E., Canman, C. E., Jackman, J., et al. (1994). WAF1/CIP1 is induced in p53-mediated G1 arrest and apoptosis. *Cancer Res, 54*(5), 1169-1174.

43. Burton, D. G. A., & Krizhanovsky, V. (2014). Physiological and pathological consequences of cellular senescence. *Cellular and Molecular Life Sciences, 71*(22), 4373-4386, doi:10.1007/s00018-014-1691-3.

44. Worman, H. J., Fong, L. G., Muchir, A., & Young, S. G. (2009). Laminopathies and the long strange trip from basic cell biology to therapy. *J Clin Invest, 119*(7), 1825-1836, doi:10.1172/jci37679.

45. Yoon, M.-H., Kang, S.-m., Lee, S.-J., Woo, T.-G., Oh, A.-Y., Park, S., et al. (2019). p53 induces senescence through Lamin A/C stabilization-mediated nuclear deformation. *Cell Death & Disease, 10*(2), 107, doi:10.1038/s41419-019-1378-7.

46. Dos Santos, Á., & Toseland, C. P. (2021). Regulation of Nuclear Mechanics and the Impact on DNA Damage. *International Journal of Molecular Sciences, 22*(6), doi:10.3390/ijms22063178.

47. Lu, X., Wang, R., Zhu, C., Wang, H., Lin, H.-Y., Gu, Y., et al. (2017). Fine-Tuned and Cell-Cycle-Restricted Expression of Fusogenic Protein Syncytin-2 Maintains Functional Placental Syncytia. *Cell Reports, 21*(5), 1150-1159, doi:10.1016/j.celrep.2017.10.019.

48. Gal, H., Lysenko, M., Stroganov, S., Vadai, E., Youssef, S. A., Tzadikevitch-Geffen, K., et al. (2019). Molecular pathways of senescence regulate placental structure and function. *The EMBO Journal, 38*(18), e100849, doi:10.15252/embj.2018100849.

49. Tuttle, C. S. L., Luesken, S. W. M., Waaijer, M. E. C., & Maier, A. B. (2021). Senescence in tissue samples of humans with age-related diseases: A systematic review. *Ageing Research Reviews, 68*, 101334, doi:https://doi.org/10.1016/j.arr.2021.101334.

50. Calcinotto, A., Kohli, J., Zagato, E., Pellegrini, L., Demaria, M., & Alimonti, A. (2019). Cellular Senescence: Aging, Cancer, and Injury. *Physiol Rev, 99*(2), 1047-1078, doi:10.1152/physrev.00020.2018.

51. Fisher, J. J., Vanderpeet, C. L., Bartho, L. A., McKeating, D. R., Cuffe, J. S. M., Holland, O. J., et al. (2021). Mitochondrial dysfunction in placental trophoblast cells experiencing gestational diabetes mellitus. *J Physiol, 599*(4), 1291-1305, doi:10.1113/jp280593.

52. Tang, Y., Zhang, X., Zhang, Y., Feng, H., Gao, J., Liu, H., et al. (2021). Senescent Changes and Endoplasmic Reticulum Stress May Be Involved in the Pathogenesis of Missed Miscarriage. *Frontiers in Cell and Developmental Biology, 9*, 656549, doi:10.3389/fcell.2021.656549.

53. Higuchi, S., Miyamoto, T., Kobara, H., Yamada, S., Asaka, R., Kikuchi, N., et al. (2019). Trophoblast type-specific expression of senescence markers in the human placenta. *Placenta, 85*, 56-62, doi:10.1016/j.placenta.2019.06.377.

54. Garcia-Martin, I., Penketh, R. J. A., Janssen, A. B., Jones, R. E., Grimstead, J., Baird, D. M., et al. (2018). Metformin and insulin treatment prevent placental telomere attrition in boys exposed to maternal diabetes. *PLoS ONE, 13*(12), e0208533, doi:10.1371/journal.pone.0208533.

55. Fischer, M. (2017). Census and evaluation of p53 target genes. *Oncogene, 36*(28), 3943-3956, doi:10.1038/onc.2016.502.

56. Storer, M., Mas, A., Robert-Moreno, A., Pecoraro, M., Ortells, M. C., Di Giacomo, V., et al. (2013). Senescence is a developmental mechanism that contributes to embryonic growth and patterning. *Cell, 155*(5), 1119-1130, doi:10.1016/j.cell.2013.10.041.

57. Yang, D.-G., Liu, L., & Zheng, X.-Y. (2008). Cyclin-dependent kinase inhibitor p16INK4a and telomerase may co-modulate endothelial progenitor cells senescence. *Ageing Research Reviews, 7*(2), 137-146, doi:https://doi.org/10.1016/j.arr.2008.02.001.

58. Molofsky, A. V., Slutsky, S. G., Joseph, N. M., He, S., Pardal, R., Krishnamurthy, J., et al. (2006). Increasing p16INK4a expression decreases forebrain progenitors and neurogenesis during ageing. *Nature, 443*(7110), 448-452, doi:10.1038/nature05091.

59. Shi, T., van Soest, D. M. K., Polderman, P. E., Burgering, B. M. T., & Dansen, T. B. (2021). DNA damage and oxidant stress activate p53 through differential upstream signaling pathways. *Free Radical Biology and Medicine, 172*, 298-311, doi:https://doi.org/10.1016/j.freeradbiomed.2021.06.013.

60. Marchenko, N. D., & Moll, U. M. (2014). Mitochondrial death functions of p53. *Mol Cell Oncol, 1*(2), e955995, doi:10.1080/23723548.2014.955995.

61. Aliouat-Denis, C. M., Dendouga, N., Van den Wyngaert, I., Goehlmann, H., Steller, U., van de Weyer, I., et al. (2005). p53-independent regulation of p21Waf1/Cip1 expression and senescence by Chk2. *Mol Cancer Res, 3*(11), 627-634, doi:10.1158/1541-7786.mcr-05-0121.

62. Abbas, T., & Dutta, A. (2009). p21 in cancer: intricate networks and multiple activities. *Nature*
Reviews Cancer, 9(6), 400-414, doi:10.1038/nrc2657.
63. Dyson, N. (1998). The regulation of E2F by pRB-family proteins. *Genes Dev, 12*(15), 2245-
2262, doi:10.1101/gad.12.15.2245.
64. Kumari, R., & Jat, P. (2021). Mechanisms of Cellular Senescence: Cell Cycle Arrest and
Senescence Associated Secretory Phenotype. *Frontiers in Cell and Developmental*
Biology, 09, 645593, doi:10.3389/fcell.2021.645593.
65. Engeland, K. (2022). Cell cycle regulation: p53-p21-RB signaling. *Cell Death & Differentiation,*
29(5), 946-960, doi:10.1038/s41418-022-00988-z.
66. Yin, M., Zhang, Y., Yu, H., & Li, X. (2021). Role of Hyperglycemia in the Senescence of
Mesenchymal Stem Cells. [Review]. *Frontiers in Cell and Developmental Biology, 9*(909),
doi:10.3389/fcell.2021.665412.
67. Wiley, C. D., & Campisi, J. (2021). The metabolic roots of senescence: mechanisms and
opportunities for intervention. *Nature Metabolism, 3*(10), 1290-1301, doi:10.1038/s42255-
021-00483-8.
68. Muralimanoharan, S., Maloyan, A., & Myatt, L. (2016). Mitochondrial function and glucose
metabolism in the placenta with gestational diabetes mellitus: role of miR-143. *Clin Sci*
(Lond), 130(11), 931-941, doi:10.1042/cs20160076.
69. Qiu, C., Hevner, K., Abetew, D., Sedensky, M., Morgan, P., Enquobahrie, D. A., et al. (2013).
Mitochondrial DNA copy number and oxidative DNA damage in placental tissues from
gestational diabetes and control pregnancies: a pilot study. *Clin Lab, 59*(5-6), 655-660,
doi:10.7754/clin.lab.2012.120227.
70. Abbade, J., Klemetti, M. M., Farrell, A., Ermini, L., Gillmore, T., Sallais, J., et al. (2020).
Increased placental mitochondrial fusion in gestational diabetes mellitus: an adaptive
mechanism to optimize feto-placental metabolic homeostasis? *BMJ Open Diabetes*
Research & Care, 8(1), e000923, doi:10.1136/bmjdrc-2019-000923.

662