Step-by-step protocol for making a knock-in *Xenopus laevis* to visualize

endogenous gene expression

Norie Kagawa[1], Yoshihiko Umesono[1], Ken-ichi T Suzuki[2] and Makoto Mochii[1]*

[1]Department of Life Science, Graduate School of Science, University of Hyogo, Ako-gun, Hyogo, 678-1297, Japan

[2]Emerging Model Organisms Facility, Trans-scale Biology Center, National Institute for Basic Biology, National Institutes of Natural Sciences, Okazaki, Aichi, 444-8585, Japan

* Correspondence: mmochii@sci.u-hyogo.ac.jp

Department of Life Science, Graduate School of Science, University of Hyogo, Ako-gun, Hyogo, 678-1297, Japan

Abstract

We established a novel knock-in technique, New and Easy *Xenopus* Targeted

integration (*NEXTi*), to recapitulate endogenous gene expression by reporter expression.

NEXTi is a CRISPR-Cas9-based method to integrate a donor DNA containing a reporter

gene (*egfp*) into target 5′ untranslated region (UTR) of *Xenopus laevis* genome. It

enables us to track eGFP expression under regulation of endogenous promoter/enhancer

activities. We obtained about 2% to 13% of knock-in vector-injected embryos showing

eGFP signal in a tissue-specific manner, targeting *krt.12.2.L*, *myod1.S*, *sox2.L* and

bcan.S loci, as previously reported. In addition, F1 embryos which show stable eGFP

signals were obtained by outcrossing the matured injected frogs with wild-type animals.

Integrations of donor DNAs into target 5′ UTRs were confirmed by PCR amplification

and sequencing. Here, we describe the step-by-step protocol for preparation of donor

DNA and single guide RNA, microinjection and genotyping of F1 animals for the

NEXTi procedure.

Key words

1. Highlight

- Intended organism: *Xenopus laevis*

- Purpose of the protocol: Efficient knock-in for visualizing endogenous target gene expression using CRISPR-Cas9 system

- Essential equipment and materials: Microinjector, fluorescence microscopy, Cas9 protein, sgRNA, donor DNA

- Features: Expression of reporter genes depends on endogenous enhancer/promoter activities.

2. Background and Features

Targeted integration or knock-in of a donor construct into a desired locus is a powerful tool which enables visualization of endogenous gene expression or manipulation of gene function in both model organisms and non-model organisms including amphibians. We previously established a simple transgenic method called New and Easy *Xenopus* Transgenesis (*NEXTrans*) in *Xenopus laevis* (Shibata et al*., 2022;* Shibata et al., 2023), by which the donor plasmid was efficiently integrated into the *tgfbr2l* loci, the proposed safe harbor sites, through the non-homologous end joining (NHEJ) repair induced by Cas9 ribonucleoprotein (RNP). Using a similar strategy to *NEXTrans* we recently

succeeded in the targeted integrations of a reporter gene, an enhanced green fluorescent

protein (eGFP) gene, into 5′ UTRs of various target genes to enable visualization of

endogenous genes' expression and termed the method New and Easy *Xenopus* Targeted

integration (*NEXTi*) (Mochii et al., 2024). In the *NEXTi* procedure, fertilized eggs were

injected with Cas9, a single guide RNA (sgRNA) for the 5′ UTR of the target gene and

a donor plasmid containing *egfp* flanked by the target genomic sequence (Fig. 1A). The

donor *egfp* was integrated into the target locus through essentially the NHEJ process

and was expressed under control of the endogenous promoter/enhancer activity (Fig.

1B). This versatile method can be applied for targeted integrations into various genome

regions and will contribute much to developmental and regeneration studies in *Xenopus*.

Advantages

NEXTi enables faithful recapitulation of the endogenous gene expression and can be

applied for any target gene in *X. laevis*. As sgRNA target sequence is located within the

5′ UTR of the target locus in this method, some insertions and/or deletions (indel) of

nucleotide at the junction between the target sequence and injected donor usually do not

disturb expression of the reporter gene. Therefore, *NEXTi* is superior to knock-in

methods targeting cording regions in which indel often causes serious effects on the reporter expression.

Limitations

As the donor integration is mediated essentially through the NHEJ process, orientation of the integration is not controlled and probably random in *NEXTi*. The reporter gene integrated in reverse orientation will be expressed incorrectly. The donor DNA should not contain the core enhancer of the target gene locus. Such the enhancer will induce reporter expression mimicking endogenous expression even in off-target insertion. Genes with a short 5′ UTR could not be targeted by *NEXTi*, since an sgRNA could not be designed within the short sequence. Integration of the donor within the 5′UTR will interfere normal expression of the targeted allele and may disturb normal development. As the efficiency of the targeted integration varies among different contexts, it may be necessary to try multiple sgRNAs for a single locus in order to achieve successful integration.

Applications

As mosaic expression of the reporter gene is frequently observed in the founders, we obtained F1 offspring with stable reporter expression in multiple lines (Mochii et al., 2024; this work). The *NEXTi* strategy can be applied for targeted integrations into various genomic regions other than 5′UTR. For example, integration of a modified exon into an intron will result in a replacement of the target coding sequence with a modified one.

3. Materials and Equipment

3.1 Solution

10 × Marc's Modified Ringers (MMR) stock solution

Reagent	Final concentration	amount
NaCl (FUJIFILM Wako, 191-01665)	1 M	58.44 g
KCl (Nacali Tesque, 28514-75)	20 mM	74.55 g
$MgCl_2 \cdot 6H_2O$ (FUJIFILM Wako,136-03995)	10 mM	2.03 g
$CaCl_2$ (FUJIFILM Wako, 039-00475)	20 mM	2.94 g
HEPES (Sigma-Aldrich, H3375-500G)	50 mM	11.92 g
10 N NaOH (FUJIFILM Wako, 198-13765)	N/A	N/A
Deionized water	N/A	Up to 1 L
Total	**N/A**	**1 L**

Adjust the solution to pH 7.5 with 10 N NaOH.

0.2% Ethyl 3-aminobenzoate methanesulfonate (MS222)

Reagent	Final concentration	Amount
MS222 (Sigma-Aldrich, A5040-100G)	0.2% (w/v)	2 g
NaHCO$_3$ (FUJIFILM Wako, 191-01305)	16.7 mM	1.4 g
Deionized water	N/A	Up to 1 L
Total	**N/A**	**1 L**

De-jellying solution

Reagent	Final concentration	Amount
L-cysteine (FUJIFILM Wako, 037-05275)	2% (w/v)	4 g
10 N NaOH (FUJIFILM Wako, 198-13765)	N/A	N/A
Deionized water	N/A	Up to 200 mL
Total	**N/A**	**200 mL**

Adjust the solution to pH 8.0 with 10N NaOH.

10x Cas9 buffer

Reagent	Final concentration	Amount
2 M KCl (Invitrogen, AM9640G)	1.5 M	15 mL
1 M HEPES (pH 7.5) (Nacalai Tesque, 15639-84)	0.2 M	4 mL
Nuclease-free water (Nacalai Tesque, 06442-95)	N/A	1 mL
Total	**N/A**	**20 mL**

3% Ficoll-0.5 x MMR (-0.1 x MMR)

Reagent	Final concentration	Amount
10 × MMR	0.5 × MMR (0.1 × MMR)	25 mL (5 mL)
Ficoll PM400 (Sigma-Aldrich, F4375) (GE Healthcare, 17-0300-50)	3% (w/v)	15 g
Kanamycin Sulfate (FUJIFILM Wako, 113-00343)	20 ng/mL	10 mg
Deionized water	N/A	up to 500 mL
Total	**N/A**	**500 mL**

0.3% methyl-cellurose-0.5 x MMR (-0.1 x MMR)

Reagent	Final concentration	Amount
10 × MMR	0.5 × MMR (0.1 × MMR)	25 mL (5 mL)
Methyl Cellulose 400 (FUJIFILM Wako, 132-05055)	0.3% (w/v)	1.5 g
Kanamycin Sulfate (FUJIFILM Wako, 113-00343)	20 ng/mL	10 mg
Deionized water	N/A	up to 500 mL
Total	**N/A**	**500**

3.2 Setup

3.2.1 Microneedle

Microneedles are made by pulling glass capillaries (Drummond Scientific, 3-000-204R)

with a puller (e.g. Narishige, PC-100). The tip of the needle is sharpened and adjusted to

an outer diameter of 20–25μm (See Tip1).

3.2.2   Injection plate

To make an injection plate a 60 mm petri dish is covered with a modeling clay,

Plastalina. Spread the clay to make it flat and make grooves in it (See Tip2).

3.3 Other reagents and equipment

PrimeSTAR MAX (Takara Bio, R010A)

PureYield Plasmid System (Promega, A2490)

Ex Taq Hot Start Version (Takara, RR006)

Wizard SV Gel and PCR clean-up system (Promega, A9282)

T7 RiboMax express large-scale RNA production system (Promega, P1320)

Alt-R S.p. Cas9 Nuclease 3NLS (Integrated DNA Technologies, 1081058)

Tris-HCl (Sigma-Aldrich, T5941)

EDTA (Sigma-Aldrich, E-6511)

Sodium lauryl sulfate (SDS) (Nacali Tesque, 31607-65)

Proteinase K (FUJIFILM Wako, 160-14001)

RNase A (Sigma-Aldrich, from bovine pancreas)

Serotropin (1,000 U/glass ampoule) (ASKA Animal Health)

Human Chorionic Gonadotropin (HCG) (5,000 U/glass ampoule) (ASKA Animal

Health)

Fluorescent stereomicroscope (Leica, MZ165FC)

Microinjector (Drummond Scientific, 3-000-204)

Micromanipulator (Drummond Scientific, 3-000-204R)

PCR thermal cycler (ASTEC, G02)

NanoDrop One (Thermo Fisher scientific, ND-ONE-W)

96 well plate (VIOLAMO, 1-1601-02)

- 0.5 U/µL Serotropin solution: dissolve Serotropin in 1 mL of 0.6% NaCl

  solution (supplied).

- 2 U/µL Human Chorionic Gonadotropin (HCG) solution: dissolve HCG in 1.6

  mL of 0.6% NaCl solution (supplied).

- 10 mM Tris-EDTA (TE) buffer (pH 7.5): dissolve 8.88 g of Tris-HCl and 3.72 g

  of EDTA in 1 L of deionized water. Adjust the pH with 5N NaOH.

4. Procedure

4.1 Preparation of *NEXTi* transgenic plasmid (donor)

NEXTi donor plasmids should contain genomic DNA fragments homologous to target sequences (Fig. 1B). While homologous fragments that ranged from 66 bp to 844 bp were used for the donors and resulted in efficient integrations (Table. 1), a longer fragment in the donor resulted in a higher integration rate at least in the case of the *krt12.2.L* locus (Mochii et al., 2024). We usually amplify genomic DNA of 500 – 900 bp including the target 5′UTR and clone it into a vector plasmid using conventional procedures.

1. Clone the DNA fragment including the 5′ UTR sequence of the target gene into a plasmid vector to construct the *NEXTi* donor. For example, an 844 bp fragment containing 698 bp upstream and 146 bp 5′ UTR sequences of *myod1.S* was amplified from the genomic DNA using a high-fidelity polymerase (e.g., PrimeSTAR Max) and cloned into a promoter-less *egfp* plasmid (Suzuki et al., 2010) by standard plasmid construction methods (Mochii et al., 2024) (See Tip 3).

2. Purify the DNA using a standard plasmid purification kit (e.g., PureYield Plasmid System, Promega).

3. (Optional) Extract the DNA with phenol-chloroform and precipitate it with ethanol.

    Dissolve the DNA in RNase-free TE buffer at 0.5 –1.0 µg/µL (See Tip 4).

4. Dilute the DNA with 1 x Cas 9 buffer at 50 ng /µL, divide it into small aliquots (5 –

    10 µL) and store them at –20˚C for working stocks.

4.2 Synthesis of sgRNA

5. The Cas9 targets should be selected within the 5´ UTR sequence cloned into the

    donor DNA (See Tip5). A PCR-based strategy was used for sgRNA preparation

    essentially as previously described (Nakayama et al., 2013; Sakane et al., 2017).

6. Prepare the sequence-specific 5´ oligonucleotides (e.g., for two target sequences of

    myod1.S 5´UTR) and the common 3´ oligonucleotide.

    *myod1.S 5´*oligo 1:

    5´- TAATACGACTCACTATAGGGGCAAGGTGACTGTGCGTG

    TTTTAGAGCTAGAAATAGCAAG -3´

    myod1.S 5´ oligo 2:

    5´- TAATACGACTCACTATAGGTTAAAAGATCCACAGCTCG

    TTTTAGAGCTAGAAATAGCAAG -3´

    Common 3´ oligo:

5´- AAAAGCACCGACTCGGTGCCACTTTTTCAAGTTGATAAC

GGACTAGCCTTATTTTAACTTGCTATTTCTAGCTCTAAAAC -3´

7. Amplify the template DNA by PCR using the oligonucleotides.

PCR master mix

Reagent	Amount
Ex Taq Hot Start Version	0.5 µL
10 x Ex Buffer	5 µL
dNTP Mixture (2.5 nM)	4 µL
5´ oligonucleotide (100 µM)	1 µL
3´ oligonucleotide (100 µM)	1 µL
Deionized water	38.5 µL
Total	**50 µL**

PCR cycling conditions

Step	Temperature	Time	Cycles
Initial Denaturation	98˚C	10 sec	1
Denaturation	98˚C	10 sec	
Annealing	60˚C	30 sec	12
Extension	72˚C	15 sec	
Final extension	72˚C	5 min	1
Hold	4˚C	forever	1

8. Check if the target PCR product is detected as a single band (117 bp) by standard

   agarose or polyacrylamide gel electrophoresis.

9. Purify the template DNA using a purification kit (e.g., Wizard SV Gel and PCR

clean- up system, Promega, A9282). Adjust the concentration to 0.1 µg/µL.

10. Synthesize and purify the sgRNA by using an *in vitro* transcription kit based on T7

RNA polymerase (e.g., Promega, P1320) following the manufacturer's instructions.

Reaction mixture for the RNA synthesis (Promega, P1320)

Reagent	Amount
Template DNA (0.2 µg)	2 µL
RiboMAX™ Express T7 2X Buffer	10 µL
Enzyme Mix, T7 Express	2 µL
Nuclease-free water	6 µL
Total	**20 µL**

a. Incubate the reaction mixture at 37°C for 3 – 4 h.

b. Add 1 µL of RNase-free DNase (1 µL/µL) and incubate at 37°C for 15

min.

c. Add 115 µL of water and 15 µL of 3M sodium acetate to stop the

reaction.

d. Extract with phenol-chloroform.

e. Add 150 µL of isopropanol and leave at –20°C for at least 15 min.

f. Centrifuge at 15,000 rpm for 15min.

g. Remove the supernatant and add 70% EtOH.

h. Centrifuge at 15,000 rpm for 15 min.

    i.   Carefully remove the 70% EtOH.

    j.   Dry up the precipitate.

    k.   Dissolve sgRNA with nuclease-free water.

    l.   Adjust the concentration to $1 - 2$ µg/µL and store at $-80°C$ (See Tip6).

11. Prepare master mix for the sgRNA aliquots.

Reagent	Amount
sgRNA	8 µg
10 x Cas9 buffer	6.2 µl
Water	X µL
Total	**62 µL**

12. Dispense 6.2 µL of the solution into a tube and store at $-80°C$ (See Tip7).

4.3 In vitro fertilization

Standard procedures for *Xenopus* embryo manipulation should follow the protocols of

each laboratory or refer to textbooks (Hoppler and Vize, 2012; Sive et al., 2000).

Researchers should obtain approval from the Animal Care and Use Committee in their

affiliated institutes and perform experiments in accordance with their guideline. In this

study, we obtained approval from the committee of University of Hyogo.

1. Inject 50 U of serotropin solution into $2 - 4$ females $2 - 3$ days before use.

    The day before use, inject 500 U of HGC solution and keep females at 18˚C.

2. Sperm preparation

    a. On the day of injection, anesthetize a male in a water tank of 0.2% MS222.

    b. After euthanasia, isolate testes and remove the fat body with a pair of fine forceps under a stereomicroscope.

    c. Keep the isolated testes in 1 x MMR containing kanamycin on ice (See Tip8).

    d. Put a third or half of the testes into a 1.5 mL tube with 1 mL of 0.5 x MMR.

    e. Mince the testis fragment in the tube using an ophthalmic scissor.

3. Collect 300–500 eggs by gently squeezing the females and keep them in separate sterile Petri dishes.

4. Add the sperm suspension onto the eggs, mix and spread the eggs gently using a pipet tip.

5. Remove as much of the liquid as possible from the dish and leave the eggs for 5min at 11˚C (See Tip9).

6. Soak the dish with 0.1 x MMR for 15 min at 11˚C (See Tip10).

7. De-jelly the embryos with de-jellying solution containing L-cysteine for 10 – 15 min

(See Tip11).

8. Rinse the embryos with 0.1 x MMR several times.

9. Keep the embryos at a low temperature (11˚C) to delay the first cleavage.

4.4 Injection

1. Dilute stock Cas9 protein to 1 µg/µL before use. For example, mix 0.5 µL of Cas9

   (10 µg/µL) with 4.5 µL of cold 1 x Cas9 buffer.

2. Prepare injection mixture consisting of the following.

Reagent	Amount
sgRNA solution	6.2 µL
donor DNA (50 ng/µL)	1 µL
Cas 9 protein (1 µg/µL)	2 µL
Total	**9.2 µL**

   After incubating the mixture at room temperature for 10 minutes, place it on ice.

3. Set a microneedle (See **3.2 Setup**) on the microinjector.

4. Fill the microneedle with 1–2 µL of injection mixture (See Tip 12).

5. Transfer 100 –150 fertilized eggs into the injection plate filled with 3-5% Ficoll-0.5

   x MMR. Gently shake the injection plate horizontally to direct the fertilized eggs to

   have their animal side up.

6. At 1 h after the in vitro fertilization, inject 9.2 nL of injection mixture into the animal

hemisphere of each fertilized egg. Complete the injection within 1 h (See Tip 13).

7. Transfer the injected eggs in another dish filled with 3% Ficoll–0.5 x MMR and incubate them at 18°C for 2–4 h.

8. Select normal embryos at 4 or 8 cell stages and transfer them into 0.1 x MMR. Unfertilized eggs and severely damaged embryos should be eliminated (see Tip 14).

9. Incubate the embryos at 18˚C for 3–4 days until the embryos develop to the late tailbud or tadpole stage (See Tip15–18).

10. Select the embryos with reporter gene expression (See Tip 19). Determine the embryo stages according to Nieuwkoop and Faber (1994).

4.5 Germline transmission

3.5.1 In vitro fertilization

1. Raise knock-in frogs to sexual maturation.

2. Outcross matured knock-in animals with wild-type male or female by in vitro fertilization.

4.6 Genotyping

1. Deeply anesthetize F1 embryos or tadpoles with MS222 and euthanatize them for

genomic DNA extraction.

2. Lyse them in 10 mM Tris-HCl (pH 8.0)-5 mM EDTA-0.5% SDS and add RNase A

    (10 µg/µL), then incubate at 37°C for 1 h.

3. Treat with Proteinase K (100 µg/µK) at 50°C for 3 h.

4. Extract DNA with phenol and precipitate with ethanol (See Tip 20).

5. Dissolve the DNA in TE buffer.

6. Amplify the junction sequence by using appropriate primer sets (See Tip 21).

7. Purify the PCR product with PCR purification system (e.g., Wizard SV Gel and PCR

    clean-up system, Promega, A9282) and sequence it.

5. Tips

1. The injection mixture is viscous. If the diameter of the needle tip is too small, the

    needle may become clogged with fluid, and it may not be possible to inject a sufficient

    volume of the mixture into the eggs. On the other hand, a needle with too-large a tip

    could seriously damage the injected embryos. We recommend a needle tip with an

    outer diameter of about 25 µm. The shape of the needle tip is also critical, as a needle

    with a flat tip damage embryos. The needle tip should be sharpened by breaking it

    with a pair of fine forceps or hitting against the edge of a glass slide. Example of the

needle tip is shown in Fig.S1.

2. Regarding use of plastalina is refer to a previous protocol (Ogino et al., 2006). Too much clay will cause the solution to overflow. Fill the Petri dish with the clay to less than half of its height. Coating an injection plate with agarose gel also works well.

3. We use the SliCE method (Okegawa and Motohashi, 2015) to clone the PCR product into the vector. Commercial kits (e.g., In-Fusion HD kit, Takara) are also recommended.

4. The donor DNA should be RNase-free.

5. We usually select two or three candidate sequences within a single target 5′ UTR using the CRISPRscan program (https://www.crisprscan.org, Moreno-Mateos et al., 2015) and prepare multiple sgRNAs for the single target gene. The rate of targeted integration is different among the different sgRNAs (Table 2).

6. Purification of sgRNA using a commercial purification kit is also recommended.

7. The solutions and tubes should be pre-chilled on ice to avoid degradation of the sgRNAs.

8. The isolated testes can be stored for several days at 4°C.

9. Excessive liquid causes floating of the eggs at the next step.

10. If the eggs float, quickly sink them using a pipet. Activation of the eggs is confirmed

by the contraction of melanin pigments. Do not use a batch of eggs in which the activation is observed in less than 50% of the eggs, because egg quality is critical for successful injection.

11. Insufficient de-jelling causes aggregation of the eggs or interferes with penetration of the microneedle.

12. Drop 1 – 2 µL of Cas9 solution on a parafilm and suck the solution into a microneedle. To prevent the tip of a microneedle from drying out, do not keep the tip in air.

13. Timing of injection is critical. Injection before 1 h or after 2 h post-fertilization results in reduced integration rate. If the rate of normal development is less than 50% after injection, reduce the injection volume, *e.g.*, to 4.6 nL per embryo. Reduced amount of injected sgRNA may reduce integration efficiency (Table 2).

14. It is important to remove unfertilized eggs and abnormally cleaved embryos which were seriously damaged by the injections. The dead or abnormal embryos exert harmful effects on the development of other embryos in the same dish.

15. Density of the embryos should be kept low. We recommend up to 100 embryos in a 100 mm Petri dish. Remove dead and abnormal embryos, and replace the medium every day.

16. Culturing a single embryo in 96-well plates increases the rate of normal development after injection. It protects healthy embryos from negative influence of dead embryos sharing the culture medium. In the case of the single embryo culture, the medium change is unnecessary until the embryos reach about NF stage 40.

17. 50-90% of the selected embryos develop into normal tadpoles in the standard condition.

18. Ficoll (3%) or methyl-cellulose (0.3%) in the culture medium increase the rate of normally developed embryos.

19. Ubiquitous or ectopic signals are frequently observed at early stages, especially at neural stage, in embryos injected with some donor constructs. Most of the signals become faint or disappeared by a later stage and are considered to be non-specific, probably due to *egfp* transcription from injected donor DNAs that were not integrated into chromosomes.

20. DNA purification using commercially available kit (*e.g.* Promega, A9282) is also recommended.

21. We usually perform the genomic PCR with primers which amplify 0.5 kb to 3 kb fragment. Primer sequences used in Fig.4 are indicated in Table.S1.

6. Anticipated results

6.1 Generation of founders

The tissue-specific eGFP signals were observed in F_0 embryos targeted for *myod1.S* (somite), *sox2.L* (central nerve system), and *krt12.2.L* loci (fin; Fig 2). The rate of embryos expressing the reporter gene in a tissue-specific pattern is varied from ~2% to ~13% (Table 1) (Mochii et al., 2024). The rate was different between different sgRNAs used for injection even with the same donor DNA. For example, injection of two different sgRNAs, *myod1.S* sg1 and sg2, resulted in 9.3% and 1.8% eGFP-expressing embryos, respectively (Table 2).

6.2 Germline transmission of integrated DNA

We confirmed germline transmission of the integrated DNAs by outcrossing matured founders and wild-type frogs by in vitro fertilization. Some F1 embryos of *sox2.L:egfp*, *bcan.S:egfp* and *krt12.2.L:egfp* founders showed eGFP signals in a tissue-specific manner like that observed in their founder embryos (Fig. 3 and data not shown) (Mochii et a., 2024). The rate of F1 embryos showing eGFP signals in a tissue-specific manner varied from 0.3% to 51.6% probably depending on the rate of the knocked-in germ cells in gonads (Table 3). On the other hand, F1 embryos with non-specific eGFP signal were

also obtained. In the case of *sox2.L:egfp* founder #2, 2.4% of F1 embryos showed

specific eGFP signal in neural tissues (Fig.3A, B), but 3.0% of the siblings showed a

weak ubiquitous signal (Fig. 3A, C).

  Genomic PCR showed the junction between the host genome and donor vector,

confirming integration into the 5´UTR of the target loci (Fig. 4). Clear bands were

detected in F1 embryos that showed eGFP signal in a tissue-specific manner for both

the upstream and downstream junction (Fig. 4B). On the other hand, the PCR band was

not detected in either junction in embryos with ubiquitous eGFP signal, suggesting an

off-target integration of the donor. The reporter gene integrated in off-target site may be

expressed with low specificity by a possible promoter contained in the donor DNA

fragment. Sequencing of the amplified fragments showed deletions in both the upstream

and downstream junction sites in many cases (Fig. 4C) but insertions in some cases

(Mochii et al., 2024).

Acknowledgments

We thank Dr. Elizabeth Nakajima for critical reading of our manuscript and Dr.

Hidefumi Orii for preparation of SLiCE. This research was supported by a JSPS

KAKENHI Grant-in-Aid for Scientific Research (C) (JP18K06266 to MM), Grant-in-

Aid for Scientific Research (B) (JP21H03829 to KTS), Special grant for research in

University of Hyogo (to MM), University of Hyogo Science and Technology Support

Foundation (to MM), NIBB Collaborative Research Program (22NIBB331, 23NIBB340

to MM), and JST, CREST, Grant Number JPMJCR 2025 (to KTS), Japan.

References

Hoppler S., Vize PD (2012). Xenopus Protocols: Post-Genomic Approaches. New Yolk: Human Press.

Mochii, M., Akiduki, K., Ossaka, H., Kagawa, N., Umesono, Y., Suzuki, K.T. (2024). A CRISPR-Cas9-mediated versatile method for targeted integration of a fluorescent protein gene to visualize endogenous gene expression in *Xenopus laevis.* Developmental Biology, 506, 42-51. https://doi.org/10.1016/j.ydbio.2023.11.010

Moreno-Mateos, M. A., Vejnar, C. E., Beaudoin, J. D., Fernandez, J. P., Mis, E. K., Khokha, M. K., & Giraldez, A. J. (2015). CRISPRscan: designing highly efficient sgRNAs for CRISPR-Cas9 targeting in vivo. Nature methods, 12, 982–988. https://doi.org/10.1038/nmeth.3543

Nakayama T, Fish MB, Fisher M, Oomen-Hajagos J, Thomsen GH, Grainger RM. (2013). Simple and efficient CRISPR/Cas9-mediated targeted mutagenesis in Xenopus tropicalis. Genesis, 51, 35-843. https://doi.org/10.1002/dvg.22720

Nieukoop, P. D. & Faber, J. (1994). Normal Table of Xenopus Laevis (Daudin).

Garland Publishing, New York.

Ogino H, McConnell WB, Grainger RM. (2006). High-throughput transgenesis in

Xenopus using I-SceI meganuclease. Nat Protoc. 1(4):1703-10.

https://doi.org/10.1038/nprot.2006.208

Okegawa, Y., Motohashi, K., 2015. A simple and ultra-low cost homemade seamless

ligation cloning extract (SLiCE) as an alternative to a commercially available seamless

DNA cloning kit. Biochem. Biophys. Rep. 4, 148–151.

https://doi.org/10.1016/j.bbrep.2015.09.005

Sakane, Y., Suzuki, K. T., & Yamamoto, T. (2017). A Simple Protocol for Loss-of-

Function Analysis in Xenopus tropicalis Founders Using the CRISPR-Cas

System. Methods in molecular biology, 1630, 189–203. https://doi.org/10.1007/978-1-

4939-7128-2_16

Shibata, Y., Suzuki, M., Hirose, N., Takayama, A., Sanbo, C., Inoue, T., Umesono, Y., Agata, K., Ueno, N., Suzuki, K. T., & Mochii, M. (2022). CRISPR/Cas9-based simple transgenesis in Xenopus laevis. Developmental biology, 489, 76–83.

https://doi.org/10.1016/j.ydbio.2022.06.001

Shibata, Y., Okumura, A., Mochii, M., & Suzuki, K. T. (2023). Protocols for transgenesis at a safe harbor site in the Xenopus laevis genome using CRISPR-Cas9. STAR protocols, 4(3), 102382. https://doi.org/10.1016/j.xpro.2023.102382

Sive HL, Grainger RM, Harland RM (2000). Early development of Xenopus laevis : a laboratory manual. New Yolk: Cold Spring Harbor Laboratory Pres.

Suzuki KT, Kashiwagi K, Ujihara M, Marukane T, Tazaki A, Watanabe K, Mizuno N, Ueda Y, Kondoh H, Kashiwagi A, Mochii M. (2010). Characterization of a novel type I keratin gene and generation of transgenic lines with fluorescent reporter genes driven by its promoter/enhancer in Xenopus laevis. Developmental Dynamics, 239:3172-81.

https://doi.org/10.1002/dvdy.22451

Suzuki, K.T., Suzuki, M., Shigeta, M., Joshua, F.D., Mawaribuchi, M., Takahashi, M., Yamamoto, T., Taira, M., Fukui, A. (2017). Clustered Xenopus keratin genes: a genomic, transcriptomic, and proteomic analysis. Developmental Biology, 426, 384–392. https://doi.org/10.1016/j.ydbio.2016.10.018

Zahn, N., James-Zorn, C., Ponferrada, V. G., Adams, D. S., Grzymkowski, J., Buchholz, D. R., Nascone-Yoder, N. M., Horb, M., Moody, S. A., Vize, P. D., & Zorn, A. M. (2022). Normal Table of Xenopus development: a new graphical resource. Development (Cambridge, England), 149, dev200356. https://doi.org/10.1242/dev.200356

Table 1. Homologous fragments cloned into donor plasmids and integration rates in founders by *NEXTi*

Target locus	Cloned fragment [a] (bp)	5´UTR sequence [b] (bp)	Rate of embryos with specific eGFP (%)	Total number of embryos	reference
krt12.2.L	537	66	13.2	167	Mochii et.al., 2024
	246	66	6.8	161	
	66	66	3.8	261	
sox2.L	787	218	2.1	1025	Mochii et.al., 2024
myod1.S	844	146	6.6	708	Mochii et.al., 2024
bcan.S	489	389	5.7	332	Mochii et.al., 2024

[a]Genomic DNA fragment cloned into donor plasmid. [b] 5´UTR sequence in cloned fragment.

Table 2. Integration efficiencies of founders using different sgRNAs or different

sgRNA doses

Target locus	sgRNA	Amount of sgRNA (pg / embryo)	Rate of embryos with specific eGFP[a] (%)	Total number of embryos	reference
myod1.S	sg1	800	9.3	86	This work
	sg2	800	1.8	112	
myod1.S	sg1	800	14.3	42	This work
	sg1	400	10	60	

Amount of donor DNA was constant. [a]Embryos with tissue-specific eGFP expression

including mosaic pattern.

Table 3. Germline transmission of transgene by *NEXTi*

Founder	Sex	Age (month)	eGFP expression	eGFP+ / total F1 embryos	Germline transmission rate (%)	Reference
sox2.L:egfp #1	Male	10	specific	86/183	47.0	Mochii et.al., 2024
sox2.L:egfp #2 [a]	Female	14	specific	19/789	2.4	This work
			ubiquitous	24/789	3.0	
bcan.S:egfp #1	Male	17	specific	80/260	30.8	This work
bcan.S:egfp #2	Female	18	specific	64/124	51.6	This work
krt12.2.L:egfp (fgk246b-egfp)	Male	14	specific	1/374	0.3	This work

[a]This founder produced two types of F1 embryos expressing eGFP, neural tissue-specific (specific) or throughout embryo (ubiquitous).

Figure Legends

Fig. 1 Overview of *NEXTi*

(A) Summary of the *NEXTi* procedure to make a knock-in founder and F1 offsprings.

The *NEXTi* donor plasmid carrying the *egfp* is injected into fertilized *Xenopus laevis*

eggs with Cas9 protein and a single guide RNA. Selected founder (F0) expressing

tissue-specific eGFP is outcrossed with wild-type animal (wt) to obtain F1 animals.

(B) Schematic representation of targeted integration by *NEXTi*.

The Cas9 RNP binds and cleaves both the target sequences in the donor plasmid and the

5´UTR of the target locus. The plasmid integrates into the target sites in the zygotic

genome via DSB/NHEJ. Insertion and/or deletion (indel) are found in the upstream and

downstream junctions.

Fig. 2 Expression of eGFP in knock-in vector-injected embryos.

Representative images of embryos targeted for *myod1.S* (A, A', NF stage 35/36),

sox2.L (B, B', NF stage 37/38) and *krt12.2.L* (C, C', NF stage 41) loci. (A-C), bright

field images. (A'-C'), fluorescence images of (A-C). Clear eGFP signal was observed in

somatic muscle (arrowhead in A'), brain and lens (arrowheads in B'), and fin

(arrowhead in C'). Scale bar = 1 mm.

Fig. 3 Expression of eGFP in F1 siblings of *sox2.L:egfp*

  Representative images of F1 embryos (NF stage 33/34) obtained by crossing a female

sox2.L:egfp founder #2 (Table. 3) and a wild-type male (A-D). (A) bright field images.

(A'), fluorescence image of (A). (B-D) magnified view of each embryo in (A'). (B)

specific eGFP signal is observed in brain (b), spinal cord (sc), optic vesicles (op),

olfactory pit (ol), and otic vesicles (ot). (C) Ubiquitous eGFP signal at low level is

observed. (D) No eGFP signal is observed. Scale bar = 1 mm.

Fig. 4 Genotyping of F1 animals

(A) Scheme of an integrated target locus. PCR was performed with genomic DNAs

obtained from F1 animals. Upstream and downstream sequences including junctions

between the donor and the target regions were amplified using donor vector primers and

gene-specific primers. Primer sequences are indicated in Table. S1. (B) Examples of

PCR analysis for *sox2.L:egfp* F1 embryos obtained from the founder #1 (1, 2) and

founder #2 (3-6). PCR products for both upstream and downstream junction fragments

were observed in DNAs from F1 embryos with specific eGFP expression (1-4) but not

from embryos with ubiquitous eGFP (5, 6) or wild-type embryos (wt). Predicted sizes of

the upstream and downstream PCR fragments are indicated as arrowheads. PCR

products with larger or smaller size indicate insertion or deletion in the junctions. (C)

Example of upstream and downstream junction sequences. The PCR products of

sox2.L:egfp F1 embryo (lane 3) were sequenced directly. Determined sequences are

indicated with wild-type partial sequence of the 5´UTR. Underline indicates the sgRNA

target sequence with PAM sequence (red underline).

Fig.1 Overview of *NEXTi*

(A) Summary of the *NEXTi* procedure to make a knock-in founder and F1 offsprings. The *NEXTi* donor plasmid carrying the *egfp* is injected into fertilized *Xenopus laevis* eggs with Cas9 protein and a single guide RNA. Selected founder (F0) expressing tissue-specific eGFP is outcrossed with wild type animal (wt) to obtain F1 animals. (B) Schematic representation of targeted integration by *NEXTi*. The Cas9 RNP digests both the target sequences in the donor plasmid and the 5′ UTR of the target locus. The plasmid integrates into the target sites in the zygotic genome via DSB/NHEJ. Insertion and/or deletion (indel) are found in the upstream and downstream junctions.

Fig.2 Expression of eGFP in vector-injected embryos targeted for *myod1.S, sox2.L, krt12.2.L* loci by *NEXTi*.

Representative images of embryos targeted for *myod1.S* (A, A', NF stage 35/36), *sox2.L* (B, B', NF stage 37/38) and *krt12.2.L* (C, C', NF stage 41) loci. (A-C), bright field images. (A'-C'), fluorescence images of (A-C). Clear eGFP signal was observed in somatic muscle (arrowhead in A'), brain and lens (arrowheads in B'), and fin (arrowhead in C'). Scale bar = 1 mm.

Fig.3 Expression of eGFP in F$_1$ siblings of *sox2.L:egfp*

Representative images of F1 embryos (NF stage 33/34) obtained by crossing a female *sox2.L:egfp* founder #2 and a wildtype male (A-D). (A) bright field images. (A'), fluorescence image of (A). (B-D) magnified view of each embryo in (A'). (B) specific eGFP signal is observed in brain (b), spinal cord (sc), optic vesicles (op), olfactory pit (ol), and otic vesicles (ot). (C) Ubiquitous eGFP signal at low level is observed. (D) No eGFP signal is observed. Scale bar = 1 mm.

A

B

C

wild type   TTGTCTATTAACTTGTTTCAAAAAAGTAGCAG<u>GACTTGTCACAAGCGCAGG</u><u style="color:red">GAGG</u>GACAGCAGTTGGCTC

Upstream   TTGTCTATTAACTTGTTTCAAAAAAGTAGCAG<u>GACTTG</u>-----<u>G-GCA</u>GG<u style="color:red">GAGG</u>GACAGCAGTTGGCTC 7b del

Downstream   TTGTCTATTAACTTGTTTCAAAAAAGTAGCAG<u>GACTTGTCACAAG</u>-----<u style="color:red">GGAGG</u>GACAGCAGTTGGCTC 5b del

Fig.4 Genotyping of F1 animals

(A) Scheme of an integrated target locus. PCR was performed with genomic DNAs obtained from F1 animals. Upstream and downstream sequences including junctions between donors and target regions were amplified using donor vector primers and gene-specific primers. Primer sequences are indicated in Table. S1. (B) Examples of PCR analysis for *sox2.L:egfp* F1 embryos obtained from the founder #1 (lane 1-2) and founder #2 (lane 3-5). PCR products for both upstream and downstream junction fragments were observed in DNAs from F1 embryos with specific eGFP expression (lane 1-4) but not from embryos with ubiquitous eGFP (lane 5-6) or wildtype embryos (wt). Predicted sizes of the upstream and downstream PCR fragments are indicated as arrowheads. PCR products with larger or smaller size indicate insertion or deletion in the junctions. (C) Example of upstream and downstream junction sequences. The PCR products of of *sox2.L:eegfp* F1 embryo (lane 3) were sequenced directly. Determined sequences are indicated with wild type sequence of the 5´UTR. Underline indicates the sgRNA target sequence with PAM sequence (red underline).

Fig.S1 Examples of microneedle tips

The sharp-tipped microneedle (A) is more suitable for the microinjection than the blunt-tipped one (B). Scale bar, 20 μm.

Primers	Sequences
sox2.L gFW	5'- TGTGCCCTTTAGTTTGATCCCT -3'
sox2.L gRV	5'- TATTCTGGTTGTTGGACGCAGA -3'
vector FW	5'- AACGACCTACACCGAACTGAGATAC -3'
egfp RV	5'- CAGGATGTTGCCGTCCTCCTTGAAGTCGAT -3'

Table.S1 primer sequences for genotyping