Combinatorial effects of multiple genes contribute to beneficial aneuploidy phenotypes

M. Sophie Koller[1,2], Claudia Himmelbauer [1], Sarah Fink[1], Madhwesh C. Ravichandran[1], Christopher S. Campbell[1@]

[1]Department of Chromosome Biology, Max Perutz Labs, University of Vienna, Vienna Biocenter (VBC), A-1030 Vienna, Austria

[2]Vienna Biocenter PhD Program, a Doctoral School of the University of Vienna and the Medical University of Vienna, A-1030 Vienna, Austria

@: Christopher S. Campbell
Email: christopher.campbell@univie.ac.at
Phone: +43-1-4277-74418
Address: Dr. Bohr-Gasse 9/5, A-1030 Vienna, Austria

Keywords: Aneuploidy; complex karyotypes; drug resistance; tubulin expression;

Abstract

Aneuploidy is one of the most common adaptive mechanisms to environmental selection in cells, yet its advantages over other genomic alterations remain unclear. We used budding yeast to determine if beneficial aneuploidy phenotypes are driven primarily by combinatorial expression changes of multiple genes. To determine the impact of complex aneuploidy on cellular fitness and drug resistance, we generated yeast collections with nearly every combination of two chromosome gains or losses. Additionally, we genetically dissected aneuploid chromosomes using partial deletions to identify chromosomal regions contributing to aneuploidy-driven drug resistance. Strong resistance phenotypes consistently came from combinations of chromosomes or chromosomal regions, with up to five regions on a chromosome contributing to resistance to a single drug. Gene ontology terms had limited predictive power in identifying the genes contributing to resistance phenotypes, as the combinatorial effects from different aneuploid regions act through multiple resistance pathways. The strongest phenotypes came from synergistic effects between copy number changes of different chromosomes or chromosomal regions, demonstrating how subtle gene expression changes of many genes combine to greatly impact cell survival.

Introduction

The accurate distribution of chromosomes during cell division is essential for maintaining genomic integrity across all eukaryotic species. Chromosome segregation errors result in alterations in chromosome copy number, which are termed aneuploidy. Aneuploidy is one of the most common genetic alterations in cancer, occurring in ~90% of solid tumors and ~70% of hematopoietic cancers (Weaver and Cleveland 2006). These high frequencies of aneuploidy are not only inherent to cancer cells but are also commonly observed within other vegetatively growing cell populations including industrial strains of baker's yeast and drug resistant strains of the human pathogens *Candida albicans* and *Leishmania* (Selmecki et al. 2006; Ubeda et al. 2008; Peter et al. 2018; Loegler et al. 2024).

Despite the frequency of aneuploidy in mitotically growing cell populations, it is generally associated with reduced cellular fitness. Engineering aneuploidy in cells invariably decreases growth rates under optimal conditions (Torres et al. 2007; Stingele et al. 2012; Hintzen et al. 2022; Zerbib et al. 2024). This decrease in fitness results from changes in gene copy numbers of many transcribed genes on the aneuploid chromosome (Bonney et al. 2015). For both yeast and human cells, gene expression levels of lab engineered aneuploidies scale with gene copy numbers for most of the aneuploid genes (Torres et al. 2007; Stingele et al. 2012; Wangsa et al. 2019; Rojas et al. 2024; Muenzner et al. 2024). These small changes in expression levels for hundreds of genes cause a wide range of deleterious phenotypes. Such general aneuploidy-associated phenotypes include increased proteotoxic stress, cell cycle delays, and increased DNA damage (Torres et al. 2007; Sheltzer et al. 2011; Thorburn et al. 2013; Oromendia et al. 2012; Beach et al. 2017; Li and Zhu 2022; Williams et al. 2008; Passerini et al. 2016; Santaguida et al. 2015; Garribba et al. 2023).

Yet, aneuploidy is one of the most frequently observed mechanisms of cellular adaptation. The first direct link between adaptive drug resistance and aneuploidy was established in the human pathogen *Candida albicans* (Selmecki et al. 2006). Adaptation via aneuploidy has since been more extensively observed in the budding yeast *Saccharomyces cerevisiae,* where specific aneuploidies consistently emerge

during experimental evolution screens under adverse environmental conditions
(Gresham et al. 2008; Yona et al. 2012; Chen et al. 2012; Voordeckers et al. 2015;
Zhang et al. 2016; Beaupere et al. 2018; Morard et al. 2019). Moreover, inducing
aneuploidy via random chromosome missegregation in both yeast and human cells
promotes resistance to a large variety of drugs (Pavelka et al. 2010; Chen et al. 2012;
Lukow et al. 2021; Ippolito et al. 2021). Aneuploidy can also act as a suppressor of
mutations that decrease cellular fitness (Rancati et al. 2008; Liu et al. 2015; Ryu et al.
2016; Ravichandran et al. 2018; Macaluso et al. 2025). In cancer, karyotype patterns
have predictive power for drug resistance, tumor progression and patient outcome
(Shukla et al. 2020).

The potential for aneuploidy as an adaptation mechanism to a large variety of selective
environments can partially explain the high prevalence of aneuploidy following long-
term vegetative growth, such as in cancer or in yeast used for alcohol production
(Peter et al. 2018). However, it still fails to explain why cells adapt through aneuploidy
over other types of mutations that don't share its inherent downsides. Different models
for the specific advantage of adaptation through aneuploidy have been proposed.
Aneuploidy has been suggested to be a rapid, but transient solution to sudden
environmental changes. In *S. cerevisiae* subjected to acute heat stress, chromosome
3 is gained early in the adaptation process but then lost again over time (Yona et al.
2012). Similarly, yeast adapting to high rates of chromosomal instability (CIN) adapted
first through the accumulation of specific aneuploidies, followed later by the
accumulation of suppressor point mutations (Clarke et al. 2023). However, there are
also many examples of specific aneuploidies that are maintained throughout long-term
adaptation, highlighting that speed is likely not the only advantage of adaptation via
aneuploidy (Selmecki et al. 2009; Ravichandran et al. 2018; Adell et al. 2023; Clarke
et al. 2023; Zhou et al. 2024; Watson et al. 2024).

An additional explanation for the advantages of aneuploidy as a mechanism of
adaptation comes from its ability to change gene expression levels of several
beneficial genes simultaneously. In yeast adaptation screens, only a handful of
examples could attribute the selective advantage of aneuploidy to changes in
expression levels of one or two specific genes. Even in these cases, they rarely
recapitulated the strength of the aneuploidy phenotype (Selmecki et al. 2009; Chen et

al. 2012; Ryu et al. 2016; Linder et al. 2017; Beaupere et al. 2018; Ravichandran et
al. 2018; Barney et al. 2021; Pavani et al. 2021; Zhou et al. 2024). In cancer,
overexpression of individual genes also fails to fully recapitulate the phenotypes of
aneuploid chromosomes (Trakala et al. 2021; Su et al. 2021; Girish et al. 2023). These
findings suggest that the challenges of identifying driver genes on aneuploid
chromosomes could be amplified by the interplay between multiple genes. It is
currently unclear how many genes on aneuploid chromosomes typically contribute to
the resistance to selective pressures.

Analyses of aneuploidy patterns in cancer genomes as well as cellular adaptation
experiments have shown strong correlations between specific chromosome or
chromosome arm combinations, indicating karyotype selection is influenced by
genetic interactions between aneuploid chromosomes (Ozery-Flato et al. 2011;
Ravichandran et al. 2018; Shukla et al. 2020; Adell et al. 2023). One such interaction
has been observed in budding yeast, where gaining an additional copy of chromosome
6 is lethal, but can be rescued with the additional gain of chromosome 13 (Torres et
al. 2007). Whether such genetic interactions between chromosome are widespread is
currently unknown. An alternative explanation for karyotype pattern formation is that
selective pressure could enrich for sets of beneficial aneuploidies, either
independently or through selection-specific chromosomal genetic interactions.

In this study, we generated a large yeast collection with nearly every possible
combination of two chromosome gains or losses to determine the frequency and
strength of genetic interactions between aneuploid chromosomes and how
combinatorial aneuploidy phenotypes impact drug resistance and sensitivity. We found
that genetic interactions between chromosomes are rare in the absence of treatment
but become much more frequent under strong selective pressure. In addition, the
strongest resistance phenotypes consistently came from combinations of aneuploid
chromosomes, demonstrating the advantage of copy number changes in multiple
genes. We genetically dissected resistant aneuploid chromosomes using partial arm
deletions to determine the number of regions that contribute to aneuploidy-driven drug
resistance on a single chromosome. We found that resistance conferred by single
disomies nearly always comes from multiple chromosomal regions. While the majority

of phenotypes are additive, the strongest resistance phenotypes of aneuploidies came

from synergistic effects between different regions or chromosomes.

Results

Generation of yeast strain libraries with combinations of chromosome gains and losses

We first determined how aneuploidy of different regions of the genome combine to

impact cellular fitness and drug response by generating two collections of yeast with

combinations of chromosome gains or losses. The collections consist of nearly all

possible single and double chromosome gains and losses (Fig 1A). To engineer

chromosome gain or loss, chromosome-specific missegregation was first induced via

a strong galactose-inducible centromere-proximal promotor, followed by selection for

the specific aneuploidy (Fig S1A). We selected for chromosome gains via

recombination events that restore the function of an auxotrophic marker. Chromosome

loss was generated by selecting against the *URA3* gene using 5-fluoroorotic acid (5-

FOA) (Fig S1A). The chromosome gain collection includes most of the possible single

(14/16) and double disomic (90/136) strains in haploid cells; each was confirmed by

qPCR (Table S1). The chromosome loss collection contains every possible single or

double monosomy in diploid cells (Fig S1B). Since chromosome loss cannot be

continuously selected for, this collection is inducible to generate newly aneuploid cells

prior to each experiment. These libraries currently constitute the largest collections of

engineered complex aneuploid cells.

We first evaluated the libraries by measuring the growth rates of the engineered

aneuploidy strains under optimal growth conditions. The high-throughput growth rate

measurements of single disomies and monosomies strongly correlate with those

previously observed (Beach et al. 2017) (Fig S1C). The previously published

measurements were performed in a strain background containing a mutation in the

SSD1 gene, which has been reported to sensitize cells to aneuploidy phenotypes

(Hose et al. 2020). However, we observe similar results in a different, wildtype *SSD1*

background (S288c), indicating that strain background does not substantially alter the

relative growth rate of aneuploid yeast. The ability to reproduce the previously

published correlations between gene number and phenotypic strength and the
differences between chromosome gain and loss attest to the quality of these strain
collections.

We next determined the impact of combining aneuploid chromosomes on cellular
fitness. Growth of complex aneuploidies negatively correlated with the number of open
reading frames (ORFs) on the aneuploid chromosomes for both chromosome gain
and loss (Fig 1B). While most of the single and double monosomies are lethal, nearly
all of the double disomies are viable, with the sole exception of those involving
chromosome 6 (Fig 1B and Fig S1B). The strongest outlier is the combination of
chromosomes 6 and 13 (Fig 1C). Chromosome 6 disomy is lethal on its own due to
the presence of an extra copy of the β-tubulin *TUB2* gene (Anders et al. 2009).
However, chromosome 6 disomy becomes viable with the additional gain of
chromosome 13. This is consistent with previous observations where selection of
chromosome 6 gain resulted in a strain that spontaneously gained the additional
disomies of chromosomes 1 and 13 (Torres et al. 2007). Chromosome 13 contains the
genes for α-tubulin, and overexpression of α-tubulin rescues the lethality of
chromosome 6 disomy (Anders et al. 2009). For the chromosome loss collection, one
of the strongest genetic interactions is again between chromosomes 6 and 13
(Fig. 1D). Chromosome 13 monosomy is lethal on its own, but the additional loss of
chromosome 6 rescues viability (Fig 1D). The heterozygous deletion of *TUB2* is
sufficient to rescue chromosome 13 monosomy, indicating that excess β-tubulin
relative to α-tubulin is the genetic basis behind these strong chromosome copy
number interactions (Fig 1D).

To identify any additional genetic interaction between chromosome gains, we
calculated growth predictions for all chromosome combinations based on the product
of the growth of the individual single aneuploidies. In general, growth of the double
disomies correlates well ($r^2 = 0.64$) with the calculated growth predictions (Fig 1C). To
measure genetic interaction between aneuploid chromosomes, we calculated an
epistasis score by subtracting predicted growth from the measured growth of the
chromosome combinations. The epistasis scores strongly adhered to a normal
distribution (Kolmogorov-Smirnov test), indicating that most combinations do not

deviate substantially from the expected growth rate (Fig S1D). Among the 90

chromosome combinations analyzed, we identified only 3 positive (z-score > 2) and 2

negative (z-score < -2) chromosome copy number interactions (Fig S1D). We

conclude that most combinations of aneuploid chromosomes show additive

phenotypes and that synergistic interactions are rare in the absence of selective

pressure.

The strongest drug resistance phenotypes come from strains with multiple

aneuploid chromosomes

We next used the chromosome gain library to determine if combinatorial interactions

between aneuploid chromosomes contribute to improved growth in stress conditions.

We measured growth in the presence of four drugs that induce cellular stresses

associated with aneuploidy: hygromycin B (ribosome inhibitor), rapamycin (TOR

pathway inhibitor), methyl methanesulfonate (MMS, DNA damage inducer) and

benomyl (microtubule depolymerizer) (Fig 2A). Although resistance mechanisms for

these drugs have been studied extensively, we aimed to identify how drug resistance

could be obtained by combining copy number changes of multiple regions in the

genome. Drug concentrations were titrated to reduce wild-type growth by ~80%.

Growth was unaffected by the solvent used to resuspend the drugs (Fig S1E).

Aneuploidy substantially altered the cellular response to hygromycin, rapamycin, and

benomyl, as growth rates no longer scaled with number of genes on aneuploid

chromosomes (Fig 2B). By contrast, MMS treatment showed fewer aneuploidy-

associated phenotypes. The percentage of strains displaying substantial relative

resistance (normalized growth in the presence/absence of the drug > 2) across the

entire library varies greatly between the different drugs ranging from 59% for benomyl

to 9% for MMS (Fig S2B and C). We conclude that the capacity for aneuploidy to

provide resistance is drug dependent.

In addition to drug resistance, we also observed frequent drug sensitivity. This was

most pronounced for hygromycin, where over 78% of strains showed a greater than

two-fold relative decrease in growth in the presence of the drug (Fig S2B). Additionally,

we measured sensitivity to two stress conditions other than drugs: 39°C for heat stress

and 18°C for cold stress. Neither of these conditions greatly influenced cell growth in

the euploid strain. However, heat stress led to a two-fold decrease in growth for 66%

of the disomic strains, which scaled with the degree of aneuploidy (Fig S2A-C). These

results suggest that aneuploid yeast are especially sensitive to hygromycin and heat

stress, which implicate translation inhibition and protein misfolding as potential

vulnerabilities in aneuploid cells. This would agree with previous reports that aneuploid

strains frequently exhibit decreased ribosome levels and proteotoxic stress

(Oromendia et al. 2012; Terhorst et al. 2020).

We next wanted to determine if combinations of aneuploid chromosomes show

stronger phenotypes than individual chromosomes. For hygromycin, rapamycin and

benomyl, specific chromosome combinations had the highest levels of drug

resistance, despite their negative impact on fitness in the absence of the drugs

(Fig 2C). Notably, absolute resistance to hygromycin was observed only for

chromosome combinations, as none of the single disomies grew better than the

euploid strain in the presence of hygromycin. We measured genetic interactions

between chromosomes in the same manner as for the untreated cells by comparing

the growth of the chromosome combinations to the predicted growth based on the

single disomies. For all four drugs, the epistasis scores no longer fit a normal

distribution, suggesting more frequent deviations from the predicted growth (Fig. S2E).

Indeed, after using the same threshold for epistasis scores as we applied to the

untreated condition, there were substantially more positive and negative genetic

interactions following drug treatment as compared to untreated cells (Fig 2D). Overall,

we conclude that the strongest drug resistance aneuploidy phenotypes typically come

from the combined effects of multiple genes on different chromosomes and that

selective pressure increases the frequency of chromosome copy number interactions.

Benomyl resistance correlates with α-Tubulin expression in aneuploid strains.

Surprisingly, a majority (59%) of disomic chromosome combinations were resistant to

the microtubule depolymerizing factor benomyl (Fig S2B and C). This high percentage

results from 6 of the 14 tested single disomic chromosomes conferring substantial

resistance to benomyl (Fig 2C and 3A). Although it has previously been reported that

many disomies lead to benomyl sensitivity (Sheltzer et al. 2011; Torres et al. 2007),

the resistance appears to outweigh the sensitivity for most chromosome combinations.

Benomyl resistance is therefore not a "general" aneuploidy phenotype but is instead

karyotype specific, despite being observed in the majority of double disomic strains.

Aneuploidy frequently leading to resistance to microtubule inhibitors was not specific

to benomyl, as four of the same aneuploid chromosomes provided resistance to

another microtubule depolymerizing drug, nocodazole (Fig S3A). We first focused on

chromosome 13, as it contains both α-tubulin genes (*TUB1* and *TUB3*) and α-tubulin

overexpression has been shown to result in a mild increase in benomyl resistance

(Schatz et al. 1986). Heterozygous deletion of *TUB3* in the chromosome 13 disomic

strain eliminated the resistance phenotype, demonstrating the contribution of this gene

to the resistance (Fig S3B). To determine if α-tubulin protein abundance contributes to

benomyl resistance in the other aneuploid strains, we measured the expression levels

in all 14 disomic strains via quantitative western blots. We found that multiple single

disomies alter α-tubulin expression levels (Fig 3B). The measured α-tubulin

expression levels correlated well with those measured by proteomics of aneuploid

yeast strains, demonstrating that these differences occur independently of the

measurement method and strain background (Fig S3C and D; (Muenzner et al. 2024;

Dephoure et al. 2014)). Elevated α-tubulin expression levels strongly correlated with

the degree of benomyl resistance for the single disomies, suggesting that this is a

major mechanism of resistance (Fig 3C and Fig S3E). Intriguingly, only chromosome

12 disomy is extremely sensitive to benomyl and is also the only disomy that shows a

substantial decrease in α-tubulin expression. The correlation with benomyl resistance

was much weaker for β-tubulin expression (Fig 3C, Fig S3F and G). These results

demonstrate that there are multiple genes across the genome that can alter tubulin

expression levels when present at an extra copy, and that these mild expression

changes can greatly alter the response to microtubule inhibitors.

To identify candidate resistance genes on the aneuploidy chromosomes, we started

by looking at gene ontology (GO) terms associated with the molecular target (tubulin),

associated phenotype (CIN), and drug sensitivity screens of the yeast deletion

collection. None of these terms showed a significant enrichment on the resistant

aneuploid chromosomes (Fig S4A). Furthermore, there were between 64 and 195

candidate genes per chromosome identified this way (Fig S4B). To further refine the

list of candidate genes, we focused on genes with substantial increases in RNA and

protein levels on the aneuploid chromosomes (Torres et al. 2007; and Dephoure et al.

2014). Applying an overexpression threshold of 1.5-fold for candidate proteins only

reduced the list by 32%. We therefore sought to identify chromosomal regions

responsible for drug resistance to decrease the number of candidate resistance

genes.

Identification of chromosome arms necessary for aneuploidy-based drug

resistance.

We next aimed to determine which regions of particular chromosomes provide

resistance when gained by a single copy. We focused primarily on benomyl resistance,

since six individual disomies led to strong resistance to the drug. This provided

multiple examples to systematically examine how many regions/genes typically

contribute to positive aneuploidy phenotypes for single aneuploid chromosomes. One

possibility is that a single strong resistance gene on each chromosome could be

sufficient to give a substantial fitness advantage under harsh drug conditions (single

gene model). Another possibility is that multiple mildly beneficial genes on single

chromosomes may act together to create strong resistance phenotypes (combinatorial

model) (Fig 4A).

To determine which regions of chromosomes have negative, positive, or neutral effects

on growth with and without benomyl, we implemented a method that deletes part of a

chromosome from the end and replaces it with a selection marker and an artificial

telomere (Linder et al., 2017) (Fig 4B). We first generated single arm disomies of all

of the benomyl resistant single disomic chromosomes. The deletion of one of the two

copies of the chromosome arms was confirmed by qPCR. As with whole chromosome

disomies, arm-level disomies have decreased growth rates proportional to the number

of aneuploid ORFs under standard conditions (Fig 4C). Benomyl resistance was

frequently associated with a single arm. For chromosomes 2, 4, 10, and 14, disomy of

the larger arm was sufficient to fully recapitulate the benomyl resistance, whereas

chromosome 13 only showed resistance with disomy of the shorter arm (Fig 4D). Both

α-tubulin genes (*TUB1 and TUB3)* are on the short (left) arm of chromosome 13.

Intriguingly, neither of the chromosome 5 arm disomies recapitulated the benomyl

resistance of the whole chromosome disomy. Chromosome 5 left arm disomy shows

no resistance, while the right arm disomy is only mildly resistant. Therefore, the

benomyl resistance for chromosome 5 results from at least one gene on each of the

two chromosome arms (Fig 4D). Overall, the correlation between growth with and
without benomyl for the arm-level disomies was not significant, demonstrating that
benomyl resistance is not the result of slow growth or a general aneuploidy phenotype
(Fig S5A). Instead, resistance likely results from the increased expression of one or
more specific genes on the aneuploid chromosome arms.

Next, we used our arm level aneuploidies to analyze resistance or sensitivity to the
three other drugs tested: hygromycin, MMS and rapamycin (Fig S6B). Similar to
benomyl, resistance to MMS, rapamycin and hygromycin was primarily associated
with single chromosome arms (3/3). By contrast, arm-level aneuploidies generally did
not fully recapitulate increased drug sensitivity phenotypes (Fig 4E and Fig S5B and
C). Instead, both arms often showed partial sensitivity. This is also the case for the
only chromosome that showed strong sensitivity to benomyl, chromosome 12 (Fig 4D).
These results suggest that aneuploidy-based drug sensitivity results from the
combined effects of more genes than drug resistance. Surprisingly, in some examples,
drug sensitivity from one chromosome arm was compensated for by the resistance
from the second chromosome arm (2/8) (Fig S5B and C).

Most aneuploidy-based drug resistance phenotypes come from the combined effect of multiple genes

Since most of the resistance phenotypes were associated with the larger arms, we
wanted to determine how many regions/genes on these arms contribute to benomyl
resistance. We therefore made multiple shorter deletions to the benomyl resistant
disomies to identify regions of drug resistance. For example, seven additional cuts
were made on the right arm of chromosome 4 to identify regions of interest. As the
size of the chromosome decreased, growth under standard conditions gradually
increased (Fig 5A). This further demonstrates that the negative effects on fitness for
aneuploid strains result from many genes across the length of a chromosome (Bonney
et al. 2015). Conversely, benomyl resistance decreased in a stepwise manner
following the deletion of three independent regions on the right arm (Fig 5A). For 5 of
the 6 resistant chromosomes, we identified at least two regions that contributed to
resistance, with the most being 5 regions for chromosome 5 (Fig 5C and Fig. S6A).
Interestingly, three of the benomyl resistant chromosomes also had regions that

increased drug sensitivity (Fig 5A and C and Fig S6A). The strength of drug resistance

is therefore a balance between regions that confer resistance and sensitivity.

To determine if the regions that lead to benomyl resistance also increase α-tubulin

protein levels, we examined the segmental deletions of chromosome 5 in the disomic

strain. Despite both arms of chromosome 5 contributing to benomyl resistance, only

the right arm was necessary for the increased α-tubulin levels. Removal of a single

region on the right arm decreased the α-tubulin levels back to euploid levels (Fig S5B).

The identified 60kb region also contributes to benomyl resistance. This region does

not contain any genes with an associated tubulin/microtubule gene ontology term.

None of the other 4 resistance regions contributing to benomyl resistance impacted α-

tubulin protein levels, indicating that these regions confer benomyl resistance through

alternative mechanisms.

We next aimed to identify the resistance genes within the identified regions. The

regions each contain between ~30 and 100 genes. We identified candidate genes by

looking for: sensitivity to benomyl in high-throughput screens of the yeast knockout

collection, essential genes, and genes that have GO Terms associated with CIN and

tubulin/microtubules (Fig S4B). Candidate genes were tested via heterozygous

deletions in the disomic strains (Fig S7A). For the two regions on chromosome 4

associated with the strongest benomyl resistance, we identified single gene deletions

that decreased resistance to a similar extent as deleting the entire chromosomal

region (Fig 5B and C). In total, we identified 9 genes that decrease the resistance of

aneuploid chromosomes when deleted heterozygously (Fig S7A and S8). In addition

to *TUB3* on chromosome 13, we identified two additional genes associated with tubulin

(*GIM4*-chr.5, *PFD1*-chr.10). Both of these genes are members of the prefoldin

complex, which is a chaperone that promotes the proper folding and assembly of α-

and β-tubulin. One resistance gene is involved in microtubule stabilization (*BIM1*-

chr.5). Interestingly, two of the resistance genes have recently been implicated in the

spindle position checkpoint (*SWR1*-chr.4, *MCK1*-chr.14) (Rathi et al. 2022; Caydasi et

al. 2023). Another resistance gene, *TOF1*-chr.14, is primarily associated with

protection of the DNA replication fork. However, it shows a synthetic lethal interaction

with *BIM1*, indicating an additional role in chromosome segregation (Tong et al. 2004;

Pan et al. 2004). Overall, we have identified many genes across the genome with a

variety of different functions that contribute to aneuploidy-based benomyl resistance.

Previous studies have only identified two or three genes responsible for resistance

phenotypes for individual perturbations, prohibiting any significant analysis of the

characteristics of genes that can contribute to aneuploidy phenotypes either in yeast

(rev. in Tsai and Nelliat 2019) or human cells (Adell et al. 2023; Ippolito et al. 2021).

We assessed whether the 9 genes identified shared common features using GO

enrichment, RNA expression levels and protein abundance. Intriguingly, GO

overrepresentation analysis of the nine identified benomyl resistance genes did not

reveal any significant enrichment terms as measured by adjusted p-values (Fig S7D).

In addition, the highest ranked hit, "microtubule cytoskeleton organization", only

contained 3 out of 9 genes. This highlights the diversity of gene functions that can

combine to contribute to aneuploidy-based drug resistance.

Different genes can have substantial differences in the degree to which their

expression increases when an additional copy is gained on an aneuploid chromosome

(Stingele et al. 2012; Dephoure et al. 2014; Muenzner et al. 2024; Hwang et al. 2021;

Sousa et al. 2019). We therefore hypothesized that genes that contribute to beneficial

aneuploidy phenotypes would be those that have their expression levels more

substantially altered by aneuploidy. Analysis of the increase in RNA expression levels

for genes on aneuploid chromosomes showed a small, yet significant increase for the

9 benomyl resistance genes when compared to the rest of the genes on these

chromosomes (Torres et al. 2007; Fig S7E) However, this may reflect a bias in the

selection of candidate genes, as there was no significant difference between the tested

genes that either did or did not contribute to benomyl resistance (Fig S7F). For the 7

benomyl resistance genes where increased protein levels on aneuploid chromosomes

have been measured, there were no significant differences in expression levels (Fig

S7E and F). We conclude that genes affecting aneuploidy phenotypes are not

enriched for genes with more substantial increases in expression on aneuploid

chromosomes.

In addition to the disomic chromosomes that lead to benomyl resistance, we also

identified chromosomal regions that lead to rapamycin resistance on chromosome 10.

Strains with disomy of chromosome 10 have previously been shown to be rapamycin
resistant (Torres et al. 2007). This was speculated to be due to the presence of the
TOR1 gene. Heterozygous deletion of *TOR1* in a chromosome 10 disomic strain only
partially decreased rapamycin resistance, indicating that there are more genes on
chromosome 10 that contribute to the phenotype (Fig S7B). Furthermore, both
chromosome arms contribute to rapamycin resistance (Fig S7C). We conclude that
the combined overexpression of multiple genes leading to strong phenotypes is a
common feature of aneuploidy.

The strongest aneuploidy phenotypes result from synergistic effects

Chromosome 5 disomy has the strongest benomyl resistance phenotype and is the
only chromosome where both arms contribute to the phenotype (Fig 4D). Additionally,
the degree of resistance for the whole chromosome was substantially higher than
either of the two individual arm-level disomies, suggesting a synergistic interaction
between the two arms. To look at this more systematically, we predicted the strength
of resistance for whole chromosome aneuploidy based on the resistance of the
individual arm-level aneuploidies (Fig 6A). Although the other single disomies
displayed resistance in line with the predictions, benomyl resistance for chromosome
5 was substantially higher than predicted. This demonstrates a strong synergistic
interaction between the two arms.

We next wanted to determine if the strongest phenotypes for double disomic strains
also result from synergistic interactions. We focused on chromosome combinations
where the single disomies on their own were benomyl resistant. We initially observed
that the combined disomies of chromosomes 2 & 4 and 2 & 13 are exceptionally
strong, showing robust resistance even at higher benomyl concentrations (Fig 6B).
These combinations are as resistant as a deletion mutation known to be highly
resistant to benomyl (Schibler et al. 2016), despite strong growth defects for aneuploid
strains in the absence of the drug (Fig 6C). These high levels of resistance do not
result from further increases in α-tubulin expression levels (Fig S9A). Looking at
combinations with chromosome 2 specifically, we observed a wide range of resistance
phenotypes, ranging from strong positive to strong negative interactions (Fig 6B and
Fig S9B). Strikingly, a strong negative genetic interaction between chromosomes 2

and 14 results in the near-complete loss of the benomyl resistance (Fig 6B and D).

Almost all combinations with chromosome 14 grow worse than the single disomies

(Fig S9C), indicating a general negative effect of chromosome 14 combinations and

benomyl resistance. We also observed a positive synergistic interaction between

chromosomes 3 and 12 for rapamycin resistance, demonstrating that such strong

genetic interactions are not benomyl-specific (Fig S9D). We conclude that most

aneuploidies display additive effects between genes, regions, and chromosomes.

However, the strongest beneficial growth phenotypes often come from synergistic

interactions.

Discussion

In this study, we generated a library of engineered single and double aneuploid yeast

strains to systematically determine the genetic drivers of aneuploidy-based drug

resistance. In addition, we used a comprehensive approach to identify regions/genes

on aneuploid chromosomes that contribute to aneuploidy phenotypes. This approach

has numerous advantages, allowing for the identification of: i) multiple regions on a

chromosome that contribute to a phenotype, ii) both resistant and sensitive regions on

a single chromosome, iii) regions that act synergistically, where each region on its own

would not have a strong phenotype. Together, these strategies provide a means to

comprehensively identify the genetic determinants of positive and negative selection

for aneuploid chromosomes either alone or in combination.

When measuring the resistance of the chromosome gain collection to four different

inhibitors, the strongest resistance phenotypes were consistently identified for

chromosome combinations. This demonstrates that altered copy numbers of multiple

genes combine to create stronger beneficial aneuploidy phenotypes. These additional

benefits are observed despite the often substantial negative growth phenotypes for

the double aneuploid strains under optimal growth conditions. Therefore, the benefits

of combining multiple aneuploid chromosomes generally outweigh the downsides

under selective conditions. Furthermore, we saw these positive combinatorial effects

for 3 of the 4 drug conditions, suggesting that it is a general property of aneuploidy.

This combinatorial behavior of aneuploid chromosomes could explain the frequent

occurrence of complex aneuploidy in both adapted yeast populations and cancer

karyotypes (Mitelman Database of Chromosome Aberrations and Gene Fusions in

Cancer, 2025. https://mitelmandatabase.isb-cgc.org, (Rancati et al. 2008;

Ravichandran et al. 2018; Peter et al. 2018; Klockner and Campbell 2024).

In addition to identifying strong phenotypes for combinations of aneuploid

chromosomes, we discovered multiple regions within individual chromosomes that

contribute to drug resistance phenotypes. We identified up to 5 regions on a single

chromosome that contribute to drug resistance, emphasizing that multiple genes are

typically responsible for strong aneuploidy phenotypes. This combinatorial action of

many genes suggests an advantage for the selection of aneuploidy over other types

of genomic alterations. Changing the expression of genes with distinct and

complementary functions may provide proliferative benefits that could not be

accomplished through single mutations. These data provide substantial support for a

"combinatorial genes" explanation for the specific benefits of aneuploidy over other

genomic alterations.

In the absence of selection, we did not observed strong genetic interactions between

chromosome gains with the exception of the combination of chromosomes 6 and 13.

However, it is worth noting that even relatively weak interactions have the potential to

affect aneuploidy patterns (Ravichandran et al. 2018). By contrast, under selection,

the number of genetic interactions between chromosomes was substantially increased

for all 4 of the drugs tested. Furthermore, increasing the degree of selection via higher

concentrations of benomyl revealed strong positive synergistic interactions between

aneuploid chromosomes. We also observed a clear synergistic interaction between

the arms of chromosome 5. These results suggest another possible advantage for

adaptation through aneuploidy, wherein the altered expression levels of two

complementary genes would not be selected for individually due to a lack of

phenotype. However, two simultaneous changes to the genome through a single

chromosome missegregation event would enable their combined selection, leading to

a strong beneficial phenotype. These results, along with the recently discovered

synergistic interaction between chromosomes 3 and 6 in multi-azole tolerance of the

human pathogen *Candida albicans* (Zhou et al. 2024), demonstrate that positive

synergistic interactions are more heavily selected for under higher stress conditions.

We also find that combining beneficial aneuploidies can occasionally lead to weaker

than expected phenotypes, highlighting that negative genetic interactions also likely
impact complex karyotype formation. In cancers, positive and negative genetic
interactions between co-occurring aneuploidies have been suggested based on
correlations between chromosome arm frequencies in aneuploidy patterns (Ozery-
Flato et al. 2011; Ravichandran et al. 2018). Such co-occurrence patterns can be
highly predictive of disease progression and chemotherapy resistance in cancer
(Shukla et al. 2020). Our results suggest that these patterns are more likely to result
from specific selective pressures, rather than more general synergies or
incompatibilities between the gain or loss of specific chromosomes arms.

The main drivers of beneficial aneuploidy phenotypes are alterations in expression
levels resulting from gene copy number changes of specific resistance-driving genes
(Selmecki et al. 2009; Chen et al. 2012; Ryu et al. 2016; Linder et al. 2017; Beaupere
et al. 2018; Ravichandran et al. 2018; Barney et al. 2021; Pavani et al. 2021; Adell et
al. 2023; Zhou et al. 2024; Watson et al. 2024). We observed a significant correlation
between growth on benomyl and altered α-tubulin expression levels. In addition to
TUB3 on chromosome 13, two members of the prefoldin complex (*GIM4*-chr. 5, *PFD1*-
chr. 10) contribute to aneuploidy-based benomyl resistance. Intriguingly, chromosome
12 disomy decreases tubulin expression levels and is very sensitive to microtubule
destabilization via benomyl. This decrease in tubulin levels for chromosome 12 disomy
could also help explain how it rescues microtubule hyperstabilization (Macaluso et al.
2025). We conclude that tubulin expression levels are not tightly regulated in yeast.
Furthermore, even relatively small changes in α-tubulin expression levels can
substantially influence resistance and sensitivity to microtubule inhibitors. Moreover,
the genes contributing to benomyl resistance were not always among the highly
expressed genes in both transcriptomics and proteomics datasets, demonstrating that
even mild overexpression can lead to substantial phenotypes. These findings
underscore the importance of recognizing subtle (<2-fold) expression changes, as
they have the capacity for causing strong phenotypes.

Although increased α-tubulin expression appears to be a common mechanism of
aneuploidy-based benomyl resistance, we also find evidence for additional
mechanisms. First, not all of the genes that contributed to benomyl resistance have

known functions related to tubulin expression (Fig S7D). We identified two
components of the spindle position checkpoint (*MCK1*-chr. 14, *SWR1*-chr. 4),
implicating this checkpoint in mitigating the effects of microtubule-targeting drugs. We
also identified a drug exporter (*SNQ2*-chr. 4) and a microtubule stabilizer (*BIM1*-chr.
5). Importantly, the genes that we identified for resistant chromosome regions were
not always the most obvious candidates (Fig. S7). *TOF1*-chr. 14 and *SAP1*-chr. 5 have
not previously been associated with microtubule functions, but both contribute to
benomyl resistance on their respective aneuploid chromosomes. Second, not all of
the resistant disomic strains had increased α-tubulin expression (e.g. Dis14, Figure
3C). Third, not every region of a disomic chromosome that contributes to benomyl
resistance increases α-tubulin expression (Fig. S6B). Finally, we find that disomy
combinations with exceptionally high benomyl resistance do not increase α-tubulin
levels further than the single disomies (Fig. S9A). We conclude that different regions
on individual chromosomes can combine for beneficial aneuploidy phenotypes
through additive or synergistic effects from combining multiple resistance pathways.
Additionally, genes that we would have expected to contribute to benomyl resistance
based on previous knockout screens often did not have an effect when heterozygously
deleted from the disomic strains. The most prominent example being a drug efflux
pump located on chromosome 2, Flr1, which did not contribute to benomyl resistance
for that chromosome. Deletion of *FLR1* leads to strong benomyl sensitivity (Brôco et
al. 1999), yet an extra copy from the disomic chromosome did not affect resistance
(Fig S7A). Together, these results demonstrate that gene functions determined
primarily via loss-of-function mutants may often not correspond with the phenotypes
that emerge from increased gene expression due to copy number gain.

While mild overexpression of specific genes can confer drug resistance, the
overexpression of many genes across an entire chromosome greatly increases
aneuploidy-induced cellular stress (Torres et al. 2007; Sheltzer et al. 2011; Oromendia
et al. 2012; Thorburn et al. 2013; Beach et al. 2017; Li and Zhu 2022). This stress
presents potential targets to selectively eliminate aneuploid cells (Cohen-Sharir et al.
2021; Ippolito et al. 2024). However, the potential to leverage aneuploidy-based
sensitivities requires a deeper understanding of the underlying mechanisms driving
these vulnerabilities. 78% of the strains in the chromosome gain collection were
sensitive to hygromycin B and 66% were sensitive to heat stress (Fig. S2C). These

results provide additional support for decreased ribosome function and proteotoxic

stress as hallmarks of aneuploidy (Oromendia et al. 2012; Terhorst et al. 2020). By

contrast, inducing DNA damage via MMS did not result in large scale sensitivity,

suggesting aneuploidy-induced genome instability may be more chromosome specific.

Despite the general sensitivity to hygromycin, we identified a few strong examples of

aneuploidy-driven hygromycin resistance. Additionally, sensitivity of one chromosome

arm can be masked by resistance of the other arm (Fig. 5C, Fig. S4B, C). Together,

these results demonstrate that regions of the genome that contribute to sensitivity or

resistance can fully or partially counteract each other. Sensitivity and negative

selection have been predicted to be strong drivers of cancer karyotype evolution,

potentially even more so than positive selection (Jubran et al. 2024).

Given the high prevalence of aneuploidy in cancer, it is an attractive target for

identifying exploitable vulnerabilities. This requires a better understanding of the

positive and negative forces underlying aneuploid chromosome selection. This work

highlights additional challenges for identifying the genes responsible for aneuploidy

phenotypes, as they are likely to result from the combined effects of multiple genes.

Material and methods

Yeast strains and media

All yeast strains and plasmids used in this study are listed in Table S2, S3. All yeast

strains are derivatives of BY4741 and BY4742. All yeast strains were cultured in

complete synthetic media (CMS) lacking histidine, uracil, lysine, and leucine and

cultured at 30 °C, unless otherwise stated. For 5-FOA (Chempur, Karlsruhe, Germany,

220141- 70-8) selection, a concentration of 1 mg/mL was used. Growth assays were

always conducted in yeast extract/peptone containing 40 µg/ml adenine-HCl (YPA)

and 2% glucose (YPAD) in liquid cultures and plate-based assays. For high-throughput

drug resistance screens, 17.5 µg/ml benomyl (Sigma-Aldrich), 40 µg/ml hygromycin B

(Carl Roth), 0.02 % MMS (Sigma-Aldrich) or 5 nM rapamycin (Santa Cruz

Biotechnology) were added to YPAD agar plates. YPAD + DMSO agar plates were

used as controls for drugs dissolved in DMSO. Genetic manipulations such as

insertions and gene deletions were conducted as previously described (Longtine et al.

1998). Strains inducible for aneuploidy were generated using a conditional

centromere, as described in Anders et al. (2009). Single disomies and double disomies

were engineered as described in (Ravichandran et al. 2018). Each disomy was

generated independently with two different pairs of auxotrophic markers (*URA3* and

HIS3; *LYS2* and *LEU2*). For engineering monosomies, one copy of the endogenous

centromere was replaced by P_{GAL1}-*CEN3 URA3*. Flow cytometry of selected strains

was used to detect potential changes in ploidy and exclude strains that spontaneously

diploidized.

PCR-mediated partial chromosome deletions

PCR-mediated partial chromosome deletions were implemented as described in

Linder et al. (2017). To generate a deletion, a 500-800 bp fragment homologous to the

intended cut-site was combined with a resistance marker (*kanMX* or *hphNT1*) and a

telomeric seed sequence. The telomeric seed sequence (6x 5'CCCCAA3') was added

at the 3' end of the primer binding the resistance cassette. The homology sequence

was PCR amplified from genomic DNA with an added 30 bp homology sequence to

the resistance cassette. Fragments were combined using overlap extension PCR.

High-throughput growth measurements

For monosomies, chromosome missegregation was induced immediately before

plating. Strains were grown for 18h in 2 ml deep-well plates in 200 µl YPAD at 30 °C

(shaking). 4 µl were transferred to 150 µl of YPA plus 1% raffinose (YPAR) and grown

to log phase (~4h). 50 µl of YPA medium containing 4% galactose were added and

grown for 3h to induce chromosome missegregation. Samples were plated on 5-FOA

agar plates using a ROTOR HDA robot for high-throughput plating (Singer

Instruments, Roadwater, UK). Images were taken after 24, 48 and 72h. The

chromosome gain collection was grown in 2 ml deep-well plates in 200 µl CSM lacking

histidine, uracil, leucine and lysine at 30 °C. After 18h, cultures were diluted 1:10 in

1 × PBS and plated with the ROTOR HDA onto condition specific YPAD plates. Strains

were grown at 30 °C (if not indicated otherwise) and images were taken at different

time points from 24 to 72 h. Timepoints were selected for quantification based on the

highest resolution of growth differences between strains.

Doubling time measurements

Overnight cultures were grown in CSM lacking histidine, uracil, leucine and lysine,

diluted in YPAD and grown for 4h to mid log-phase. Cultures were diluted to an

optical density at 600 nm of 0.02 in 200 µl YPAD in 96-well plates. OD

measurements were taken every 15 min for 24 h using a Synergy S1 plate reader

(BioTek Instruments, Inc., Luzern, Switzerland).

Serial dilutions

Strains were grown over night in CSM lacking histidine, uracil, leucine and lysine.

Cultures were diluted in fresh media (OD ~ 0.05) and grown to log phase (OD

~0.45), normalized by OD, diluted in 1 x PBS (1 : 10) and plated on condition specific

agar plates.

High-throughput growth assay quantification and analysis

Images of yeast colonies on plates were quantified using ImageJ (National Institute of

Health). Integrated densities of each spot area and surrounding area were quantified.

Background subtraction was done for each spot area individually by subtracting the

surrounding intensity from the colony intensity (Supplemental data file). For comparing

growth of chromosome loss strains to chromosome gain strains, 10 representative

chromosome loss strains were spotted on YPAD after selection on 5-FOA plates and

measured for colony intensity. These values were then used to adjust the growth

measurements for the other chromosome loss strains. Growth rate prediction of

double disomies and double monosomies in high throughput experiments was

predicted by multiplying the growth of the two individual single disomies/monosomies.

Epistasis scores were calculated by subtracting predicted growth from measured

growth. This was done for each treatment individually. Determination of genetic

interactions between disomic chromosomes on high drug concentrations were

calculated from dilution series (Fig 6A and C) by adding the measurements of the

individual single disomies (DisA + DisB - 1). Growth rates greater/smaller than additive

prediction +/- 30% were classified as synergistic genetic interaction. The same method

was used to identify genetic interactions between chromosome arms.

qPCR

For DNA extraction, cells were taken from plates and lysed in 20 µl of 0.02 M NaOH

for 10 min at 100 °C. Lysates were centrifuged for 10 min on a mini-centrifuge (Thermo

Fisher Scientific) to pellet cellular debris. Each 20 µl reaction contained 10 µl of Luna

qPCR Mastermix (New England BioLabs Inc.), 1 µl of lysate and 140 nM of each

primer. Measurements were made on a RealPlex2 qPCR machine (Eppendorf).

Cycling conditions were: activation/denaturation for 5 min at 95 °C followed by

40 cycles of 15 sec at 95 °C, 1 min at 60 °C with a subsequent fluorescence

measurement. Melting curves were generated to determine primer specificity. *Ct*

values were determined using the automatic thresholding of the RealPlex2 software.

Chromosome copy numbers were calculated using a slightly modified ΔΔCt method

(Schmittgen 2001). The Ct values from duplicates were averaged and used to obtain

the ΔCt, which was raised to the negative power of 2 to give the fold change. The ratio

of fold change of the test strains to that of a wild-type strain was calculated to obtain

the chromosome copy numbers. Karyotyping primers are in Table S5.

Protein extraction and Western blotting

For protein extraction, saturated ON cultures were diluted to an OD of 0.02 into YPAD. Cells were grown to an OD of ~0.6, normalized by OD and harvested. Proteins were extracted by pelleting cells, resuspending them in 100 µl ddH$_2$O, adding 100 µl 0.2 M NaOH and incubating them at room temperature for 5 min. Cells were pelleted and resuspended in 1x sample buffer before incubating them at 95° C for 3 min. Samples were stored at -20°C. Membranes were stained with 0.1 % Ponceau S (Sigma Aldrich) in 5% acetic acid. Total protein amount was detected with the ChemiDocMP imaging system (Bio-Rad). Membranes were washed for 30 min in Tris-buffered saline with 0.01% Tween (TBS-T) with three buffer exchanges, blocked for 1 h in 5% BSA (Sigma-Aldrich) or 5% skim milk (Gerbu Biotechnik GmbH) in TBS-T, depending on the primary antibody (see below). Primary antibodies diluted in their respective blocking buffers were incubated overnight at 4°C on an orbital shaker. Membranes were washed for 30 min in TBS-T with three buffer exchanges before and after secondary antibody incubation. Horseradish peroxidase (HRP)-conjugated antimouse IgG secondary antibody (1:10,000; Cell Signaling Technology 7076) diluted in the respective blocking buffers was incubated for 1 h at room temperature. Chemiluminescence was developed using the Amersham ECL Prime detection reagent (Cytiva) and detected with the ChemiDocMP. Protein bands were quantified using ImageJ and normalized to the total protein amount quantified from the Ponceau S staining. The following primary antibodies were used: mouse anti-Pgk1 (in BSA and skim milk–TBST, respectively; 1:20.000; Thermo Fisher Scientific 22C5D8), anti-α-Tubulin (in skim milk–TBS; 1:20,000; mouse; Sigma-Aldrich T6074), anti-β-Tubulin (in BSA; 1:5.000; mouse; Developmental Studies Hybridoma Bank E7).

Flow Cytometry

Cultures were grown to log-phase, pelleted and resuspended in 300 µl H$_2$O. 700 µl of 100 % EtOH were added during vertexing, and cells were fixed for 1h at room temperature. Cells were pelleted and resuspended in 50 mM sodium citrate, sonicated to disperse cell clumps and treated with 250 µg/mL RNase A (Sigma-Aldrich, R6513) and 1 mg/mL Proteinase K (Sigma-Aldrich, P2308) overnight at 37°C. Cells were resuspended in 50 mM sodium citrate solution containing 1 µM SYTOX green (Thermo Fisher, 10768273). Samples were run on BD LSRFortessa Cell Analyzer equipped

with a 15-mW 488-nm laser. Maximum count peaks for fluorescence intensities were

calculated using the BD FACSDiva 8.0.1 software.

Gene ontology enrichment analysis

For GO enrichment tests, the GOrilla gene ontology tool (http://cbl-gorilla.cs.

technion.ac.il, accessed in September 2025) was used. The target gene list was

composed of all genes identified as resistance genes, and the background list was all

6,002 genes annotated in the *S. cerevisiae* genome. All lists of candidate genes were

assembled from the Saccharomyces Genome Database, based on specific phenotype

annotations. The list of Benomyl-related genes was based on the following

classifications after benomyl treatment: resistance to chemicals: decreased;

chromosome/plasmid maintenance: decreased; viability: decreased. 799 genes

contain at least one of these terms. The list of CIN-related genes was assembled using

the following GO terms: colony sectoring: increased; chromosome segregation:

abnormal; chromosome/plasmid maintenance: decreased rate; chromosome/plasmid

maintenance: abnormal; chromosome segregation (Clarke et al. 2023). 718 different

genes contain at least one of these terms. The list of tubulin/microtubule related genes

was based on: cytoplasmic microtubule organization; gammatubulin binding; microtubule

cytoskeleton organization; microtubule depolymerization; microtubule nucleation;

microtubule polymerization; microtubule based process; mitotic spindle assembly; mitotic

spindle elongation; tubulin binding; tubulin complex; tubulin complex assembly. 43 genes

contain at least one of these terms.

RNA expression and protein abundance analysis

Data sets from Torres et al. 2007, Dephoure et al 2014 and Muenzner et al. 2024 were

used to analyze RNA and protein levels. Due to the high protein sequence homology

between Tub1 and Tub3 (>90% identical), their protein levels where added together.

Expression data from Muenzner et al. was excluded from the analysis of benomyl

resistance driver genes, as only 4 out of 9 proteins were identified in their data set.

Statistical analysis

Statistical analysis was performed using GraphPad Prism software. Details for the

statistical tests used in a particular experiment are reported in the figure legends.

Competing interest statement

The authors declare no competing interests.

Acknowledgements

The authors thank the Campbell and Dammermann laboratories for helpful discussions and comments. We thank the Kraft lab and Kirk Anders for gifts of strains and plasmids and Alexander Dammermann and Sean Montgomery for careful reading of the manuscript. We acknowledge the service of the FACS Facility.

This work was supported by a Max Perutz Labs Fellowship to M.S.K and the Austrian Science Fund (FWF) grant F 8802-B to C.S.C.

Author Contributions

M.S.K conducted the majority of the experiments. M.S.K., S.F., M.C.R and C.H., generated the chromosome gain collection. C.H. and S.F. generated the chromosome loss collection. C.H. performed growth measurements and analysis of the chromosome loss collection. M.S.K. and C.S.C conceptualized the research, analyzed the data, and wrote the manuscript. All authors reviewed and approved the manuscript.

References

Adell MAY, Klockner TC, Höfler R, Wallner L, Schmid J, Markovic A, Martyniak A, Campbell CS. 2023. Adaptation to spindle assembly checkpoint inhibition through the selection of specific aneuploidies. *Genes Dev* **37**: 171–190.

Anders KR, Kudrna JR, Keller KE, Kinghorn BAA, Miller EM, Pauw D, Peck AT, Shellooe CE, Strong IJT. 2009. A strategy for constructing aneuploid yeast strains by transient nondisjunction of a target chromosome. *BMC Genet* **10**: 1–11.

Barney JB, Chandrashekarappa DG, Soncini SR, Schmidt MC. 2021. Drug resistance in diploid yeast is acquired through dominant alleles, haploinsufficiency, gene duplication and aneuploidy ed. A.M. Dudley. *PLOS Genet* **17**: e1009800.

Beach RR, Ricci-Tam C, Brennan CM, Moomau CA, Hsu P, Hua B, Silberman RE, Springer M, Amon A. 2017. Aneuploidy Causes Non-genetic Individuality. *Cell* **169**: 229-242.e21.

Beaupere C, Dinatto L, Wasko BM, Chen RB, VanValkenburg L, Kiflezghi MG, Lee MB, Promislow DEL, Dang W, Kaeberlein M, et al. 2018. Genetic screen identifies adaptive aneuploidy as a key mediator of ER stress resistance in yeast. *Proc Natl Acad Sci* **115**: 9586–9591.

Bonney ME, Moriya H, Amon A. 2015. Aneuploid proliferation defects in yeast are not driven by copy number changes of a few dosage-sensitive genes. *Genes Dev* **29**: 898–903.

Brôco N, Tenreiro S, Viegas CA, Sá-Correia I. 1999. FLR1 gene (ORF YBR008c) is required for benomyl and methotrexate resistance inSaccharomyces cerevisiae and its benomyl-induced expression is dependent on Pdr3 transcriptional regulator. *Yeast* **15**: 1595–1608.

Caydasi AK, Khmelinskii A, Darieva Z, Kurtulmus B, Knop M, Pereira G. 2023. SWR1 chromatin remodeling complex prevents mitotic slippage during spindle position checkpoint arrest ed. K. Bloom. *Mol Biol Cell* **34**: ar11.

Chen G, Bradford WD, Seidel CW, Li R. 2012. Hsp90 stress potentiates rapid cellular adaptation through induction of aneuploidy. *Nature* **482**: 246–250.

Clarke MN, Marsoner T, Adell MAY, Ravichandran MC, Campbell CS. 2023. Adaptation to high rates of chromosomal instability and aneuploidy through multiple pathways in budding yeast. *EMBO J* **42**: e111500.

Cohen-Sharir Y, McFarland JM, Abdusamad M, Marquis C, Bernhard SV, Kazachkova M, Tang H, Ippolito MR, Laue K, Zerbib J, et al. 2021. Aneuploidy renders cancer cells vulnerable to mitotic checkpoint inhibition. *Nature* **590**: 486–491.

Dephoure N, Hwang S, O'Sullivan C, Dodgson SE, Gygi SP, Amon A, Torres EM.
2014. Quantitative proteomic analysis reveals posttranslational responses to
aneuploidy in yeast. *eLife* **3**: e03023.

Garribba L, De Feudis G, Martis V, Galli M, Dumont M, Eliezer Y, Wardenaar R,
Ippolito MR, Iyer DR, Tijhuis AE, et al. 2023. Short-term molecular
consequences of chromosome mis-segregation for genome stability. *Nat*
Commun **14**: 1353.

Girish V, Lakhani AA, Thompson SL, Scaduto CM, Brown LM, Hagenson RA,
Sausville EL, Mendelson BE, Kandikuppa PK, Lukow DA, et al. 2023.
Oncogene-like addiction to aneuploidy in human cancers. *Science* **381**:
eadg4521.

Gresham D, Desai MM, Tucker CM, Jenq HT, Pai DA, Ward A, DeSevo CG, Botstein
D, Dunham MJ. 2008. The Repertoire and Dynamics of Evolutionary
Adaptations to Controlled Nutrient-Limited Environments in Yeast ed. M.
Snyder. *PLoS Genet* **4**: e1000303.

Hintzen DC, Soto M, Schubert M, Bakker B, Spierings DCJ, Szuhai K, Lansdorp PM,
Kluin RJC, Foijer F, Medema RH, et al. 2022. The impact of monosomies,
trisomies and segmental aneuploidies on chromosomal stability ed. D. Cimini.
PLOS ONE **17**: e0268579.

Hose J, Escalante LE, Clowers KJ, Dutcher HA, Robinson D, Bouriakov V, Coon JJ,
Shishkova E, Gasch AP. 2020. The genetic basis of aneuploidy tolerance in
wild yeast. *eLife* **9**.

Hwang S, Cavaliere P, Li R, Zhu LJ, Dephoure N, Torres EM. 2021. Consequences
of aneuploidy in human fibroblasts with trisomy 21. *Proc Natl Acad Sci* **118**:
e2014723118.

Ippolito MR, Martis V, Martin S, Tijhuis AE, Hong C, Wardenaar R, Dumont M, Zerbib
J, Spierings DCJ, Fachinetti D, et al. 2021. Gene copy-number changes and
chromosomal instability induced by aneuploidy confer resistance to
chemotherapy. *Dev Cell* **56**: 2440-2454.e6.

Ippolito MR, Zerbib J, Eliezer Y, Reuveni E, Viganò S, Taglietti S, Scorzoni S,
Gianotti S, Martin S, Muenzner J, et al. 2024. Increased RNA and protein
degradation is required for counteracting transcriptional burden and
proteotoxic stress in human aneuploid cells. *Cancer Discov* **14**.

Jubran J, Slutsky R, Rozenblum N, Rokach L, Ben-David U, Yeger-Lotem E. 2024.
Machine-learning analysis reveals an important role for negative selection in
shaping cancer aneuploidy landscapes. *Genome Biol* **25**: 95.

Kaboli S, Miyamoto T, Sunada K, Sasano Y, Sugiyama M, Harashima S. 2016.
Improved stress resistance and ethanol production by segmental
haploidization of the diploid genome in Saccharomyces cerevisiae. *J Biosci*
Bioeng **121**: 638–644.

Klockner TC, Campbell CS. 2024. Selection forces underlying aneuploidy patterns in cancer. *Mol Cell Oncol* **11**: 2369388.

Li R, Zhu J. 2022. Effects of aneuploidy on cell behaviour and function. *Nat Rev Mol Cell Biol* **23**: 250–265.

Linder RA, Greco JP, Seidl F, Matsui T, Ehrenreich IM. 2017. The Stress-Inducible Peroxidase *TSA2* Underlies a Conditionally Beneficial Chromosomal Duplication in *Saccharomyces cerevisiae*. *G3 GenesGenomesGenetics* **7**: 3177–3184.

Liu X, Wang X, Yang X, Liu S, Jiang L, Qu Y, Hu L, Ouyang Q, Tang C. 2015. Reliable cell cycle commitment in budding yeast is ensured by signal integration. *eLife* **4**: e03977.

Loegler V, Friedrich A, Schacherer J. 2024. Overview of the *Saccharomyces cerevisiae* population structure through the lens of 3,034 genomes ed. G. Brown. *G3 Genes Genomes Genet* jkae245.

Longtine MS, McKenzie A, Demarini DJ, Shah NG, Wach A, Brachat A, Philippsen P, Pringle JR. 1998. Additional modules for versatile and economical PCR-based gene deletion and modification in Saccharomyces cerevisiae. *Yeast Chichester Engl* **14**: 953–61.

Lukow DA, Sausville EL, Suri P, Chunduri NK, Wieland A, Leu J, Smith JC, Girish V, Kumar AA, Kendall J, et al. 2021. Chromosomal instability accelerates the evolution of resistance to anti-cancer therapies. *Dev Cell* **56**: 2427-2439.e4.

Macaluso F, Bos T, Chiroli E, Bonaiuti P, Apuan JC, Gross F, Pompei S, Rice LM, Ciliberto A. 2025. Evolutionary adaptation to hyperstable microtubules selectively targets tubulins and is empowered by the spindle assembly checkpoint. *Cell Rep* **44**: 115323.

Morard M, Macías LG, Adam AC, Lairón-Peris M, Pérez-Torrado R, Toft C, Barrio E. 2019. Aneuploidy and Ethanol Tolerance in Saccharomyces cerevisiae. *Front Genet* **10**: 82.

Muenzner J, Trébulle P, Agostini F, Zauber H, Messner CB, Steger M, Kilian C, Lau K, Barthel N, Lehmann A, et al. 2024. Natural proteome diversity links aneuploidy tolerance to protein turnover. *Nature* **630**: 149–157.

Oromendia AB, Dodgson SE, Amon A. 2012. Aneuploidy causes proteotoxic stress in yeast. *Genes Dev* **26**: 2696–2708.

Ozery-Flato M, Linhart C, Trakhtenbrot L, Izraeli S, Shamir R. 2011. Large-scale analysis of chromosomal aberrations in cancer karyotypes reveals two distinct paths to aneuploidy. *Genome Biol* **12**: R61.

Pan X, Yuan DS, Xiang D, Wang X, Sookhai-Mahadeo S, Bader JS, Hieter P, Spencer F, Boeke JD. 2004. A Robust Toolkit for Functional Profiling of the Yeast Genome. *Mol Cell* **16**: 487–496.

Passerini V, Ozeri-Galai E, de Pagter MS, Donnelly N, Schmalbrock S, Kloosterman WP, Kerem B, Storchova Z. 2016. The presence of extra chromosomes leads to genomic instability. *Nat Commun* **7**: 10754.

Pavani M, Bonaiuti P, Chiroli E, Gross F, Natali F, Macaluso F, Póti Á, Pasqualato S, Farkas Z, Pompei S, et al. 2021. Epistasis, aneuploidy, and functional mutations underlie evolution of resistance to induced microtubule depolymerization. *EMBO J* **40**: e108225.

Pavelka N, Rancati G, Zhu J, Bradford WD, Saraf A, Florens L, Sanderson BW, Hattem GL, Li R. 2010. Aneuploidy confers quantitative proteome changes and phenotypic variation in budding yeast. *Nature* **468**: 321–325.

Peter J, De Chiara M, Friedrich A, Yue J-X, Pflieger D, Bergström A, Sigwalt A, Barre B, Freel K, Llored A, et al. 2018. Genome evolution across 1,011 Saccharomyces cerevisiae isolates. *Nature* **556**: 339–344.

Rancati G, Pavelka N, Fleharty B, Noll A, Trimble R, Walton K, Perera A, Staehling-Hampton K, Seidel CW, Li R. 2008. Aneuploidy Underlies Rapid Adaptive Evolution of Yeast Cells Deprived of a Conserved Cytokinesis Motor. *Cell* **135**: 879–893.

Rathi S, Polat I, Pereira G. 2022. The budding yeast GSK-3 homologue Mck1 is an essential component of the spindle position checkpoint. *Open Biol* **12**: 220203.

Ravichandran MC, Fink S, Clarke MN, Hofer FC, Campbell CS. 2018. Genetic interactions between specific chromosome copy number alterations dictate complex aneuploidy patterns. *Genes Dev* **32**: 1485–1498.

Rojas J, Hose J, Dutcher HA, Place M, Wolters JF, Hittinger CT, Gasch AP. 2024. Comparative modeling reveals the molecular determinants of aneuploidy fitness cost in a wild yeast model. *Cell Genomics* **4**: 100656.

Ryu H-Y, Wilson NR, Mehta S, Hwang SS, Hochstrasser M. 2016. Loss of the SUMO protease Ulp2 triggers a specific multichromosome aneuploidy. *Genes Dev* **30**: 1881–1894.

Santaguida S, Vasile E, White E, Amon A. 2015. Aneuploidy-induced cellular stresses limit autophagic degradation. *Genes Amp Dev* **29**: 2010–2021.

Schatz PJ, Solomon F, Botstein D. 1986. Genetically essential and nonessential alpha-tubulin genes specify functionally interchangeable proteins. *Mol Cell Biol* **6**: 3722–3733.

Schibler A, Koutelou E, Tomida J, Wilson-Pham M, Wang L, Lu Y, Cabrera AP, Chosed RJ, Li W, Li B, et al. 2016. Histone H3K4 methylation regulates deactivation of the spindle assembly checkpoint through direct binding of Mad2. *Genes Dev* **30**: 1187–1197.

Schmittgen TD. 2001. Real-Time Quantitative PCR. *Methods* **25**: 383–385.

Selmecki A, Forche A, Berman J. 2006. Aneuploidy and Isochromosome Formation
in Drug-Resistant *Candida albicans*. *Science* **313**: 367–370.

Selmecki AM, Dulmage K, Cowen LE, Anderson JB, Berman J. 2009. Acquisition of
Aneuploidy Provides Increased Fitness during the Evolution of Antifungal
Drug Resistance ed. H.D. Madhani. *PLoS Genet* **5**: e1000705.

Sheltzer JM, Blank HM, Pfau SJ, Tange Y, George BM, Humpton TJ, Brito IL,
Hiraoka Y, Niwa O, Amon A. 2011. Aneuploidy Drives Genomic Instability in
Yeast. *Science* **333**: 1026–1030.

Shukla A, Nguyen THM, Moka SB, Ellis JJ, Grady JP, Oey H, Cristino AS, Khanna
KK, Kroese DP, Krause L, et al. 2020. Chromosome arm aneuploidies shape
tumour evolution and drug response. *Nat Commun* **11**: 449.

Sousa A, Gonçalves E, Mirauta B, Ochoa D, Stegle O, Beltrao P. 2019. Multi-omics
Characterization of Interaction-mediated Control of Human Protein
Abundance levels. *Mol Cell Proteomics* **18**: S114–S125.

Stingele S, Stoehr G, Peplowska K, Cox J, Mann M, Storchova Z. 2012. Global
analysis of genome, transcriptome and proteome reveals the response to
aneuploidy in human cells. *Mol Syst Biol* **8**: 608.

Su XA, Ma D, Parsons JV, Replogle JM, Amatruda JF, Whittaker CA, Stegmaier K,
Amon A. 2021. *RAD21* is a driver of chromosome 8 gain in Ewing sarcoma to
mitigate replication stress. *Genes Dev* **35**: 556–572.

Sugiyama M, Ikushima S, Nakazawa T, Kaneko Y, Harashima S. 2005. PCR-
Mediated Repeated Chromosome Splitting in *Saccharomyces* Cerevisiae.
BioTechniques **38**: 909–914.

Sugiyama M, Nakazawa T, Murakami K, Sumiya T, Nakamura A, Kaneko Y,
Nishizawa M, Harashima S. 2008. PCR-mediated one-step deletion of
targeted chromosomal regions in haploid Saccharomyces cerevisiae. *Appl
Microbiol Biotechnol* **80**: 545–53.

Terhorst A, Sandikci A, Keller A, Whittaker CA, Dunham MJ, Amon A. 2020. The
environmental stress response causes ribosome loss in aneuploid yeast cells.
Proc Natl Acad Sci U S A **117**: 17031–17040.

Thorburn RR, Gonzalez C, Brar GA, Christen S, Carlile TM, Ingolia NT, Sauer U,
Weissman JS, Amon A. 2013. Aneuploid yeast strains exhibit defects in cell
growth and passage through START ed. D.J. Lew. *Mol Biol Cell* **24**: 1274–
1289.

Tong AHY, Lesage G, Bader GD, Ding H, Xu H, Xin X, Young J, Berriz GF, Brost RL,
Chang M, et al. 2004. Global Mapping of the Yeast Genetic Interaction
Network. *Science* **303**: 808–813.

Torres EM, Sokolsky T, Tucker CM, Chan LY, Boselli M, Dunham MJ, Amon A. 2007.
Effects of Aneuploidy on Cellular Physiology and Cell Division in Haploid
Yeast. *Science* **317**: 916–924.

Trakala M, Aggarwal M, Sniffen C, Zasadil L, Carroll A, Ma D, Su XA, Wangsa D, Meyer A, Sieben CJ, et al. 2021. Clonal selection of stable aneuploidies in progenitor cells drives high-prevalence tumorigenesis. *Genes Dev* **35**: 1079–1092.

Tsai H-J, Nelliat A. 2019. A Double-Edged Sword: Aneuploidy is a Prevalent Strategy in Fungal Adaptation. *Genes* **10**: 787.

Ubeda J-M, Légaré D, Raymond F, Ouameur AA, Boisvert S, Rigault P, Corbeil J, Tremblay MJ, Olivier M, Papadopoulou B, et al. 2008. Modulation of gene expression in drug resistant Leishmania is associated with gene amplification, gene deletion and chromosome aneuploidy. *Genome Biol* **9**: R115.

Voordeckers K, Kominek J, Das A, Espinosa-Cantú A, De Maeyer D, Arslan A, Van Pee M, Van Der Zande E, Meert W, Yang Y, et al. 2015. Adaptation to High Ethanol Reveals Complex Evolutionary Pathways ed. J. Zhang. *PLOS Genet* **11**: e1005635.

Wangsa D, Braun R, Stuelten CH, Brown M, Bauer KM, Emons G, Weston LA, Hu Y, Yang HH, Vila-Casadesús M, et al. 2019. Induced Chromosomal Aneuploidy Results in Global and Consistent Deregulation of the Transcriptome of Cancer Cells. *Neoplasia* **21**: 721–729.

Watson EV, Lee JJ-K, Gulhan DC, Melloni GEM, Venev SV, Magesh RY, Frederick A, Chiba K, Wooten EC, Naxerova K, et al. 2024. Chromosome evolution screens recapitulate tissue-specific tumor aneuploidy patterns. *Nat Genet* **56**: 900–912.

Weaver BA, Cleveland DW. 2006. Does aneuploidy cause cancer? *Curr Opin Cell Biol* **18**: 658–667.

Williams BR, Prabhu VR, Hunter KE, Glazier CM, Whittaker CA, Housman DE, Amon A. 2008. Aneuploidy Affects Proliferation and Spontaneous Immortalization in Mammalian Cells. *Science* **322**: 703–709.

Yona AH, Manor YS, Herbst RH, Romano GH, Mitchell A, Kupiec M, Pilpel Y, Dahan O. 2012. Chromosomal duplication is a transient evolutionary solution to stress. *Proc Natl Acad Sci* **109**: 21010–21015.

Zerbib J, Ippolito MR, Eliezer Y, De Feudis G, Reuveni E, Savir Kadmon A, Martin S, Viganò S, Leor G, Berstler J, et al. 2024. Human aneuploid cells depend on the RAF/MEK/ERK pathway for overcoming increased DNA damage. *Nat Commun* **15**: 7772.

Zhang K, Zhang L-J, Fang Y-H, Jin X-N, Qi L, Wu X-C, Zheng D-Q. 2016. Genomic structural variation contributes to phenotypic change of industrial bioethanol yeast *Saccharomyces cerevisiae* ed. Z. Zhao. *FEMS Yeast Res* **16**: fov118.

Zhou X, Hilk A, Solis NV, Scott N, Beach A, Soisangwan N, Billings CL, Burrack LS, Filler SG, Selmecki A. 2024. Single-cell detection of copy number changes reveals dynamic mechanisms of adaptation to antifungals in Candida albicans. *Nat Microbiol* **9**: 2923–2938.

Figure 1: Strong genetic interactions between aneuploid chromosomes are rare
(A) Schematic overview of the chromosome gain and chromosome loss collections. (B) Quantified growth of the chromosome gain and chromosome loss collections. The averages of four independent high-throughput growth assays were normalized to WT. Colony intensities were quantified after 24h on YPAD plates for chromosome gains, and 72h on plates containing 5-Fluoroorotic Acid (5-FOA) for chromosome losses. Single disomies/monosomies are depicted in grey, double disomies/monosomies in black. r^2 values are from quadratic fitts. (C) Correlation between growth of the double disomies and growth predictions based on the single disomies (p < 0.0001). Growth was predicted by calculating the product of single disomy growth. r^2 values are from simple linear regression and p-values are from *F*-tests. (D) Quantification of colony sizes on YPAD plates after 48h normalized to a euploid diploid strain. (n=3).

Figure 2: The strongest drug resistance phenotypes come from complex aneuploidy

Figure 2: The strongest drug resistance phenotypes come from complex aneuploidy

(A) Schematic of the work-flow for plate-based high throughput growth assays under selective conditions. Cells were grown in liquid cultures before being transferred to condition specific agar plates. Images for colony intensities were taken after 24, 36 , or 48h. (B) Averaged growth of two independent high-throughput growth assays. Control plates were quantified after 24h; hygromycin (40 µg/ml) and rapamycin (5 nM) plates after 48h; and MMS (0.02%) and benomyl (17.5 µg/ml) plates after 36h. All quantifications were normalized to a haploid WT. (C) Growth measurements of single disomies (Single) and double disomies (Double) across different treatments. Mean of each group is indicated by a red line. The YPAD control is an average of YPAD and YPAD + DMSO plate growth. The dashed line indicates WT growth. (D) Epistasis scores were calculated by subtracting the predicted growth (the product of the growth of the single disomies) from the measured growth. Genetic interactions are based on the normal distribution of epistasis scores without selective pressure. Positive genetic interactions have an epistasis score greater than the median plus 2 standard deviations and negative interactions have an epistasis score less than the median minus 2 standard deviations.

Figure 3: Benomyl resistance correlates with alpha-tubulin expression in aneuploid strains

Fiure. 3: Benomyl resistance correlates with alpha-tubulin expression in aneuploid strains
(A) 10-fold serial dilution analysis of benomyl resistant single disomies on control (YPAD + DMSO, 24h) and treated (YPAD + 20 µg/ml benomyl, 48h) plates. (B) Western blot analysis of α-tubulin expression levels of single disomies. *Pgk1* is shown as a loading control. (C) Western blot quantifications of α-tubulin (n = 3) and β-tubulin (n = 3) expression levels for all single disomies. Strains were grown in YPAD and harvested in log-phase. Quantified bands were normalized to the Ponceau stained membrane and the haploid WT. Relative growth on benomyl denotes the average growth of single disomies from two independent high throughput colony size assays. r² values are from simple linear regression and p-values are from *F*-tests.

Figure 4: Identification of chromosome arms necessary for aneuploidy-based drug resistance

Figure 4: Identification of chromosome arms necessary for aneuploidy-based drug resistance.

(A) Schematic of two models for the genetic basis behind aneuploidy driven drug resistance. (B) Method of chromosome arm deletion. Chromosome arms are deleted by homologous recombination with a 500 bp homology sequence connected to a resistance marker and a telomeric seed sequence. (C) Doubling times of whole chromosome and single arm disomies for chromosomes 4, 5, 10 and 14. ORFs depict the remaining aneuploid ORFs after chromosome arm deletions. (D) 10-fold dilution series and quantification of whole chromosome and single arm disomies (n=3). Control plates (YPAD + DMSO) were imaged after 24h and treated plates (YPAD + 25 µg/ml benomyl) after 48h. * Indicates a lower benomyl concentration (20 µ/ml benomyl). The second dilution was quantified and normalized to WT, with the exceptions of chromosome 13 (third dilution) and chromosome 12 (first dilution). p-values are from one-way ANOVA, uncorrected Fisher's LSD test. (E) Quantification of 10-fold serial dilutions treated with MMS (0.04 %), rapamycin (10 nM) or hygromycin (60 µg/ml). Relative growth was calculated as the ratio of normalized growth on treated plates (48h) to normalized growth on control plates (24h). Representative examples for single arm sensitivity, combinatorial sensitivity, compensatory sensitivity, resistance, and no interactions are shown. Images and quantifications of all 10-fold serial dilutions are depicted in Supplemental Fig. S4A, B.

Figure 5: Most aneuploidy-based drug resistance phenotypes come from the combined effects of multiple genes.

Figure 5: Most aneuploidy-based drug resistance phenotypes come from the combined effects of multiple genes.

(A) Representative image of 10-fold serial dilution and quantification of partial chromosome arm deletions of the right arm of chromosome 4. For control plates (YPAD + DMSO) the 2rd spot was quantified after 24h and for treated plates (YPAD + 25µg/ml benomyl) the 2nd and 3rd spots were quantified and averaged after 48h (n=3). p-values are from one-way ANOVA, uncorrected Fisher's LSD test. Growth on DMSO was normalized to a haploid WT and growth on benomyl to a strain disomic for chromosome 4. (B) 10-fold serial dilution and quantification of partial chromosome arm deletions and heterozygous deletions of genes contributing to benomyl resistance. For treated plates (YPAD + 25µg/l benomyl) the 3rd spot was quantified after 48h and normalized to the strain disomic for chromosome 4 (n=3). p-values are from one-way ANOVA, uncorrected Fisher's LSD test. (C) Schematic depiction of benomyl resistant and sensitive chromosome regions. Identified resistance genes are indicated in orange at their approximate chromosomal location.

Figure 6: Synergistic effects contribute to the strongest aneuploidy phenotypes.

Figure 6: Synergistic effects contribute to the strongest aneuploidy phenotypes.
(A) Single disomy growth on benomyl compared to growth predictions based on single arm disomies. Growth was quantified from 10-fold serial dilutions in Figure 4D. The area highlighted in magenta indicates additive genetic interactions (+/-30% of predicted growth). Disomies that fall below this threshold are considered a negative synergistic interaction and those above are considered a positive synergistic interaction. (B) 10-fold serial dilution of single and double disomies. Control plates (YPAD + DMSO) were imaged after 24h and treated plates (YPAD + 30 µg/ml benomyl) after 48h. (C) 10-fold serial dilutions of benomyl resistant *set1Δ* haploid and two synergistically interacting double disomies (2&4, 2&13). Benomyl plates contained 30 µg/ml benomyl. (D) Quantification of the 10-fold serial dilutions of the double disomies on plates with 30 µg/ml benomyl compared to predicted growth based on the additive resistance of the single disomies (for images, see Sup. Figure 8A). Combinations in the highlighted area (magenta) are additive genetic interactions (+/-30% of predicted growth). Disomy combinations that fall below this threshold are considered a negative synergistic interaction and those above are considered a positive synergistic interaction.