Combinatorial effects of multiple genes contribute to beneficial aneuploidy phenotypes

M. Sophie Koller[1,2], Claudia Himmelbauer [1], Sarah Fink[1], Madhwesh C. Ravichandran[1], Christopher S. Campbell[1@]

[1]Department of Chromosome Biology, Max Perutz Labs, University of Vienna, Vienna Biocenter (VBC), A-1030 Vienna, Austria
[2]Vienna Biocenter PhD Program, a Doctoral School of the University of Vienna and the Medical University of Vienna, A-1030 Vienna, Austria

@: Corresponding author
Email: christopher.campbell@univie.ac.at
Phone: +43-1-4277-74418
Address: Dr. Bohr-Gasse 9/5, A-1030 Vienna, Austria

Key words: Aneuploidy; complex karyotypes; drug resistance; tubulin expression;

Abstract

Aneuploidy is one of the most common mechanisms of adaptation to environmental selection in vegetatively growing cells. However, it is still unclear what advantages aneuploidy provides over other types of genomic alterations. In this study, we aimed to determine if beneficial aneuploidy phenotypes are driven primarily by combinatorial expression changes of multiple genes using budding yeast as a model system. To determine the impact of complex aneuploidy on cellular fitness and resistance to a variety of drug treatments, we generated yeast collections with nearly every possible combination of two chromosome gains or losses. In addition, we genetically dissected aneuploid chromosomes using partial deletions to identify chromosomal regions that contribute to aneuploidy-driven drug resistance. Strong resistance phenotypes consistently came from combinations of chromosomes or chromosomal regions. We identified up to five regions on a single chromosome that contribute to resistance to a single drug treatment. Finally, we found that the strongest aneuploidy phenotypes came from synergistic effects between copy number changes of different chromosomes or chromosomal regions. These results provide insights into how combining subtle gene expression changes across the genome impacts aneuploidy phenotypes.

Introduction

The accurate distribution of chromosomes during cell division is essential for maintaining genomic integrity across all eukaryotic species. Chromosome segregation errors result in alterations in chromosome copy number, which are termed aneuploidy. Aneuploidy is one of the most common genetic alterations in cancer, occurring in ~90% of solid tumors and ~70% of hematopoietic cancers (Weaver and Cleveland 2006). These high frequencies of aneuploidy are not only inherent to cancer cells but are also commonly observed within other vegetatively growing cell populations including drug resistant strains of the human pathogen *Candida albicans* and industrial strains of baker's yeast (Selmecki et al. 2006; Peter et al. 2018; Loegler et al. 2024).

Despite the frequency of aneuploidy in mitotically growing cell populations, it is generally associated with reduced cellular fitness. Engineering aneuploidy in cells invariably decreases growth rates under optimal conditions (Torres et al. 2007; Stingele et al. 2012). This decrease in fitness results from changes in gene copy numbers of many transcribed genes on the aneuploid chromosome (Bonney et al. 2015). For both yeast and human cells, gene expression levels of lab engineered aneuploidies scale with gene copy numbers for most of the aneuploid genes (Torres et al. 2007; Stingele et al. 2012; Wangsa et al. 2019; Rojas et al. 2024; Muenzner et al. 2024). These small changes in expression levels for hundreds of genes cause a wide range of deleterious phenotypes. Such general aneuploidy-associated phenotypes include increased proteotoxic stress, cell cycle delays, and increased DNA damage (Torres et al. 2007; Sheltzer et al. 2011; Thorburn et al. 2013; Oromendia et al. 2012; Beach et al. 2017; Li and Zhu 2022).

Yet, aneuploidy is one of the most frequently observed mechanisms of cellular adaptation. The first direct link between adaptive drug resistance and aneuploidy was established in the human pathogen *Candida albicans* (Selmecki et al. 2006). Adaptation via aneuploidy has since been more extensively observed in the budding yeast *Saccharomyces cerevisiae,* where specific aneuploidies consistently emerge during experimental evolution screens under adverse environmental conditions (Gresham et al. 2008; Yona et al. 2012; Chen et al. 2012; Voordeckers et al. 2015; Zhang et al. 2016; Beaupere et al. 2018; Morard et al. 2019). Moreover, inducing

aneuploidy via random chromosome missegregation in both yeast and human cells
promotes resistance to a large variety of drugs (Pavelka et al. 2010; Chen et al. 2012;
Lukow et al. 2021; Ippolito et al. 2021). Aneuploidy can also act as a suppressor of
mutations that decrease cellular fitness (Rancati et al. 2008; Liu et al. 2015; Ryu et al.
2016; Ravichandran et al. 2018; Macaluso et al. 2025). In cancer, karyotype patterns
have predictive power for drug resistance, tumor progression and patient outcome
(Shukla et al. 2020).

The potential for aneuploidy as an adaptation mechanism to a large variety of selective
environments can partially explain the high prevalence of aneuploidy following long-
term vegetative growth, such as in cancer or in yeast used for alcohol production
(Peter et al. 2018). However, it still fails to explain why cells adapt through aneuploidy
over other types of mutations that don't share its inherent downsides. Different models
for the specific advantage of adaptation through aneuploidy have been proposed.
Aneuploidy has been suggested to be a rapid, but transient solution to sudden
environmental changes. In *S. cerevisiae* subjected to acute heat stress, chromosome
3 is gained early in the adaptation process but then lost again over time (Yona et al.
2012). Similarly, yeast adapting to high rates of chromosomal instability (CIN) adapted
first through the accumulation of specific aneuploidies, followed later by the
accumulation of suppressor point mutations (Clarke et al. 2023). However, there are
also many examples of specific aneuploidies that are maintained throughout long-term
adaptation, highlighting that speed is likely not the only advantage of adaptation via
aneuploidy (Selmecki et al. 2009; Ravichandran et al. 2018; Adell et al. 2023; Clarke
et al. 2023; Zhou et al. 2024; Watson et al. 2024).

An additional explanation for the advantages of aneuploidy as a mechanism of
adaptation comes from its ability to change gene expression levels of several
beneficial genes simultaneously. In yeast adaptation screens, only a handful of
examples could attribute the selective advantage of aneuploidy to changes in
expression levels of one or two specific genes. Even in these cases, they rarely
recapitulated the strength of the aneuploidy phenotype (Selmecki et al. 2009; Chen et
al. 2012; Ryu et al. 2016; Linder et al. 2017; Beaupere et al. 2018; Ravichandran et
al. 2018; Barney et al. 2021; Pavani et al. 2021; Zhou et al. 2024). In cancer,
overexpression of individual genes also fails to fully recapitulate the phenotypes of

aneuploid chromosomes (Trakala et al. 2021; Su et al. 2021; Girish et al. 2023). These

findings suggest that the challenges of identifying driver genes on aneuploid

chromosomes could be amplified by the interplay between multiple genes. It is

therefore important to determine if multiple genes acting cooperatively is a general

characteristic of aneuploid chromosomes that confer an adaptive advantage.

In this study, we generated a large yeast collection with nearly every possible

combination of two chromosome gains or losses to determine how combinatorial

aneuploidy phenotypes impact drug resistance and sensitivity. We found that the

strongest resistance phenotypes consistently came from combinations of aneuploid

chromosomes, demonstrating the advantage of copy number changes in multiple

genes. Additionally, we genetically dissected resistant aneuploid chromosomes using

partial arm deletions to determine the number of regions that contribute to aneuploidy-

driven drug resistance on a single chromosome. We found that resistance conferred

by single disomies nearly always comes from multiple chromosomal regions. While

the majority of phenotypes are additive, the strongest resistance phenotypes of

aneuploidies came from synergistic effects between different regions or

chromosomes.

Results

**Generation of yeast strain libraries with combinations of chromosome gains
and losses**

We first determined how aneuploidy of different regions of the genome combine to

impact cellular fitness and drug response by generating two collections of yeast with

combinations of chromosome gains or losses. The collections consist of nearly all

possible single and double chromosome gains and losses (Fig. 1A). To engineer

chromosome gain or loss, chromosome-specific missegregation was first induced via

a strong galactose-inducible centromere-proximal promotor, followed by selection for

the specific aneuploidy (Supplemental Fig. S1A). We selected for chromosome gains

via recombination events that restore the function of an auxotrophic marker.

Chromosome loss was generated by selecting against the *URA3* gene using 5-

fluoroorotic acid (5-FOA) (Supplemental Fig. S1A). The chromosome gain collection

includes most of the possible single (14/16) and double disomic (90/136) strains in

haploid cells; each was confirmed by qPCR (Supplemental Table S1). The

chromosome loss collection contains every possible single or double monosomy in

diploid cells (Supplemental Fig. S1B). Since chromosome loss cannot be continuously

selected for, this collection is inducible to generate newly aneuploid cells prior to each

experiment. These libraries currently constitute the largest collections of engineered

complex aneuploid cells.

We first evaluated the libraries by measuring the growth rates of the engineered

aneuploidy strains under optimal growth conditions. Similar to previous observations

with single aneuploidies (Beach et al. 2017), the growth of complex aneuploidies

scaled with the number of aneuploid open reading frames (ORFs) and chromosome

loss is more detrimental to cell growth than chromosome gain (Fig. 1B). Furthermore,

the high-throughput growth rate measurements of single disomies and monosomies

strongly correlate with those previously observed (Supplemental Fig. S1C). The

previously published measurements were performed in a strain background containing

a mutation in the *SSD1* gene, which has been reported to sensitize cells to aneuploidy

phenotypes (Hose et al. 2020). However, we observe similar results in a different,

wildtype *SSD1* background (S288c), indicating that strain background does not

substantially alter the relative growth rate of aneuploid yeast.

We next determined the impact of combining aneuploid chromosomes on cellular

fitness. Most of the single and double monosomies are lethal. By contrast, nearly all

of the double disomies are viable, with the sole exception of those involving

chromosome 6 (Fig. 1B, Supplemental Fig. S1B). To identify potential genetic

interactions between aneuploid chromosomes, we calculated growth predictions for

all chromosome combinations based on the product of the two single aneuploidies. In

general, growth of the double disomies correlates well (r^2 = 0.64) with the calculated

growth predictions. The strongest outlier is the combination of chromosomes 6 and 13

(Fig. 1C). Chromosome 6 disomy is lethal on its own due to the presence of an extra

copy of the β-tubulin *TUB2* gene (Anders et al. 2009). However, chromosome 6 disomy

becomes viable with the additional gain of chromosome 13. This is consistent with

previous observations where selection of chromosome 6 gain resulted in a strain that

spontaneously gained the additional disomies of chromosomes 1 and 13 (Torres et al.
2007). Chromosome 13 contains the genes for α-tubulin, and overexpression of α-
tubulin rescues the lethality of chromosome 6 disomy (Anders et al. 2009). For the
chromosome loss collection, the strongest genetic interaction is again between
chromosomes 6 and 13 (Fig. 1D). Chromosome 13 monosomy is lethal on its own, but
the additional loss of chromosome 6 rescues viability (Fig. 1D). The heterozygous
deletion of *TUB2* is sufficient to rescue chromosome 13 monosomy, indicating that
excess β-tubulin relative to α-tubulin is the genetic basis behind these strong
chromosome copy number interactions (Figure 1D).

To find additional genetic interactions between chromosome gains, we identified
chromosome combinations whose growth significantly differed from the prediction
based on single disomies. The data adhered to a normal distribution, enabling the
calculation of z-scores (Kolmogorov-Smirnov test). Significant interactions were
defined as having z-scores >1 or <-1. Among the 90 chromosome combinations
analyzed, we identified 11 positive (12.2%) and 11 negative (12.2%) chromosome
copy number interactions (Supplemental Fig. S1D). Importantly, these interactions are
substantially stronger than interactions previously demonstrated to heavily impact
karyotype pattern formation (Ravichandran et al. 2018). We conclude that, although
most combinations of aneuploid chromosomes show additive phenotypes,
chromosome copy number interactions are a common feature of aneuploidy
combinations.

The strongest drug resistance phenotypes come from strains with multiple
aneuploid chromosomes

We next used the chromosome gain library to determine if combinatorial interactions
between aneuploid chromosomes contribute to improved growth in stress conditions.
We measured growth in the presence of four drugs that induce cellular stresses
associated with aneuploidy: hygromycin B (ribosome inhibitor), rapamycin (TOR
pathway inhibitor), methyl methanesulfonate (MMS, DNA damage inducer) and
benomyl (microtubule depolymerizer) (Fig. 2A). Drug concentrations were titrated to
reduce wild-type growth by ~80%. Growth was unaffected by the solvent used to
resuspend the drugs (Supplemental Fig. S1E). Aneuploidy substantially altered the

cellular response to hygromycin, rapamycin, and benomyl, as growth rates no longer

scaled with number of genes on aneuploid chromosomes (Fig. 2B). By contrast, MMS

treatment showed fewer aneuploidy-associated phenotypes. The percentage of

strains displaying substantial relative resistance (normalized growth in the

presence/absence of the drug > 2) across the entire library varies greatly between the

different drugs ranging from 59% for benomyl to 9% for MMS (Supplemental Fig. S2B,

C). We conclude that the capacity for aneuploidy to provide resistance is drug

dependent.

In addition to drug resistance, we also observed frequent drug sensitivity. This was

most pronounced for hygromycin, where over 78% of strains showed a greater than

two-fold relative decrease in growth in the presence of the drug. Additionally, we

measured sensitivity to two stress conditions other than drugs: 39°C for heat stress

and 18°C for cold stress. Neither of these conditions greatly influenced cell growth in

the euploid strain. However, heat stress led to a two-fold decrease in growth for 66%

of the disomic strains, which scaled with the degree of aneuploidy (Supplemental

Fig. S2A-C). These results suggest that aneuploid yeast are especially sensitive to

hygromycin and heat stress, which implicate translation inhibition and protein

misfolding as potential vulnerabilities in aneuploid cells. This would agree with

previous reports that aneuploid strains frequently exhibit decreased ribosome levels

and proteotoxic stress (Oromendia et al. 2012; Terhorst et al. 2020).

We next wanted to determine if combinations of aneuploid chromosomes show

stronger phenotypes than individual chromosomes. For hygromycin, rapamycin and

benomyl, specific chromosome combinations had the highest levels of drug

resistance, despite their decreased fitness in the absence of the drugs (Fig. 2C).

Notably, absolute resistance to hygromycin was observed only for chromosome

combinations, as none of the single disomies grew better than the euploid strain in the

presence of hygromycin. To determine if the strength of resistance for chromosome

combinations can be predicted based on the single disomies, we compared the growth

of double disomies to the product of the single disomies (Supplemental Fig. S2D).

Similar to what we observed under standard growth conditions, the predicted and

measured growth correlate well for all four drug treatments. This suggests that the

majority of the phenotypes for chromosome combinations are additive, and that

increased expression of additional resistance genes outweighs the negative effects

from the increased expression of all of the other genes when aneuploidies are

combined. However, there were some substantial outliers, suggesting that

chromosome copy number interactions also play a role in drug resistance/sensitivity

(Supplemental Fig. S2D). Overall, we conclude that the strongest drug resistance

aneuploidy phenotypes typically come from the combined effects of multiple genes on

different chromosomes.

Benomyl resistance correlates with α-Tubulin expression in aneuploid strains.

Surprisingly, a majority (59%) of disomic chromosome combinations were resistant to

the microtubule depolymerizing factor benomyl (Supplemental Fig. S2B, C). This high

percentage results from 6 of the 14 tested single disomic chromosomes conferring

substantial resistance to benomyl (Fig. 2C, 3A). Aneuploidy frequently leading to

resistance to microtubule inhibitors was not specific to benomyl, as four of the same

aneuploid chromosomes provided resistance to another microtubule depolymerizing

drug, nocodazole (Supplemental Fig. S3A). We first focused on chromosome 13, as it

contains both α-tubulin genes (*TUB1* and *TUB3*). Heterozygous deletion of *TUB3* in

the chromosome 13 disomic strain eliminated the resistance phenotype,

demonstrating that this gene contributes to the resistance (Supplemental Fig. S3B).

To determine if α-tubulin protein abundance contributes to benomyl resistance in the

other aneuploid strains, we measured the expression levels in all 14 disomic strains

via quantitative western blots. We found that multiple single disomies alter α-tubulin

expression levels (Fig. 3B). The measured α-tubulin expression levels correlated well

with those measured by high-throughput proteomics of aneuploid yeast strains,

demonstrating that these differences occur independently of the measurement method

and strain background (Supplemental Fig. S3C, (Muenzner et al. 2024)). Elevated α-

tubulin expression levels strongly correlated with the degree of benomyl resistance for

the single disomies, suggesting that this is a major mechanism of resistance (Fig. 3C).

Intriguingly, only chromosome 12 disomy is extremely sensitive to benomyl and is also

the only disomy that shows a substantial decrease in α-tubulin expression. The

correlation with benomyl resistance was much weaker for β-tubulin expression (Fig.

3C, Supplemental Fig. S3D). These results demonstrate that there are multiple genes

across the genome that can alter tubulin expression levels when present at an extra

copy, and these mild expression changes can greatly alter the response to microtubule

inhibitors.

Identification of chromosome arms necessary for aneuploidy-based drug

resistance.

We next aimed to determine which regions of particular chromosomes provide

resistance when gained by a single copy. We focused primarily on benomyl resistance,

since six individual disomies led to strong resistance to the drug. This provided

multiple examples to systematically examine how many regions/genes typically

contribute to positive aneuploidy phenotypes for single aneuploid chromosomes. One

possibility is that a single strong resistance gene on each chromosome could be

sufficient to give a substantial fitness advantage under harsh drug conditions (single

gene model). Another possibility is that multiple mildly beneficial genes on single

chromosomes may act together to create strong resistance phenotypes (combinatorial

model) (Fig. 4A).

To determine which regions of chromosomes have negative, positive, or neutral effects

on growth with and without benomyl, we implemented a method that deletes part of a

chromosome from the end and replaces it with a selection marker and an artificial

telomere (Linder et al., 2017) (Fig. 4B). We first generated single arm disomies of all

of the benomyl resistant single disomic chromosomes. The deletion of one of the two

copies of the chromosome arms was confirmed by qPCR. As with whole chromosome

disomies, arm-level disomies have decreased growth rates proportional to the number

of aneuploid ORFs under standard conditions (Fig. 4C). Benomyl resistance was

frequently associated with a single arm. For chromosomes 2, 4, 10, and 14, disomy of

the larger arm was sufficient to fully recapitulate the benomyl resistance, whereas

chromosome 13 only showed resistance with disomy of the shorter arm (Fig. 4D). Both

α-tubulin genes (*TUB1 and TUB3)* are on the short (left) arm of chromosome 13.

Intriguingly, neither of the chromosome 5 arm disomies recapitulated the benomyl

resistance of the whole chromosome disomy. Chromosome 5 left arm disomy shows

no resistance, while the right arm disomy is only mildly resistant. Therefore, the

benomyl resistance for chromosome 5 results from at least one gene on each of the

two chromosome arms (Fig. 4D). Overall, the correlation between growth with and

without benomyl for the arm-level disomies was not significant, demonstrating that

benomyl resistance is not the result of slow growth or a general aneuploidy phenotype
(Supplemental Fig. S4A). Instead, resistance likely results from the increased
expression of one or more specific genes on the aneuploid chromosome arms.

Next, we used our arm level aneuploidies to analyze resistance or sensitivity to the
three other drugs tested: hygromycin, MMS and rapamycin (Supplemental Fig. S4B).
Similar to benomyl, resistance to MMS, rapamycin and hygromycin was primarily
associated with single chromosome arms (3/3). By contrast, arm-level aneuploidies
generally did not fully recapitulate increased drug sensitivity phenotypes (Fig. 4E,
Supplemental Fig. S4B, C). Instead, both arms often showed partial sensitivity. This is
also the case for the only chromosome that showed strong sensitivity to benomyl,
chromosome 12 (Fig. 4D). These results suggest that aneuploidy-based drug
sensitivity results from the combined effects of more genes than drug resistance.
Surprisingly, in some examples, drug sensitivity from one chromosome arm was
compensated for by the resistance from the second chromosome arm (2/8)
(Supplemental Fig. S4B, C).

Most aneuploidy-based drug resistance phenotypes come from the combined
effect of multiple genes

Since most of the resistance phenotypes were associated with the larger arms, we
wanted to determine how many regions/genes on these arms contribute to benomyl
resistance. We therefore made multiple shorter deletions to the benomyl resistant
disomies to identify regions of drug resistance. For example, seven additional cuts
were made on the right arm of chromosome 4 to identify regions of interest. As the
size of the chromosome decreased, growth under standard conditions gradually
increased (Fig. 5A). This further demonstrates that the negative effects on fitness for
aneuploid strains result from many genes across the length of a chromosome (Bonney
et al. 2015). Conversely, benomyl resistance decreased in a stepwise manner
following the deletion of three independent regions on the right arm (Fig. 5A). For 5 of
the 6 resistant chromosomes, we identified at least two regions that contributed to
resistance, with the most being 5 regions for chromosome 5 (Fig. 5C,
Supplemental Fig. S5). Interestingly, three of the benomyl resistant chromosomes
also had regions that increased drug sensitivity (Fig. 5A, C, Supplemental Fig. S5A).

The strength of drug resistance is therefore a balance between regions that confer
resistance and sensitivity.
318
We next aimed to identify the resistance genes within the identified regions. The
regions each contain between ~30 and 100 genes. We identified candidate genes by
looking for: sensitivity to benomyl in high-throughput screens of the yeast knockout
collection, essential genes, and genes that have GO Terms associated with
tubulin/microtubules. Candidate genes were tested via heterozygous deletions in the
disomic strains (Supplemental Fig. S6A). For the two regions on chromosome 4
associated with the strongest benomyl resistance, we identified single gene deletions
that decreased resistance to a similar extent as deleting the entire chromosomal
region (Fig. 5B, C). In total, we identified 9 genes that decrease the resistance of
aneuploid chromosomes when deleted heterozygously (Supplemental Fig. S6A, S7).
In addition to *TUB3* on chromosome 13, we identified two additional genes associated
with tubulin expression (*GIM4*-chr.5, *PFD1*-chr.10). Both of these genes are members
of the prefoldin complex, which is a chaperone that promotes the proper folding of α-
and β-tubulin. One resistance gene is involved in microtubule stabilization (*BIM1*-
chr.5). Interestingly, two of the resistance genes have recently been implicated in the
spindle position checkpoint (*SWR1*-chr.4, *MCK1*-chr.14) (Rathi et al. 2022; Caydasi et
al. 2023). Another resistance gene, *TOF1*-chr.14, is primarily associated with
protection of the DNA replication fork. However, it shows a synthetic lethal interaction
with *BIM1*, indicating an additional role in chromosome segregation (Tong et al. 2004;
Pan et al. 2004). Overall, we have identified many genes across the genome that
contribute to aneuploidy-based benomyl resistance.
340
In addition to the disomic chromosomes that lead to benomyl resistance, we also
identified chromosomal regions that lead to rapamycin resistance on chromosome 10.
Strains with disomy of chromosome 10 have previously been shown to be rapamycin
resistant (Torres et al. 2007). This was speculated to be due to the presence of the
TOR1 gene. Heterozygous deletion of *TOR1* in a chromosome 10 disomic strain only
partially decreased rapamycin resistance, indicating that there are more genes on
chromosome 10 that contribute to the phenotype (Supplemental Fig. S6B).
Furthermore,   both   chromosome   arms   contribute   to   rapamycin   resistance

(Supplemental Fig. S6C). We conclude that the combined overexpression of multiple

genes leading to strong phenotypes is a common feature of aneuploidy.

The strongest aneuploidy phenotypes result from synergistic effects

Chromosome 5 disomy has the strongest benomyl resistance phenotype and is the

only chromosome where both arms contribute to the phenotype (Fig. 4D). Additionally,

the degree of resistance for the whole chromosome was substantially higher than

either of the two individual arm-level disomies, suggesting a synergistic interaction

between the two arms. To look at this more systematically, we predicted the strength

of resistance for whole chromosome aneuploidy based on the resistance of the

individual arm-level aneuploidies (Fig. 6A). Although the other single disomies

displayed resistance in line with the predictions, benomyl resistance for chromosome

5 was substantially higher than predicted. This demonstrates a strong synergistic

interaction between the two arms.

We next wanted to determine if the strongest phenotypes for double disomic strains

also result from synergistic interactions. We focused on chromosome combinations

where the single disomies on their own were benomyl resistant. We initially observed

that the combined disomies of chromosomes 2 & 4 and 2 & 13 are exceptionally

strong, showing robust resistance even at higher benomyl concentrations (Fig. 6B).

These combinations are as resistant as a deletion mutation known to be highly

resistant to benomyl (Schibler et al. 2016), despite strong growth defects for aneuploid

strains in the absence of the drug (Fig. 6C). Looking at combinations with chromosome

2 specifically, we observed a wide range of resistance phenotypes, ranging from

strong positive to strong negative interactions (Fig. 6B, Supplemental Fig. S8A).

Strikingly, a strong negative genetic interaction between chromosomes 2 and 14

results in the near-complete loss of the benomyl resistance (Fig. 6B, D). Almost all

combinations with chromosome 14 grow worse than the single disomies

(Supplemental Fig. S8B), indicating a general negative effect of chromosome 14

combinations and benomyl resistance. We also observed a positive synergistic

interaction between chromosomes 3 and 12 for rapamycin resistance, demonstrating

that such interactions are not benomyl-specific (Supplemental Fig. S8C). We conclude

that although most aneuploidies display additive effects between genes, regions, and

chromosomes, the strongest beneficial growth phenotypes often come from

synergistic interactions.

Discussion

In this study, we generated a library of engineered single and double aneuploid yeast

strains to systematically determine the genetic drivers of aneuploidy-based drug

resistance. In addition, we used a comprehensive approach to identify regions/genes

on aneuploid chromosomes that contribute to aneuploidy phenotypes. This approach

has numerous advantages, allowing for the identification of: i) multiple regions on a

chromosome that contribute to a phenotype, ii) both resistant and sensitive regions on

a single chromosome, iii) regions that act synergistically, where each region on its own

would not have a strong phenotype. Together, these strategies provide a means to

comprehensively identify the genetic determinants of positive and negative selection

for aneuploid chromosomes either alone or in combination.

When measuring the resistance of the chromosome gain collection to different

inhibitors, the strongest resistance phenotypes were consistently identified for

chromosome combinations. This demonstrates that altered copy numbers of multiple

genes combine to create stronger beneficial aneuploidy phenotypes. These additional

benefits are observed despite the often substantial negative growth phenotypes for

the double aneuploid strains under optimal growth conditions. Therefore, the benefits

of combining multiple aneuploid chromosomes generally outweigh the downsides

under selective conditions. This combinatorial behavior of aneuploid chromosomes

could explain the frequent occurrence of complex aneuploidy in both adapted yeast

populations and cancer karyotypes (Mitelman Database of Chromosome Aberrations

and Gene Fusions in Cancer, 2025. https://mitelmandatabase.isb-cgc.org, Rancati et

al. 2008; Ravichandran et al. 2018; Peter et al. 2018; Klockner and Campbell 2024).

In addition to identifying strong phenotypes for combinations of aneuploid

chromosomes, we discovered multiple regions within individual chromosomes that

contribute to drug resistance phenotypes. We identified up to 5 regions on a single

chromosome that contribute to drug resistance, emphasizing that multiple genes are

typically responsible for strong aneuploidy phenotypes. This combinatorial action of

many genes suggests an advantage for the selection of aneuploidy over other types

of genomic alterations. Changing the expression of genes with distinct and

complementary functions may provide proliferative benefits that could not be

accomplished through single mutations. These data provide substantial support for a

"combinatorial genes" explanation for the specific benefits of aneuploidy over other

genomic alterations.

Increasing the degree of selection via higher concentrations of benomyl revealed

strong positive synergistic interactions between aneuploid chromosomes. We also

observed a clear synergistic interaction between the arms of chromosome 5. These

results suggest another possible advantage for adaptation through aneuploidy,

wherein the altered expression levels of two complementary genes would not be

selected for individually due to a lack of phenotype. However, two simultaneous

changes to the genome through a single chromosome missegregation event would

enable their combined selection, leading to a strong beneficial phenotype. These

results, along with the recently discovered synergistic interaction between

chromosomes 3 and 6 in multi-azole tolerance of the human pathogen *Candida*

albicans (Zhou et al. 2024), demonstrate that positive synergistic interactions are more

heavily selected for under higher stress conditions. We also find that combining

beneficial aneuploidies can occasionally lead to weaker than expected phenotypes,

highlighting that negative genetic interactions also likely impact complex karyotype

formation. Positive and negative genetic interactions between co-occurring

aneuploidies have been suggested based on correlations between chromosome arm

frequencies in cancer aneuploidy patterns (Ravichandran et al. 2018). Such co-

occurrence patterns can be highly predictive of disease progression and

chemotherapy resistance in cancer (Shukla et al. 2020).

The main drivers of beneficial aneuploidy phenotypes are alterations in expression

levels resulting from gene copy number changes of specific resistance-driving genes

(Selmecki et al. 2009; Chen et al. 2012; Ravichandran et al. 2018; Barney et al. 2021;

Ryu et al. 2016; Zhou et al. 2024; Pavani et al. 2021; Beaupere et al. 2018; Linder et

al. 2017; Adell et al. 2023; Watson et al. 2024). We observed a significant correlation

between growth on benomyl and altered α-tubulin expression levels. In addition to

TUB3 on chromosome 13, two members of the prefoldin complex (*GIM4*-chr. 5, *PFD1*-

chr. 10) contribute to aneuploidy-based benomyl resistance. Intriguingly, chromosome

12 disomy decreases tubulin expression levels and is very sensitive to microtubule

destabilization via benomyl. This decrease in tubulin levels for chromosome 12 disomy

could also help explain how it rescues microtubule hyperstabilization (Macaluso et al.

2025). We conclude that tubulin expression levels are not tightly regulated in yeast.

Furthermore, even relatively small changes in α-tubulin expression levels can

substantially influence resistance and sensitivity to microtubule inhibitors. These

findings underscore the importance of recognizing subtle (<2-fold) expression

changes, as they have the capacity for causing strong phenotypes.

Not all of the genes that contributed to benomyl resistance have functions related to

tubulin expression. We identified two components of the spindle position checkpoint

(*MCK1*-chr. 14, *SWR1*-chr. 4), implicating this checkpoint in mitigating the effects of

microtubule-targeting drugs. We also identified a drug exporter (*SNQ2*-chr. 4) and a

microtubule stabilizer (*BIM1*-chr. 5). Importantly, the genes that we identified for

resistant chromosome region were not always the most obvious candidates

(Supplemental Fig. S7). *TOF1*-chr. 14 and *SAP1*-chr. 5 have not previously been

associated with microtubule functions, but both contribute to benomyl resistance on

their respective aneuploid chromosomes. These results highlight the need for

unbiased approaches to determine the genes responsible for aneuploidy phenotypes,

as phenotypes resulting from mild changes in expression could greatly differ from

those that come from inhibition/deletion.

While mild overexpression of specific genes can confer drug resistance, the

overexpression of many genes across an entire chromosome greatly increases

aneuploidy-induced cellular stress (Torres et al. 2007; Sheltzer et al. 2011; Oromendia

et al. 2012; Thorburn et al. 2013; Beach et al. 2017; reviewed in Li and Zhu 2022).

This stress presents potential targets to selectively eliminate aneuploid cells (Cohen-

Sharir et al. 2021; Ippolito et al. 2024). However, the potential to leverage aneuploidy-

based sensitivities requires a deeper understanding of the underlying mechanisms

driving these vulnerabilities. 78% of the strains in the chromosome gain collection were

sensitive to hygromycin B and 66% were sensitive to heat stress (Supplemental Fig.

S2C). These results provide additional support for decreased ribosome function and

proteotoxic stress as hallmarks of aneuploidy (Oromendia et al. 2012; Terhorst et al.
2020). By contrast, inducing DNA damage via MMS did not result in large scale
sensitivity, suggesting aneuploidy-induced genome instability may be more
chromosome specific. Despite the general sensitivity to hygromycin, we identified a
few strong examples of aneuploidy-driven hygromycin resistance. Additionally,
sensitivity of one chromosome arm can be masked by resistance of the other arm
(Supplemental Fig. S4B, C). Together, these results demonstrate that regions of the
genome that contribute to sensitivity or resistance can fully or partially counteract each
other. Sensitivity and negative selection have been predicted to be strong drivers of
cancer karyotype evolution, potentially even more so than positive selection (Jubran
et al. 2024).

Given the high prevalence of aneuploidy in cancer, it is an attractive target for
identifying exploitable vulnerabilities. Initially, focus has been placed on vulnerabilities
resulting from aneuploidy in general, as opposed to ones created by specific
chromosomes. Currently, one of the leading candidates for targeting a general
vulnerability of aneuploidy is proteotoxic stress, as it has been found in both yeast and
human cells with a wide variety of aneuploid karyotypes (Tang et al. 2011; Oromendia
et al. 2012; Stingele et al. 2012). Our results showing a general sensitivity to heat
stress across dozens of engineered aneuploid strains would agree with proteotoxic
stress as a good candidate and may additionally suggest that localized hyperthermia
therapy (reviewed in Hannon et al. 2021) may be especially effective on highly
aneuploid tumors. We also observed a general sensitivity to the ribosome inhibitor
hygromycin B for the majority of aneuploid strains, implicating protein synthesis as an
additional candidate for targeting aneuploid cells. However, specific chromosome
disomies were not sensitive to the drug and could even provide resistance. These
chromosome-specific differences suggest that even for effective general aneuploidy-
targeting treatments, certain karyotypes may not respond well and that linking cancer
karyotypes to drug responses could greatly improve treatment effectiveness and
choice. Additional efforts have recently been made to identify vulnerabilities of specific
aneuploid chromosomes (Cai et al. 2016; Huth et al. 2023; Girish et al. 2023). Our
work also supports this strategy, as we identified frequent chromosome-specific drug
sensitivities.

Material and methods

Yeast strains and media

All yeast strains and plasmids used in this study are listed in Supplemental Table S2, S3. All yeast strains are derivatives of BY4741 and BY4742. All yeast strains were cultured in complete synthetic media (CMS) lacking histidine, uracil, lysine, and leucine and cultured at 30 °C, unless otherwise stated. For 5-FOA (Chempur, Karlsruhe, Germany, 220141- 70-8) selection, a concentration of 1 mg/mL was used. Growth assays were always conducted in yeast extract/peptone containing 40 µg/ml adenine-HCl (YPA) and 2% glucose (YPAD) in liquid cultures and plate-based assays. For high-throughput drug resistance screens, 17.5 µg/ml benomyl (Sigma-Aldrich), 40 µg/ml hygromycin B (Carl Roth), 0.02 % MMS (Sigma-Aldrich) or 5 nM rapamycin (Santa Cruz Biotechnology) were added to YPAD agar plates. YPAD + DMSO agar plates were used as controls for drugs dissolved in DMSO. Genetic manipulations such as insertions and gene deletions were conducted as previously described (Longtine et al. 1998). Strains inducible for aneuploidy were generated using a conditional centromere, as described in Anders et al. (2009). Single disomies and double disomies were engineered as described in (Ravichandran et al. 2018). Each disomy was generated independently with two different pairs of auxotrophic markers (*URA3* and *HIS3*; *LYS2* and *LEU2*). For engineering monosomies, one copy of the endogenous centromere was replaced by P_{GAL1}-*CEN3 URA3*. Flow cytometry of selected strains was used to detect potential changes in ploidy and exclude strains that spontaneously diploidized.

PCR-mediated partial chromosome deletions

PCR-mediated partial chromosome deletions were implemented as described in Linder et al. (2017). To generate a deletion, a 500-800 bp fragment homologous to the intended cut-site was combined with a resistance marker (*kanMX* or *hphNT1*) and a telomeric seed sequence. The telomeric seed sequence (6x 5'CCCCAA3') was added at the 3' end of the primer binding the resistance cassette. The homology sequence was PCR amplified from genomic DNA with an added 30 bp homology sequence to the resistance cassette. Fragments were combined using overlap extension PCR.

High-throughput growth measurements

For monosomies, chromosome missegregation was induced immediately before plating.

Strains were grown for 18h in 2 ml deep-well plates in 200 µl YPAD at 30 °C (shaking). 4 µl

were transferred to 150 µl of YPA plus 1% raffinose (YPAR) and grown to log phase (~4h).

50 µl of YPA medium containing 4% galactose were added and grown for 3h to induce

chromosome missegregation. Samples were plated on 5-FOA agar plates using a ROTOR

HDA robot for high-throughput plating (Singer Instruments, Roadwater, UK). Images were

taken after 24, 48 and 72h. The chromosome gain collection was grown in 2 ml deep-well

plates in 200 µl CSM lacking histidine, uracil, leucine and lysine at 30 °C. After 18h,

cultures were diluted 1:10 in 1 × PBS and plated with the ROTOR HDA onto condition

specific YPAD plates. Strains were grown at 30 °C (if not indicated otherwise) and images

were taken at different time points from 24 to 72 h.

Doubling time measurements

Overnight cultures were grown in CSM lacking histidine, uracil, leucine and lysine,

diluted in YPAD and grown for 4h to mid log-phase. Cultures were diluted to an optical

density at 600 nm of 0.02 in 200 µl YPAD in 96-well plates. OD measurements were

taken every 15 min for 24 h using a Synergy S1 plate reader (BioTek Instruments, Inc.,

Luzern, Switzerland).

Serial dilutions

Strains were grown over night in CSM lacking histidine, uracil, leucine and lysine.

Cultures were diluted in fresh media (OD ~ 0.05) and grown to log phase (OD ~0.45),

normalized by OD, diluted in 1 x PBS (1 : 10) and plated on condition specific agar

plates.

High-throughput growth assay quantification and analysis

Images of yeast colonies on plates were quantified using ImageJ (National Institute of

Health). Integrated densities of each spot area and surrounding area were quantified.

Background subtraction was done for each spot area individually by subtracting the

surrounding intensity from the colony intensity (Supplemental data file). For comparing

growth of chromosome loss strains to chromosome gain strains, 10 representative

chromosome loss strains were spotted on YPAD after selection on 5-FOA plates and

measured for colony intensity. These values were then used to adjust the growth

measurements for the other chromosome loss strains. Growth rate prediction of double

disomies and double monosomies in high throughput experiments was predicted by

multiplying the growth of the two individual single disomies/monosomies. This was done

for each treatment individually. Determination of genetic interactions between disomic

chromosomes were calculated from dilution series by adding the measurements of the

individual single disomies (DisA + DisB - 1). Growth rates greater/smaller than additive

prediction +/- 30% were classified as synergistic genetic interaction. The same method

was used to identify genetic interactions between chromosome arms.

qPCR

For DNA extraction, cells were taken from plates and lysed in 20 µl of 0.02 M NaOH for

10 min at 100 °C. Lysates were centrifuged for 10 min on a mini-centrifuge (Thermo Fisher

Scientific) to pellet cellular debris. Each 20 µl reaction contained 10 µl of Luna qPCR

Mastermix (New England BioLabs Inc.), 1 µl of lysate and 140 nM of each primer.

Measurements were made on a RealPlex2 qPCR machine (Eppendorf). Cycling

conditions were: activation/denaturation for 5 min at 95 °C followed by 40 cycles of

15 sec at 95 °C, 1 min at 60 °C with a subsequent fluorescence measurement. Melting

curves were generated to determine primer specificity. *Ct* values were determined using

the automatic thresholding of the RealPlex2 software. Chromosome copy numbers were

calculated using a slightly modified ΔΔCt method (Schmittgen 2001). The Ct values from

duplicates were averaged and used to obtain the ΔCt, which was raised to the negative

power of 2 to give the fold change. The ratio of fold change of the test strains to that of a

wild-type strain was calculated to obtain the chromosome copy numbers. Karyotyping

primers are in Supplemental Table S5.

Protein extraction and Western blotting

For protein extraction, saturated ON cultures were diluted to an OD of 0.02 into YPAD. Cells were grown to an OD of ~0.6, normalized by OD and harvested. Proteins were extracted by pelleting cells, resuspending them in 100 µl ddH$_2$O, adding 100 µl 0.2 M NaOH and incubating them at room temperature for 5 min. Cells were pelleted and resuspended in 1x sample buffer before incubating them at 95° C for 3 min. Samples were stored at -20°C. Membranes were stained with 0.1 % Ponceau S (Sigma Aldrich) in 5% acetic acid. Total protein amount was detected with the ChemiDocMP imaging system (Bio-Rad). Membranes were washed for 30 min in Tris-buffered saline with 0.01% Tween (TBS-T) with three buffer exchanges, blocked for 1 h in 5% BSA (Sigma-Aldrich) or 5% skim milk (Gerbu Biotechnik GmbH) in TBS-T, depending on the primary antibody (see below). Primary antibodies diluted in their respective blocking buffers were incubated overnight at 4°C on an orbital shaker. Membranes were washed for 30 min in TBS-T with three buffer exchanges before and after secondary antibody incubation. Horseradish peroxidase (HRP)-conjugated antimouse IgG secondary antibody (1:10,000; Cell Signaling Technology 7076) diluted in the respective blocking buffers was incubated for 1 h at room temperature. Chemiluminescence was developed using the Amersham ECL Prime detection reagent (Cytiva) and detected with the ChemiDocMP. Protein bands were quantified using ImageJ and normalized to the total protein amount quantified from the Ponceau S staining. The following primary antibodies were used: mouse anti-Pgk1 (in BSA and skim milk–TBST, respectively; 1:20.000; Thermo Fisher Scientific 22C5D8), anti-α-Tubulin (in skim milk–TBS; 1:20,000; mouse; Sigma-Aldrich T6074), anti-β-Tubulin (in BSA; 1:5.000; mouse; Developmental Studies Hybridoma Bank E7).

Flow Cytometry

Cultures were grown to log-phase, pelleted and resuspended in 300 µl H$_2$O. 700 µl of 100 % EtOH were added during vertexing, and cells were fixed for 1h at room temperature. Cells were pelleted and resuspended in 50 mM sodium citrate, sonicated to disperse cell clumps and treated with 250 µg/mL RNase A (Sigma-Aldrich, R6513) and 1 mg/mL Proteinase K (Sigma-Aldrich, P2308) overnight at 37°C. Cells were resuspended in 50 mM sodium citrate solution containing 1 µM SYTOX green (Thermo Fisher,

10768273). Samples were run on BD LSRFortessa Cell Analyzer equipped with a 15-mW

488-nm laser. Maximum count peaks for fluorescence intensities were calculated using

the BD FACSDiva 8.0.1 software.

Statistical analysis

Statistical analysis was performed using GraphPad Prism software. Details for the

statistical tests used in a particular experiment are reported in the figure legends.

642
643
644

Competing interest statement

The authors declare no competing interests.

Acknowledgements

The authors thank the Campbell and Dammermann laboratories for helpful discussions and

comments. We thank the Kraft lab and Kirk Anders for gifts of strains and plasmids and

Alexander Dammermann and Sean Montgomery for careful reading of the manuscript. We

acknowledge the service of the FACS Facility.

This work was supported by a Max Perutz Labs Fellowship to M.S.K and the Austrian Science

Fund (FWF) grant F 8802-B to C.S.C.

Author Contributions

M.S.K conducted the majority of the experiments. M.S.K., S.F., M.C.R and C.H., generated

the chromosome gain collection. C.H. and S.F. generated the chromosome loss collection.

C.H. performed growth measurements and analysis of the chromosome loss collection. M.S.K.

and C.S.C conceptualized the research, analyzed the data, and wrote the manuscript. All

authors reviewed and approved the manuscript.

References

Adell MAY, Klockner TC, Höfler R, Wallner L, Schmid J, Markovic A, Martyniak A,
Campbell CS. 2023. Adaptation to spindle assembly checkpoint inhibition
through the selection of specific aneuploidies. *Genes Dev* **37**: 171–190.

Anders KR, Kudrna JR, Keller KE, Kinghorn BAA, Miller EM, Pauw D, Peck AT, Shellooe CE,
Strong IJT. 2009. A strategy for constructing aneuploid yeast strains by transient
nondisjunction of a target chromosome. *BMC Genet* **10**: 1–11.

Barney JB, Chandrashekarappa DG, Soncini SR, Schmidt MC. 2021. Drug resistance in
diploid yeast is acquired through dominant alleles, haploinsufficiency, gene
duplication and aneuploidy. *PLOS Genet* **17**: e1009800.

Beach RR, Ricci-Tam C, Brennan CM, Moomau CA, Hsu P, Hua B, Silberman RE,
Springer M, Amon A. 2017. Aneuploidy Causes Non-genetic Individuality. *Cell*
169: 229-242.e21.

Beaupere C, Dinatto L, Wasko BM, Chen RB, VanValkenburg L, Kiflezghi MG, Lee MB,
Promislow DEL, Dang W, Kaeberlein M, et al. 2018. Genetic screen identifies
adaptive aneuploidy as a key mediator of ER stress resistance in yeast. *Proc Natl*
Acad Sci **115**: 9586–9591.

Bonney ME, Moriya H, Amon A. 2015. Aneuploid proliferation defects in yeast are not
driven by copy number changes of a few dosage-sensitive genes. *Genes Dev* **29**:
898–903.

Cai Y, Crowther J, Pastor T, Abbasi Asbagh L, Baietti MF, De Troyer M, Vazquez I, Talebi A,
Renzi F, Dehairs J, et al. 2016. Loss of Chromosome 8p Governs Tumor
Progression and Drug Response by Altering Lipid Metabolism. *Cancer Cell* **29**:
751–766.

Caydasi AK, Khmelinskii A, Darieva Z, Kurtulmus B, Knop M, Pereira G. 2023. SWR1
chromatin remodeling complex prevents mitotic slippage during spindle position
checkpoint arrest. *Mol Biol Cell* **34**: ar11.

Chen G, Bradford WD, Seidel CW, Li R. 2012. Hsp90 stress potentiates rapid cellular
adaptation through induction of aneuploidy. *Nature* **482**: 246–250.

Clarke MN, Marsoner T, Adell MAY, Ravichandran MC, Campbell CS. 2023. Adaptation to
high rates of chromosomal instability and aneuploidy through multiple pathways
in budding yeast. *EMBO J* **42**: e111500.

Cohen-Sharir Y, McFarland JM, Abdusamad M, Marquis C, Bernhard SV, Kazachkova M,
Tang H, Ippolito MR, Laue K, Zerbib J, et al. 2021. Aneuploidy renders cancer
cells vulnerable to mitotic checkpoint inhibition. *Nature* **590**: 486–491.

Girish V, Lakhani AA, Thompson SL, Scaduto CM, Brown LM, Hagenson RA, Sausville EL,
Mendelson BE, Kandikuppa PK, Lukow DA, et al. 2023. Oncogene-like addiction
to aneuploidy in human cancers. *Science* **381**: eadg4521.

Gresham D, Desai MM, Tucker CM, Jenq HT, Pai DA, Ward A, DeSevo CG, Botstein D,
Dunham MJ. 2008. The Repertoire and Dynamics of Evolutionary Adaptations to
Controlled Nutrient-Limited Environments in Yeast. *PLoS Genet* **4**: e1000303.

Hannon G, Tansi FL, Hilger I, Prina-Mello A. 2021. The Effects of Localized Heat on the
Hallmarks of Cancer. *Adv Ther* **4**: 2000267.

Hose J, Escalante LE, Clowers KJ, Dutcher HA, Robinson D, Bouriakov V, Coon JJ,
Shishkova E, Gasch AP. 2020. The genetic basis of aneuploidy tolerance in wild
yeast. *eLife* **9**.

Huth T, Dreher EC, Lemke S, Fritzsche S, Sugiyanto RN, Castven D, Ibberson D, Sticht C,
Eiteneuer E, Jauch A, et al. 2023. Chromosome 8p engineering reveals increased
metastatic potential targetable by patient-specific synthetic lethality in liver
cancer. *Sci Adv* **9**: eadh1442.

Ippolito MR, Martis V, Martin S, Tijhuis AE, Hong C, Wardenaar R, Dumont M, Zerbib J,
Spierings DCJ, Fachinetti D, et al. 2021. Gene copy-number changes and
chromosomal instability induced by aneuploidy confer resistance to
chemotherapy. *Dev Cell* **56**: 2440-2454.e6.

Ippolito MR, Zerbib J, Eliezer Y, Reuveni E, Viganò S, Taglietti S, Scorzoni S, Gianotti S,
Martin S, Muenzner J, et al. 2024. Increased RNA and protein degradation is
required for counteracting transcriptional burden and proteotoxic stress in
human aneuploid cells. *Cancer Discov* **14**.

Jubran J, Slutsky R, Rozenblum N, Rokach L, Ben-David U, Yeger-Lotem E. 2024.
Machine-learning analysis reveals an important role for negative selection in
shaping cancer aneuploidy landscapes. *Genome Biol* **25**: 95.

Kaboli S, Miyamoto T, Sunada K, Sasano Y, Sugiyama M, Harashima S. 2016. Improved
stress resistance and ethanol production by segmental haploidization of the
diploid genome in Saccharomyces cerevisiae. *J Biosci Bioeng* **121**: 638–644.

Klockner TC, Campbell CS. 2024. Selection forces underlying aneuploidy patterns in
cancer. *Mol Cell Oncol* **11**: 2369388.

Li R, Zhu J. 2022. Effects of aneuploidy on cell behaviour and function. *Nat Rev Mol Cell*
Biol **23**: 250–265.

Linder RA, Greco JP, Seidl F, Matsui T, Ehrenreich IM. 2017. The Stress-Inducible
Peroxidase *TSA2* Underlies a Conditionally Beneficial Chromosomal Duplication
in *Saccharomyces cerevisiae*. *G3 GenesGenomesGenetics* **7**: 3177–3184.

Liu X, Wang X, Yang X, Liu S, Jiang L, Qu Y, Hu L, Ouyang Q, Tang C. 2015. Reliable cell
cycle commitment in budding yeast is ensured by signal integration. *eLife* **4**:
e03977.

Loegler V, Friedrich A, Schacherer J. 2024. Overview of the *Saccharomyces cerevisiae*
population structure through the lens of 3,034 genomes. *G3 Genes Genomes*
Genet jkae245.

Longtine MS, McKenzie A, Demarini DJ, Shah NG, Wach A, Brachat A, Philippsen P,
Pringle JR. 1998. Additional modules for versatile and economical PCR-based
gene deletion and modification in Saccharomyces cerevisiae. *Yeast Chichester*
Engl **14**: 953–61.

Lukow DA, Sausville EL, Suri P, Chunduri NK, Wieland A, Leu J, Smith JC, Girish V, Kumar
AA, Kendall J, et al. 2021. Chromosomal instability accelerates the evolution of
resistance to anti-cancer therapies. *Dev Cell* **56**: 2427-2439.e4.

Macaluso F, Bos T, Chiroli E, Bonaiuti P, Apuan JC, Gross F, Pompei S, Rice LM, Ciliberto
A. 2025. Evolutionary adaptation to hyperstable microtubules selectively targets
tubulins and is empowered by the spindle assembly checkpoint. *Cell Rep* **44**:
115323.

Mitelman Database of Chromosome Aberrations and Gene Fusions in Cancer (2025).
Mitelman F, Johansson B and Mertens F (Eds.), https://mitelmandatabase.isb-
cgc.org

Morard M, Macías LG, Adam AC, Lairón-Peris M, Pérez-Torrado R, Toft C, Barrio E. 2019.
Aneuploidy and Ethanol Tolerance in Saccharomyces cerevisiae. *Front Genet* **10**:
82.

Muenzner J, Trébulle P, Agostini F, Zauber H, Messner CB, Steger M, Kilian C, Lau K,
Barthel N, Lehmann A, et al. 2024. Natural proteome diversity links aneuploidy
tolerance to protein turnover. *Nature* **630**: 149–157.

Oromendia AB, Dodgson SE, Amon A. 2012. Aneuploidy causes proteotoxic stress in
yeast. *Genes Dev* **26**: 2696–2708.

Pan X, Yuan DS, Xiang D, Wang X, Sookhai-Mahadeo S, Bader JS, Hieter P, Spencer F,
Boeke JD. 2004. A Robust Toolkit for Functional Profiling of the Yeast Genome.
Mol Cell **16**: 487–496.

Pavani M, Bonaiuti P, Chiroli E, Gross F, Natali F, Macaluso F, Póti Á, Pasqualato S,
Farkas Z, Pompei S, et al. 2021. Epistasis, aneuploidy, and functional mutations
underlie evolution of resistance to induced microtubule depolymerization.
EMBO J **40**: e108225.

Pavelka N, Rancati G, Zhu J, Bradford WD, Saraf A, Florens L, Sanderson BW, Hattem
GL, Li R. 2010. Aneuploidy confers quantitative proteome changes and
phenotypic variation in budding yeast. *Nature* **468**: 321–325.

Peter J, De Chiara M, Friedrich A, Yue J-X, Pflieger D, Bergström A, Sigwalt A, Barre B,
Freel K, Llored A, et al. 2018. Genome evolution across 1,011 Saccharomyces
cerevisiae isolates. *Nature* **556**: 339–344.

Rancati G, Pavelka N, Fleharty B, Noll A, Trimble R, Walton K, Perera A, Staehling-
779   Hampton K, Seidel CW, Li R. 2008. Aneuploidy Underlies Rapid Adaptive
780   Evolution of Yeast Cells Deprived of a Conserved Cytokinesis Motor. *Cell* **135**:
781   879–893.

782   Rathi S, Polat I, Pereira G. 2022. The budding yeast GSK-3 homologue Mck1 is an
783   essential component of the spindle position checkpoint. *Open Biol* **12**: 220203.

784   Ravichandran MC, Fink S, Clarke MN, Hofer FC, Campbell CS. 2018. Genetic
785   interactions between specific chromosome copy number alterations dictate
786   complex aneuploidy patterns. *Genes Dev* **32**: 1485–1498.

787   Rojas J, Hose J, Dutcher HA, Place M, Wolters JF, Hittinger CT, Gasch AP. 2024.
788   Comparative modeling reveals the molecular determinants of aneuploidy fitness
789   cost in a wild yeast model. *Cell Genomics* **4**: 100656.

790   Ryu H-Y, Wilson NR, Mehta S, Hwang SS, Hochstrasser M. 2016. Loss of the SUMO
791   protease Ulp2 triggers a specific multichromosome aneuploidy. *Genes Dev* **30**:
792   1881–1894.

793   Schibler A, Koutelou E, Tomida J, Wilson-Pham M, Wang L, Lu Y, Cabrera AP, Chosed RJ,
794   Li W, Li B, et al. 2016. Histone H3K4 methylation regulates deactivation of the
795   spindle assembly checkpoint through direct binding of Mad2. *Genes Dev* **30**:
796   1187–1197.

797   Schmittgen TD. 2001. Real-Time Quantitative PCR. *Methods* **25**: 383–385.

798   Selmecki A, Forche A, Berman J. 2006. Aneuploidy and Isochromosome Formation in
799   Drug-Resistant *Candida albicans*. *Science* **313**: 367–370.

800   Selmecki AM, Dulmage K, Cowen LE, Anderson JB, Berman J. 2009. Acquisition of
801   Aneuploidy Provides Increased Fitness during the Evolution of Antifungal Drug
802   Resistance. *PLoS Genet* **5**: e1000705.

803   Sheltzer JM, Blank HM, Pfau SJ, Tange Y, George BM, Humpton TJ, Brito IL, Hiraoka Y,
804   Niwa O, Amon A. 2011. Aneuploidy Drives Genomic Instability in Yeast. *Science*
805   **333**: 1026–1030.

806   Shukla A, Nguyen THM, Moka SB, Ellis JJ, Grady JP, Oey H, Cristino AS, Khanna KK,
807   Kroese DP, Krause L, et al. 2020. Chromosome arm aneuploidies shape tumour
808   evolution and drug response. *Nat Commun* **11**: 449.

809   Stingele S, Stoehr G, Peplowska K, Cox J, Mann M, Storchova Z. 2012. Global analysis of
810   genome, transcriptome and proteome reveals the response to aneuploidy in
811   human cells. *Mol Syst Biol* **8**: 608.

812   Su XA, Ma D, Parsons JV, Replogle JM, Amatruda JF, Whittaker CA, Stegmaier K, Amon A.
813   2021. *RAD21* is a driver of chromosome 8 gain in Ewing sarcoma to mitigate
814   replication stress. *Genes Dev* **35**: 556–572.

Tang Y-C, Williams BR, Siegel JJ, Amon A. 2011. Identification of Aneuploidy-Selective
Antiproliferation Compounds. *Cell* **144**: 499–512.

Terhorst A, Sandikci A, Keller A, Whittaker CA, Dunham MJ, Amon A. 2020. The
environmental stress response causes ribosome loss in aneuploid yeast cells.
Proc Natl Acad Sci U S A **117**: 17031–17040.

Thorburn RR, Gonzalez C, Brar GA, Christen S, Carlile TM, Ingolia NT, Sauer U,
Weissman JS, Amon A. 2013. Aneuploid yeast strains exhibit defects in cell
growth and passage through START. *Mol Biol Cell* **24**: 1274–1289.

Tong AHY, Lesage G, Bader GD, Ding H, Xu H, Xin X, Young J, Berriz GF, Brost RL, Chang
M, et al. 2004. Global Mapping of the Yeast Genetic Interaction Network. *Science*
303: 808–813.

Torres EM, Sokolsky T, Tucker CM, Chan LY, Boselli M, Dunham MJ, Amon A. 2007.
Effects of Aneuploidy on Cellular Physiology and Cell Division in Haploid Yeast.
Science **317**: 916–924.

Trakala M, Aggarwal M, Sniffen C, Zasadil L, Carroll A, Ma D, Su XA, Wangsa D, Meyer A,
Sieben CJ, et al. 2021. Clonal selection of stable aneuploidies in progenitor cells
drives high-prevalence tumorigenesis. *Genes Dev* **35**: 1079–1092.

Voordeckers K, Kominek J, Das A, Espinosa-Cantú A, De Maeyer D, Arslan A, Van Pee M,
Van Der Zande E, Meert W, Yang Y, et al. 2015. Adaptation to High Ethanol
Reveals Complex Evolutionary Pathways. *PLOS Genet* **11**: e1005635.

Wangsa D, Braun R, Stuelten CH, Brown M, Bauer KM, Emons G, Weston LA, Hu Y, Yang
HH, Vila-Casadesús M, et al. 2019. Induced Chromosomal Aneuploidy Results in
Global and Consistent Deregulation of the Transcriptome of Cancer Cells.
Neoplasia **21**: 721–729.

Watson EV, Lee JJ-K, Gulhan DC, Melloni GEM, Venev SV, Magesh RY, Frederick A, Chiba
K, Wooten EC, Naxerova K, et al. 2024. Chromosome evolution screens
recapitulate tissue-specific tumor aneuploidy patterns. *Nat Genet* **56**: 900–912.

Weaver BA, Cleveland DW. 2006. Does aneuploidy cause cancer? *Curr Opin Cell Biol*
18: 658–667.

Yona AH, Manor YS, Herbst RH, Romano GH, Mitchell A, Kupiec M, Pilpel Y, Dahan O.
2012. Chromosomal duplication is a transient evolutionary solution to stress.
Proc Natl Acad Sci **109**: 21010–21015.

Zhang K, Zhang L-J, Fang Y-H, Jin X-N, Qi L, Wu X-C, Zheng D-Q. 2016. Genomic
structural variation contributes to phenotypic change of industrial bioethanol
yeast *Saccharomyces cerevisiae*. *FEMS Yeast Res* **16**: fov118.

Zhou X, Hilk A, Solis NV, Scott N, Beach A, Soisangwan N, Billings CL, Burrack LS, Filler
SG, Selmecki A. 2024. Single-cell detection of copy number changes reveals

dynamic mechanisms of adaptation to antifungals in Candida albicans. *Nat*
Microbiol **9**: 2923–2938.

Figure 1

Figure 1: Strong genetic interactions between aneuploid chromosomes are rare
(A) Schematic overview of the chromosome gain and chromosome loss collections. (B) Quantified growth of the chromosome gain and chromosome loss collections. The averages of four independent high-throughput growth assays were normalized to WT. Colony intensities were quantified after 24h on YPAD plates for chromosome gains, and 72h on plates containing 5-Fluoroorotic Acid (5-FOA) for chromosome losses. Single disomies/monosomies are depicted in grey, double disomies/monosomies in black. (C) Correlation between growth of the double disomies and growth predictions based on the single disomies (p < 0.0001). Growth was predicted by calculating the product of single disomy growth. r^2 values are from simple linear regression and p-values are from *F*-tests. (D) Quantification of a serial dilution on 5-FOA plates after 72h normalized to a diploid WT.

Figure 2

Figure 2: The strongest drug resistance phenotypes come from complex aneuploidy
(A) Schematic of the work-flow for plate-based high throughput growth assays under selective conditions. Cells were grown in liquid cultures before being transferred to condition specific agar plates. Images for colony intensities were taken after 24, 36, or 48h. (B) Averaged growth of two independent high-throughput growth assays. Control plates were quantified after 24h; hygromycin (40 µg/ml) and rapamycin (5 nM) plates after 48h; and MMS (0.02%) and benomyl (17.5 µg/ml) plates after 36h. All quantifications were normalized to a haploid WT. (C) Growth measurements of single disomies (Single) and double disomies (Double) across different treatments. The YPAD control is an average of YPAD and YPAD + DMSO plate growth. The dashed line indicates WT growth.

Figure 3

Figure 3: Benomyl resistance correlates with alpha-tubulin expression in aneuploid strains
(A) 10-fold serial dilution analysis of benomyl resistant single disomies on control (YPAD + DMSO, 24h) and treated (YPAD + 20 µg/ml benomyl, 48h) plates. (B) Western blot analysis of α-tubulin expression levels of single disomies. *Pgk1* is shown as a loading control. (C) Western blot quantifications of α-tubulin (n = 3) and β-tubulin (n = 1-2) expression levels for all single disomies. Strains were grown in YPAD and harvested in log-phase. Quantified bands were normalized to the Ponceau stained membrane and the haploid WT. Relative growth on benomyl denotes the average growth of single disomies from two independent high throughput colony size assays. r^2 values are from simple linear regression and p-values are from *F*-tests.

Figure 4

Figure 4: Identification of chromosome arms necessary for aneuploidy-based drug resistance.

(A) Schematic of two models for the genetic basis behind aneuploidy driven drug resistance. (B) Method of chromosome arm deletion. Chromosome arms are deleted by homologous recombination with a 500 bp homology sequence connected to a resistance marker and a telomeric seed sequence. (C) Doubling times of whole chromosome and single arm disomies for chromosomes 4, 5, 10 and 14. ORFs depict the remaining aneuploid ORFs after chromosome arm deletions. (D) 10-fold dilution series and quantification of whole chromosome and single arm disomies. Control plates (YPAD + DMSO) were imaged after 24h and treated plates (YPAD + 25 µg/ml benomyl) after 48h. * Indicates a lower benomyl concentration (20 µ/ml benomyl). The second dilution was quantified and normalized to WT, with the exceptions of chromosome 13 (third dilution) and chromosome 12 (first dilution). (E) Quantification of 10-fold serial dilutions treated with MMS (0.04 %), rapamycin (10 nM) or hygromycin (60 µg/ml). Relative growth was calculated as the ratio of normalized growth on treated plates (48h) to normalized growth on control plates (24h). Representative examples for single arm sensitivity, combinatorial sensitivity, compensatory sensitivity, resistance, and no interactions are shown. Images and quantifications of all 10-fold serial dilutions are depicted in Supplemental Fig. S4A, B.

Figure 5

Figure 5: Most aneuploidy-based drug resistance phenotypes come from the combined effects of multiple genes.

(A) 10-fold serial dilution and quantification of partial chromosome arm deletions of the right arm of chromosome 4. For control plates (YPAD + DMSO) the 3rd spot was quantified after 24h and for treated plates (YPAD + 25µg/ml benomyl) the 2nd and 3rd spots were quantified and averaged after 48h. Growth was normalized to a haploid WT. (B) 10-fold serial dilution and quantification of partial chromosome arm deletions and heterozygous deletions of genes contributing to benomyl resistance. For treated plates (YPAD + 25µg/l benomyl) the 2nd spot was quantified after 48h and normalized to the haploid wt. (C) Schematic depiction of benomyl resistant and sensitive chromosome regions. Identified resistance genes are indicated in orange at their approximate chromosomal location.

Figure 6

Figure 6: Synergistic effects contribute to the strongest aneuploidy phenotypes.
(A) Single disomy growth on benomyl compared to growth predictions based on single arm disomies. Growth was quantified from 10-fold serial dilutions in Figure 4D. The area highlighted in magenta indicates additive genetic interactions (+/-30% of predicted growth). Disomies that fall below this threshold are considered a negative synergistic interaction and those above are considered a positive synergistic interaction. (B) 10-fold serial dilution of single and double disomies. Control plates (YPAD + DMSO) were imaged after 24h and treated plates (YPAD + 30 µg/ml benomyl) after 48h. (C) 10-fold serial dilutions of benomyl resistant *set1*Δ haploid and two synergistically interacting double disomies (2&4, 2&13). Benomyl plates contained 30 µg/ml benomyl. (D) Quantification of the 10-fold serial dilutions of the double disomies on plates with 30 µg/ml benomyl compared to predicted growth based on the additive resistance of the single disomies (for images, see Sup. Figure 8A). Combinations in the highlighted area (magenta) are additive genetic interactions (+/-30% of predicted growth). Disomy combinations that fall below this threshold are considered a negative synergistic interaction and those above are considered a positive synergistic interaction.