Two-Stage CD8+ CAR T-Cell Differentiation in Patients with Large B-Cell Lymphoma

Guoshuai Cao[1,#], Yifei Hu[2,#], Tony Pan[1,#], Erting Tang[1], Nick Asby[1], Thomas Althaus[6], Jun

Wan[7], Peter A. Riedell[4,5,6], Michael R. Bishop[5,6], Justin P. Kline[3,4,5,6], Jun Huang[1,3,4,*]

[1]Pritzker School of Molecular Engineering, University of Chicago, Chicago, IL 60637, USA

[2]Pritzker School of Medicine, University of Chicago, Chicago, IL 60637, USA

[3]Committee on Immunology, University of Chicago, Chicago, IL 60637, USA

[4]Committee on Cancer Biology, University of Chicago, Chicago, IL 60637, USA

[5]Department of Medicine, University of Chicago, Chicago, IL 60637, USA

[6]The David and Etta Jonas Center for Cellular Therapy, University of Chicago, Chicago, IL

60637, USA

[7]Department of Medical and Molecular Genetics, Indiana University School of Medicine,

Indianapolis, IN 46202, USA

[#]These authors contributed equally.

*Correspondence to: huangjun@uchicago.edu

SUMMARY

Chimeric antigen receptor (CAR) T-cell therapy has expanded therapeutic options for patients

with diffuse large B-cell lymphoma (DLBCL). However, progress in improving clinical outcomes

has been limited by an incomplete understanding of CAR T-cell differentiation in patients. To

comprehensively investigate CAR T-cell differentiation in vivo, we performed single-cell, multi-

modal, and longitudinal analyses of CD28-costimulated CAR T cells from infusion product and

peripheral blood (day 8-28) of patients with DLBCL who were successfully treated with

axicabtagene ciloleucel. Here, we show that $CD8^+$ CAR T cells undergo two distinct waves of

clonal expansion. The first wave is dominated by CAR T cells with an exhausted-like effector

memory phenotype during the peak expansion period (day 8-14). The second wave is dominated

by CAR T cells with a terminal effector phenotype during the post-peak persistence period (day

21-28). Importantly, the two waves have distinct ontogeny and are biologically uncoupled.

Furthermore, lineage tracing analysis via each CAR T cell's endogenous TCR clonotype

demonstrates that the two waves originate from different effector precursors in the infusion

product. Precursors of the first wave exhibit more effector-like signatures, whereas precursors of

the second wave exhibit more stem-like signatures. These findings suggest that pre-infusion

heterogeneity mediates the two waves of in vivo clonal expansion. Our findings provide evidence

against the intuitive idea that the post-peak contraction in CAR abundance is solely apoptosis or

extravasation of short-lived CAR T cells from peak expansion. Rather, our findings demonstrate

that CAR T-cell expansion and persistence are mediated by clonally, phenotypically, and

ontogenically distinct CAR T-cell populations that serve complementary clinical purposes.

Key Words: large B-cell lymphoma, chimeric antigen receptor (CAR), axicabtagene ciloleucel,

CD28-costimulated CAR, CAR T-cell differentiation, two-stage differentiation

INTRODUCTION

Diffuse large B-cell lymphoma (DLBCL), the most common non-Hodgkin's lymphoma in the United States, is characterized by diffusely proliferating and malignant B cells at nodal or extranodal sites.[1] Although up-front chemoimmunotherapy is often curative, patients with DLBCL that is refractory to up-front treatment or relapse following remission (r/r DLBCL) have limited treatment options and poor outcomes.[2,3] Effective treatment options were limited until the United States Food and Drug Administration approved autologous CD19-directed chimeric antigen receptor (CAR) T-cell therapies for r/r DLBCL in 2017. Autologous CD19-directed CAR T-cell therapy involves virally transducing a patient's T cells ex vivo with a CD19-directed CAR—an engineered receptor consisting of an extracellular anti-CD19 single-chain variable fragment, a hinge/transmembrane region, and intracellular costimulatory (CD28 or 4-1BB) and activation (CD3ζ) domains. CAR-transduced T cells (i.e., CAR T cells) are cultured and infused into the patient, where they lyse CD19$^+$ lymphoma cells.[4] CD19-directed CAR T-cell therapy has achieved complete response rates of 40-54%[5–8] for r/r DLBCL, but non-response rates[6,9] and treatment toxicities[10] remain as challenges.[11]

Development of CAR T-cell formulations with higher response rates and fewer toxicities requires a thorough understanding of how CAR T cells differentiate in patients with r/r DLBCL. The pioneering ZUMA-1 trial (NCT02348216) for axicabtagene ciloleucel (autologous CD28-costimulated CAR T cells) demonstrated that peripheral blood CAR T cells expand, contract, and sometimes persist.[5] Greater expansion predicted higher response rates, but also higher likelihood of developing treatment toxicities.[5,12] Longer persistence predicted durable remission and long-term immunosurveillance with leukemias, but its relevance for preventing DLBCL relapse remains obscure.[9,13] Although many factors are associated with expansion and persistence (including CAR design[14,15] and CAR T-cell phenotypes[16–20], among others[21–23]), how and why CAR T cells differentiate into expansive or persistent phenotypes in vivo is still an open question. Addressing

this question requires a single-cell approach that integrates CAR T-cell phenotypes and clonal

kinetics over longitudinal timepoints. However, existing studies have either focused on CAR T-

cell phenotypes[19,24,25] or clonal kinetics[26] without integrating both data modalities, or lacked the

temporal resolution required to decipher cell fates over longer periods.[12,27] Consequently, a

complete and longitudinal understanding of CAR T-cell differentiation in vivo has remained

elusive.

To comprehensively elucidate CAR T-cell differentiation in vivo, we performed single-cell,

multi-modal (paired RNA-seq/CITE-seq/TCR-seq), and longitudinal analyses of CD28-

costimulated CAR T cells from infusion product and peripheral blood of seven patients with r/r

DLBCL who were complete responders under treatment with axicabtagene ciloleucel. Peripheral

blood CAR T cells were sorted using CD19 antigen-tetramers.[28] Importantly, we report that the

CD8[+] CAR T cells observed during the peak expansion period (day 8-14) have distinct clonotypic

repertoires compared to the later CD8[+] CAR T cells observed during the post-peak persistence

period (day 21-28). This discovery led us to analyze how these two CAR T-cell populations differ

with respect to phenotypes, transcriptional profiles, regulatory networks, and infusion product

precursors. Our findings not only offer a finer understanding of CAR T-cell biology in vivo, but also

inform efforts to develop CAR T cells with improved expansion and persistence.

RESULTS

Study design and clinical findings.

To interrogate CD28-costimulated CAR T-cell differentiation in vivo, we longitudinally interrogated

the phenotypes and clonal dynamics of CAR T cells from seven patients (P1-7) who achieved

complete responses under treatment with axicabtagene ciloleucel (**Figure 1a**). Patients were

diagnosed with r/r DLBCL and treated at the University of Chicago Medicine between 2019 and

2021 (table S1). Clinical response was determined by positron emission tomography/computed

tomography imaging 30 days after infusion product administration. Complete response was

defined as no detectable lymphoma (figure S1). To capture phenotypic heterogeneity and clonal

dynamics longitudinally, we performed single-cell multi-modal analyses (paired RNA-seq/CITE-

seq/TCR-seq via the 10x Genomics platform) on infusion product and peripheral blood

biospecimens at three timepoints: peak expansion (T_{exp}, day 8-14), early post-peak persistence

(T_{per1}, day 21), and late post-peak persistence (T_{per2}, day 28) (table S2). The three peripheral

blood timepoints were available for all patients, except P3 (only T_{exp} and T_{per1}). CAR T cells were

sorted from peripheral blood using CD19 antigen-tetramers (representative staining in **Figure 1b**,

figure S2).[28] Compared to CD19 antigen-tetramer-negative T cells, CD19 antigen-tetramer-

positive T cells specifically expressed the CAR transgene, which validates our sorting strategy

(**Figure 1c**).

CD8$^+$ CAR T cells undergo a clonotypic shift between T_{exp} and T_{per}.

To analyze CAR T-cell population dynamics, we tracked CAR abundance (% CAR$^+$ of CD3$^+$ T

cells) in peripheral blood throughout the course of therapy. CAR abundance at peak expansion

ranged from 11% to 73% (**Figure 1d**). Peak expansion occurred at day 8-14, which is consistent

with prior clinical findings.[5] For each timepoint, we quantified proportions of CD8$^+$ and CD4$^+$ T

cells within the total CAR T-cell population by single-cell RNA-seq and CITE-seq. CAR T cells

were predominantly $CD8^+$ across most patients and timepoints (figure S3).

We next analyzed the dynamics of the CAR T-cell clonal repertoire across longitudinal

timepoints (T_{exp} at day 8-14, T_{per1} at day 21, T_{per2} at day 28) using single-cell TCR-seq. Clone

sizes were similar across timepoints (figure S4a). Clonotypes did not overlap between patients

(figure S4b). Repertoire overlap analysis indicated that T_{exp} clonotypes were significantly distinct

from T_{per1} or T_{per2} clonotypes (**Figure 1e-f**, left). In sharp contrast, T_{per1} and T_{per2} clonotypes

overlapped substantially more. As a control, we also analyzed endogenous (non-CAR) T cells

from matched timepoints. Unlike with CAR T cells, endogenous T cells did not show distinctive

clonotypic patterns (**Figure 1e-f**, right), indicating that the shift in clonotypes between T_{exp} and

T_{per} is CAR-specific. This CAR-specific clonotypic shift was consistent across patients (figure

S4b). Moreover, the distinction between T_{exp} and T_{per} clonotypes is driven by $CD8^+$ T cells, and

not by $CD4^+$ T cells (figure S4c). Collectively, these findings indicate that $CD8^+$ CAR T cells

undergo a clonotypic shift between T_{exp} and T_{per}.

$CD8^+$ CAR T cells undergo a phenotypic shift from exhausted-like EM to TE.

Having shown a shift in $CD8^+$ CAR T-cell clonotypes between T_{exp} and T_{per}, we hypothesized that

unique $CD8^+$ CAR T-cell phenotypes dominate T_{exp} and T_{per}. To test this hypothesis, we filtered

$CD8^+$ CAR T cells for Uniform Manifold Approximation and Projection (UMAP) and identified six

T-cell clusters (**Figure 2a**) based on gene and protein markers (**Figure 2b**, expanded marker set

in figure S5a). No cluster was patient-specific (figure S5b). All clusters expressed CAR transgene

and CD8α, validating our sorting and filtering processes, respectively. All clusters expressed

CXCR3, a chemokine receptor that demarks activated T cells.

Examination of gene and protein markers revealed that one of the six clusters represented

proliferating T cells ($MKI67^+TOP2A^+$). The remaining five non-proliferating clusters were

annotated as central memory (CM, $TCF7^+TBX21^-$), effector memory (EM, $TCF7^+TBX21^+$), or

terminal effector (TE, $TCF7^-TBX21^+$) T cells. The $TCF7^+TBX21^-$ CM cluster exhibited markers of

stemness (*IL7R*, high CD127) and minimal markers of effectorness (*GZMB*, *CX3CR1*). The two

$TCF7^+TBX21^+$ EM clusters occupied the lower half of the UMAP and uniquely expressed *CXCR6*,

a chemokine receptor that facilitates trafficking into solid tumors.[29] One of these EM clusters

upregulated markers consistent with early exhaustion (*NR4A2*, *TOX*, *GZMK*, low TIM-3), hence it

was designated "exhausted-like EM". Lastly, the two $TCF7^-TBX21^+$ TE clusters occupied the

upper half of the UMAP and uniquely downregulated *GZMK*. One of these TE clusters upregulated

markers consistent with late exhaustion (*TOX*, *PDCD1*, high TIM-3), hence it was designated

"exhausted-like TE". The other TE cluster was highly clonal (some clone sizes > 100), suggesting

expansion through proliferation (figure S7a). Overall, most CD8$^+$ CAR T cells were $TBX21^+$ EM

or TE, which is consistent with the established link between CD28 costimulation and effector

memory (rather than central memory) differentiation.[14,15]

Phenotypic compositions of CD8$^+$ CAR T cells at T_{exp} and T_{per} were compared. CAR T

cells at T_{exp} were predominantly exhausted-like EM (64%) whereas CAR T cells at T_{per} were

predominantly TE (63% for T_{per1}, 77% for T_{per2}) (**Figure 2c**). These findings were statistically

significant and consistent across all seven patients (**Figure 2d**, figure S5c). Moreover, the large

clone sizes within the T_{per}-specific TE cluster (figure S7a) suggest active TE proliferation at T_{per}.

CAR T cells at T_{per1} were enriched for EM. From T_{per1} to T_{per2}, the EM proportion decreased while

the TE proportion increased, suggesting progressive differentiation from EM to TE. However, the

overall phenotypic compositions at T_{per1} and T_{per2} were more similar than different, which is

concordant with findings from repertoire overlap analysis (**Figure 1e-f**). Moreover, we observed

decreasing proliferating proportions and increasing CM proportions over time (**Figure 2d**), though

this was not always statistically significant. These changing proportions may suggest some CAR

T cells were returning from an activated to a resting phenotype. In conclusion, CD8$^+$ CAR T cells

phenotypically shifted from exhausted-like EM to TE between T_{exp} and T_{per}.

CD4$^+$ CAR T cells maintain a memory phenotype with CAR Treg persistence.

We next performed clustering, annotation, and longitudinal analyses of CD4$^+$ CAR T cells to

investigate phenotypes at T_{exp} and T_{per}. We identified five T-cell clusters, all expressing CAR

transgene and CD4, based on gene and protein markers (**Figure 2e-f**; expanded marker set in

figure S6a). No cluster was patient-specific (figure S6b). All clusters expressed *CXCR3*, indicating

T-cell activation and type 1 helper polarization.

One of the five clusters represented proliferating T cells (*MKI67$^+$TOP2A$^+$*). The remaining

four non-proliferative clusters were annotated as memory (Mem, *TBX21$^-$FOXP3$^-$*), type 1 helper

(Th1, *TBX21$^+$FOXP3$^-$*), or regulatory (Treg, *TBX21$^-$FOXP3$^+$*) T cells. The two *TBX21$^-$FOXP3$^-$*

Mem clusters at the center of the UMAP comprised most of the cells. One of the Mem clusters

upregulated *IRF7* and interferon-stimulated genes (*MX1*, *OAS1*, *ISG15*), indicating response to

type I interferon signaling. This signature suggests dynamic interferon secretion in vivo and is in

concordance with type I interferon's role in memory CD4$^+$ T-cell differentiation.[30] The

TBX21$^+$FOXP3$^-$ Th1 cluster along the lower half of the UMAP upregulated cytolytic genes (*PRF1*,

GNLY, *GZMK*, *NKG7*) and tissue-homing chemokine receptors (*CX3CR1*, *CXCR6*), resembling

the cytotoxic CD4$^+$ CAR T cells described by Melenhorst et al.[19] Lastly, the *TBX21$^-$FOXP3$^+$* Treg

cluster expressed classic Treg markers (*IL2RA*, CD25, low *IL7R*, low CD127) and upregulated

IKZF2, indicating a suppressive phenotype.[31] CAR Tregs inhibit conventional CAR T cells[27,32] and

portend progressive disease.[12,27]

Phenotypic compositions of CD4$^+$ CAR T cells at T_{exp} and T_{per} were compared. CD4$^+$ CAR

T cells predominantly exhibited a Mem phenotype (49-82%) across all timepoints (**Figure 2g**).

This observation was consistent across all patients (figure S6c). Between T_{exp} and T_{per1}, the Mem

proportion significantly decreased while the Th1 proportion significantly increased, indicating that

CD4$^+$ CAR T cells may be increasingly polarized during the early contraction period (**Figure 2h**).

Moreover, across all patients, the Treg proportion steadily increased from T_{exp} (5%) to T_{per1} (11%)

to T_{per2} (21%), indicating that CAR Tregs were maintained in peripheral blood after peak

expansion. The CAR Treg cluster was predominantly non-clonal (figure S7b), suggesting CAR

Tregs were maintained through persistence rather than proliferation. While previous studies have

only investigated CAR Tregs at day 7[12,27], this current study affirms and extends the presence of

CAR Tregs until at least day 28. Importantly, persistence of CAR Tregs points towards their

possible involvement in decreasing acute inflammation and restoring immune homeostasis after

peak expansion. In conclusion, we discovered that CD4$^+$ CAR T cells did not exhibit an abrupt

T_{exp}-to-T_{per} phenotypic shift. Rather, they consistently exhibited a memory phenotype with CAR

Tregs persisting over the course of therapy.

Integration of clonotypic and phenotypic shifts in CD8$^+$ CAR T cells supports a two-stage

differentiation model.

Having observed shifts in both CD8$^+$ CAR T-cell clonotypes and phenotypes between T_{exp} and

T_{per}, we hypothesized the existence of two distinct waves of in vivo clonal expansion. To test this

hypothesis, we linked the clonotype and phenotype of single CD8$^+$ CAR T cells in our dataset

using their cell barcodes as unique indices. After this linking process, we confirmed that the

proportions of T cells with exhausted-like EM (EM-exh) and terminal effector (TE) phenotypes

(**Figure 3a**) are consistent with prior results (**Figure 2c**). A phenotype label was subsequently

assigned to each CD8$^+$ CAR T-cell clone based on its predominant phenotype at each timepoint.

For each patient, we tracked the total abundances of clones sharing a common phenotype label

across timepoints. Notably, the abundance of EM-exh clones at T_{exp} was significantly reduced at

T_{per1} and T_{per2} (**Figure 3b**, left), while the abundance of TE clones at T_{per1} and T_{per2} was

significantly reduced at T_{exp} (**Figure 3b**, middle and right). In addition to redemonstrating the shifts

in CD8$^+$ CAR T-cell clonotypes and phenotypes, these findings support the existence of two

distinct waves of in vivo clonal expansion. (**Figure 3c**).

Next, we evaluated the explanatory power of our hypothesis for two distinct clonal waves.

To minimize noise, we filtered for the top 500 largest clones (mean size 10.8, minimum size 4).

Resulting clones were annotated with the following three definitions, based on our hypothesis for

two waves: "Wave 1" (phenotypically EM-exh with highest abundance at T_{exp}), "Wave 2"

(phenotypically TE with highest abundance at T_{per1}/T_{per2}), or "Other" (not Wave 1 or Wave 2).

Wave 1 and Wave 2 clones together accounted for a significant majority (76%) of the top 500

clones (**Figure 3d**). In accordance with our definitions, Wave 1 clones were maximally abundant

at T_{exp}, while Wave 2 clones were maximally abundant at T_{per} (**Figure 3e**). Wave 1 clones

predominantly exhibited an EM-exh phenotype at T_{exp} and sharply decreased in abundance at

T_{per} (**Figure 3f**, left). In contrast, Wave 2 clones predominantly exhibited a TE phenotype at T_{per}

and sharply decreased in abundance at T_{exp} (**Figure 3f**, right). To investigate whether these

findings are dependent on clone size, we expanded our analysis to the top 3000 largest clones

(mean size 3.7, minimum size 2). Under this filter, Wave 1 and Wave 2 clones together continued

to account for a significant majority (79%) of the top 3000 clones (figure S8a) and reaffirm the

expected clonal and phenotypic dynamics (figure S8b-c). Therefore, our hypothesis for two

distinct waves of in vivo clonal expansion has high explanatory power and is robust to clone size.

In conclusion, the two distinct waves of in vivo clonal expansion strongly substantiate a

two-stage CD8[+] CAR T-cell differentiation model (**Figure 3g**). Under this model, some CAR T

cells (Wave 1) with an exhausted-like EM phenotype expand earlier to dominate the peak

expansion timeframe (T_{exp}), while other CAR T cells (Wave 2) with a TE phenotype expand later

to dominate the post-peak persistence timeframe (T_{per}). Of note, the two-stage differentiation

model uncouples CD8[+] CAR T cells from peak expansion and post-peak persistence by

designating these two waves as separate lineages. Moreover, these findings provide evidence

against the intuitive idea that the post-peak contraction in CAR abundance is solely apoptosis or

extravasation of short-lived CAR T cells from peak expansion. Rather, even as total CAR

abundance contracts after peak expansion, a distinct subset of CAR T-cell clones simultaneously

expands to eventually dominate the peripheral blood CAR T-cell repertoire.

T_{exp}- and T_{per}-specific transcriptional signatures and regulatory networks.

We set out to identify the molecular determinants underlying CD8$^+$ CAR T cells at T_{exp} and T_{per}

using gene set enrichment analysis (GSEA), differential gene expression analysis (DGEA), and

regulatory network analysis. For internal validation, the average expression of timepoint-specific

genes (see Methods) for each sample (patient by timepoint) were hierarchically clustered and

compared via correlation (**Figure 4a**). Samples largely clustered by timepoint, validating the

existence of patient-independent molecular signatures. We then validated our dataset against

external data from Maus et al., which consisted of CD8$^+$ CAR T cells from patients with large B-

cell lymphoma (**Figure 4a-b**, colored in tan and brown, six in total after filtering).[27] Based on their

sample collection timing (day 7), external data from Maus et al. should resemble our samples at

T_{exp}, rather than at T_{per}. Consistent with expectations, external data resembled T_{exp} transcriptomes

on both pseudo-bulk (**Figure 4a**, via correlation) and single-cell (**Figure 4b**, via label transfer)

levels, which increases the external validity of our findings.

To elucidate the immunological processes at each timepoint, CD8$^+$ CAR T-cell

transcriptomes at T_{exp} and T_{per} from our seven patients were compared via GSEA (**Figure 4c** and

table S3 for single-cell method, figure S9a and table S4 for pseudo-bulk method) and DGEA

(**Figure 4d**, figure S9b, tables S5-6). CAR T cells at T_{exp} upregulated gene sets for cell cycling

and apoptosis, consistent with a short-lived phenotype. Upregulation of the cell cycling gene set

remained true even when only comparing the proliferating cell cluster at T_{exp} and T_{per} (figure S9d).

This indicates that, while proliferating cells are present at both T_{exp} and T_{per}, those at T_{exp} are more

proliferative than those at T_{per}. We also observed T_{exp}-specific and cluster-independent

upregulation of genes related to T-cell activation (including *CD69*, *CD44*, *CD74*) and exhaustion

(including *PRDM1*, *CTLA4*, *NR4A2*), implicating antigen-engagement and CAR-mediated

signaling during T_{exp}. On the other hand, CAR T cells at T_{per} upregulated gene sets for cytotoxicity

and immune regulation. Expression of cytotoxicity-related genes (including *KLRB1*, *FCGR3A*,

PRF1, *GZMB*, *NKG7*) was restricted to TE and exhausted-like TE clusters, indicating that this

gene signature arises from T_{per}-specific preponderance of the terminal effector phenotype. We

also observed T_{per}-specific upregulation of regulatory genes, including GTPase immune-

associated proteins (*GIMAP5*, *GIMAP7*) which predict long-term persistence[33], as well as

sphingosine-1-phosphate receptors (*S1PR1*, *S1PR5*) which indicate blood localization[34]. These

signatures suggest that CAR T cells at T_{per} are functional and capable of long-term persistence in

peripheral blood. Lastly, downregulation of genes related to T-cell activation and exhaustion

implicate decreased in vivo antigen load and/or CAR-mediated signaling during T_{per}.

274   Interestingly, some differentially expressed gene signatures between CD8$^+$ CAR T cells

at T_{exp} and T_{per} pertained to cytokine signaling (**Figure 4c-d**, see figure S9c). CAR T cells at T_{exp}

upregulated TNF response genes (including *NFKBIA*, *NFKBIE*, *DUSP4*, *RELB*). TNF can be

secreted by CAR T cells for autocrine signaling[20], predicts efficacious CAR T cells[20], and

contributes to CRS.[35] On the other hand, CAR T cells at T_{per} upregulated type I interferon (IFN-I)

response genes (including *STAT1*, *IFITM1*, *IFITM2*, *IRF7*). Many IFN-I response genes have

known antiviral functions. The temporally specific and cluster-independent upregulation of TNF

and IFN-I response genes indicates a dynamic in vivo cytokine environment during the CAR T-

cell immune response.

283   Next, we investigated T_{exp}- and T_{per}-specific regulatory networks using a machine-learning

model (**Figure 4e**, see Methods). Single-cell transcriptomes were transformed into regulomes to

calculate regulon scores. A machine-learning model was trained using the regulon scores to

classify CD8$^+$ CAR T cells at T_{exp} and T_{per}. This strategy resulted in high (~90%) classification

accuracy across patients (figure S10a). Top T_{exp}- and T_{per}-determining regulons were identified

based on each regulon's importance (SHAP value) for the model's prediction (**Figure 4f**, table

S7). Sizes of timepoint-determining regulons span approximately 10–1000 genes (figure S10b).

Regulon expression was compared between cell clusters and timepoints (**Figure 4g**). T_{exp}- and

T_{per}-determining regulons were broadly upregulated at T_{exp} and T_{per}, respectively. However,

expression of some regulons was cluster-specific (e.g., *FLI1* regulon among EM clusters).

The top three T_{exp}-determining regulons (for *JUND, RELB, BHLHE40*) exhibited higher

expression among non-proliferating clusters and included co-regulated inflammation-associated

genes (including *NFKBIA, TNFAIP3, DUSP5*) (**Figure 4h**, see figure S11a).[37] The *RELB* and

BHLHE40 genes themselves are included in the *JUND* regulon, indicating they may be directly

upregulated by *JUND*. These three regulons were upregulated at T_{exp} and suggest response to

pro-inflammatory cytokines (such as TNF, discussed above) or CAR signaling. *BHLHE40* rewires

mitochondrial metabolism for tissue residency[38], which may contribute to CAR T-cell fitness after

entering the lymphoma. Shared genes between the *JUND* and *BHLHE40* regulons included AP-

1 transcription factors (*JUN, JUNB, FOSL2*), which have been implicated in T-cell and CAR T-cell

exhaustion[39,40], and may contribute to the exhausted-like EM phenotype that predominates the

first clonal wave at T_{exp}. Among remaining T_{exp}-determining regulons, *FLI1* is notable for

antagonizing effector T-cell differentiation.[41] Consistent with known biology, expression of the

FLI1 regulon was upregulated among EM clusters, downregulated among TE clusters, and did

not significantly change between T_{exp} and T_{per} within TE clusters.

On the other hand, T_{per}-determining regulons were more obscure (**Figure 4i**, see figure

S11b). Among top T_{per}-determining regulons, *KLF13* regulates T-cell apoptosis[42], *BPTF* activates

a stemness gene expression program[43] and maintains peripheral T-cell homeostasis[44], and *ELF4*

promotes memory CD8$^+$ T cells by regulating quiescence[45]. Altogether, the putative functions of

the *KLF13*, *BPTF*, and *ELF4* regulons may underlie the delayed expansion and persistence of

the second clonal wave at T_{per}. Moreover, these functions indicate restoration of immune

homeostasis during T_{per}. The *TBX21* regulon was notable for effector-associated genes (including

NKG7, ZEB2) and was upregulated within non-CM clusters, consistent with preponderance of the

TE phenotype at T_{per}. The *TBX21* regulon also includes *S1PR5*, which is a known T-BET target[46]

and retains T cells in peripheral blood[34]. Lastly, the *STAT1* regulon was notable for interferon-

associated antiviral genes (including *OAS2, ISG20*), in concordance with upregulation of IFN-I

signaling gene sets at T_{per}.

CD8$^+$ CAR T cells at T$_{exp}$ and T$_{per}$ originate from distinct precursors in the infusion product.

The distinct clonotypes, phenotypes, transcriptional signatures, and regulatory networks

underlying the two waves of CD8$^+$ CAR T cells led us to hypothesize that they originate from

distinct precursors in the infusion product. To test this hypothesis, we first investigated infusion

product phenotypes. After filtering for infusion product CD8$^+$ T cells, we identified four clusters

based on gene expression density (**Figure 5a-b**, figure S12b) and levels (**Figure 5c**, figure S12a):

proliferating (high *MKI67*), activated naïve-like (naïve-act, high *TCF7*), type 1 effector (EFF-Tc1,

high *TBX21*), and type 2 effector (EFF-Tc2, high *GATA3*) T cells. The *TCF7*^{hi} naïve-act cluster

upregulated markers of naiveness (*IL7R*, *LEF1*, *CCR7*, CD45RA) and T-cell activation (*CD38*,

CD95), consistent with an early activated or stem cell-like memory phenotype. The two EFF

clusters upregulated cytolytic molecules (*GZMB*, *PRF1*, *GNLY*) and downregulated markers of

naiveness (*IL7R*, *LEF1*). EFF-Tc1 and EFF-Tc2 were distinguished by expression density of

lineage-specific transcription factors (*TBX21* for Tc1, *GATA3* for Tc2) and receptors (*KLRG1* for

Tc1, *CCR4* for Tc2). The EFF-Tc1 cluster exhibited lower CAR transgene expression than the

EFF-Tc2 cluster (figure S12b-c), which may impact their functional phenotypes in vivo.[47,48] The

three non-proliferating clusters upregulated *IRF7* and anti-viral genes (*ISG20*, *IFITM1*, *IFITM2*),

indicating type I interferon signaling during ex vivo transduction and/or expansion. Cluster

proportions varied between patients (figure S12d). In general, proliferating T cells (54%) were the

most prevalent, while proportions of non-proliferating phenotypes (naïve-act, 14%; EFF-Tc1,

11%; EFF-Tc2, 21%) were mutually comparable (**Figure 5e**, top row).

Having established infusion product phenotypes, we next performed TCR lineage tracing

analysis to developmentally link infusion product CAR T cells with peripheral blood CAR T cells

using their endogenous TCR clonotypes as unique indices. This method delineates infusion

product precursors of CD8$^+$ CAR T cells at T$_{exp}$ ("Pre-T$_{exp}$") and at T$_{per}$ ("Pre-T$_{per}$") (**Figure 5d**).

Both Pre-T$_{exp}$ and Pre-T$_{per}$ comprised all four infusion product clusters (**Figure 5e**, bottom rows)

and largely overlapped on the UMAP (**Figure 5f**). Hence, we conclude that coarse cluster

phenotype alone cannot accurately predict in vivo differentiation. Compared to T cells not linked

to peripheral blood ("IP only"), Pre-T_{exp} and Pre-T_{per} cells upregulated expression of effectorness

(including *GZMB*, *TNF*, *GNLY*) and type 1 polarization (including *IFNG*, *KLRG1*) genes, as well

as lower expression of stemness (including *SELL, IL7R*) and type 2 polarization (including

GATA3, IL4R) genes (**Figure 5g**). The significance of type 1 versus type 2 polarized CD8$^+$ CAR

T cells is not well-understood in the literature and warrants future studies. Pathway analysis

indicated that Pre-T_{exp} and Pre-T_{per} upregulated gene sets for T-cell effector function ("PD-1

signaling", "T-cell-mediated cytotoxicity"), response to interferon γ ("interferon γ signaling",

"antigen processing and presentation"), and DNA replication, whereas non-linked CAR T cells

upregulated gene sets for stemness (**Figure 5h**, table S8). Our discovery of CD8$^+$ CAR T-cell

precursors with an effector-like phenotype is corroborated by findings from Thomas et al., who

also identified effector-like precursors from CAR T-cell patients treated for B-cell acute

lymphoblastic leukemia.[49] Possible fates for the non-linked, naïve-like, IP-only CAR T cells include

extravasation into the lymphoma, expansion in peripheral blood outside the T_{exp}-T_{per} window (day

8-28), or failure to persist in vivo.

Lastly, we directly compared transcriptomic signatures of Pre-T_{exp} and Pre-T_{per} via GSEA.

Although both Pre-T_{exp} and Pre-T_{per} broadly exhibited an effector phenotype (**Figure 5g-h**), Pre-

T_{exp} upregulated effectorness-associated gene sets (including TCR, PD-1, interferon γ, and IL-1

signaling), whereas Pre-T_{per} upregulated stemness-associated gene sets (**Figure 5i**). In

concordance, DGEA showed that Pre-T_{exp} upregulated effector molecules (*GZMB*, *GZMK*,

FCGR3A, *NKG7*, *KLRG1*), AP-1 transcription factors (*JUN*, *JUND*, *FOS*), and MHC class II

expression (*HLA-DRB1*, *HLA-DRB5*, *HLA-DRA*), whereas Pre-T_{per} upregulated naïve-like

markers (*SELL*, *IL7R*, *S1PR1*) (figure S12e, table S9). Upregulation of *JUN* in Pre-T_{exp} confers

exhaustion resistance[39], potentially underlying their eventual efficacy at T_{exp}. Interestingly, Pre-

T_{per} exhibited greater CAR transgene expression (figure S12e), which may reflect transduction

differences among apheresis precursors or a link between CAR expression and in vivo

differentiation.[48] Next, we compared Pre-T_{exp} and Pre-T_{per} using transcriptomic signatures from a

human CD8[+] differentiation atlas from Giles et al.[50] Pre-T_{exp} more closely resembled an effector

memory state, whereas Pre-T_{per} more closely resembled either a stem-cell memory or central

memory state (**Figure 5j**). Altogether, these patterns indicate that, while both Pre-T_{exp} and Pre-

T_{per} exhibited an effector phenotype in the infusion product, Pre-T_{exp} were more differentiated with

greater effectorness, whereas Pre-T_{per} were less differentiated with greater stemness.

In conclusion, TCR lineage tracing analysis supports our hypothesis that CD8[+] CAR T

cells at T_{exp} and T_{per} originate from different infusion product precursors. Integrating these findings

into the two-stage differentiation model paints a more complete picture of CD8[+] CAR T cells over

the month following CAR T-cell administration (**Figure 5k**). Effector CD8[+] CAR T cells exist along

a gradient of effectorness and stemness in the infusion product. Following infusion into the patient,

effector CAR T cells with greater effectorness rapidly expand until peak expansion (T_{exp}, days 8-

12), adopting a functional and cytotoxic EM phenotype with exhaustion-like characteristics upon

antigen stimulation. Lymphoma infiltration and in vivo killing[51], CRS[5], and ICANS[5] coincide with

the T_{exp} timeframe, suggesting that these CAR T cells mediate tumor clearance and side effects.

Subsequently, this first wave of expanded CAR T cells diminishes through apoptosis or

extravasation. Simultaneously, the remaining effector CAR T cells with greater stemness from the

infusion product expand during the post-peak persistence timeframe (T_{per}, day 21-28). These

newer and longer-lived CAR T cells adopt effector characteristics and persist in vivo, where they

may ensure a durable response through long-term immunosurveillance.

CD8[+] Exhausted-like EM CAR T cells at T_{exp} exhibit characteristics of early exhaustion.

Discovery of the predominant exhausted-like EM phenotype in peripheral blood at T_{exp} may be

surprising since exhaustion was conventionally defined in lymphoid organs or tumor-infiltrating

lymphocytes.[52] To verify this annotation, we further characterized the molecular signatures of the

exhausted-like EM cluster. The exhausted-like EM cluster was densely situated at the lower half

of the UMAP (**Figure 6a-b**). We used RNA-seq/CITE-seq to examine gene and protein expression

density, respectively (**Figure 6c**, figure S13a for additional markers). The exhausted-like EM

cluster highly expressed exhaustion-associated transcription factors (*NR4A2*, *TOX*, *IRF4*) and

inhibitory receptors (*ENTPD1*, *PDCD1*, *TIGIT*, *LAG3*, *CTLA4*). Intermediate expression of

memory (*TCF7*, *LEF1*, *CD27*, *IL7R*) and effector (*TBX21*, *GZMB*, *PRF1*, *GNLY*, *IFNG*, *NKG7*)

genes was also observed. Additionally, the cluster exhibited low expression of *CX3CR1* (effector

lineage marker) and *B3GAT1* (senescence marker). CITE-seq measurements of protein

expression for cell surface receptors (including CD39, PD-1, TIGIT, LAG3, CD27, CD57, CD127,

TIM-3, CXCR3) were largely consistent with corresponding gene expression. Sequencing data

indicated that high expression of CD39 (protein for *ENTPD1*) and low expression of CD57 (protein

for *B3GAT1*) most clearly differentiated the exhausted-like EM cluster from other clusters. To

validate these characteristics, flow cytometry was employed to analyze a set of longitudinal

patient PBMCs. Consistent with sequencing data, flow cytometry data demonstrated that the

CD39$^+$CD57$^-$ phenotype was highest among CD8$^+$ CAR T cells at T_{exp} (75%) compared to at T_{per1}

(33%) and T_{per2} (18%) (**Figure 6d**, top row). Furthermore, flow cytometry showed decreased

CX3CR1 expression at T_{exp}, and increased expression at T_{per1} and T_{per2}, in agreement with

sequencing data (**Figure 6d**, bottom row). This mixed expression pattern of exhaustion, memory,

and effector markers resembles the circulating PD1$^+$CD39$^+$ T_{ex} cells described by the Wherry

group's human T-cell differentiation atlas.[50]

Next, we conducted GSEA using gene sets from Wherry et al. that identified T-cell

differentiation states in the Armstrong/clone 13 LCMV model (**Figure 6e**).[53] These gene sets

constitute comprehensive references for defining T-cell exhaustion because they (1) incorporate

standing knowledge from the entire field, (2) integrate single-cell RNA-seq and ATAC-seq data

from serial timepoints, and (3) originate from the Wherry group where comprehensive

transcriptional signatures of T-cell exhaustion and most exhausted subsets were first defined. The

two EM clusters (EM and exhausted-like EM) in our dataset correlated with different states from

the Wherry group's model. Our EM cluster resembled the "transitional I" T cells from the

Armstrong model, indicating that these cells are not exhausted. Conversely, our exhausted-like

EM cluster resembled the "precursor exhausted" T cells from the clone 13 model (**Figure 6f**, left).

Simultaneously, the exhausted-like EM cluster did not resemble the "terminally exhausted" T cells

from the same model (**Figure 6f**, right), which instead correlated with our exhausted-like TE

cluster. Given clear differences between the EM and exhausted-like EM clusters, we directly

compared their transcriptomic signatures through DGEA and GSEA. The exhausted-like EM

cluster upregulated genes associated with T-cell activation (*CD69*, *CD38*, *DUSP4*, *NFKBIA*) and

exhaustion (*TOX*, *NR4A2*, *PRDM1*, *CTLA4*) (**Figure 6g**, table S10). In contrast, the EM cluster

upregulated genes associated with effector function (*TBX21*, *CX3CR1*, *GZMB*, *PRF1*, *FCGR3A*),

memory (*IL7R*, *S1PR1*), and type I interferon signaling (*STAT1*, *IFITM1*, *IFITM3*). GSEA revealed

that the exhausted-like EM cluster exhibited more exhaustion signatures and fewer effector

signatures compared to the EM cluster (**Figure 6h**, left column, see also figure S13b). GSEA also

confirmed that the exhausted-like EM cluster resembled the "exhausted progenitor" and

"exhausted precursor" subsets from Wherry et al. (whereas the EM cluster resembled the

"transitional I" subset) (**Figure 6h**, middle and right columns). Our findings suggest that the

exhausted-like EM cluster exhibits robust gene and protein signatures of early exhaustion

(justifying the "exhausted-like" annotation), yet is also phenotypically distinct from effector

memory T cells.

DISCUSSION

In this study, we use a combination of single-cell RNA-seq/CITE-seq/TCR-seq and longitudinal analyses to investigate CD28-costimulated CAR T-cell differentiation in seven patients with r/r DLBCL. Our findings are summarized in our two-stage model for CAR T-cell differentiation. Specifically, CD8+ CAR T cells undergo two distinct clonal expansion waves (at peak expansion and post-peak persistence timeframes), as revealed by clonotypic, phenotypic, and linked clonotypic-phenotypic analyses. The two waves are dominated by exhausted-like effector memory and terminal effector phenotypes, respectively. The exhausted-like effector memory annotation is supported by a CD39+CD57- flow phenotype, low CX3CR1 expression, and an early exhaustion signature. We also identify transcription factors and regulatory networks associated with the first wave (including *JUND*, *RELB*, *BHLHE40*, *FLI1* regulons) and second wave (including *KLF13*, *BPTF*, *ELF4*, *TBX1*, *STAT1* regulons). Lastly, lineage tracing analysis determined that CD8+ CAR T cells from both waves derive from effector precursors in the infusion product. However, effector precursors of the first wave exhibit more effector-like signatures, whereas effector precursors of the second wave exhibit more stem-like signatures, suggesting that pre-infusion heterogeneity mediates two-stage differentiation. Our two-stage model implies that manipulating the phenotypic composition of the infusion product may allow more precise control over in vivo CAR T-cell differentiation for modulating therapeutic efficacy.

Although two-stage differentiation is a novel phenomenon that has not yet been reported in the literature, the individual elements of our model are consistent with current understanding in the field. Previous studies on 4-1BB-costimulated CAR T-cell differentiation described bursts of CAR T-cell clonal expansion that lead to significant changes in the CAR T-cell clonal repertoire over time.[19,54] Our study not only generalizes this observation to CD28-costimulated CARs, but also directly pairs changes in the clonal repertoire with changes in T-cell phenotypes. Furthermore, previous reports have described CAR T-cell clones with memory-like phenotypes

that exhibit delayed expansion after infusion[23], which corroborates with the second clonal

expansion wave at T_{per}. Finally, the T_{exp}- and T_{per}-specific phenotypes described in our model are

consistent with prior phenotyping studies.[25,27,49,55,56] We also report agreement between our

transcriptomes at T_{exp} and external data from Maus et al.[27] By integrating our findings from multiple

data modalities with established studies in the CAR T-cell field, we present a more complete and

coherent two-stage model for in vivo CD28-costimulated CAR T-cell differentiation.

The two-stage differentiation model we present in this study has an important implication:

CD28-costimulated CD8$^+$ CAR T cells from the peak expansion and post-peak persistence

timeframes are biologically uncoupled. This uncoupling significantly informs how we understand

CAR T-cell expansion and persistence. Expansion facilitates rapid tumor clearance but can also

cause CRS and ICANS[5,17], while persistence facilitates long-term immunosurveillance but can

also cause B-cell aplasia and hypogammaglobulinemia.[9,13,57] Our findings suggest that expansion

and persistence, both of which serve complementary clinical purposes, are mediated by distinct

CD8$^+$ CAR T-cell populations. Effective and personalized CAR T-cell therapies balance expansion

and persistence, taking into consideration each patient's tumor burden, tolerance for side effects,

and risk of relapse. Our findings suggest that expansion and persistence may be independently

tuned to meet a patient's needs. On the other hand, our findings also suggest that engineering

CD8$^+$ CAR T cells that simultaneously expand and persist competently may be challenging, given

that these characteristics originate from uncoupled CAR T-cell populations in vivo.

In addition to clonal kinetics and phenotypic heterogeneity, we characterized T_{exp}- and

T_{per}-specific upregulation of TNF and IFN-I response gene sets and regulons, respectively.

Notably, TNF-secreting CD8$^+$ CAR T cells have been associated with complete responders[20],

suggesting that TNF may be signaling in an autocrine or paracrine manner during T_{exp}.

Upregulation of IFN-I response genes during T_{per} was intriguing. IFN-I response signatures in the

apheresis of patients with B-cell acute lymphoblastic leukemia predict poor CAR T-cell

persistence[58], but IFN-I signaling may also enhance CAR T-cell efficacy in vivo.[59] Complicating

this story further, we also identified IFN-I response signatures in the infusion product, which

implies that interferons are secreted by an unknown source during ex vivo transduction and/or

expansion. Hence, the possible roles of IFN-I in CAR T-cell differentiation are likely complex,

context-dependent, or time-dependent, and warrant further mechanistic investigation.

We concluded our study by using single-cell TCR-seq to link CD8$^+$ CAR T cells at T_{exp} and

T_{per} with effector precursors in the infusion product that exhibit more effector-like or more stem-

like signatures, respectively. Discovery of effector precursors is externally corroborated by

findings from Thomas et al[49]. Importantly, this linkage suggests that the magnitude or duration of

peak expansion and post-peak persistence can be modulated by manipulating the relative

quantities of precursors in the infusion product. However, we cannot rule out the possibility that

the in vivo peripheral blood environment in a patient with DLBCL also plays an instrumental role

in determining whether a CAR T cell clonally expands at T_{exp} or T_{per}. Future studies can investigate

this possibility by analyzing interactions between CAR T cells and other blood cells (such as

myeloid cells, B cells, or NK cells) at T_{exp} and T_{per}.

The present study focuses on CD28-costimulated CAR T cells from complete responders,

but two-stage differentiation may be more widely applicable. Others forms of adoptive cell therapy,

such as 4-1BB-costimulated CAR T cells for B-cell leukemias[60] or systemic lupus

erythematosus[61], may also exhibit a two-stage differentiation pattern in vivo. Indeed, our

transcriptomic signatures at T_{exp} were highly consistent with external data from day 7

tisagenlecleucel CD8$^+$ CAR T cells from Maus et al.[27], hinting that two-stage differentiation may

be more general. Different forms of adoptive cell therapy and disease contexts may exhibit

different temporal dynamics or favor one wave over the other. Two-stage differentiation may also

be aberrant in non-responders. Future studies are needed to explore potential differences in

temporal dynamics, cell phenotypes, or major regulatory pathways at longitudinal timepoints in

alternative therapeutic contexts.

Although this current study represents an advance for the CAR T-cell therapy field, we

acknowledge four limitations. Firstly, we analyzed CAR T cells in two locations: peripheral blood

and infusion product. However, CAR T cells are also found elsewhere, including lymph nodes and

lymphoma foci. Importantly, CAR T-cell differentiation may be influenced by interactions with

antigen-presenting cells in lymph nodes or with tumor cells in the lymphoma. Our study was

ultimately limited by sample availability. Future studies can focus on less accessible locations and

determine CAR T cell fates beyond peripheral blood. Secondly, we studied CAR T-cell intrinsic

factors (RNAs, proteins, clonotypes) that can influence differentiation. However, differentiation

may also depend on interactions between CAR T cells and a complex in vivo environment that

includes other leukocytes (myeloid cells, B cells, NK cells), lymphoma cells, and cytokines.

Thirdly, we focused on CAR T cells longitudinally between peak expansion and post-peak

persistence timeframes. Hence, we cannot rule out additional waves of clonal expansion in the

near (<1 week after infusion) or long (>1 month after infusion) terms. Lastly, we lack knowledge

on whether two-stage differentiation applies to other contexts, such as non-responders, 4-1BB-

costimulated CAR T cells, or TCR-transduced T cells. Future studies can explore alternative

clinical contexts and test the generality of the two-stage differentiation model.

METHODS

Patient biospecimen collection.

Deidentified biospecimens were obtained from a cellular therapy biobank in accordance with the institutional review board at the University of Chicago Medicine. Ethical guidelines were followed.

Longitudinal peripheral blood mononuclear cells (PBMCs) were collected from peripheral blood biospecimens by Ficoll-Paque PLUS (Cytiva, 95021-205) and stored in freezing media (RPMI, 10% FBS, 10% DMSO). Residual infusion product cells were collected from the patient's spent infusion product bag and stored in CELLBANKER 1 (Amsbio, 11888). Cells were cryopreserved in liquid-phase nitrogen.

Generation of CD19-tetramers.

Tetramers[28] were constructed from AviTag-biotinylated human His-tagged CD19 (Acro Biosystems, CD9-H82E9) and Alexa Fluor 647-labeled streptavidin (BioLegend, 405237). Biotinylated CD19 was added to the tetrameric streptavidin at a 4:1 molar ratio for 30 minutes at 4°C in the dark. This mixture was diluted with PBS to convenient concentrations for staining. For each batch of biotinylated CD19, quality controls were performed by generating CD19-tetramers and quantifying their ability to stain anti-CD19 CAR-transduced Jurkat T cells.

Single-cell RNA-seq/CITE-seq/TCR-seq.

Cryopreserved biospecimens were thawed (RPMI, 10% FBS) and washed with cold FACS buffer (PBS, 2% BSA, 0.05% sodium azide). Fc receptors were blocked with Human TruStain FcX (BioLegend, 422301) at 1:50 dilution for 5 minutes at 4°C. Then, cells were incubated for 30 minutes at 4°C in the dark with a staining solution containing CITE-seq antibodies (described below), BV421-labeled anti-CD3ε (clone SK7, BioLegend, 344833), and AF647-labeled CD19-tetramers (3 nM final concentration) for CAR T-cell phenotyping and detection. Subsequently,

stained cells were conjugated with LIVE/DEAD Fixable Near-IR viability dye (Invitrogen, L34975)

at 1:1000 dilution in PBS for 5 minutes at room temperature. Finally, cells were washed three

times in cold cell media (RPMI, 10% FBS) before fluorescence-activated cell sorting (BD

Biosciences, FACSAria Fusion). CAR$^+$ sorting gates were drawn based on fluorescence of

PBMCs from a similarly stained healthy donor.

Sorted endogenous T cells (CD3$^+$CAR$^-$) and CAR T cells (CD3$^+$CAR$^+$) were separately

partitioned into droplets for single-cell RNA-seq/CITE-seq/TCR-seq via Chromium Next GEM

Single-Cell 5'Kit v2 (10x Genomics, 1000263). RNA-seq libraries were prepared according to

manufacturer protocols. CITE-seq libraries were prepared via the 5' Feature Barcode Kit (10x

Genomics, 1000256). TCR-seq libraries were prepared via the Chromium Single-Cell Human

TCR Amplification Kit (10x Genomics, 1000252). All libraries (RNA-seq, CITE-seq, TCR-seq)

were quantified via the Qubit dsDNA HS Assay Kit (Invitrogen, Q32851), quality-checked for

fragment sizes via high-sensitivity D5000 screentapes (Agilent, 5067-5592), pooled, and

sequenced (Illumina, Novaseq-6000 and Nextseq-550).

CITE-seq antibody preparation.

Thirty-one human "Cellular Indexing of Transcriptomes and Epitopes by Sequencing" (CITE-

seq[62]) antibodies were obtained from BioLegend (TotalSeq-C reagents): anti-CD3ε (clone

UCHT1, 300479), anti-CD5 (clone UCHT2, 300637), anti-TCRα/β (clone IP26, 306743), anti-

TCRγ/δ (clone B1, 331231), anti-CD4 (clone SK3, 344651), anti-CD8α (clone SK1, 344753), anti-

CD45RA (clone HI100, 304163), anti-CD45RO (clone UCHL1, 304259), anti-CCR7 (clone

G043H7, 353251), anti-CD95 (clone DX2, 305651), anti-CD57 (clone QA17A04, 393321), anti-

CD25 (clone BC96, 302649), anti-CD127 (clone A019D5, 351356), anti-CD103 (clone Ber-ACT8,

350233), anti-CXCR3 (clone G025H7, 353747), anti-CCR4 (clone L291H4, 359425), anti-CCR6

(clone G034E3, 353440), anti-PD-1 (clone EH12.2H7, 329963), anti-TIM-3 (clone F38-2E2,

345049), anti-LAG-3 (clone 11C3C65, 369335), anti-CD39 (clone A1, 328237), anti-TIGIT (clone

A15153G, 372729), anti-CD27 (clone O323, 302853), anti-CD40L (clone 24-31, 310849), anti-

GITR (clone 108-17, 371227), anti-OX40 (clone Ber-ACT35, 350035), anti-4-1BB (clone 4B4-1,

309839), anti-CD28 (clone CD28.2, 302963), mouse IgG1-κ isotype control (clone MOPC-21,

400187), mouse IgG2a-κ isotype control (clone MOPC-173, 400293), and mouse IgG2b-κ isotype

control (clone MPC-11, 400381). Residual infusion product cells were stained with all 31 CITE-

seq antibodies. Patient peripheral blood mononuclear cells were stained with all CITE-seq

antibodies minus anti-CD3ε (30 in total) because T cells sorted from these samples were stained

with BV421-labeled anti-CD3ε. To prepare the CITE-seq staining solution, the antibody pool was

constructed and centrifuged at 14000 × g for 10 minutes at room temperature in FACS buffer

(PBS, 2% BSA, 0.05% sodium azide) to remove aggregates. The antibody supernatant was

extracted and diluted with fresh FACS buffer to appropriate staining concentrations.

Analysis of CAR T-cell phenotype by flow cytometry.

Cryopreserved biospecimens were thawed (RPMI, 10% FBS) and washed with cold FACS buffer

(PBS, 2% BSA, 0.05% sodium azide). Fc receptors were blocked with Human TruStain FcX

(BioLegend, 422301) at 1:50 dilution for 5 minutes at 4°C. Then, cells were incubated for 30

minutes at 4°C in the dark with a staining solution containing AF647-labeled CD19-tetramers (3

nM final concentration), BV421-labeled anti-CD3ε (clone SK7, BioLegend, 344833), AF488-

labeled anti-CD8α (clone SK1, BioLegend, 344716), PE-labeled anti-CX3CR1 (clone 2A9-1,

BioLegend, 341603), PE/Cy7-labeled anti-CD39 (clone A1, BioLegend, 328211), and BV605-

labeled anti-CD57 (clone QA17A04, BioLegend, 393303). Subsequently, stained cells were

conjugated with LIVE/DEAD Fixable Near-IR viability dye (Invitrogen, L34975) at 1:1000 dilution

in PBS for 5 minutes at room temperature. Finally, cells were washed three times in cold FACS

buffer (RPMI, 10% FBS) before flow cytometer analysis (BD Biosciences, FACSAria Fusion).

Single-cell RNA-seq data processing.

RNA-seq and paired CITE-seq reads were aligned to the GRCh38 reference genome, which was

modified with the CAR transgene sequence used in axicabtagene ciloleucel, and quantified using

the cellranger count (10x Genomics, version 7.0.0).[63] Only filtered gene/feature-barcode matrices

that contained barcodes with unique molecular identifier (UMI) counts that passed the quality

control were used for downstream analyses.

UMAP analysis and clustering on single-cell RNA-seq data.

UMAP analysis and clustering were performed using the Seurat package (Version 4.3.0).[64] Raw

count matrices were first converted to Seurat objects before being further merged into one Seurat

object, with protein expression added as the antibody-derived tag (ADT) assay. Cells with <300

genes detected or >10% mitochondrial RNA content were excluded from further analysis.

The raw count was log-normalized using the *NormalizeData* function with default options.

The top 5,000 variable features were then identified using the *FindVariableFeatures* function with

the default "vst" method. The data were centered and scaled using the *ScaleData* function, with

additional regression against 1) the percent of mitochondrial RNA content and 2) difference in cell

cycle S-phase score and G2M-phase score. Scaled data were then used as input for principal

component analysis (PCA) based on variable genes using the *RunPCA* function. Data

harmonization to remove patient-specific effects was performed on the principal components

using the Harmony package (Version 1.0) through the *RunHarmony* function.[65] Next, UMAP was

constructed based on the first 30 harmony components. The same harmony components were

used to construct the shared nearest neighbor (SNN) graph using the *FindNeighbors* function,

which was then partitioned to identify clusters using the *FindClusters* function with default Louvain

algorithm. These clusters were manually aggregated and classified as T-cell subsets based on

known markers. CD8 and CD4 T-cell classifications were based on both gene and protein

expression of CD4, CD8α, and CD8β.

For subset analysis, the corresponding subsets were extracted from the master Seurat

object using the *subset* function. The above detailed preprocessing steps were repeated to

generate the corresponding UMAP and subset annotations.

Gene expression visualization.

Gene expression was visualized in three ways: violin plots, heatmaps, and density plots. Violin

plots were made based on normalized expression. Heatmaps were made based on average

scaled expression, both using Seurat internal functions. Density plots were constructed using the

Nebulosa package (version 1.6.0) to enhance visibility and mitigate sparsity of gene expression

on UMAP. This is necessary for datasets with large number of cells (e.g. our CD8[+] CAR-T dataset

has ~38k cells).

Differential gene expression (DEG) analysis on single-cell RNA-seq data.

DEG analyses were by default performed using the *FindMarkers* function in Seurat package, with

default parameters and the appropriate 'ident.1" and "ident.2" set as contrast. Unless otherwise

stated, the results were filtered with p_val_adj < 0.05 and abs(avg_log2FC) > 0.25. Moreover, we

also performed additional DEG analyses through a pseudo-bulk approach to better control for

patient-specific effects, using the LibraDEG package (version 1.0.0)[66] with default parameters

(edgeR with LRT method). This was mainly performed when extracting timepoint-specific genes.

Areas where this pseudo-bulk method is used are clearly stated in the figure legend.

Timepoint-specific gene signature validation.

To validate the timepoint-specific molecular signatures shared across patients and against

external data, we adopted two independent and complementary approaches: 1) correlation of

timepoint-specific gene expression, 2) label transfer from our data to external data.

For the correlation approach, we first conducted pseudo-bulk DEG analyses of T_{exp} vs T_{per}

across all patients and kept all the statistically significant genes (padj <0.05) as timepoint-specific

genes. Note that we did not filter by log fold-change to include even weak timepoint-specific

signatures. Essentially, the goal here is to remove all non-timepoint-specific genes that may

confound correlation analysis. The normalized gene expressions of these timepoint-specific

genes were extracted from both our data and the external data. Following that, average

expression was calculated for each sample (patient by timepoint). Correlations were then

calculated pairwise for all samples.

For the label transfer approach, we adapted the Seurat standard data integration and label

transfer workflow with default parameters. This is a complementary approach where, rather than

at patient-level expression, individual cells in the external data and our data could be compared

and similar timepoint-labels could be assigned.

GSEA and pathway enrichment analysis.

Gene Set Enrichment Analysis (GSEA) and pathway enrichment analysis were carried out using

the clusterProfiler (version 4.4.4)[67] package based on the msigdb database built in msigdbr

package (version 7.5.1).[68]

Exhaustion signatures from reference datasets.

Besides standard msigdb exhaustion-related pathways, we obtained T-cell subset-specific

signatures from Wherry et al. (see Supplementary Table 1 in their study).[53] The reference study

provides the most updated and comprehensive CD8[+] (exhausted) T cell subsets, in the standard

LCMV/Cl13 mouse models where T_{ex} was first defined. To refine the signature, we converted the

mouse genes to their human orthologous genes and filtered the list to retain only the top 500 (by

descending avg_log2_FC) statistically significant (p_val_adj < 0.05) genes for each subset.

Regulon analysis.

To construct regulons in a specific cell subset, we first extracted the raw count matrix of that

subset. Then, we supplied it as input for the pySCENIC package (version 0.10.4) and ran through

the workflow as detailed in its documentation and publication.[69,70] More specifically, the base script

was adapted from the "PBMC10K" example script on the pyScenic website.[71] In addition, the

arboreto component was used to speed up analysis. Each resultant regulon is a gene list with a

central transcription factor and all its putative target genes determined through the SCENIC

algorithm, and each set of regulons is sample/subset-specific and represents a putative regulatory

unit in that specific sample/subset.

Regulon or general gene set signature scoring.

Individual cells were scored using the AUCell package (version 1.10.0)[69] for a particular gene set

from the msidb database or from a single-cell-derived regulon as follows. The normalized gene

expression was first used as input into the *AUCell_buildRankings* function to score each cell for

gene set enrichment and to build a ranking matrix. The signature score was then calculated as

an AUC score using the *AUCell_calcAUC* function with all default parameters. In later revisions,

Ucell package (version 2.0.1)[72] was used in some re-analyses, as it is computationally more

efficient and generates similar results as the AUCell package.

Machine learning classification on regulon scores.

To classify CD8$^+$ CAR T cells at T_{exp} and T_{per}, we first built a regulon set for each subset

independently, assuming that regulatory networks for each timepoint may be different. Then, we

merged all regulon sets into a master regulon set. We calculated the regulon scores using the

master regulon set for all cells from the two subsets, resulting in a regulon score matrix. Each cell

then served as an observation and each regulon as a feature for machine learning.

The machine learning classification models were built in the caret framework using the

caret package (version 6.0-90).[73] Briefly, we divided the input data into training (75%), validation

(15%), and test sets (10%). We trained and optimized xgboost models[74] through grid

hyperparameter search and with a 5-fold cross-validation until reaching a best accuracy of >85%

(refer to figures for specific accuracies).

We interrogated the final models for feature importance evaluation using the Shapley

Additive Explanation (SHAP) method[75] implemented in the SHAPforxgboost package (version

0.1.1).[76] Average SHAPley values were used to rank the regulons. Top timepoint-determining

regulons were visualized and further analyzed.

Single-cell TCR-seq data processing.

The TCR-seq reads from each sample were aligned to the 10x curated GRCh38 vdj reference

genome and quantified using the cellranger vdj (10x Genomics, version 6.0.0). The results were

then aggregated using the cellranger aggr function, with the source patient (donor) and timepoint

(origin) of each sample supplied in the metadata to guide the clonotype assignment. The resultant

clonotype and filtered contig annotation data were used for downstream analyses.

Single-cell TCR-seq tracing and phenotype linking.

Each unique clonotype is defined by the amino acid and nucleotide sequence of CDR3 regions

for paired productive TCRα and TCRβ chains. TCR clonality overlap and tracing analyses were

carried out using the immunarch package (version 0.9.0).[77] Clones with the same TCR CDR3

sequences from the same patient at different timepoints were re-grouped and traced from infusion

product to the post-infusion samples.

Phenotype-to-clonotype linking: To identify T_{exp} and T_{per} precursors, clonotypes among

post-infusion T_{exp} and T_{per} cell populations were identified respectively and traced back to the

infusion product cells. When linking the TCR clonotype to the corresponding cell phenotype, we

keep only data with matching cell barcodes from both the TCR-seq and RNA-seq data.

Abundance tracing: Clonal abundance for each CAR T clone is calculated as the

percentage of all CAR T cells occupied by a clone at a given timepoint. Each clone was assigned

a phenotype at each timepoint based on the most frequently occurring phenotype. T_{exp} clones that

were predominantly EM-exh had their abundances tracked at T_{per1}/T_{per2}. T_{per1}/T_{per2} clones that

were predominantly TE had their abundances tracked at T_{exp}.

CAR transgene mapping.

The axicabtagene ciloleucel CAR design is documented.[78] Its sequence was confirmed by Sanger

sequencing of genomic DNA extracted from axicabtagene ciloleucel infusion products. The CAR

sequence was then added to the GRCh38 FASTA and GTF files accordingly. A custom reference

for cellranger was built from these annotation files using cellranger mkref (10x Genomics, version

7.0.0). The resultant custom reference was used for CAR transgene mapping through cellranger

count.

Data availability.

All data are available from the authors upon reasonable request.

Code availability.

For reproducibility, code used for our analysis and figure-making is available at github

(https://github.com/DanielGuoshuaiCao/TwoStageCART).

Inclusion and Ethics Statement.

Roles and responsibilities were agreed amongst collaborators in advance. Our findings are not

expected to result in stigmatization, incrimination, or discrimination.

ACKNOWLEDGEMENTS

ACKNOWLEDGEMENTS

We thank Phi Beta Psi, the Ullman Fund in Cancer Immunology, the Hoogland Lymphoma

Research Pilot Projects, Chicago Immunoengineering Innovation Center, American Cancer

Society Scholar Award (SG-22-136-01-1BCD) and NIH New Innovator award (1DP2AI144245)

for financial support (to J.H.). Y.H. was supported by the University of Chicago MSTP Training

Grant (T32GM007281). T.P. was supported by the University of Chicago CSTR training grant

(T32HL007381). N.A. was supported by the University of Chicago MTCR training grant

(T32CA009594). This project was supported by the National Center for Advancing Translational

Sciences of the National Institutes of Health through Grant Number 5UL1TR002389-02 that funds

the Institute for Translational Medicine. We thank the UChicago Human Immunologic Monitoring

Facility for patient biospecimen cryopreservation, as well as the Pritzker School of Molecular

Engineering Single-Cell Immunophenotyping Core and UChicago Genomics Facility for

instrumentation to generate single-cell sequencing data. This work was completed in part with

resources provided by the University of Chicago's Research Computing Center. The authors

thank Dr. Karen M. Watters for scientific editing of the manuscript.

AUTHOR CONTRIBUTIONS

Y.H., G.C., J.P.K., and J.H. conceived and designed all experiments. J.H. supervised the project.

T.A., P.A.R., and M.R.B. coordinated patient biospecimens for these studies. G.C. and T.P.

performed all single-cell and machine-learning analyses. Y.H. organized execution of all

experiments. Y.H. and E.T. sorted CAR T cells and generated single-cell data. G.C. and Y.H.

jointly interpreted data. N.A. and Y.H. designed cartoons. E.T. performed flow cytometry

validation. Y.H. and G.C. prepared the manuscript. J.H. and J.P.K. edited the manuscript. All

authors reviewed the manuscript.

COMPETING INTERESTS

P.A.R. reports Research Support/Funding: BMS, Kite Pharma, Inc./Gilead, MorphoSys, Calibr,

Tessa Therapeutics, Fate Therapeutics, Xencor, and Novartis Pharmaceuticals Corporation.

Speaker's Bureau: Kite Pharma, Inc./Gilead; Consultancy on advisory boards: AbbVie, Novartis

Pharmaceuticals Corporation, BMS, Janssen, BeiGene, Karyopharm Therapeutics Inc., Takeda

Pharmaceutical Company, Kite Pharma, Inc./Gilead, Sana Biotechnology, Nektar Therapeutics,

Nurix Therapeutics, Intellia Therapeutics, and Bayer. Honoraria: Novartis Pharmaceuticals

Corporation. M.R.B. reports Membership on an Advisory Board or Consultancy for Kite/Gilead,

Novartis, CRISPR Therapeutics, Autolus Therapeutics, BMS, Incyte, Sana

Biotechnology, Iovance Biotherapeutics. He has served on a Speakers Bureau for BMS,

Kite/Gilead, Agios, and Incyte. J.P.K. receives research support from Merck, Verastem, and

iTeos; has served on a speaker's bureau for Kite/Gilead; and has served on advisory boards for

Verastem, Seattle Genetics, MorphoSys, and Karyopharm.

REFERENCES

1.   Diffuse Large B-Cell Lymphoma - Cancer Stat Facts.
      https://seer.cancer.gov/statfacts/html/dlbcl.html.

2.   Liu, Y. & Barta, S. K. Diffuse large B-cell lymphoma: 2019 update on diagnosis, risk
      stratification, and treatment. *Am. J. Hematol.* **94**, 604–616 (2019).

3.   Longo, D. L., Sehn, L. H. & Salles, G. Diffuse Large B-Cell Lymphoma. *https://doi-
      org.proxy.uchicago.edu/10.1056/NEJMra2027612* **384**, 842–858 (2021).

4.   June, C. H., O'Connor, R. S., Kawalekar, O. U., Ghassemi, S. & Milone, M. C. CAR T cell
      immunotherapy for human cancer. *Science* vol. 359 1361–1365 (2018).

5.   Neelapu, S. S. *et al.* Axicabtagene Ciloleucel CAR T-Cell Therapy in Refractory Large B-
      Cell Lymphoma. *N. Engl. J. Med.* **377**, 2531–2544 (2017).

6.   Schuster, S. J. *et al.* Tisagenlecleucel in adult relapsed or refractory diffuse large B-cell
      lymphoma. *N. Engl. J. Med.* **380**, 45–56 (2019).

7.   Abramson, J. S. *et al.* Pivotal Safety and Efficacy Results from Transcend NHL 001, a
      Multicenter Phase 1 Study of Lisocabtagene Maraleucel (liso-cel) in Relapsed/Refractory
      (R/R) Large B Cell Lymphomas. *Blood* **134**, 241–241 (2019).

8.   Abramson, J. S. *et al.* Lisocabtagene maraleucel for patients with relapsed or refractory
      large B-cell lymphomas (TRANSCEND NHL 001): a multicentre seamless design study.
      Lancet **396**, 839–852 (2020).

9.   Locke, F. L. *et al.* Long-term safety and activity of axicabtagene ciloleucel in refractory
      large B-cell lymphoma (ZUMA-1): a single-arm, multicentre, phase 1–2 trial. *Lancet
      Oncol.* **20**, 31–42 (2019).

10.   Schubert, M. L. *et al.* Side-effect management of chimeric antigen receptor (CAR) T-cell
      therapy. *Ann. Oncol.* **32**, 34–48 (2021).

11.   Milone, M. C. *et al.* Engineering-enhanced CAR T cells for improved cancer therapy. *Nat.
      Cancer 2021 28* **2**, 780–793 (2021).

12.   Good, Z. *et al.* Post-infusion CAR TReg cells identify patients resistant to CD19-CAR
      therapy. *Nat. Med.* **28**, 1860–1871 (2022).

13.   Gupta, A. & Gill, S. CAR-T cell persistence in the treatment of leukemia and lymphoma.
      https://doi-org.proxy.uchicago.edu/10.1080/10428194.2021.1913146 **62**, 2587–2599
      (2021).

14.   Kawalekar, O. U. *et al.* Distinct Signaling of Coreceptors Regulates Specific Metabolism
      Pathways and Impacts Memory Development in CAR T Cells. *Immunity* **44**, 380–390
      (2016).

15.   Salter, A. I. *et al.* Phosphoproteomic analysis of chimeric antigen receptor signaling
      reveals kinetic and quantitative differences that affect cell function. *Sci. Signal.* **11**, 1–18
      (2018).

16.   Fraietta, J. A. *et al.* Determinants of response and resistance to CD19 chimeric antigen
      receptor (CAR) T cell therapy of chronic lymphocytic leukemia. *Nat. Med.* **24**, 563–571
      (2018).

17.   Rossi, J. *et al.* Preinfusion polyfunctional anti-CD19 chimeric antigen receptor T cells are
      associated with clinical outcomes in NHL. *Blood* **132**, 804–814 (2018).

18.   Deng, Q. *et al.* Characteristics of anti-CD19 CAR T cell infusion products associated with
      efficacy and toxicity in patients with large B cell lymphomas. *Nat. Med.* **26**, 1878–1887
      (2020).

19.   Melenhorst, J. J. *et al.* Decade-long leukaemia remissions with persistence of CD4+ CAR
      T cells. *Nature* **602**, 503–509 (2022).

20.   Finney, O. C. *et al.* CD19 CAR T cell product and disease attributes predict leukemia

remission durability. *J. Clin. Invest.* **129**, 2123–2132 (2019).

21.   Neelapu, S. S. CAR-T efficacy: is conditioning the key? *Blood* **133**, 1799–1800 (2019).

22.   Turtle, C. J. *et al.* CD19 CAR–T cells of defined CD4+:CD8+ composition in adult B cell
ALL patients. *J. Clin. Invest.* **126**, 2123–2138 (2016).

23.   Fraietta, J. A. *et al.* Disruption of TET2 promotes the therapeutic efficacy of CD19-
targeted T cells. *Nature* 1 (2018) doi:10.1038/s41586-018-0178-z.

24.   Michelozzi, I. M. *et al.* The Enhanced Functionality of Low-Affinity CD19 CAR T Cells Is
Associated with Activation Priming and Polyfunctional Cytokine Phenotype. *Blood* **136**,
52–53 (2020).

25.   Jackson, Z. *et al.* Sequential single cell transcriptional and protein marker profiling
reveals TIGIT as a marker of CD19 CAR-T cell dysfunction in patients with non-Hodgkin's
lymphoma. *Cancer Discov.* (2022) doi:10.1158/2159-8290.CD-21-
1586/696778/SEQUENTIAL-SINGLE-CELL-TRANSCRIPTIONAL-AND-PROTEIN.

26.   Sheih, A. *et al.* Clonal kinetics and single-cell transcriptional profiling of CAR-T cells in
patients undergoing CD19 CAR-T immunotherapy. *Nat. Commun.* **11**, 1–13 (2020).

27.   Haradhvala, N. J. *et al.* Distinct cellular dynamics associated with response to CAR-T
therapy for refractory B cell lymphoma. *Nat. Med.* **28**, 1848–1859 (2022).

28.   Hu, Y. *et al.* Antigen multimers: Specific, sensitive, precise, and multifunctional high-
avidity CAR-staining reagents. *Matter* **4**, 1–24 (2021).

29.   Lesch, S. *et al.* T cells armed with C-X-C chemokine receptor type 6 enhance adoptive
cell therapy for pancreatic tumours. *Nat. Biomed. Eng. 2021 511* **5**, 1246–1260 (2021).

30.   Davis, A. M., Ramos, H. J., Davis, L. S. & Farrar, J. D. Cutting Edge: A T-bet-
Independent Role for IFN-α/β in Regulating IL-2 Secretion in Human CD4 + Central
Memory T Cells. *J. Immunol.* **181**, 8204–8208 (2008).

31.   Kim, H. J. *et al.* Stable inhibitory activity of regulatory T cells requires the transcription
factor Helios. *Science (80-.).* **350**, 334–339 (2015).

32.   Lee, J. C. *et al.* In vivo inhibition of human CD19-targeted effector T cells by natural T
regulatory cells in a xenotransplant murine model of B cell malignancy. *Cancer Res.* **71**,
2871–2881 (2011).

33.   Ciucci, T. & Bosselut, R. Gimap and T cells: A matter of life or death. *Eur. J. Immunol.* **44**,
348–351 (2014).

34.   Evrard, M. *et al.* Sphingosine 1-phosphate receptor 5 (S1PR5) regulates the peripheral
retention of tissue-resident lymphocytes. *J. Exp. Med.* **219**, (2021).

35.   Zhang, L. *et al.* Etanercept as a new therapeutic option for cytokine release syndrome
following chimeric antigen receptor T cell therapy. *Exp. Hematol. Oncol.* **10**, 4–7 (2021).

36.   Lukhele, S. *et al.* The transcription factor IRF2 drives interferon-mediated CD8+ T cell
exhaustion to restrict anti-tumor immunity. *Immunity* **55**, 2369-2385.e10 (2022).

37.   Daniels, M. A., Luera, D. & Teixeiro, E. NFκB signaling in T cell memory. *Front. Immunol.*
14, 1–12 (2023).

38.   Park, S. L. & Mackay, L. K. Bhlhe40 Keeps Resident T Cells Too Fit to Quit. *Immunity* **51**,
418–420 (2019).

39.   Lynn, R. C. *et al.* c-Jun overexpression in CAR T cells induces exhaustion resistance.
Nature **576**, 293–300 (2019).

40.   Stelekati, E. *et al.* Long-Term Persistence of Exhausted CD8 T Cells in Chronic Infection
Is Regulated by MicroRNA-155. *Cell Rep.* **23**, 2142–2156 (2018).

41.   Chen, Z. *et al.* In vivo CD8+ T cell CRISPR screening reveals control by Fli1 in infection
and cancer. *Cell* **184**, 1262-1280.e22 (2021).

42.   Zhou, M., Clayberger, C. & Krensky, A. M. Krüppel-Like Transcription Factor 13
Regulates T Lymphocyte Survival In Vivo. *J. Immunol.* **178**, 5496–5504 (2007).

43.   Xu, B. *et al.* The Chromatin Remodeler BPTF Activates a Stemness Gene-Expression
Program Essential for the Maintenance of Adult Hematopoietic Stem Cells. *Stem Cell*

905    *Reports* **10**, 675–683 (2018).

906   44.   Wu, B. *et al.* BPTF Is Essential for T Cell Homeostasis and Function. *J. Immunol.* **197**,
907     4325–4333 (2016).

908   45.   Mamonkin, M., Puppi, M. & Lacorazza, H. D. Transcription factor ELF4 promotes
909     development and function of memory CD8+ T cells in Listeria monocytogenes infection.
910     *Eur. J. Immunol.* **44**, 715–727 (2014).

911   46.   Jenne, C. N. *et al.* T-bet-dependent S1P5 expression in NK cells promotes egress from
912     lymph nodes and bone marrow. *J. Exp. Med.* **206**, 2469–2481 (2009).

913   47.   Walker, A. J. *et al.* Tumor Antigen and Receptor Densities Regulate Efficacy of a
914     Chimeric Antigen Receptor Targeting Anaplastic Lymphoma Kinase. *Mol. Ther.* **25**,
915     2189–2201 (2017).

916   48.   Rodriguez-Marquez, P. *et al. CAR density influences antitumoral efficacy of BCMA CAR*
917     *T cells and correlates with clinical outcome. Sci. Adv* vol. 8 https://www.science.org
918     (2022).

919   49.   Wilson, T. L. *et al.* Common trajectories of highly effective CD19-specific CAR T cells
920     identified by endogenous T cell receptor lineages. *Cancer Discov.* (2022)
921     doi:10.1158/2159-8290.CD-21-1508/3166017/cd-21-1508.pdf.

922   50.   Giles, J. R. *et al.* Human epigenetic and transcriptional T cell differentiation atlas for
923     identifying functional T cell-specific enhancers. *Immunity* **55**, 557-574.e7 (2022).

924   51.   Chen, P. H. *et al.* Activation of CAR and non-CAR T cells within the tumor
925     microenvironment following CAR T cell therapy. *JCI Insight* **5**, (2020).

926   52.   Blank, C. U. *et al.* Defining 'T cell exhaustion'. *Nat. Rev. Immunol.* **19**, 665–674 (2019).

927   53.   Giles, J. R. *et al.* Shared and distinct biological circuits in effector, memory and
928     exhausted CD8+ T cells revealed by temporal single-cell transcriptomics and epigenetics.
929     *Nat. Immunol.* **23**, 1600–1613 (2022).

930   54.   Shah, N. N. *et al.* Clonal expansion of CAR T cells harboring lentivector integration in the
931     CBL gene following anti-CD22 CAR T-cell therapy. *Blood Adv.* **3**, 2317–2322 (2019).

932   55.   Goldberg, L. *et al.* Single-cell analysis by mass cytometry reveals CD19 CAR T cell
933     spatiotemporal plasticity in patients. *Oncoimmunology* **11**, (2022).

934   56.   Li, X. *et al.* Single-Cell Transcriptomic Analysis Reveals BCMA CAR-T Cell Dynamics in a
935     Patient with Refractory Primary Plasma Cell Leukemia. *Mol. Ther.* **29**, 645–657 (2021).

936   57.   Chavez, J. C., Bachmeier, C. & Kharfan-Dabaja, M. A. CAR T-cell therapy for B-cell
937     lymphomas: clinical trial results of available products. *Ther. Adv. Hematol.* **10**,
938     2040620719841581 (2019).

939   58.   Chen, G. M. *et al.* Integrative bulk and single-cell profiling of pre-manufacture T-cell
940     populations reveals factors mediating long-term persistence of CAR T-cell therapy.
941     *Cancer Discov.* candisc.1677.2020 (2021) doi:10.1158/2159-8290.cd-20-1677.

942   59.   Zhao, Z. *et al.* Structural Design of Engineered Costimulation Determines Tumor
943     Rejection Kinetics and Persistence of CAR T Cells. *Cancer Cell* **28**, 415–428 (2015).

944   60.   Maude, S. L. *et al.* Tisagenlecleucel in Children and Young Adults with B-Cell
945     Lymphoblastic Leukemia. *N. Engl. J. Med.* **378**, 439–448 (2018).

946   61.   Mackensen, A. *et al.* Anti-CD19 CAR T cell therapy for refractory systemic lupus
947     erythematosus. *Nat. Med.* **28**, 2124–2132 (2022).

948   62.   Stoeckius, M. *et al.* Simultaneous epitope and transcriptome measurement in single cells.
949     *Nat. Methods 2017 149* **14**, 865–868 (2017).

950   63.   What is Cell Ranger? -Software -Single Cell Gene Expression -Official 10x Genomics
951     Support. https://support.10xgenomics.com/single-cell-gene-
952     expression/software/pipelines/latest/what-is-cell-ranger.

953   64.   Hao, Y. *et al.* Integrated analysis of multimodal single-cell data. *Cell* **184**, 3573-3587.e29
954     (2021).

955   65.   Korsunsky, I. *et al.* Fast, sensitive and accurate integration of single-cell data with

956   Harmony. *Nat. Methods* **16**, 1289–1296 (2019).

957   66. Squair, J. W. *et al.* Confronting false discoveries in single-cell differential expression. *Nat.*
958   *Commun.* **12**, (2021).

959   67. Wu, T. *et al.* clusterProfiler 4.0: A universal enrichment tool for interpreting omics data.
960   *Innov.* **2**, (2021).

961   68. Subramanian, A. *et al.* Gene set enrichment analysis: A knowledge-based approach for
962   interpreting genome-wide expression profiles. *Proc. Natl. Acad. Sci. U. S. A.* **102**, 15545–
963   15550 (2005).

964   69. Aibar, S. *et al.* SCENIC: Single-cell regulatory network inference and clustering. *Nat.*
965   *Methods* **14**, 1083–1086 (2017).

966   70. Van de Sande, B. *et al.* A scalable SCENIC workflow for single-cell gene regulatory
967   network analysis. *Nat. Protoc. 2020 157* **15**, 2247–2276 (2020).

968   71. PBMC10k_downstream-analysis.
969   http://htmlpreview.github.io/?https://github.com/aertslab/SCENICprotocol/blob/master/not
970   ebooks/PBMC10k_downstream-analysis.html.

971   72. Andreatta, M. & Carmona, S. J. UCell: Robust and scalable single-cell gene signature
972   scoring. *Comput. Struct. Biotechnol. J.* **19**, 3796–3798 (2021).

973   73. Kuhn, M. Building Predictive Models in R Using the caret Package. *J. Stat. Softw.* **28**, 1–
974   26 (2008).

975   74. Tianqi, C. & Guestrin, C. XGBoost: A Scalable Tree Boosting System. *KDD* (2016).

976   75. Lundberg, S. M. & Lee, S. I. A Unified Approach to Interpreting Model Predictions. *Adv.*
977   *Neural Inf. Process. Syst.* **2017-December**, 4766–4775 (2017).

978   76. Liu, Y. & Just, A. SHAPforxgboost: SHAP Plots for 'XGBoost'. (2020).

979   77. ImmunoMind Team. immunarch: An R Package for Painless Bioinformatics Analysis of T-
980   Cell and B-Cell Immune Repertoires. (2019) doi:10.5281/zenodo.3367200.

981   78. Kochenderfer, J. N. *et al.* Construction and preclinical evaluation of an anti-CD19
982   chimeric antigen receptor. *J. Immunother.* **32**, 689–702 (2009).

 MAIN FIGURES AND LEGENDS

Figure 1. CD8⁺ CAR T cells from complete responders undergo a clonotypic shift in vivo.
(**a**) Schematic depicting sorting strategy, data generation, and single-cell multi-modal analysis of
CAR T cells in peripheral blood from seven CAR T-cell therapy patients (P1-7) who exhibited
complete responses with axicabtagene ciloleucel. (**b**) Representative flow plots depicting anti-
CD3 and CD19 antigen-tetramer staining of peripheral blood mononuclear cells from P3 versus
a healthy donor. CD3⁺Tet⁺ ("CAR") and CD3⁺Tet⁻ ("ENDO") patient cells were sorted from the
indicated gates. (**c**) Violin plots depicting normalized CAR transgene mRNA expression of sorted
CAR and ENDO T cells, split by CD4⁺ (left) and CD8⁺ (right) subsets. Expression levels were
compared by Wilcoxon Rank-Sum test, whereby **** indicates p<0.0001. (**d**) Line plots depicting
expansion and contraction of peripheral blood CAR abundance over the course of therapy. (**e-f**)
Bar graphs and heatmaps depicting overlap coefficients for TCR clonotypes comparing CAR (left)
and ENDO (right) repertoires between T_{exp}, T_{per1}, and T_{per2}. Overlap coefficients were compared
by *t*-test, whereby * indicates p<0.05 and ns indicates not significant.

Figure 2. Phenotypic heterogeneity of peripheral blood CD8$^+$ and CD4$^+$ CAR T cells. (**a**, **e**)
UMAPs depicting single-cell transcriptomes of CD8$^+$ (**a**) and CD4$^+$ (**e**) CAR T cells colored by cell
cluster. Inset depicts distribution of transcriptomes across timepoints. (**b**, **f**) Violin plots depicting
normalized expression levels of key genes and proteins for annotating and phenotyping CD8$^+$ (**b**)
and CD4$^+$ (**f**) CAR T cells. For extended versions, see figure S5a and S6a. (**c**, **g**) Stacked bar
graphs depicting proportions of each CD8$^+$ (**c**) and CD4$^+$ (**g**) CAR T-cell phenotype at different
timepoints. (**d**, **h**) Boxplots depicting proportion of CD8$^+$ (**d**) and CD4$^+$ (**h**) CAR T cells of a given
phenotype at different timepoints. Each dot represents a measurement from a single patient.
Proportions are compared between timepoints by Wilcoxon Rank-Sum test, whereby *** indicates
p<0.001, ** indicates p<0.01, * indicates p<0.05, and ns indicates not significant. Abbreviations:
CM, central memory; EM, effector memory; TE, terminal effector; Mem, memory; ISG, interferon
stimulated genes; Th1, type 1 helper-like

Figure 3. Integration of clonotypic and phenotypic shifts supports a two-stage differentiation model. All figure panels are based on the clonotype-phenotype linked dataset (n=32432 cells). (**a**) Proportion of CAR T cells with an exhausted-like effector memory (EM-exh), terminal effector (TE), or other phenotypes at each timepoint. (**b**) Total abundance of all clones that were predominantly EM-exh at T_{exp} (left), TE at T_{per1} (middle), or TE at T_{per2} (right), measured across timepoints. A clone's predominant phenotype was defined as the phenotype with the greatest representation. Each point represents clones from a patient (n=6). Abundance across timepoints were compared by repeated measures ANOVA with post-hoc t-tests. (**c**) Overlay of clonal abundance dynamics with 95% confidence intervals for clones annotated as EM-exh from T_{exp} (red) or TE from T_{per1}/T_{per2} (blue). (**d**) Pie chart depicting proportion of the top 500 clones classified as Wave 1, Wave 2, or Other. (**e**) Heatmap depicting normalized CAR abundance across timepoints for the top 500 largest clones. (**f**) Clonal dynamics and phenotypic distribution for the top 500 clones, grouped into Wave 1 and Wave 2. Left panels depict the change in clonal abundance across timepoints, while right panels show the predominant phenotype distribution at each timepoint. (**g**) Cartoon summarizing the two-stage model for CAR T-cell differentiation. Bulk CAR T-cell expansion and contraction (black line) masks the dynamics of Wave 1 (EM-exh, expansion phase timeframe, red) and Wave 2 (TE, persistence phase timeframe, blue) clones.

1030

Figure 4. Transcriptional signatures and regulatory networks of CD8$^+$ CAR T cells at T$_{exp}$ and T$_{per}$. (**a**) Heatmap depicting correlation between pseudo-bulk transcriptome of each sample (patient by timepoint). Samples were ordered along columns and rows by hierarchical clustering. Transcriptomes of day 7 samples from Maus et al. were added for external validation. (**b**) Stacked bar graph depicting label transfer of timepoint (T$_{exp}$ or T$_{per}$) from this study's dataset onto single-cell transcriptomes of samples from Maus et al. (**c**) Gene set enrichment analysis comparing CAR T cells between T$_{exp}$ and T$_{per}$. Gene sets were ordered by direction of upregulation and magnitude of enrichment. (**d**) Tile map depicting normalized expression of genes (columns) among different cell clusters at T$_{exp}$ and T$_{per}$ (rows). Genes were manually grouped into modules according to known functions. (**e**) Schematic for regulon construction and T$_{exp}$ versus T$_{per}$ classification. After transcriptomes were transformed into regulomes, regulon scores were calculated to train a machine-learning model to classify CAR T cells from T$_{exp}$ and T$_{per}$. Key regulons were identified based on importance for the model's predictions. (**f**) Bar graph depicting the SHapley Additive exPlanation (SHAP) values for the top eight regulons underlying T$_{exp}$ and T$_{per}$ predictions. (**g**) Tile map depicting normalized signature scores of regulons (columns) among different cell clusters at T$_{exp}$ and T$_{per}$ (rows). Regulons were grouped as T$_{exp}$-determining (left) and T$_{per}$-determining (right). (**h**, **i**) Target networks for the top eight T$_{exp}$-determining (**h**) and T$_{per}$-determining regulons (**i**). In each regulatory network, only the top differentially expressed genes are depicted. Each gene is colored according to log$_2$ fold-change between expression in T$_{exp}$ (red) and T$_{per}$ (blue).

Figure 5. Infusion product precursors of peripheral blood CD8⁺ CAR T cells. (**a**) UMAP
depicting single-cell transcriptomes of infusion product CD8⁺ T cells colored by cell cluster. (**b**, **c**)
Density maps (**b**) and violin plots (**c**) depicting expression levels of key genes and proteins for
annotation and phenotyping. For extended version, see figure S11a-b. (**d**) Cartoon depicting
identification of Pre-T$_{exp}$ and Pre-T$_{per}$ using endogenous TCR clonotypes as unique indices. (**e**)
Stacked bar graph depicting proportion distribution of all infusion product (top row) or precursors
of peripheral blood CD8⁺ CAR T cells (bottom two rows) among the four cell clusters. (**f**) Colored
UMAPs depicting distribution of Pre-T$_{exp}$, Pre-T$_{per}$, and non-linked infusion product cells ("IP only")
on the overall UMAP. (**g**) Tile map depicting normalized expression of genes (columns) among
Pre-T$_{exp}$, Pre-T$_{per}$, and IP only infusion product cells (rows). Genes were manually grouped into
modules according to known functions. (**h**) Violin plots depicting expression levels of select
differentially expressed gene sets between Pre-T$_{exp}$, Pre-T$_{per}$, and IP only infusion product cells.
Expression levels were compared by Wilcoxon Rank-Sum test, whereby **** indicates $p<0.0001$
and ** indicates $p<0.01$. (**i**) Gene set enrichment analysis comparing Pre-T$_{exp}$ and Pre-T$_{per}$. Gene
sets were ordered according to direction of upregulation and magnitude of enrichment. (**j**)
Enrichment plots for select gene sets differentially expressed between Pre-T$_{exp}$ and Pre-T$_{per}$
infusion product cells. (**k**) Cartoon depicting fates of CD8⁺ CAR T cells over the entire course of
therapy, from infusion product precursors to peripheral blood CAR T cells at T$_{exp}$ and T$_{per}$.
Abbreviations: NES, normalized enrichment score

1070

Figure 6. Transcriptomic signatures of exhausted-like EM CD8$^+$ CAR T cells. (**a**, **b**) UMAP
depicting single-cell transcriptomes of peripheral blood CD8$^+$ T cells colored by cell cluster (**a**) or
density contours of each cluster (**b**). The exhausted-like EM cluster is located on the lower half of
the UMAP. (**c**) Density maps depicting expression levels of major T-cell genes and proteins,
divided into categories. In the "receptor" category, proteins are placed directly beneath the
corresponding gene. (**d**) Flow plots for validating transcriptomic data. Plots depict expression of
CD39 and CD57 (top row) or CX3CR1 (bottom row) in CD8$^+$ CAR T cells at each timepoint from
P2. Exhausted-like EM CAR T cells were expected to be T_{exp}-specific with high CD39 and low
CD57/CX3CR1 expression. (**e**, **f**) Heat map depicting normalized expression of major gene sets
from Wherry et al. (**e**). Expression of two of the gene sets were depicted as violin plots (**f**), ordered
by decreasing expression level per cluster. Expression levels were compared to that of the cluster
with highest expression via Wilcoxon Rank-Sum test, with p-values adjusted for multiple
hypotheses testing using the Benjamini-Hochberg method, whereby **** indicates p<0.0001, ***
indicates p<0.001. (**g**) Volcano plot depicting differentially expressed genes between EM and
exhausted-like EM CD8$^+$ CAR T cells. Genes were colored according to direction of upregulation.
(**h**) Enrichment plots for select gene sets differentially expressed between EM and exhausted-like
EM CD8$^+$ CAR T cells.