1   **Detection of dedifferentiated stem cells in *Drosophila* testis**

2

3   Muhammed Burak Bener[1], Boris M. Slepchenko[1, 2]* and Mayu Inaba[1]*

4   1.   Department of Cell Biology, University of Connecticut Health, Farmington, CT 06030

5   2.   Richard D. Berlin Center for Cell Analysis and Modeling, University of Connecticut School

6   of Medicine, Farmington, CT 06030

7   * Correspondence: inaba@uchc.edu, boris@uchc.edu

8

9   **Abstract**

10

11   Tissue homeostasis relies on the stable maintenance of the stem cell pool throughout an

12   organism's lifespan. Dedifferentiation, a process in which partially or terminally differentiated

13   cells revert to a stem cell state, has been observed in a wide range of stem cell systems, and it has

14   been implicated in the mechanisms for stem cell maintenance. Dedifferentiated stem cells are

15   morphologically indistinguishable from original stem cells, making them challenging to identify.

16   Therefore, whether dedifferentiated stem cells have any distinguishable characteristics compared

17   with original stem cells is poorly understood. The *Drosophila* testis provides a well-established

18   model to study dedifferentiation. While our previous live imaging analyses have identified

19   dedifferentiation events constantly occurring at steady state, existing genetic marking methods

20   fail to detect most of the dedifferentiated stem cells and thus significantly underestimate the

21   frequency of dedifferentiation events. Here, we established a genetic tool with improved

22   sensitivity and used live imaging and mathematical modeling to evaluate the system. Our

findings indicate that the specificity of lineage-specific promoters is critical for successfully

identifying dedifferentiated stem cells.

Introduction

Tissue homeostasis and repair rely on a stable stem cell pool, and deficiencies in stem

cells lead to tissue degeneration, dysfunction and disease. Therefore, it is essential to maintain

the stem cell pool throughout the lifespan of an organism. Dedifferentiation, where partially or

terminally differentiated cells revert to the stem cell state, is one mechanism proposed to support

stem cell maintenance across various tissues during both steady-state and regeneration conditions

[1-9]. Given its role in stem cell maintenance, studying dedifferentiation and identifying

dedifferentiated stem cells is essential. However, the underlying mechanisms of dedifferentiation

remain poorly understood.

A key challenge in identifying dedifferentiated stem cells is that they are often

morphologically and molecularly similar to the original stem cells [4]. Genetic marking, such as

a lineage tracing technique, has been employed to distinguish dedifferentiated stem cells from

the original stem cells [3-5, 10-14]. Genetic marking often relies on promoters active in

differentiated cells, but not in stem cells, to induce recombination of reporter cassettes, resulting

in the permanent marking of these cells. Ideally, recombination should remain completely silent

in stem cells while being sufficiently high in targeted cells for accurate marking. However, this

has been proven to be difficult because many cell-stage-specific markers exhibit overlapping

expression across stages (reviewed in [2]), resulting in either leaky reporter activation in stem

cells due to basal recombinase expression or insufficient reporter activation in differentiating

cells, reducing the accuracy of tracing tools [15].

The *Drosophila* testis serves as an excellent model for studying *in vivo* stem cell maintenance [16]. Each testis contains 8 to 12 germline stem cells (GSCs) around a cluster of post-mitotic cells called the hub [17]. GSCs typically divide asymmetrically, producing one daughter cell that remains attached to the hub and retains its stem cell identity, while the other daughter cell is detached from the hub (called gonialblasts or GBs) and starts the differentiation process [18]. A GB undergoes four rounds of mitosis to form cysts of 2-, 4-, 8-, and 16-cell spermatogonia (SG) [19].

Dedifferentiation can be experimentally induced using forced differentiation of stem cells in *Drosophila* testis. When the differentiation factor, Bag of Marbles (Bam), is temporarily expressed, all GSCs start differentiation and physically leave the hub. After the treatment, cells detached from the hub start migrating back to the hub and regaining stem cell identity [1]. This phenomenon was first described by Brawley et al., in 2004 [1]. However, how frequently dedifferentiation occurs under steady-state conditions is still unclear.

Current genetic marking tools detect dedifferentiated GSCs at low frequencies [10, 11]. All previous tools utilized the bamGal4 driver to drive FLP recombinase (UAS-FLP) expression and permanently mark differentiating cells (Figure 1B-D). The low efficiency of marking is likely due to the delay of FLP expression, which may not reach the threshold for effective recombination until the 4-cell SG stage or later. Since GBs are the primary source of dedifferentiation [9], such delay significantly affects the marking efficiency. Consequently, these existing tools may fail to capture dedifferentiated GSCs across the fly's lifespan fully [10, 11].

In this study, we aimed to develop tools to improve efficiency. Specifically, we utilized FLP directly driven by the Bam promoter combined with FRT-flanked cassettes. We tested two FRT cassettes, each exhibiting a distinct threshold sensitivity to FLP dose. To evaluate the

efficiency and specificity of this system, we conducted long-term live imaging. We found that

the new FLP transgene has better efficiency, albeit with incomplete marking of dedifferentiated

stem cells in the *Drosophila* testicular niche.

Results

Improvement of genetic marking system for identifying dedifferentiated GSCs.

In our previous study, we investigated the mechanisms of GSC replenishment in the

Drosophila testis niche under steady-state conditions using long-term live imaging (~16 hours)

[9]. Among all the events observed in the niche (~6 total events per recording), 72% were

asymmetric cell division (ACD), 14% were losses (symmetric differentiation or stem cell loss

without division), 4% were symmetric renewal, and 10% were dedifferentiation (Figure 1A).

Given this 10% dedifferentiation rate observed over 16-hour imaging period (~0.6

dedifferentiation events per recording), we estimate that roughly ~1 GSC will be supplied via

dedifferentiation every day [9].

Previous studies have employed lineage-tracing tools to detect dedifferentiated GSCs [10,

11]. These systems rely on bamGal4-driven Flippase (FLP) expression that removes FRT-

flanked stop codons located between the promoter and the downstream reporter coding sequence.

When FLP-recombined cells dedifferentiate and become GSCs, these cells will be positive for

LacZ or GFP [10, 11]. We first tested this previous system using nos-FRT-mCherry-stop-FRT-

Gal4 cassette with UAS-GFP (hereafter referred to as nos>>GFP). On average, GFP-positive

GSCs comprised only approximately 10% of the total GSC pool at 28 days post-eclosion (Figure

1B-D), similar to the previous report (~14% shown in [11]). Considering the frequency observed

in live imaging (1~2/day [9]), this system might miss most dedifferentiation events. We

speculate that this issue was likely due to delays in FLP expression when the Gal4/UAS binary

system was used. Therefore, we sought to establish a new system.

To improve marking efficiency, we generated a transgenic fly to express FLPD5 (an FLP

variant with enhanced activity [20]) directly driven by the Bam promoter so that we can bypass

the Gal4-mediated UAS induction to minimize delays in inducing FLP recombinase activity

(Figure 1E).

100   Bam-FLPD5 causes spontaneous flipping events in GSCs, which obscures dedifferentiation

101   marking.

The newly established Bam-FLPD5 transgene showed a high frequency of GFP-positive

cells after introducing the same nos>>GFP reporter cassette as in the previous system (Figure

2A-D). To evaluate whether the marked GFP-positive GSC frequency reflects the

dedifferentiation rate, we conducted long-term live imaging of the niche to monitor GFP

activation. Testes expressing both the Bam-FLPD5, nos>>GFP, and Vasa-mCherry (a germline

cell marker) were imaged for approximately 12 hours using a z-stack to cover the entire niche. In

27 niches, each imaged for 12 hours, we observed eleven GSCs becoming GFP positive within

the niche (Figure 2E, Movie S1), even though the Bam promoter used to drive FLP expression

was expected to be active specifically in GB/SGs and not in GSCs. This is likely due to a

leaky/nonspecific driver activity in GSCs, resulting in unintended flipping events. Within the

same time period, the number of GB/SGs that became GFP-positive was twelve, indicating

almost no difference in the frequency of flipping events in the GSC and GB/SG pools. Given the

near-equal Bam-FLPD5 activity in GSC and GB/SG populations, we conclude that the majority

of the GFP-positive GSCs were not generated by dedifferentiation events, and therefore this

system is unsuitable for tracing dedifferentiated cells.

UAS-FRT-stop-FRT-mCD8-GFP cassette shows reduced spontaneous flipping rate in

GSCs.

We next strategized to overcome the issue of the nonspecific marking. Since the activity

of Bam promoter in GSCs is expected to be weak [21, 22], Bam-FLPD5 likely yields low levels

of FLP recombinase in the stem cells in the niche. On the other hand, the substrate constructs

with FRT sites may differ in their sensitivity to FLP recombinase activity [15]. Therefore, we

attempted to use an alternative reporter cassette with a higher threshold for FLP mediated

recombination to minimize spontaneous GFP activation in GSCs, so that it responds linearly to

FLP activity differences.

First, to test the recombination rates of different FRT cassettes, we compared the

recombination frequency of nos>>GFP to an alternative UAS-FRT-stop-FRT-mCD8-GFP,

together with nosGal4 (hereafter referred to as nos>>mCD8GFP) using heat-shock activatable

promoter-FLP (hs-FLP). We performed a single 30-min heat-shock treatment at 37°C to induce

FLP expression, then examined GFP positive GSC frequency after 24 hours [23, 24]. The

nos>>mCD8GFP yielded a significantly lower percentage of GFP-positive GSCs than

nos>>GFP indicating that nos>>mCD8GFP has a higher threshold to FLP and may be a good

candidate for Bam-FLPD5 substrate for marking of dedifferentiated cells (Figure S2A-C).

Then, we examined the frequency of GFP-positive GSCs using a combination of Bam-

FLPD5 and nos>>mCD8GFP and found that it indeed marks GSCs at a lower rate (Figure 3A-C)

than the Bam-FLPD5, nos>>GFP system (Figure 2D). To assess the recombination rate of Bam-

FLPD5 with nos>>mCD8GFP cassette in GSCs and in GB/SGs, we again employed a live

imaging approach. During imaging, we observed frequent GFP activation in differentiating cells,

including twelve GBs and four SGs, across 25 niches imaged for ~12 hours each (Figure 3D,

Movie S2). Notably, from these 25 recordings, we observed only one instance of GFP activation

that occurred in GSCs without prior dedifferentiation events (Figure 3D).

The observed reduction in spontaneous GFP activation in GSCs and efficient GFP

activation in GB/SGs suggests that the Bam-FLPD5 in combination with nos>>mCD8GFP

detects differentiating cells with higher specificity.

Quantifying specificity and efficiency of lineage marking

To estimate the number of GFP-positive cells that reflect true dedifferentiation events, we

formulated a computational model using observed rates of key cellular processes in our previous

live imaging study, including: asymmetric division (p), symmetric self-renewal (q), symmetric

differentiation (r), dedifferentiation (k_{dd}). Then, we defined flippase recombination rate in

GB/SGs to be (α) and recombination occurring in GSCs (α*) as the spontaneous background

event, disregarding stage-specific promoter activity. Because this background recombination

must also occur at the same rate in GB/SGs, the stage-specific recombination rate in GB/SGs

(α^{True}) is determined as ($\alpha^{True} = \alpha - \alpha$*) (all rates are per cell cycle, Figure S1A, B).

Using a publicly available software *Virtual Cell (*VCell*)* [25, 26], we first simulated the

time-dependent changes in the fraction of GFP-positive GSCs and validated the model by fitting

the curve to the experimentally observed GFP rates. For the data obtained by using both

nos>>GFP and nos>>mCD8GFP cassettes, the model outputs closely matched the experimental

data (Figure 4A, B), yielding values of α* that are consistent with the estimates obtained directly

from the experimental data, see methods section "Estimation of flippase recombination rate in

GSCs (α*)" for details. These validate the model and indicate its ability to accurately recapitulate

the frequency of marked cells in the niche.

Stage specificity of the recombination determines the feasibility of the detection system

We next used the model to estimated how many of GFP positive cells are true

dedifferentiation cells. When the nos>>GFP cassette was used, spontaneous flippase

recombination in GSCs occurs at a rate nearly equal to the flippase recombination in GB/SGs,

$\alpha^* \approx \alpha$, indicating that the flippase activity is entirely not stage specific. In this case, the model

revealed that dedifferentiation rate k_{dd} had no discernible effect on the frequency of GFP-

positive GSCs, as expected. Indeed, the predicted frequency of GFP-positive GSCs remained

essentially same across a range of k_{dd} values (0, 0.1, and 0.2) (Figure 4C), indicating that the

observed GFP-positive rate is not reporting the dedifferentiation rate at all.

In contrast, when nos>>mCD8GFP cassette was used, which has a significantly lower

flippase recombination rate in GSCs than in GB/SGs, $\alpha^* << \alpha$, the GFP positive rate showed the

clear sensitivity to the dedifferentiation rate k_{dd}, as the percentage of GFP-positive GSCs

significantly increases with the increasing k_{dd} (Figure 4D). These results highlight the

importance of the stage specificity of flippase excision events in marking the cells for estimating

dedifferentiation frequency.

An important aspect of lineage tracing is distinguishing true lineage-marked cells from

those marked due to non-specific flipping events (false positives). The model estimates the rate

of 'false positives' as GFP-positive GSCs originating from two sources: 1) GSCs that

spontaneously activate GFP within the niche and 2) dedifferentiated GFP-positive GSCs whose

lineage traces back to a spontaneous GFP activation event within and outside the niche. With the

Bam-FLPD5, nos>>GFP system, the model revealed that essentially all (~100%) of the observed

GFP-positive GSCs at all time points were false positives (Figure 4E, green line). In contrast,

with the nos>>mCD8GFP cassette, the fraction of false positives was reduced to about 45%

across all time points (Figure 4E, orange line). While still substantial, this represents a significant

improvement over the Bam-FLPD5, nos>>GFP system.

To further explore the relationship between the stage specificity of the flipping event and

false positive rates, we systematically varied the "flipping" ratio ($\alpha*/\alpha^{True}$) – the ratio of

background flippase activity in GSCs ($\alpha*$) to the stage-specific activity in GB/SGs (α^{True}) –

while keeping other parameters constant. We performed this analysis at a representative time

point (t=12 cycles), given that the probability of false positives is largely time-independent

(Figure 4E). As expected, low flipping ratios yield low false-positive rates approaching 0, but as

the flipping ratio increases, the probability of false grows rapidly, showing a sigmoid semi-log

dependence (Figure 4F). These findings indicate that minimizing background flippase activity is

a critical factor for the genetic marking system to be successful.

In summary, our modeling analysis demonstrates that the Bam-FLPD5, nos>>mCD8GFP

system reports dedifferentiation rates much more efficiently than the previous system, although

approximately half of the GFP positive GSCs are still false positives. The expected "true"

dedifferentiation rate, assuming zero background recombination in GSCs, and 100%

recombination in GB/SGs, is represented by the blue line in Figure 4B. This curve indicates that

the dedifferentiated cells rapidly occupy the entire niche within ~14 days of fly's age (Figure

4B). The difference between the blue and orange curves represents the false negative rate that

needs to be improved in future studies.

Experimentally increasing the dedifferentiation rates results in a high frequency of marked

GSCs

To experimentally test the model's predicted sensitivities to dedifferentiation, we

introduced a heat-shock-inducible Bam (hs-Bam) transgene into flies carrying either the Bam-

FLPD5, nos>>GFP or Bam-FLPD5, nos>>mCD8GFP lineage marking system. The hs-Bam

transgene is commonly used to induce dedifferentiation; a heat-shock treatment temporarily

forces GSCs to differentiate, and when treatment stops, GSCs are replenished through

dedifferentiation [27].

Our model predicts that the Bam-FLPD5, nos>>GFP system, with its high $\alpha*/\alpha^{True}$ ratio,

would be insensitive to changes in k_{dd} (Figure 4C), while the Bam-FLPD5, nos>>mCD8GFP

system, with its lower $\alpha*/\alpha^{True}$ ratio, would show an increase in GFP-positive GSCs with

increasing k_{dd} (Figure 4D).

At 7 days after eclosion, we observed a significant increase in GFP-positive GSCs in flies

carrying the hs-Bam transgene with the Bam-FLPD5, nos>>mCD8GFP system (Figure 5A, B,

E). This is likely due to the basal activity of the heat-shock promoter at room temperature,

leading to continuous Bam expression. This continuous Bam expression causes GSC loss, which

subsequently induces the replacement events likely through dedifferentiation. Notably, this

increase in GFP-positive GSCs is consistent with the model's prediction that Bam-FLPD5,

nos>>mCD8GFP system is sensitive to changes in the dedifferentiation rate.

In contrast, and as predicted by our model, increasing dedifferentiation frequency in flies

carrying the hs-Bam transgene with Bam-FLPD5, nos>>GFP system, did not result in a

significant change in GFP-positive GSCs under the same conditions (Figure 5C-E). These data

suggest that the observed increase in GFP-positive GSCs in the Bam-FLPD5, nos>>mCD8GFP

system reflects the dedifferentiation events.

Interestingly, the standard heat-shock protocol to induce dedifferentiation with hs-Bam at

37°C followed by a 3 to 7-day recovery period, did not result in a significant change in GFP-

positive GSCs with both nos>>GFP and nos>>mCD8GFP cassettes (Figure S3). We speculate

that high levels of Bam production during heat-shock treatments may cause negative feedback

mechanisms that suppress transcription from the Bam promoter used for the FLPD5 (Bam-

FLPD5), thereby reducing the efficiency of marking differentiating cells.

Discussion

Lineage tracing is a powerful tool for tracking cellular dynamics *in vivo*, including

proliferation, differentiation, and dedifferentiation. Recombinase-based systems, such as Cre-

loxP and FLP-FRT are widely used in lineage tracing. However, their reliance on tissue-specific

promoters makes them susceptible to "leaky" or off-target recombination. Studies in mice have

demonstrated the non-specificity of various Cre lines [28, 29]. Ideally, a lineage-specific

promoter should be distinctly active ("ON") in one target lineage and inactive ("OFF") in all

other lineages.

In this study, we developed an enhanced genetic marking system (Bam-FLPD5) to

identify dedifferentiated GSCs with improved sensitivity. By addressing key limitations of

previous lineage tracing methods, including delayed reporter activation that reduces marking

efficiency, our Bam-FLPD5 system offers a more accurate approach for monitoring

dedifferentiation events. While genetic marking is a valuable tool for studying dedifferentiation,

live imaging remains essential for confirming these events and distinguishing them from artifacts

like spontaneous reporter activation in off-target cells. Several past studies have utilized live

imaging to observe stem cell dynamics in real time, including studies in mouse spermatogonial

stem cells [30], and *Drosophila* male GSCs [31]. Through real-time tracing of cells within intact

tissues, these studies have provided insights into cellular plasticity and ability of partially or

terminally differentiated cells to revert to a stem cell state. In this study, we integrated live

imaging with our genetic marking approach to further enhance the accuracy and provide an

additional layer of validation.

Our findings emphasize spontaneous recombination's significant and previously

underestimated impact on lineage tracing studies. Spontaneous recombination likely occurs in

other lineage tracing systems but has often been overlooked due to the lack of real-time

validation methods like live imaging. Our simulations showed a strong positive correlation

between the false positive rates and the FLP recombination ratio ($\alpha*/\alpha$) (Figure 4F). For

example, the Bam-FLPD5, nos>>GFP system, where the estimated $\alpha*/\alpha$ ratio was ~1 (almost no

stage specificity of the recombination), our simulation showed essentially all of GFP-positive

GSCs (~100%) across 56 cycles are false positives as we expected. This suggests that none of

the observed GFP-positive GSCs in this system represent the true dedifferentiation events

(Figure 4E, green line). In contrast, the Bam-FLPD5, nos>>mCD8GFP system, with a lower

$\alpha*/\alpha$ ratio, exhibited a reduced false positive rate of approximately 45% (Figure 4E, orange line),

implying that roughly 55% of GFP-positive GSCs likely represent true dedifferentiation. This

enables the use of this system to monitor the dedifferentiation rate for future studies.

Although our new genetic marking system improved the sensitivity of previously

established systems, it still does not capture all dedifferentiated cells. A comparison between the

observed percentages of dedifferentiated GSCs in the Bam-FLPD5 system (Figure 4B, blue bars)

and the expected curve derived from our mathematical model (Figure 4B, blue line), which

assumes 100% recombination efficiency in all differentiating cells, reveals that a large portion of

differentiating cells remains unmarked with currently achieved recombination efficiency. There

is still a need for further research focused on identifying gene promoters with both high activity

and high specificity.

With the advances in next-generation sequencing and single-cell technologies,

computational-based lineage tracing tools have emerged (reviewed in [32, 33]). These include

Polylox DNA barcoding, which uses Cre recombinase to create diverse combinations barcodes to

uniquely label cells of interest for lineage tracing (although promoter specificity of Cre

expression remains a consideration) [34, 35]. Other approaches leverage naturally occurring

somatic mutations, such as those in mitochondrial DNA (mtDNA) [36], or engineered mutations

introduced via CRISPR-Cas9 systems [37] to infer cellular relationships. While DNA barcoding

and marker-independent methods offer valuable insights, they often lack the ability to directly

validate the lineage tracing events in real-time. A key advantage of our approach is the

integration of long-term live imaging with genetic marking. This combination allows for direct

assessment of accuracy, a capability lacking in many newer techniques.

GSCs typically divide asymmetrically, with several intrinsic factors non-randomly

segregated between the stem cell and its differentiating daughter cell [38-40]. However,

dedifferentiation disrupts this intrinsic asymmetry, and whether this results in differences

between original and dedifferentiated stem cells is not well understood. Moreover,

dedifferentiated stem cells have been implicated as a cellular origin of tumorigenesis in several

tissues [41, 42]. Lineage tracing tools, such as the Bam-FLPD5 system developed in this study,

provide a valuable approach to investigate these questions.

Materials and methods

Fly husbandry and strains

Flies were raised on standard Bloomington medium at 25°C (unless temperature control was required). 0- to 7-day-old adults were used for all experiments unless specific ages of flies were noted. The following fly stocks were obtained from Bloomington Stock Center (BDSC); nosGal4 (BDSC64277) [43]; 20XUAS-FLPD5 (BDSC55805); UAS(FRT.stop)mCD8-GFP (BDSC30032); hs-flp (BDSC1929) and hs-bam (BDSC24636). pVas-Vasa mCherry (FBtp0065762) [44] was a kind gift from Yukiko Yamashita. nos-FRT-mCherry-stop-FRT-Gal4, UAS-GFP [23] was from Inaba lab stock.

Immunofluorescence staining

Testes were dissected in 1x phosphate-buffered saline (PBS) (Fisher Scientific) and fixed in 4% formaldehyde (Electron Microscopy Sciences) in PBS for 30 minutes. Following fixation, testes were washed in PBS containing 0.2% TritonX-100 (Fisher Scientific) (PBST) for a minimum of 60 minutes (three washes of 20 minutes each). They were then incubated overnight at 4°C with primary antibody in 3% bovine serum albumin (BSA) (Sigma) in PBST. The following day, samples were washed in PBST for 60 minutes (three washes of 20 minutes each), incubated with secondary antibody in 3% BSA in PBST at room temperature for 2 hours, and then washed again in PBST for 60 minutes (three washes of 20 minutes each). Finally, samples were mounted using VECTASHIELD with 4',6-diamidino-2-phenylindole (DAPI) (Vector Lab). The primary antibodies used were as follows: rat anti-Vasa (1:20; developed by A. Spradling and D. Williams, obtained from Developmental Studies Hybridoma Bank (DSHB); mouse anti-hu-li

tai shao (Hts) (1:20, 1B1; DSHB) and mouse-anti- Fasciculin III (FasIII) (1:40, 7G10; DSHB).

AlexaFluor-conjugated secondary antibodies (Abcam) were used at a dilution of 1:400.

Long-term live imaging

Long-term live imaging was conducted using a previously established method [31]. Testes were

dissected in 1X Becker Ringer's [31] solution and mounted onto 35mm Glass Bottom Dishes

(Nunc). To prepare the imaging dish, 500 µL of 1 mg/mL poly-L-lysine (Sigma) was applied to

the coverslip area of the dish and incubated at room temperature for 5- to 7-hours. Following

incubation, the poly-L-lysine solution was replaced with the Becker Ringer's solution. Testes

were then dissected in Becker Ringer's solution and mounted on the poly-L-lysine coated surface

with the tips oriented downward. Next, Becker Ringer's solution was carefully removed, and

pre-warmed (room temperature) Schneider's *Drosophila* medium (supplemented with 10% fetal

bovine serum and glutamine–penicillin–streptomycin, Sigma) was added to the dish.

Z-stack images were acquired at 2 µm intervals (10-15 stacks total) using the Zeiss LSM800

Airyscan microscope with a 63× oil immersion objective (NA = 1.4) every 10 minutes over a 16-

hour period. The preset tiling function was utilized for automated image acquisition.

Image analyses

Airyscan time-lapse images were processed with Zen software (Zeiss). Each z-stack was

manually analyzed to track cells, record their GFP-expression status, and identify

dedifferentiation events. The hub area was identified as a circular, autofluorescent region located

near the testis tip, surrounded by fluorescently labeled germline stem cells. Dedifferentiation

events were recorded when germline cells, initially positioned more than one cell layer away

from the hub's edge reattached to the hub and maintained their attachment for at least 30

minutes. The moment a dedifferentiated cell observed at the hub edge was marked as the

"reattachment" time point. Movies were annotated using Veed.io.

Estimation of flippase recombination rate in GSCs (α*)

The flippase recombination rate in GSCs (α*) was determined directly from live-imaging data.

For the BamFLPD5, nos>>GFP system, 27 testes were imaged for 12 hours each (Figure 2E).

Each testis contains an average of 9 GSCs. Across all testes, 11 GSCs were observed to become

GFP-positive during the imaging period. Therefore, α* was calculated by dividing the total

number of flipping events in GSCs (11) by the total number of GSC cell cycles observed (27

testes*9 GSCs/testis=243 GSC cycles), yielding a value of 0.045 flipping events per cell cycle. A

similar procedure was followed for the Bam-FLPD5, nos>>mCD8GFP system, using its

respective experimental data in Figure 3D. Here, α* was calculated by dividing the total number

of flipping events in GSCs (1) by the total number of GSC cell cycles observed (25 testes*9

GSCs/testis=225 GSC cycles), yielding a value of 0.004 flipping events per cell cycle.

Short-term live imaging

Short-term live imaging was performed for static image acquisition of testes to avoid potential

signal loss or loss of extracellularly located fluorescent protein (GFP) during fixation and

permeabilization. Dissected testes were placed in Schneider's Drosophila medium (supplemented

with 10% fetal bovine serum and glutamine–penicillin–streptomycin, Sigma), then placed onto

two etched rings on Gold Seal Rite-On Micro Slides with media and covered with coverslips. Z-

stack images were acquired at 1 μm intervals (15-20 stacks total) using the Zeiss LSM800

Airyscan microscope with a 63× oil immersion objective (NA = 1.4). For these short-term live

imaging experiments, all images were taken within 30 minutes of dissection, without time-lapse

imaging.

GSC depletion and recovery

GSC depletion and recovery were performed as previously described, with modifications [27].

Adult flies (0- to 3-days old) carrying the hs-Bam transgene (BDSC24636) were raised at 22°C.

They underwent six heat shock treatments in a 37°C water bath for 30 minutes, twice daily,

within vials with fly food. Between treatments, the vials were maintained in a 29°C incubator.

Following the last heat shock treatment, the flies were returned to 22°C for the recovery phase.

Testes were dissected after a 5-day recovery period.

Generation of Bam-FLPD5 transgenic line

The minimum bam promoter sequence (-198 to transcription start site, TSS) [45] was amplified

from genomic DNA isolated from wild-type flies using the following primers:

SphI KpnI bam -198 Forward: 5'-

AGCGGATCCAAGCTTGCATGCGGTACCCCAAATCAGTGTGTATAATT-3'

Bam -198 Reverse: 5'-TATTCTTAAGTTAAATCACACAAATCACTCGAT-3'

Separately, the FLPD5 sequence, excluding the PEST sequence, was amplified from the genomic

DNA of the 20XUAS-FLPD5 strain (BDSC55805) using the following primers:

FLPD5 Forward: 5'-

GTGATTTGTGTGATTTAACTTAAGAATAATGCCGCAGTTTGATATCCTCTG-3'

FLPD5 Reverse: 5'-GTACCCTCGAGCCGCTTAAATACGGCGATTGATGTAGGAGCTCA-

3'

pUAST-attB vector was digested by SphI and NotI restriction enzymes to remove the UASt

promoter. The amplified bam -198 and FLPD5 fragments were assembled with the digested

vector using a Gibson Assembly kit (NEB) and sequenced. Transgenic flies were generated using

strain attP2 by PhiC31 integrase-mediated transgenesis (BestGene).

Mathematical modeling

Model

The model was developed for the analysis of efficiency, specificity, and sensitivity of

genetic constructs designed for the identification of dedifferentiated germline stem cells (GSCs)

in *Drosophila* testes.

The model outputs must mimic experimental readouts - time-dependent fractions of

marked dedifferentiated GSCs. Correspondingly, we formulate our model in terms of numbers of

unmarked and marked GSCs in the niche and corresponding GBs and SGs available for

dedifferentiation in the vicinity of the niche. The stem cell pool is presented in the model by the

following variables: the number of the innate unmarked stem cells, $n_0(t)$; the number of the

dedifferentiated unmarked stem cells, $n_1(t)$; the number of the dedifferentiated stem cells

marked as a result of the promoter activation and subsequent flippase recombination, $n_2(t)$; and

the number of the stem cells marked by spontaneous nonspecific GFP activation (false positives),

$n_3(t)$. The GB/SGs pool is described by the following variables: the number of unmarked

GB/SGs, $m_1(t)$; the number of GB/SGs marked due to the promoter activation, $m_2(t)$; and the

number of GB/SGs marked by spontaneous GFP activation, $m_3(t)$.

The variables described above are governed by the following key processes included in

the model (Figure S1A,B): asymmetric division (characterized by rate constant p), symmetric

renewal (q), symmetric differentiation/loss (r), dedifferentiation of GB/SGs (k_{dd}), differentiation

of GB/SGs (β), flippase recombination in GB/SGs (α), flippase recombination in GSCs (α^*), and

stage-specific flippase recombination in GB/SGs (α^{True}) ($\alpha^{\mathrm{True}} = \alpha - \alpha^*$) (all rates are per cell

cycle).

We assume that the processes involving unmarked and marked cells are described by

equivalent parameter values. We further assume that the niche tightly controls the total number

of GSCs, $N = n_0(t) + n_1(t) + n_2(t) + n_3(t)$, so that it remains constant at all times,

$$n_0(t) + n_1(t) + n_2(t) + n_3(t) = N = const. \tag{1}$$

Experimental fractions of GFP-positive cells were obtained from the pools containing large

numbers of cells for each time point (on the order of a hundred or more), so that the fluctuations

of their averages were relatively small. This allowed us to ignore the discreteness of the variables

and approximate the dynamics of the corresponding fractions, $s_i(t) = \frac{n_i(t)}{N}, i = 0,1,2,3$, and $g_j(t)$

$= \frac{m_j(t)}{N}, j = 1,2,3$, by differential equations:

$$\partial_t s_0 = -(r - q + \alpha^*)s_0$$

$$\partial_t s_1 = -(r - q + \alpha^*)s_1 + k_{dd}g_1, \tag{2}$$

$$\partial_t s_2 = -(r - q)s_2 + k_{dd}g_2,$$

$$\partial_t s_3 = -(r - q)s_3 + \alpha^*(s_0 + s_1) + k_{dd}g_3;$$

and

$$\partial_t g_1 = \nu(s_0 + s_1) - (\alpha + \alpha^* + \beta + k_{dd})g_1,$$

$$\partial_t g_2 = \nu s_2 + \alpha g_1 - (\beta + k_{dd})g_2, \tag{3}$$

$$\partial_t g_3 = \nu s_3 - (\beta + k_{dd})g_3 + \alpha^* g_1.$$

Parameter ν in Eqs (3) represents the stem-cell mitotic rate, $\nu = 1$ per cell cycle.

The system of Eqs (2) and (3) was solved using VCell (www.vcell.org). The VCell

implementation of the model can be found in the VCell database of public MathModels under

username "boris"; the model name is "Inaba_dedifferentiation_model_public".

Model Parameters

We constrained model parameters using the results of long-term live imaging, along with

the analysis of the model's salient features.

Not all parameters of the model are independent. Experimental observations indicate that

by the time of eclosion, the pool of differentiated cells available for dedifferentiation has

approached its steady-state size equaling N, i.e. $m_1(t) + m_2(t) + m_3(t) = N = const$ and

$\sum_{j=1}^{3} g_j(t) = 1$. This makes the mechanism of replenishing of GSCs by dedifferentiation

mathematically equivalent to symmetric renewal with an effective $q_{eff} = k_{dd} + q = r$ and $p +$

$q_{eff} + r = \nu$. Indeed, from Eq (1), $\sum_{i=0}^{3} s_i(t) = 1$. Then summing up Eqs (2) yields $k_{dd} =$

$\frac{(r-q)}{\sum_{j=1}^{3} g_j(t)} = r - q$, and since $\sum_{j=1}^{3} g_j(t) = 1$,

$$k_{dd} + q = r, \text{ and } p + 2r = \nu. \tag{4}$$

Eqs (4) are consistent with the data cited at the beginning of the section *Improvement of genetic*

marking system for identifying dedifferentiated GSCs. Correspondingly, $p = 0.72$/cycle and k_{dd}

$= 0.1$/cycle, then $r = \frac{1}{2}(\nu - p) = 0.14$/cycle and $q = r - k_{dd} = 0.04$/cycle.

Similarly, summing up Eqs (3) yields $v = \beta + k_{dd}$, and $\beta = 0.9$/cycle.

Fitting the model to experimental data

To determine parameters α^{True} and α^*, we fitted our model to experimental time

dependences of fractions of the GFP-positive GSCs obtained for a given genetic marker with an

experimentally constrained ratio $\alpha^*/\alpha^{\text{true}}$. For this, Eqs (2) and (3) should be solved with varying

α^{true} and initial conditions, to minimize the deviation between the solution of the model and the

experimental data. As the experimental data are available beginning with the fly's eclosion,

trying possible values of the seven model variables at the time of eclosion would be unwieldy.

Instead, we moved the zero time point to the time in the larvae stage at which stem cells start to

divide. It is reasonable to assume that at that time, all GCSs are unmarked, and the numbers of

differentiated cells are zero. This allowed us to solve the model with same initial conditions in all

trials and only vary the time of eclosion, τ_{ecl}. One caveat is that $k_{dd}(t) = \frac{(r-q)}{\sum_{j=1}^{3} g_j(t)}$ becomes

now a function of time, since $\sum_{j=1}^{3} g_j(t)$ is not at steady state at $t \leq \tau_{ecl}$. To avoid division by

zero in the equation for $k_{dd}(t)$, the initial conditions used in all our trials were as follows: $s_0(0)$

$= 1$, $s_i(0) = 0$ ($i = 1,2,3$), $g_0(0) = \varepsilon$, $g_j(0) = 0$ ($j = 2,3$), where ε is a small number (in our

simulations, $\varepsilon = 10^{-6}$). The results were then offset by τ_{ecl} for comparison with the experimental

data.

We characterized discrepancy between the simulation results and experimental data by

the root-mean-square deviation (RMSD) defined as $\delta = (n^{-1} \sum_{i=1}^{n} (y_{exp}(t_i) - y_{theor}(t_i))^2)^{1/2}$,

where n is the number of time points, for which we performed the comparison, $y_{exp}(t_i)$ are the

experimental values at times t_i, and $y_{theor}(t_i) = s_2(t_i) + s_3(t_i) \equiv s_+(t_i)$. Good fits,

characterized by $\delta \sim$ 0.02-0.04, were obtained with $\tau_{ecl} =$ 6 cycles.

Once fully parameterized, the model was used to explore sensitivities of the marking

systems to k_{dd} and α^{true}, and to compute the probability of false positives, $\frac{s_3(t)}{s_+(t)} = \frac{s_3(t)}{s_2(t) + s_3(t)}$.

Statistical analysis and graphing

No statistical methods were used to predetermine the sample size. The experiment values were

not randomized. The investigators were not blinded to allocation during experiments and

outcome assessment. All experiments were independently repeated at least 3 times to confirm the

results. Statistical analysis and graphing were performed using GraphPad prism9. The

illustrations were generated by using Adobe Illustrator or BioRender. All data are shown as

means ± s.d.

Acknowledgments

We thank Yukiko M. Yamashita, Michael Buszczak and the Bloomington *Drosophila* Stock

Center, and the Developmental Studies Hybridoma Bank for reagents. This research is supported

by R35GM128678 from the National Institute for General Medical Sciences and start-up funds

from UConn Health (to M.I.). B.M.S. was supported in part by R24GM137787 from National

Institute for General Medical Sciences.

Author Contributions

M.I. and M.B.B. conceived the project, designed and executed experiments and analyzed data.

B.M.S. and M.B.B. conducted mathematical modeling and data fitting using *Virtual Cell*. All

authors wrote and edited the manuscript.

Declaration of Interests

The authors declare no competing interests.

Figures

Figure 1. Improvement of genetic marking system for identifying dedifferentiated GSCs.

A) A schematic diagram illustrating the estimated rates of asymmetric division, symmetric renewal, differentiation, and dedifferentiation of GSCs under steady-state conditions within the testicular niche shown in our previous study [9]. **B)** An illustration of the bamGal4-driven FLP lineage tracing system. The bamGal4 driver induces FLP expression in differentiating cells (4-cell and later-stage SGs), where FLP excises the mCherry-stop cassette. If cells with the removed stop cassette revert to the GSC state, they can be identified by the GFP expression under the control of nosGal4 driver. **C**) Representative immunofluorescence images showing the testicular niche in bamGal4-driven FLP lineage tracing system in the males at 0 and 14 days post-eclosion. The testes were stained with Vasa (blue), FasIII (red), and Hts (red) antibodies. GFP-positive GSCs (green) are encircled by white dotted lines. **D**) A graph showing the percentage of GFP-positive GSCs per testis at 0, 7, 14, and 28 days post-eclosion. **E)** An illustration of the lineage tracing system where FLP recombinase is directly expressed under a promoter and excises the FRT sites from the reporter cassette. The reporter gene is expressed under a separate promoter. All scale bars are 10 μm. Asterisks in the images indicate the approximate location of the hub. "n" indicates the number of testes scored.

Figure 2. Bam-FLPD5 causes spontaneous flipping events in GSCs, which obscures dedifferentiation marking.

A) An illustration of the Bam-FLPD5 lineage tracing system. Here, FLPD5 expression is driven directly under the Bam promoter sequence (-198 to transcription start site, TSS) [45]. The Bam-FLPD5 is crossed with nos-FRT-mCherry-stop-FRT-Gal4, UAS-GFP cassette (nos>>GFP) **B, C)** Representative images of live testes of indicated ages of the Bam-FLPD5-driven lineage-tracing system. GFP and mCherry channels are shown in separate panels in **B'** (for GFP) and **B''** (for mCherry). **D)** A graph showing the scored percentage of GFP-positive GSCs per testis on days 0, 7, 14, 21, and 28 post-eclosion. **E)** Selected frames of time-lapse images from a 12-hour recording (interval:10min) of a niche showing the GFP-activation in a native GSC (corresponding to Movie S1). Germline cells are visualized by expression of Vasa-mCherry (magenta). White dotted lines encircle a GSC, becoming GFP-positive. Asterisks in the images indicate the approximate location of the hub. The table (right panel) shows the number of times GFP was observed to activate for each indicated cell stage (GSC, GB, and SG) during the 12-hour live imaging recording of 27 testes.

All scale bars are 10 μm. Asterisks in the images indicate the approximate location of the hub. "n" indicates the number of testes scored.

Figure 3. UAS-FRT-stop-FRT-mCD8-GFP cassette is efficiently flipped in GB/SGs but not

in GSCs.

A) An illustration of the Bam-FLPD5 lineage tracing system. Here, Bam-FLPD5 is combined

with the UAS-FRT-stop-FRT-mCD8-GFP cassette and nosGal4 (nos>>mCD8GFP). FLPD5

removes the stop codon from this cassette. If cells once removed stop cassette in the past become

GSCs, they can be identified by the GFP expression under the control of nosGal4 driver. **B)**

Representative immunofluorescence images of the testicular niche of the Bam-FLPD5 marking

system in the males at 0 and 14 days post-eclosion. The testes were stained with Vasa (blue) and

FasIII (red) antibodies. GFP-positive GSCs (green) are outlined by white dotted lines. **C)** A

graph showing the percentage of GFP-positive GSCs per testis at 0, 7, 14, and 28 days post-

eclosion. **D)** Selected frames from a 12-hour time-lapse recording (interval:10 min) of a niche

showing the GFP-activation in a GB (corresponding to Movie S2). Germline cells are visualized

by expression of Vasa-mCherry (magenta). White dotted lines encircle a GB becoming GFP

positive. The table (right panel) shows the number of times GFP was observed to activate for

each indicated cell stage (GSC, GB, and SG) during 12-hour live imaging recording of 25 testes.

All scale bars are 10 μm. Asterisks in the images indicate the approximate location of the hub.

"n" indicates the number of testes scored.

Figure 4. Quantifying specificity and efficiency of lineage marking

A) Model fitting for the Bam-FLPD5, nos>>GFP system. Experimental data showing the percentage of GFP-positive GSCs over time (blue bars; data from Figure 2D) are compared to the best-fit model output (orange curve). The model incorporates key cellular processes (see Figure S1A, B and Materials and Methods for details). The root mean squared deviation (RMSD) between the model and the experimental data is 0.026. Two cycles are equivalent to one day [46]. Fitted parameters (α^{True} and α^*) are shown in the graph. See Source Data file for numerical values and all parameter values.

B) Model fitting for the Bam-FLPD5, nos>>mCD8GFP system. Experimental data showing the percentage of GFP-positive GSCs over time (blue bars; data from Fig 3C) are compared to the best-fit model output (orange curve). The root mean squared deviation (RMSD) between the model and the experimental data is 0.035. The blue line represents the expected percentage of GFP-positive GSCs if the recombination efficiency was 100% in all differentiating cells ($\alpha^{True}=\infty$) and with no spontaneous GFP activation ($\alpha^*=0$)

C) Predicted effect of dedifferentiation rate (k_{dd}) on GFP-positive GSCs in the Bam-FLPD5, nos>>GFP system. Model predictions for the percentage of GFP-positive GSCs over time are shown for different dedifferentiation rates (k_{dd} = 0, 0.1, and 0.2 per cell cycle). The overlapping lines demonstrate that the system is insensitive to changes in k_{dd} due to high ratio of α^*/α^{True} = 11.

D) Predicted effect of dedifferentiation rate (k_{dd}) on GFP-positive GSCs in the Bam-FLPD5, nos>>mCD8GFP system. Model predictions for the percentage of GFP-positive GSCs over time are shown for different dedifferentiation rates (k_{dd} = 0, 0.1, and 0.2 per cell cycle). The lines demonstrate the system's sensitivity to changes in k_{dd} due to lower ratio of α^*/α^{True} = 0.0667.

E) Predicted probability of false positive GSCs over time. The model-predicted probabilities of a GFP-positive GSC being a false positive are shown for the Bam-FLPD5, nos>>GFP system (green line) and the Bam-FLPD5, nos>>mCD8GFP system (orange line). The nos>>GFP system shows a near 100% false positive probability, while the nos>>mCD8GFP system shows a significantly lower probability of approximately 45%.

F) Predicted probability of false positive GSCs as a function of the flipping ratio over time for two different marking systems with varying flipping ratios (α^*/α^{True}). The analysis was performed at a representative time point (t=12 cycles).

Figure 5. Introduction of a hs-Bam transgene results in high frequency of marked GSCs

A, B) Representative immunofluorescence images of the testis tip from 7-day-old flies with

Bam-FLPD5 marking system with the nos>>mCD8GFP cassette alone **(A)** or with an additional

single copy of the hs-Bam transgene **(B)**. Testes were stained with Vasa (blue) and FasIII (red)

antibodies. GFP-positive GSCs (green) are outlined with white dotted lines.

C, D) Representative images of live testis tip from 7-day-old flies with Bam-FLPD5 marking

system with the nos>>GFP cassette alone **(C)** or a combination with a single copy of hs-Bam

(D).

E) The graph shows the quantification of GFP-positive GSCs per testis in 7-day-old flies without

heat-shock treatment for the indicated genotypes.

All scale bars are 10 μm. Asterisks in the images indicate the approximate location of the hub. p-

values were calculated by Šídák's multiple comparisons test. "n" indicates the number of testes

scored.

References

1.   Brawley, C. and E. Matunis, *Regeneration of male germline stem cells by spermatogonial dedifferentiation in vivo.* Science, 2004. **304**(5675): p. 1331-4.
2.   Liu, Y. and Y.G. Chen, *Intestinal epithelial plasticity and regeneration via cell dedifferentiation.* Cell Regen, 2020. **9**(1): p. 14.
3.   Sun, Q., et al., *Dedifferentiation maintains melanocyte stem cells in a dynamic niche.* Nature, 2023. **616**(7958): p. 774-782.
4.   Tata, P.R., et al., *Dedifferentiation of committed epithelial cells into stem cells in vivo.* Nature, 2013. **503**(7475): p. 218-223.
5.   Tetteh, Paul W., et al., *Replacement of Lost Lgr5-Positive Stem Cells through Plasticity of Their Enterocyte-Lineage Daughters.* Cell Stem Cell, 2016. **18**(2): p. 203-213.
6.   Murata, K., et al., *Ascl2-Dependent Cell Dedifferentiation Drives Regeneration of Ablated Intestinal Stem Cells.* Cell Stem Cell, 2020. **26**(3): p. 377-390.e6.
7.   Merrell, A.J. and B.Z. Stanger, *Adult cell plasticity in vivo: de-differentiation and transdifferentiation are back in style.* Nat Rev Mol Cell Biol, 2016. **17**(7): p. 413-25.
8.   Kai, T. and A. Spradling, *Differentiating germ cells can revert into functional stem cells in Drosophila melanogaster ovaries.* Nature, 2004. **428**(6982): p. 564-569.
9.   Bener, M.B., A. Twillie, and M. Inaba, *Dedifferentiating germ cells regain stem-cell specific polarity checkpoint prior to niche reentry.* bioRxiv, 2023: p. 2023.04.26.538507.
10.   Herrera, S.C. and E.A. Bach, *JNK signaling triggers spermatogonial dedifferentiation during chronic stress to maintain the germline stem cell pool in the Drosophila testis.* eLife, 2018. **7**: p. e36095.
11.   Cheng, J., et al., *Centrosome misorientation reduces stem cell division during ageing.* Nature, 2008. **456**(7222): p. 599-604.
12.   Malagola, E., et al., *Isthmus progenitor cells contribute to homeostatic cellular turnover and support regeneration following intestinal injury.* Cell, 2024. **187**(12): p. 3056-3071.e17.
13.   Nagai, H., et al., *Nutrient-driven dedifferentiation of enteroendocrine cells promotes adaptive intestinal growth in Drosophila.* Developmental Cell, 2023. **58**(18): p. 1764-1781.e10.
14.   Van Es, J.H., et al., *Dll1+ secretory progenitor cells revert to stem cells upon crypt damage.* Nature cell biology, 2012. **14**(10): p. 1099-1104.
15.   Álvarez-Aznar, A., et al., *Tamoxifen-independent recombination of reporter genes limits lineage tracing and mosaic analysis using CreER(T2) lines.* Transgenic Res, 2020. **29**(1): p. 53-68.
16.   Fuller, M.T. and A.C. Spradling, *Male and female Drosophila germline stem cells: two versions of immortality.* Science, 2007. **316**(5823): p. 402-4.
17.   Greenspan, L.J., M. de Cuevas, and E. Matunis, *Genetics of gonadal stem cell renewal.* Annu Rev Cell Dev Biol, 2015. **31**: p. 291-315.
18.   Yamashita, Y.M., D.L. Jones, and M.T. Fuller, *Orientation of asymmetric stem cell division by the APC tumor suppressor and centrosome.* Science, 2003. **301**(5639): p. 1547-50.
19.   de Cuevas, M. and E.L. Matunis, *The stem cell niche: lessons from the Drosophila testis.* Development, 2011. **138**(14): p. 2861-9.

20.   Nern, A., et al., *Multiple new site-specific recombinases for use in manipulating animal genomes.* Proceedings of the National Academy of Sciences, 2011. **108**(34): p. 14198-14203.

21.   Cherry, C.M. and E.L. Matunis, *Epigenetic regulation of stem cell maintenance in the Drosophila testis via the nucleosome-remodeling factor NURF.* Cell Stem Cell, 2010. **6**(6): p. 557-67.

22.   Insco, M.L., et al., *Accumulation of a differentiation regulator specifies transit amplifying division number in an adult stem cell lineage.* Proceedings of the National Academy of Sciences, 2009. **106**(52): p. 22311-22316.

23.   Salzmann, V., et al., *Lineage tracing quantification reveals symmetric stem cell division in Drosophila male germline stem cells.* Cell Mol Bioeng, 2013. **6**(4): p. 441-448.

24.   Bener, M.B., B.M. Slepchenko, and M. Inaba, *Asymmetric stem cell division maintains the genetic heterogeneity of tissue cells.* bioRxiv, 2024: p. 2024.05.16.594576.

25.   Resasco, D.C., et al., *Virtual Cell: computational tools for modeling in cell biology.* Wiley Interdiscip Rev Syst Biol Med, 2012. **4**(2): p. 129-40.

26.   Slepchenko, B.M. and L.M. Loew, *Use of virtual cell in studies of cellular dynamics.* Int Rev Cell Mol Biol, 2010. **283**: p. 1-56.

27.   Sheng, X.R., C.M. Brawley, and E.L. Matunis, *Dedifferentiating Spermatogonia Outcompete Somatic Stem Cells for Niche Occupancy in the Drosophila Testis.* Cell Stem Cell, 2009. **5**(2): p. 191-203.

28.   Payne, S., S. De Val, and A. Neal, *Endothelial-Specific Cre Mouse Models.* Arteriosclerosis, Thrombosis, and Vascular Biology, 2018. **38**(11): p. 2550-2561.

29.   Magnuson, M.A. and A.B. Osipovich, *Pancreas-specific Cre driver lines and considerations for their prudent use.* Cell metabolism, 2013. **18**(1): p. 9-20.

30.   Nakagawa, T., et al., *Functional hierarchy and reversibility within the murine spermatogenic stem cell compartment.* Science, 2010. **328**(5974): p. 62-67.

31.   Sheng, X.R. and E. Matunis, *Live imaging of the Drosophila spermatogonial stem cell niche reveals novel mechanisms regulating germline stem cell output.* Development, 2011. **138**(16): p. 3367-76.

32.   Wu, S.S., J.H. Lee, and B.K. Koo, *Lineage Tracing: Computational Reconstruction Goes Beyond the Limit of Imaging.* Mol Cells, 2019. **42**(2): p. 104-112.

33.   Wagner, D.E. and A.M. Klein, *Lineage tracing meets single-cell omics: opportunities and challenges.* Nature Reviews Genetics, 2020. **21**(7): p. 410-427.

34.   Pei, W., et al., *Polylox barcoding reveals haematopoietic stem cell fates realized in vivo.* Nature, 2017. **548**(7668): p. 456-460.

35.   Pei, W., et al., *Using Cre-recombinase-driven Polylox barcoding for in vivo fate mapping in mice.* Nature Protocols, 2019. **14**(6): p. 1820-1840.

36.   Ludwig, L.S., et al., *Lineage Tracing in Humans Enabled by Mitochondrial Mutations and Single-Cell Genomics.* Cell, 2019. **176**(6): p. 1325-1339.e22.

37.   Kalhor, R., et al., *Developmental barcoding of whole mouse via homing CRISPR.* Science, 2018. **361**(6405).

38.   Yamashita, Y.M., *Nonrandom sister chromatid segregation of sex chromosomes in Drosophila male germline stem cells.* Chromosome Research, 2013. **21**(3): p. 243-254.

39.   Tran, V., et al., *Asymmetric division of Drosophila male germline stem cell shows*
asymmetric histone distribution. Science, 2012. **338**(6107): p. 679-82.

40.   Yamashita, Y.M., et al., *Asymmetric inheritance of mother versus daughter centrosome*
in stem cell division. Science, 2007. **315**(5811): p. 518-21.

41.   Schwitalla, S., et al., *Intestinal tumorigenesis initiated by dedifferentiation and*
acquisition of stem-cell-like properties. Cell, 2013. **152**(1-2): p. 25-38.

42.   Friedmann-Morvinski, D., et al., *Dedifferentiation of neurons and astrocytes by*
oncogenes can induce gliomas in mice. Science, 2012. **338**(6110): p. 1080-4.

43.   Van Doren, M., A.L. Williamson, and R. Lehmann, *Regulation of zygotic gene expression*
in Drosophila primordial germ cells. Curr Biol, 1998. **8**(4): p. 243-6.

44.   Lerit, D.A. and E.R. Gavis, *Transport of germ plasm on astral microtubules directs germ*
cell development in Drosophila. Curr Biol, 2011. **21**(6): p. 439-48.

45.   Ridwan, S.M., et al., *Diffusible fraction of niche BMP ligand safeguards stem-cell*
differentiation. Nature Communications, 2024. **15**(1): p. 1166.

46.   Gadre, P., et al., *The rates of stem cell division determine the cell cycle lengths of its*
lineage. iScience, 2021. **24**(11): p. 103232.

797