High-resolution HIV-1 m^{6}A epitranscriptome reveals splicing-dependent methylation clusters and unique 2-LTR transcript modifications

Delphine Naquin*, Sandra Blanchet*, Erwin van Dijk, Lisa Bertrand, Sylvie Grégoire, Cecilia Bertha Ramirez, Rania Ouazahrou, Yan Jaszczyszyn, Arnaud Moris, Claude Thermes, Céline Hernandez, Olivier Namy

ABSTRACT

The N^6-methyladenosine (m^{6}A) modification of HIV-1 RNAs plays an essential role in regulating viral infection. This modification has been widely studied but the number and precise positions of the m^{6}A sites remain unclear due to the lack of precision of detection methods. We used the latest Nanopore chemistry and direct m^{6}A base-calling option to identify 18 m^{6}As, 14 of which were located at the 3' end of the genome, the other four being located in central regions. Our data reveal differential methylation of these positions between splicing isoforms. Eleven of these sites are clustered in two short segments (197 nt and 43 nt) with peak-shaped methylation profiles. Single-molecule analysis revealed that a very small number of transcripts were unmethylated in both clusters (≤ 1.5% of spliced transcripts and ≤ 5.4% of unspliced viral RNA). We also identified a ~732 nt RNA species resulting from the transcription of non-integrated viral DNA circles closed by two long terminal repeats (2-LTR circles). These transcripts started in the first LTR, terminated at the polyA site of the second LTR and harbored six m^{6}A sites. Five of these sites were present in other transcripts and, remarkably, had the highest methylation rates. The sixth site was methylated only in this transcript, suggesting a role for this RNA in HIV-1 infection. These findings reveal a new landscape of HIV m^{6}A transcriptome modifications and pave the way for studies deciphering their role in the viral life cycle.

INTRODUCTION

Epitranscriptomic RNA modifications play crucial roles in diverse cellular and viral processes and diseases (1). The N6-methyladenosine (m^{6}A) modification occurs frequently, in all types of cellular RNA, and consists of the covalent addition of a methyl group to the N6 position of adenosine. This methyl group addition is catalyzed by

a multi-subunit methyltransferase complex with a catalytic core composed of methyltransferase-like 3

(METTL3) and methyltransferase-like 14 (METTL14) (2). This writer complex preferentially targets adenosines

in the consensus sequence 5′-DRACH-3′ (D = A/G/U, R = A/G, H = U/A/C) in mammalian mRNAs (3). DRACH

motifs are frequent in mRNA but are not always targeted, as only about 10% of these motifs were found to be

m^6A-methylated in cells. This suggests that there are probably other context-dependent signals for the

methylation of specific DRACH motifs. The m^6A modification is reversible and can be removed by erasers,

such as alkB homolog 5 (ALKBH5) in particular (4). Reader proteins recognize m^6A and determine the

functional consequences of the modification for the fate of the RNA. The readers capable of binding m^6A

include the YTH domain-containing family of proteins, the most frequently studied members of which are

YTHDC-1, YTHDF1,2,3 (5, 6). However, discrepancies remain concerning the function of these proteins (7). In

mammals, m^6A is the most prevalent modification involved in gene expression regulation and RNA

metabolism, including mRNA transport, localization, stability, translation efficiency and processing (8-13), but

this modification is also present in viral RNA (14). Several studies have detected methylated regions (m^6A) in

the HIV-1 genome with techniques based on methylated RNA immunoprecipitation (15-21). However, it was

not possible to localize the methylated positions with single-nucleotide precision because the resolution of the

techniques used ranged from 20 nt to more than 200 nt. There were also differences between studies, with

some reporting peaks distributed along most of the length of the genome (15, 18), whereas others detected

peaks only at the 5' and 3' ends (17, 19) or only at the 3' end (16). These discrepancies may reflect differences

between the cell types or infection protocols (cell transfection with proviral plasmids or infection with viruses)

used or a lack of reproducibility of antibody-based methods (22). In principle, Oxford Nanopore Technologies

(ONT) direct RNA sequencing (23) would allow the detection of modified RNA nucleotides at single-molecule

level. Several studies have provided diverse bioinformatics tools for the detection of RNA modifications based

on various strategies (base-call errors induced by modifications, altered current intensity, modification-aware

base-calling models, reviewed in (24)). However, improvements in data quality and recovery rates for full-

length transcripts are required to enable direct base-calling for the modified nucleotides (25). Here, we

considered the full-length genome, including the 5'LTR and 3'LTR sequences. A recent study using the

RNA002 kit resulted in the development of a modified library protocol and major improvements in the recovery

of full-length RNA molecules (26). Combining several bioinformatics tools based on different properties of the

ionic current, the authors identified three m^6A sites at the 3' end of the viral genome. We made use of an

advance developed by ONT — RNA004 chemistry (new nanopore and new motor protein) — that greatly

improves overall yield and data quality (27), and new base-calling models making it possible to call m^{6}As

directly with the four standard bases. Using these developments, we characterized the HIV-1 m^{6}A methylation

pattern of HIV-1 mRNAs in infected CD4$^+$ T cells, including a CD4$^+$ T cell line (SupT1) and, importantly,

primary CD4$^+$ T cells. We extended the detection to 18 m^{6}A sites at single-nucleotide resolution, single-

molecule level and in different splicing isoforms, providing the most complete and precise description of the

landscape of HIV-1 m^{6}A methylation to date.

RESULTS

Nanopore sequencing of full-length HIV-1 transcripts and identification of splicing isoforms

For the detection of HIV-1 m^{6}A sites, cellular transcripts from SupT1 and primary CD4$^+$ T cells infected with

HIV-1 were extracted and sequenced by ONT direct RNA sequencing (DRS, Materials and Methods). It has

been reported that it is very difficult to obtain reads corresponding to full-length HIV RNA molecules with the

standard DRS library preparation protocol (26). In our previous DRS experiments with the standard protocol,

we obtained less than 0.008% and 0.06% of total reads for our first two samples, corresponding to the ~9.6 kb

unspliced HIV-1 genome. In this study, we used a modified protocol described elsewhere (26) to increase

recovery rates for full-length transcripts. We used two biological replicates of SupT1 cells (rep1 and rep2)

harvested 20 h and 30 h post-infection and primary CD4$^+$ T cells isolated from the blood of two different blood

donors collected 48 h and 72 h post-infection (Figure S1). PolyA RNA was isolated from each of these six

infected samples and sequenced to generate 7.3 to 28.9 million total reads (Table S1) and between 33,587

and 164,360 reads corresponding to full-length HIV transcripts (Table S1). Full-length reads were examined for

the presence of splice junctions. We identified eight known donor sites (four major sites — D1, D2, D3, D4 —

together with D1c, D2b, D4a and D4b) and eleven acceptor sites (four major sites — A1, A2, A5, A7 —

together with A1b, A3, A4a, A4b, A4c, A5a and A5c) (28-31) (Table S2A). The different isoforms were identified

by the succession of donor and acceptor sites (Materials and Methods); we identified 51 isoforms, as

previously described (31, 32) (Table S2B). These isoforms are typically classified into three groups on the

basis of size: (i) unspliced ~9 kb transcripts (US), (ii) partially spliced transcripts (~4 kb, PS, lacking intron 1

and retaining intron 4) including Vif2, Vpr3,4, Env/Vpu and Tat5, and (iii) completely spliced transcripts (~2 kb,

CS, lacking introns 1 and 4) including Vif1, Vpr1,2, Tat1-4, Rev and Nef. The relative abundances of the

various isoforms were similar in all samples (correlation R>0.86 in all comparisons, Table S3).

Detection of m^6A sites in all HIV-1 transcripts

The first step was the detection of m^6A sites in all HIV-1 reads. The positions of N6-methyladenosine

nucleotides were determined directly at read level during the base-calling process, providing a first estimate of

methylation rates at each position in a set of transcripts. These methylation rates were compared with those

for the corresponding positions in the unmodified RNAs transcribed *in vitro* (Figure 1A). We performed this

analysis for all HIV-1 transcripts from the six samples, resulting in a set of 14 putative m^6A sites, all located in

the 3' region of the HIV-1 genome common to all transcripts (Tables 1, S4; Figure 1B). The numbering used in

this paper corresponds to that of the full-length genome, including the 5'LTR and 3'LTR sequences

(transcription start site in position +454). The corresponding methylation landscape presented two clusters of

sites. The first cluster consisted of eight sites (A8436 to A8632) with two highly methylated central sites (64-

80%), and the second cluster consisted of three sites (A9400, A9428, A9442) with a highly methylated central

site (79-84%). Three additional sites displaying low rates of methylation were detected between these two

clusters. Remarkably, the segment containing the first cluster (8436 to 8632) was found to harbor five DRACH

sequences (3, 33), all of which were methylated. The other three sites of this segment correspond to 1-nt

variants of the DRACH motif (found here at positions A8487, A8555, and A8572) interspersed with the

canonical DRACH sites (Tables 1, S4; Figure 1B). It should be noted that m^6As have also been detected in

non-DRACH motifs, especially the variant GGAC<u>G</u> (found here at positions A8487 and A8572), which has

been found to be highly methylated in human mRNAs (34, 35).

m^6A position	Sequence	Methylation rate (%)
A8436	AGACA	9..7
A8442	AGACA	10.3
A8487	GGAC<u>G</u>	30
A8532	AGACT	79.3
A8555	GGA<u>TT</u>	19.3
A8563	GAACT	79.3
A8572	GGAC<u>G</u>	23.7
A8632	GAACT	10.3
A9024	TGACT	7.3
A9074	GGACT	8.7
A9163	GAACT	17.3
A9400	GAACT	29.7
A9428	GGACT	84.3
A9442	GGACT	53.3

107 **Table 1.** List of the 14 m^{6}A sites detected in the 3' region (8360-9700) common to all transcripts. Methylation

108 rates were measured in the SupT1_20h_pi rep1 sample. The underlined nucleotides are single-nucleotide

109 variants of the consensus DRACH sequence.

111
Figure 1. Detection of m^{6}A sites in all HIV-1 transcripts. **(A)** The IGV screenshot shows the methylation rate

measured in the cluster 1 region of the HIV-1 genome including sites A8436 to A8632. Line 1, methylation rate

measured directly during the m^{6}A base-call of SupT1_20h_pi rep1. Line 2, methylation rate measured directly

for the in vitro sample in the same region. Line 3, Difference (line 1) - (line 2); red arrows indicate the eight

sites detected; negative values are set to zero; sites with rates < 3% were not retained (black arrow). Vertical

lines, positions of As in the five DRACH sequences of the region. **(B)** Methylation profile of all HIV-1 transcripts

in the 3' region (8360-9700). Closed red circles, SupT1_20h_pi rep1; empty red circles, SupT1_30h_pi rep1;

closed green circles, SupT1_20h_pi rep2; empty green circles, SupT1_30h_pi rep2; closed black circles,

CD4_48h_pi; empty black circles, CD4_72h_pi. Vertical lines, positions of As in the 33 DRACH sequences of

the region.

123 **Efficiency of m⁶A detection in synthetic oligonucleotides**

124 We assessed the efficiency of m⁶A site detection by subjecting two synthetic RNA oligonucleotides with

125 sequences corresponding to HIV-1 segments but harbouring m⁶A at positions 8532 and 8563 (oli1-mod) and

126 at positions 9400, 9428 and 9432 (oli2-mod) to the detection procedure. As controls, we analyzed the same

127 HIV RNA oligonucleotides without m⁶A modifications. Remarkably, all but one (position A9400) of the

128 methylated As were identified (Figure 2, blue bars). The m⁶A site at position A9400 is located too close (9 nt)

129 to the 5' end of oli2-mod, which prevents its accurate detection (25). The methylated A8563, A9428 and A9442

130 positions were detected with a methylation rate of ~100% in both oligomers (Figure 2). The lower methylation

131 rate (~75%) observed for A8532 is probably due to its position close (13 nt away) to the 5' end of oli1-mod.

132 These observations confirm the specificity and high level of efficiency of m⁶A detection with RNA004 chemistry

133 and the m⁶A base-calling option. We further assessed the quality of m⁶A detection by treating oli1-mod and

134 oli2-mod harboring m⁶A modifications with the ALKBH5 demethylase and then performing the detection

135 process. Methylation rates were significantly lower at all sites, confirming the efficiency of the m⁶A detection

136 process (Figure 2, red bars).

137

138

139 *Figure 2. Efficiency of m⁶A site detection with methylated synthetic oligonucleotides. In blue, methylation*

140 *rates, measured at the indicated m⁶A positions, comparing unmodified and m⁶A-modified oli1-nmod and oli2-*

141 *nmod. In red, the same measurement after the demethylation of synthetic m⁶A-modified oligonucleotides with*

142 *ALKBH5 (Materials and Methods). *, p-value < 10⁻⁶; ***, p-value < 10⁻¹⁴⁹.*

143

144 **Validation of the candidate m⁶A sites with the ALKBH5 demethylase**

For further validation of the detection of m^{6}A sites in viral RNAs from HIV-1-infected cells, replicates 1 and 2 of

SupT1 cells infected for 20 h (SupT1_20h_pi rep 1 and rep2) were treated with the ALKBH5 demethylase,

sequenced and the results compared with those of the untreated samples. Remarkably, all 14 sites had

significantly lower methylation rates (Table S5). Overall, these results, obtained with synthetic oligonucleotides

and viral mRNA from HIV-1 infected cells, highlight the quality of our detection method and confirm that the 14

m^{6}A modification sites identified in HIV-1 transcripts are *bona fide* m^{6}A modification sites.

Detection of m^{6}A sites in HIV-1 splicing isoforms

We then investigated the distribution of m^{6}A sites in each of the different splicing isoforms. The 14 m^{6}A sites

found in the pool of all transcripts were also detected in each of the 51 isoforms identified in the six samples

(Figure 3A, Table S6, Figures S2A-F, S3A-E). The methylation rate of these sites was similar to that for all

transcripts, characterized by cluster 1 centered around the two highly methylated A8532 and A8563 sites and

cluster 2 centered around the highly methylated A9428. An unsupervised clustering of viral transcripts based

on the methylation rates observed at the 14 m^{6}A sites showed that the isoforms clustered into three groups.

The first group corresponded to the US transcripts. The second group consisted mostly of PS transcripts and

the third group consisted of the CS transcripts (Figure 3A, Figure S3A-E). Methylation rates differed

significantly between these classes and were highest for CS and lowest for US (CS rates > PS rates > US

rates, Figure 3B, Figure S4A-E). These results are consistent with previous observations showing that

methylation levels are higher in CS than in PS transcripts, and higher in PS than in US transcripts (17, 19, 26).

A

B

165
166

Figure 3. *Methylation rates for the 14 m^{6}A sites detected in SupT1_20h_pi rep1. **(A)** Unsupervised heatmap of*

the methylation rates of HIV-1 transcripts (Materials and Methods). Isoforms are clustered based on the

In parallel to the 14 sites common to all viral mRNA isoforms, we detected four additional m^{6}A sites specific to

only a few isoforms. Three of these sites, A5613, A5654 and A5703, were detected in the Vif and Vpr coding

regions (between the D3 and A3 splice sites) and are specific to the isoforms encoding Vif and Vpr. The fourth

site, A5887, was detected in the Tat coding region (between A3 and A4c) and is specific to the Vif, Vpr, Tat and

SORF1 isoforms (Figure 4A, Table 2, Table S7). The size of the corresponding RNAs depends strongly on the

absence (CS, -i4) or presence (PS, +i4) of intron 4 (i4: D4A7, 2325 nt-long). As for the global trend observed

for the 14 sites common to all isoforms presented in Figure 3, the methylation rates of these four additional

m^{6}A sites was higher for the CS (-i4) than for the PS (+i4) isoforms, but this difference was more pronounced

(Tables 2, S7). For example, in the SupT1_20h_pi rep1 sample, the methylation rates of the four sites in the

Vif1 isoform (-i4) were 3 to 9 times higher than those in the Vif2 isoform (+i4) (Table 2). Similarly, the

methylation rates in Vpr1 and Vpr2 were much higher than those in Vpr3 and Vpr4, respectively. There was

one notable exception to this tendency: the rate of A5887 methylation was significantly higher in Tat 5 (PS,

+i4) than in Tat1 (CS, -i4) (Figure 4B). Importantly, all four sites were unmethylated or had very low rates of

methylation in unspliced transcripts (≤ 1.3%). A similar trend was observed for the other five SupT1 and CD4

samples, for the CS and PS transcripts of the Vif, Vpr and Tat isoforms (Table S7).

187
188
189

190
191

192 **Figure 4.** *Position and methylation rates of the four isoform-specific m⁶A sites. (A) The A5613, A5654 and*

193 *A5703 sites are common to the Vif and Vpr isoforms; A5887 is common to the Vif, Vpr, Tat and SORF1*

194 *isoforms. (B) Methylation rates in SupT1_20h_pi rep1, calculated for the four sites in the CS (- intron 4, Vif1,*

195 *Vpr1, Vpr2), PS (+ intron 4, Vif2, Vpr3, Vpr4) and US isoforms. (C) Methylation rates calculated for the A5887*

196 *site for the CS (Tat1, 2, 3, 4, SORF1), PS (Tat5) and US isoforms.*

Isoform		SupT1_20h_pi rep1			
CS - i4	PS + i4	% A5613 GGACA	% A5654 AGACA	% A5703 AAACT	% A5887 AAACT
Vif1		35.2	2.4	11.4	18.6
	Vif2	9.3	0.7	3.3	2
Vpr1		58	9.3	24	25.3
	Vpr3	12.3	1	4	0.3
Vpr2		37.3	4.6	10	15.9
	Vpr4	5.7	2.3	3.7	0
Tat1					18.3
	Tat5				28.7
Tat2					12.3
Tat3					12.3
Tat4					9.5
SORF1					16.7
US		1.3	0	0	0

Table 2. *Methylation rates measured for the A5613, A5654, A5703 and A5887 m^6A sites in the Vif, Vpr, Tat and sORF1 isoforms in the SupT1_20h_pi rep1 sample. The first two columns correspond to the isoforms completely spliced (CS, - i4) or partially spliced (PS, + i4). The last row indicates the methylation rates in the unspliced transcripts.*

Single-molecule analysis of m^6A sites

We investigated the distribution of the methylated m^6A sites in individual transcripts by counting, for US, PS and CS transcripts, the numbers of methylated As in the eight sites of cluster 1 (from 0 to 8 methylated As) (Figure 5A) and the three sites of cluster 2 (from 0 to 3 methylated As) (Figure 5B) of the 3'UTR region of HIV-1, as defined in Figure 1. We examined the two clusters either separately (Figure 5A, 5B, Figure S6A-D) or simultaneously on each read (Figure 5C-E, Figure S7A-O, Table S8). The distribution of m^6A sites was similar for rep1 and rep2 samples, for the SupT1 samples at 20 h and 30 h post-infection, and also in SupT1 and primary CD4[+] T cells. We found that the number of sites modified per molecule increased from US to PS to CS. When considering the two clusters simultaneously, we observed that the peaks for the US transcripts corresponded to two methylated sites in cluster 1 and one or two methylated sites in cluster 2 (Figure 5A,B, Figure S6A-D). By contrast, for the PS and CS transcripts, the peaks corresponded to two or three methylated

216 sites in cluster 1 and two methylated sites in cluster 2. In the six samples, a small proportion (3.7% to 5.4%) of

217 US transcripts were totally unmethylated simultaneously in the two clusters. The proportion of unmethylated

218 transcripts decreased to 1.5% for PS transcripts and 0.3% for CS transcripts (Table S8).

220
221

222
223

Figure 5. *Single-molecule analysis of methylated m^6A sites. **(A)** Cluster 1. **(B)** Cluster 2. **(C-E)** Correlation*

between the number of methylated sites in cluster 1 and in cluster 2 of the same transcript from the

SupT1_20h_pi rep1 sample. The intensity of the color on the heatmaps indicates the number of reads showing

the indicated numbers of methylated As in cluster 1 (0 to 8) and in cluster 2 (0 to 3) (Materials and Methods).

Detection and analysis of 2-LTR transcripts

During the analysis of the splicing isoforms, we identified a class of RNA transcripts mapping to the long

terminal repeats (LTR, constituted by the U3-R-U5 segments) located at either end of the viral genome. They

result from the transcription of 2-LTR circles, a non-integrated circular form of the reverse-transcribed viral

DNA formed early after infection during the conversion of the viral genome into DNA by the junction of the two

LTRs located at the ends of the genome (36) (Figure 6A, 6B). They begin at the promoter at the 5' end of the R

segment of the 3' LTR and terminate at the polyA site at the 3' end of the R segment of the 5' LTR. They create

a precise join between the 3' end of the RU5 segment of the 3' LTR and the 5' end of the U3R segment of the

5' LTR. Studies of the 2-LTR U5-U3 junction (the junction between the two LTRs) have shown this junction to

be highly heterogeneous in infected cells and between infected individuals, frequently due to insertions (37,

38). Transcripts encompassing this junction have been detected by RT-PCR, but were not precisely identified.

A recent study of the HIV-1 transcriptome by Nanopore cDNA sequencing identified RNA species that could

correspond to 2-LTR transcripts and were detected in infected CD4$^+$ T and HeLa cells (31). Here, the 2-LTR

transcripts were detected in both infected SupT1 and infected primary CD4$^+$ T cells (0.16% to 1.69 % of HIV-1

transcripts, Table S2B). They displayed either the precise U5-U3 junction or a number of variants mostly

carrying short (1-nt to 10-nt) insertions. The detection of m^{6}A sites in these 2-LTR transcripts revealed the

presence of six sites that were detected in all six samples (Figure 6B, 6D, Table S9). The first site, 2L125, is

homologous to the A578 position of the viral genome (Figure 6C) common to all HIV-1 transcripts.

Remarkably, this 2L125 site was methylated in 24 to 38% of 2-LTR transcripts (Table S9) but was not detected

as methylated in any other HIV-1 transcript, in all six samples. The other five sites are located in the U3

segment of the 5' LTR and are homologous to m^{6}A sites A9074, A9163, A9400, A9428 and A9442, which are

common to all splicing isoforms (Fig. 6B-D). Remarkably, in the 2-LTR transcripts, the methylation rates of all

five of these sites were higher than those of the homologous sites in the viral genome (Table S9). For

example, the 2L537 site attained methylation rates of 91.3% and 94.1% in SupT1 and CD4$^+$ T-cell samples,

respectively (Table S9), whereas the homologous site A9428 had methylation rates of 84.3% and 79.7% in the

corresponding samples (Table S4).

A

B

2-LTR transcript: R-U5-U3-R

```
1
R-->
GGGTCTCTCTGGTTAGACCAGATCTGAGCCTGGGAGCTCTCTGGCTAACTAGGGAACCCACTGCTTAAGCCTCAATAAAGCTT

        <--R U5-->                          2L125(578)
GCCTTGAGTGCTCAA AGTAGTGTGTGCCCGTCTGTTGTGTGACTCTGGTAACTAGAGATCCCTCAGACCCTTTTAGTCAGTGT

        <--U5||U3-->
                2L183(9074*)
GGAAAATCTCTAGCAGACTGGAAGGGCTAATTCACTCCCAAAGAAGACAAGATATCCTTGATCTGTGGATCTACCACACACAA

        2L272(9163)
GGCTACTTCCCTGATTGGCAGAACTACACACCAGGGCCAGGGGTCAGATATCCACTGACCTTTGGATGGTGCTACAAGCTAGT

ACCAGTTGAGCCAGATAAGGTAGAAGAGGCCAATAAAGGAGAGAACACCAGCTTGTTACACCCTGTGAGCCTGCATGGAATGG

ATGACCCTGAGAGAGAAGTGTTAGAGTGGAGGTTTGACAGCCGCCTAGCATTTCATCACGTGGCCCGAGAGCTGCATCCGGAG

        2L509(9400)              2L537(9428)   2L551(9442)
TACTTCAAGAACTGCTGACATCGAGCTTGCTACAAGGGACTTTCCGCTGGGGACTTTCCAGGGAGGCGTGGCCTGGGCGGGAC

                                        <--U3 R-->
TGGGGAGTGGCGAGCCCTCAGATGCTGCATATAAGCAGCTGCTTTTTGCCTGTACT GGGTCTCTCTGGTTAGACCAGATCTGA

                                                R<-- -->polyA
GCCTGGGAGCTCTCTGGCTAACTAGGGAACCCACTGCTTAAGCCTCAATAAAGCTTGCCTTGAGTGCTTCAA AAAAAAA...
```

Figure 6. *Transcription and methylation of the 2-LTR transcripts.* **(A)** *Diagram of the 2-LTR circles and of the transcripts joining the RU5 segment of the 3' LTR to the U3R segment of the 5' LTR. Several examples of insertion variants at the U5-U3 junction are shown.* **(B)** *Sequence of the 2-LTR transcripts. The successive transcribed regions — R, U5, U3, R — are indicated; m^6A sites are shown in bold typeface and underlined and are named according to the position of the methylated A in the 2-LTR transcript; in brackets, the homologous position in the HIV-1 reference genome; for example, for 2L125(578), 125 is the position of the m^6A in the 2-LTR transcript and (578) is the homologous position in the HIV-1 reference genome; * indicates that the sequence context of 2L183 in the 2-LTR transcript (AG\underline{A}CT, see B) differs from that at the homologous position 9074 in the reference genome (GG\underline{A}CT).* **(C)** *Correspondence table for a position in the 2-LTR transcript and the homologous position in the reference genome.* **(D)** *Sites detected in the 2-LTR transcripts of infected SupT1 and CD4$^+$ T-cell samples (Materials and Methods). Large red circles, SupT1_20h_pi rep1; orange circles, SupT1_30h_pi rep1; blue circles, SupT1_20h_pi rep2; green circles, SupT1_30h_pi rep2; yellow circles, CD4_48h_pi; black circles, CD4_72h_pi. Vertical lines indicate the positions of the As in all the DRACH motifs (21) of the transcript.*

DISCUSSION

We report here the detection of m^6A nucleotides in HIV-1 transcripts from infected human cells, with comparison to previous studies (15-19, 26). We included cells of two different origins: CD4$^+$ T cells from the SupT1 cell line and CD4$^+$ T cells from the peripheral blood mononuclear cells (PBMC) of blood donors. These cells, which are natural HIV-1 targets, were infected for different periods of time (20 h and 30 h for SupT1

cells; 48 h and 72 h for blood donor cells). Using the new RNA004 chemistry and a new base-calling model,

we detected 14 m^{6}A sites common to the unspliced HIV-1 RNA and to all the splicing isoforms encoding viral

proteins.

The m^{6}A positions were clustered in two regions of the 3' end of the HIV-1 genome, both having a methylation

profile featuring highly methylated central sites. These sites are partly included in the methylated regions

identified in previous studies based on the immunoprecipitation of m^{6}A-containing RNA (reviewed in (9)) or on

m^{6}A-SAC-seq (39). Unlike these studies, we found no methylation sites in the 5' region of the reference HIV-1

genome, either in the unspliced RNA or in the diverse splicing isoforms. Overall, we observed a remarkable

specificity of the 14 m^{6}A sites in the 245 DRACH sites of the HIV-1 genome. The saturation of the five DRACH

sites located in the 200 nt cluster 1 and the additional methylation of three single-nucleotide DRACH variants

in the same region further increase the specificity of HIV-1 methylation. However, despite the proposal of

hypotheses involving the recruitment of m^{6}A writers by RNA-binding proteins or by RNA polymerase II close to

DRACH sites during transcription (7, 40), there is currently no explanation for the specificity of the methylation

of some DRACH sites but not others. Interestingly, on examination of the secondary structure of the HIV-1

RNA, most of the 14 m^{6}A sites were found to be located in single-stranded portions of the structure, or within

the final nucleotides at the 3' end of a helix (Figure S8). This finding is fully consistent with studies showing

that RRACH sequences located in a double-stranded RNA duplex prevent efficient m^{6}A methylation by the

METTL3–METTL14 complex (41). The 14 m^{6}A sites include three sites (A8532, A9428, A9442) previously

detected as peaks in RNA immunoprecipitation experiments (16-18) and as methylated nucleotides (m^{6}As) by

direct RNA sequencing (26). This last study also identified A8563 as a weakly methylated m^{6}A site, contrasting

with our results, as A8563 was one of the highly methylated sites of cluster 1 (Figure 1). We found that the 14

m^{6}A sites had higher methylation rates in CS isoforms than in PS isoforms and US transcripts (CS rates > PS

rates > US rates, Figure 3B, Figure S4A-E). At single-molecule level, we observed that most CS transcripts

had three methylated sites in cluster 1, and two in cluster 2. By contrast, most US transcripts had two

methylated sites in cluster 1, and one in cluster 2. In addition, only very small proportions of CS transcripts had

no methylated site in either cluster (0.3%). This proportion increased only slightly, from 3.7% to 5.4%, in US

transcripts. These observations are consistent with a previous study (26) of three methylation sites (A8532,

A9428, A9442) reporting that 2.5% and 8.5% of CS and US transcripts, respectively, were unmethylated.

Overall, these results are consistent with previous observations that reducing m^{6}A methylation inhibits splicing

and stimulates genomic RNA packaging (17, 19, 26). They also strongly suggest that the methylation of these

sites plays an important role in splicing regulation and, ultimately, in virus infectivity.

Four additional sites were detected specifically in several isoforms (Figure 4A, Tables 2, S7). The methylation

rates of these additional sites were higher for CS than for PS isoforms, but the ratio of CS transcript

methylation to PS transcript methylation was much higher than for the 14 sites common to all isoforms (Figure

4B and Figure 3B). This finding provides additional support for a role for methylation in splicing regulation. It

also suggests that m^6A sites located at different positions within a specific isoform may be involved in different

processes, such as translation or the maintenance of stability. Further experiments involving the mutation of

one or several of these sites would help to address these questions.

In addition to the splicing isoforms, we detected ≈736 nt-long RNAs resulting from the transcription of 2-LTR

unintegrated DNA circles joining the R segments of the two LTRs (RU5-U3R, Figure 6A). These transcripts

contained six m^6A sites, five of which were common to all spliced isoforms and had higher methylation rates

than any of the other transcripts. An additional site, 2L125, was identified in the U5 segment, with a

methylation rate ranging from 24% to 35%. We observed that (i) positions homologous to this position (2L125)

in the other transcripts were not considered methylated by our detection process and (ii) the other five sites

had methylation rates higher than that at the homologous sites in all the other transcripts. These results

strongly suggest that, despite not encoding canonical viral proteins, 2-LTR transcripts play a functional role in

the viral life cycle. Consistent with this idea, 2-LTR circles were initially thought to be dead-end products in the

viral life cycle (42), but many studies subsequently showed that they could be transcribed and might constitute

a reserve supply of HIV-1 genomes for proviral integration (43-48). In addition, the large difference in 2L125

methylation rate between 2-LTR transcripts and the other transcripts (same sequence context) confirms that

this site is not an artifact of the detection process but, indeed, a *bona fide* methylation site.

It is generally though that methylation occurs cotranscriptionally and that exclusion of m^6A from splice-site

proximal regions by the exon junction complex results in low methylation rates close to splice junctions (49,

50). This hypothesis is not consistent with our observation that CS transcripts have higher methylation levels

than PS and US transcripts. It has recently been shown that m^{6}A methylation can occur posttranscriptionally, especially in transcripts that are highly methylated (51). This raises the possibility that the methylated sites observed at highly specific positions within HIV-1 transcripts are methylated posttranscriptionally and are potentially associated with an epitranscriptomic role in the virus life cycle.

In conclusion, we provide an unbiased assessment of m^{6}A modifications in HIV-1 transcripts isolated from infected CD4$^+$ T cells. We reveal the existence of novel m^{6}A sites in the 3' end common to all transcripts, in specific isoforms, and in 2-LTR transcripts. This study also highlights the complex interplay between m^{6}A modifications, splicing events and the fate of HIV-1 transcripts. These results pave the way for efforts to decipher the role of m^{6}A modifications in the HIV life cycle.

MATERIALS AND METHODS

Cell infection

VSV-G pseudotyped HIV$_{NL4-3\ XCS}$ was produced by transfection of 293T cells using CAPHOS kit (Sigma-Aldrich) as in (52). Viruses were pseudotyped to favor viral entry and to achieve a high infection rate within a short time. SupT1 CD4$^+$ T cells were cultured with RPMI GlutaMax 1640 (Gibco) complemented with 10% FBS (Dutscher) and 1% penicillin/streptomycin.

2 x 10^8 SupT1 CD4$^+$ T cells were infected with 800 ng of VSV-G-HIV$_{NL4-3\ XCS}$ HIV-Gagp24 for 3h in IMDM plus 10mM HEPES (Gibco/ThermoFisher Scientific) supplemented with 2 µg/ml of DEAE-dextran (Sigma). Two biological replicates were performed corresponding to 100 million cells infected at time 0. Eighteen hours post-infection a fraction of the cells (2 x 10^5) was harvested, stained with CD4-Vio770 antibody (Miltenyi) and a Live/Dead (LVD) fixable violet dye (Invitrogen), fixed with 4% paraformaldehyde (ChemCruz) and permeabilized (PBS1X, BSA 0,5%, saponin 0,05%). Cells were then stained with an HIV-1 Gag-p24 specific antibody (KC57-PE, Beckman Coulter). Sample acquisition and analysis were performed using a BD LSRFortessa and FlowJo v10.8 Software, respectively (both from BD Life Sciences). Mock-infected cells were used as negative control. Twenty and thirty hours post-infection, cells were harvested, washed in cold PBS and lysed at 4°C in RLT (from RNeasy Mini Kit (Qiagen) buffer containing ß-mercaptoethanol and 1% NP40.

PBMCs from anonymous blood donors (n=2, donor 864 and 880) were purchased from the blood bank at EFS (Établissement Français du sang) under the agreement number 15/EFS/022. PBMCs were isolated using a Ficoll gradient and monocytes depleted by overnight incubation in lying cell culture flasks containing

RPMI+10%FBS supplemented with 10 IU/ml of recombinant human IL-2 (rhIL-2, Miltenyi Biotec). CD4$^+$ T cells

were then isolated using CD4$^+$ magnetic beads (Miltenyi Biotec) and activated with Phytohemagglutinin (PHA,

1 µg/ml, PAA Laboratories) in RPMI GlutaMax 1640 complemented with 10% FBS, 1% penicillin/streptomycin,

nonessential amino acids and sodium pyruvate (all from Gibco/ThermoFisher Scientific), so called T-cell

medium. 48 h post-activation, CD4$^+$ T cells were harvested and resuspended in T-cell medium containing 100

IU/ml of recombinant human IL-2 (rhIL-2, Miltenyi Biotec). 7-days post activation, 300×10^6 and 130×10^6

CD4$^+$ T cells, were infected (for donor 864 and donor 880, respectively) at a multiplicity of infection (moi) 0,01

with VSV-G-HIV$_{NL4-3\ XCS}$ (representing around 1000 ng of HIVGagp24 / 10^7 cells) for 3h in IMDM plus 10mM

HEPES supplemented with 2 µg/ml of DEAE-dextran. CD4$^+$ T cells were then maintained in T-cell medium

containing 100 IU/ml of rhIL-2 and HIV infection was monitored using flow cytometry, as for SupT1 CD4$^+$ T

cells. For donor 864 and donor 880, cells were harvested and lysed 48 h and 72 h post-infection, respectively.

In vitro transcription

HIV-1 genomic RNA was transcribed in two parts, from the beginning of 5'LTR to the end of Pol (5124 bases)

and from the end of Pol to the end of 3'LTR (4829 bases), with 225 overlapping bases. Each part of the

genome was PCR amplified from pNL4.3XCS (53) and cloned in a pUC19 vector containing T7 promoter,

using NEBuilder cloning kit (New England Biolabs). The plasmids were then linearized with SmaI restriction

enzyme during 3 hours at 30°C, to perform run-off transcription. In vitro transcription was performed using

RiboMAX T7 Large Scale RNA Production System (Promega) with incubation 4 hours at 37°C , followed by

DNAseI treatment during 15 min and RNA column clean-up (ZymoResearch).

RNA purification

Total RNA was extracted from the cell pellets using RNeasy Mini Kit (Qiagen), according to the manufacturer's

protocol. Briefly, cells were lysed in RLT buffer containing b-mercaptoethanol and homogenized using

QIAshredder homogenizer. The lysate was then purified on RNeasy spin columns and RNA was eluted in

RNase-free water. PolyA RNA were then purified from the total RNA using two rounds on Oligo(dT)$_{25}$

Dynabeads (Thermo Fisher Scientific). In vitro transcripts were polyadenylated using E. coli PolyA polymerase

kit from NEB and purified on Oligo(dT)$_{25}$ Dynabeads.

Library preparation and sequencing

403 The polyA RNA was used for ONT direct RNA sequencing using the RNA004 kit. The reverse transcription

404 (RT) step was modified as described by (26); a pool of 111 oligonucleotides complementary to the HIV-1

405 genome was added during library preparation in an estimated 30-fold molar excess to the viral RNA, which

406 resulted in a much more efficient reverse transcription process.

407 Sequencing was processed on P2solo through the GridION (MinKNOW version: 24.02.10 for SupT1_20h_pi

408 rep1, 24.02.16 for SupT1_30h_pi rep1 and IVT, 24.06.15 for SupT1_20h_pi+demeth rep1 and rep2,

409 SupT1_20h_pi rep2, SupT1_30h_pi rep2, CD4_48h_pi and CD4_72h_pi, IVT). After sequencing, base-calling

410 has been done with dorado-0.7.2 and rna004_130bps_sup@v5.0.0 model with –modified-bases m^6A

411 parameter (for all samples).

413 **Identification of HIV-1 splicing isoforms**

414 Pass reads were mapped with minimap2 (v2.17-r941) (with -x splice and –secondary=no options) on the

415 PNL4-3XCS HIV-1 genome. BamToBed tools with -splice option have been used to select full-length

416 transcripts using a homemade script python. With this script, each read is analyzed and its splice junctions

417 must match (with a tolerance of one base) the splice junctions of a specific isoform. In each full-length read we

418 identified the donor (D) and acceptor (A) sites that match the already known HIV-1 splice sites (eight donor

419 and eleven acceptor sites, Nguyen Quang et al., Emery et al., Ocwieja et al., O'Reilly et al., Widera et al.)

420 (Table S2A). Some rare isoforms were detected with only few reads and only those detected in at least five of

421 the six samples were considered. Finally, 51 isoforms were identified by their succession of splice junctions

422 and were named as in (Nguyen Quang et al. 2020) (Table S2B). Between 4.4% (CD4_48h_pi) and 8.0%

423 (SupT1_20h_pi rep1) of full-length reads did not fully match these sites and were not considered.

425 **Detection of m^6A sites, computation of methylation rates**

426 Nucleotide position of m^6A's was obtained thanks to the specific base-calling process that can detect m^6A's

427 (version 5.1.0). For each sample, the methylation rate is computed by modkit pileup with –filter-threshold

428 A:0.8, --mod-thresholds a:0.9 and –edge-filter 20. The nucleotides that are called as m^6A are counted at each

429 position in a set of reads to determine the corresponding modification rate. The methylation rates computed at

430 each position of the transcripts in a sample were compared to those computed at the corresponding positions

431 of the IVT sample using the [DMR model and scoring details - Modkit](#) command that computes a final

432 modification rate and a corresponding p-value at each position. The positions presenting p-values $p \leq 10^{-5}$ and

a number of methylated reads ≥ 30 were retained as putative m[6]A sites. For the clustering analyses in Figure

3 and Figure S3, only the isoforms identified by more than 100 full-length reads were retained.

Single molecule analysis of methylated m[6]A sites

Modkit extract enables access to read-level base modification calls. Parameters –mapped-only and –include-

bed were used (bed file containing eight positions from cluster1 and three positions from cluster2). The output

tsv files were parsed with the R language. During this analysis, the threshold of probability of modification to

call 6mA was 0.9 (as with modkit pileup).

Synthetic oligonucleotides

We used two synthetic RNA oligonucleotides identical to HIV-1 regions with methylated As and the same two

oligonucleotides differing only at extremities, without methylated As.

Oli2-mod harbors m[6]A at positions 9400, 9428 et 9432:

CTTCAAGA(m[6]A)CTGCTGACATCGAGCTTGCTACAAGGG(m[6]A)CTTTCCGCTGGGG(m[6]A)CTTTCCAGG

(60 nt-long).

Oli2-nmod harbors non-modified A's and 6nt tag at 3' end (**TCATAG**):

CTTCAAGAACTGCTGACATCGAGCTTGCTACAAGGGACTTTCCGCTGGGGACTTTCCAGG**TCATAG** (66 nt-

long).

Oli1-mod harbors m[6]A at positions 8532 et 8563 + 5nt tag at 3' end (**TATGC**):

CCGCTTGAGAG(m[6]A)CTTACTCTTGATTGTAACGAGGATTGTGGA(m[6]A)CTTCTGGGAC**TATGC** (58 nt-

long).

Oli1-nmod harbors non-modified A's and 5nt tag at 3' end (**GTACG**):

CCGCTTGAGAGACTTACTCTTGATTGTAACGAGGATTGTGGAACTTCTGGGAC**GTACG** (58 nt-long).

Oligonucleotides demethylation and sequencing

As recommended by the manufacturer, less than 3 μM of m[6]A was incubated with 1 μM ALKBH5 at room

temperature. The composition of the reaction mixture was as follows: 3 μL enzyme (0.9 μg, 19.8 pmol), a mix

of four oligos at 0.13 μM each (oli1-mod, oli1-nmod, oli2-mod and oli2-nmod). A total of (3 x 0.13) + (2 x 0.13)

= 0.65 μM m[6]A was used in this step; the buffer contained 100 mM KCl, 2 mM MgCl2, 2 mM L-ascorbic acid,

300 μM alpha-ketoglutarate, 150 μM (NH4)2Fe((SO4)2 and 50 mM Hepes buffer (pH 6.5), in a total volume of

20 µL. The reactions were stopped after 1.5 h incubation by adding EDTA to a final concentration of 5 mM; the oligos were purified using SPRIselect beads following a modified protocol as follows. Water was added to a 25 µL total volume, followed by the addition of 45 µL SPRI beads and mixing. After 5 minutes of room temperature incubation, the beads were magnetized and the supernatant containing the small RNA was transferred to a new tube containing 45 µL SPRI beads and 135 µL isopropanol, and mixed. After another 5 minutes of room temperature incubation, the beads were magnetized and the supernatant was discarded. The beads were washed twice with 86% ethanol, allowed to dry for 5 minutes and resuspended in 10 µL water. The purified oligos were then used for ONT direct RNA004 library preparation and sequenced on two MinION flow cells (MinKNOW version: 24.06.14). Distinction between oligos has been made by mapping to the reference with these four oligos added by eight As in 3'.

ACKNOWLEDGMENTS

This work was granted by the Agence Nationale de Recherche sur le SIDA (ANRS-Maladies infectieuses émergentes) for fundings to A.M. and O.N., S.B salary was supported by ANRS. L.B PhD salary was supported by ANRS-MIE. This study was initiated thanks to fundings from the cross-disciplinary thematic axes of I2BC through the epiRNA programs. The HTS facility of the I2BC acknowledges the support of France Génomique (funded by the French National Program "Investissement d'Avenir" ANR-10-INBS-09).

A

HIV-1$_{NL4-3}$
(VSV-G pseudotyped for
the first cycle of infection)

flow cytometry

analysis

assessment
of cell viability &
infection rate

Viral mRNA

Cell lysis

Total RNA

Poly A purification

Nanopore sequencing

RNA sequence analysis
and mapping on HIV genome

AGTCCCTGAATCGA

SupT1 cell line
or
primary CD4$^+$ T cells

B

MOCK | HIV$_{NL4-3}$

94.8

0.45 | 22.2
0.01 | 73.3

20h pi

81.5

0.02 | 1.34
0 | 97.5

30h pi

SSC | Live dead

CD4-APC | Gag-PE

C

MOCK | HIV$_{NL4-3}$

0.03 | 61.6
0.05 | 11.8

Donor 864
48h pi

82.8

0.12 | 43.5
0.02 | 11.9

Donor 880
72h pi

SSC | Live dead

CD4-APC | Gag-PE

Figure S1. (A) Outline of the infection and Nanopore sequencing protocols. SupT1 T cells or primary

CD4$^+$ T cells were infected with HIV$_{NL4-3}$ pseudotyped with VSV-G to enhance viral infection. At 20 or 30 hours

post-infection (pi) for HIV $_{NL4-3}$-infected SupT1 cells and at 48 h or 72 h pi for HIV $_{NL4-3}$-infected primary CD4$^+$ T

cells, we collected a fraction of the cell culture. The viability and percentages of HIV-Gag$^+$ and CD4$^+$ cells were

assessed by flow cytometry. During HIV infection, cells start expressing HIV-Gag protein and progressively

lose their membrane expression of CD4 due to expression of the HIV accessory proteins Nef and Vpu, and

HIV-Env. The cells were harvested and polyA RNA was purified and sequenced with Nanopore technology.

Reads were mapped onto the HIV$_{NL4-3}$ genome sequence. Created with BioRender. Moris, A. (2025)

https://BioRender.com/o40u811 **(B) Analysis of the infection rate of HIV$_{NL4-3}$-infected CD4$^+$ SupT1 cells.**

After 20 h or 30 h, cells infected with HIV$_{NL4-3}$ or mock-treated were labeled with a viability dye and stained

with antibodies specific for CD4 and HIV-Gag. Left panels: the percentage viable cells. Middle and right

panels: percentages of CD4$^-$HIV-Gag$^+$ and CD4$^+$HIV-Gag$^+$ cells among mock-treated and HIV$_{NL4-3}$-infected

CD4$^+$ SupT1 cells, respectively. **(C) Analysis of the infection rate of HIV$_{NL4-3}$-infected primary CD4$^+$ T**

cells. After 48 h or 72 h, cells from donors 864 and 880 were harvested and stained as in (B). Left panels: the

percentage viable cells. Middle and right panels: percentages of CD4$^-$HIV-Gag$^+$ and CD4$^+$HIV-Gag$^+$ cells

among mock-treated and HIV$_{NL4-3}$-infected primary CD4$^+$ T cells, respectively, from donors 864 (top panels)

and 880 (bottom panels). Numbers indicated the percentage of cells within the quadrant. MOCK: cells mock-

treated without infection used as a negative control.

A

504
505
506
507   B

C

D

528 E

529
530
531 F

532
533
534 **Figure S2.** m[6]A sites detected in splicing isoforms. **(A)** SupT1_20h_pi rep 1 sample; **(B)** SupT1_30h_pi rep1

535 sample; **(C)** SupT1_20h_pi rep2 sample; **(D)** SupT1_30h_pi rep2 sample; **(E)** CD4_48h_pi sample; **(F)**

536 CD4_72h_pi sample. Only positions with p-values $\leq 1 \times 10^{-5}$ and a number of modified reads ≥ 30 are

537 displayed.

538 A - SupT1_30h_pi rep1

539
540
541 B - SupT1_20h_pi rep2   C - SupT1_30h_pi rep2

542
543
544 D - CD4_48h_pi   E - CD4_72h_pi

545
546

547 **Figure S3.** Heatmap of the methylation levels of the 14 m^{6}A sites detected in the various splicing isoforms. **(A-**

548 **E)**, results for the samples indicated. Methylation level was calculated for each set of full-length reads

549 corresponding to each isoform represented by ≥100 reads (Materials and Methods). Isoforms are clustered

550 based on the methylation level of the 14 m^{6}A sites and the clusters obtained correspond essentially to US, PS

551 and CS isoforms.

Figure S4. Methylation levels of m⁶A sites in CS, PS and US transcripts. (**A-E**), results for the samples indicated.

SupT1_20h_pi rep2

A

SupT1_30h_pi rep1

C

SupT1_30h_pi rep2

E

586
587
588 **CD4_48h_pi**
589
590 G
591

H

592
593
594
595 **CD4_72h_pi**
596
597 I

J

598
599 **Figure S5.** Methylation rates of the four isoform-specific m^{6}A sites in the samples indicated. The A5613,
600 A5654 and A5703 sites are common to the Vif and Vpr isoforms; A5887 is common to the Vif, Vpr, Tat and
601 SORF1 isoforms. **(A,C,E,G,I)** Methylation rates calculated for the four sites in CS (- intron 4, Vif1, Vpr1, Vpr2),
602 PS (+ intron 4, Vif2, Vpr3, Vpr4) and US isoforms. **(B,D,F,H,J)** Methylation rates calculated for the A5887 site
603 for the CS (Tat1, 2, 3, 4, SORF1), PS (Tat5) and US isoforms.
604

606
607
608
609

610
611
612 **Figure S6**. Single-molecule analysis of methylated m[6]A sites. **(A)** Cluster 1 and **(B)** cluster 2 of the indicated

613 SupT1 samples. **(C)** Cluster 1 and **(D)** cluster 2 of the indicated CD4 samples.

SupT1_30h_pi rep1

A (US)   B (PS)   C (CS)

SupT1_20h_pi rep2

D (US)   E (PS)   F (CS)

SupT1_30h_pi rep2

G (US)   H (PS)   I (CS)

636 **CD4_48h_pi**
637
638 J (US)   K (PS)   L (CS)

639
640
641 **CD4_72h_pi**
642
643 M (US)   N (PS)   O (CS)

644
645
Figure S7. Single-molecule analysis of methylated m^{6}A sites illustrating the correlation between the

methylation patterns for clusters 1 and 2 of the same transcripts. The intensity of the color on the heatmap

indicates the number of reads, with the indicated numbers of methylated As in cluster 1 (0 to 8 m^{6}As) and

cluster 2 (0 to 3 m^{6}As). US, unspliced; PS partially spliced; CS, completely spliced. One m^{6}A of one read is

considered methylated if the probability of base modification is ≥ 0.5 (value in modkit extract), Materials and

Methods).

A

B

C

Figure S8. Secondary structures of HIV-1 RNA around the 14 m[6]A sites detected in all isoforms. **(A)** Region of the Rev/Env sequence containing the eight m[6]A sites of cluster 1; this structure is common to the US and PS transcripts as both retain the end of the unspliced intron 4 at the A7 acceptor site, in green (intron 3' end). In red, the two highly methylated sites (64-80%). **(B)** Region in the Nef sequence containing the methylated sites A9024 and A9074. **(C)** Region in the 3'LTR containing the three m[6]A sites of cluster 2. In red, the highly methylated site (79-84%). The structure predictions are adapted from (54).

SUPPLEMENTARY TABLES

	SupT1_20h _pi rep1	SupT1_20h _pi rep2	SupT1_30h_ pi rep1	SupT1_30h _pi rep2	CD4_48h _pi	CD4_72h _pi
Total reads	7317460	27288259	15758351	27289373	26172498	28942428
HIV-1 reads	386852	287479	1074370	345351	166733	393666
Infection rate (%)	5.5	1.31	7.1	1.53	0.77	1.9
Full-length reads	80230	134086	164360	127964	33587	39278
Full length (%)	20.74	46.64	15.3	37.05	20.14	9.98

Table S1. Run statistics. Infection rate, number of HIV reads divided by the total of HIV and human reads.

A

Donor sites	Acceptor sites
D1: 743	
D1c: 747	
	A1: 4913
	A1b: 4919
D2: 4962	
D2b: 5058	
	A2: 5390
D3 : 5463	
	A3: 5777
	A4c: 5936
	A4a: 5954
	A4b: 5960
	A5: 5976
	A5a: 5980
	A5c: 5992
D4: 6044	
D4a: 6048	
D4b: 6051	
	A7: 8369

B

		SupT1_20h_pi rep1	%	SupT1_20h_pi rep2	%	SupT1_30h_pi rep1	%	SupT1_30h_pi rep2	%	CD4_48h_pi	%	CD4_72h_pi	%
US		610	0.83	10494	8.21	2387	1.57	16566	13.75	2286	7.12	3280	8.76
envvpu1	D1A5	3971	5.38	17206	13.47	12515	8.22	16904	14.04	3699	11.52	4054	10.83
envvpu2	D1A4b	214	0.29	757	0.59	659	0.43	901	0.75	216	0.67	151	0.4
envvpu3	D1A4a	139	0.19	462	0.36	370	0.24	451	0.37	269	0.84	348	0.93
envvpu5	D1A1D2A5	281	0.38	756	0.59	672	0.44	943	0.78	111	0.35	93	0.25
envvpu9	D1A2D3A5	619	0.84	1345	1.05	1304	0.86	2367	1.97	397	1.24	195	0.52
envvpu10	D1A2D3A4b	46	0.06	61	0.05	54	0.04	143	0.12	30	0.09	9	0.02
envvpu13	D1A1D2A2D3A5	76	0.1	147	0.12	140	0.09	250	0.21	38	0.12	21	0.06
envvpu17	D1A5a	92	0.12	488	0.38	350	0.23	466	0.39	91	0.28	100	0.27
envvpu18	D1cA4b	2	0	6	0	3	0	9	0.01	3	0.01	2	0.01
envvpu19	D1cA5	49	0.07	127	0.1	120	0.08	140	0.12	30	0.09	40	0.11
nef1	D1A7	286	0.39	463	0.36	663	0.44	492	0.41	175	0.55	228	0.61
nef2	D1A5D4A7	29546	40.01	49676	38.88	65277	42.86	34533	28.67	8482	26.42	13658	36.47

nef3	D1A1D2A5D4A7	6934	9.39	7760	6.07	13894	9.12	7378	6.13	1066	3.32	1214	3.24
nef4	D1A2D3A5D4A7	7816	10.59	6156	4.82	11672	7.66	7242	6.01	1591	4.95	931	2.49
nef5	D1A1D2A2D3A5D4A7	2290	3.1	1512	1.18	3781	2.48	1897	1.58	376	1.17	162	0.43
nef9	D1A5aD4A7	690	0.93	1382	1.08	1540	1.01	857	0.71	205	0.64	357	0.95
nef12	D1cA5D4A7	214	0.29	331	0.26	496	0.33	268	0.22	65	0.2	105	0.28
nef13	D1A5D4aA7	25	0.03	35	0.03	58	0.04	27	0.02	8	0.02	10	0.03
nef14	D1A2D3A5D4aA7	8	0.01	8	0.01	11	0.01	8	0.01	2	0.01	0	0
nef16	D1cA5D4A7	10	0.01	10	0.01	23	0.02	6	0	1	0	3	0.01
nef17	D1A5D4bA7	8	0.01	9	0.01	22	0.01	9	0.01	2	0.01	2	0.01
rev1	D1A4bD4A7	3290	4.46	5743	4.5	7876	5.17	4360	3.62	1645	5.12	1350	3.61
rev2	D1A4aD4A7	3748	5.08	6522	5.11	8662	5.69	4546	3.77	3284	10.23	3979	10.63
rev3	D1A4cD4A7	213	0.29	348	0.27	539	0.35	252	0.21	396	1.23	584	1.56
rev4	D1A1D2A4bD4A7	363	0.49	504	0.39	830	0.55	445	0.37	169	0.53	110	0.29
rev5	D1A1D2A4aD4A7	199	0.27	326	0.26	464	0.3	218	0.18	137	0.43	111	0.3
rev7	D1A2D3A4bD4A7	786	1.06	641	0.5	1111	0.73	742	0.62	387	1.21	90	0.24
rev8	D1A2D3A4aD4A7	717	0.97	524	0.41	1006	0.66	626	0.52	507	1.58	190	0.51
rev9	D1A2D3A4cD4A7	33	0.04	38	0.03	52	0.03	35	0.03	51	0.16	34	0.09
rev10	D1A1D2A2D3A4bD4A7	191	0.26	154	0.12	315	0.21	169	0.14	81	0.25	23	0.06
rev11	D1A1D2A2D3A4aD4A7	202	0.27	161	0.13	335	0.22	151	0.13	130	0.4	42	0.11
rev13	D1cA4bD4A7	23	0.03	29	0.02	38	0.02	29	0.02	8	0.02	11	0.03
SORF1	D1A1D2bA3D4A7	218	0.3	145	0.11	358	0.24	155	0.13	66	0.21	84	0.22
SORF3	D1A1D2bA2D3A3D4A7	7	0.01	10	0.01	35	0.02	13	0.01	6	0.02	4	0.01
tat1	D1A3D4A7	4534	6.14	3746	2.93	4210	2.76	2101	1.74	1584	4.93	2379	6.35
tat2	D1A1D2A3D4A7	2354	3.19	2181	1.71	3641	2.39	1914	1.59	745	2.32	1019	2.72
tat3	D1A2D3A3D4A7	484	0.66	262	0.21	568	0.37	332	0.28	171	0.53	97	0.26
tat4	D1A1D2A2D3A3D4A7	171	0.23	109	0.09	294	0.19	120	0.1	73	0.23	53	0.14
tat5	D1A3	230	0.31	278	0.22	251	0.16	270	0.22	250	0.78	281	0.75
tat6	D1A1D2A3	16	0.02	31	0.02	39	0.03	32	0.03	34	0.11	39	0.1
tat7	D1A2D3A3	7	0.01	15	0.01	10	0.01	13	0.01	12	0.04	7	0.02
tat9	D1cA3D4A7	11	0.01	8	0.01	6	0	10	0.01	9	0.03	6	0.02
vif1	D1A1D4A7	98	0.13	267	0.21	389	0.26	280	0.23	39	0.12	112	0.3
vif2	D1A1	677	0.92	3199	2.5	2108	1.38	5234	4.35	547	1.7	591	1.58
vif3	D1cA1b	2	0	4	0	10	0.01	7	0.01	2	0.01	3	0.01
vif4	D1cA1	0	0	11	0.01	9	0.01	22	0.02	2	0.01	2	0.01

vif5	D1A1b	3	0	10	0.01	15	0.01	19	0.02	0	0	3	0.01
vpr1	D1A2D4A7	332	0.45	290	0.23	435	0.29	426	0.35	139	0.43	186	0.5
vpr2	D1A1D2A2D4A7	61	0.08	50	0.04	148	0.1	76	0.06	21	0.07	16	0.04
vpr3	D1A2	880	1.19	2793	2.19	2386	1.57	5651	4.69	2340	7.29	1022	2.73
vpr4	D1A1D2A2	64	0.09	166	0.13	142	0.09	335	0.28	131	0.41	54	0.14
Total isoforms		73840		127756		152293		120440		32109		37445	
2LTR*		1273	1.69	566	0.44	2391	1.55	878	0.72	52	0.16	109	0.29

679
680
681 **Table S2. (A)** List of the HIV-1 donor and acceptor splice sites. **(B)** Absolute and relative amounts of the HIV-1
682 transcript isoforms. %, amount calculated relative to all identified splicing isoforms (not including 2LTR); *,
683 amount calculated relative to all HIV-1 isoforms including 2LTR. Gray lines, the seven most abundant
684 isoforms, accounting for about 80% of all isoforms.
685

	SupT1_20h_pi rep2	SupT1_30h_pi rep1	SupT1_30h_pi rep2	CD4_48h_pi	CD4_72h_pi
SupT1_20h_pi rep1	0.946	0.991	0.861	0.890	0.925
SupT1_20h_pi rep2		0.972	0.964	0.952	0.978
SupT1_30h_pi rep1			0.891	0.911	0.946
SupT1_30h_pi rep2				0.941	0.933
CD4_48h_pi					0.971

Table S3. Pearson correlation coefficients for the relationship between the relative amounts of the splicing isoforms (Table S2B).

m^{6}A position	Sequence	SupT1-20h-pi rep1	SupT1-20h-pi rep2	SupT1-30h-pi rep1	SupT1-30h-pi rep2	CD4-48h-pi	CD4-72h-pi
A8436	AGACA	9.7	6.3	8	5.7	4.7	3.6
A8442	AGACA	10.3	7.0	9.7	6.7	6.3	5.3
A8487	GGAC<u>G</u>	30	22.7	27	22	19.7	18
A8532	AGACT	79.3	72.7	79.3	72	68.3	63.7
A8555	GGA<u>T</u>T	19.3	13	17.3	13	11.7	9.3
A8563	GAACT	79.3	74	79.7	72.7	70.3	66.3
A8572	GGAC<u>G</u>	23.7	27	23.3	26	25.7	18.7
A8632	GAACT	10.3	6.7	9.3	6.7	5.7	3.7
A9024	TGACT	7.3	5.3	6.0	5.0	4.3	3.6
A9074	GGACT	8.7	7.7	9.7	6.3	5.7	5.7
A9163	GAACT	17.3	13	15	11.7	10.3	9.7
A9400	GAACT	29.7	28.0	32.0	25.0	21.0	18.7
A9428	GGACT	84.3	82.7	84.3	80.7	79.7	79.3
A9442	GGACT	53.3	49	52.3	46	41	41.3

Table S4. Rates of methylation of the 14 m^{6}A sites detected in all HIV-1 transcripts in each of the 6 samples. Underlined, 1-nt variants of the consensus DRACH sequences. Gray, the three most methylated sites, A8532, A8563, A9428.

Position	Sequence	SupT1_20h_pi rep1	SupT1_20h_pi+demeth rep1		SupT1_20h_pi rep2	SupT1_20h_pi+demeth rep2	
		% methyl.	% methyl.	p-value	% methyl.	% methyl.	*p*-value
A8436	AGACA	9.7	1	2.00E+101	6.3	2	9.60E-38
A8442	AGACA	10.3	1.7	7.90E-147	7	2.7	2.40E-36
A8487	GGAC<u>G</u>	30	7.3	1.60E-265	22.7	11.7	3.90E-75
A8532	AGACT	79.3	49.3	0	72.7	51.7	6.70E-191
A8555	GGA<u>TT</u>	19.3	11.3	4.00E-56	13	9	2.50E-14
A8563	GAACT	79.3	60	5.30E-174	74	60.7	8.60E-82
A8572	GGAC<u>G</u>	23.7	12.3	1.80E-24	27	17.3	1.70E-15
A8632	GAACT	10.3	3	1.10E-80	6.7	2.3	3.30E-36
A9024	TGACT	7.3	1.3	3.00E-87	5.3	1.3	1.00E-43
A9074	GGACT	8.7	5	1.10E-22	7.7	5.3	3.20E-09
A9163	GAACT	17.3	3.3	1.00E-165	13	5.3	4.30E-58
A9400	GAACT	29.7	18.3	3.20E-93	28	21.7	6.90E-24
A9428	GGACT	84.3	22.3	0	82.7	25.3	0
A9442	GGACT	53.3	18.3	0	49	22.3	1.5e-311

724
725
Table S5. Demethylation efficiency for the 14 m^{6}A sites detected in all transcripts of SupT1_20h_pi rep1 and SupT1_20h_pi rep2. SupT1_20h_pi+demeth is the SupT1_20h_pi sample demethylated with ALKBH5 (Materials and Methods). *P*-values are given for the difference between the numbers of methylated reads in SupT1_20h_pi and in SupT1_20h_pi+demeth (Materials and Methods).

A - SupT1_20h_pi rep1

	8436	8442	8487	8532	8555	8563	8572	8632	9024	9074	9163	9400	9428	9442
envvpu1	9.00	11.00	31.00	85.33	18.00	84.67	24.33	8.67	7.67	9.00	18.00	32.33	88.67	59.00
envvpu2	8.12	13.37	33.63	86.14	18.83	84.77	35.48	12.00	10.58	9.33	21.21	31.54	87.42	59.77
envvpu3	11.45	14.48	37.81	89.22	18.87	88.22	22.58	11.31	10.83	12.97	18.33	33.51	91.33	52.72
envvpu5	14.94	11.89	43.11	93.81	24.92	92.54	28.20	12.85	7.77	10.32	26.73	42.67	90.46	65.72
envvpu9	14.33	12.00	36.67	89.00	26.67	88.33	30.82	11.33	9.00	9.67	27.67	33.67	89.67	62.67
nef1	14.18	12.26	39.08	92.82	28.73	92.46	36.00	13.40	10.44	9.64	22.14	36.62	90.65	68.79
nef12	13.61	15.46	42.78	88.73	27.75	93.39	41.97	11.77	7.23	12.10	26.31	38.24	89.33	61.10
nef2	14.00	14.00	41.33	92.67	27.33	92.33	33.67	14.33	9.33	11.00	22.00	38.00	91.00	63.00
nef3	14.33	16.00	44.67	93.67	30.33	93.00	36.00	14.33	11.00	11.33	25.67	41.33	90.33	66.33
nef4	16.00	15.67	45.67	93.67	31.67	93.33	37.33	16.33	12.67	10.67	27.67	40.00	91.00	66.33
nef5	18.33	17.00	49.67	94.67	35.67	94.00	42.33	19.33	14.67	11.67	29.00	42.00	92.00	69.33
nef9	13.00	15.00	41.67	92.00	25.00	92.67	37.44	13.67	11.33	11.33	23.00	36.33	90.33	66.33
rev1	12.67	14.00	37.00	92.00	24.33	90.67	32.00	13.00	9.33	9.00	22.67	37.67	91.00	62.00
rev10	10.75	18.48	41.39	92.68	31.41	90.54	33.72	15.48	13.17	16.88	27.25	45.02	91.37	65.40
rev11	15.10	15.91	41.05	95.12	31.18	94.15	52.88	14.10	16.66	15.40	26.08	44.85	91.55	71.86
rev2	12.67	12.67	37.33	90.33	24.33	90.67	31.67	12.00	9.67	10.67	21.33	36.67	90.00	60.67
rev3	10.95	16.66	30.68	88.50	24.78	90.60	33.52	8.55	6.79	11.39	19.84	35.14	88.76	61.13
rev4	13.67	12.33	45.00	90.67	30.00	94.00	29.37	12.67	13.67	12.67	25.33	39.00	90.00	60.67
rev5	11.64	13.28	33.21	94.26	28.68	93.90	33.63	16.46	9.68	9.06	23.33	37.76	90.90	61.83
rev7	14.67	16.00	41.00	93.00	29.67	93.33	42.52	18.00	13.00	12.33	30.33	38.33	91.00	65.00
rev8	18.33	17.33	42.33	93.67	32.67	93.67	34.26	13.33	13.00	11.00	28.67	39.67	92.00	64.67
SORF1	15.61	17.37	45.83	95.40	27.48	92.54	21.55	17.93	13.46	14.60	31.49	42.18	89.19	65.47
tat1	13.33	13.33	38.67	90.67	22.00	91.33	34.00	13.67	10.00	12.33	23.67	31.33	91.00	62.33
tat2	18.00	18.33	47.67	93.67	32.33	93.33	35.67	16.00	13.33	11.33	28.67	39.67	91.00	67.67
tat3	19.00	17.67	49.00	94.33	32.33	93.33	43.70	18.00	12.67	11.00	27.67	41.33	92.00	68.67
tat4	25.31	16.43	51.76	91.88	39.37	96.89	51.67	21.15	23.18	12.30	34.00	40.95	93.42	69.13

tat5	10.23	10.98	27.91	82.54	22.81	85.05	21.41	9.95	9.60	13.55	26.76	26.27	89.54	61.77
us	4.33	5.67	19.00	68.67	10.33	70.00	12.99	5.67	5.33	5.00	11.33	17.33	81.33	44.33
vif2	11.33	10.33	30.67	83.67	18.00	82.00	21.81	10.33	10.00	8.00	18.00	33.00	85.33	59.67
vpr1	12.00	20.67	47.00	92.33	30.00	92.67	37.22	18.33	14.00	16.33	30.73	35.13	91.29	67.36
vpr3	9.33	8.33	29.33	80.33	16.33	80.00	19.42	10.33	6.33	9.00	17.33	27.00	88.67	53.00

731
732
733 **B - SupT1_30h_pi rep1**

	8436	8442	8487	8532	8555	8563	8572	8632	9024	9074	9163	9400	9428	9442
envvpu1	7.00	8.67	27.00	82.33	16.00	82.67	23.00	8.33	6.67	9.33	15.00	32.00	88.00	55.67
envvpu13	13.71	10.78	44.99	89.45	36.43	91.19	41.19	15.18	14.74	15.49	25.68	49.71	92.21	75.56
envvpu17	7.33	10.00	27.67	81.67	13.33	82.00	23.12	7.00	7.00	10.67	15.00	30.67	86.67	54.33
envvpu19	12.84	12.06	29.28	87.28	18.09	86.83	19.76	8.01	3.64	5.53	8.43	33.11	91.81	53.41
envvpu2	8.33	8.67	28.00	84.33	15.67	81.00	23.06	11.33	5.33	8.33	17.00	34.67	86.67	51.33
envvpu3	7.33	9.67	28.33	81.00	19.67	84.00	22.68	9.33	7.00	8.67	17.33	36.67	88.00	54.00
envvpu5	9.67	10.33	32.00	90.33	23.00	88.67	31.31	11.67	9.67	14.67	19.00	40.00	91.67	65.67
envvpu9	8.67	9.33	31.00	86.00	18.00	86.33	27.33	10.00	7.00	10.67	20.00	38.00	87.67	56.67
nef1	12.00	17.00	40.33	93.00	24.33	90.67	35.21	12.00	9.67	12.00	19.67	45.00	89.00	62.33
nef12	12.67	11.67	35.67	91.00	25.00	89.00	42.24	13.33	6.33	11.00	19.00	39.00	87.67	59.00
nef2	11.00	13.33	37.00	92.00	24.67	91.00	31.67	12.67	7.67	11.67	19.00	39.67	90.00	62.00
nef3	11.00	12.67	37.00	91.00	26.33	92.33	34.00	13.67	9.00	12.67	21.33	43.00	88.67	64.33
nef4	11.00	12.00	36.67	91.33	25.00	91.33	31.33	13.33	9.33	12.33	21.00	41.00	89.67	63.33
nef5	12.00	13.00	40.33	91.33	29.33	92.67	35.00	16.33	10.67	14.33	24.33	45.67	89.00	67.67
nef9	11.67	12.33	38.00	91.67	23.33	91.33	30.67	10.67	8.33	11.00	20.00	38.33	89.67	63.33
rev1	10.00	11.67	32.67	90.00	22.00	89.67	31.33	12.33	7.67	12.00	19.00	38.67	89.67	61.00
rev10	11.60	16.33	37.67	90.33	26.00	93.67	35.32	18.87	14.94	18.45	25.64	46.29	89.19	63.75
rev11	11.00	8.67	35.00	89.67	30.33	92.33	31.67	15.00	10.00	13.67	27.63	45.33	90.67	68.48
rev2	10.00	11.00	32.00	89.67	22.33	90.00	32.67	11.67	7.67	10.67	19.00	39.33	89.33	60.33
rev3	10.33	12.33	29.67	91.67	21.00	89.67	31.43	10.00	8.67	11.33	18.33	37.00	88.67	63.00

rev4	8.67	11.67	33.33	90.67	25.00	93.00	33.20	12.33	6.33	13.33	20.33	44.00	88.33	65.33
rev5	9.00	11.00	29.67	90.33	23.67	91.67	42.43	13.33	8.00	12.67	22.33	43.67	87.33	62.67
rev7	9.00	11.33	35.67	90.00	26.33	91.67	32.33	15.67	9.67	13.67	21.67	40.33	90.00	62.33
rev8	10.00	11.67	35.00	91.00	24.67	91.67	32.08	14.00	9.33	14.67	22.67	42.67	91.00	63.33
SORF1	12.00	14.00	33.33	90.33	26.33	92.00	29.76	13.67	11.00	16.67	23.00	37.33	89.33	64.00
tat1	15.67	17.67	41.00	91.67	25.33	91.67	32.67	15.00	9.67	14.67	23.00	40.33	91.33	63.67
tat2	15.00	17.67	40.67	91.67	28.33	92.33	33.67	16.00	10.67	15.00	23.33	43.00	90.00	65.67
tat3	16.67	18.00	42.67	91.67	30.00	91.67	34.77	15.67	11.00	15.33	24.33	40.00	89.67	63.67
tat4	18.01	17.09	50.58	93.05	33.07	90.31	39.89	20.77	15.18	22.33	31.67	47.94	90.13	74.81
tat5	5.98	9.13	30.40	83.34	17.25	83.01	29.00	17.10	10.15	11.25	20.41	37.17	87.73	53.42
us	4.33	5.00	16.00	66.33	8.67	66.67	13.00	6.00	3.67	5.33	9.67	21.00	77.33	43.33
vif1	20.00	19.00	51.33	95.33	40.33	95.33	47.83	21.67	15.67	16.00	28.67	44.00	88.67	71.33
vif2	8.33	8.00	27.33	80.67	17.00	81.67	25.33	10.33	7.67	9.67	19.00	31.00	88.00	53.67
vpr1	20.33	17.67	47.00	92.33	34.00	91.67	31.09	21.33	14.33	13.00	27.00	43.33	93.67	68.33
vpr2	24.48	17.03	47.23	93.24	34.02	92.19	47.05	19.74	18.19	21.23	36.13	49.17	91.18	67.63
vpr3	6.67	6.67	24.00	77.33	14.33	77.33	23.00	9.00	6.33	8.33	14.33	30.00	85.67	52.33
vpr4	10.45	10.04	33.81	90.38	22.91	85.46	29.29	12.18	9.84	15.12	22.33	35.52	88.84	61.83

734
735
736 **C - SupT1_20h_pi rep2**

	8436	8442	8487	8532	8555	8563	8572	8632	9024	9074	9163	9400	9428	9442
envvpu1	7.00	9.33	27.67	83.33	15.67	82.33	32.67	7.33	7.00	10.00	16.33	33.67	89.67	56.67
envvpu13	5.51	11.81	40.84	83.11	23.31	93.33	28.77	12.52	11.52	12.71	23.43	40.38	90.21	67.64
envvpu17	8.33	8.00	30.33	82.33	15.33	84.00	34.27	5.33	8.67	11.33	17.33	35.00	90.67	54.00
envvpu19	7.36	15.50	19.51	85.73	17.85	83.70	21.41	3.94	5.55	9.42	11.26	34.93	91.77	56.19
envvpu2	7.67	9.00	26.67	81.33	16.00	81.00	27.90	8.00	6.67	7.67	16.00	33.67	87.33	59.33
envvpu3	4.67	8.33	26.67	81.67	14.33	80.33	32.57	6.33	6.67	10.67	18.00	31.33	88.67	56.33
envvpu5	10.00	13.67	33.67	92.00	21.33	89.67	45.98	9.33	9.33	15.33	24.67	43.33	92.33	65.33
envvpu9	8.00	9.67	29.33	87.33	18.00	86.00	35.67	9.00	8.00	11.67	18.67	36.00	90.67	61.00

nef1	17.33	20.67	50.67	93.00	24.33	90.33	40.24	11.67	12.00	17.33	25.33	42.00	93.33	64.33
nef12	11.57	15.67	39.67	90.33	26.67	90.33	49.08	6.67	8.33	8.67	16.73	44.47	88.07	56.91
nef2	11.00	13.67	38.33	92.00	21.00	90.67	42.67	11.00	7.67	11.00	18.33	39.67	91.67	61.67
nef3	9.67	12.33	39.00	92.00	23.00	91.33	43.67	11.33	8.67	11.67	19.00	42.00	90.67	62.67
nef4	10.00	11.33	37.33	91.33	22.67	91.00	42.67	10.67	8.00	10.67	17.67	40.67	90.67	62.00
nef5	10.67	12.00	41.33	92.33	24.33	91.67	46.33	12.33	8.67	11.00	20.67	43.67	90.33	64.33
nef9	12.00	16.00	39.00	92.00	20.67	90.33	43.33	10.33	6.67	11.33	18.67	40.00	91.67	63.33
rev1	9.67	11.33	34.67	91.00	20.67	89.33	45.67	10.33	7.67	11.00	19.67	39.67	92.33	61.67
rev10	7.36	14.90	36.50	92.38	16.01	87.69	54.58	15.88	11.23	10.62	26.01	46.63	90.64	65.18
rev11	8.94	12.32	34.53	90.17	20.57	91.44	40.44	12.65	8.76	16.33	16.61	37.49	87.40	66.10
rev2	9.00	11.67	33.33	91.00	20.33	90.00	46.00	11.00	8.00	11.00	18.67	38.00	91.67	61.67
rev3	8.67	11.33	29.33	88.67	20.00	85.67	42.78	8.67	9.67	11.33	16.67	37.33	90.33	66.33
rev4	9.67	11.33	35.33	91.67	23.00	90.67	33.33	12.67	9.00	14.00	18.33	43.33	89.33	64.33
rev5	12.33	11.67	41.33	90.33	19.33	91.67	49.08	11.33	10.00	14.00	20.99	41.33	92.00	67.01
rev7	7.33	10.33	31.00	90.33	20.00	93.00	41.47	10.67	9.33	13.00	19.67	45.67	92.00	65.67
rev8	8.00	10.00	33.67	90.67	22.33	88.67	47.60	11.00	6.33	9.33	18.00	37.33	90.00	59.67
SORF1	12.90	15.75	39.56	89.18	19.86	89.88	55.00	10.23	10.16	16.21	18.23	39.04	92.70	70.88
tat1	14.33	17.00	40.67	93.00	21.33	91.33	42.33	11.67	9.00	12.67	22.00	40.67	92.67	65.00
tat2	13.00	16.67	43.67	93.67	22.67	91.33	45.67	11.33	9.67	14.33	23.67	42.33	92.00	66.33
tat3	17.22	17.31	38.92	92.27	24.60	92.52	40.71	9.86	8.90	12.84	24.96	45.84	92.98	68.09
tat4	10.88	21.89	40.90	92.19	28.41	92.94	45.39	15.05	13.82	11.10	24.26	45.05	90.82	62.67
tat5	9.10	9.26	33.33	76.50	18.85	85.86	36.79	8.72	6.38	10.99	20.31	37.02	90.14	61.92
us	4.00	5.33	18.33	66.33	10.33	68.33	20.33	5.33	4.33	6.00	10.00	23.00	79.33	43.00
vif1	14.69	12.97	43.05	88.56	24.58	89.12	65.00	12.22	12.64	7.87	23.27	39.49	89.69	64.71
vif2	7.00	8.67	26.67	79.67	15.00	79.00	29.33	7.00	6.33	9.33	15.67	31.67	87.67	56.67
vpr1	13.90	14.56	44.85	91.54	20.35	91.78	52.28	13.95	12.57	14.77	22.17	34.28	89.81	62.40
vpr3	6.00	7.67	25.00	78.67	14.67	78.67	30.67	8.67	6.00	7.67	16.67	31.33	86.00	53.67
vpr4	9.92	10.10	32.13	84.33	20.74	87.87	44.67	10.92	7.21	13.30	27.57	34.37	93.30	70.14

D - SupT1_30h_pi rep2

	8436	8442	8487	8532	8555	8563	8572	8632	9024	9074	9163	9400	9428	9442
envvpu1	8.00	8.67	29.00	83.33	16.33	82.67	31.00	7.33	6.67	7.33	16.00	31.67	87.33	54.67
envvpu10	10.04	13.43	42.38	84.45	20.74	84.34	28.64	10.04	9.16	9.37	15.80	44.52	90.33	52.34
envvpu13	11.25	10.50	37.82	88.60	23.48	86.85	34.33	9.75	11.90	10.92	27.77	38.29	91.43	67.07
envvpu17	7.00	9.00	25.67	82.00	16.67	83.00	25.00	8.00	6.00	7.33	14.00	34.67	90.67	53.67
envvpu19	11.76	12.37	21.95	80.58	18.20	74.26	12.62	5.77	10.69	6.07	14.18	31.64	81.57	55.50
envvpu2	6.33	11.00	33.33	82.00	17.33	82.67	31.83	7.00	6.33	8.00	16.33	33.33	86.33	55.33
envvpu3	10.00	9.67	31.00	82.00	12.67	82.67	40.27	5.33	6.67	6.67	13.67	31.00	85.33	54.67
envvpu5	11.00	13.00	34.33	87.67	22.33	89.67	38.43	10.33	8.67	11.33	19.67	38.33	87.67	64.00
envvpu9	9.67	11.33	32.67	86.33	20.00	87.00	37.67	8.33	7.33	7.67	18.00	33.33	88.00	58.67
nef1	13.33	16.33	39.67	93.33	25.67	90.33	42.67	9.00	10.67	10.33	21.67	44.00	90.67	60.67
nef12	12.86	16.06	40.30	89.27	26.76	86.66	30.48	11.27	8.87	9.13	14.84	35.55	89.22	53.35
nef2	11.67	14.00	37.67	90.67	23.00	90.33	44.00	10.33	8.00	9.67	18.00	37.33	88.67	60.00
nef3	11.00	13.00	36.67	90.33	25.00	90.67	46.33	11.33	8.33	9.67	20.00	38.00	86.33	62.00
nef4	10.67	12.33	37.00	90.67	26.33	90.67	47.33	11.33	9.33	9.33	20.33	37.67	89.00	60.33
nef5	10.67	14.00	38.67	90.67	25.67	91.67	44.00	11.00	10.00	12.67	21.33	40.67	86.67	62.00
nef9	9.33	12.33	38.67	90.33	22.00	90.67	44.72	10.67	7.33	10.67	17.00	37.00	88.00	60.67
rev1	9.67	11.67	33.67	90.33	22.00	89.67	48.00	10.67	7.67	9.00	18.00	36.67	89.00	58.33
rev10	13.59	13.33	45.58	85.73	28.45	92.12	51.27	11.54	9.16	15.19	31.03	42.81	89.40	64.77
rev11	10.55	11.91	37.78	87.59	30.72	92.47	31.67	6.56	12.77	13.86	17.80	45.54	87.64	66.33
rev2	10.00	11.33	32.33	90.00	22.67	89.33	44.33	9.67	8.00	10.67	18.67	35.67	88.00	58.67
rev3	8.03	12.48	32.47	88.74	21.95	88.51	29.44	7.93	6.87	6.84	15.89	40.58	87.56	58.70
rev4	9.33	12.33	33.67	89.67	25.00	89.00	45.07	9.33	10.33	11.00	18.67	37.67	86.33	61.67
rev5	11.11	13.88	39.50	88.82	20.74	87.76	46.06	12.71	3.55	7.06	21.04	34.63	85.67	57.48
rev7	11.00	12.00	35.33	88.67	25.67	90.33	41.65	11.67	10.00	10.00	19.00	38.67	89.00	62.33
rev8	9.67	12.00	35.67	90.33	25.00	92.67	50.03	12.33	12.67	11.33	20.67	37.00	88.67	59.33

SORF1	12.42	16.89	32.11	94.40	24.17	88.26	40.44	7.05	8.57	10.78	23.63	39.87	90.45	68.20
tat1	16.00	17.33	41.67	90.33	23.67	90.00	47.00	12.00	10.67	11.00	22.67	36.67	89.00	62.67
tat2	16.33	18.00	42.00	90.67	27.00	91.33	50.33	10.67	10.00	11.67	24.00	36.67	88.67	64.33
tat3	16.33	20.67	45.33	91.67	33.33	91.00	42.08	13.00	12.00	13.33	24.00	39.67	88.39	61.75
tat4	19.67	15.32	48.11	91.97	31.46	86.39	57.16	12.60	12.49	12.06	26.10	38.97	87.02	71.48
tat5	9.51	7.87	32.99	83.32	17.43	78.74	30.08	8.93	9.43	11.52	16.67	29.45	89.08	61.24
us	4.00	4.67	18.33	67.67	11.67	68.00	21.33	5.00	4.33	5.33	9.67	22.00	79.33	43.00
vif1	13.63	10.94	39.15	88.35	25.39	88.06	49.40	7.70	7.30	9.21	19.00	36.98	89.76	61.41
vif2	7.67	8.00	27.33	81.67	17.33	81.00	32.67	7.67	7.33	7.00	15.67	29.67	85.33	55.00
vpr1	14.67	16.00	45.67	91.00	26.67	93.33	45.95	15.33	10.00	9.33	21.67	35.67	92.00	65.67
vpr3	7.00	7.33	26.67	78.00	15.67	79.33	32.00	8.33	6.33	7.33	14.33	28.33	85.33	52.00
vpr4	10.00	8.67	36.33	85.00	18.67	88.00	44.12	14.67	9.67	7.67	25.46	39.67	91.33	58.33

E - CD4_48h_pi

	8436	8442	8487	8532	8555	8563	8572	8632	9024	9074	9163	9400	9428	9442
envvpu1	5.00	8.00	25.67	79.67	14.67	79.67	30.67	5.33	5.00	7.33	13.33	26.00	86.33	48.67
envvpu2	6.88	9.81	19.92	74.43	11.32	77.99	23.33	5.86	3.03	5.14	16.86	27.63	85.65	53.03
envvpu3	4.56	9.51	22.62	75.73	12.23	74.77	36.21	7.90	5.92	11.27	11.24	26.61	85.33	46.48
envvpu5	7.14	11.93	45.08	85.65	20.36	83.00	31.67	8.24	6.46	10.88	23.21	29.81	86.67	55.11
envvpu9	5.67	6.67	29.67	83.33	15.00	81.67	31.26	6.67	7.33	11.00	14.00	26.67	86.67	50.33
nef1	10.71	14.12	38.38	87.90	22.53	85.38	52.33	10.98	6.18	6.73	17.40	32.87	90.02	61.40
nef2	9.00	12.00	36.00	89.33	20.33	88.67	41.67	8.33	6.67	10.00	15.33	32.33	87.67	54.67
nef3	9.00	12.67	36.00	89.33	22.33	88.33	39.82	7.33	6.33	9.33	19.67	35.33	86.67	57.00
nef4	8.33	13.67	37.67	88.33	21.33	87.67	39.67	8.67	6.67	8.67	16.00	31.67	88.33	55.67
nef5	15.67	10.33	38.67	90.33	22.33	88.33	50.71	9.33	8.67	13.67	19.67	37.67	88.33	61.00
nef9	7.44	11.73	39.31	85.17	27.79	88.61	48.33	6.30	9.77	9.36	13.97	31.99	85.01	52.10
rev1	8.00	12.00	30.33	87.33	18.00	85.67	42.33	7.33	6.00	9.33	16.00	30.33	86.00	53.33

rev11	10.41	10.87	40.27	89.91	31.81	91.85	52.50	14.30	12.47	13.49	20.85	44.22	89.89	58.00
rev2	8.00	10.33	33.00	89.33	19.00	87.33	38.33	8.33	5.33	9.00	15.33	32.67	88.33	54.00
rev3	8.67	10.00	29.67	87.00	14.67	85.33	46.68	6.33	5.67	10.00	17.67	29.67	85.67	51.67
rev4	6.54	9.91	30.21	82.20	20.12	82.49	40.27	5.80	5.09	9.10	16.12	30.78	86.17	53.41
rev5	11.88	10.11	28.55	82.50	24.56	84.00	48.33	10.19	8.26	12.17	20.49	31.64	86.95	56.69
rev7	11.00	9.67	35.00	88.00	21.33	88.00	51.56	11.00	7.00	11.33	17.67	37.33	90.00	53.67
rev8	8.00	10.33	33.33	90.00	22.00	88.00	41.19	9.33	6.67	9.00	19.67	33.67	88.00	55.33
tat1	11.67	16.67	38.67	88.67	19.67	88.00	38.67	10.00	7.33	11.33	17.00	31.67	89.67	54.00
tat2	12.00	15.67	41.67	90.67	23.67	89.33	40.12	9.00	8.67	15.00	19.00	32.33	89.00	61.67
tat3	12.41	15.42	39.63	90.03	21.29	91.69	42.78	9.31	10.10	14.32	14.13	29.71	89.46	52.37
tat5	6.56	9.33	29.17	80.74	14.61	83.35	36.52	7.36	5.60	8.60	17.60	22.48	85.54	54.73
us	3.67	4.67	17.67	64.33	10.33	69.33	23.67	5.00	4.00	4.00	7.33	19.00	78.33	39.00
vif2	6.67	5.67	24.00	76.67	16.67	77.33	36.54	9.67	4.67	7.67	13.00	24.33	82.33	51.00
vpr1	7.48	11.70	36.88	92.20	20.21	87.73	40.27	10.35	9.30	14.51	22.44	35.42	84.85	56.12
vpr3	7.33	9.33	28.67	76.33	15.67	79.00	33.33	7.00	6.33	7.00	15.00	22.00	83.00	46.33
vpr4	9.58	16.33	37.28	88.04	19.93	91.14	58.94	9.19	11.48	9.34	16.39	33.22	85.62	53.77

744
745
746 **F - CD4_72h_pi**

	8436	8442	8487	8532	8555	8563	8572	8632	9024	9074	9163	9400	9428	9442
envvpu1	4.67	8.00	26.67	78.00	11.33	77.00	21.33	4.67	4.33	8.67	12.33	25.67	88.00	52.33
envvpu17	3.04	5.98	23.47	79.26	10.34	77.79	10.83	2.76	4.93	9.04	16.20	30.97	86.44	49.67
envvpu2	9.81	5.26	33.25	79.40	8.52	79.24	31.67	6.47	3.75	9.94	15.54	26.23	86.61	56.40
envvpu3	7.00	7.33	25.67	78.33	12.67	76.67	21.22	4.33	5.00	8.00	10.33	23.33	85.67	47.67
envvpu9	6.45	6.54	22.60	83.44	14.22	82.11	28.89	3.37	5.13	9.50	13.86	26.36	83.67	49.67
nef1	10.19	11.43	33.13	86.54	11.09	77.04	23.33	6.09	3.35	8.38	7.84	34.98	89.44	54.64
nef12	5.22	14.82	35.96	92.74	7.08	88.78	31.67	0.71	1.73	10.30	15.33	28.93	92.42	67.41
nef2	7.33	11.00	33.33	89.00	13.67	86.00	32.33	6.33	4.67	9.00	13.00	29.67	92.00	58.00
nef3	7.67	10.33	33.00	89.33	14.33	86.67	34.60	8.67	4.00	8.67	15.33	30.33	92.33	60.00

nef4	6.33	11.00	33.33	87.33	16.67	86.00	31.96	7.67	6.67	11.00	16.67	31.00	90.33	58.00
nef5	14.43	14.90	29.77	90.81	17.71	88.27	37.22	14.41	8.11	17.00	11.50	38.55	92.33	56.31
nef9	4.33	10.00	29.67	88.67	15.67	86.00	27.37	7.67	3.00	9.00	15.67	24.33	94.67	59.67
rev1	6.33	9.33	29.00	86.00	14.33	84.67	32.95	6.33	4.33	7.67	12.67	26.00	91.00	56.67
rev2	7.33	10.00	29.00	87.00	13.67	83.67	31.33	6.00	5.00	9.67	13.00	28.33	91.67	57.00
rev3	6.00	10.33	33.00	88.00	13.33	83.00	33.01	7.67	4.33	9.67	12.67	26.00	91.00	56.33
rev4	5.27	9.10	36.86	88.56	8.62	82.52	15.72	5.38	-0.33	6.33	11.00	26.69	92.42	59.45
rev5	4.30	15.41	36.27	89.67	13.95	87.63	45.00	3.37	6.21	10.24	17.81	34.24	90.67	65.05
rev8	7.62	9.22	26.06	87.77	13.13	81.19	28.64	5.19	4.04	8.19	15.77	24.06	91.15	57.47
tat1	9.00	14.67	35.33	88.00	14.67	86.33	32.67	7.00	6.00	12.67	15.67	27.00	92.67	59.67
tat2	9.33	14.67	34.67	90.00	16.67	88.00	33.57	7.33	7.33	12.00	18.00	32.67	91.67	63.00
tat5	4.34	7.47	29.48	73.75	14.33	79.96	22.06	7.59	7.59	9.82	16.89	22.01	88.27	57.73
us	2.67	4.67	16.33	60.00	7.33	62.33	16.33	4.00	3.00	5.33	8.67	17.33	77.67	38.67
vif1	9.57	14.21	32.94	75.81	21.43	81.91	28.33	11.70	14.35	8.08	22.00	29.61	81.69	48.69
vif2	7.67	6.67	23.33	79.00	15.33	77.33	23.50	5.33	10.00	9.00	16.67	21.00	86.33	50.67
vpr1	12.31	8.51	38.77	80.33	16.22	83.73	36.11	7.49	8.46	8.05	20.18	26.76	84.21	58.97
vpr3	5.33	8.67	23.00	73.67	13.00	74.33	24.84	5.00	5.00	7.00	16.67	22.00	84.00	48.33

Table S6. Rates of methylation of the 14 m^6A sites detected in all transcripts of the splicing isoforms found in the samples indicated. **A-F**, rates for the isoforms represented by ≥ 100 full-length reads (Materials and Methods).

Isoform		SupT1_30h_pi rep1			
CS - i4	PS + i4	%methyl (5613)	%methyl (5654)	%methyl (5703)	%methyl (5887)
Vif1		21	4	5.7	10.7
	Vif2	8.7	0	2.0	1
Vpr1		40.3	5	12	13.7
	Vpr3	10.3	0.3	1.3	0.3
Vpr2		39.0	8.0	18.5	15
	Vpr4	12.6	0.5	6.1	2.0
Tat1					16.7
	Tat5				18.8
Tat2					13
Tat3					13.3
Tat4					18.3
SORF1					16
US		1.3	0	0	0

Isoform		SupT1_20h_pi rep2			
CS - i4	PS + i4	%methyl (5613)	%methyl (5654)	%methyl (5703)	%methyl (5887)
Vif1		15.7	1.2	5.9	5.5
	Vif2	7.3	0.3	1.7	1.3
Vpr1		40.1	4.7	15.8	14.8
	Vpr3	8.7	0.3	2	1
Vpr2		32	6.7	19.2	16.3
	Vpr4	13.3	0.3	0.9	2.1
Tat1					17
	Tat5				24.3
Tat2					11.7
Tat3					16.6
Tat4					16.6
SORF1					16.7

US	1.0	0	0	0.3

Isoform		SupT1_30h_pi rep2			
CS - i4	PS + i4	%methyl (5613)	%methyl (5654)	%methyl (5703)	%methyl (5887)
Vif1		18.7	1.4	9.4	7.0
	Vif2	7	0.7	1.7	1
Vpr1		42.3	7.7	17	13.3
	Vpr3	9.3	1	2	0.3
Vpr2		41.0	5.5	13.8	6.4
	Vpr4	10	2.1	3	1.3
Tat1					16.7
	Tat5				19.7
Tat2					12.0
Tat3					14.3
Tat4					14.3
SORF1					16.1
US		1.0	0.3	0.3	0

756
757

Isoform		CD4_48h_pi			
CS - i4	PS + i4	%methyl (5613)	%methyl (5654)	%methyl (5703)	%methyl (5887)
Vif1		15.1	5.3	5.2	10.8
	Vif2	6.7	0.3	1.3	2
Vpr1		43.5	5.1	8.8	14.0
	Vpr3	8.7	0.7	1	0.7
Vpr2		42.2	12.3	18.7	33.0
	Vpr4	8.3	0	1.2	2.8
Tat1					19.3
	Tat5				23.5
Tat2					13.7

					13.5
Tat3					13.5
Tat4					16.6
SORF1					15.3
US	1.0	0	0		0.3

Isoform		CD4_72h_pi			
CS - i4	PS + i4	%methyl (5613)	%methyl (5654)	%methyl (5703)	%methyl (5887)
Vif1		9.8	1.0	4.5	6.0
	Vif2	8	1.0	2.3	1.3
Vpr1		27.1	4.4	13.9	9.1
	Vpr3	8	0.7	3	1
Vpr2		32.7	0	0	0
	Vpr4	13.3	0	3.4	3.6
Tat1					13
	Tat5				21.1
Tat2					8
Tat3					9.1
Tat4					20.4
SORF1					16.7
US		1.3	0.3	0	0.3

762 **Table S7.** Methylation rates for the A5613, A5654, A5703 and A5887 m^{6}A sites in the Vif, Vpr, Tat and SORF1

763 isoforms, as measured in the samples indicated. The first two columns indicate the isoforms completely

764 spliced (CS, lacking intron 4, - i4) or partially spliced (PS, containing intron 4, + i4).

A

SupT1_20h_pi rep1

	US				PS				CS			
	0	1	2	3	0	1	2	3	0	1	2	3
0	19	26	11	1	34	75	39	7	86	82	72	12
1	16	61	24	3	65	313	274	54	149	876	1066	290
2	9	80	75	22	109	679	864	261	561	4310	6522	2136
3	7	33	47	9	54	540	854	313	655	5241	9068	3503
4	1	8	16	1	26	208	433	202	284	2878	5581	2679
5	0	3	4	2	5	45	115	84	100	861	2005	1203
6	0	0	0	0	0	13	24	14	18	162	427	303
7	0	0	0	0	0	2	1	0	1	19	50	57
8	0	0	0	0	0	0	0	0	0	2	4	4

B

SupT1_20h_pi rep2

	US				PS				CS			
	0	1	2	3	0	1	2	3	0	1	2	3
0	341	503	167	18	194	396	220	34	147	143	124	29
1	366	1076	644	95	293	1373	1320	287	251	1508	1962	570
2	252	1280	1298	282	346	2877	3872	1212	722	6740	11069	4025
3	78	566	737	191	213	1826	3268	1234	739	6966	13411	5547
4	13	135	222	77	50	605	1268	615	332	3078	6830	3278

5	3	24	41	15	12	137	279	183	74	719	1696	982
6	0	5	7	5	1	19	26	30	4	75	247	168
7	0	0	0	1	0	3	4	0	0	4	19	18
8	0	0	0	0	0	0	0	0	0	0	0	0

772 **C**
773 **SupT1_30h_pi rep1**

	US				PS				CS			
	0	1	2	3	0	1	2	3	0	1	2	3
0	71	110	50	4	134	259	165	21	138	170	194	53
1	91	238	156	24	251	1125	961	208	375	2112	2709	881
2	52	308	278	71	316	2232	2757	804	1318	9691	14648	5491
3	29	134	157	31	156	1460	2359	938	1304	10289	17984	7611
4	5	28	52	12	58	481	955	452	596	4745	9458	4762
5	2	3	7	2	13	128	240	142	135	1211	2667	1634
6	0	0	1	0	0	19	43	22	17	170	442	349
7	0	0	0	0	0	1	1	4	1	19	52	47
8	0	0	0	0	0	0	0	0	0	0	3	5

775 **D**
776 **SupT1_30h_pi rep2**

	US				PS				CS			
	0	1	2	3	0	1	2	3	0	1	2	3

0	496	740	257	25	265	432	264	42	122	139	143	31
1	609	1681	1048	158	440	1736	1520	300	237	1301	1541	423
2	428	2110	1999	389	558	3630	4691	1300	854	5717	8325	2608
3	160	945	1174	284	319	2405	4023	1292	792	5991	9961	3600
4	22	215	344	94	105	847	1596	643	328	2685	5025	2192
5	6	32	53	27	17	152	371	160	28	584	1332	649
6	0	3	3	3	0	11	60	16	11	68	202	134
7	0	0	0	0	1	0	1	3	0	7	14	11
8	0	0	0	0	0	0	0	0	0	1	0	0

778 **E**
779 **CD4_48h_pi**

	US				PS				CS			
	0	1	2	3	0	1	2	3	0	1	2	3
0	81	95	35	7	94	151	84	16	51	78	52	23
1	90	253	134	12	123	562	370	59	120	623	613	125
2	64	293	277	45	172	1015	1097	211	320	2203	2606	736
3	14	156	155	26	103	688	746	215	275	2048	2930	901
4	2	30	34	7	24	208	306	106	104	790	1242	453
5	2	10	9	4	9	36	56	19	18	158	322	152
6	0	0	1	0	1	5	10	6	4	17	42	29
7	0	0	0	0	0	0	1	2	0	0	3	4

| 8 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 0 | 0 | 0 | 0 |

781 **F**
782 **CD4_72h_pi**

	US				PS				CS			
	0	1	2	3	0	1	2	3	0	1	2	3
0	148	250	73	4	78	165	77	10	55	111	91	16
1	129	418	214	32	108	513	425	68	168	1000	1021	200
2	82	419	366	68	116	902	943	256	402	3123	4221	1138
3	26	178	170	48	65	510	762	189	273	2264	3585	1129
4	7	31	43	12	9	126	211	78	67	700	1273	508
5	0	3	6	1	5	15	40	16	10	118	244	141
6	0	2	0	0	1	0	4	2	4	10	21	28
7	0	0	0	0	0	0	0	1	0	0	2	1
8	0	0	0	0	0	0	0	0	0	0	1	0

784 **Table S8.** Single-molecule analysis of methylated m^6A sites illustrating the correlation between the methylation
785 of clusters 1 and 2 of the same transcript, by sample. **(A-F)** Number of reads in the indicated samples with
786 defined numbers of methylated As in cluster 1 (by line: 0 to 8 m^6As) and in cluster 2 (by column: 0 to 3 m^6As).
787 US, unspliced; PS partially spliced; CS, completely spliced. One m^6A of one read is considered methylated if
788 the probability of base modification is ≥ 0.5 (value in modkit extract, Materials and Methods).

Position	Sequence	SupT1_20h_pi rep1	SupT1_20h_pi rep2	SupT1_30h_pi rep1	SupT1_30h_pi rep2	CD4_48h_pi	CD4_72h_pi
		% methyl.	% methyl.	% methyl.	% methyl.	% methyl.	% methyl.
2L125	UGACU	35.3	30.7	33.3	35	38.4 (19)	24.4 (25)
2L183	AGACU	11.7	9	13.7	9	6.5 (3)	4.1 (4)
2L272	GAACU	27.7	30.7	30.3	29	21.9 (11)	26 (24)
2L509	GAACU	32.7	37.3	35	32.7	26.8 (14)	19.7 (21)
2L537	GGACU	90.3	91.3	89.3	90	94.1	79.2
2L551	GGACU	66.3	66.7	68.7	70	69.2	64

Table S9. Rates of methylation of the six m^{6}A sites detected in the 2-LTR transcripts. The names of the sites refer to the position of the corresponding A in the HIV-1 reference genome. * indicates that the sequence context of the A9074 in the 2-LTR transcript (AGACT, Figure 6B) differs from the context of the same position, 9074, in the reference genome (GGACT). The number of methylated reads is indicated in brackets if below 30.

REFERENCES

1. Y. Saletore et al., The birth of the Epitranscriptome: deciphering the function of RNA modifications. Genome Biol **13**, 175 (2012).

2. J. Liu et al., A METTL3-METTL14 complex mediates mammalian nuclear RNA N6-adenosine methylation. Nat Chem Biol **10**, 93-95 (2014).

3. D. Dominissini et al., Topology of the human and mouse m6A RNA methylomes revealed by m6A-seq. Nature **485**, 201-206 (2012).

4. G. Zheng et al., ALKBH5 is a mammalian RNA demethylase that impacts RNA metabolism and mouse fertility. Mol Cell **49**, 18-29 (2013).

5. I. A. Roundtree et al., YTHDC1 mediates nuclear export of N(6)-methyladenosine methylated mRNAs. Elife **6** (2017).

6.   S. Zaccara, S. R. Jaffrey, A Unified Model for the Function of YTHDF Proteins in Regulating m(6)A-
Modified mRNA. *Cell* **181**, 1582-1595 e1518 (2020).

7.   S. Zaccara, R. J. Ries, S. R. Jaffrey, Reading, writing and erasing mRNA methylation. *Nat Rev Mol*
Cell Biol **20**, 608-624 (2019).

8.   S. U. Madugalle, K. Meyer, D. O. Wang, T. W. Bredy, RNA N(6)-Methyladenosine and the Regulation
of RNA Localization and Function in the Brain. *Trends Neurosci* **43**, 1011-1023 (2020).

9.   S. Phillips, T. Mishra, S. Huang, L. Wu, Functional Impacts of Epitranscriptomic m(6)A Modification on
HIV-1 Infection. *Viruses* **16** (2024).

10.   I. A. Roundtree, M. E. Evans, T. Pan, C. He, Dynamic RNA Modifications in Gene Expression
Regulation. *Cell* **169**, 1187-1200 (2017).

11.   X. Wang *et al.*, N6-methyladenosine-dependent regulation of messenger RNA stability. *Nature* **505**,
117-120 (2014).

12.   X. Wang *et al.*, N(6)-methyladenosine Modulates Messenger RNA Translation Efficiency. *Cell* **161**,
1388-1399 (2015).

13.   X. Zhao *et al.*, FTO-dependent demethylation of N6-methyladenosine regulates mRNA splicing and is
required for adipogenesis. *Cell Res* **24**, 1403-1419 (2014).

14.   S. Lavi, A. J. Shatkin, Methylated simian virus 40-specific RNA from nuclei and cytoplasm of infected
BSC-1 cells. *Proc Natl Acad Sci U S A* **72**, 2012-2016 (1975).

15.   G. Lichinchi *et al.*, Dynamics of the human and viral m(6)A RNA methylomes during HIV-1 infection of
T cells. *Nat Microbiol* **1**, 16011 (2016).

16.   E. M. Kennedy *et al.*, Posttranscriptional m(6)A Editing of HIV-1 mRNAs Enhances Viral Gene
Expression. *Cell Host Microbe* **19**, 675-685 (2016).

17.   N. Tirumuru *et al.*, N(6)-methyladenosine of HIV-1 RNA regulates viral infection and HIV-1 Gag protein
expression. *Elife* **5** (2016).

18.   S. Cristinelli, P. Angelino, A. Janowczyk, M. Delorenzi, A. Ciuffi, HIV Modifies the m6A and m5C
Epitranscriptomic Landscape of the Host Cell. *Frontiers in Virology* **1** (2021).

19.   C. Pereira-Montecinos *et al.*, Epitranscriptomic regulation of HIV-1 full-length RNA packaging. *Nucleic*
Acids Res **50**, 2302-2318 (2022).

20.   S. Chen *et al.*, N6-methyladenosine modification of HIV-1 RNA suppresses type-I interferon induction
in differentiated monocytic cells and primary macrophages. *PLoS Pathog* **17**, e1009421 (2021).

21.   K. Tsai *et al.*, Epitranscriptomic addition of m(6)A regulates HIV-1 RNA stability and alternative
splicing. *Genes Dev* **35**, 992-1004 (2021).

22.   A. B. R. McIntyre *et al.*, Limits in the detection of m(6)A changes using MeRIP/m(6)A-seq. *Sci Rep* **10**,
6590 (2020).

23.   D. R. Garalde *et al.*, Highly parallel direct RNA sequencing on an array of nanopores. *Nat Methods* **15**,
201-206 (2018).

24.   O. Begik, J. S. Mattick, E. M. Novoa, Exploring the epitranscriptome by native RNA sequencing. *RNA*
28, 1430-1439 (2022).

25.   M. Jain, R. Abu-Shumays, H. E. Olsen, M. Akeson, Advances in nanopore direct RNA sequencing. *Nat*
Methods **19**, 1160-1164 (2022).

26.   A. Baek *et al.*, Single-molecule epitranscriptomic analysis of full-length HIV-1 RNAs reveals functional
roles of site-specific m(6)As. *Nat Microbiol* **9**, 1340-1355 (2024).

27.   W. Liu-Wei *et al.*, Sequencing accuracy and systematic errors of nanopore direct RNA sequencing.
BMC Genomics **25**, 528 (2024).

28.   D. F. Purcell, M. A. Martin, Alternative splicing of human immunodeficiency virus type 1 mRNA
modulates viral protein expression, replication, and infectivity. *J Virol* **67**, 6365-6378 (1993).

29.   K. E. Ocwieja *et al.*, Dynamic regulation of HIV-1 mRNA populations analyzed by single-molecule
enrichment and long-read sequencing. *Nucleic Acids Res* **40**, 10345-10355 (2012).

30.   A. Emery, S. Zhou, E. Pollom, R. Swanstrom, Characterizing HIV-1 Splicing by Using Next-Generation
Sequencing. *J Virol* **91** (2017).

31.   N. Nguyen Quang *et al.*, Dynamic nanopore long-read sequencing analysis of HIV-1 splicing events
during the early steps of infection. *Retrovirology* **17**, 25 (2020).

32.   W. Li *et al.*, Characterizing transcripts of HIV-1 different substrains using direct RNA sequencing.
Heliyon **10**, e39474 (2024).

33.   K. D. Meyer *et al.*, Comprehensive analysis of mRNA methylation reveals enrichment in 3' UTRs and
near stop codons. *Cell* **149**, 1635-1646 (2012).

34.   C. Liu *et al.*, Absolute quantification of single-base m(6)A methylation in the mammalian transcriptome
using GLORI. *Nat Biotechnol* **41**, 355-366 (2023).

35.   Y. Zhou *et al.*, m6A sites in the coding region trigger translation-dependent mRNA decay. *Mol Cell* **84**,
4576-4593 e4512 (2024).

36.   H. E. Varmus, Viruses, genes, and cancer. I. The discovery of cellular oncogenes and their role in
neoplasia. *Cancer* **55**, 2324-2328 (1985).

37.   J. S. Smith, S. Y. Kim, M. J. Roth, Analysis of long terminal repeat circle junctions of human
immunodeficiency virus type 1. *J Virol* **64**, 6286-6290 (1990).

38.   S. Jurriaans, A. de Ronde, J. Dekker, J. Goudsmit, M. Cornelissen, Analysis of human
immunodeficiency virus type 1 LTR-LTR junctions in peripheral blood mononuclear cells of infected
individuals. *J Gen Virol* **73 (Pt 6)**, 1537-1541 (1992).

39.   T. Mishra *et al.*, Epitranscriptomic m(6)A modifications during reactivation of HIV-1 latency in CD4(+) T
cells. *mBio* 10.1128/mbio.02214-24, e0221424 (2024).

40.   B. Slobodin *et al.*, Transcription Impacts the Efficiency of mRNA Translation via Co-transcriptional N6-
adenosine Methylation. *Cell* **169**, 326-337 e312 (2017).

41.   N. Meiser, N. Mench, M. Hengesbach, RNA secondary structure dependence in METTL3-METTL14
mRNA methylation is modulated by the N-terminal domain of METTL3. *Biol Chem* **402**, 89-98 (2020).

42.   J. M. Coffin, S. H. Hughes, H. E. Varmus, "The Interactions of Retroviruses and their Hosts" in
Retroviruses*,* J. M. Coffin, S. H. Hughes, H. E. Varmus, Eds. (Cold Spring Harbor (NY), 1997).

43.   A. Brussel, P. Sonigo, Evidence for gene expression by unintegrated human immunodeficiency virus
type 1 DNA species. *J Virol* **78**, 11263-11271 (2004).

44.   J. Kelly *et al.*, Human macrophages support persistent transcription from unintegrated HIV-1 DNA.
Virology **372**, 300-312 (2008).

45.   R. D. Sloan, M. A. Wainberg, The role of unintegrated DNA in HIV infection. *Retrovirology* **8**, 52 (2011).

46.   S. Thierry *et al.*, Opposite transcriptional regulation of integrated vs unintegrated HIV genomes by the
NF-kappaB pathway. *Sci Rep* **6**, 25678 (2016).

47.   C. Richetta *et al.*, Two-long terminal repeat (LTR) DNA circles are a substrate for HIV-1 integrase. *J
Biol Chem* **294**, 8286-8295 (2019).

48.   X. Wei *et al.*, HIV Replication Under High-Level Cabotegravir Is Associated with the Appearance of 3'-
PPT Mutations, Circular DNA Transcription and Recombination. *Viruses* **16** (2024).

49.   A. Uzonyi *et al.*, Exclusion of m6A from splice-site proximal regions by the exon junction complex
dictates m6A topologies and mRNA stability. *Mol Cell* **83**, 237-251 e237 (2023).

50.   P. C. He *et al.*, Exon architecture controls mRNA m(6)A suppression and gene expression. *Science*
379, 677-682 (2023).

51.   P. Tang *et al.*, Nuclear retention coupled with sequential polyadenylation dictates post-transcriptional
m(6)A modification in the nucleus. *Mol Cell* **84**, 3758-3774 e3710 (2024).

52.   L. Bertrand *et al.*, Unveiling conserved HIV-1 open reading frames encoding T cell antigens using
ribosome profiling. *Nat Commun* **16**, 1707 (2025).

53.   F. Bouchonnet *et al.*, Quantification of the effects on viral DNA synthesis of reverse transcriptase
mutations conferring human immunodeficiency virus type 1 resistance to nucleoside analogues. *J Virol*
79, 812-822 (2005).

54.   J. M. Watts *et al.*, Architecture and secondary structure of an entire HIV-1 RNA genome. *Nature* **460**,
711-716 (2009).

908