Conserved strategies underpin the replicated evolution of aridity tolerance in trees

Ben Halliwell[1,2,5], Chris J. Blackman[1,2], Rachael V. Gallagher[1,3], Luke A. Yates[1,2], Travis G. Britton[1,3], Rebecca C. Jones[1,2], Dean Nicolle[4], Gabrielle E. Hartill[1,2], Ian J. Wright[1,3], Barbara R. Holland[1,2] & Timothy J. Brodribb[1,2]

[1]ARC Centre of Excellence for Plant Success in Nature and Agriculture, University of Queensland, Brisbane, QLD, Australia

[2]School of Natural Sciences, University of Tasmania, Hobart, Tasmania, Australia

[3]Hawkesbury Institute for the Environment, Western Sydney University, Penrith, NSW, Australia

[4]Currency Creek Arboretum, Currency Creek, SA, Australia

[5]Corresponding author. Email: benjamin.halliwell@utas.edu.au

Abstract

Predicting forest responses to climate change requires a detailed understanding of trait-environment coordination. Adaptation to environment comprises both conserved and labile components of trait variation. However, few studies explore the decomposition of trait-environment relationships in a rigorous phylogenetic framework. Combining a revised phylogeny with trait, climate and soil data to achieve unprecedented species coverage (~85%), we identified patterns of replicated evolution that allowed the evergreen tree genus, *Eucalyptus*, to rapidly radiate across Australia in response to aridification. Eucalypts from arid regions are short, produce dense wood, and have small, physically robust leaves with high nitrogen content, promoting hydraulic safety and economies in photosynthetic water use. Phylogenetic modelling reveals strong niche conservatism, with adaptation to aridity occurring primarily via clade-level divergences, followed by phylogenetically independent adjustments to local conditions. Ancestral state reconstructions accounting for trends in the paleoclimate record indicate that transitions in aridity tolerance are associated with distinct signals of environmental filtering and directional selection on functional trait variation. However, astonishing repeatability of trait changes in different clades reveals a narrow optimal solution to water availability, opening a path to predict future species distributions from phylogenetically structured trait data and signalling major implications for functional and species diversity under progressive climate change.

Keywords: functional traits, plant hydraulics, plant strategy, water-use-efficiency, leaf economics, heat, drought, phylogenetic comparative analysis, replicated evolution, tree, Eucalyptus.

Introduction

Global increases in tree mortality[1,2] and forest die-back events[3-6] bring an urgency to understand the process of climatic adaptation in woody plants[7,8]. To predict ecosystem responses to climate change, we must understand which species traits confer resilience to temperature and drought stress, including the decomposition of factors that contribute to this functional trait variation across species and environments[9-11]. This knowledge is critical to effective planning and protection of natural and managed forest systems into the future, both of which are vulnerable to climate induced increases in tree mortality[12,13].

Comparative analyses emphasise unifying trait spectra that relate plant functional variation to species environmental niches[14-17]. Trait coordination can be understood in terms of economic and physiological trade-offs, such as between tissue lifespan and productivity[17,18], and the safety and efficiency of water transport[19,20]. Interspecific variation in functional traits therefore reflects the evolutionary outcomes of these trade-offs for fitness across environmental gradients[21,22]. In woody species, adaptation to aridity is thought to involve coordination among trait spectra, leading to whole-plant strategies that integrate evolutionary responses in hydraulic, economic and photosynthetic function[23-26]. However, a major challenge for predicting forest responses to environmental change is that trait-environment relationships vary considerably[27] due to inherent biological and biogeographic differences between plant clades [28-31]. One consequence is that broad-scale trait-environment relationships are often poor predictors of local-scale plant responses[9,32]. Phylogenetic models of trait-environment coordination within prominent clades offer a promising avenue for overcoming these challenges of model transferability[33,34]. However, few studies have attempted to decompose trait-trait and trait-environment relationships in such a rigorous phylogenetic framework.

Genus *Eucalyptus* presents a model system to study adaptation to aridity in evergreen trees. Emerging 50-60 Mya in the warm and humid Paleogene[35], fossil evidence suggests eucalypts became increasingly abundant during the Miocene (23-5 Mya), coincident with the onset of progressive climatic drying and depletion of soil nutrients across the Australian continent[36,37]. This rapid environmental change imposed strong selection and environmental filtering on ancestral eucalypts, leading to multiple independent transitions into arid environments in different lineages (figure 2). These recent, replicated evolutionary events, along with the functional and ecological diversity of eucalypts, present an outstanding opportunity to decompose drivers of trait coordination during adaptation to aridity[38]. Recent drought-induced canopy die-back of eucalypt forest in Australia[39-41], promoting catastrophic wildfires[42,43], underscores the urgency of predicting future changes in eucalypt forest to mitigate impacts of climate change on both natural and managed systems.

Drawing on the latest phylogenetic evidence for *Eucalyptus*[44], we apply multivariate phylogenetic models to assess trait-trait and trait-environment coordination. We leverage a dataset of >22,500 trait observations covering ~85% of *Eucalyptus* species[45], by combining AusTraits[46] data records with targeted sampling of >200 species (table 1). We combine these trait data with environmental data observed at two spatial scales (species mean and site-level) in a single analysis to simultaneously assess 1) the independent effects of climate and soils on trait variation; 2) the decomposition of trait-trait and trait-environment relationships across

phylogenetic and non-phylogenetic variance components; 3) whether past changes in trait
values represent causes or consequences of transitions to more arid environments.

Results

Adaptation to aridity represents a whole-plant strategy

We focus on 6 traits with well-established links to hydraulic, economic and photosynthetic
function: leaf area, leaf mass per area (LMA), leaf N per area (N_{area}), discrimination between ^{13}C
and ^{12}C carbon isotopes in leaf tissue (δ^{13}C), wood density, and species maximum potential
height (max height); and 4 key dimensions of species environmental niche: mean annual
temperature (temperature), mean aridity index (moisture), mean soil phosphorus concentration
(soil P) and mean soil bulk density (bulk density).

Two principal components explained 66.7% of the phenotypic variation across the 6 traits and 4
environmental niche variables (figure 1; table S1; S2). PC1 (49.3%) captured variation in
moisture, soil P and a suite of functional traits, whereas PC2 (17.4%) primarily captured
variation in temperature, soil bulk density and leaf N_{area}. In contrast to results from global
analyses reporting independent axes of variation in size and economic traits[14,15], our targeted
analysis of *Eucalyptus* reveals extensive coordination between these traits, consistent with
variation along a whole-plant fast-slow economic spectrum (figure 1). Eucalypts occupying hot
dry environments are characterised by short stature, dense wood, small leaves, as well as high
LMA, N_{area} and δ^{13}C (figure 1). The phylogenetic distribution of functional trait variation indicates
that divergence along this trait spectrum has occurred repeatedly during adaptation to aridity
gradients within different eucalypt clades (figure 2).

Decomposing trait-environment relationships

Understanding the process of climatic adaptation at different biological scales requires that we
separate trait variance due to local environmental conditions from that due to conserved trait-
environment relationships[47]. The multi-response phylogenetic mixed model (MR-PMM)[48]
presented here leverages an unprecedented dataset for *Eucalyptus*, both in terms of
phylogenetic coverage and within-species replication of functional traits (table 1), to jointly
estimate and partition these effects (figure 3). These analyses allow us to decompose
correlations between species traits and environmental niche variables into phylogenetic and
non-phylogenetic components. Furthermore, for trait observations matched with
environmental data from the site individuals were sampled, we calculated environmental
anomalies (standardized differences between species mean environmental niche values and
site-level environmental conditions) for each species-site combination. By fitting species-
specific environmental anomalies as fixed effects in the MR-PMM, we can separate the
influence of site-level environmental conditions on trait variation *within-species*, when
estimating correlations between traits and niche variables *between-species*. We first consider
the effect of environmental anomalies, then explore the decomposition of trait-trait and trait-
environment correlations within components of variance that remain unexplained by these site-
level effects.

Responses to local environment

Within-species variation in functional traits was significantly related to site-level environmental anomalies (figure 3A), indicating adjustments in trait values to deviations from species mean environmental conditions. On average, individuals from warmer sites have higher density wood, larger leaves, lower N_{area} and more negative $\delta^{13}C$, implying less efficient water use during photosynthesis. However, elevated temperature had similar effects to elevated moisture on trait values. For example, eucalypts respond to both hotter and wetter sites by decreasing N_{area} (figure 3A). This suggests a partly compensatory mix of trait responses to hotter and drier (i.e., more arid) sites, which predominate given strong negative covariance between temperature and moisture across the bioclimatic niche space of *Eucalyptus* (figure 2, figure 3C). Climate effects were particularly pronounced for leaf $\delta^{13}C$, highlighting adjustments in photosynthetic properties as a key response to site-level temperature and water availability. Soil effects were detected for LMA, N_{area} and wood density, indicating a shift to more conservative growth strategies in challenging soil conditions. Specifically, eucalypts respond to low relative soil P by increasing LMA and leaf N_{area}, and to high relative soil bulk density by increasing wood density (figure 3A).

Drivers of trait variation

Fixed effects of environmental anomalies explained considerably more variance in wood density and leaf $\delta^{13}C$ than any other functional traits (figure 3B). However, these within-species, site-level effects explained only a tiny fraction (1-3%) of total trait variance (figure 3B; table S3). Study effects consistently explained a greater proportion of trait variance than fixed effects, likely due to characteristics of study sites not captured by our fixed effects (e.g., micro-climate, topography, biotic factors), as well as differences in methodology or measurement technique between studies. Non-phylogenetic between-species effects (i.e., effects of species identity after accounting for phylogenetic relationships) were small (5-11%) and generally attributed less variance than either study effects or residual errors, indicating that divergences between closely related species contribute relatively little to observed trait variation. In contrast, for every trait except leaf $\delta^{13}C$, the phylogenetic component of variance was greater than all other variance components combined (figure 3B). Indeed, phylogenetic signal ($0 < \lambda < 1$) for functional traits ranged from 0.48 (leaf $\delta^{13}C$) to 0.87 (max height), and was extremely high for species mean environmental variables ($\lambda > 0.85$), reflecting strong niche conservatism.

Phylogenetic versus non-phylogenetic correlations

Decomposing between-species trait correlations into phylogenetic and non-phylogenetic components confirms that similar relationships operate on both levels (figure 3C), broadly reflecting patterns observed in PCA of phenotypic data (figure 1). This indicates a relatively scale-free evolutionary response to environmental gradients, with similar patterns of trait coordination in the climatic divergence of closely related species as in the climatic divergence of major clades over deeper time (figure 2, figure S1). However, phylogenetic correlations were considerably stronger (figure 3C, lower triangular), emphasising the ecological significance of clade-level divergences (figure 2).

Identifying a core network of integrated trait responses

To identify candidate causal relationships between response variables within the major phylogenetic component of trait variance, we computed partial correlations from the fitted phylogenetic covariance matrix. These correlations revealed a highly integrated network of trait-trait and trait-environment relationships, consistent with a conserved, whole-plant response to environmental gradients (figure 3D). Trait coordination can be described as a trade-off between two modules of positively covarying traits: a 'fast' acquisitive trait module comprising size-related traits (leaf size and max height), and a 'slow' conservative trait module comprising traits with hydraulic, economic and photosynthetic functions (wood density, LMA, N_{area}, $\delta^{13}C$, figure 3D). This trade-off defines a major axis of ecological strategy in *Eucalyptus*, separating species across a continuum between productivity and persistence in a whole-plant 'fast-slow' economic spectrum.

Moisture, temperature, and soil P each displayed (non-zero) partial correlations with a range of traits, indicating that adaptation to these environmental factors involves direct responses in hydraulic, economic and photosynthetic function (figure 3D). Moisture was strongly negatively associated to traits within the 'slow' trait module (wood density, LMA, N_{area}, and $\delta^{13}C$), where-as evidence for positive associations with the 'fast' trait module (max height, leaf size) varied depending on the taxonomic scale at which phylogenetic effects were modelled (species- vs series-level topology, figure S2D). Pairwise partial correlations generally supported a direct functional link between moisture and max height (table S4, figure S2D), but suggest a relative decoupling between moisture and leaf size. We find particularly strong partial correlations between temperature and each of wood density, leaf area, N_{area}, and $\delta^{13}C$, suggesting tight coordination between these traits and the temperature niche of eucalypt lineages. Partial correlations between soil bulk density and both max height and wood density highlight adjustments in hydraulic properties as a primary mechanism of adaptation to reduced soil water holding capacity (figure 3D; figure S2D). Taken together, these results emphasise a coordinated strategy favouring hydraulic safety, conservative growth economics, and high photosynthetic water-use-efficiency (WUE) as a consistent feature of adaptation to hot, dry, and poor soil environments in *Eucalyptus*.

Ancestral state reconstructions

Coordination between plant traits and environment can arise by filtering on standing variation in trait values, i.e., during colonisation or rapid environmental change (environmental filtering)[49], or by directional selection on trait values, i.e., adaptation *in situ* (local adaptation)[50]. To distinguish the relative importance of these processes, we performed ancestral state reconstructions that account for historical shifts in the Australian paleoclimate to evaluate whether changes in trait values are likely to represent a cause (environmental filtering) or consequence (local adaptation) of transitions to more arid environments (see methods). To investigate whether similar processes have acted on functional trait variation at different points along the aridity continuum, we compared the inferred sequence of trait changes during transitions between mesic (M), semi-arid (S) and arid (A) environments. Transitions from M (moisture index > 0.5) to S (moisture index < 0.5), and from S to A (moisture index < 0.2), environments have occurred many times independently across different clades within our

broad phylogenetic sample of *Eucalyptus* (figure 4A; table S5). This analysis of replicated
evolution reveals that selection has repeatedly produced the same solution during transitions
to more arid environments. Node categories representing transition types between different
aridity states were strongly associated with variation in reconstructed trait values (figure S3),
reflecting a progressive shift toward slower leaf and wood economic strategies, and higher
photosynthetic WUE, with increasing aridity (figure 4B).

To test for differences in reconstructed trait values between node categories while accounting
for phylogenetic relationships, we computed contrasts from posterior samples of an MR-PMM
specifying node category as a fixed effect when estimating ancestral states. Figure 4C presents
results for wood density, LMA, and $\delta^{13}C$, traits that each show partial correlations with moisture
(figure 3D), and relate broadly to hydraulic, economic and photosynthetic functions (results for
other traits shown in figure S4). Evidence for environmental filtering during transitions from M to
S environments (M_S versus M_M) was limited to a subset of traits; M ancestors giving rise to S
descendants (M_S) had lower max height, as well as marginally higher wood density and LMA,
than M ancestors giving rise only to M descendants (M_M) (figure 4C; figure S4). For most traits,
evidence was stronger for directional changes during persistence in S environments (S_S versus
M_M), consistent with local adaptation to water limitation (figure 4C; figure S4). However,
results differed when considering transitions from S to A environments, representing tolerance
of more extreme water limitation. We find stronger evidence for environmental filtering,
particularly on LMA, leaf N_{area} and $\delta^{13}C$, during transitions from S to A environments (S_A versus
S_S). We also find strong evidence for directional changes in trait values during persistence in A
environments (A_A versus S_S). Finally, we investigated whether reversals to less arid
environments were associated with opposing changes in the same network of traits. Consistent
with this, we observed coordinated reversion of trait values toward a faster, riskier, and more
acquisitive strategy during reversals from A to S, and from S to M, environments (figure 4C,
figure S4).

Table 1. Sample size information and summary statistics for each functional trait and environmental niche variable within the dataset used to fit MR-PMM. N_{obs} = total number of observations across species, $N_{species}$ = number of species with data. The mean number of observations per species (and per taxonomic series) for each trait demonstrates good within-species replication across the 767 species included in analyses.

	N_{obs}	$N_{species}$	proportion of species	mean number of obs per species	mean number of obs per series	min value	max value	units
leaf area	8727	766	0.99	11.4	72.1	155	16422	mm^3
LMA	6820	625	0.81	10.9	60.9	42	932	g/m^2
N_{area}	2521	504	0.66	5	25.2	0.17	6.19	g/m^2
$\delta^{13}C$	1860	505	0.66	3.7	18.8	-31.9	-23.2	‰
wood density	1930	385	0.50	5	19.5	0.4	1.19	mg/mm^3
max height	761	761	0.99	-	-	1	99.6	m
temperature	767	767	1	-	-	6.3	27.8	°C
moisture	767	767	1	-	-	0.06	2.25	unitless
soil P	767	767	1	-	-	0.007	0.033	‰
soil bulk density	767	767	1	-	-	1.08	1.7	g/cm^3

Figure 1. **Adaptation to aridity in *Eucalyptus* represents a whole-plant strategy.** Principal components analysis of functional traits and environmental factors using data gap-filled with predictions from the fitted MR-PMM. PC1 (49.3% of variance) primarily captured trade-offs in functional traits along an axis of variation in moisture index and soil variables. PC2 (17.4% of variance) primarily captured variation in temperature, soil bulk density and N_{area}. PC loadings and cumulative variance are shown in tables S1; S2.

Figure 2. **Conserved strategies underpin the replicated evolution of aridity tolerance in different *Eucalyptus* clades**. Phylogenetic distribution of functional trait values and environmental niche variables for *Eucalyptus* ($N_{species}$=767) using gap-filled data imputed by MR-PMM. Taxonomic series-level topology (left) coloured by subgenera is plotted against a matrix of series-level means (centre) for each trait and environmental niche variable. Scatterplots (centre) show series-level means for each trait and niche variable ordered by moisture. The phylogenetic distribution of trait values reveals that multiple clades have evolved the same set of trait values to cope with dry climates, evident as blocks of similar trait values among clades characterised by low moisture. Remarkable convergence in whole-plant phenotype to moisture availability is observed across multiple independent origins of aridity tolerance in different clades (right): a) *E. deserticola*, b) *E. repullulans*, c) *E. gamophylla*, d) *E. grandifolia,* e) *E. pileata,* f) *E. erythrocorys,* g) *E. miniata,* h) *E. leiocarpa,* i) *E. gummifera,* j) *E. microcorys,* k) *E. globulus,* l) *E. regnans* (photos by Dean Nicolle).

Figure 3. Within-species, traits vary with site-level environmental conditions, but effects are relatively weak. Decomposition of the remaining trait (co)variance across species reveals strong phylogenetic signal underlying a conserved network of trait-trait and trait-environment relationships. Results from MR-PMM fit to 100 candidate series-level topologies ($N_{species}$ = 767): **A** Posterior fixed effect estimates of site-level environmental anomalies for each functional trait associated with site-level data. **B** Variance decomposition for each functional trait and environmental niche variable. Traits that are constant for a species are partitioned into phylogenetic and residual components, while traits varying within-species are partitioned into phylogenetic, non-phylogenetic, study, fixed, and residual components. **C** Phylogenetic (lower triangular) and non-phylogenetic (upper triangular) between-species correlations between response variables from MR-PMM. Values represent the posterior mean estimate for each correlation coefficient. **D** Network graphs showing significant (90% and 95% CI) partial phylogenetic correlations (positive = blue edges, negative = red edges) between functional traits (green), climate (light blue) and soil (orange) niche variables. Edge widths are proportional to partial correlation coefficients. All correlation estimates derive from the same MR-PMM model fit but are grouped and plotted in separate networks for illustrative purposes. Note that partial phylogenetic correlations between max height and environmental niche variables were markedly stronger when phylogenetic random effects were modelled using the species-level topology (figure S2D).

Figure 4. **Ancestral state reconstructions suggest that adaptation to moisture gradients in *Eucalyptus* has involved both environmental filtering and directional selection on functional trait values, with more stringent filtering during the colonisation of arid environments. A** Species level phylogeny ($N_{species}$ = 631) showing major Eucalypt clades. Branch lengths have been transformed[51] by δ = 10 for plotting. Branch colour maps an ancestral state reconstruction of species mean moisture index (MI) according to a shift model that accounts for strong directional change toward drier climates in Australia since the Miocene (23 Mya). Internal nodes are classified according to their own reconstructed values of MI (mesic = MI > 0.5, semi-arid = 0.2 < MI < 0.5, arid = MI < 0.2) and those of their descendants (see methods). Posterior mean counts for each node category are indicated in parentheses within the legend. Tip labels indicate whether the corresponding species MI value classifies it as mesic (purple), semi-arid (yellow) or arid (red). **B** Posterior distributions of trait values at internal nodes for each transition category from ancestral state reconstruction. **C** Contrasts between node transition categories representing different stages in the evolution of aridity tolerance (see methods). Contrasts are computed and reported for transitions between mesic and semi-arid environments (green), as well as between semi-arid and arid environments (cream). The mean (point), 50% (heavy whisker) and 95% (light whisker) CI of each posterior distribution are shown in black.

Discussion

Drawing on data for 85% of *Eucalyptus* species, we find strong and consistent functional trait
responses during adaptation to environmental gradients. Species respond to site-level
deviations from mean environmental niche values with finely tuned adjustments in a range of
functional traits (Figure 3A). However, these effects are generally small and insufficient to
explain broad trait-environment relationships across species (Figure 3B). Instead, strong
phylogenetic signal reveals that these relationships are driven primarily by conserved
components of trait covariance, reflecting a replicated pattern of trait changes during the
climatic divergence of eucalypt clades. Traits associated with hydraulic, economic and
photosynthetic functions showed strong relationships with species moisture niche, consistent
with an integrated whole-plant response to aridity gradients (figure 1; figure 3C).

Pairwise trait relationships were broadly consistent with predictions from leaf economic
theory[17], least-cost theory[52,53], pace of life syndromes[26] and the allometry of hydraulic traits[54,55]
(figure 3A). These include positive correlations between LMA and wood density (linking leaf and
wood economic spectra), N_{area} and $\delta^{13}C$ (linking carboxylation capacity and stomatal limitation
on photosynthesis), and leaf size and max height (linking variation in the size of plant organs), as
well as negative correlations consistent with trait trade-offs, including between wood density
and max height (stress-tolerance versus growth rate). A strong positive relationship between
LMA and N_{area} emphasises coordination between leaf economics and the manner in which
nitrogen and water use in photosynthesis is optimised in relation to water limitation[53,56], but
also contributions of nitrogenous compounds other than RUBISCO, especially cell walls, which
also increase LMA[57,58].

More broadly, the functional trait network we identify indicates coordination within and
between plant tissues; opposing modules of traits (figure 3D) define a spectrum of whole-plant
strategy from productivity to persistence that integrates tissue economics, hydraulic
vulnerability, and photosynthetic WUE[16,26]. These results are consistent with detailed
comparative ecophysiology studies of *Eucalyptus*[59-62], which report extensive coordination
between economic, hydraulic and photosynthetic functions. Indeed, the conserved trait
network we identify may have been key to the ecological success of *Eucalyptus*, by aligning
major axes of functional variation[63] with the harsh aridity gradients that have come to
characterise the Australian continent.

The trait-environment relationships we find for *Eucalyptus* accord well with those reported for
other woody taxa, indicating species from arid environments tend to be short[55], produce dense
wood[16], and have small, robust leaves with more active stomatal regulation[23]. Of the traits
examined here, wood density and max height have been linked most directly to the capacity of
trees to tolerate acute water-stress[64-68]. Previous studies considering a smaller number of
Eucalyptus species have linked wood density to moisture availability[69] ($N_{species}$ = 28), as well as
directly to measures of hydraulic vulnerability[70] ($N_{species}$ = 20). We confirm a strong association
between wood density and aridity tolerance across a much larger sample of *Eucalyptus* in this
study ($N_{species}$ = 767), including partial correlations with both the temperature and moisture axes
of species' environmental niches. Our analyses also support strong associations between

species max height and environmental niche variables (figure 1; figure 3C), consistent with
previous work on *Eucalyptus*[62,71]. Evidence for partial correlations between max height and
climate varied depending on the scale at which phylogenetic effects were modelled, with
stronger evidence for these effects when using the species-level phylogeny (figure 3D; figure
S2D). This may suggest that conserved relationships between max height and climate can to
some extent be explained by covariance with other functional traits and environmental factors.
However, as max height is defined at the species level, a likely explanation for this model
dependence is that lower order divergences in the phylogeny (i.e., species-level relationships)
provide the necessary resolution to detect conserved relationships between max height and
environmental factors. Indeed, when using the species-level phylogeny, max height and wood
density showed equally strong partial phylogenetic correlations with moisture (figure S2D).

Across all analyses, LMA showed a strong relationship with the moisture niche of *Eucalyptus*
species. This contrasts with global analyses, which indicate that LMA is only weakly associated
with climatic variation at both the species and community trait level[14,72] (though see[9]). Our
analyses show that, within phylogenetically constrained species groups, LMA may be tightly
coupled with climatic variation. Indeed, ancestral state reconstructions identified both
environmental filtering and directional selection on LMA as features of transitions into arid
environments (Figure 4C). This suggests that, along with traits that confer resilience to acute
drought stress, successful evolutionary strategies involve optimising leaf economics toward
slow, conservative growth[26]. Extreme phylogenetic signal in LMA (λ = 0.85) further indicates an
evolutionary rigidity in leaf economic strategies, with individual clades occupying narrow
regions of the economic spectrum. LMA was also strongly negatively related to soil P content
(figure 3D), agreeing with global analyses that link leaf economics to soil fertility, in particular
the availability and cost of acquiring essential soil nutrients[14].

Both N_{area} and $\delta^{13}C$ displayed partial correlations with moisture and temperature niche,
indicating fundamental roles of WUE and stomatal regulation in the capacity of eucalypt
species to persist in arid environments. Higher N_{area} implies greater investment in the
carboxylating enzyme RUBISCO, meaning that, at a given level of stomatal conductance, leaf-
internal CO_2 during photosynthesis is lower – leading to more efficient water use (higher WUE)
for a given photosynthetic rate per unit leaf area (A_{area}) and site VPD[53]. Lower ci:ca is reflected in
leaf $\delta^{13}C$[73,74], which may explain the exceptionally strong positive correlations between N_{area} and
$\delta^{13}C$ observed in our analyses (figure 3C, figure 3D). While WUE is clearly crucial for plant
performance in hot dry conditions, the negative partial correlation between N_{area} and
temperature can be explained by reduced investments in RUBISCO with increasing efficiency of
carboxylation at higher temperatures[52,56,75].

Of all functional traits, leaf area showed the weakest relationships with both species mean
environmental niche variables and site-level environmental conditions (figure 3A). Despite this,
leaf area was strongly associated with max height, wood density and $\delta^{13}C$, consistent with a
role of leaf size in plant allometric relationships affecting hydraulic demand and water-use
strategies[54].

Ancestral state reconstructions indicate that coordinated trait changes represent both a cause
and a consequence of the colonisation of water-limiting environments in *Eucalyptus* (Figure 4B,

C). Evidence for environmental filtering on trait variation was generally stronger during
transitions from semi-arid to arid environments (S_A) compared with transitions from mesic to
semi-arid environments (M_S), particularly for leaf traits (figure 4C, figure S4). This supports the
notion that extreme environments impose greater constraints on viable trait combinations,
leading to more stringent trait filtering[76]. A notable exception was max height, which showed
stronger effects during transitions from mesic to semi-arid environments (figure S4), consistent
with the drastic reduction in tree height observed between wet- and dry-sclerophyll eucalypt
forest (figure 2). Our analyses further suggest that, following transitions to more water-limiting
environments, ancestral eucalypts persisting in those environments underwent strong
directional evolutionary change, becoming markedly shorter, with higher wood density, LMA,
N_{area}, and $\delta^{13}C$ (figure 4; S2). Finally, during reversals to less water-limiting environment (S_M
and A_S), we observed an opposing pattern of trait changes consistent with environmental
filtering on the same coordinated suite of traits. The astonishing repeatability of trait changes
across multiple independent transitions in aridity tolerance suggests a narrow optimal solution
to water availability in *Eucalyptus*[38]. Indeed, similar trait spectra characterise drought tolerant
species across many clades and biogeographic regions, suggesting convergence on a
fundamental archetype of water stress tolerance in trees[9,77].

Most variation in functional traits and species environmental niche can be attributed to
phylogenetic effects (figure 3D), reflecting major ecological divergences between eucalypt
clades (figure 2). This emphasizes phylogenetic niche conservatism[47,78] as a major feature of
climatic adaptation in *Eucalyptus*, mirroring findings for woody species at broader phylogenetic
scales[79,80]. Indeed, the progressive emergence of aridity gradients in Australia since the
Miocene (23 Mya)[36,37] is consistent with a biogeographic history of species filtering processes
followed by rapid adaptation of aridity tolerant lineages, invoking coordinated changes in traits
that jointly maximised fitness in these novel environments; a scenario well supported by our
ancestral state reconstructions (figure 4C). Niche conservatism demonstrates that organisms
tend to establish and persist in environments to which they are already well adapted[81].
Examples of arid niche conservatism in *Eucalyptus* include the speciose sections of subgenus
Symphyomyrtus; *Bisectae*, *Dumaria* and *Glandulosae* (figure S1). These clades radiated very
recently (5-10 Mya), together accounting for the majority (78%) of arid distributed *Eucalyptus*
species and are each characterised by the same suite of traits conferring slow growth and
stress tolerance (figure S1). Such clades are likely to expand in distribution as climate change
progresses, with profound implications for the functional and phylogenetic diversity of
Australian forests[82], including fire regimes[83] and ecosystem services[84]. Specifically, our results
suggest that the initial impact of increases in semi-arid habitat with climate change[85] is likely to
be strong species filtering based on tree height, wood density and LMA, promoting a rapid
transition from tall, closed-canopy forest to sparse, open woodland, with consequences for
primary productivity, carbon sequestration, and the frequency and intensity of wildfires.

The remarkable strength of phylogenetic signal (figure 3D) in functional traits and
environmental niche variables we report for *Eucalyptus* confirms that phylogenetic information
is severely underutilised in global predictive vegetation models[86-88]. Indeed, species hydraulic,
economic, photosynthetic, and phenological traits are strongly conserved at multiple
phylogenetic scales[89-91], presenting a tremendous opportunity to improve prediction of future

species distributions and abundances[87], as well as trait-related ecosystem services such as
primary productivity and carbon sequestration[92]. For *Eucalyptus*, a paleoclimatic trend of
increasing aridity induced the adaptive radiation and competitive dominance of the drought
tolerant *Symphyomyrtus* clade (figure 2), while reducing many humid adapted Gondwanan
lineages to relictual species[93] (figure 2j). As climate change progresses, further declines in the
viability of 'fast', hydraulically risky, economic strategies across multiple biomes puts entire
clades at risk of extirpation and extinction[94], with major consequences for the functional and
phylogenetic diversity of ecosystems[95,96]. Similar patterns of conserved hydraulic strategy
within prominent woody clades such as *Quercus*[97], *Juniper*[98], *Magnolia*[99] and Fabaceae[100], as
well as more broadly across both angiosperms and gymnosperms[90,101], indicate a concentration
of drought tolerance within a limited subset of current tree species diversity, raising the alarm
for catastrophic, phylogenetically-biased biodiversity loss as we pass critical thresholds of
global aridification[102].

Encouragingly, strong phylogenetic correlations between functional traits and environmental
variables also highlight opportunities for imputation of scarce physiological data[103,104], vastly
expanding our capacity to assess species hydraulic risk[89,105]. Using MR-PMM on a global plant
dataset, Sanchez-Martinez et al.[103] demonstrate improved prediction of leaf hydraulic traits by
leveraging phylogenetic correlations with leaf economic traits. As trait-trait and trait-
environment relationships are often tighter within phylogenetically constrained species
groups[19,27,60], such correlations may be particularly valuable for predicting hydraulic traits within
speciose, phylogenetically structured clades, such as *Eucalyptus*. This approach has
enormous potential for parameterizing species-specific hydraulic models of drought-induced
tree mortality[106], aiding the assessment of species vulnerability and ecosystem resilience under
climate change scenarios. A clear direction for future work is therefore to clarify the strength of
relationships between hard (physiological) and soft (morpho-anatomical) traits both within and
between prominent woody clades, so that phylogenetic information can be leveraged most
effectively at a global scale. Discovery and refinement of soft traits that correspond strongly to
physiological thresholds of water stress tolerance (i.e., cavitation resistance and hydraulic
safety margins), while being cost effective to observe and scalable across diverse taxa (e.g.,
xylem area per unit leaf area (XLA)[70]), is a particularly valuable avenue for future research.

Adaptation to aridity in *Eucalyptus* has proceeded via multivariate shifts along a whole-plant
'fast-slow' economic spectrum, prioritizing hydraulic safety and photosynthetic WUE, over
resource acquisition and rate of return on tissue investment. These findings support a general
trade-off between productivity and persistence in plant strategy[26], and link variation along this
trait spectrum to multiple independent transitions in aridity tolerance in a major tree genus.
Strong conservatism in functional traits and environmental niche variables of woody lineages
worldwide suggests that species responses to climate change will be phylogenetically biased,
putting entire clades at risk of rapid decline and threatening to disrupt the structure and
function of ecosystems. Out of this adversity comes a valuable opportunity to leverage the
phylogenetic revolution to better predict species responses to environmental change.

Methods

Trait Data

We compiled all available observations of leaf mass per area (LMA), leaf area, leaf nitrogen per area (N_{area}), leaf $\delta^{13}C$, maximum potential height (max height), and wood density for *Eucalyptus* species in the AusTraits[46] database (figure S5). We augmented this dataset by 1) adding records from observations made by authors in the field (N = 1321) and at Currency Creek Arboretum, South Australia (N = 2107) between 2022-2024; 2) adding records from recent publications[60,70,107] yet to be incorporated into AusTraits; and 3) deriving trait values when missing observations could be calculated from available trait data, e.g., calculating N_{area} from N_{mass} and LMA, or leaf area from leaf length and width measurements (see below). The combined dataset contained 22,650 observations on our 6 focal traits across 767 eucalypt species. Total sample sizes for each trait, and the proportion of eucalypt species for which data were available for each trait are shown in table 1.

Data cleaning and taxonomic assignment

Details of data cleaning and exclusion are summarised in figure S5. We excluded records derived from experimental studies and those reporting observations on juvenile or immature specimens. We included records reporting raw trait values (i.e., observations on an individual specimen) as well as species mean trait values estimated from multiple observations within- or between-sites. We restricted data on height to records of maximum species height, taking the maximum value if multiple values were reported for a given species.

In AusTraits, leaf dimension measurements (length and width) were available for many more *Eucalyptus* species than leaf area measurements (96% vs 68% of species). To leverage these data in our study, we used leaf length and width measurements for each species to calculate leaf area as,

leaf length x leaf width x CF,

where CF = 0.68 is the correction factor (CF) applicable to *Myrtaceae* based on Shrader et al. (2021). To confirm the accuracy of this method, we regressed observed leaf area values against those calculated from length and width measurements (where both were available) and confirmed a tight linear relationship (Pearsons's correlation coefficient = 0.97).

Climate and Soil Data

To separate the effects of local environmental conditions, phylogenetic history, and study effects on functional trait variation, we integrated climate (mean annual temperature, mean moisture index) and soil (phosphorus content, bulk density) data (table 1) observed at two different spatial scales (species mean values and site-level values) into a single model.

For species mean values, climate and soil characteristics of each species range were extracted by intersecting point locations associated with digitised herbarium collections (latitude and longitude) with long-term gridded datasets of climate and modelled soil variables and extracting values. Climate data was sourced from the CHELSA dataset[109] where current climate is represented by average conditions between 1979-2013, and soil data from the National Soil

and Landscape Grid of Australia (https://esoil.io/TERNLandscapes/Public/Pages/SLGA/). This
provided a distribution of values across raster cells for each environmental variable for each
species, from which we calculated species mean values, representing the average climatic and
soil conditions experienced by each species across their geographic range. These species
mean environmental variables (values constant within-species) were then fit alongside
functional trait variables (values variable within-species) in a multi-response (MR) phylogenetic
mixed model (PMM), allowing us to decompose trait-environment correlations across different
levels in the model hierarchy.

For site-level environmental data, we used the procedure described above to extract point-
location climate and soil values for each trait observation reporting latitude and longitude of the
collection site (N = 4120). For each observation, we then subtracted the relevant species mean
environmental niche values from the site-level environment values, yielding a set of variables
expressing the differences between site-level environmental conditions and species mean
environmental niche values, i.e., environmental anomalies. Records for which site-level data
were unavailable (N = 6872) were given a value of 0 for these anomaly variables and were thus
assumed to have been collected at sites representing species mean environmental conditions.
By fitting species-specific environmental anomalies as fixed effects in the MR-PMM, this
approach allowed us to separate and test the influence of site-level environmental conditions
on functional trait variation within-species, when estimating correlations between niche
characteristics and functional traits between-species.

Phylogeny

Hybridisation and introgression among closely related species complicate the phylogenetic
inference of *Eucalyptus*[110,111], leading to difficulties in confidently assigning sister species
relationships. To account for the potential impacts of this phylogenetic uncertainty, we
integrated the latest phylogenetic evidence[44] and taxonomic revisions[45,112] to augment the
largest time-calibrated phylogeny currently available for *Eucalyptus*[35]. Then, we used this
augmented tree to model phylogenetic structure at two different taxonomic scales (species-
level and series-level). This approach allowed us to maximise the number of species and trait
observations included in analyses, while explicitly incorporating phylogenetic uncertainty into
parameter estimates. For all analyses, we follow the recent taxonomic revision proposed by
Nicolle et al. (2024a; 2024b), in which all previously recognised eucalypt genera (i.e.,
Eucalyptus, *Corymbia*, *Angophora* and *Blakella*) are treated as subgenera of a unified genus
Eucalyptus.

First, we used the recent phylogeny of Crisp et al. (2024) to infer a strict consensus tree of
Eucalyptus at the taxonomic section level. We began by generating 1000 candidate section-
level topologies from each of the two maximum likelihood partition trees presented in Crisp et
al. (2024) by randomly sampling one species from each taxonomic section and pruning the
respective tree to those taxa. We combined these 2000 candidate section-level topologies into
a single set to infer a strict consensus tree. We then augmented the "ML1" phylogeny presented
in Thornhill et al. (2019) to match this backbone topology by 1) removing taxa that were
discordant with this consensus tree, 2) re-locating subgenus *Eudesmia* to be sister to

subgenera *Eucalyptus* and *Symphyomyrtus*, in line with Crisp et al. (2024); and 3) excluding taxa
identified as inter-sectional or inter-series hybrids according to Nicolle et al. (2024).

Ancestral state reconstructions require species-level topology to leverage information on trait
differences between closely related species that have nonetheless diverged in their climatic
preferences. However, for inferring and decomposing correlations between traits, it may be
preferable to collapse the phylogeny to a higher taxonomic rank to include as much data as
possible in analyses (see below). We therefore used our augmented species-level tree to
further generate candidate series-level topologies, by sampling one species from each
taxonomic series and pruning the tree to those taxa. The resulting trees represent hypotheses
of phylogenetic relationships between *Eucalyptus* taxonomic series, rather than species,
providing several advantages for downstream analyses. In particular, by treating species as
replicates of a taxonomic series we: 1) avoid strict assumptions about sister-species
relationships that may be complicated by hybridisation and introgression; 2) include data for all
species with resolved series-level taxonomy, including those that do not feature in the species-
level phylogeny; 3) are able to directly propagate phylogenetic uncertainty into our analyses by
fitting models over a sample of candidate topologies and combining posterior estimates for
inference.

Statistical Analyses

Multi-Response Phylogenetic Mixed Models (MR-PMM)

We used Bayesian MR-PMM to decompose correlations between species functional traits and
environmental niche variables into phylogenetic and non-phylogenetic components[48].
Specifically, we fit all functional traits and environmental niche characteristics jointly as
response variables, removing the global intercept to estimate separate intercepts for each
response variable, and using correlated random effects to specify phylogenetic and non-
phylogenetic between-species covariance matrices. We modelled multi-level structure in the
data by fitting random intercepts across species (accounting for within-species replication) and
studies (accounting for study effects). Retaining observation-level data allowed us to estimate
within-species variances for each functional trait, allowing us to more accurately estimate
phylogenetic and non-phylogenetic trait (co)variances[113]. All MR-PMM were fit using the
MCMCglmm package[114] in R.

Phylogenetic imputation

Traits varied in the extent of missing data, ranging from 50% missingness at the species level for
wood density to 1% missingness for leaf area and max height (table 1). Missing response values
are permitted in MCMCglmm, with missing values imputed conditional on the full phylogenetic
covariance structure of the model. In our case, this means that both phylogenetic and non-
phylogenetic between-species trait correlations inform the imputation of missing values. One
benefit of this approach over multiple imputation procedures is that imputation uncertainty is
naturally propagated through to the posterior distribution of parameter estimates. We used the
gap-filled dataset predicted from the MR-PMM fit to conduct PCA (figure 1) and visualise the
phylogenetic distribution of functional trait values (figure 2).

Model Settings

For all analyses based on series-level phylogeny (see above), we incorporated phylogenetic
uncertainty by fitting models over a sample of 100 candidate topologies, and pooling MCMC
samples across fits prior to inference. We assessed convergence by 1) visually inspecting
traces of the MCMC posterior estimates; and 2) confirming potential scale reduction factors ($\hat{r}$),
a convergence diagnostic test that compares within- and between-chain variance[115], were
≤1.01 for all parameter estimates. Estimates for all variance components converged
successfully using parameter expanded priors with (variance) V = I$_m$ (an identity matrix of
dimension equal to the number of response traits, k), and (degree of belief) v = 1k+1 for random
effects[116], nitts = 11000, burn-in = 1000, and thin = 10. We used the default independent normal
priors with (mean) = 0 and V = 10^{10} for fixed effects. We performed model validation via
predictive assessment using approximate leave-one-out (LOO) cross-validation (CV)
procedures[117,118] (figure S8) and posterior predictive checks (figure S9).

Parameter estimates from models are reported as posterior means and 95% credible intervals
(table S4). For hypothesis tests of the significance of effects estimated by MR-PMM (see below),
we checked whether the 95% CI of the posterior distribution of estimates crossed zero.

Phylogenetic signal

We calculated posterior samples of the phylogenetic signal (λ) in each trait and environmental
factor from the fitted MR-PMM as

$$
624 \qquad \lambda_i \; = \; \frac{\Sigma_{ii}^{\text{phy}}}{\Sigma_{ii}^{\text{phy}} + \Sigma_{ii}^{\text{non}} + \Sigma_{ii}^{\text{stu}} + \Sigma_{ii}^{\text{res}}} \, , \qquad\qquad (1)
$$

where $\Sigma_{ii}^{\text{phy}}, \Sigma_{ii}^{\text{non}}, \Sigma_{ii}^{\text{stu}}$ and Σ_{ii}^{res} are the estimated phylogenetic (between-species), non-
phylogenetic (between-species), study and residual variances for trait i, respectively.

Correlation estimates

Based on the MR-PMM decomposition, we estimated correlations between traits, and between
traits and environmental factors on both the phylogenetic and non-phylogenetic level. We
calculated the phylogenetic correlation between two response variables as

$$
631 \qquad \rho_{ij}^{\text{phy}} = \frac{\Sigma_{ij}^{\text{phy}}}{\sqrt{\Sigma_{ii}^{\text{phy}}\Sigma_{jj}^{\text{phy}}}} \, , \qquad\qquad (2)
$$

Where ρ_{ij}^{phy} is the phylogenetic correlation between traits i and j, Σ_{ij}^{phy} is the phylogenetic
covariance between traits i and j, and Σ_{ii}^{phy} is the phylogenetic variance in trait i. Non-
phylogenetic between-species correlations were calculated similarly by substituting elements
of $\mathbf{\Sigma}^{\text{ind}}$ into eq. 2.

As for interpretation, phylogenetic correlations represent correlations between traits that are
conserved over evolutionary time, i.e., the tendency for traits to change together (and be
conserved) during the divergence of phylogenetic clades. Non-phylogenetic correlations
represent correlations between traits that are independent of phylogenetic structure, i.e., the
tendency for traits to covary after accounting for conserved relationships. Thus, by comparing

phylogenetic and residual correlations, we can evaluate the extent to which trait (co)variation
can be explained by conserved differences between clades versus differences between
individual species that are independent of phylogenetic affiliation.

Partial Correlation Estimates

To identify candidate causal relationships between response variables, we calculated partial
correlations and associated credible intervals for all pairs of response variables from the
precision of the phylogenetic trait covariance matrix[48,119]. The precision matrix (inverse of the
covariance matrix) estimates the undirected conditional dependency among a set of variables,
from which we can derive partial correlations; pairwise relationships between variables after
accounting for covariances with all other variables[120,121]. For example, if we obtain data on three
covarying species traits x, y and z, the conditional dependency between x and y is the
component of covariance between these traits that cannot be explained by z. If we assume no
important omitted variables, significant partial correlations are consistent with direct
relationships between traits, as opposed to relationships manifest from cross-correlation with
other variables. We used pcor.shrink() from the corpcor[122] R package to compute partial
correlations using the default regularisation settings. By conditioning on all other response
variables, (non-zero) partial correlations between species traits provide evidence for direct
functional or mechanistic associations, rather than correlations arising from trait responses to
the environment[48]. Similarly, partial correlations between traits and environmental factors
provide evidence for direct associations between trait values and the ability to persist and
compete across environmental gradients, rather than correlations due to relationships with
other traits.

Ancestral State Reconstruction

Given strong phylogenetic signal in all traits and environmental factors (figure 3B), we used
ancestral state reconstruction to evaluate whether changes in trait values represent causes or
consequences of transitions to arid environments, i.e., to assess hypotheses relating to the
sequence of evolutionary events. We did this using a two-step process following the approach
of Cornwallis et al. (2017).

For step one, we used an MR-PMM fit to the species-level phylogeny to estimate ancestral
states for moisture index (MI) at all internal nodes. These estimates represent the expected MI
of ancestral species, after accounting for model multilevel structure, according to a Brownian
motion model of evolution. We then adjusted these Brownian ancestral state estimates
according to a shift model, which assumes that all ancestral *Eucalyptus* lineages occupied
humid (MI > 0.65) niches prior to the Miocene boundary (23 Mya), before experiencing a
directional trend toward more arid niches into the present day (figure S6). Specifically, we
adjusted ancestral states of internal nodes prior to the Miocene boundary (node depth > 23
Mya) by an amount equal to the difference between the root state estimated by Brownian
motion (MI = 0.46) and the MI defining humid environments (MI = 0.65). We adjusted ancestral
states of internal nodes following the Miocene boundary (node depth < 23 Mya) by an amount
equal to (23 – D)*R, where D is the node depth and R is the rate of directional change implied by
the supplied root state. This procedure yielded a set of ancestral state estimates for MI that
represent Brownian evolution from a humid root state (MI = 0.65) up until 23 Mya, followed by a

directional trend toward lower MI values between 23 Mya-present (figure S6). This scenario
more accurately reflects the paleoclimatic history of Australia, which is characterised by
progressive aridification since the Miocene, due to northward drift of the Australian continent
and establishment of the Antarctic Circumpolar Current[36,37]. Nonetheless, sensitivity analyses
confirmed that results from contrasts (see below) were similar whether we applied the shift
model or simple Brownian motion (figure S10).

For step two, we used these reconstructed MI values to compare reconstructed functional trait
values between parent nodes that did and did not give rise to more arid descendants (figure S7).
Specifically, for each posterior sample of ancestral MI estimates, we first classified each
internal node as mesic (M) if the estimated MI > 0.5, semi-arid (S) if the estimated MI was 0.5 > X
> 0.2, or arid (A) if the estimated MI < 0.2. We then re-classified all internal nodes by adding
information about the state of their descendant nodes. For example, a parent node was
classified as M_M if it was M and gave rise only to M descendants (child nodes); M_S if it was M
and gave rise to at least one S descendant (representing an origin, or transition, from mesic to
semi-arid environments); S_S if it was S and gave rise only to S descendants; and S_M if it was S
and gave rise to at least one M descendant (representing a reversal, or transition, to mesic
environments). We used this procedure to classify all internal nodes in the phylogeny. We then
fit an MR-PMM taking only functional traits as response variables, specifying a trait-level fixed
effect of node classification (possible levels: M_M, M_S, M_A, S_S, S_A, S_M, A_A, A_S, A_M),
and a phylogenetic random effect linked to ancestral nodes, to estimate the influence of node
class on reconstructed trait values. Importantly, the model was fit only to trait data at the tips,
which treats tips as replicate observations (descendants) of parental nodes when estimating
the influence of node classification on reconstructed trait values. This approach means that
uncertainty about reconstructed trait values at more ancient evolutionary divergences is
naturally propagated through to the fixed effect estimates of node classification.

To test whether reconstructed trait values differed for nodes representing **origins** of transitions
to more arid environments (i.e., nodes directly preceding a transition to a more arid
environment, i.e., M_S and S_A) consistent with environmental filtering, we constructed
contrasts from the posterior samples of the fitted model by subtracting the relevant posterior
distribution of effect sizes for each transition type. Specifically, by subtracting fixed effect
estimates associated with no change in aridity state (e.g., M_M) from those associated with
transition to a more arid state among descendants (e.g., M_S), and checking whether the 95%
CI of the resulting distribution of contrast values crossed zero. These contrasts allowed us to
test for differences in reconstructed trait values associated with the initial transition to more
arid environments. We then used this technique to create additional contrasts that tested for
changes in trait values associated with **persistence** in more arid environments (S_S – M_M and
A_A – S_S) consistent with adaptation *in situ*, and filtering during **reversals** to less arid
environments (S_M – S_S, and A_S – A_A). Parameter uncertainty from the MR-PMM fit was
propagated into ancestral state reconstructions by classifying nodes and calculating contrasts
for each posterior sample.

Acknowledgments

We thank Brad M. Potts for helpful discussions regarding eucalypt phylogenetics. We thank Vinod Jacob and Emma E. Sumner for assistance with field data collection. This work was funded by The Australian Research Council Centre of Excellence for Plant Success in Nature and Agriculture (CE200100015).

Data Accessibility

Data and R code sufficient to replicate all analyses are available at
https://github.com/Benjamin-Halliwell/Eucalyptus_Replicated_Evolution

References

Brienen, R. J. *et al.* Long-term decline of the Amazon carbon sink. *Nature* **519**, 344-348 (2015). https://doi.org:10.1038/nature14283

van Mantgem, P. J. *et al.* Widespread increase of tree mortality rates in the western United States. *Science* **323**, 521-524 (2009). https://doi.org:10.1126/science.1165000

Senf, C., Buras, A., Zang, C. S., Rammig, A. & Seidl, R. Excess forest mortality is consistently linked to drought across Europe. *Nat Commun* **11**, 6200 (2020). https://doi.org:10.1038/s41467-020-19924-1

Nolan, R. H. *et al.* Hydraulic failure and tree size linked with canopy die-back in eucalypt forest during extreme drought. *New Phytol* **230**, 1354-1365 (2021). https://doi.org:10.1111/nph.17298

Lewis, S. L., Brando, P. M., Phillips, O. L., van der Heijden, G. M. & Nepstad, D. The 2010 Amazon drought. *Science* **331**, 554 (2011). https://doi.org:10.1126/science.1200807

Breshears, D. D. *et al.* Regional vegetation die-off in response to global-change-type drought. *Proc Natl Acad Sci U S A* **102**, 15144-15148 (2005). https://doi.org:10.1073/pnas.0505734102

McDowell, N. G. *et al.* Mechanisms of woody-plant mortality under rising drought, CO2 and vapour pressure deficit. *Nat Rev Earth Env* **3**, 294-308 (2022). https://doi.org:10.1038/s43017-022-00272-1

Brodribb, T. J., Powers, J., Cochard, H. & Choat, B. Hanging by a thread? Forests and drought. *Science* **368**, 261-266 (2020).

Li, J. & Prentice, I. C. Global patterns of plant functional traits and their relationships to climate. *Commun Biol* **7**, 1136 (2024). https://doi.org:10.1038/s42003-024-06777-3

Madani, N. *et al.* Future global productivity will be affected by plant trait response to climate. *Sci Rep* **8**, 2870 (2018). https://doi.org:10.1038/s41598-018-21172-9

Dawson, T. P., Jackson, S. T., House, J. I., Prentice, I. C. & Mace, G. M. Beyond predictions: biodiversity conservation in a changing climate. *Science* **332**, 53-58 (2011). https://doi.org:10.1126/science.1200303

Adams, H. D. *et al.* Temperature response surfaces for mortality risk of tree species with future drought. *Environmental Research Letters* **12** (2017). https://doi.org:10.1088/1748-9326/aa93be

McDowell, N. G. *et al.* Multi-scale predictions of massive conifer mortality due to chronic temperature rise. *Nature Climate Change* **6**, 295-300 (2015). https://doi.org:10.1038/nclimate2873

Joswig, J. S. *et al.* Climatic and soil factors explain the two-dimensional spectrum of global plant trait variation. *Nat Ecol Evol* **6**, 36-50 (2022). https://doi.org:10.1038/s41559-021-01616-8

Diaz, S. *et al.* The global spectrum of plant form and function. *Nature* **529**, 167-171 (2016). https://doi.org:10.1038/nature16489

Chave, J. *et al.* Towards a worldwide wood economics spectrum. *Ecol Lett* **12**, 351-366 (2009). https://doi.org:10.1111/j.1461-0248.2009.01285.x

Wright, I. J. *et al.* The worldwide leaf economics spectrum. *Nature* **428**, 821-827 (2004).

King, D. A., Davies, S. J., Tan, S. & Noor, N. S. M. The role of wood density and stem support costs in the growth and mortality of tropical trees. *Journal of Ecology* **94**, 670-680 (2006). https://doi.org:10.1111/j.1365-2745.2006.01112.x

Gleason, S. M. *et al.* Weak tradeoff between xylem safety and xylem-specific hydraulic efficiency across the world's woody plant species. *New Phytol* **209**, 123-136 (2016). https://doi.org:10.1111/nph.13646

20   Hacke, U. G., Sperry, J. S., Wheeler, J. K. & Castro, L. Scaling of angiosperm xylem structure
with safety and efficiency. *Tree Physiol* **26**, 689-701 (2006).
https://doi.org:10.1093/treephys/26.6.689

21   McGill, B. J., Enquist, B. J., Weiher, E. & Westoby, M. Rebuilding community ecology from
functional traits. *Trends Ecol Evol* **21**, 178-185 (2006).
https://doi.org:10.1016/j.tree.2006.02.002

22   Westoby, M. & Wright, I. J. Land-plant ecology on the basis of functional traits. *Trends Ecol*
Evol **21**, 261-268 (2006). https://doi.org:10.1016/j.tree.2006.02.004

23   Jin, Y. *et al.* Aridity-dependent sequence of water potentials for stomatal closure and
hydraulic dysfunctions in woody plants. *Glob Chang Biol* **29**, 2030-2040 (2023).
https://doi.org:10.1111/gcb.16605

24   Wang, X. *et al.* Plant trait networks reveal adaptation strategies in the drylands of China.
BMC Plant Biol **23**, 266 (2023). https://doi.org:10.1186/s12870-023-04273-0

25   Flores-Moreno, H. *et al.* Robustness of trait connections across environmental gradients and
growth forms. *Global Ecology and Biogeography* **28**, 1806-1826 (2019).
https://doi.org:10.1111/geb.12996

26   Reich, P. B. & Cornelissen, H. The world-wide 'fast–slow' plant economics spectrum: a traits
manifesto. *Journal of Ecology* **102**, 275-301 (2014). https://doi.org:10.1111/1365-
2745.12211

27   Anderegg, L. D. L. Why can't we predict traits from the environment? *New Phytol* **237**, 1998-
2004 (2023). https://doi.org:10.1111/nph.18586

28   Andrew, S. C. *et al.* Functional diversity of the Australian flora: Strong links to species
richness and climate. *Journal of Vegetation Science* **32** (2021).
https://doi.org:10.1111/jvs.13018

29   Šímová, I. *et al.* Spatial patterns and climate relationships of major plant traits in the New
World differ between woody and herbaceous species. *Journal of Biogeography* **45**, 895-916
(2018). https://doi.org:10.1111/jbi.13171

30   Li, D., Ives, A. R. & Waller, D. M. Can functional traits account for phylogenetic signal in
community composition? *New Phytol* **214**, 607-618 (2017).
https://doi.org:10.1111/nph.14397

31   Wright, I. J. *et al.* Assessing the generality of global leaf trait relationships. *New Phytol* **166**,
485-496 (2005). https://doi.org:10.1111/j.1469-8137.2005.01349.x

32   Butler, E. E. *et al.* Increasing Functional Diversity in a Global Land Surface Model Illustrates
Uncertainties Related to Parameter Simplification. *Journal of Geophysical Research:*
Biogeosciences **127** (2022). https://doi.org:10.1029/2021jg006606

33   Anderegg, L. D. L. *et al.* Representing plant diversity in land models: An evolutionary
approach to make "Functional Types" more functional. *Glob Chang Biol* **28**, 2541-2554
(2022). https://doi.org:10.1111/gcb.16040

34   Yates, K. L. *et al.* Outstanding Challenges in the Transferability of Ecological Models. *Trends*
Ecol Evol **33**, 790-802 (2018). https://doi.org:10.1016/j.tree.2018.08.001

35   Thornhill, A. H. *et al.* A dated molecular perspective of eucalypt taxonomy, evolution and
diversification. *Australian Systematic Botany* **32**, 29-48 (2019).

36   Fujioka, T. & Chappell, J. History of Australian aridity: chronology in the evolution of arid
landscapes. *Geological Society, London, Special Publications* **346**, 121-139 (2010).
https://doi.org:10.1144/sp346.8

37   Martin, H. A. Cenozoic climatic change and the development of the arid vegetation in
Australia. *Journal of Arid Environments* **66**, 533-563 (2006).
https://doi.org:10.1016/j.jaridenv.2006.01.009

38   James, M. E., Brodribb, T., Wright, I. J., Rieseberg, L. H. & Ortiz-Barrientos, D. Replicated
Evolution in Plants. *Annu Rev Plant Biol* **74**, 697-725 (2023).
https://doi.org:10.1146/annurev-arplant-071221-090809

39   Bendall, E. R. *et al.* Demographic change and loss of big trees in resprouting eucalypt forests
exposed to megadisturbance. *Global Ecology and Biogeography* **33** (2024).
https://doi.org:10.1111/geb.13842

40   Keppel, G. *et al.* Population decline in a Pleistocene refugium: Stepwise, drought-related
dieback of a South Australian eucalypt. *Sci Total Environ* **876**, 162697 (2023).
https://doi.org:10.1016/j.scitotenv.2023.162697

41   Matusick, G. *et al.* Chronic historical drought legacy exacerbates tree mortality and crown
dieback during acute heatwave-compounded drought. *Environmental Research Letters* **13**
(2018). https://doi.org:10.1088/1748-9326/aad8cb

42   Boer, M. M., de Dios, V. R. & Bradstock, R. A. Unprecedented burn area of Australian mega
forest fires. *Nature Climate Change* **10**, 171-172 (2020). https://doi.org:10.1038/s41558-
020-0716-1

43   Nolan, R. H. *et al.* Causes and consequences of eastern Australia's 2019-20 season of mega-
fires. *Glob Chang Biol* **26**, 1039-1041 (2020). https://doi.org:10.1111/gcb.14987

44   Crisp, M. D. *et al.* Perianth evolution and implications for generic delimitation in the
eucalypts (Myrtaceae), including the description of the new genus, Blakella. *Journal of*
Systematics and Evolution **62**, 942-962 (2024). https://doi.org:10.1111/jse.13047

45   Nicolle, D. Classification of the eucalypts, genus Eucalyptus. (2024).

46   Falster, D. *et al.* AusTraits, a curated plant trait database for the Australian flora. *Scientific*
Data **8**, 1-20 (2021).

47   Westoby, M., Yates, L., Holland, B. & Halliwell, B. Phylogenetically conservative trait
correlation: Quantification and interpretation. *Journal of Ecology* **111**, 2105-2117 (2023).
https://doi.org:10.1111/1365-2745.14150

48   Halliwell, B., Holland, B. R., Yates, L. A. Multi-response phylogenetic mixed models: concepts
and application. *Biological Reviews* (in press).

49   Kraft, N. J. B. *et al.* Community assembly, coexistence and the environmental filtering
metaphor. *Functional Ecology* **29**, 592-599 (2015). https://doi.org:10.1111/1365-2435.12345

50   Christmas, M. J., Breed, M. F. & Lowe, A. J. Constraints to and conservation implications for
climate change adaptation in plants. *Conservation Genetics* **17**, 305-320 (2015).
https://doi.org:10.1007/s10592-015-0782-5

51   Pagel, M. Inferring the historical patterns of biological evolution. *Nature* **401**, 877-884
(1999).

52   Prentice, I. C., Dong, N., Gleason, S. M., Maire, V. & Wright, I. J. Balancing the costs of carbon
gain and water transport: testing a new theoretical framework for plant functional ecology.
Ecol Lett **17**, 82-91 (2014). https://doi.org:10.1111/ele.12211

53   Wright, I. J., Reich, P. B. & Westoby, M. Least-cost input mixtures of water and nitrogen for
photosynthesis. *Am Nat* **161**, 98-111 (2003). https://doi.org:10.1086/344920

54   Mencuccini, M. *et al.* Leaf economics and plant hydraulics drive leaf : wood area ratios. *New*
Phytol **224**, 1544-1556 (2019). https://doi.org:10.1111/nph.15998

55   Olson, M. E. *et al.* Plant height and hydraulic vulnerability to drought and cold. *Proc Natl*
Acad Sci U S A **115**, 7551-7556 (2018). https://doi.org:10.1073/pnas.1721728115

56   Dong, N. *et al.* Leaf nitrogen from the perspective of optimal plant function. *J Ecol* **110**, 2585-
2602 (2022). https://doi.org:10.1111/1365-2745.13967

57   Dong, N. *et al.* Leaf nitrogen from first principles: field evidence for adaptive variation with
climate. *Biogeosciences* **14**, 481-495 (2017). https://doi.org:10.5194/bg-14-481-2017

58   Onoda, Y. *et al.* Physiological and structural tradeoffs underlying the leaf economics
spectrum. *New Phytol* **214**, 1447-1463 (2017). https://doi.org:10.1111/nph.14496

59   Li, X. *et al.* Tree hydraulic traits are coordinated and strongly linked to climate-of-origin
across a rainfall gradient. *Plant Cell Environ* **41**, 646-660 (2018).
https://doi.org:10.1111/pce.13129

60 Peters, J. M. R. *et al.* Living on the edge: A continental-scale assessment of forest
vulnerability to drought. *Glob Chang Biol* **27**, 3620-3641 (2021).
https://doi.org:10.1111/gcb.15641
61 Barotto, A. J. *et al.* Wood density and anatomy of three Eucalyptus species: implications for
hydraulic conductivity. *Forest Systems* **26** (2017). https://doi.org:10.5424/fs/2017261-10446
62 Smith, D. D. *et al.* Ecophysiological adaptations shape distributions of closely related trees
along a climatic moisture gradient. *Nat Commun* **14**, 7173 (2023).
https://doi.org:10.1038/s41467-023-42352-w
63 Hellmann, J. J. & Pineda-Krch, M. Constraints and reinforcement on adaptation under
climate change: Selection of genetically correlated traits. *Biological Conservation* **137**, 599-
609 (2007). https://doi.org:10.1016/j.biocon.2007.03.018
64 Liang, X., Ye, Q., Liu, H. & Brodribb, T. J. Wood density predicts mortality threshold for
diverse trees. *New Phytol* **229**, 3053-3057 (2021). https://doi.org:10.1111/nph.17117
65 Stovall, A. E. L., Shugart, H. & Yang, X. Tree height explains mortality risk during an intense
drought. *Nat Commun* **10**, 4385 (2019). https://doi.org:10.1038/s41467-019-12380-6
66 Anderegg, W. R. *et al.* Meta-analysis reveals that hydraulic traits explain cross-species
patterns of drought-induced tree mortality across the globe. *Proc Natl Acad Sci U S A* **113**,
5024-5029 (2016). https://doi.org:10.1073/pnas.1525678113
67 Bennett, A. C., McDowell, N. G., Allen, C. D. & Anderson-Teixeira, K. J. Larger trees suffer
most during drought in forests worldwide. *Nat Plants* **1**, 15139 (2015).
https://doi.org:10.1038/nplants.2015.139
68 Hacke, U. G., Sperry, J. S., Pockman, W. T., Davis, S. D. & McCulloh, K. A. Trends in wood
density and structure are linked to prevention of xylem implosion by negative pressure.
Oecologia **126**, 457-461 (2001). https://doi.org:10.1007/s004420100628
69 Pfautsch, S. *et al.* Climate determines vascular traits in the ecologically diverse genus
Eucalyptus. *Ecol Lett* **19**, 240-248 (2016). https://doi.org:10.1111/ele.12559
70 Blackman, C. J., Halliwell, B., Hartill, G. E. & Brodribb, T. J. Petiole XLA (xylem to leaf area
ratio) integrates hydraulic safety and efficiency across a diverse group of eucalypt leaves.
Plant, Cell & Environment **47**, 49-58 (2024).
71 Givnish, T. J., Wong, S. C., Stuart-Williams, H., Holloway-Phillips, M. & Farquhar, G. D.
Determinants of maximum tree height in Eucalyptus species along a rainfall gradient in
Victoria, Australia. *Ecology* **95**, 2991-3007 (2014).
72 Bruelheide, H. *et al.* Global trait-environment relationships of plant communities. *Nat Ecol*
Evol **2**, 1906-1917 (2018). https://doi.org:10.1038/s41559-018-0699-8
73 Farquhar, G. D., Ehleringer, J. R., & Hubick, K. T. Carbon isotope discrimination and
photosynthesis. *Annual review of plant physiology and plant molecular biology* **40**, 503-537
(1989).
74 Cernusak, L. A. *et al.* Environmental and physiological determinants of carbon isotope
discrimination in terrestrial plants. *New Phytol* **200**, 950-965 (2013).
https://doi.org:10.1111/nph.12423
75 Reich, P. B. & Oleksyn, J. Global patterns of plant leaf N and P in relation to temperature and
latitude. *Proc Natl Acad Sci U S A* **101**, 11001-11006 (2004).
https://doi.org:10.1073/pnas.0403588101
76 Cornwell, W. K., Schwilk, D. W. & Ackerly, D. D. A Trait-Based Test for Habitat Filtering:
Convex Hull Volume. *Ecology* **87**, 1465-1471 (2006). https://doi.org:10.1890/0012-
9658(2006)87[1465:Attfhf]2.0.Co;2
77 Fontes, C. G. *et al.* Convergent evolution of tree hydraulic traits in Amazonian habitats:
implications for community assemblage and vulnerability to drought. *New Phytol* **228**, 106-
120 (2020). https://doi.org:10.1111/nph.16675
78 Wiens, J. J. *et al.* Niche conservatism as an emerging principle in ecology and conservation
biology. *Ecol Lett* **13**, 1310-1324 (2010). https://doi.org:10.1111/j.1461-0248.2010.01515.x

79 Li, F. *et al.* Evolutionary history shapes variation of wood density of tree species across the
world. *Plant Divers* **46**, 283-293 (2024). https://doi.org:10.1016/j.pld.2024.04.002
80 Coelho de Souza, F. *et al.* Evolutionary heritage influences Amazon tree ecology. *Proc Biol Sci*
283 (2016). https://doi.org:10.1098/rspb.2016.1587
81 Ackerly, D. D. Community assembly, niche conservatism, and adaptive evolution in changing
environments. *Int J Plant Sci* **164**, S165-S184 (2003). https://doi.org:Doi 10.1086/368401
82 Booth, T. H. *et al.* Native forests and climate change: Lessons from eucalypts. *Forest Ecology*
and Management **347**, 18-29 (2015). https://doi.org:10.1016/j.foreco.2015.03.002
83 Cunningham, C. X. *et al.* Pyrogeography in flux: Reorganization of Australian fire regimes in a
hotter world. *Glob Chang Biol* **30**, e17130 (2024). https://doi.org:10.1111/gcb.17130
84 García-Valdés, R., Bugmann, H., Morin, X. & Zurell, D. Climate change-driven extinctions of
tree species affect forest functioning more than random extinctions. *Diversity and*
Distributions **24**, 906-918 (2018). https://doi.org:10.1111/ddi.12744
85 Huang, J. *et al.* Global semi-arid climate change over last 60 years. *Climate Dynamics* **46**,
1131-1150 (2015). https://doi.org:10.1007/s00382-015-2636-8
86 Penone, C. *et al.* Imputation of missing data in life-history trait datasets: which approach
performs the best? *Methods in Ecology and Evolution* **5**, 961-970 (2014).
https://doi.org:10.1111/2041-210x.12232
87 James, T. D., Salguero-Gomez, R., Jones, O. R., Childs, D. Z. & Beckerman, A. P. Bridging gaps
in demographic analysis with phylogenetic imputation. *Conserv Biol* **35**, 1210-1221 (2021).
https://doi.org:10.1111/cobi.13658
88 Urban, M. C. *et al.* Improving the forecast for biodiversity under climate change. *Science* **353**
(2016). https://doi.org:10.1126/science.aad8466
89 Knighton, J., Sanchez-Martinez, P. & Anderegg, L. A global dataset of tree hydraulic and
structural traits imputed from phylogenetic relationships. *Sci Data* **11**, 1336 (2024).
https://doi.org:10.1038/s41597-024-04254-4
90 Avila-Lovera, E., Winter, K. & Goldsmith, G. R. Evidence for phylogenetic signal and
correlated evolution in plant-water relation traits. *New Phytol* **237**, 392-407 (2023).
https://doi.org:10.1111/nph.18565
91 Yang, X. *et al.* Variations of leaf N and P concentrations in shrubland biomes across northern
China: phylogeny, climate, and soil. *Biogeosciences* **13**, 4429-4438 (2016).
https://doi.org:10.5194/bg-13-4429-2016
92 Mo, L. *et al.* The global distribution and drivers of wood density and their impact on forest
carbon stocks. *Nat Ecol Evol* **8**, 2195-2212 (2024). https://doi.org:10.1038/s41559-024-
02564-9
93 Ladiges, P. Y., Udovicic, F. & Nelson, G. Australian biogeographical connections and the
phylogeny of large genera in the plant family Myrtaceae. *Journal of Biogeography* **30**, 989-
998 (2003). https://doi.org:10.1046/j.1365-2699.2003.00881.x
94 Willis, C. G., Ruhfel, B., Primack, R. B., Miller-Rushing, A. J. & Davis, C. C. Phylogenetic
patterns of species loss in Thoreau's woods are driven by climate change. *Proc Natl Acad Sci*
U S A **105**, 17029-17033 (2008). https://doi.org:10.1073/pnas.0806446105
95 Li, D., Miller, J. E. D. & Harrison, S. Climate drives loss of phylogenetic diversity in a grassland
community. *Proc Natl Acad Sci U S A* **116**, 19989-19994 (2019).
https://doi.org:10.1073/pnas.1912247116
96 van Tiel, N. *et al.* Regional uniqueness of tree species composition and response to forest
loss and climate change. *Nat Commun* **15**, 4375 (2024). https://doi.org:10.1038/s41467-024-
48276-3
97 Skelton, R. P. *et al.* Evolutionary relationships between drought-related traits and climate
shape large hydraulic safety margins in western North American oaks. *Proc Natl Acad Sci U S*
A **118** (2021). https://doi.org:10.1073/pnas.2008987118

98   Willson, C. J., Manos, P. S. & Jackson, R. B. Hydraulic traits are influenced by phylogenetic
history in the drought-resistant, invasive genus Juniperus (Cupressaceae). *Am J Bot* **95**, 299-
314 (2008). https://doi.org:10.3732/ajb.95.3.299

99   Liu, H. *et al.* Strong phylogenetic signals and phylogenetic niche conservatism in
ecophysiological traits across divergent lineages of Magnoliaceae. *Sci Rep* **5**, 12246 (2015).
https://doi.org:10.1038/srep12246

100   Liu, H. *et al.* Phylogeny and climate explain contrasting hydraulic traits in different life forms
of 150 woody Fabaceae species. *Journal of Ecology* **112**, 741-754 (2024).
https://doi.org:10.1111/1365-2745.14266

101   Laughlin, D. C. *et al.* Climatic limits of temperate rainforest tree species are explained by
xylem embolism resistance among angiosperms but not among conifers. *New Phytol* **226**,
727-740 (2020). https://doi.org:10.1111/nph.16448

102   Guo, W. Y. *et al.* Climate change and land use threaten global hotspots of phylogenetic
endemism for trees. *Nat Commun* **14**, 6950 (2023). https://doi.org:10.1038/s41467-023-
42671-y

103   Sanchez-Martinez, P. *et al.* A framework to study and predict functional trait syndromes
using phylogenetic and environmental data. *Methods in Ecology and Evolution* **15**, 666-681
(2024).

104 Debastiani, V. J., Bastazini, V. A. G. & Pillar, V. D. Using phylogenetic information to impute
missing functional trait values in ecological databases. *Ecological Informatics* **63**, 101315
(2021).

105 Sanchez-Martinez, P. *et al.* Increased hydraulic risk in assemblages of woody plant species
predicts spatial patterns of drought-induced mortality. *Nat Ecol Evol* **7**, 1620-1632 (2023).
https://doi.org:10.1038/s41559-023-02180-z

106 De Kauwe, M. G. *et al.* Towards species-level forecasts of drought-induced tree mortality
risk. *New Phytol* **235**, 94-110 (2022). https://doi.org:10.1111/nph.18129

107 Portelli, A. M. *et al.* From mallees to mountain ash, specific leaf area is coordinated with
eucalypt tree stature, resprouting, stem construction, and fruit size. *Australian Journal of
Botany* **71**, 506-522 (2023). https://doi.org:10.1071/bt23028

108 Schrader, J. *et al.* Leaf size estimation based on leaf length, width and shape. *Ann Bot* **128**,
395-406 (2021). https://doi.org:10.1093/aob/mcab078

109 Karger, D. N. *et al.* Climatologies at high resolution for the earth's land surface areas. *Sci
Data* **4**, 170122 (2017). https://doi.org:10.1038/sdata.2017.122

110 Jones, R. C., Nicolle, D., Steane, D. A., Vaillancourt, R. E. & Potts, B. M. High density, genome-
wide markers and intra-specific replication yield an unprecedented phylogenetic
reconstruction of a globally significant, speciose lineage of Eucalyptus. *Mol Phylogenet Evol*
105, 63-85 (2016). https://doi.org:10.1016/j.ympev.2016.08.009

111 Rutherford, S., Wilson, P. G., Rossetto, M. & Bonser, S. P. Phylogenomics of the green ash
eucalypts (Myrtaceae): a tale of reticulate evolution and misidentification. *Australian
Systematic Botany* **28** (2015). https://doi.org:10.1071/sb15038

112 Nicolle, D. *et al.* The genus problem – Eucalyptus as a model system for minimising
taxonomic disruption. *Taxon* (2024). https://doi.org:10.1002/tax.13240

113 Cinar, O., Nakagawa, S. & Viechtbauer, W. Phylogenetic multilevel meta-analysis: A
simulation study on the importance of modelling the phylogeny. *Methods in Ecology and
Evolution* **13**, 383-395 (2022).

114 Hadfield, J. D. MCMC methods for multi-response generalized linear mixed models: the
MCMCglmm R package. *Journal of Statistical Software* **33**, 1-22 (2010).

115 Vehtari, A., Gelman, A., Simpson, D., Carpenter, B. & Bürkner, P.-C. Rank-Normalization,
Folding, and Localization: An Improved R̂ for Assessing Convergence of MCMC (with
Discussion). *Bayesian Analysis* **16** (2021). https://doi.org:10.1214/20-ba1221

116   de Villemereuil, P. Quantitative genetic methods depending on the nature of the phenotypic
1033   trait. *Annals of the New York Academy of Sciences* **1422**, 29-47 (2018).
1034   117   Bürkner, P.-C., Gabry, J. & Vehtari, A. Efficient leave-one-out cross-validation for Bayesian
1035   non-factorized normal and Student-t models. *Computational Statistics* **36**, 1243-1261 (2021).
1036   https://doi.org:10.1007/s00180-020-01045-4
1037   118   Vehtari, A., Gelman, A. & Gabry, J. Practical Bayesian model evaluation using leave-one-out
1038   cross-validation and WAIC. *Statistics and Computing* **27**, 1413-1432 (2017).
1039   https://doi.org:10.1007/s11222-016-9696-4
1040   119   Popovic, G. C., Warton, D. I., Thomson, F. J., Hui, F. K. C. & Moles, A. T. Untangling direct
1041   species associations from indirect mediator species effects with graphical models. *Methods
1042   in Ecology and Evolution* **10**, 1571-1583 (2019).
1043   120   Warton, D. I. *Eco-Stats: Data Analysis in Ecology: From t-tests to Multivariate Abundances*.
1044   (Springer Nature, 2022).
1045   121   Ebert-Uphoff, I. & Deng, Y. Causal Discovery for Climate Research Using Graphical Models.
1046   *Journal of Climate* **25**, 5648-5665 (2012). https://doi.org:10.1175/jcli-d-11-00387.1
1047   122   Schafer, J. & Strimmer, K. A shrinkage approach to large-scale covariance matrix estimation
1048   and implications for functional genomics. *Stat Appl Genet Mol Biol* **4**, Article32 (2005).
1049   https://doi.org:10.2202/1544-6115.1175
1050   123   Cornwallis, C. K. *et al.* Cooperation facilitates the colonization of harsh environments. *Nature
1051   Ecology & Evolution* **1**, 1-10 (2017).

1052

Supplementary figures

1053

1054

1055

1056

Figure S1. Phylogenetic distribution of functional trait values and environmental niche variables in *Eucalyptus* subgenus *Symphyomyrtus* using gap-filled data imputed by MR-PMM. Species-level topology (left) coloured by section is plotted against a matrix of species-level means (centre) for each trait and niche variable. Scatterplots (right) show species-level means for each trait and niche variable ordered by moisture niche

Figure S2. Results from MR-PMM fit to species-level phylogeny (N$_{species}$ = 631): **A** Posterior fixed effect estimates of site-level environmental anomalies for each functional trait associated with site-level data. **B** Variance decomposition for each functional trait and environmental niche variable. Traits that are constant for a species are partitioned into phylogenetic and residual components, while traits varying within-species are partitioned into phylogenetic, non-phylogenetic, study, fixed, and residual components. **C** Phylogenetic (lower triangular) and non-phylogenetic (upper triangular) between-species correlations between response variables from MR-PMM. Values represent the posterior mean estimate for each correlation coefficient. **D** Network graphs showing significant (90% and 95% CI) partial phylogenetic correlations (positive = blue edges, negative = red edges) between functional traits (green), climate (light blue) and soil (orange) niche variables. Edge widths are proportional to partial correlation coefficients. All correlation estimates derive from the same MR-PMM model fit but are grouped and plotted in separate networks for illustrative purposes.

1088

1089

1090 **Figure S3.** Posterior distributions of estimated trait values at internal nodes for each transition
1091 category with transition classes coloured as in figure 4. The mean (point), 50% (heavy whisker) and
1092 95% (light whisker) CI of each posterior distribution are shown in black.

1093

Figure S4. Posterior distributions of contrasts between node categories from an MR-PMM ancestral state reconstruction adjusted according to a shift model (see methods). Contrasts are computed and reported for transitions between mesic and semi-arid environments (green), as well as between semi-arid and arid environments (cream). The mean (point), 50% (heavy whisker) and 95% (light whisker) CI of each posterior distribution are shown in black.

1099

1100 **Figure S5**. Flow diagram of data compiled for MR-PMM analyses. Data records from AusTraits[46] were
1101 filtered to focal traits then cleaned by matching species nomenclature with the phylogeny[35] and
1102 excluding records based on the criteria shown. Additional trait observations from targeted field work
1103 by the authors (n=3428) and a recent study[60,70,107] not yet included in AusTraits were added. Finally,
1104 records for superfluous variables were removed.

1105

1106 **Figure S6**. Ancestral state reconstructions of moisture index for *Eucalyptus* species according to **A** a
1107 simple Brownian motion and **B** a shift model in which the ancestral moisture niche of *Eucalyptus* is
1108 assumed to be humid (moisture index > 0.65), with a directional trend toward more arid (lower
1109 moisture index) environments from the Miocene boundary 23 Mya (dotted line).

Figure S7. Simplified diagrammatic example of the method used to define aridity transition categories for internal nodes. In step 1, continuous values of moisture index at the tips (species mean moisture) are used to classify tips into discrete moisture niche categories and reconstruct ancestral states along internal branches. In step 2, reconstructed values are used to categorise internal nodes. In step 3, internal nodes are re-classified according to their own moisture niche state and that of their descendant (child) nodes, defining transition categories that distinguish internal nodes that did and did not give rise to more / less arid descendants. In step 4, reconstructed functional trait values are mapped onto transition categories, facilitating comparisons between transition categories in reconstructed values (figure 4B) as well as contrasts between posterior distributions of fixed effects for different transition categories (figure 4C).

1127

Figure S8. Leave-one-out (LOO) predictive checks for functional traits and environmental niche
variables from MR-PMM. Three-point summaries show the mean (black points) and 0.95 CI (grey
bars) of model predictions for left-out species. Predictions are made conditional on all random
effects, with observations ordered by the predictive mean. Observed species mean values for each
trait and niche variable are superimposed and coloured based on whether the observed value falls
within the predictive interval. Species for which the model had greater predictive uncertainty (wide
grey bars) are generally those with few close relatives in the phylogeny, e.g. monotypic subgenera,
highlighting the importance of phylogeny for predictive performance. Species-level traits
(environmental niche variables and max height) tend to show poorer predictive performance, with
more observations falling outside the predictive interval. For some variables, predictive
performance varies across the range of data values, e.g., poorer predictive performance at lower
values of soil BD.

1140
1141 **Figure S9**. Posterior predictive checks for functional traits and environmental niche variables from
1142 MR-PMM. Posterior predictive checks compare simulated data from the fitted model (blue lines) to
1143 the observed data distribution (black line) to test both the adequacy of the fit to the data and the
1144 plausibility of model predictions.

1145
1146 **Figure S10.** Posterior distributions of contrasts between node categories from an MR-PMM
1147 ancestral state reconstruction based on Brownian Motion. Contrasts are computed and reported for
1148 transitions between mesic and semi-arid environments (green), as well as between semi-arid and
1149 arid environments (cream). The mean (point), 50% (heavy whisker) and 95% (light whisker) CI of
1150 each posterior distribution are shown in black.