Heterocellular and homocellular electrically coupled networks of the thalamus and cortex revealed by focal photomapping

Abbreviated title: Opto-δL maps coupled networks

Authors: Mitchell J. Vaughn[1], Kevin J. Bender[2] and Julie S. Haas[1]*

Affiliations:

1. Department of Biological Sciences, Lehigh University, 111 Research Drive, Bethlehem, PA 18015 USA

2. Weill Institute for Neurosciences and Department of Neurology, University of California, San Francisco, San Francisco, CA, USA

Author Contributions: JSH and KJB designed the method and experiments. MJV optimized the method and performed the electrophysiological research. MJV and JSH wrote and revised the manuscript. JSH supervised and acquired funding.

Correspondence: Julie S. Haas, juliehaas@lehigh.edu

Number of figures: 4 figures

Number of tables: 0 tables

Number of multimedia: 0 multimedia

Number of words for abstract: 179 words

Number of words for Introduction: 621 words

Number of words for Discussion: 816 words

Acknowledgements: We thank all members of the Haas and Bender laboratories for input and discussion.

Conflict of interest: The authors report no conflict of interest

Funding: NIH R01 NS128713, Whitehall Grant in Aid to JSH

Abstract

Electrical synapses are present widely across the mammalian brain and are crucial

components of active neural circuitry and connectomes. Identification of electrically coupled

networks in living tissue has been limited by technical demands of multiplexed recordings, and

no dyes, fluorescent reporters, or genetic labels are currently able to fill the gap. Here, we

introduce a novel method of identifying and measuring electrical synapses: opto-δL combines

focal photostimulation of soma-targeted opsins and a novel method for computing the strength of

electrical synapses, based on spike timing, to rapidly measure and map electrically coupled

networks *in vitro*. We measured electrical synapses and characterized coupled networks in

mature thalamic reticular nucleus (TRN) and in cortex. We used opto-δL to show that TRN

neurons form functional networks that extend as far as 100 μm, and that TRN neurons synapse

promiscuously, coupling both matching and disparate genetic subtypes of neurons. Future

applications of opto-δL will allow identification and characterization of coupled networks across

the brain and enable circuit and systems-level interrogations of the identity and roles of electrical

synapses in circuitry, behavior, and cognition.

Introduction

Chemical and electrical synapses, the twins of the foundational Golgi-Cajal debate, together compose the basis of interneuronal communication across the brain. Neurotransmitter-based chemical synapses underlie communication that is primarily unidirectional and stereotyped in timecourses of activation and decay. In contrast, electrical synapses formed by gap junctions (GJs) relay neuronal signals of any size, polarity, and timecourse bidirectionally between coupled cells, and they do so without the requirement or cost of a spike for signal transduction. Most mature mammalian GJs comprise paired hexameric channels of connexin36[1-3] , which is expressed widely in GABAergic neurons throughout the brain[4]. While chemical synapses and the neuronal networks they form have been described in exquisite detail, descriptions of electrical synaptic transmission and, importantly, their functional networks have been much more limited.

Electrical synapses are increasingly appreciated for the diverse roles they play throughout the brain. They are most widely characterized as mediating synchronization of oscillatory activity in neuronal circuits[5-15]. Beyond synchronization, electrical synapses directly mediate feedforward and lateral excitation, inhibition, reduce signal to noise ratio, sharpen coincidence detection, and facilitate object discrimination[16-32]. GJs create uniquely offset receptive fields in retinal ganglion cells, where they also contribute to tuning responses, enhanced signal saliency, sensitivity, and enhanced coding of visual features[16, 33-36].

Our understanding of the roles electrical synapses and networks play within neuronal networks is sharply limited by the tools currently available. The gold standard for measuring electrical synapses is through paired recordings, in which current is injected into one neuron and a hyperpolarizing deflection can be measured in a coupled neuron[37]; the ratio of deflections indicates synapse strength. While this method is the most direct, it is time-intensive, low-

throughput, and limited to single synapses *in vitro*. Only one experiment has successfully

recorded from a small sample of coupled neurons *in vivo*[38]. Dye-coupling methods[39-46] confound

primary, secondary, or tertiary connections, fail to quantify the strength of synapses, and have

been limited to fixed tissue, although modern variations of dye-coupling have used exogenously

applied molecules that diffuse across a GJ, rendered more specific by limiting entry[47] or

fluorescence[48] to genetically labeled cells. For these reasons, electrical synapses and networks

remain neglected relative to their chemical counterparts in our modern understanding of the

brain.

The thalamic reticular nucleus (TRN) is a nucleus of GABAergic neurons[49] that provide the

majority of inhibition to sensory thalamic relay areas and are hypothesized to focus the

searchlight of attention to the sensory surround[50]. TRN is composed of distinct neuronal subtypes

with spatially and genetically distinct expression[51-53], and our previous paired recordings

indicated that homocellular and heterocellular electrical synapses (coupling same and disparate

cell subtypes) of the TRN[54] may provide the only thalamic link between distinct but parallel

thalamocortical relay channels[51-53, 55]. Because visualization of TRN neurons is impeded by

myelination of thalamocortical fibers after eye opening, all paired-patch experiments

characterizing electrical synapses in TRN have been limited to juveniles[12, 54, 56-62]. While

connexin36 (Cx36) expression persists into adulthood[63] and spikelets have been reported in

adults[64], the possibility that the contributions of GJ networks to thalamocortical relay are limited

to developmental stages remains.

Here, we leverage focal photostimulation of soma-targeted opsins to detect and measure

electrical synapses via changes in spike timing and to map electrically coupled networks in adult

thalamic reticular nucleus (TRN). [49-54, 63, 64] and cortex. Our maps of electrically coupled

networks in adult TRN demonstrate that coupled networks comprise both heterocellular and

homocellular electrical synapses that range from 1 to 4 synapses in size and extend as far as 100

μm of distance. In addition to providing a new and efficient toolset for measuring and mapping

electrical synapses and networks across the living brain, our results provide insight into the

neural circuits that underlie attentional focus on the sensory surround.

Results

Precision of neuronal spike timing is a tightly controlled fundamental process[65] that is

crucial for fidelity of information processing, and is often a parameter of interest in neural

coding[66-70]. Electrical synapses are important contributors in determining spike times[30, 31, 58];

activity in a coupled neighbor can shift spike times of a neuron responding by tens of ms[58].

Leveraging this powerful effect of electrical synapses, we previously compared spike timing

differences due to electrical synapses in two cases using paired recordings: 1) rheobase

stimulation in one neuron alone (t_1) and 2) rheobase stimulation with simultaneous activation of

a coupled neighboring neuron (t_2). We calculated the percentage change in latency of evoked

spikes, δL, as:

108
$$\delta\mathrm{L} = \frac{100 * (t_1 - t_2)}{t_1}$$

A. IR image of a patched pair of TRN juvenile neurons (red and blue circles). Right: voltage responses (V_m) from both neurons during current injection (I_{app}) into cell 2. The coupling coefficient (cc) is the ratio of voltage deflections in both cells; cc = 0.08. Scale bars 100 ms, 2 mV.

$$cc = \frac{\Delta V_{m1}}{\Delta V_{m2}}$$

$$\delta L = \frac{100 * (t_1 - t_2)}{t_1}$$

Figure 1: **Electrical coupling measured by paired patching and opto-δL in the same pair**.
A. IR image of a patched pair of TRN juvenile neurons (red and blue circles). Right: voltage responses (V_m) from both neurons during current injection (I_{app}) into cell 2. The coupling coefficient (cc) is the ratio of voltage deflections in both cells; cc = 0.08. Scale bars 100 ms, 2 mV.
B. Left: Live fluorescence image of the same pair showing st-ChroME-mRuby expression in both neurons. Orange circle indicates area of focal photostimulation of cell 2. Scale bar: 12.5 μm Right: Spiking in neuron 1 for 100 pA applied current (dark blue) alone, and with simultaneous focal photostimulation of neuron 2 (light blue). Dots mark the time of the first spike in each interleaved trial. δL was calculated from the difference in mean spike latencies with (t_2) and without (t_1) photostimulation; δL = 43.4%;
C. Left: Same image as in B. Orange oval marks the area of focal photostimulation of a blank spot that is equidistant from neuron 1. Right: Spiking in neuron 1 for 110 pA applied current alone (dark blue), and with simultaneous photostimulation of the blank spot (lighter blue) from interleaved trials. Dots mark the time of the first spike in each trial. Scale bars 50 ms, 20 mV. For the blank spot used in trial **C**, δL = 16.3%. Differences between δL for trials in **B** and **C** were significant (p < 0.01, Wilcoxon signed-rank test).
D. In two separately patched neurons from adult tissue, increasing amplitudes of current injection into the cell revealed spikelets (arrows) driven by shared excitation of an unrecorded coupled neuron. Scale bars: 25 ms, 10 mV. **E.** Spikelets in a third patched neuron.
F. IR image of the patched neuron from **E** with fluorescence overlay from two depths.
G. Two opto-δL measurements in the patched cells during photostimulation of the cells circled and numbered in
F. Darker traces are responses from photostimulation of equidistant control locations, and light green traces are from photstimulation of the corresponding numbered cell in **G**. We identified that the cell labeled 1 was one of two powerful synapses (δL = 25%) generating the spikelets in **E**.
H. Diagram of the coupled network embedded within all tested neurons from **F**. Scale bar: 25 μm.

This quantity is well-correlated with traditional measures of the strength of electrical

synapses[71]. Here, we combined measurements of δL with focal photostimulation of a soma-

targeted opsin, st-ChroME[72], in potentially coupled nearby neurons, thereby reducing the

requirement for recording an electrical synapse to one patched neuron and simultaneously widely

expanding the number of potential synapses tested. We refer to this method as opto-δL.

To map electrically coupled networks using opto-δL we injected AAV9-CAG-DIO-

ChroME-st-P2A-H2B-mRuby into the TRN of PV-Cre or SOM-Cre mice, and after at least two

weeks, prepared acute brain slices from those mice. We first validated opto-δL in a patched pair

of neurons, with both neurons expressing st-ChroME, in juvenile tissue. We measured the

coupling coefficient for the synapse between these neurons using the traditional small

hyperpolarizing current injections to one of the neurons (Fig. 1A); the coupling coefficient (cc) is

the ratio of voltage deflections in both neurons in response to current injection into one. This pair

had a mean cc of 0.075, close to the average value for coupling between pairs of TRN neurons[73].

Next, we used a digital mirror device (Polygon) to deliver 1-photon focal (34 - 50 μm^2) 560 nm

photostimulation to nearby regions of interest, including the neighboring neuron, during trials of

rheobase-drive spiking in the patched neuron, and computed δL. For all targeted regions of

interest, we used a light intensity sufficient to drive spiking when focused directly on the soma of

a patched neuron (see Methods; Supp. Fig. 1). Because both 1-photon and 2-photon

photostimulation result in light scattering in live tissue[74-76], we measured spike times in cell 1

when photostimulating a blank spot that was equally distanced from cell 1 as the neighboring

neuron (cell 2) as a control for any light scattering-induced excitation of cell 1. Driven by

rheobase current (100 pA) alone, cell 1 spiked 176 ± 3 ms after stimulus onset (Fig. 1B, dark

blue traces). In interleaved trials, we applied rheobase stimulation to cell 1 and focal

photoexcitation of the neighboring neuron; for those trials, cell 1 spiked with a latency of 99 ±

0.3 ms. (Fig. 1B, lighter blue traces), and we computed δL for the synapse: $\delta L = 43.4\%$. To

compared δL for cell 2 to δL driven by scatter excitation to a blank spot, we applied concurrent

photostimulation of the blank spot and injection of rheobase current to cell 1, which resulted in

spikes with latency of 148 ± 3 ms (Fig. 1C, light blue traces) and produced δL of 16.3%. δL was

significantly larger (by 27.1%; $p < 0.01$, Wilcoxon signed-rank test) for photostimulation of the

neighboring cell 2 compared to the blank spot, allowing us to conclude based on spike latencies

that the neighboring cell 2 was coupled to the patched cell 1. These measurements validate the

use of a single electrode and focal optogenetics to identify and characterize an electrical synapse

and provide a foundation for its application to studying electrical synapses and networks in

heavily myelinated adult TRN tissue that precludes paired patching but where spikelets provide

evidence of electrical synapses (Fig. 1D). Indeed, observations of spikelets in adult tissue pointed

us towards identification of electrically coupled neighbors through δL (Fig. 1E-H).

Genetically and anatomically distinct sets of neurons form distinct sets of relay pathways

through the thalamocortical system[51-53, 55]; electrical synapses in the TRN may be the only

thalamic link between these information channels. We previously used paired recordings in

juveniles to test whether TRN electrical synapses interconnect neurons of distinct cell types[54].

Those experiments necessitated the use of crossed Cre reporter mice to label neurons for juvenile

recordings, presenting an additional possible confound of mislabeling neurons due to Cre off-

target recombination [77]. To identify, measure and characterize homocellular and heterocellular

electrically coupled networks in adult TRN, we injected Cre-dependent st-ChroME into adult

mice to genetically identify subsets of neurons (Fig. 2). To map homocellular electrical synapses,

we patched ChroME-expressing mRuby+ neurons in SOM-Cre mice (SOM+), and measured δL

induced by focally photostimulating other nearby mRuby+ neurons (Fig. 2A). To map

heterocellular electrical synapses, we patched mRuby- neurons in SOM-Cre mice (SOM-), and

measured δL induced by focally photostimulating nearby mRuby+ neurons (Fig. 2B). We also

measured homocellular

Figure 2: **Maps of coupled networks between TRN neurons**.
A. Representative homocellular network amongst SOM+ neurons. Overlay of IR and fluorescence images from live recording of st-ChroME-mRuby-expressing neurons in SOM-Cre mouse. Fluorescence is pseudocolored magenta. The patched neuron is circled with white dashes. Scale bar: 50 μm.
B. Diagram of the coupled network embedded within all tested neurons from **A**.
C. Spiking responses used to measure δL and identify network in **A**. Dots mark the times of first spikes in the patched neuron from each trial. Black traces are responses of the patched neuron during rheobase-only trials and colored traces are responses during added target cell photostimulation. Traces are vertically offset for clarity. Each set of traces is numbered by the corresponding cell in A. Scale bars: 5 ms, 20 mV. p values (Wilcoxon signed-rank test) for δLs: 1) < 0.01, 2) 0.70 3) 0.34, 4) < 0.01.
D - F: Representative heterocellular network between SOM- and SOM+ neurons., as described for **A-C**. p values (Wilcoxon signed-rank test) for δL: 0) 1.0, 1) < 0.01 2) 0.56, 3) < 0.01.
G – I: Example homocellular network between PV+ neurons, as described for **A-C**. p values (Wilcoxon signed-rank test) for δL: 1) 0.98, 2) < 0.01 3) <0.01, 4) 0.03.

networks in PV+ neurons (Fig. 2C). To compensate for excitation from scatter in these

experiments, we developed correction procedures for each recorded neuron (Supp. Fig. 3; see

Methods); scatter effects depend on factors including light intensity, opsin expression, and input

resistance, as well as distance but diminished to insignificant levels by 50 - 100 μm (Supp. Fig.

2). We note that mapping heterocellular networks has the advantage of targeting opsin-lacking

neurons for recording, thereby averting the need for compensation in those experiments.

We implemented opto-δL in a population of n = 398 cells comprising 64 possible coupled

networks. Overall, we measured values of δL that were significantly different from 0 within all

combinations of genetic subtypes tested (Fig. 3A). We measured significant values of δL

between SOM+ and other SOM+ neurons in 16 of 35 possible homocellular SOM networks. The

mean coupling rate was 15.1% across all tested connections (n = 206 possible connections), and

40.3% across all tested connections within networks that contained at least one electrical synapse

(n = 67 possible connections). The mean δL between SOM+ neurons was 25.0 ± 3.2%, and the

mean homocellular SOM network contained 1.7 ± 0.1 direct electrical synapses (Fig. 3B).

Between SOM+ and SOM- neurons (Fig. 3C) we measured significant values of δL in 4 of 12

possible heterocellular networks. The mean coupling rate was 7.3% across all tested connections

(n = 85 possible synapses) and 25.8% across all tested connections within networks that

contained at least one electrical synapse (n = 31 possible synapses). The mean δL for SOM+ to

SOM- connections was 16.4 ± 2.4%, and the mean heterocellular network contained was 2.0 ±

0.2 direct electrical synapses (Fig. 3C). For connections between PV+ and PV+ neurons, we

measured significant values of δL in 13 of 17 homocellular networks. The mean coupling rate

was 25.9% across all tested connections (n = 108 possible synapses), and 39.1% across all tested

connections within networks that contained at least one electrical synapse (n = 74 connections).

The mean δL for PV+ to PV+ electrical synapses was 18.9 ± 2.8% (Fig. 3D), and the mean

network size was 2.2 ± 0.1 direct electrical synapses (Fig. 3D). Across networks, there were no

significant differences in δL between subtypes (Fig. 3B-3D, p > 0.05). The mean δL across all

subtypes was 22.8% ± 1.9% (n = 67 synapses), and 50% of synapses had a δL between 11.0%

and 33.3%. The largest δL was 68.4%. We observed several instances in which no nearby opsin-

expressing neuron had any electrical synapses to the recorded neuron. It was rarer for there to be

no electrical synapses between PV neurons (Fig. 3B-D, p < 0.01, chi-squared test). PV neurons

had no electrical synapses to nearby opsin-expressing cells in 23.5% of recorded neurons, while

SOM+ neurons had no connections 54.3% of the time, and SOM- neurons had no connections

66.7% of the time. There was no difference between SOM+ and SOM- neurons in the frequency

of cells that had no nearby electrical synapses (p = 0.07, chi-squared test).

Measuring electrical synapses by patching pairs introduces spatial selection biases to the

pairs that are ultimately tested in experiments; this effect may have contributed to discrepancy

between results from rat TRN suggesting that electrical synapses occur within intersomatic

distances less than 40 μm[73], while dye-coupling results indicate a median intersomatic distance

of 80-120 μm[62]. We used opto-δL to more exhaustively sample the spatial extent of coupling for

distances up to 250 μm. In our data, across all neuron types grouped together, we confirmed that

Figure 3: Distributions of electrical synapses over strength and space for TRN networks.
A. Latencies (+/- SEM) used to compute δL for all δL values determined to be significant.
B. Top: δL for homocellular SOM+ synapses (mean δL = 25.0 ± 3.2%, n = 31 synapses). Bottom: network sizes for SOM+/SOM+ networks (mean size: 1.7 ± 0.1 synapses per map).
C. As in **B,** for heterocellular SOM-/SOM+ synapses (mean δL = 16.4 ± 2.4%, n = 8 synapses; mean size 2.0 ± 0.2 synapses per network).
D. As in **B** for PV+/PV+ synapses (mean δL = 18.9 ± 2.8%, n = 28 synapses; mean size 2.2 ± 0.1 synapses per network, n = 13 maps).
E. Distribution of δL across intersomatic distance for each subtype in TRN (n = 67 synapses). Fit: single exponent.
F. Number of photostimulated neurons that were (black) or were not (grey) coupled to a patched neuron, plotted against distance from the patched neuron (n = 398 neurons).
G. Probability of coupling versus intersomatic distance (n = 398 neurons).

TRN neurons that were closer together were stronger (Fig. 3E) and more likely to be coupled

(Fig. 3F-G), confirming previous reports that δL decreases with intersomatic distance.

Collectively, our data showed that electrically coupled networks for all cell types link neurons as

far as 100 μm apart (Fig. 3F).

To establish whether opto-δL is a viable tool for mapping coupled networks across the brain,

we used opto-δL to identify and map coupled networks in somatosensory cortex using st-

ChroME-mRuby expression to identify possible members of the network. In a SOM-Cre mouse

(Fig. 4A), we identified one network that comprised four neurons, with average δL of 51.4% and

Figure 4: Maps of coupled networks between cortical GABAergic neurons.
A. Network between SOM+ cortical neurons. A) Overlay of IR and fluorescence images of st-ChroME-mRuby-expressing neurons in SOM-Cre mouse. Fluorescence is pseudocolored magenta. The patched neuron is circled with white dashes. Scale bar 50 μm.
B. Diagram of the coupled network embedded within all tested neurons from A.
C. Spiking responses used to measure δL and identify network in A. Dots mark the times of first spikes in the patched neuron from each trial. Black traces are responses during control trials (Iapp only) and colored traces are responses during target cell photostimulation. Traces are vertically offset for clarity. Each set of traces is numbered by the corresponding cell in A. Scalebars are 5 ms, 20 mV. p values (Wilcoxon signed-rank test) for δL: 0) 0.13, 1) < 0.01 2) 0.01, 3) 0.01.
D – F: Example network between PV+ neurons, as described for **A - C**. p values (Wilcoxon signed-rank test)

208   average intersomatic distance of 43.8 μm. In a PV-cre mouse (Fig. 4B), we identified one

209   network that was limited to two neurons, with a δL of 9.0% and intersomatic distance of 43.5

210   μm.

Discussion

Electrical synapses are widely expressed throughout the adult mammalian brain, yet

functional measurements and maps of their networks remain entirely absent in live tissue. Here

we introduce opto-δL, which leverages focal photostimulation of opsin-expressing neurons

combined with measurements of changes in spike timing, including tests of significance,

resulting from excitation across electrical synapses. Using opto-δL, we show that GABAergic

TRN neurons of differing molecular subtypes are coupled in homocellular and heterocellular

electrical networks and extend further across tissue than previously identified by dual-cell

patching[73]. Further, we identify functional network size, which is limited to far fewer neurons

than dye coupling had previously suggested[62].

In our experiments in TRN, we observed PV neurons without electrical synapses less

frequently than among SOM neurons (Fig. 3B). We hypothesize that reflects that PV expression

is more of an umbrella than a distinct marker for a subset of TRN neurons, as has long been

believed[52, 53, 78-82], and therefore, our maps of PV neurons might include both heterocellular and

homocellular electrical synapses. However, other experiments suggest that PV neurons are a

distinct subset of TRN neurons that primarily form connections with first-order thalamus[51, 83],

similar to subsets identified by the protein CB[53] and the transcription factor SPP1[52], and distinct

from SOM+ neurons. If the latter were the case, our results imply that TRN neurons that project

to first-order thalamus form more electrical synapses than SOM neurons. In turn, that difference

would imply that coordination between larger numbers of TRN neurons is more prevalent for

first-order thalamus.

The electrical connections we observed in networks emanating from one neuron were small,

ranging from 1 to 4 electrical synapses; these are the first measurements of the number of direct

electrical synapses composing a network. A previous dye-coupling study in juvenile TRN stained

networks with an average of 7 neurons, and as large as 10 to 20 neurons[62]. Dye-coupling allows

spread between neurons that are not directly coupled to the originally patched neuron, but it is

unlikely that an indirectly coupled neuron would affect spike timing by the magnitudes we

regularly observed. Our observations of networks of 1-4 neurons are consistent with results

inferring that TRN neurons form small clusters of coupled neurons[73].

As is the case with all focal optogenetic stimuli, opto-δL must account for the effects of off-

target stimulation due to photons scattering in tissue. Previous work measuring chemical

postsynaptic responses used post-hoc filtering of direct photostimulation artifacts, leveraging

distinctions in channel and photocurrent kinetics[76, 84]. Opto-δL accounts for the problem of light

scatter by comparing shifts in spike latencies caused by photostimulating a neuron to spike

latencies when an equidistant location is photostimulated (Fig. 1, Supp Fig. 3, see methods).

Opto-δL can be implemented in other brain regions, as shown by the network maps we

created in cortex (Figure 4). Unlike TRN, cortex is a mix of inhibitory and excitatory neurons,

and potentially coupled inhibitory neurons may be separated by longer distances. In cortex, an

additional barrier to identifying and measuring coupling by paired patching is the need to

visually identify GABAergic neurons. Use of fluorescent labels in opto-δL mitigates this barrier,

although it remains for targeting heterocellular networks.

Our experiments likely undersampled electrical synapses and networks for several reasons.

First, slicing brain tissue cuts some connections, causing some neuron loss. Second, our previous

experiments determined that heterocellular synapses are common[54], but we were unable to

photostimulate neurons that did not express opsin, which represent another pool of possibly

undetected coupling. Failure of virus transfection for some neurons could be a source of

misidentification in our experiments. Furthermore, due to experimental time constraints, we only

tested a subset of nearby neurons for each patched neuron; it is possible that we missed some

untested connections in experiments. Our method may also underestimate connections due to a

floor effect from light scatter compensation and differences in sensitivity between cc and δL to

very weak synapses. Therefore, while our experiments failed to detect any electrical synapses for

26 neurons, we expect that we overestimated the number of TRN neurons without electrical

synapses. It is possible that a subset of TRN neurons exists without electrical synapses, which

would be consistent with the observation that some TRN neurons do not stain for Cx36[56].

Likewise, we interpret our report of 1 to 4 electrical synapses for TRN networks as establishing a

floor for electrical network size.

Measurements of δL can in principle be used *in vivo* with focal holographic stimulation to

map electrical synapses, and we suggest that this would be a valuable next step to establish more

accurate TRN networks fully in three dimensions. As a tool, opto-δL opens the possibility for

more easily detecting and measuring coupled networks throughout the brain. For example, one

could examine how a network changes in response to different plasticity paradigms.

Alternatively, combining opto-δL with calcium imaging can be informative as to how members

of a network respond to different inputs.

Methods

Animals

SOM-Cre (Jax: 028532) or PV-Cre (Jax: 013044) mice were used for experiments. Mice

were at least P28 for surgeries and P42 for electrophysiological recordings. For one juvenile

experiment (Fig. 1), a calbindin-Cre (Jax: 028532) mouse was injected at P0 and

electrophysiological recordings occurred at P14.

Electrophysiology

Mice were transcardially perfused with NMDG ACSF solution (in mM): 92 NMDG, 2.5

KCl, 1.25 NaH_2PO_4, 30 $NAHCO_3$, 20 HEPES, 25 glucose, 2 thiourea, 5 NA ascorbate, 3 NA

pyruvate, 0.5 $CaCl_2$, and 10 $MgSO_4 \cdot 7H_2O$ (315 mOsm L^{-1}, 7.3-7.4 pH). Horizontal brain slices

250 μm thick were cut and incubated in NMDG solution. Slices were incubated at 34°C for 12-

15 min in NMDG following cutting and returned to HEPES ACSF at room temperature until

recording. The HEPES ACSF solution comprised in mM: 92 NaCl, 2.5 KCl, 1.25 NaH_2PO_4, 30

$NaHCO_3$, 20 HEPES, 25 glucose, 2 Thiourea, Na ascorbate, NA pyruvate, 2 CaCl2, and 2

$MgSO_4 \cdot 7H_2 0$ (315 mOsm L^{-1}, 7.3-7.4 pH). The ACSF bath during recording contained (in mM):

126 NaCl, 3 KCl, 1.25 NaH_2PO_4, 2 $MgSO_4$, 26 $NaHCO_3$, 10 dextrose, and 2 CaCl2 (315–320

mOsm L^{-1}, saturated with 95% O_2/5% CO_2). The submersion recording chamber was held at

34°C (TC-324B, Warner Instruments). Electrodes were filled with (in mM): 135 potassium

gluconate, 2 KCl, 4 NaCl, 10 Hepes, 0.2 EGTA, 4 ATP-Mg, 0.3 GTP-Tris, and 10

phosphocreatine-Tris (pH 7.25, 295 mOsm L^{-1}). 1 M KOH was used to adjust pH of the internal

solution. The approximate bath flowrate was 2 ml min^{-1} and the recording chamber held

approximately 5 ml solution.

The TRN was visualized under 4x magnification, and pairs of adjacent TRN cells from any

sensory sector were identified and patched under 40x IR-DIC optics (SliceScope, Scientifica,

Uckfield, UK). Cell type was determined by mRuby epifluorescence in the patched neurons.

Cells with low fluorescent contrast were avoided. The mRuby reporter was excited by a 560 nm

diode delivered through the objective (CoolLED pE-300). Voltage signals were amplified and

lowpass filtered at 8 kHz (MultiClamp, Axon Instruments, Molecular Devices, Sunnyvale, CA,

USA), digitized at 20 kHz with custom Matlab routines controlling a National Instruments

(Austin, TX, USA, USB6221 DAQ board), and data were stored for offline analysis in Matlab

(Mathworks, R2023b, Natick, MA, USA). Recordings were made in whole-cell current-clamp

mode. Negative current was used to maintain cells at −70 mV to -60 mV during recordings.

Pipette resistances were 5 - 9 MΩ before bridge balance; recordings were discarded if access

resistance exceeded 25 MΩ. Voltages are reported uncorrected for the liquid junction potential.

Stereotaxic surgery

SOM-Cre (Jax: 028532) or PV-Cre (Jax: 013044) mice were anesthetized via inhaled

isoflurane (0.8-1.2%) and 0.1 mL of 0.015 mg/mL buprenorphine via intraparietal injection.

Mice were positioned on a heated pad and in a stereotax (Kopf). After exposure of the skull, a

dental drill was used to open a hole above the injection site. Then, AAV9-CAG-DIO-ChroME-st-

P2A-H2B-mRuby (titer 2.7 x 10^{13} GC/mL) was injected at a rate of 100 nL/min (WPI UMP-3).

The coordinates for injection were -1.35 AP, ± 2.1 ML, -3.0 and -3.2 DV for TRN and -0.5 AP, ±

3.0 ML, -2.0 and -2.6 DV for somatosensory cortex. We injected a total of 600 nL of virus for

TRN and 800 nL of virus for somatosensory cortex. Post surgery and scalp suturing, mice were

given an intraperitoneal injection of 0.1 mL of 0.5 mg/mL carprofen and 1 mL saline, and

monitored for recovery. Mice were given at least two weeks for viral expression before

experiments. For the juvenile surgery, a calbindin-Cre (Jax: 028532) mouse was anesthetized via

hypothermia on a cold metal plate, and the coordinates for injection targeting TRN were -1.4 AP,

± 1.9 ML, -1.4 and -1.9 DV.

Image processing

All fluorescent images were modified in Fiji and psuedocolored to facilitate visualization.

Optogenetic δL

For the rheobase current necessary to measure δL, we used the smallest stimuli that drove

spiking at least 80% of the time. Trials were excluded if latency to spike was less than 20 ms

when the neuron was given a rheobase stimulus.

To perform single-cell optogenetic stimulation, we used a 560 nm LED (Mightex) and

digital mirror device (Mightex Polygon 1000) to direct light to an area the size of an individual

soma (7 - 8 μm diameter circle). The area size used mostly covered the entire soma without

extending beyond the edge of the soma. Prior to any experiments, we patched an opsin-

expressing cell to determine an intensity of light that was sufficient to drive bursts for a 50 ms

stimulus, and then used a photostimulus that was 50% stronger for all subsequent experiments in

that slice. Light intensity ranged from 3.7 - 99.8 mW/mm^2 with most experiments occurring in

6.2 - 25.0 mW/mm^2 range.

Single-cell optogenetic stimulation is confounded by off-target excitation by light scatter for

both 1 photon and 2 photon stimulation paradigms[74-76]. For the purposes of mapping synaptic

connections, scatter-induced photostimulation of the patched neuron is a potential confound.

Depolarization from photostimulation caused by light scatter is directly proportional to the

distance between the patched neuron and focal stimulation (Supp. Fig. 2). Opto-δL is well-suited

to overcome the problem of unintended photostimulation to the patched neuron because the

method quantifies latency of spikes rather than voltage changes. Consequently, comparing

latency of spikes when a nearby neuron is photostimulated to latency when an equidistant empty

control location is photostimulated allowed us to distinguish the contribution of GJs to spike

times.

After patching TRN neurons from any sensory sector, we used one of two methods to first

make measurements that would control for the effects of light scattering, and then to compute

δL.

i)   Corrections to δL: We delivered a rheobase stimulus to the patched neuron alone 10

times. Interleaved between those stimulations, we gave the neuron a rheobase

stimulus while also focally photostimulating either a nearby mRuby+ neuron or an

equidistant empty spot. Offline, we measured the latency of spikes from stimulation

onset for each condition. We report δL as the change in latency induced by focal

photostimulation of the nearby neuron minus the change in latency induced by

photostimulation of an empty area.

ii)   Correction to Iapp: We applied focal photostimulation ten times to an empty location

equidistant from the patched neuron as the nearby neuron of interest. We measured

the average depolarizing current that resulted from the light scattering. We then

adjusted rheobase Iapp by average current caused by light scatter (Supp. Fig. 3) and

used that adjusted Iapp while photostimulating the nearby mRuby+ or an equidistant

362   empty location. Offline, we calculated δL by comparing latency changes between

363   when light targeted the empty control location (t1) compared to light targeting the

364   mRuby+ neuron (t1).

365  We used corrected δL directly for 30 homocellular network measurements, and we used

366  correction to I_{app} for 29 homocellular network measurements.

367  *Numerical analysis*

368  To calculate δL we used the average spike latency from control trials (t_1) and average spike

369 latency when a nearby neuron was photostimulated (t_2):

370
$$\delta L = \frac{100 * (t_1 - t_2)}{t_1}$$

371  We tested δLs for significance with a Wilcoxon signed-rank test. We compared δLs between

372 different neuronal subtypes via Mann-Whitney U test, and we used Chi-squared tests to compare

373 electrical synapses counts per network between different genetic subtypes. Statistical tests were

374 considered significant when $p < 0.05$. In our analysis of network size, we only included

375 experiments where three or more possible connections were tested.

A SOM-st-ChroMe-mRuby3

B

Supplementary Figure 1: Validation of single-cell focal photostimulation. A. Expression pattern of st-ChoME-mRuby3 via AAV injection in a SOM-Cre mouse (fluorescence pseudocolored magenta). Expression was restricted to the shell of TRN. **B.** Left, spiking driven by increasing area of light in a spot (3.14, 7.07, 14.14, 22.0, 31.4, 47.2 μm^2 circles). Right, spiking driven by increasing light intensity (24.4, 37.1, 59.2, 66.2, 80.1, 87.0, 105.7 mW/mm^2). Scale bar 25 ms, 10 mV. $V_m = -61$ mV.

Supplementary Figure 2: **Example of excitation from light scattering. A.** Overlay of IR and fluorescence images from live recording of st-ChroME-mRuby-expressing neurons in PV-Cre mouse. Empty locations were serially photostimulated by increasing 10 μm intervals measured from the edge of the soma from the patched neuron. The power of the photostimulation was 10.5 mW/mm^2 and was sufficient to drive the neuron to burst when targeted on the soma. Scale bar: 50 μm. **B.** Voltage responses of the patched neuron from focally stimulating the corresponding numbered locations in A. Scale bar: 10 ms, 1 mV. **C.** Shifts in in spike latency caused by focally photostimulating empty locations in A. **C1.** $\delta L = 13.4\%$, $p < 0.01$. **C2.** $\delta L = 9.2\%$, $p < 0.01$. **C3.** $\delta L = 5.2\%$, $p < 0.01$. **C4.** $\delta L = 3.9\%$, $p = 0.04$. **C5.** $\delta L = 3.5\%$, $p = 0.07$. **C6.** $\delta L = 1.8\%$, $p = 0.38$. Scale bar: 5 ms, 20 mV. Asterix mark significance ($p < 0.05$, Wilcoxon signed-rank test).

Supplementary Figure 3: **Compensation for effects of scatter-induced photoexcitation on spike times.**
A. Overlay of IR and fluorescent images of ChroME-mRuby-expressing neurons in a SOM-Cre mouse.
Fluorescence is pseudocolored magenta. Scale bar: 50 μm. The patched neuron is circled with white dashes.
Empty regions of interests (ROIs) (e.g. orange circles) at different distances were photostimulated to
determine the compensation for scatter to rheobase for the patched neuron. **B.** Spiking responses for
rheobase (51 pA) applied current alone (black traces; mean latency 38.6 ms) or during simultaneous
photostimulation of ROI1 (green traces, mean latency 31.2 ms). Dots mark the times of first spikes from
each trial. Uncompensated scatter from ROI 1 caused a false δL of 19%. **C.** Spiking responses for
compensated rheobase (22 pA) and simultaneous photostimulation of ROI 2 (blue traces; mean latency 36.6
ms), and for compensated rheobase during photostimulation of ROI 1 (green traces; mean latency 36.5 ms).
D. Overlay of IR and fluorescence (pseudocolor: magenta) images of ChroME-mRuby-expressing neurons
in a slice from a SOM-Cre mouse. Scale bar 50 μm. The patched neuron is circled with white dashes.
Without any photostimuli, rheobase was determined to be 65 pA. Two non-fluorescing ROIs, circled in
orange, were selected for photostimulation. **E.** Spiking in the recorded neuron after correction of rheobase
(square pulse, lower) for scatter to 46 pA during photostimulation of ROI 1 (orange bar). The corrected
mean latency of the first spike was 35 ms. **F.** Spiking in the recorded neuron after correction of rheobase for
scatter to 62 pA (square pulse, below) during photostimulation of ROI 2 (red bar). The mean corrected
latency of the first spike was 34 ms. Scale bars are 25 ms, 10 mV.

1.   Belluardo N, Mudò G, Trovato-Salinaro A, Le Gurun S, Charollais A, Serre-Beinier V, Amato G, Haefliger J-A, Meda P, Condorelli DF. Expression of connexin36 in the adult and developing rat brain. Brain research. 2000;865(1):121-38.

2.   Condorelli DF, Belluardo N, Trovato-Salinaro A, Mudò G. Expression of Cx36 in mammalian neurons. Brain Research Reviews. 2000;32(1):72-85.

3.   Rash J, Staines W, Yasumura T, Patel D, Furman C, Stelmack G, Nagy J. Immunogold
evidence that neuronal gap junctions in adult rat brain and spinal cord contain connexin-36 but not
connexin-32 or connexin-43. Proceedings of the National Academy of Sciences. 2000;97(13):7573-
8.
4.   Bennett MV, Zukin RS. Electrical coupling and neuronal synchronization in the Mammalian
brain. Neuron. 2004;41(4):495-511. Epub 2004/02/26. doi: 10.1016/s0896-6273(04)00043-1.
PubMed PMID: 14980200.
5.   Abbott L, Marder E, Hooper SL. Oscillating networks: control of burst duration by electrically
coupled neurons. Neural Computation. 1991;3(4):487-97.
6.   Traub RD, Kopell N, Bibbig A, Buhl EH, LeBeau FE, Whittington MA. Gap junctions between
interneuron dendrites can enhance synchrony of gamma oscillations in distributed networks.
Journal of Neuroscience. 2001;21(23):9478-86.
7.   Gutierrez GJ, O'Leary T, Marder E. Multiple mechanisms switch an electrically coupled,
synaptically inhibited neuron between competing rhythmic oscillators. Neuron. 2013;77(5):845-58.
8.   Destexhe A. Spike-and-wave oscillations based on the properties of GABAB receptors.
Journal of Neuroscience. 1998;18(21):9099-111.
9.   Draguhn A, Traub R, Schmitz D, Jefferys J. Electrical coupling underlies high-frequency
oscillations in the hippocampus in vitro. Nature. 1998;394(6689):189-92.
10.   Whittington MA, Traub RD. Interneuron diversity series: inhibitory interneurons and network
oscillations in vitro. Trends in neurosciences. 2003;26(12):676-82.
11.   Chow CC, Kopell N. Dynamics of spiking neurons with electrical coupling. Neural Comput.
2000;12(7):1643-78. Epub 2000/08/10. doi: 10.1162/089976600300015295. PubMed PMID:
10935921.
12.   Haas JS, Landisman CE. State-dependent modulation of gap junction signaling by the
persistent sodium current. Front Cell Neurosci. 2011;5:31. Epub 2012/02/10. doi:
10.3389/fncel.2011.00031. PubMed PMID: 22319469; PMCID: PMC3263475.
13.   Pernelle G, Nicola W, Clopath C. Gap junction plasticity as a mechanism to regulate
network-wide oscillations. PLoS computational biology. 2018;14(3):e1006025.
14.   Pfeuty B, Mato G, Golomb D, Hansel D. The combined effects of inhibitory and electrical
synapses in synchrony. Neural Computation. 2005;17(3):633-70.
15.   Pfeuty B, Mato G, Golomb D, Hansel D. Electrical synapses and synchrony: the role of
intrinsic currents. Journal of Neuroscience. 2003;23(15):6280-94.
16.   DeVries SH, Qi X, Smith R, Makous W, Sterling P. Electrical coupling between mammalian
cones. Current Biology. 2002;12(22):1900-7.
17.   Medvedev GS. Electrical coupling promotes fidelity of responses in the networks of model
neurons. Neural computation. 2009;21(11):3057-78.
18.   Zhu P, Frank T, Friedrich RW. Equalization of odor representations by a network of
electrically coupled inhibitory interneurons. Nature neuroscience. 2013;16(11):1678-86.
19.   Galarreta M, Hestrin S. Spike transmission and synchrony detection in networks of
GABAergic interneurons. Science. 2001;292(5525):2295-9.
20.   Gibson JR, Beierlein M, Connors BW. Functional properties of electrical synapses between
inhibitory interneurons of neocortical layer 4. Journal of neurophysiology. 2005;93(1):467-80.
21.   Russo G, Nieus TR, Maggi S, Taverna S. Dynamics of action potential firing in electrically
connected striatal fast-spiking interneurons. Frontiers in cellular neuroscience. 2013;7:209.
22.   Apostolides PF, Trussell LO. Control of interneuron firing by subthreshold synaptic
potentials in principal cells of the dorsal cochlear nucleus. Neuron. 2014;83(2):324-30.

23.   Trenholm S, McLaughlin AJ, Schwab DJ, Awatramani GB. Dynamic tuning of electrical and
chemical synaptic transmission in a network of motion coding retinal neurons. Journal of
Neuroscience. 2013;33(37):14927-38.
24.   Deans MR, Volgyi B, Goodenough DA, Bloomfield SA, Paul DL. Connexin36 is essential for
transmission of rod-mediated visual signals in the mammalian retina. Neuron. 2002;36(4):703-12.
25.   Christie JM, Westbrook GL. Lateral excitation within the olfactory bulb. Journal of
Neuroscience. 2006;26(8):2269-77.
26.   Ren Y, Liu Y, Luo M. Gap Junctions Between Striatal D1 Neurons and Cholinergic
Interneurons. Frontiers in Cellular Neuroscience. 2021;15:191.
27.   Veruki ML, Hartveit E. Electrical synapses mediate signal transmission in the rod pathway of
the mammalian retina. Journal of Neuroscience. 2002;22(24):10558-66.
28.   Alcami P. Electrical synapses enhance and accelerate interneuron recruitment in response
to coincident and sequential excitation. Frontiers in cellular neuroscience. 2018;12:156.
29.   Rabinowitch I, Chatzigeorgiou M, Schafer WR. A gap junction circuit enhances processing
of coincident mechanosensory inputs. Current Biology. 2013;23(11):963-7.
30.   Pham T, Haas JS. Electrical synapses regulate both subthreshold integration and population
activity of principal cells in response to transient inputs within canonical feedforward circuits. PLoS
computational biology. 2019;15(2):e1006440.
31.   Pham T, Haas JS. Electrical synapses between inhibitory neurons shape the responses of
principal neurons to transient inputs in the thalamus: a modeling study. Scientific reports.
2018;8(1):1-12.
32.   Hoehne A, McFadden MH, DiGregorio DA. Feed-forward recruitment of electrical synapses
enhances synchronous spiking in the mouse cerebellar cortex. Elife. 2020;9:e57344.
33.   Kuo SP, Schwartz GW, Rieke F. Nonlinear spatiotemporal integration by electrical and
chemical synapses in the retina. Neuron. 2016;90(2):320-32.
34.   Szarka G, Ganczer A, Balogh M, Tengölics ÁJ, Futácsi A, Kenyon G, Pan F, Kovács-Öller T,
Völgyi B. Gap Junctions Fine-tune Ganglion Cell Signals to Equalize Response Kinetics within a
Given Electrically Coupled Array. iScience. 2024.
35.   Cooler S, Schwartz GW. An offset ON–OFF receptive field is created by gap junctions
between distinct types of retinal ganglion cells. Nature neuroscience. 2021;24(1):105-15.
36.   Wang S, Borst A, Zaslavsky N, Tishby N, Segev I. Efficient encoding of motion is mediated by
gap junctions in the fly visual system. PLoS computational biology. 2017;13(12):e1005846.
37.   Bennett MV. Physiology of electrotonic junctions. Ann N Y Acad Sci. 1966;137(2):509-39.
Epub 1966/07/14. doi: 10.1111/j.1749-6632.1966.tb50178.x. PubMed PMID: 5229812.
38.   van Welie I, Roth A, Ho SS, Komai S, Häusser M. Conditional spike transmission mediated
by electrical coupling ensures millisecond precision-correlated activity among interneurons in vivo.
Neuron. 2016;90(4):810-23.
39.   Stewart WW. Functional connections between cells as revealed by dye-coupling with a
highly fluorescent naphthalimide tracer. Cell. 1978;14(3):741-59.
40.   Connors B, Benardo L, Prince D. Coupling between neurons of the developing rat neocortex.
Journal of Neuroscience. 1983;3(4):773-82.
41.   Hampson EC, Weiler R, Vaney DI. pH-gated dopaminergic modulation of horizontal cell gap
junctions in mammalian retina. Proc Biol Sci. 1994;255(1342):67-72. Epub 1994/01/22. doi:
10.1098/rspb.1994.0010. PubMed PMID: 8153138.
42.   Hampson E, Vaney DI, Weiler R. Dopaminergic modulation of gap junction permeability
between amacrine cells in mammalian retina. Journal of Neuroscience. 1992;12(12):4911-22.
43.   Peinado A, Yuste R, Katz LC. Extensive dye coupling between rat neocortical neurons during
the period of circuit formation. Neuron. 1993;10(1):103-14.

44. Onn S, Grace AA. Dye coupling between rat striatal neurons recorded in vivo: compartmental organization and modulation by dopamine. Journal of Neurophysiology. 1994;71(5):1917-34.

45. Mills SL, Xia XB, Hoshi H, Firth SI, Rice ME, Frishman LJ, Marshak DW. Dopaminergic modulation of tracer coupling in a ganglion-amacrine cell network. Vis Neurosci. 2007;24(4):593-608. Epub 2007/08/23. doi: 10.1017/S0952523807070575. PubMed PMID: 17711603; PMCID: PMC2213423.

46. Ribelayga C, Cao Y, Mangel SC. The circadian clock in the retina controls rod-cone coupling. Neuron. 2008;59(5):790-801. Epub 2008/09/13. doi: 10.1016/j.neuron.2008.07.017. PubMed PMID: 18786362; PMCID: PMC5581203.

47. Qiao M, Sanes JR. Genetic method for labeling electrically coupled cells: application to retina. Frontiers in molecular neuroscience. 2016;8:81.

48. Tian L, Yang Y, Wysocki LM, Arnold AC, Hu A, Ravichandran B, Sternson SM, Looger LL, Lavis LD. Selective esterase–ester pair for targeting small molecules with cellular specificity. Proceedings of the National Academy of Sciences. 2012;109(13):4756-61.

49. Pinault D. The thalamic reticular nucleus: structure, function and concept. Brain Res Brain Res Rev. 2004;46(1):1-31. Epub 2004/08/07. doi: 10.1016/j.brainresrev.2004.04.008. PubMed PMID: 15297152.

50. Crick F. Function of the thalamic reticular complex: the searchlight hypothesis. Proc Natl Acad Sci U S A. 1984;81(14):4586-90. Epub 1984/07/01. doi: 10.1073/pnas.81.14.4586. PubMed PMID: 6589612; PMCID: PMC345636.

51. Clemente-Perez A, Makinson SR, Higashikubo B, Brovarney S, Cho FS, Urry A, Holden SS, Wimer M, Dávid C, Fenno LE. Distinct thalamic reticular cell types differentially modulate normal and pathological cortical rhythms. Cell reports. 2017;19(10):2130-42.

52. Li Y, Lopez-Huerta VG, Adiconis X, Levandowski K, Choi S, Simmons SK, Arias-Garcia MA, Guo B, Yao AY, Blosser TR. Distinct subnetworks of the thalamic reticular nucleus. Nature. 2020:1-6.

53. Martinez-Garcia RI, Voelcker B, Zaltsman JB, Patrick SL, Stevens TR, Connors BW, Cruikshank SJ. Two dynamically distinct circuits drive inhibition in the sensory thalamus. Nature. 2020;583(7818):813-8.

54. Vaughn MJ, Laswick Z, Wang H, Haas JS. Functionally distinct circuits are linked by heterocellular electrical synapses in the thalamic reticular nucleus. eneuro. 2023;11(1).

55. Hartley ND, Krol A, Choi S, Rome N, Levandowski K, Pasqualoni S, Jones C, Tian J, Lee S, Lee H. Distinct structural and functional connectivity of genetically segregated thalamoreticular subnetworks. Cell Reports. 2024;43(12).

56. Landisman CE, Long MA, Beierlein M, Deans MR, Paul DL, Connors BW. Electrical synapses in the thalamic reticular nucleus. J Neurosci. 2002;22(3):1002-9. Epub 2002/02/05. PubMed PMID: 11826128; PMCID: PMC6758490.

57. Haas JS, Zavala B, Landisman CE. Activity-dependent long-term depression of electrical synapses. Science. 2011;334(6054):389-93. Epub 2011/10/25. doi: 10.1126/science.1207502. PubMed PMID: 22021860.

58. Sevetson J, Haas JS. Asymmetry and modulation of spike timing in electrically coupled neurons. J Neurophysiol. 2015;113(6):1743-51. Epub 2014/12/30. doi: 10.1152/jn.00843.2014. PubMed PMID: 25540226.

59. Landisman CE, Connors BW. Long-term modulation of electrical synapses in the mammalian thalamus. Science. 2005;310(5755):1809-13.

60.   Deleuze C, Huguenard JR. Distinct electrical and chemical connectivity maps in the
thalamic reticular nucleus: potential roles in synchronization and sensation. Journal of
Neuroscience. 2006;26(33):8633-45.
61.   Lam Y-W, Nelson CS, Sherman SM. Mapping of the functional interconnections between
thalamic reticular neurons using photostimulation. Journal of neurophysiology. 2006;96(5):2593-
600.
62.   Lee S-C, Patrick SL, Richardson KA, Connors BW. Two functionally distinct networks of gap
junction-coupled inhibitory neurons in the thalamic reticular nucleus. Journal of Neuroscience.
2014;34(39):13170-82.
63.   Degen J, Meier C, Van Der Giessen RS, Söhl G, Petrasch-Parwez E, Urschel S, Dermietzel R,
Schilling K, De Zeeuw CI, Willecke K. Expression pattern of lacZ reporter gene representing
connexin36 in transgenic mice. Journal of Comparative Neurology. 2004;473(4):511-25.
64.   Blethyn KL, Hughes SW, Crunelli V. Evidence for electrical synapses between neurons of the
nucleus reticularis thalami in the adult brain in vitro. Thalamus & related systems. 2008;4(1):13-20.
65.   Mainen ZF, Sejnowski TJ. Reliability of spike timing in neocortical neurons. Science.
1995;268(5216):1503-6.
66.   Gütig R. To spike, or when to spike? Current opinion in neurobiology. 2014;25:134-9.
67.   VanRullen R, Guyonneau R, Thorpe SJ. Spike times make sense. Trends in neurosciences.
2005;28(1):1-4.
68.   Stüttgen MC, Schwarz C. Psychophysical and neurometric detection performance under
stimulus uncertainty. Nature neuroscience. 2008;11(9):1091-9.
69.   Gollisch T, Meister M. Rapid neural coding in the retina with relative spike latencies.
science. 2008;319(5866):1108-11.
70.   Kilinc D, Demir A. Spike timing precision of neuronal circuits. Journal of computational
neuroscience. 2018;44:341-62.
71.   Haas JS. A new measure for the strength of electrical synapses. Frontiers in cellular
neuroscience. 2015;9:378.
72.   Mardinly AR, Oldenburg IA, Pégard NC, Sridharan S, Lyall EH, Chesnov K, Brohawn SG,
Waller L, Adesnik H. Precise multimodal optical control of neural ensemble activity. Nature
neuroscience. 2018;21(6):881-93.
73.   Long MA, Landisman CE, Connors BW. Small clusters of electrically coupled neurons
generate synchronous rhythms in the thalamic reticular nucleus. Journal of Neuroscience.
2004;24(2):341-9.
74.   Shemesh OA, Tanese D, Zampini V, Linghu C, Piatkevich K, Ronzitti E, Papagiakoumou E,
Boyden ES, Emiliani V. Temporally precise single-cell-resolution optogenetics. Nature
neuroscience. 2017;20(12):1796-806.
75.   Hage TA, Bosma-Moody A, Baker CA, Kratz MB, Campagnola L, Jarsky T, Zeng H, Murphy GJ.
Synaptic connectivity to L2/3 of primary visual cortex measured by two-photon optogenetic
stimulation. Elife. 2022;11:e71103.
76.   Antin B, Sadahiro M, Gajowa M, Triplett MA, Adesnik H, Paninski L. Removing direct
photocurrent artifacts in optogenetic connectivity mapping data via constrained matrix
factorization. PLOS Computational Biology. 2024;20(5):e1012053.
77.   Hu H, Cavendish JZ, Agmon A. Not all that glitters is gold: off-target recombination in the
somatostatin–IRES-Cre mouse line labels a subset of fast-spiking interneurons. Frontiers Media SA;
2013. p. 195.
78.   Frassoni C, Bentivoglio M, Spreafico R, Sánchez MP, Puelles L, Fairen A. Postnatal
development of calbindin and parvalbumin immunoreactivity in the thalamus of the rat.
Developmental brain research. 1991;58(2):243-9.

79.   Mitrofanis J. Patterns of antigenic expression in the thalamic reticular nucleus of developing
rats. Journal of Comparative Neurology. 1992;320(2):161-81.
80.   Hou G, Smith AG, Zhang ZW. Lack of Intrinsic GABAergic Connections in the Thalamic
Reticular Nucleus of the Mouse. J Neurosci. 2016;36(27):7246-52. Epub 2016/07/08. doi:
10.1523/JNEUROSCI.0607-16.2016. PubMed PMID: 27383598; PMCID: PMC4938865.
81.   Thankachan S, Katsuki F, McKenna JT, Yang C, Shukla C, Deisseroth K, Uygun DS, Strecker
RE, Brown RE, McNally JM. Thalamic reticular nucleus parvalbumin neurons regulate sleep
spindles and electrophysiological aspects of schizophrenia in mice. Scientific reports. 2019;9(1):1-
16.
82.   Seto-Ohshima A, Emson P, Berchtold M, Heizmann C. Localization of parvalbumin mRNA in
rat brain by in situ hybridization histochemistry. Experimental brain research. 1989;75(3):653-8.
83.   Hoseini MS, Higashikubo B, Cho FS, Chang AH, Clemente-Perez A, Lew I, Ciesielska A,
Stryker MP, Paz JT. Gamma rhythms and visual information in mouse V1 specifically modulated by
somatostatin+ neurons in reticular thalamus. Elife. 2021;10:e61437.
84.   Printz Y, Patil P, Mahn M, Benjamin A, Litvin A, Levy R, Bringmann M, Yizhar O. Determinants
of functional synaptic connectivity among amygdala-projecting prefrontal cortical neurons in male
mice. Nature communications. 2023;14(1):1667.

638