Cooperative Behaviors and Social Interactions in the Carnivorous Bat *Vampyrum spectrum*

Marisa Tietge[1,2] *, Eduardo Artavia Durán[3], Mirjam Knörnschild[1,2,4]

[1] Museum for Natural History, Leibniz Institute for Evolution and Biodiversity Science, 10115 Berlin, Germany

[2] Institute for Biology, Humboldt-Universität zu Berlin, 10099 Berlin, Germany

[3] Manzú Conservación de Murciélagos, 50102, Costa Rica

[4] Smithsonian Tropical Research Institute, 0843-03092, Panama

*Corresponding author: Marisa Tietge

Keywords: Biparental care; Chiroptera; Food provisioning; Neotropical; Phyllostomidae; Prey sharing; Roost; Social behavior

Highlights

Extensive parental care: Most likely both male and female spectral bats participate in parental care, e.g. food provisioning, a rare behavior among mammals

Prey sharing as a social cooperative behavior: Detailed observations of prey sharing within the roost, highlighting social cooperation and resource allocation

Extended dependency period for pups: A prolonged developmental period allows pups to receive care and acquire essential survival skills until they achieve full independence

Juvenile play: Pups engaged predominantly in social play (mock fights), but also played with objects or food items, a rare behavior among bats

Abstract

Bats exhibit a diverse array of social behaviors, yet detailed studies on the intricacies of these interactions, particularly in rare species like the spectral bat (*Vampyrum spectrum*), remain scarce. This study presents the first comprehensive description of prey sharing and other social behaviors in a wild social group of *V. spectrum*. Over several months, we conducted extensive video recordings in a hollow tree, used as a day- and night-roost, to document these behaviors, aiming to elucidate key social interactions among the bats and their role within their ecological niche. We observed various remarkable social behaviors, including food sharing between family members. Our findings support the hypothesis that prey sharing as a form of biparental care may serve as a method for adults to transition young bats from milk to a carnivorous diet, ensuring adequate food intake and allowing them to practice how to handle large prey items. We also challenge the notion that *V. spectrum* forages exclusively solitarily, as we documented several instances of synchronized roost departures and returns and thus presumably cooperative foraging. This indicates a more complex social structure and behavioral ecology than previously understood. Our comprehensive analysis of observational data enhances our understanding of the social dynamics of *V. spectrum*, providing new insights into the evolution of cooperative behaviors in bats.

Introduction

Cooperative behaviors are observed across many animal species and play a crucial role in

their survival and social structures (1–3). Such behaviors often involve individuals working

together to achieve shared goals, like defending territories, caring for young, or securing

resources (4, 3, 5). These interactions are especially well-documented in social animals,

where cooperation can range from kinship-based behaviors to complex alliances among

unrelated individuals (6–8). In addition to kinship, alliances among non-relatives are

widespread; for instance, in species like baboons and dolphins, unrelated individuals

cooperate to hunt more effectively, which enhances foraging success (9–12). This behavior

allows access to larger or more challenging prey than would be possible alone, illustrating

how collaborative behaviors can enhance survival and fitness within an ecological niche (13).

Bats, with over 1,480 species (14), exhibit cooperative behaviors on many levels. As one of

the most ecologically significant and diverse orders of mammals, bats occupy a broad range

of habitats and exhibit various dietary preferences – including insectivory, frugivory,

nectarivory, piscivory, and even carnivory (15, 16). Their social structures are equally varied,

spanning solitary roosting, complex communal arrangements, such as harems, social

monogamy, and promiscuity (17, 18), making them an exceptional case for studying the

dynamics of cooperation across different social systems. Most bat species are highly

gregarious, presumably due to ecological constraints like roost limitation, physiological

demands like social thermoregulation, and their longevity (18). Bats exhibit complex

communication, group-level recognition, and various other social interactions among

individuals (19). Some remarkable social behaviors have evolved, like roost sharing and

defense in tent-making bats where individuals construct tents from large leaves. These

shelters provide protection from predators and harsh weather. Bats within these groups

cooperate in the maintenance and defense of their roosts (20–22). Previous studies have

shown that bats from various social systems can show cooperative behavior when it comes

to the availability of food and prey capture using auditory, visual, and olfactory cues,

especially when living in living in fission-fusion systems (23–29). Behavioral evidence also

suggests that bats use calls to distinguish between familiar and unfamiliar conspecifics, like

the Spix's disk-winged bats (*Thyroptera tricolor*) that uses social calls to stay in touch with

group members while foraging (30). This behavior is similar to "grooming at a distance" in

primates, where individuals that frequently groom each other maintain vocal contact when

separated during foraging (reviewed in (31). Maintaining contact while foraging may help

bats ensure they later roost with preferred group members and reinforce cooperative

behavior (30).

Understanding the social behaviors of bats is crucial for several reasons. Social interactions

can impact reproductive success, foraging efficiency, predator avoidance, and disease

transmission (32–40). These behaviors are also integral to the conservation and

management of bat populations, many of which are threatened by habitat loss, climate

change, and human activities (41). Bat species developed a lot of different social behaviors,

most of the them are still unknown because bats are often hard to observe due to their

nocturnal and highly mobile lifestyle. Especially bat species that are rare to find and/or are

critically endangered are understudied.

The spectral bat (*Vampyrum spectrum*) is widely recognized as the largest carnivorous bat

species in the New World, with individuals weighing approximately 180 grams and boasting

a wingspan of up to 900 millimeters (42). This rare species can be found in lowland tropical

dry forests, evergreen forests, and occasionally in cloud forests, deciduous forests, or

swampy areas (43). *Vampyrum spectrum* is considered near threatened by the International

Union for Conservation of Nature (IUCN) and is listed as a species of special concern or

endangered in several countries throughout its range with a decreasing population trend

(44). Belonging to the family of leaf-nosed bats (Phyllostomidae), these formidable predators

have a diverse diet that includes a variety of birds, rodents, and small mammals, including

other bat species (42, 45). *Vampyrum spectrum* typically roosts in small groups within hollow

trees or caves. These groups exhibit a socially monogamous structure, usually comprising a

single male, a single female, and their recent offspring who have not yet dispersed (43). This

socially monogamous social structure is relatively rare among mammals, making this species

a particularly interesting subject for study.

The primary goals of this study were to get insights into the social behaviors of *Vampyrum*

spectrum within the context of their ecological niche as carnivorous bats. By conducting

extensive video recordings in their roost, we aimed to document and analyze key behaviors,

with a focus on social interactions and cooperative behaviors. This research not only aims to

shed light on the social behaviors of *V. spectrum* but also contributes to the broader

understanding of social behavior in bats.

Methods

Study site

The study was conducted in the tropical dry forest of "La Estación Experimental Forestal

Horizontes" (N 10° 42' 47.883973, E -85° 35' 43.759147) in Guanacaste, Costa Rica. The focal

roost is located within a partially hollow yet living tree of the species *Manilkara chicle*,

amidst typical dry forest vegetation and adjacent to a dried riverbed. This roost was first

identified by M.T. in December 2022, housing four *V. spectrum* individuals (Fig. 1), and we

began systematically monitoring them approximately one year later. The *V. spectrum* bats

occupy the highest section of the hollow tree, while a colony of 15 smaller bats, *Saccopteryx*

bilineata, inhabits a lower section near the entrance of the tree hole. Although we did not

individually mark the four *V. spectrum* bats we are confident that this colony consisted of a

female and male bat pair and their respective two young for several reasons: 1) this species

lives in a socially monogamous social structure that either constitutes a single roosting bat

or family group (42), 2) the bats in the video displayed noticeable size differences and the

teats of the adult, post-lactating female were visible on occasion, and 3) additional video

recordings in 2024 showed a newly born pup being nursed by its mother and made it

possible to identify the other large individual as male (because the penis was visible).

Data collection

At the beginning of the dry season in November 2023, a wildlife camera (Snapshot Mini

Black 30MP 4K, Dörr GmbH) was installed inside the roost at a height of approximately 10

meters, using a small camera mount (CAMVATE photography mount with 1/4"-20 thread).

The camera was positioned about 2 meters from the primary roosting site of the bats,

oriented upwards to capture optimal footage from below. To ensure minimal disturbance to

the bats, the camera was carefully installed during a period of low activity. Observations

following the installation indicated that the bats' behavior remained unaffected by the

presence of the equipment. Over the course of three months, the camera was regularly

checked, cleaned, equipped with a new SD card, and fitted with fresh batteries before being

reinstalled at the fixed location within the tree. The camera recorded automatically triggered

infrared videos with a maximum duration of one minute. Initially, continuous recording was

achieved each night; however, as battery power depleted, the camera's built-in power-saving

function reduced the length of recorded videos.

Over the course of three months, we video-recorded the four bats for a total of 60 days. On

some days, only a few videos were captured, while on other days, up to 40 videos were

recorded, depending on how frequently the camera sensor was triggered by bat activity. In

total, 502 videos were collected. Most of these videos (203 out of 502) were triggered by

bats either leaving or entering the roost. The other bat species, *S. bilineata,* was very seldom

recorded, as the camera was installed above their roosting spot. Out of the 502 recorded

videos, 73 contained social behaviors or other interesting activity. By visual examination of

the videos, we classified eight different behavioral categories: "social roosting", "greeting

behavior", "presumably sexual behavior", "bringing prey into the roost", "eating a prey item",

"prey sharing", "food checks", and "play behavior". All observed behaviors were recorded

during the night between 5pm and 6am and occurred with no specific distribution pattern

(see Fig. 2). In order to ensure long-term monitoring of this social group, further videos are

recorded in the colony in 2024/2025. All data presented in this study are based on video

recordings collected during the 2023/2024 season. However, one additional video from the

2024/2025 season was included as an anecdotal observation due to its particularly

interesting and relevant nature.

Individual distinction by size estimation

Due to the absence of individual markings on the animals, we employed supplementary

methods to support our observations and interpretations of behaviors, particularly in prey-

sharing interactions. We measured the distance between the eyes of the bat entering the

roost with prey and the bat approaching to take over the prey. This was done using

screenshots from video frames capturing prey-sharing behaviors, where both animals were

positioned at approximately the same height and their eyes were clearly visible (N=8 prey

sharing interactions). The screenshots were analyzed in GIMP (GIMP 2.10.38), utilizing the

tape measure function to determine the eye distance in pixels. These measurements were

subsequently compared using a non-parametric paired samples Wilcoxon test. Additionally,

we used a spaghetti plot to visualize the results (see figure 4).

All statistical analyses were performed using the program Sypder (Spyder IDE, The Scientific

Python Development Environment, Python 3.12.4)

Results

Based on the 73 infrared videos, we created an ethogram that contained eight different

behavioral categories exhibited by *V. spectrum* bats in their roost. Video examples of the

respective behaviors can be found in the electronic supplementary material.

Social roosting (16 out of 73 videos): This behavior involves two or more bats roosting in

close proximity with body contact, establishing a ball-like formation. At least one bat wraps

its wings around the other bat(s). This behavior is often accompanied by allogrooming

and/or social vocalizations and was observed very frequently.

Greeting Behavior (4 out of 73 videos): This behavior constitutes of a hugging-like interaction

between a bat already in the roost and a newly arrived bat. The resident bat may actively

approach or greet the newcomer as it reaches close proximity in the main roosting area. The

greeting behavior is comparable to the initiation to social roosting, where at least one bat

wraps its wings around the other, establishing a ball-like formation for several seconds. This

behavior may be accompanied by social vocalizations.

Presumably sexual behavior (2 out of 73 videos): This behavior involves two bats, with one

bat—presumably the male—positioned behind the female, aligning belly to back while

wrapping its wings around her. This interaction often includes frequent position changes and

licking of various lower body parts. This kind of behavior was only recorded twice and

therefore we can only assume that this observed behavior is indeed produced in a sexual

context.

Bringing prey into the roost (4 out of 73 videos): A bat enters the roost after foraging,

carrying a prey item in its mouth (normally a bird or rodent). The prey is always held by its

head and appears already dead. The bat carries the prey item in its mouth and crawls

backward to the main roosting area in the hollow tree, with only the back or tail of the prey

visible.

Eating a prey item (18 out of 73 videos): A bat suspends itself at a specific location within the

roost, positioning the prey item between its wings and securing it with its thumb claws while

biting into it. Audible chewing noises are a distinctive feature of this process. We estimate

that consuming a single prey item may takes approximately 30-40 minutes, based on

sequential video recordings showing bats feeding on the same prey item. Nevertheless, this

duration remains unverified due to the current limitation of our video equipment, which

recorded only 1-minute segments. A video sequence was classified as "eating a prey item"

when there was no immediate social interaction with another bat preceding or following the

behavior. In 12 videos, we could not observe the bat entering the roost, only the eating

process was recorded, while in six videos a bat entered the roost with prey, chose not to

share it, and consumed it alone.

Prey sharing (12 out of 73 videos): This behavior involves a bat entering the roost with a prey

item held in its mouth. In many instances, as soon as the prey-carrying bat climbs to the

main roosting spot, normally only one other bat approaches while emitting vocalizations.

The approaching bat comes very close to the prey-carrying bat, typically from the back

and/or side. The approaching bat attempts to bite into the upper body of the prey. In most

cases (92%), the prey-carrying bat voluntarily releases the prey item quickly. Subsequently,

both bats crawl up to the main roosting location where the bat with the prey usually begins

to consume the prey item immediately (see Fig. 3). On a single occasion, a recorded

interaction between two bats involved a fight over prey brought back into the roost,

resembling a tug-of-war scenario. Additionally, we recorded an anecdotal observation in a

video from December 2024, where a bat—presumed to be the mother—was present in the

roost alongside a small pup when another bat entered carrying prey. The mother

approached the incoming bat, swiftly and silently took the prey, and then returned to the

primary roosting spot next to the pup with the prey to consume it.

Food checks (4 out of 73 videos): One bat enters the roost, and another bat approaches,

seemingly to check if the entering bat has brought back food from foraging. The approaching

bat displays behaviors similar to prey-sharing interactions, though in these cases, no prey is

present. After realizing this, both bats typically crawl up to the main roosting spot.

Play (13 out of 73 videos): This bat species exhibits notable curiosity and playful behavioral

traits. Various play behaviors were observed and categorized as object play (n= 4), social play

(n=5), or food play (n=4). One individual was observed interacting with the wildlife camera,

occasionally altering its position while emitting vocalizations which we categorized as object

play. In other videos, bats were attempting to catch passing cockroaches, classified as food

play. Additional footage displayed playful interactions between two bats, characterized by

playful fighting, which we categorized as social play.

Figure 1: Roost with the four *Vampyrum spectrum* individuals (presumably male, female, and

two pups).

Prey determination

In 30 videos, a bat carrying and/or eating a prey item was observed. In 50% of the instances,

the prey was a small mammal, primarily mice and rats. In 30% of the cases, the prey was

visually identified as a bird. In 20% of the cases, the prey item could not be identified due to

low video quality and/or the bats being too far from the camera.

Entering/leaving the roost

A significant portion of the videos (203 in total) captured by the camera showed the bats

entering and leaving the roost, as this is when they passed the camera and triggered the

sensor. Although it appears that bats usually foraged alone, on 40 occasions, at least two

bats were observed leaving or returning to the roost together, sometimes even all four bats.

In six instances, two bats were observed leaving the roost simultaneously and returning

together after an average of about 40 minutes.

Figure 2: Scatter plot illustrating the temporal occurrence of the different observed social

behaviors in a *V. spectrum* night-roost between 5 pm and 6 am (N= 60 nights).

Figure 3: A + B: Illustrations of prey sharing behavior between two bats; C: Spaghetti plot

illustrating the difference of eye distances between the prey carrying bat and the

approaching bat, showing a significant difference in eye distance (as a proxy for body size).

Eye distance measurements

We detected a clear difference in eye distance between bats involved in prey sharing

interactions. The bat that brought the prey into the roost had a significantly larger eye

distance than the bat receiving the prey (paired Wilcoxon test, Z= -2.527; n= 8, exact p:

0.0078; Fig. 4). Assuming that eye distance can be used as a proxy for body size, our data

suggests that the bat sharing prey is larger than the bat receiving prey, which would be

expected if adults share prey with their offspring.

Discussion

Here, we provide the first comprehensive account of prey sharing and other social behaviors

in the spectral bat *V. spectrum*. Prey sharing was a clearly cooperative social behavior

wherein a bat successfully captured prey, brought it to the roost where group members

were present, and willingly transferred the prey to another bat. Our video observations

consistently showed that upon entering the roost with prey, the carrying bat was often

approached by another bat attempting to acquire the prey. When other bats were present,

they typically remained calm and inactive, suggesting they recognized the intended recipient

of the prey, often a bat already positioned to receive it. In most instances of prey sharing,

the providing bat voluntarily and promptly handed over the prey to the receiving bat.

Occasionally, conflicts over prey arose, as evidenced by one video depicting a tug-of-war

scenario where an approaching bat successfully wrestled the prey away. Previous

observations of a *V. spectrum* pair with a pup in captivity suggest that male bats may

occasionally bring prey to the roost, potentially to provision females (Bradbury, personal

observation via (43). This behavior aligns with an anecdotal observation from December

2024, where one bat, presumably the lactating female, remained in the roost with the pup

while two other bats were out foraging. The prey transfer between the prey-carrying bat and

the presumed mother, which was remarkably silent comparted to other prey transfers that

were accompanied by social vocalizations supports the hypothesis of males provisioning

females under certain circumstances.

To date, there has been scarce evidence of prey sharing among bats. In the closely related

sister species *Chrotopterus auritus,* individuals reportedly bring back prey into the roost and

then share it with the other members (Rodrigo Medellín, personal communication, (46). Two

studies in captivity provide additional evidence for parental food sharing in carnivorous

species. In the fish-eating *Noctilio albiventris*, (47) observed that just prior to weaning at

three months, captive mothers fed juveniles with masticated fish or mealworms from their

cheek pouches. Similarly, in *Megaderma lyra*, mothers reportedly transferred either entire

or partly consumed frogs to their young. These food transfers ceased when the young bats

reached approximately 74 days of age (48). It appears that lactating females of *M. lyra*

provision their offspring by supplementing milk with solid food, akin to the behavior

hypothesized for *V. spectrum* (43). The only well-documented instance of prey-sharing

behavior observed in the wild among bats occurs in *Micronycteris microtis*. In this species,

weaned pups are provisioned with large insects by their mothers for an additional five

months. This extended provisioning likely helps pups learn to handle challenging prey and

refine the acoustic skills essential for effective hunting. Although *M. microtis* primarily

forages on insects, it has a remarkably broad diet that also includes lizards, making it the

smallest known carnivorous bat to date (49, 50). Another prominent example of food

sharing behavior in bats is the well-studied regurgitation of blood in vampire bats; however,

here, food is shared between kin as well as non-kin group members that are in critical

physical conditions (51–53), rather than on a regular basis and/or for learning purposes.

Nectar-feeding bats also regurgitate food, but only for their own offspring, presumably to

substitute their diet on their way to independence (54).

Extended pup dependency & presumed biparental care

Observations of a captive *V. spectrum* pair with a pup suggest that males may occasionally

bring prey to the roost to provision females or nourish young pups not yet fully capable of

independent hunting (Bradbury, via (43)). During the nursing period, pups likely rely on milk

supplemented with meat, which facilitates their transition to a carnivorous diet and helps

them develop prey-handling skills (43). Our data supports this strategy as a way to ensure

adequate nutrition and skill development in young bats. Based on eye-distance

measurements, two smaller individuals in our roost likely represent pups from different

years, as twins have never been reported in *V. spectrum*. Pups are reported to be born at the

end of the dry season (42, 43) and the roost with these four bats was already discovered by

us in November 2022, so the younger offspring must have been born at least 1.5 years ago,

and the older one approximately 2.5 years ago. This indicates a prolonged period of parental

care, which is relatively uncommon in bats. This extended investment is likely tied to the

species' socially monogamous structure, a rarity among mammals (55, 56) and to the

challenges associated with carnivory. Parental investment, defined as post-birth behaviors

that enhance offspring reproductive success (57), is influenced by the costs and benefits to

parents (58). In species with low mate competition and high paternity certainty, such as *V.*

spectrum, males are more likely to invest in parental care. This aligns with the species' low

population density, large territory requirements, and limited roost availability in dry forests

(59). Biparental care, hypothesized by Vehrencamp et al. (43), likely enhances offspring

survival and success compared to uniparental care (60). While biparental care is common in

90% of bird species (61), it is rare in mammals, where females alone provide care in 90% of

species (58). Notably, some of the social behaviors of the spectral bat seem to have quite

some commonalties with the big-eared woolly bat (*Chrotopterus auritus)* that also exhibits a

socially monogamous social group structure, shares a similar habitat and comparable diet

(62, 63). Therefore, it can be surmised that similar factors act on *C. auritus* and it would be

very likely that this species has also evolved some kind of biparental care (46) .

Prey items and Foraging Dynamics

In addition to prey-sharing behavior, several videos documented bats consuming prey within

the roost. Due to the 1-minute recording limitation, it was often unclear whether the prey

was brought into the roost by the bat itself or transferred prior to the recording. However,

some videos showed bats entering with prey and consuming it alone without sharing. In one

instance, a bat carrying prey crawled to its roosting spot and ate it despite a nearby bat

approaching and gently stretching its head toward the prey, possibly begging. The prey-

carrying bat appeared undisturbed and continued eating. Vehrencamp et al. (43) proposed

that bats might bring prey to the roost when caught late at night, close to sunrise, to avoid

daylight exposure. However, our observations recorded such behavior at various times

during the night, suggesting that when prey is captured near the roost, it provides a

convenient, safe location for consumption. Visual classifications identified small rodents as

the primary prey brought back to the roost, with only 9 of 30 videos showing birds being

brought in. A study in 2022 based on fecal sample analysis from a *V. spectrum* roost in

361 Nicaragua found birds to be the most common prey group, followed by bats and then small

362 rodents (59). Interestingly, no videos documented *V. spectrum* bringing smaller bat species

363 back to the roost, suggesting that bats may be hunted midair, unlike diurnal birds, which are

364 likely located in their nests through olfaction (42, 43). While previous studies reported *V.*

365 *spectrum* foraging solitarily, with individuals leaving and returning to the roost separately

366 (43), our data recorded 40 instances where two or more bats left or returned

367 simultaneously, including 6 cases where two bats left and returned together after

368 approximately 40 minutes. These observations suggest that while *V. spectrum* typically

369 forages alone, exceptions occur, particularly when subadults may forage with adults to

370 practice and develop hunting skills.

372 Further social behaviors

373 Play behavior may contribute to shaping adult skill sets. Observed examples in our videos

374 include chasing cockroaches, exploring wildlife cameras, and play-fighting, likely among

375 subadults. Play is often viewed as a "non-serious" behavior, where actions are performed for

376 their own sake rather than practical outcomes. This allows youngsters to explore

377 environments and develop responses in a low-risk manner (64, 65). Play and exploration are

378 documented in humans and non-human animals, including birds, frogs, and cephalopods,

379 but are most prevalent in mammals, especially those with extended parental care (66, 65).

380 Species with greater parental care often exhibit more play, which helps juveniles develop

381 adaptive behaviors for current and future challenges (67). Play may also influence gene

382 expression and evolution indirectly (65). Our data highlights the presence of play behavior in

383 bats, emphasizing its potential developmental importance, despite limited research to date.

The pronounced social roosting behaviors of *V. spectrum*, including hugging-like actions—

characterized by embracing, wrapping, or holding closely with limbs, wings, or body—and

greeting interactions, were observed daily. Most bats are highly social, roosting in close

contact and relying on strong bonds for survival (8). Such bonds may offer cooperative

benefits, such as information transfer within the group. Similar to allogrooming, a behavior

that helps maintain social bonds in dynamic systems (31, 68), these close-contact behaviors

in *V. spectrum* likely reinforce social ties. While all bats exhibit autogrooming, allogrooming

is less common (69–71). In *V. spectrum*, hugging and greeting likely strengthen bonds within

their small social groups, which is crucial for both parent-offspring and male-female

relationships.

Conclusion

In conclusion, our findings revealed that prey sharing in *V. spectrum* is a common and mostly

voluntary behavior, with bats bringing prey back to the roost and handing it over to other

roost-mates, particularly those that may be less capable of hunting for themselves. This

behavior supports the hypothesis that adults assist in transitioning young bats from milk to a

carnivorous diet, ensuring adequate food intake and offering them the opportunity to

practice how to handle large prey items. This study contributes to the broader

understanding of bat sociality by highlighting the unique cooperative behaviors of *V.*

spectrum and provides a foundation for further research into the ecological and evolutionary

implications of these behaviors. The insights gained from this study not only enhance our

knowledge of *V. spectrum* but also contribute to the general understanding of social

evolution and cooperative behaviors in mammals.

Acknowledgments

We would like to express our thanks to all the workers from "La Estación Experimental

Forestal Horizontes", especially Marlon Gonzalez. We are also grateful to Emma Dittrich for

her drawings.

Roles of contribution:

Marisa Tietge: Discoverer of the roost, Conceptualization, Methodology, Investigation,

Formal analysis, Data curation, Visualization, Project administration, Writing- Original draft

preparation, Writing- Reviewing and Editing

Mirjam Knörnschild: Study Design, Supervision, Writing- Reviewing and Editing

Eduardo Artavia Durán: Local cooperation partner, Writing- Reviewing and Editing

Competing interests: The authors state no conflict of interest. The authors have no financial

or proprietary interests in any material discussed in this article.

Research funding: This work was supported by a grant from the Leibniz Foundation

(P122/2020) to M.K.

Supplementary material

Example videos from above-described key social behaviors:

- Prey sharing behavior (small mammal): <u>Vampyrum spectrum prey sharing</u>

<u>mammal.mp4</u>

- Prey sharing behavior (bird): <u>Vampyrum_spectrum_prey_sharing_bird.mp4</u>

- Fight about food: <u>Vampyrum spectrum_fight (over food).mp4</u>

- Social roosting: <u>Vampyrum_spectrum_social roosting.mp4</u>

- Greeting behavior: <u>Vampyrum_spectrum_greeting_behavior.mp4</u>

- Play (object): <u>Vampyrum_spectrum_play (object).mp4</u>

- Play (food): <u>Vampyrum_spectrum_play (food).mp4</u>

- Play (social): <u>Vampyrum_spectrum_play (social).mp4</u>

Literature Cited

1. Buchan JC, Alberts SC, Silk JB, Altmann J. True paternal care in a multi-male primate
society. Nature 2003; 425(6954):179–81.

2. Engh AL, Siebert ER, Greenberg DA, Holekamp KE. Patterns of alliance formation and postconflict aggression indicate spotted hyaenas recognize third-party relationships. Animal Behaviour 2005; 69(1):209–17.

3. Komdeur J, Eikenaar C, Brouwer L, Richardson DS. The Evolution and Ecology of Cooperative Breeding in Vertebrates. In: The Evolution and Ecology of Cooperative Breeding in Vertebrates. Wiley; 2008.

4. Krams I, Krama T, Igaune K, Mänd R. Experimental evidence of reciprocal altruism in the pied flycatcher. Behav Ecol Sociobiol 2008; 62(4):599–605.

5. Nowak MA. Evolving cooperation. J Theor Biol 2012; 299:1–8.

6. Dugatkin LA. Cooperation in animals: An evolutionary overview. Biol Philos 2002; 17(4):459–76.

7. Lacy EA, Sherman PW. Cooperative breeding in mammals: Cooperative Breeding In Naked Mole-Rats: Implications for Vertebrate and Invertebrate Sociality. Cambridge: Cambridge University Press; 2007.

8. Carter GG, Wilkinson GS. Food sharing in vampire bats: reciprocal help predicts donations more than relatedness or harassment. Proc. R. Soc. B. 2013; 280(1753):20122573.

9. Connor RC, editor. Group living in whales and dolphins. Chicago: The University of Chicago Press; 2000. (Cetacean societies: field studies of dolphins and whales).

10. Bailey I, Myatt JP, Wilson AM. Group hunting within the Carnivora: physiological, cognitive and environmental influences on strategy and cooperation. Behav Ecol Sociobiol 2013; 67(1):1–17.

11. Nichols TC. Cooperative hunting of Canada Geese (*Branta canadensis*) by Gray Wolves (*Canis lupus*) in northern Quebec. Can Field Nat 2015; 129(3):290.

12. Harcourt A, De Waal FB. Coalitions and alliances in humans and other animals [Alexander Harcourt and Frans B.M. de Waal]. American J Phys Anthropol 1993; 1992(4):533–5.

13. Dugatkin LA. The Evolution of Cooperation. BioScience 1997; 47(6):355–62.

14. Simmons NB, Cirranello AL. Bat Species of the World: A taxonomic and geographic database; 2024 [cited Accessed on 2024 Dec 10]. Available from: URL: https://batnames.org.

15. Kunz T. H, Fenton M. B. Bat Ecology. Journal of Mammalogy 2004; 85(2):366–7.

16. Altringham JD. Bats: From evolution to conservation. 2nd ed. New York: Oxford University Press; 2011.

17. McCracken GF, Wilkinson GS. Bat Mating Systems. In: McCracken GF, Wilkinson GS, editors. Reproductive Biology of Bats. Elsevier; 2000. p. 321–62.

18. Kerth G. Causes and Consequences of Sociality in Bats. BioScience 2008; 58(8):737–46.

19. Chaverri G, Ancillotto L, Russo D. Social communication in bats. Biol Rev Camb Philos Soc
2018; 93(4):1938–54.

20. Barbour T. A Peculiar Roosting Habit of Bats. The Quarterly Review of Biology 1932;
7(3):307–12.

21. Kunz TH, editor. Ecology of Bats. Boston, MA: Springer US; 1982.

22. Chaverri G, Kunz TH. Ecological Determinants of Social Systems. In: Behavioral ecology of
tropical animals. Elsevier; 2010. p. 275–318 (Advances in the Study of Behavior).

23. Gaudet CL, Fenton MB. Observational learning in three species of insectivorous bats
(Chiroptera). Animal Behaviour 1984; 32(2):385–8.

24. Ratcliffe JM, Hofstede HM ter. Roosts as information centres: social learning of food
preferences in bats. Biol Lett 2005; 1(1):72–4.

25. Page RA, Ryan MJ. Social transmission of novel foraging behavior in bats: frog calls and
their referents. Curr Biol 2006; 16(12):1201–5.

26. Dechmann DKN, Heucke SL, Giuggioli L, Safi K, Voigt CC, Wikelski M. Experimental
evidence for group hunting via eavesdropping in echolocating bats. Proc. R. Soc. B. 2009;
276(1668):2721–8.

27. Dechmann DKN, Kranstauber B, Gibbs D, Wikelski M. Group hunting-a reason for sociality
in molossid bats? PLoS One 2010; 5(2):e9012.

28. Wright GS, Wilkinson GS, Moss CF. Social Learning of a Novel Foraging Task by Big Brown
Bats (Eptesicus fuscus). Animal Behaviour 2011; 82(5):1075–83.

29. O'Mara MT, Dechmann DKN, Page RA. Frugivorous bats evaluate the quality of social
information when choosing novel foods. Behavioral Ecology 2014; 25(5):1233–9.

30. Chaverri G, Gillam EH, Kunz TH. A call-and-response system facilitates group cohesion
among disc-winged bats. Behavioral Ecology 2013; 24(2):481–7.

31. Dunbar R, Shultz S. Bondedness and sociality. Behav 2010; 147(7):775–803.

32. Wilkinson GS. Information transfer at evening bat colonies. Animal Behaviour 1992;
44:501–18.

33. Wilkinson GS, Boughman JW. Social calls coordinate foraging in greater spear-nosed bats.
Animal Behaviour 1998; 55(2):337–50.

34. Willis CKR, Brigham RM. Social thermoregulation exerts more influence than
microclimate on forest roost preferences by a cavity-dwelling bat. Behav Ecol Sociobiol 2007;
62(1):97–108.

35. Ortega J, Guerrero JA, Maldonado JE. Aggression and Tolerance by Dominant Males of
Artibeus jamaicensis : Strategies to Maximize Fitness in Harem Groups. Journal of
Mammalogy 2008; 89(6):1372–8.

36. Nagy M, Knörnschild M, Voigt CC, Mayer F. Male greater sac-winged bats gain direct
fitness benefits when roosting in multimale colonies. Behavioral Ecology 2012; 23(3):597–
606.

37. Gager Y. Information transfer about food as a reason for sociality in bats. Mammal
Review 2019; 49(2):113–20.

38. Ripperger SP, Stockmaier S, Carter GG. Tracking sickness effects on social encounters via
continuous proximity sensing in wild vampire bats. Behavioral Ecology 2020; 31(6):1296–
302.

39. Moreno KR, Weinberg M, Harten L, Salinas Ramos VB, Herrera M LG, Czirják GÁ et al. Sick
bats stay home alone: fruit bats practice social distancing when faced with an immunological
challenge. Ann N Y Acad Sci 2021; 1505(1):178–90.

40. McWilliam AN. Emergence behaviour of the bat Tadarida (Chaerephon) pumila
(Chiroptera: Molossidae) in Ghana, West Africa. Journal of Zoology 1989; 219(4):698–701.

41. Frick WF, Kingston T, Flanders J. A review of the major threats and challenges to global
bat conservation. Ann N Y Acad Sci 2020; 1469(1):5–25.

42. Greenhall AM. Notes on the Behavior of the False Vampire Bat. Journal of Mammalogy
1968; 49(2):337.

43. Vehrencamp SL, Stiles FG, Bradbury JW. Observations on the Foraging Behavior and Avian
Prey of the Neotropical Carnivorous Bat, Vampyrum spectrum. Journal of Mammalogy 1977;
58(4):469–78.

44. Solari S. Vampyrum spectrum. The IUCN Red List of Threatened Species 2018; 2018.

45. Gual-Suárez F, Medellín RA. We eat meat: a review of carnivory in bats. Mammal Review
2021; 51(4):540–58.

46. Trujillo LA, Gual-Suárez F, Trujillo RE, Medellín RA. Arachnids that feed on vertebrate
carrion: necrophagy by the whip spider Paraphrynus raptator (Amblypygi: Phrynidae) and its
relation to the feeding behavior of the woolly false vampire bat Chrotopterus auritus
(Chiroptera: Phyllostomidae). The Journal of Arachnology 2021; 49(3).

47. Brown PE, Brown TW, Grinnell AD. Echolocation, development, and vocal communication
in the lesser bulldog bat, Noctilio albiventris. Behav Ecol Sociobiol 1983; 13(4):287–98.

48. Raghuram H, Marimuthu G. Maternal feeding of offspring with vertebrate prey in captive
Indian false vampire bat, Megaderma lyra. Acta Chiropterologica 2007; Volume 9(2):437-
443(7).

49. Santana SE, Geipel I, Dumont ER, Kalka MB, Kalko EKV. All you can eat: high performance
capacity and plasticity in the common big-eared bat, Micronycteris microtis (Chiroptera:
Phyllostomidae). PLoS One 2011; 6(12):e28584.

50. Geipel I, Jung K, Kalko EKV. Perception of silent and motionless prey on vegetation by
echolocation in the gleaning bat Micronycteris microtis. Proc. R. Soc. B. 2013;
280(1754):20122830.

51. Wilkinson GS. Reciprocal food sharing in the vampire bat. Nature 1984; 308(5955):181–
4.

52. Elizalde-Arellano C, López-Vidal JC, Arroyo-Cabrales J, Medellín RA, Laundré JW. Food
sharing behavior in the hairy-legged vampire bat Diphylla ecaudata. Acta Chiropterologica
2007; 9(1):314–9.

53. Wilkinson GS, Carter GG, Bohn KM, Adams DM. Non-kin cooperation in bats. Philos Trans
R Soc Lond B Biol Sci 2016; 371(1687):20150095.

54. Rose A, Wöhl S, Bechler J, Tschapka M, Knörnschild M. Maternal mouth-to-mouth
feeding behaviour in flower-visiting bats, but no experimental evidence for transmitted
dietary preferences. Behav Processes 2019; 165:29–35.

55. Orians GH. On the Evolution of Mating Systems in Birds and Mammals. The American
Naturalist 1969; 103(934):589–603.

56. Alexander RD. The Evolution of Social Behavior. Annu. Rev. Ecol. Syst. 1974; 5(1):325–83.

57. Klug, Hope, Alonzo SH, Bonsall, Michael B. Theoretical foundations of parental care 2012.

58. Clutton-Brock TH. The evolution of parental care. Princeton, N.J.: Princeton University
Press; 1991. (Monographs in behavior and ecology).

59. Martínez-Fonseca JG, Mau R, Walker FM, Medina-Fitoria A, Yasuda K, Chambers CL.
Vampyrum spectrum (Phyllostomidae) movement and prey revealed by radio-telemetry and
DNA metabarcoding. PLoS One 2022; 17(4):e0265968.

60. Barta Z, Székely T, Liker A, Harrison F. Social role specialization promotes cooperation
between parents. The American Naturalist 2014; 183(6):747–61.

61. Cockburn A. Prevalence of different modes of parental care in birds. Proc. R. Soc. B.
2006; 273(1592):1375–83.

62. Medellín RA. Chrotopterus auritus. Mammalian species 1989; 343:1–5.

63. Esbérard CEL, Motta AG, Almeida JC, Ferreira LCS, Costa LM. Reproduction of
Chrotopterus auritus (Peters) in captivity (Chiroptera, Phyllostomidae). Braz J Biol 2006;
66(3):955–6.

64. Spinka M, Newberry RC, Bekoff M. Mammalian play: training for the unexpected. The
Quarterly Review of Biology 2001; 76(2):141–68.

65. Burghardt GM. The Genesis of Animal Play. The MIT Press; 2005.

66. Zylinski S. Fun and play in invertebrates. Curr Biol 2015; 25(1):R10-2.

67. Bjorklund D. Mother knows best: Epigenetic inheritance, maternal effects, and the
evolution of human intelligence. Developmental Review 2006; 26(2):213–42.

68. Russell YI, Phelps S. How do you measure pleasure? A discussion about intrinsic costs
and benefits in primate allogrooming. Biol Philos 2013; 28(6):1005–20.

69. Wilkinson GS. Social grooming in the common vampire bat, Desmodus rotundus. Animal
Behaviour 1986; 34(6):1880–9.

70. Fleming TH, Nelson AA, Dalton VM. Roosting Behavior of the Lesser Long-Nosed Bat,
Leptonycteris curasoae. Journal of Mammalogy 1998; 79(1):147–55.

71. Ortega J, Maldonado JE. Female interactions in harem groups of the Jamaican fruit-
eating bat, Artibeus jamaicensis (Chiroptera: Phyllostomidae). Acta Chiropterologica 2006;
8(2):485–95.