Cooperative Behaviors and Social Interactions in the Carnivorous Bat *Vampyrum spectrum*

Marisa Tietge[1,2] *, Eduardo Artavia Durán[3], Mirjam Knörnschild[1,2,4]

[1] Museum for Natural History, Leibniz Institute for Evolution and Biodiversity Science, 10115 Berlin, Germany

[2] Institute for Biology, Humboldt-Universität zu Berlin, 10099 Berlin, Germany

[3] Manzú Conservación de Murciélagos, 50102, Costa Rica

[4] Smithsonian Tropical Research Institute, 0843-03092, Panama

*Corresponding author: Marisa Tietge

Keywords: Biparental care; Chiroptera; Food provisioning; Neotropical; Phyllostomidae; Prey sharing; Roost; Social behavior

Highlights

Extensive parental care: Most likely both male and female spectral bats participate in parental care, e.g. food provisioning, a rare behavior among mammals

Prey sharing as a social cooperative behavior: Detailed observations of prey sharing within the roost, highlighting social cooperation and resource allocation

Extended dependency period for pups: A prolonged developmental period allows pups to receive care and acquire essential survival skills until they achieve full independence

Juvenile play: Pups engaged predominantly in social play (mock fights), but also played with objects or food items, a rare behavior among bats

Abstract

Bats exhibit a diverse array of social behaviors, yet detailed studies on the intricacies

of these interactions, particularly in rare species like the spectral bat (*Vampyrum*

spectrum), remain scarce. This study presents the first comprehensive description of

prey sharing and other social behaviors in a wild social group of *V. spectrum*. Over

several months, we conducted extensive video recordings in a hollow tree, used as a

day- and night-roost, to document these behaviors, aiming to elucidate key social

interactions among the bats and their role within their ecological niche. We observed

various remarkable social behaviors, including food sharing between family

members. Our findings support the hypothesis that prey sharing as a form of

biparental care may serve as a method for adults to transition young bats from milk

to a carnivorous diet, ensuring adequate food intake and allowing them to practice

how to handle large prey items. We also challenge the notion that *V. spectrum*

forages exclusively solitarily, as we documented several instances of synchronized

roost departures and returns and thus presumably cooperative foraging. This

indicates a more complex social structure and behavioral ecology than previously

understood. Our comprehensive analysis of observational data enhances our

understanding of the social dynamics of *V. spectrum*, providing new insights into the

evolution of cooperative behaviors in bats.

Introduction

Cooperative behaviors are observed across many animal species and play a crucial

role in their survival and social structures (1–3). Such behaviors often involve

individuals working together to achieve shared goals, like defending territories,

caring for young, or securing resources (4, 3, 5). These interactions are especially

well-documented in social animals, where cooperation can range from kinship-based

behaviors to complex alliances among unrelated individuals (6–8). In addition to

kinship, alliances among non-relatives are widespread; for instance, in species like

baboons and dolphins, unrelated individuals cooperate to hunt more effectively,

which enhances foraging success (9–12). This behavior allows access to larger or

more challenging prey than would be possible alone, illustrating how collaborative

behaviors can enhance survival and fitness within an ecological niche (13).

Bats, with over 1,480 species (14), exhibit cooperative behaviors on many levels. As

one of the most ecologically significant and diverse orders of mammals, bats occupy

a broad range of habitats and exhibit various dietary preferences – including

insectivory, frugivory, nectarivory, piscivory, and even carnivory (15, 16). Their social

structures are equally varied, spanning solitary roosting, complex communal

arrangements, such as harems, social monogamy, and promiscuity (17, 18), making

them an exceptional case for studying the dynamics of cooperation across different

social systems. Most bat species are highly gregarious, presumably due to

ecological constraints like roost limitation, physiological demands like social

thermoregulation, and their longevity (18). Bats exhibit complex communication,

group-level recognition, and various other social interactions among individuals (19).

Some remarkable social behaviors have evolved, like roost sharing and defense in

tent-making bats where individuals construct tents from large leaves. These shelters

provide protection from predators and harsh weather. Bats within these groups

cooperate in the maintenance and defense of their roosts (20–22). Previous studies

have shown that bats from various social systems can show cooperative behavior

when it comes to the availability of food and prey capture using auditory, visual, and

olfactory cues, especially when living in living in fission-fusion systems (23–29).

Behavioral evidence also suggests that bats use calls to distinguish between familiar

and unfamiliar conspecifics, like the Spix's disk-winged bats (*Thyroptera tricolor*) that

uses social calls to stay in touch with group members while foraging (30). This

behavior is similar to "grooming at a distance" in primates, where individuals that

frequently groom each other maintain vocal contact when separated during foraging

(reviewed in (31). Maintaining contact while foraging may help bats ensure they later

roost with preferred group members and reinforce cooperative behavior (30).

Understanding the social behaviors of bats is crucial for several reasons. Social

interactions can impact reproductive success, foraging efficiency, predator

avoidance, and disease transmission (32–40). These behaviors are also integral to

the conservation and management of bat populations, many of which are threatened

by habitat loss, climate change, and human activities (41). Bat species developed a

lot of different social behaviors, most of the them are still unknown because bats are

often hard to observe due to their nocturnal and highly mobile lifestyle. Especially bat

species that are rare to find and/or are critically endangered are understudied.

The spectral bat (*Vampyrum spectrum*) is widely recognized as the largest

carnivorous bat species in the New World, with individuals weighing approximately

180 grams and boasting a wingspan of up to 900 millimeters (42). This rare species

can be found in lowland tropical dry forests, evergreen forests, and occasionally in

cloud forests, deciduous forests, or swampy areas (43). *Vampyrum spectrum* is

considered near threatened by the International Union for Conservation of Nature

(IUCN) and is listed as a species of special concern or endangered in several

countries throughout its range with a decreasing population trend (44). Belonging to

the family of leaf-nosed bats (Phyllostomidae), these formidable predators have a

diverse diet that includes a variety of birds, rodents, and small mammals, including

other bat species (42, 45). *Vampyrum spectrum* typically roosts in small groups

within hollow trees or caves. These groups exhibit a socially monogamous structure,

usually comprising a single male, a single female, and their recent offspring who

have not yet dispersed (43). This socially monogamous social structure is relatively

rare among mammals, making this species a particularly interesting subject for

study.

The primary goals of this study were to get insights into the social behaviors of

Vampyrum spectrum within the context of their ecological niche as carnivorous bats.

By conducting extensive video recordings in their roost, we aimed to document and

analyze key behaviors, with a focus on social interactions and cooperative

behaviors. This research not only aims to shed light on the social behaviors of *V.*

spectrum but also contributes to the broader understanding of social behavior in

bats.

Methods

Study site

The study was conducted in the tropical dry forest of "La Estación Experimental

Forestal Horizontes" (N 10° 42' 47.883973, E -85° 35' 43.759147) in Guanacaste,

Costa Rica. The focal roost is located within a partially hollow yet living tree of the

species *Manilkara chicle*, amidst typical dry forest vegetation and adjacent to a dried

riverbed. This roost was first identified by M.T. in December 2022, housing four *V.*

spectrum individuals (Fig. 1), and we began systematically monitoring them

approximately one year later. The *V. spectrum* bats occupy the highest section of the

hollow tree, while a colony of 15 smaller bats, *Saccopteryx bilineata,* inhabits a lower

section near the entrance of the tree hole. Although we did not individually mark the

four *V. spectrum* bats we are confident that this colony consisted of a female and

male bat pair and their respective two young for several reasons: 1) this species

lives in a socially monogamous social structure that either constitutes a single

roosting bat or family group (42), 2) the bats in the video displayed noticeable size

differences and the teats of the adult, post-lactating female were visible on occasion,

and 3) additional video recordings in 2024 showed a newly born pup being nursed by

its mother and made it possible to identify the other large individual as male

(because the penis was visible).

Data collection

At the beginning of the dry season in November 2023, a wildlife camera (Snapshot

Mini Black 30MP 4K, Dörr GmbH) was installed inside the roost at a height of

approximately 10 meters, using a small camera mount (CAMVATE photography

mount with 1/4"-20 thread). The camera was positioned about 2 meters from the

primary roosting site of the bats, oriented upwards to capture optimal footage from

below. To ensure minimal disturbance to the bats, the camera was carefully installed

during a period of low activity. Observations following the installation indicated that

the bats' behavior remained unaffected by the presence of the equipment. Over the

course of three months, the camera was regularly checked, cleaned, equipped with a

new SD card, and fitted with fresh batteries before being reinstalled at the fixed

location within the tree. The camera recorded automatically triggered infrared videos

with a maximum duration of one minute. Initially, continuous recording was achieved

each night; however, as battery power depleted, the camera's built-in power-saving

function reduced the length of recorded videos.

Over the course of three months, we video-recorded the four bats for a total of 60

days. On some days, only a few videos were captured, while on other days, up to 40

videos were recorded, depending on how frequently the camera sensor was

triggered by bat activity. In total, 502 videos were collected. Most of these videos

(203 out of 502) were triggered by bats either leaving or entering the roost. The other

bat species, *S. bilineata,* was very seldom recorded, as the camera was installed

above their roosting spot. Out of the 502 recorded videos, 73 contained social

behaviors or other interesting activity. By visual examination of the videos, we

classified eight different behavioral categories: "social roosting", "greeting behavior",

"presumably sexual behavior", "bringing prey into the roost", "eating a prey item",

"prey sharing", "food checks", and "play behavior". All observed behaviors were

recorded during the night between 5pm and 6am and occurred with no specific

distribution pattern (see Fig. 2). In order to ensure long-term monitoring of this social

group, further videos are recorded in the colony in 2024/2025. All data presented in

this study are based on video recordings collected during the 2023/2024 season.

However, one additional video from the 2024/2025 season was included as an

anecdotal observation due to its particularly interesting and relevant nature.

Individual distinction by size estimation

Due to the absence of individual markings on the animals, we employed

supplementary methods to support our observations and interpretations of

behaviors, particularly in prey-sharing interactions. We measured the distance

between the eyes of the bat entering the roost with prey and the bat approaching to

take over the prey. This was done using screenshots from video frames capturing

prey-sharing behaviors, where both animals were positioned at approximately the

same height and their eyes were clearly visible (N=8 prey sharing interactions). The

screenshots were analyzed in GIMP (GIMP 2.10.38), utilizing the tape measure

function to determine the eye distance in pixels. These measurements were

subsequently compared using a non-parametric paired samples Wilcoxon test.

Additionally, we used a spaghetti plot to visualize the results (see figure 4).

All statistical analyses were performed using the program Sypder (Spyder IDE, The

Scientific Python Development Environment, Python 3.12.4)

Results

Based on the 73 infrared videos, we created an ethogram that contained eight

different behavioral categories exhibited by *V. spectrum* bats in their roost. Video

examples of the respective behaviors can be found in the electronic supplementary

material.

Social roosting (16 out of 73 videos): This behavior involves two or more bats

roosting in close proximity with body contact, establishing a ball-like formation. At

least one bat wraps its wings around the other bat(s). This behavior is often

accompanied by allogrooming and/or social vocalizations and was observed very

frequently.

Greeting Behavior (4 out of 73 videos): This behavior constitutes of a hugging-like

interaction between a bat already in the roost and a newly arrived bat. The resident

bat may actively approach or greet the newcomer as it reaches close proximity in the

main roosting area. The greeting behavior is comparable to the initiation to social

roosting, where at least one bat wraps its wings around the other, establishing a ball-

like formation for several seconds. This behavior may be accompanied by social

vocalizations.

Presumably sexual behavior (2 out of 73 videos): This behavior involves two bats,

with one bat—presumably the male—positioned behind the female, aligning belly to

back while wrapping its wings around her. This interaction often includes frequent

position changes and licking of various lower body parts. This kind of behavior was

only recorded twice and therefore we can only assume that this observed behavior is

indeed produced in a sexual context.

Bringing prey into the roost (4 out of 73 videos): A bat enters the roost after foraging,

carrying a prey item in its mouth (normally a bird or rodent). The prey is always held

by its head and appears already dead. The bat carries the prey item in its mouth and

crawls backward to the main roosting area in the hollow tree, with only the back or

tail of the prey visible.

Eating a prey item (18 out of 73 videos): A bat suspends itself at a specific location

within the roost, positioning the prey item between its wings and securing it with its

thumb claws while biting into it. Audible chewing noises are a distinctive feature of

this process. We estimate that consuming a single prey item may takes

approximately 30-40 minutes, based on sequential video recordings showing bats

feeding on the same prey item. Nevertheless, this duration remains unverified due to

the current limitation of our video equipment, which recorded only 1-minute

segments. A video sequence was classified as "eating a prey item" when there was

no immediate social interaction with another bat preceding or following the behavior.

In 12 videos, we could not observe the bat entering the roost, only the eating

process was recorded, while in six videos a bat entered the roost with prey, chose

not to share it, and consumed it alone.

Prey sharing (12 out of 73 videos): This behavior involves a bat entering the roost

with a prey item held in its mouth. In many instances, as soon as the prey-carrying

bat climbs to the main roosting spot, normally only one other bat approaches while

emitting vocalizations. The approaching bat comes very close to the prey-carrying

bat, typically from the back and/or side. The approaching bat attempts to bite into the

upper body of the prey. In most cases (92%), the prey-carrying bat voluntarily

releases the prey item quickly. Subsequently, both bats crawl up to the main roosting

location where the bat with the prey usually begins to consume the prey item

immediately (see Fig. 3). On a single occasion, a recorded interaction between two

bats involved a fight over prey brought back into the roost, resembling a tug-of-war

scenario. Additionally, we recorded an anecdotal observation in a video from

December 2024, where a bat—presumed to be the mother—was present in the roost

alongside a small pup when another bat entered carrying prey. The mother

approached the incoming bat, swiftly and silently took the prey, and then returned to

the primary roosting spot next to the pup with the prey to consume it.

Food checks (4 out of 73 videos): One bat enters the roost, and another bat

approaches, seemingly to check if the entering bat has brought back food from

foraging. The approaching bat displays behaviors similar to prey-sharing

interactions, though in these cases, no prey is present. After realizing this, both bats

typically crawl up to the main roosting spot.

Play (13 out of 73 videos): This bat species exhibits notable curiosity and playful

behavioral traits. Various play behaviors were observed and categorized as object

play (n= 4), social play (n=5), or food play (n=4). One individual was observed

interacting with the wildlife camera, occasionally altering its position while emitting

vocalizations which we categorized as object play. In other videos, bats were

attempting to catch passing cockroaches, classified as food play. Additional footage

displayed playful interactions between two bats, characterized by playful fighting,

which we categorized as social play.

Figure 1: Roost with the four Vampyrum spectrum individuals (presumably male,

female, and two pups).

Prey determination

In 30 videos, a bat carrying and/or eating a prey item was observed. In 50% of the

instances, the prey was a small mammal, primarily mice and rats. In 30% of the

cases, the prey was visually identified as a bird. In 20% of the cases, the prey item

could not be identified due to low video quality and/or the bats being too far from the

camera.

Entering/leaving the roost

A significant portion of the videos (203 in total) captured by the camera showed the

bats entering and leaving the roost, as this is when they passed the camera and

triggered the sensor. Although it appears that bats usually foraged alone, on 40

occasions, at least two bats were observed leaving or returning to the roost together,

sometimes even all four bats. In six instances, two bats were observed leaving the

roost simultaneously and returning together after an average of about 40 minutes.

Figure 2: Scatter plot illustrating the temporal occurrence of the different observed

social behaviors in a V. spectrum night-roost between 5 pm and 6 am (N= 60 nights).

Figure 3: A + B: Illustrations of prey sharing behavior between two bats; C: Spaghetti

plot illustrating the difference of eye distances between the prey carrying bat and the

approaching bat, showing a significant difference in eye distance (as a proxy for

body size).

Eye distance measurements

We detected a clear difference in eye distance between bats involved in prey sharing

interactions. The bat that brought the prey into the roost had a significantly larger eye

distance than the bat receiving the prey (paired Wilcoxon test, Z= -2.527; n= 8, exact

p: 0.0078; Fig. 4). Assuming that eye distance can be used as a proxy for body size,

our data suggests that the bat sharing prey is larger than the bat receiving prey,

which would be expected if adults share prey with their offspring.

Discussion

Here, we provide the first comprehensive account of prey sharing and other social

behaviors in the spectral bat *V. spectrum*. Prey sharing was a clearly cooperative

social behavior wherein a bat successfully captured prey, brought it to the roost

where group members were present, and willingly transferred the prey to another

bat. Our video observations consistently showed that upon entering the roost with

prey, the carrying bat was often approached by another bat attempting to acquire the

prey. When other bats were present, they typically remained calm and inactive,

suggesting they recognized the intended recipient of the prey, often a bat already

positioned to receive it. In most instances of prey sharing, the providing bat

voluntarily and promptly handed over the prey to the receiving bat. Occasionally,

conflicts over prey arose, as evidenced by one video depicting a tug-of-war scenario

where an approaching bat successfully wrestled the prey away. Previous

observations of a *V. spectrum* pair with a pup in captivity suggest that male bats may

occasionally bring prey to the roost, potentially to provision females (Bradbury,

personal observation via (43). This behavior aligns with an anecdotal observation

from December 2024, where one bat, presumably the lactating female, remained in

the roost with the pup while two other bats were out foraging. The prey transfer

between the prey-carrying bat and the presumed mother, which was remarkably

silent comparted to other prey transfers that were accompanied by social

vocalizations supports the hypothesis of males provisioning females under certain

circumstances.

To date, there has been scarce evidence of prey sharing among bats. In the closely

related sister species *Chrotopterus auritus,* individuals reportedly bring back prey

into the roost and then share it with the other members (Rodrigo Medellín, personal

communication, (46). Two studies in captivity provide additional evidence for

parental food sharing in carnivorous species. In the fish-eating *Noctilio albiventris*,

(47) observed that just prior to weaning at three months, captive mothers fed

juveniles with masticated fish or mealworms from their cheek pouches. Similarly, in

Megaderma lyra, mothers reportedly transferred either entire or partly consumed

frogs to their young. These food transfers ceased when the young bats reached

approximately 74 days of age (48). It appears that lactating females of *M. lyra*

provision their offspring by supplementing milk with solid food, akin to the behavior

hypothesized for *V. spectrum* (43). The only well-documented instance of prey-

sharing behavior observed in the wild among bats occurs in *Micronycteris microtis*. In

this species, weaned pups are provisioned with large insects by their mothers for an

additional five months. This extended provisioning likely helps pups learn to handle

challenging prey and refine the acoustic skills essential for effective hunting.

Although *M. microtis* primarily forages on insects, it has a remarkably broad diet that

also includes lizards, making it the smallest known carnivorous bat to date (49, 50).

Another prominent example of food sharing behavior in bats is the well-studied

regurgitation of blood in vampire bats; however, here, food is shared between kin as

well as non-kin group members that are in critical physical conditions (51–53), rather

than on a regular basis and/or for learning purposes. Nectar-feeding bats also

regurgitate food, but only for their own offspring, presumably to substitute their diet

on their way to independence (54).

Extended pup dependency & presumed biparental care

Observations of a captive *V. spectrum* pair with a pup suggest that males may

occasionally bring prey to the roost to provision females or nourish young pups not

yet fully capable of independent hunting (Bradbury, via (43)). During the nursing

period, pups likely rely on milk supplemented with meat, which facilitates their

transition to a carnivorous diet and helps them develop prey-handling skills (43). Our

data supports this strategy as a way to ensure adequate nutrition and skill

development in young bats. Based on eye-distance measurements, two smaller

individuals in our roost likely represent pups from different years, as twins have

never been reported in *V. spectrum*. Pups are reported to be born at the end of the

dry season (42, 43) and the roost with these four bats was already discovered by us

in November 2022, so the younger offspring must have been born at least 1.5 years

ago, and the older one approximately 2.5 years ago. This indicates a prolonged

period of parental care, which is relatively uncommon in bats. This extended

investment is likely tied to the species' socially monogamous structure, a rarity

among mammals (55, 56) and to the challenges associated with carnivory. Parental

investment, defined as post-birth behaviors that enhance offspring reproductive

success (57), is influenced by the costs and benefits to parents (58). In species with

low mate competition and high paternity certainty, such as *V. spectrum*, males are

more likely to invest in parental care. This aligns with the species' low population

density, large territory requirements, and limited roost availability in dry forests (59).

Biparental care, hypothesized by Vehrencamp et al. (43), likely enhances offspring

survival and success compared to uniparental care (60). While biparental care is

common in 90% of bird species (61), it is rare in mammals, where females alone

provide care in 90% of species (58). Notably, some of the social behaviors of the

spectral bat seem to have quite some commonalties with the big-eared woolly bat

(*Chrotopterus auritus)* that also exhibits a socially monogamous social group

structure, shares a similar habitat and comparable diet (62, 63). Therefore, it can be

surmised that similar factors act on *C. auritus* and it would be very likely that this

species has also evolved some kind of biparental care (46) .

Prey items and Foraging Dynamics

In addition to prey-sharing behavior, several videos documented bats consuming

prey within the roost. Due to the 1-minute recording limitation, it was often unclear

whether the prey was brought into the roost by the bat itself or transferred prior to the

recording. However, some videos showed bats entering with prey and consuming it

alone without sharing. In one instance, a bat carrying prey crawled to its roosting

spot and ate it despite a nearby bat approaching and gently stretching its head

toward the prey, possibly begging. The prey-carrying bat appeared undisturbed and

continued eating. Vehrencamp et al. (43) proposed that bats might bring prey to the

roost when caught late at night, close to sunrise, to avoid daylight exposure.

However, our observations recorded such behavior at various times during the night,

suggesting that when prey is captured near the roost, it provides a convenient, safe

location for consumption. Visual classifications identified small rodents as the

primary prey brought back to the roost, with only 9 of 30 videos showing birds being

brought in. A study in 2022 based on fecal sample analysis from a *V. spectrum* roost

in Nicaragua found birds to be the most common prey group, followed by bats and

then small rodents (59). Interestingly, no videos documented *V. spectrum* bringing

smaller bat species back to the roost, suggesting that bats may be hunted midair,

unlike diurnal birds, which are likely located in their nests through olfaction (42, 43).

While previous studies reported *V. spectrum* foraging solitarily, with individuals

leaving and returning to the roost separately (43), our data recorded 40 instances

where two or more bats left or returned simultaneously, including 6 cases where two

bats left and returned together after approximately 40 minutes. These observations

suggest that while *V. spectrum* typically forages alone, exceptions occur, particularly

when subadults may forage with adults to practice and develop hunting skills.

Further social behaviors

Play behavior may contribute to shaping adult skill sets. Observed examples in our

videos include chasing cockroaches, exploring wildlife cameras, and play-fighting,

likely among subadults. Play is often viewed as a "non-serious" behavior, where

actions are performed for their own sake rather than practical outcomes. This allows

youngsters to explore environments and develop responses in a low-risk manner

(64, 65). Play and exploration are documented in humans and non-human animals,

including birds, frogs, and cephalopods, but are most prevalent in mammals,

especially those with extended parental care (66, 65). Species with greater parental

care often exhibit more play, which helps juveniles develop adaptive behaviors for

current and future challenges (67). Play may also influence gene expression and

evolution indirectly (65). Our data highlights the presence of play behavior in bats,

emphasizing its potential developmental importance, despite limited research to

date.

The pronounced social roosting behaviors of *V. spectrum*, including hugging-like

actions—characterized by embracing, wrapping, or holding closely with limbs, wings,

or body—and greeting interactions, were observed daily. Most bats are highly social,

roosting in close contact and relying on strong bonds for survival (8). Such bonds

may offer cooperative benefits, such as information transfer within the group. Similar

to allogrooming, a behavior that helps maintain social bonds in dynamic systems (31,

68), these close-contact behaviors in *V. spectrum* likely reinforce social ties. While all

bats exhibit autogrooming, allogrooming is less common (69–71). In *V. spectrum*,

hugging and greeting likely strengthen bonds within their small social groups, which

is crucial for both parent-offspring and male-female relationships.

Conclusion

In conclusion, our findings revealed that prey sharing in *V. spectrum* is a common

and mostly voluntary behavior, with bats bringing prey back to the roost and handing

it over to other roost-mates, particularly those that may be less capable of hunting for

themselves. This behavior supports the hypothesis that adults assist in transitioning

young bats from milk to a carnivorous diet, ensuring adequate food intake and

offering them the opportunity to practice how to handle large prey items. This study

contributes to the broader understanding of bat sociality by highlighting the unique

cooperative behaviors of *V. spectrum* and provides a foundation for further research

into the ecological and evolutionary implications of these behaviors. The insights

gained from this study not only enhance our knowledge of *V. spectrum* but also

contribute to the general understanding of social evolution and cooperative

behaviors in mammals.

Acknowledgments

We would like to express our thanks to all the workers from "La Estación

Experimental Forestal Horizontes", especially Marlon Gonzalez. We are also grateful

to Emma Dittrich for her drawings.

Roles of contribution:

Marisa Tietge: Discoverer of the roost, Conceptualization, Methodology,

Investigation, Formal analysis, Data curation, Visualization, Project administration,

Writing- Original draft preparation, Writing- Reviewing and Editing

Mirjam Knörnschild: Study Design, Supervision, Writing- Reviewing and Editing

Eduardo Artavia Durán: Local cooperation partner, Writing- Reviewing and Editing

Competing interests: The authors state no conflict of interest. The authors have no

financial or proprietary interests in any material discussed in this article.

Research funding: This work was supported by a grant from the Leibniz Foundation

(P122/2020) to M.K.

Supplementary material

Example videos from above-described key social behaviors:

o   Prey sharing behavior (small mammal)

o   Prey sharing behavior (bird)

o   Fight about food

o   Social roosting

o   Greeting behavior

o   Play (object)

o   Play (food)

o   Play (social)

Literature Cited

1. Buchan JC, Alberts SC, Silk JB, Altmann J. True paternal care in a multi-male primate society. Nature 2003; 425(6954):179–81.

2. Engh AL, Siebert ER, Greenberg DA, Holekamp KE. Patterns of alliance formation and postconflict aggression indicate spotted hyaenas recognize third-party relationships. Animal Behaviour 2005; 69(1):209–17.

3. Komdeur J, Eikenaar C, Brouwer L, Richardson DS. The Evolution and Ecology of Cooperative Breeding in Vertebrates. In: The Evolution and Ecology of Cooperative Breeding in Vertebrates. Wiley; 2008.

4. Krams I, Krama T, Igaune K, Mänd R. Experimental evidence of reciprocal altruism in the pied flycatcher. Behav Ecol Sociobiol 2008; 62(4):599–605.

5. Nowak MA. Evolving cooperation. J Theor Biol 2012; 299:1–8.

6. Dugatkin LA. Cooperation in animals: An evolutionary overview. Biol Philos 2002; 17(4):459–76.

7. Lacy EA, Sherman PW. Cooperative breeding in mammals: Cooperative Breeding In Naked Mole-Rats: Implications for Vertebrate and Invertebrate Sociality. Cambridge: Cambridge University Press; 2007.

8. Carter GG, Wilkinson GS. Food sharing in vampire bats: reciprocal help predicts donations more than relatedness or harassment. Proc. R. Soc. B. 2013; 280(1753):20122573.

9. Connor RC, editor. Group living in whales and dolphins. Chicago: The University of Chicago Press; 2000. (Cetacean societies: field studies of dolphins and whales).

10. Bailey I, Myatt JP, Wilson AM. Group hunting within the Carnivora: physiological, cognitive and environmental influences on strategy and cooperation. Behav Ecol Sociobiol 2013; 67(1):1–17.

11. Nichols TC. Cooperative hunting of Canada Geese (*Branta canadensis*) by Gray Wolves (*Canis lupus*) in northern Quebec. Can Field Nat 2015; 129(3):290.

12. Harcourt A, De Waal FB. Coalitions and alliances in humans and other animals [Alexander Harcourt and Frans B.M. de Waal]. American J Phys Anthropol 1993; 1992(4):533–5.

13. Dugatkin LA. The Evolution of Cooperation. BioScience 1997; 47(6):355–62.

14. Simmons NB, Cirranello AL. Bat Species of the World: A taxonomic and geographic database; 2024 [cited Accessed on 2024 Dec 10]. Available from: URL: https://batnames.org.

15. Kunz T. H, Fenton M. B. Bat Ecology. Journal of Mammalogy 2004; 85(2):366–7.

16. Altringham JD. Bats: From evolution to conservation. 2nd ed. New York: Oxford
University Press; 2011.

17. McCracken GF, Wilkinson GS. Bat Mating Systems. In: McCracken GF,
Wilkinson GS, editors. Reproductive Biology of Bats. Elsevier; 2000. p. 321–62.

18. Kerth G. Causes and Consequences of Sociality in Bats. BioScience 2008;
58(8):737–46.

19. Chaverri G, Ancillotto L, Russo D. Social communication in bats. Biol Rev Camb
Philos Soc 2018; 93(4):1938–54.

20. Barbour T. A Peculiar Roosting Habit of Bats. The Quarterly Review of Biology
1932; 7(3):307–12.

21. Kunz TH, editor. Ecology of Bats. Boston, MA: Springer US; 1982.

22. Chaverri G, Kunz TH. Ecological Determinants of Social Systems. In: Behavioral
ecology of tropical animals. Elsevier; 2010. p. 275–318 (Advances in the Study of
Behavior).

23. Gaudet CL, Fenton MB. Observational learning in three species of insectivorous
bats (Chiroptera). Animal Behaviour 1984; 32(2):385–8.

24. Ratcliffe JM, Hofstede HM ter. Roosts as information centres: social learning of
food preferences in bats. Biol Lett 2005; 1(1):72–4.

25. Page RA, Ryan MJ. Social transmission of novel foraging behavior in bats: frog
calls and their referents. Curr Biol 2006; 16(12):1201–5.

26. Dechmann DKN, Heucke SL, Giuggioli L, Safi K, Voigt CC, Wikelski M.
Experimental evidence for group hunting via eavesdropping in echolocating bats.
Proc. R. Soc. B. 2009; 276(1668):2721–8.

27. Dechmann DKN, Kranstauber B, Gibbs D, Wikelski M. Group hunting-a reason
for sociality in molossid bats? PLoS One 2010; 5(2):e9012.

28. Wright GS, Wilkinson GS, Moss CF. Social Learning of a Novel Foraging Task
by Big Brown Bats (Eptesicus fuscus). Animal Behaviour 2011; 82(5):1075–83.

29. O'Mara MT, Dechmann DKN, Page RA. Frugivorous bats evaluate the quality of
social information when choosing novel foods. Behavioral Ecology 2014;
25(5):1233–9.

30. Chaverri G, Gillam EH, Kunz TH. A call-and-response system facilitates group
cohesion among disc-winged bats. Behavioral Ecology 2013; 24(2):481–7.

31. Dunbar R, Shultz S. Bondedness and sociality. Behav 2010; 147(7):775–803.

32. Wilkinson GS. Information transfer at evening bat colonies. Animal Behaviour
1992; 44:501–18.

33. Wilkinson GS, Boughman JW. Social calls coordinate foraging in greater spear-
nosed bats. Animal Behaviour 1998; 55(2):337–50.

34. Willis CKR, Brigham RM. Social thermoregulation exerts more influence than
microclimate on forest roost preferences by a cavity-dwelling bat. Behav Ecol
Sociobiol 2007; 62(1):97–108.

35. Ortega J, Guerrero JA, Maldonado JE. Aggression and Tolerance by Dominant
Males of Artibeus jamaicensis : Strategies to Maximize Fitness in Harem Groups.
Journal of Mammalogy 2008; 89(6):1372–8.

36. Nagy M, Knörnschild M, Voigt CC, Mayer F. Male greater sac-winged bats gain
direct fitness benefits when roosting in multimale colonies. Behavioral Ecology 2012;
23(3):597–606.

37. Gager Y. Information transfer about food as a reason for sociality in bats.
Mammal Review 2019; 49(2):113–20.

38. Ripperger SP, Stockmaier S, Carter GG. Tracking sickness effects on social
encounters via continuous proximity sensing in wild vampire bats. Behavioral
Ecology 2020; 31(6):1296–302.

39. Moreno KR, Weinberg M, Harten L, Salinas Ramos VB, Herrera M LG, Czirják
GÁ et al. Sick bats stay home alone: fruit bats practice social distancing when faced
with an immunological challenge. Ann N Y Acad Sci 2021; 1505(1):178–90.

40. McWilliam AN. Emergence behaviour of the bat Tadarida (Chaerephon) pumila
(Chiroptera: Molossidae) in Ghana, West Africa. Journal of Zoology 1989;
219(4):698–701.

41. Frick WF, Kingston T, Flanders J. A review of the major threats and challenges
to global bat conservation. Ann N Y Acad Sci 2020; 1469(1):5–25.

42. Greenhall AM. Notes on the Behavior of the False Vampire Bat. Journal of
Mammalogy 1968; 49(2):337.

43. Vehrencamp SL, Stiles FG, Bradbury JW. Observations on the Foraging
Behavior and Avian Prey of the Neotropical Carnivorous Bat, Vampyrum spectrum.
Journal of Mammalogy 1977; 58(4):469–78.

44. Solari S. Vampyrum spectrum. The IUCN Red List of Threatened Species 2018;
2018.

45. Gual-Suárez F, Medellín RA. We eat meat: a review of carnivory in bats.
Mammal Review 2021; 51(4):540–58.

46. Trujillo LA, Gual-Suárez F, Trujillo RE, Medellín RA. Arachnids that feed on
vertebrate carrion: necrophagy by the whip spider Paraphrynus raptator (Amblypygi:
Phrynidae) and its relation to the feeding behavior of the woolly false vampire bat
Chrotopterus auritus (Chiroptera: Phyllostomidae). The Journal of Arachnology 2021;
49(3).

47. Brown PE, Brown TW, Grinnell AD. Echolocation, development, and vocal
communication in the lesser bulldog bat, Noctilio albiventris. Behav Ecol Sociobiol
1983; 13(4):287–98.

48. Raghuram H, Marimuthu G. Maternal feeding of offspring with vertebrate prey in
captive Indian false vampire bat, Megaderma lyra. Acta Chiropterologica 2007;
Volume 9(2):437-443(7).

49. Santana SE, Geipel I, Dumont ER, Kalka MB, Kalko EKV. All you can eat: high
performance capacity and plasticity in the common big-eared bat, Micronycteris
microtis (Chiroptera: Phyllostomidae). PLoS One 2011; 6(12):e28584.

50. Geipel I, Jung K, Kalko EKV. Perception of silent and motionless prey on
vegetation by echolocation in the gleaning bat Micronycteris microtis. Proc. R. Soc.
B. 2013; 280(1754):20122830.

51. Wilkinson GS. Reciprocal food sharing in the vampire bat. Nature 1984;
308(5955):181–4.

52. Elizalde-Arellano C, López-Vidal JC, Arroyo-Cabrales J, Medellín RA, Laundré
JW. Food sharing behavior in the hairy-legged vampire bat Diphylla ecaudata. Acta
Chiropterologica 2007; 9(1):314–9.

53. Wilkinson GS, Carter GG, Bohn KM, Adams DM. Non-kin cooperation in bats.
Philos Trans R Soc Lond B Biol Sci 2016; 371(1687):20150095.

54. Rose A, Wöhl S, Bechler J, Tschapka M, Knörnschild M. Maternal mouth-to-
mouth feeding behaviour in flower-visiting bats, but no experimental evidence for
transmitted dietary preferences. Behav Processes 2019; 165:29–35.

55. Orians GH. On the Evolution of Mating Systems in Birds and Mammals. The
American Naturalist 1969; 103(934):589–603.

56. Alexander RD. The Evolution of Social Behavior. Annu. Rev. Ecol. Syst. 1974;
5(1):325–83.

57. Klug, Hope, Alonzo SH, Bonsall, Michael B. Theoretical foundations of parental
care 2012.

58. Clutton-Brock TH. The evolution of parental care. Princeton, N.J.: Princeton
University Press; 1991. (Monographs in behavior and ecology).

59. Martínez-Fonseca JG, Mau R, Walker FM, Medina-Fitoria A, Yasuda K,
Chambers CL. Vampyrum spectrum (Phyllostomidae) movement and prey revealed
by radio-telemetry and DNA metabarcoding. PLoS One 2022; 17(4):e0265968.

60. Barta Z, Székely T, Liker A, Harrison F. Social role specialization promotes
cooperation between parents. The American Naturalist 2014; 183(6):747–61.

61. Cockburn A. Prevalence of different modes of parental care in birds. Proc. R.
Soc. B. 2006; 273(1592):1375–83.

62. Medellín RA. Chrotopterus auritus. Mammalian species 1989; 343:1–5.

63. Esbérard CEL, Motta AG, Almeida JC, Ferreira LCS, Costa LM. Reproduction of
Chrotopterus auritus (Peters) in captivity (Chiroptera, Phyllostomidae). Braz J Biol
2006; 66(3):955–6.

64. Spinka M, Newberry RC, Bekoff M. Mammalian play: training for the unexpected.
The Quarterly Review of Biology 2001; 76(2):141–68.

65. Burghardt GM. The Genesis of Animal Play. The MIT Press; 2005.

66. Zylinski S. Fun and play in invertebrates. Curr Biol 2015; 25(1):R10-2.

67. Bjorklund D. Mother knows best: Epigenetic inheritance, maternal effects, and
the evolution of human intelligence. Developmental Review 2006; 26(2):213–42.

68. Russell YI, Phelps S. How do you measure pleasure? A discussion about
intrinsic costs and benefits in primate allogrooming. Biol Philos 2013; 28(6):1005–20.

69. Wilkinson GS. Social grooming in the common vampire bat, Desmodus rotundus.
Animal Behaviour 1986; 34(6):1880–9.

70. Fleming TH, Nelson AA, Dalton VM. Roosting Behavior of the Lesser Long-
Nosed Bat, Leptonycteris curasoae. Journal of Mammalogy 1998; 79(1):147–55.

71. Ortega J, Maldonado JE. Female interactions in harem groups of the Jamaican
fruit-eating bat, Artibeus jamaicensis (Chiroptera: Phyllostomidae). Acta
Chiropterologica 2006; 8(2):485–95.

Fig 1: Photo Vampyrum in roost

Fig 2: behavior occurence

Fig 3: eye distance measurement