Tumbleweed: an artificial motor protein that walks along a DNA track

Authors: Patrik Nilsson[1,2†], Neil O. Robertson[3,4†], Nils Gustafsson[1,2†], Roberta B. Davies[3†], Chu Wai Liew[3,4], Aaron Lyons[5], Ralf Eichhorn[6], Cassandra S. Niman[1,2], Gerhard A. Blab[5‡], Elizabeth H.C. Bromley[7], Andrew E. Whitten[8], Anthony P. Duff[8], Jason P. Beech[1,2], Peter Jönsson[1,9], Till Böcking[10,11], Birte Höcker[12], Derek N. Woolfson[13,14,15], Nancy R. Forde[5], Heiner Linke[1,2*], Paul M. G. Curmi[3,4*]

Affiliations:

[1]NanoLund, Lund University; Box 118, 221 00 Lund, Sweden.

[2]Solid State Physics, Lund University; Box 118, 221 00 Lund, Sweden

[3]School of Physics, University of New South Wales; Sydney, NSW 2052, Australia.

[4]School of Biotechnology and Biomolecular Sciences, University of New South Wales; Sydney, NSW 2052, Australia.

[5]Department of Physics, Simon Fraser University Burnaby; BC, V5A 1S6, Canada.

[6]Nordita, Royal Institute of Technology and Stockholm University; SE-106 91, Stockholm, Sweden.

[7]Department of Physics, University of Durham; Durham, United Kingdom.

[8]Australian Nuclear Science and Technology Organisation; Lucas Heights, NSW 2234, Australia.

[9]Physical Chemistry, Lund University; P.O. Box 124, SE-22100 Lund, Sweden.

[10]Department of Molecular Medicine, School of Biomedical Sciences, University of New South Wales; Sydney, NSW 2052, Australia.

[11]EMBL Australia Node in Single Molecule Science, School of Biomedical Sciences, University of New South Wales; Sydney, NSW 2052, Australia.

[12]Department of Biochemistry, University of Bayreuth; 95447 Bayreuth, Germany.

[13]School of Chemistry, University of Bristol; Bristol BS8 ITS, United Kingdom.

[14]School of Biochemistry, University of Bristol; Bristol BS8 ITD, United Kingdom.

[15]Max Planck-Bristol Centre for Minimal Biology, University of Bristol; Bristol BS8 ITS, United Kingdom.

*Corresponding author. Email: heiner.linke@ftf.lth.se and p.curmi@unsw.edu.au

† These authors contributed equally to this work.

‡Present address: Debye Institute for Nanomaterials Science and Department of Physics, Utrecht University; The Netherlands.

Abstract:

Nature has used proteins to evolve molecular motors that facilitate life. Although small-molecule- and DNA-based molecular motors have been synthesized, the creation of an artificial motor protein has remained a goal of synthetic biology. Here we describe a modular approach to create an artificial motor protein, the Tumbleweed (TW). TW has three legs, each with a ligand-gated DNA binding domain that controls binding to a DNA track. TW works via a Brownian ratchet mechanism where steps are effected by diffusion and then rectified by the controlling ligands. Using single-molecule fluorescence assays and a microfluidic device, we show that TW steps directionally along a DNA track when the ligand concentrations are altered in sequence.

Introduction:

Life is characterized by directed motion on all length scales. Nature has evolved numerous molecular machines that produce directed motion, usually powered by free energy gained from ATP hydrolysis. Several of these machines can translate along linear tracks, which are made of either protein polymers (e.g. myosin on actin (*1, 2*); kinesin (*3*) and dynein (*4*) on microtubules) or polynucleic acids (e.g. helicases and polymerases on DNA (*5*); and the ribosome on mRNA (*6*)). The sophisticated and nuanced functions of these and other biological machines are afforded by the chemical and structural complexity of proteins, which are key to their success in evolution.

Inspired by biological motors, and to help understand how they work, biological motor proteins have been re-engineered (*7*), proteins have been repurposed as active components of large motors (*8*), and artificial molecular motors have been created based on small molecules (*9*) as well as more sophisticated DNA-based motors and machines (*10*). The complexity of proteins offers the prospect of even more advanced artificial motors but also makes their successful design a significant, unsolved challenge.

Here, we build an artificial motor protein designed to walk along a DNA track. Tumbleweed (TW) is a three-legged molecular walker created using a modular approach (*11-13*) that combines the use of biological DNA-binding proteins and structural components coupled via a SpyTag/SpyCatcher (*14*) hub (Fig. 1A). TW is a clocked walker that is powered by periodically changing the concentrations of controlling ligands. We use TW to explore one physical mechanism of mechanochemical free energy transduction: a Brownian ratchet mechanism where all movement is facilitated by thermal fluctuations that are rectified by external ligand control in combination with an asymmetric track (*15*). Single-molecule Förster resonance energy transfer (FRET) assays show that TW can step along DNA under microfluidic ligand control of both timing and direction of stepping.

TW mechanism

TW is designed to step directionally along a specific, asymmetric DNA track, clocked by control of externally regulated ligand concentrations (*13*). TW has three distinct feet each made from different repressor proteins that bind to cognate DNA sites in the presence of their respective ligand, namely TrpR and its ligand L-tryptophan (Trp), DtxR and Co^{2+}, and MetJ and S-adenosyl-L-methionine (SAM). The three feet are connected to a central hub by α-helical coiled-coil legs (Fig. 1A). To ensure unidirectional motion, the DNA track is designed to present specific binding sites for the TW feet in a unique order: *trpR*, *dtxR*, *metJ*, which repeats for the

length of the track (Fig. 1B). Consider the case where TW is clocked to walk in this order (towards the right in Fig. 1C). When TW is bound to the DNA track via one foot, for example TrpR bound to *trpR* in the presence of Trp, it can reach and bind to the adjacent site, *dtxR*, when the next ligand, Co^{2+}, is introduced (Fig. 1C, top). However, its legs are designed to be too short to reach the previous *dtxR* site, which is separated from the TW-bound *trpR* site by an intervening *metJ* site (Fig. 1B). Thus, by bathing TW in solutions comprising pairs of controlling ligands in a cyclical, time-dependent manner with (Trp + Co^{2+}), then (Co^{2+} + SAM), then (SAM + Trp), TW will move from a state where it is bound to the track via TrpR and DtxR, next by DtxR and MetJ, and then MetJ and TrpR (Fig. 1C). In between these relatively stable, two-foot-bound states, TW will transit more vulnerable, single-foot-bound states (Fig. 1C), where there is a higher probability of detaching from the DNA track. The direction of motion can be controlled by altering the order in which ligand solutions are presented.

The free energy powering this unidirectional motion comes from the controlling ligands and their chemical potentials, which are externally controlled in a time-dependent manner. Each DNA-binding module will bind its respective ligand when the concentration of the ligand is high (high chemical potential) and will eventually dissociate and release its controlling ligand when the ligand concentration is low (low chemical potential). Because each step is enabled by diffusive motion, TW is a Brownian ratchet (*15*).

Results

Construction of a three-legged TW

Three-legged, heterotrimeric proteins are not common in nature. Therefore, we designed and assembled such a hub using the SpyTag/SpyCatcher system (*14*) as follows. TW was made in two parts, TW1 and TW2, each produced by expressing synthetic genes in *Escherichia coli* and purifying the His-tagged proteins using immobilized metal affinity chromatography (IMAC; Figs 1A and 2A). TW1 comprises DtxR, the α-helical domain from the geminin coiled-coil, and SpyCatcher, each connected by flexible Gly-Ser linkers. TW2 comprises TrpR, the geminin coiled-coil α helix, SpyTag, a second copy of the geminin coiled-coil α helix and MetJ, each connected by Gly-Ser linkers. TW1 and TW2 form homodimeric proteins that were purified to homogeneity (Fig. 2A). The size of each molecule determined by mass photometry was consistent with the expected mass (Fig. 2B; Table S1). Mature TW was generated by mixing TW1 and TW2, with a slight excess of TW1 (1.2:1). The covalent TW product was confirmed by SDS-PAGE (Fig. 2A). This was purified by size-exclusion chromatography (SEC) with the size determined by mass photometry consistent with the expected mass (Fig. 2B; Table S1).

Characterization of TW structure

To verify the structural integrity of TW, we collected SEC-coupled small-angle X-ray scattering (SEC-SAXS) data using the Australian Synchrotron (Fig. 2C). The radius of gyration (R_g Guinier 67.7 ± 0.6 Å; R_g $P(r)$ 69.1 ± 0.4 Å) was consistent with an extended protein (Fig. S1). Given the engineered hinge flexibility of TW, we anticipated an ensemble of structures in solution. We were able to fit the experimental data using the program Multi-FoXS (*16*), where we allowed the program to bend AlphaFold2-generated models (*17*) of TW via six hinge points linking the three legs to both the SpyTag/SpyCatcher hub and to each of the DNA binding modules. The best fit was obtained with two conformers: an L-shaped conformer (72.5%; Model

1, Fig. 2C top left) and an extended conformer (27.5%; Model 2, Fig. 2C bottom left). Adding further conformations only marginally improved the fit of the model to the data and the new conformers clustered with the original two.

Design of DNA tracks

DNA tracks were designed to space binding sites, *trpR*, *dtxR* and *metJ*, such that each protein foot would bind on the same side of the DNA (Fig. 1B). The spacing between adjacent sites was set so that two feet of TW could bind adjacent sites but the stride of TW was not long enough to bind simultaneously to non-adjacent sites, assuming the DNA to be straight. The spacing between *trpR* and *dtxR* sites was made slightly longer as both TrpR and DtxR bind to the major groove of DNA whereas MetJ binds to the minor groove (Fig. 1B).

Visualizing TW on an eight-repeat track

A track was generated with eight copies of *metJ-trpR-dtxR* inserted into the pYIC plasmid (*18*). This was linearized and mixed with TW in the presence of 10 mM Trp and visualized using atomic force microscopy (AFM) after drying on a mica substrate. Images revealed eight bright, evenly spaced features along the DNA (Fig. 2D). The contour lengths of the DNA flanking the eight bound TWs are consistent with the location of the inserted (*metJ-trpR-dtxR*)$_8$ track.

Binding of TW to DNA is specific for its cognate site and controlling ligand

For TW to function as intended, we needed to ensure that each foot of TW could bind specifically to its cognate DNA sites with high affinity (K_d 1-100 nM) only with its respective controlling ligand. We used surface plasmon resonance (SPR) to characterize the binding of TW to various DNA tracks (Fig. 3A; Figs. S2-S4) and to optimize both the intervening sequences in the DNA track and the solution conditions (controlling ligand concentrations, pH and ionic strength) for binding.

SPR sensorgrams for TW binding to DNA oligonucleotides containing single cognate binding sites showed that in the absence of ligands, TW bound minimally to DNA (Fig. 3A, top row). However, with each ligand, TW bound to the appropriate cognate DNA site as intended, with minimal offsite non-specific binding. The sensorgrams were used to calculate the dissociation constants, K_d, for TW to single-site DNA oligonucleotides, which were found to be in the 10–30 nM range (Fig. 3B; Table S2). TW dissociation constants were very similar albeit greater than those of purified repressor proteins DtxR and MetJ, while TW had a slightly stronger affinity for *trpR* oligonucleotides in the presence of Trp compared to purified TrpR repressor (Fig. 3B, Fig. S2, Table S2).

The specificity of each TW foot for its cognate DNA site was determined by titrating TW in the presence of a single controlling ligand on an SPR chip with two-site DNA tracks (Fig. S3). In the absence of ligands, TW did not bind to any two-site track where [TW] was varied from 2.5 to 40 nM (Fig. S3A). In this concentration range, no non-specific binding was observed to a track containing *metJ* and *trpR* when Co^{2+} was present (Fig. S3C). A low level of non-specific binding was observed in the presence of SAM on a track containing *trpR* and *dtxR* sites when [TW] > 20 nM (Fig. S3B). The highest degree of non-specific binding by TW was observed in

the presence of the Trp ligand (Fig. S3D). Titration of TW onto a track containing *metJ* and *dtxR* sites showed clear, specific binding of TW in the presence of appropriate ligands, either SAM or Co^{2+} (Fig. S3B,C,E), while increasing levels of non-specific binding were observed in the presence of non-cognate ligand Trp (Fig. S3D,E). However, the apparent affinity of off-target binding to the *dtxR-metJ* track in the presence of Trp was more than threefold lower than the on-target affinities (Table S2), and for concentrations of TW below 10 nM, non-specific binding to this track was minimal compared with specific binding (Fig. S3D,E).

The intended stoichiometry of TW:DNA complexes was confirmed by mass photometry. TW bound all single cognate binding sites in a 1:1 fashion (Fig. S5A, Table S1). Importantly, TW also formed a 1:1 complex with DNA containing two cognate sites in the presence of both corresponding ligands, even when the DNA was added in 4–10 fold molar excess to TW (Fig. 3C, Fig. S5B, Table S1). In these experiments, we used 25 nM TW with up to 250 nM DNA, indicating TW preferentially binds on a single DNA species at least up to the nanomolar concentration range.

Binding kinetics of TW to DNA

Timescales are critical for the operation of a processive clocked walker (*11, 13, 19, 20*). For TW, the time that it spends in a particular two-ligand solution condition must be short compared to the half-life of the corresponding two-foot-bound state, while the release of single feet when ligands are removed must be fast compared to this timescale. To determine these timescales, the dissociation kinetics of TW:DNA complexes were studied by an SPR displacement assay, where during the dissociation phase, a large excess of free specific DNA was introduced to prevent rebinding of TW to surface-bound DNA. This allowed the determination of apparent off-rates in different ligand conditions (Fig. 3D, Fig. S4, Table S3). The half-lives for TW following the removal of controlling ligands were < 2 s. In the presence of single controlling ligands, the half-lives were ~ 0.6 – 4 s. In both the absence and presence of controlling ligands, the MetJ foot was the fastest to dissociate from DNA, and the DtxR foot the slowest. The off-rates decreased dramatically in the presence of two controlling ligands, with half-lives in the ~30–50 s range. These measurements set the boundaries for clocking a processive TW.

Single-molecule observation of clocked TW stepping along a DNA track in real time

With the designed features and functions of each component of the assembled TW confirmed, we examined its ability to walk along the DNA track. Using single-molecule fluorescence microscopy we showed that TW can take multiple steps forward and backward along a DNA track in response to microfluidically controlled ligand exchanges.

A short DNA track (Fig. 4A) containing four binding sites (from bottom to top: *trpR, dtxR, metJ,* and *trpR*), was attached at one end through streptavidin to a PLL-PEG-biotin polymer–coated coverslip. The track's two *trpR* sites were labelled with two different FRET acceptors, namely ATTO647N (red emission) at the top and ATTO565 (yellow emission) at the bottom. The corresponding FRET donor, AlexaFluor 488 (green emission), was used to label the TrpR foot of TW close to the DNA binding site (Ser107) to enable FRET when the TrpR foot binds to either of the two *trpR* sites. This allows each of the three different, two-foot-bound, states to be uniquely identified by the wavelength of emission. Starting closest to the coverslip

we expected: yellow FRET emission with TW bound to the bottom-*trpR* site and the *dtxR* site (Fig. 4B); green emission (*i.e.*, no FRET) with TW bound to the *dtxR* and *metJ* sites (Fig. 4C); and red FRET emission with TW bound to the *metJ* and top-*trpR* sites (Fig. 4D).

To control TW stepping by changing the ligand concentrations in real time, we used a custom-designed microfluidic device (Fig. 4E). This allowed switching between three different controlling solutions with minimal drag in arbitrary temporal order, with a time resolution of less than ~ 0.2s (*21*). Ligand-pair solutions were changed at 7 s intervals, long enough to collect several fluorescence frames during each ligand step and to allow the slowest foot, DtxR, to dissociate (Fig. S4; Table S3). Once one foot is detached from the track, TW will diffuse within a larger volume around the remaining bound foot. This allows the next foot in the step series to get close enough to bind to the next cognate binding site on a subsecond timescale (*19, 20*). Given that two-foot-bound states have a half-life of ~30 s (Fig. S4; Table S3), we expected to observe an average of > 4 steps per TW molecule using a repeating sequence of solution exchanges with an exchange time of 7 s.

We report two types of experiment (Fig. 4F-I and Fig. 4J-M, respectively) that together allow us to fully characterize TW stepping up and down the four-site track. To generate stepping, we sequentially changed the ligand-pair solutions to induce TW to step down the track, then two steps up, then two steps down, and repeating for a total of 11 steps as indicated in Figs 4F and 4J. In each experiment, DNA:TW complexes were identified by colocalization, and time traces of background-subtracted fluorescence intensity from each individual complex were extracted.

Experiment 1 shows that the TrpR foot of TW binds and unbinds the *trpR* sites as the ligands are changed by the fluidic device, while TW remains attached to the DNA track (Fig. 4F-I). 1 nM TW was added to the fluidic device in the presence of Co^{2+}+SAM ligands to induce TW to bind to the central *dtxR* and *metJ* sites of the DNA (Fig. 4C). We collected light simultaneously in two wavelength bands, the first suitable to observe green emission from the TW TrpR foot and the second suitable for emission from both red and yellow fluorophores. As such, in this experiment both FRET positions (at the bottom and the top of the track) were observed in the same channel and signal from the TW was always detectable as either donor emission or FRET from either acceptor. In synchrony with the solution changes, the donor (shown in green in Fig. 4G, H)) and acceptors (shown in red in Fig. 4G, H) signals alternated out of phase between low and high emission, as shown in the example trace (Fig. 4G), until the signal was lost through bleaching or full detachment of TW. Overlaid kymographs for each of the two observation channels (Fig. 4H) from one single field of view show that there are 322 complexes where the FRET state of the TW:DNA complex is alternating in synchrony with the change in controlling solution, indicating stepping. The sum of these individual traces (Fig. 4I) demonstrates the same anti-correlated behavior in phase with the changes in controlling solution in the context of reducing total signal over time as bleaching and TW dissociation occur. Thus, Experiment 1 confirms that the TrpR foot binds and unbinds to/from the *trpR* sites on the track as dictated by the ligands. However, it cannot discriminate between TrpR binding to the intended *trpR* site versus the *trpR* site at the opposite end of the track.

Experiment 2 (Fig. 4J-M) confirms that TW successfully visits both the top and bottom of the track as intended by the controlling ligands. To achieve this, we used two-color simultaneous imaging of the bottom and top FRET acceptors recorded in separate channels. Examples of the anti-correlated FRET signals from *trpR* sites on the top and bottom of the track are shown in Fig. 4K and Fig. S6 (magenta = top of track; cyan = bottom of track, see Fig. 4J). Overlaid kymographs for each of the two observation channels from four separate fields of view

show that there are 236 complexes that exhibit dynamic behavior correlated with the controlling ligand exchange. The dark bands in these kymographs coincide with periods where TW is expected to be bound to the central *metJ* and *dtxR* sites (Fig. 4C). The sum of the individual traces (Fig. 4M) shows that TW starts at the top *trpR* site, moves down through the dark central site and continues to step. The alternation between dark and bright states clearly confirms stepping, and the presence of magenta and cyan signals confirms that TW reaches both the top and the bottom of the tracks.

In individual traces (Fig. 4K; Fig. S6), we observe that when the controlling solution contains Trp, single TWs show highly dynamic behavior, switching on a timescale of seconds between being bound to the top-*trpR* and the bottom-*trpR* sites (Fig. 4K, L; Fig. S6). This indicates that in these experiments TW can "overstep", by binding to the non-adjacent *trpR* site. By analyzing all traces during the first three solution conditions (Fig. 4L; Fig. S6), we find that the TW TrpR foot binds the intended (adjacent) *trpR* site ~65% ± 7.7% of the time (correct stepping), whereas it binds the unintended, distal *trpR* site ~35% ± 6.3% of the time (overstepping). Importantly, as we show below, modeling indicates that the motor moves forward directionally on average regardless of overstepping.

The kymographs in Figs 4H and L were sorted by time of the last observed fluorescence signal; this occurs when the TW either bleached or dissociated from the track. Because our experiments were designed to make maximum use of the available photon budget during the expected bound time, the two possible events (bleaching and dissociation) occur on similar timescales in our experiments and are thus indistinguishable. The observed bound half-life of ~30-40 s (Fig. 4H and L) agrees well with the expected bound time of a TW based on the dissociation rates measured in SPR. The coincidence of the observed TW track-bound half-life with that of the three possible two-foot-bound states (Table S3) indicates that TW only transiently populates the more vulnerable single-foot-bound states during ligand exchange.

Finally, control experiments show that no stepping by TW on the four-site oligonucleotide track occurs when switching solutions with the same ligands (keeping ligand concentrations constant while microfluidically switching between identical solutions), confirming that stepping is caused by changing ligand concentrations, and that any hydrodynamic forces due to the microfluidics play no role in the observed motion (Supplementary Text 1, Fig. S7). When ligands are removed, we observe that TW falls off the tracks as expected (Supplementary Text 1, Figs S8-S9).

Coarse-grain modeling of TW stepping

To understand how the observed stepping behavior translates into TW's ability to walk along a long DNA track that contains many *trpR - dtxR - metJ* repeats, we created a coarse-grained model based on Master equations governing the binding and unbinding events between the three TW feet and the track (Supplementary Text 2). Based on the experimentally observed binding and unbinding rates (Table S3), and including the observed likelihood to overstep, we predict that a TW on an infinite track would move directionally at a speed of about 0.5 track periods per solution cycle (Fig. S10), or ~1.2 nm/s using the 21 s experimental cycle time. The processivity is ~ 50 – 100 nm (Figs S11 and S12) and is governed by the observed detachment time in the presence of two ligands.

Discussion

We have constructed a three-legged TW where each foot binds to its cognate DNA site in the presence of its controlling ligand. TW and its DNA track have been optimized through multiple rounds of binding experiments (SPR and mass photometry) to produce a clocked Brownian walker that binds specifically to its track as a monomer with nanomolar affinity. Single-molecule FRET experiments show that under external microfluidic control, TW takes multiple 15 nm steps along its DNA track.

Taken together, our single-molecule Experiments 1 and 2 demonstrate that individual TWs successfully take up to 11 steps up and down a short track before dissociating. The dynamic behavior also indicates that in the presence of the Trp ligand, TW can "overstep", sometimes binding to the non-adjacent *trpR* site instead of the intended *trpR* site that is adjacent to the to the bound foot. Most likely this overstepping occurs because of inherent flexibility of the DNA track (Supplementary Text 3) and the TW linker regions. Similar dynamic, stochastic behavior, including occasional mis-stepping, is well known from single-molecule studies of biological motors (*22-24*). The main consequence of overstepping is variability in the displacement of individual TWs (Fig. S11) and is insensitive to the precise probability of overstepping over a wide parameter range (Figs S10-S12). In the absence of overstepping, which may be achievable by using DNA tracks based on rigid origami nanotubes (*25*), the model predicts that TW approaches a speed of one-track period per solution cycle. This corresponds to 2.3 nm/s using the 21 s experimental cycle time and may be sped up by reducing the solution cycle time. While speed is also limited by the timescale for detachment of TW feet, a speed of 10 – 20 nm/s is likely achievable.

By creating a protein walker from non-motor components, we have shown that a modular protein engineering approach can deliver a new, designed function. This approach to protein design is analogous to how new functional proteins evolve in biology through the combination and assembly of existing protein domains to render emergent properties. The eukaryotic cytoskeletal motors myosin and kinesin evolved around the time of eukaryogenesis, where eukaryotes emerged from archaea. Current studies show that Asgard archaea have acquired cytoskeletal tracks – actin and microtubules – plus small GTPases that are signature eukaryotic proteins (*26, 27*), however, they do not appear to have evolved the eukaryotic motors – myosin and kinesin – even though the motor domain putatively arose from the small GTPase domain (*26, 28*). The future discovery of a primordial myosin or kinesin will be a major landmark in understanding motor evolution.

Like the initial step to eukaryogenesis, we have taken a step towards creating an artificial motor engineered from non-motor proteins. The job is not complete, it is only just started. TW is a clocked walker that offers the advantage of external control, including the possibility for stepping along more complex tracks with a degree of programmability. Another motivation for creating artificial motor proteins is to understand how biological motors work. This will require the mastery of allostery in protein design and will begin to address the remaining challenge in the field: to achieve autonomous protein motors that use chemical energy but require no external control (*11*).

References and Notes

1. Z. Wang, S. Raunser, Structural Biochemistry of Muscle Contraction. Annu Rev Biochem 92, 411-433 (2023).

2. J. Robert-Paganin, O. Pylypenko, C. Kikuti, H. L. Sweeney, A. Houdusse, Force Generation by Myosin Motors: A Structural Perspective. Chem Rev 120, 5-35 (2020).

3. A. Yildiz, Mechanism and regulation of kinesin motors. Nature reviews. Molecular cell biology, 26, 86-103 (2025).

4. S. L. Reck-Peterson, W. B. Redwine, R. D. Vale, A. P. Carter, The cytoplasmic dynein transport machinery and its many cargoes. Nature reviews. Molecular cell biology 19, 382-398 (2018).

5. C. Bustamante, Y. R. Chemla, N. R. Forde, D. Izhaky, Mechanical processes in biochemistry. Annu Rev Biochem 73, 705-748 (2004).

6. V. Ramakrishnan, The ribosome emerges from a black box. Cell 159, 979-984 (2014).

7. N. V. DelRosso, N. D. Derr, Exploiting molecular motors as nanomachines: the mechanisms of de novo and re-engineered cytoskeletal motors. Current opinion in biotechnology 46, 20-26 (2017).

8. C. S. Korosec et al., Motility of an autonomous protein-based artificial motor that operates via a burnt-bridge principle. Nat Commun 15, 1511 (2024).

9. M. Baroncini et al., Making and Operating Molecular Machines: A Multidisciplinary Challenge. ChemistryOpen 7, 169-179 (2018).

10. T. Zhang, H. Liu, The motive forces in DNA-enabled nanomachinery. iScience 27, 109453 (2024).

11. P. F. Nilsson et al., Walking by design: how to build artificial molecular motors made of proteins. Nano Futures 8, 042501 (2024).

12. H. Linke, B. Höcker, K. Furuta, N. R. Forde, P. M. G. Curmi, Synthetic biology approaches to dissecting linear motor protein function: towards the design and synthesis of artificial autonomous protein walkers. Biophys Rev 12, 1041-1054 (2020).

13. E. H. C. Bromley et al., The Tumbleweed: towards a synthetic protein motor. Hfsp Journal 3, 204-212 (2009).

14. B. Zakeri et al., Peptide tag forming a rapid covalent bond to a protein, through engineering a bacterial adhesin. Proceedings of the National Academy of Sciences of the United States of America 109, E690-697 (2012).

15. R. D. Astumian, Thermodynamics and kinetics of a Brownian motor. Science 276, 917-922 (1997).

16. D. Schneidman-Duhovny, M. Hammel, J. A. Tainer, A. Sali, FoXS, FoXSDock and MultiFoXS: Single-state and multi-state structural modeling of proteins and their complexes based on SAXS profiles. Nucleic Acids Res 44, W424-429 (2016).

17. J. Jumper et al., Highly accurate protein structure prediction with AlphaFold. Nature 596, 583-589 (2021).

18. S. Kovacic et al., Design and Construction of a One- Dimensional DNA Track for an Artificial Molecular Motor. J Nanomater, 109238 (2012).

19. N. J. Kuwada et al., Tuning the performance of an artificial protein motor. Physical Review E 84, 031922 (2011).

20. N. J. Kuwada, G. A. Blab, H. Linke, A classical Master equation approach to modeling an artificial protein motor. Chem Phys 375, 479-485 (2010).

21. C. S. Niman et al., Controlled microfluidic switching in arbitrary time-sequences with low drag. Lab Chip 13, 2389-2396 (2013).

22. S. Sudhakar et al., Germanium nanospheres for ultraresolution picotensiometry of kinesin motors. Science 371, (2021).

23. T. Yanagida, M. Iwaki, Y. Ishii, Single molecule measurements and molecular motors. Philos Trans R Soc Lond B Biol Sci 363, 2123-2134 (2008).

24. S. L. Reck-Peterson et al., Single-molecule analysis of dynein processivity and stepping behavior. Cell 126, 335-348 (2006).

25. F. Mathieu et al., Six-helix bundles designed from DNA. Nano Lett 5, 661-665 (2005).

26. A. Spang et al., Complex archaea that bridge the gap between prokaryotes and eukaryotes. Nature 521, 173-179 (2015).

27. J. Vosseberg et al., The emerging view on the origin and early evolution of eukaryotic cells. Nature 633, 295-305 (2024).

28. F. J. Kull, R. D. Vale, R. J. Fletterick, The case for a common ancestor: kinesin and myosin motor proteins and G proteins. Journal of muscle research and cell motility 19, 877-886 (1998).

29. L. Li, J. O. Fierer, T. A. Rapoport, M. Howarth, Structural analysis and optimization of the covalent association between SpyCatcher and a peptide Tag. J Mol Biol 426, 309-317 (2014).

30. C. S. Chen, A. White, J. Love, J. R. Murphy, D. Ringe, Methyl groups of thymine bases are important for nucleic acid recognition by DtxR. Biochemistry 39, 10397-10407 (2000).

31. M. Thepaut et al., Crystal structure of the coiled-coil dimerization motif of geminin: structural and functional insights on DNA replication regulation. J Mol Biol 342, 275-287 (2004).

32. C. W. Garvie, S. E. Phillips, Direct and indirect readout in mutant Met repressor-operator complexes. Structure 8, 905-914 (2000).

33. Y. Y. He et al., Structural and functional studies of an intermediate on the pathway to operator binding by Escherichia coli MetJ. J Mol Biol 320, 39-53 (2002).

34. E. Tully, R. O'Kennedy, in Encyclopedia of Microfluidics and Nanofluidics, D. Li, Ed. (Springer US, Boston, MA, 2008), pp. 737-749.

35. N. M. Kirby et al., A low-background-intensity focusing small-angle X-ray scattering undulator beamline. Journal of Applied Crystallography 46, 1670-1680 (2013).

36. T. M. Ryan et al., An optimized SEC-SAXS system enabling high X-ray dose for rapid SAXS assessment with correlated UV measurements for biomolecular structure analysis. Journal of Applied Crystallography 51, 97-111 (2018).

37. K. Manalastas-Cantos et al., ATSAS 3.0: expanded functionality and new tools for small-angle scattering data analysis. J Appl Crystallogr 54, 343-355 (2021).

38. M. V. Petoukhov et al., New developments in the ATSAS program package for small-angle scattering data analysis. J Appl Crystallogr 45, 342-350 (2012).

39. M. Mirdita et al., ColabFold: making protein folding accessible to all. Nat Methods 19, 679-682 (2022).

40. E. F. Pettersen et al., UCSF Chimera--a visualization system for exploratory research and analysis. J Comput Chem 25, 1605-1612 (2004).

41. M. Jeeves, P. D. Evans, R. A. Parslow, M. Jaseja, E. I. Hyde, Studies of the Escherichia coli Trp repressor binding to its five operators and to variant operator sequences. Eur J Biochem 265, 919-928 (1999).

42. R. Sharma, A. S. Patelli, L. De Bruin, J. H. Maddocks, cgNA+web : A Visual Interface to the cgNA+ Sequence-dependent Statistical Mechanics Model of Double-stranded Nucleic Acids. J Mol Biol 435, 167978 (2023).

43. L. De Bruin, J. H. Maddocks, cgDNAweb: a web interface to the cgDNA sequence-dependent coarse-grain model of double-stranded DNA. Nucleic Acids Res 46, W5-w10 (2018).

44. W. L. DeLano, The PyMOL User's Manual. (DeLano Scientific, San Carlos, CA, USA, 2002).

45. N. Rezaei, A. Lyons, N. R. Forde, Environmentally Controlled Curvature of Single Collagen Proteins. Biophysical journal 115, 1457-1469 (2018).

46. S. Hertel et al., The stability and number of nucleating interactions determine DNA hybridization rates in the absence of secondary structure. Nucleic Acids Res 50, 7829-7841 (2022).

47. P. Schuck, H. Zhao, The role of mass transport limitation and surface heterogeneity in the biophysical characterization of macromolecular binding processes by SPR biosensing. Methods Mol Biol 627, 15-54 (2010).

48. S. Wang, G. M. Poon, W. D. Wilson, Quantitative Investigation of Protein-Nucleic Acid Interactions by Biosensor Surface Plasmon Resonance. Methods Mol Biol 1334, 313-332 (2015).

49. P. G. Stockley, B. Persson, Surface plasmon resonance assays of DNA-protein interactions. Methods Mol Biol 543, 653-669 (2009).

50. T. D. Pollard, A guide to simple and informative binding assays. Mol Biol Cell 21, 4061-4067 (2010).

51. E. C. Hulme, M. A. Trevethick, Ligand binding assays at equilibrium: validation and interpretation. Br J Pharmacol 161, 1219-1237 (2010).

52. D. Wu, G. Piszczek, Standard protocol for mass photometry experiments. Eur Biophys J 50, 403-409 (2021).

53. S. van der Walt et al., scikit-image: image processing in Python. PeerJ 2, e453 (2014).

54. T. Lindeberg, Scale Selection Properties of Generalized Scale-Space Interest Point Detectors. Journal of Mathematical Imaging and Vision 46, 177-210 (2013).

55. P. Virtanen et al., SciPy 1.0: fundamental algorithms for scientific computing in Python. Nat Methods 17, 261-272 (2020).

56. J. F. Marko, S. Cocco, The micromechanics of DNA. Physics World 16, 37 (2003).

Acknowledgments:

This project has evolved over two decades with input from many scientists. We acknowledge the contributions made by group members**:** Martin Zuckermann, Olivier Laprévote, Nathan Kuwada, Ben Lopez, Aimee Boyle, Drew Thompson, Laleh Samii, Suzana Kovacic, Ivan Unksov, Hasti Iranmanesh, James Walsh, Christopher Angstmann, Jowa Chan, Richard Sessions, Marc Bruning, Lara Small, Ben Clifton, Martina Balaz and Michael Konopik. We also thank Nicholas Dixon, Lisanne Spenkelink, Daniela Stock, Jonas Tegenfeldt, Lawrence Lee, Sophie Hertel and Slobodan Jergic for advice and guidance at various stages of this project. The authors acknowledge the use of facilities in the Structural Biology Facility within the Mark Wainwright Analytical Centre-UNSW. We thank UNSW Single Molecule Science for generous access to SPR facilities.

Funding:

Human Frontiers in Science Program grant RGP0031/2007 (HL, PMGC, NRF, DNW)

European Union's Horizon 2020 research and innovation programme under grant agreement No 951375 (ArtMotor) (HL, BH, PMGC)

Australian Research Council Discovery Project grant DP170103153 (PMGC, TB)

Natural Sciences and Engineering Research Council of Canada (NSERC) grants RGPIN-2015-05545, RGPIN-2020-04680 (NRF)

Swedish Research Council project numbers 2015-03824 (HL), 2020-04226 (HL), 2020-05266 (RE), 638-2013-9243 (RE).

Author contributions:

Conceptualization: RBD, DNW, NRF, HL, PMGC

Methodology: PN, NOR, NG, RBD, CWL, AL, RE, CSN, GAB, EHCB, AEW, APD, JPB, PJ, TB, BH, DNW, NRF, HL, PMGC

Investigation: PN, NOR, NG, RBD, CWL, AL, RE, AEW, APD, BH, DNW, NRF, HL, PMGC

Visualization: PN, NOR, NG, RBD, CWL, AL, RE, AEW, APD, NRF, HL, PMGC

Funding acquisition: RE, TB, BH, DNW, NRF, HL, PMGC

Project administration: BH, DNW, NRF, HL, PMGC

Supervision: BH, DNW, NRF, HL, PMGC

Writing – original draft: PN, NOR, NG, CWL, HL, PMGC

Writing – review & editing: PN, NOR, NG, RBD, CWL, AL, RE, CSN, GAB, EHCB, AEW, APD, JPB, PJ, TB, BH, DNW, NRF, HL, PMGC

Competing interests: Authors declare that they have no competing interests.

Data and materials availability: SAXS data have been deposited with the Small Angle Scattering Biological Data Bank (SASBDB) with the accession code SASDWM8. All other data are available in the main text or the supplementary materials.

Supplementary Materials

Materials and Methods

Supplementary Text S1 to S3

Figs. S1 to S20

Tables S1 to S8

References (*1–56*)

Fig. 1 The Tumbleweed (TW). (**A**) The construction of TW from two homodimeric proteins, TW1 and TW2. TW1 comprises the DtxR repressor followed by a geminin helix and finally the SpyCatcher protein. TW2 comprises the TrpR repressor followed by a geminin helix, the SpyTag, another geminin helix and finally the MetJ repressor. On combining TW1 and TW2, the SpyTag/SpyCatcher system covalently assembles TW. The 10 nm scale bar only refers to (**A**). (**B**) A DNA track for TW has an ordered series of cognate binding sites for each of the three TW feet (shown docked): *trpR-dtxR-metJ*, which can repeat indefinitely. The sites are arranged so that TW remains on the same face of the DNA track. (**C**) The mechanism of TW. The top panel shows TW bound by the TrpR foot (cyan) in the presence of tryptophan (Trp, blue triangle). On adding Co^{2+} (pink square), the DtxR foot (magenta) also binds to the track adjacent to the TrpR foot (second panel). As the concentration of Trp decreases, the TrpR foot releases the track (third panel). On addition of S-adenosyl methionine (SAM; orange circle), the MetJ foot (brown) binds adjacent to the DtxR foot (fourth panel). As the concentration of Co^{2+} decreases, the DtxR foot is released (fifth panel). Finally, on re-addition of Trp, the TrpR foot binds adjacent to the MetJ foot (bottom panel). Note, TW and the track are designed so only one site is reachable for each binding foot (e.g. near but not far *trpR* site in the bottom panel).

Fig. 2 Assembly and characterization of TW. (**A**) SDS-PAGE showing the covalent TW assembly and TW assembly after size exclusion chromatography (TW sized). TW sized protein was used for all remaining experiments. (**B**) Mass photometry characterization of purified TW1, TW2 and TW. Inset are the theoretical and observed molecular weights (mean ± SEM, $n = 4$) for each protein species. (**C**) SEC-SAXS analysis of TW. Top right shows the SAXS intensity as a function of q (momentum transfer in Å^{-1}). A Guinier plot of the data is shown in the inset. The data were fit to an ensemble of models using the program Multi-FoXS. The best fit (top right, red line) as determined by χ^2 was obtained using two conformers, one L-shaped (top left) and one extended (bottom left), weighted at 0.725 and 0.275, respectively. The error-weighted residual plot (lower right panel) shows no significant systematic deviations between the model and the experimental data. (**D**) AFM image showing eight TWs bound in the presence of Trp to eight repeats of a *metJ-trpR-dtxR* track inserted between flanking sequences.

Fig. 3 TW-DNA track interaction. (**A**) A panel of SPR sensorgrams titrating 5, 10, 20, 40 and 80 nM TW onto DNA oligonucleotides containing single cognate sites (*metJ*, *dtxR* and *trpR* sites) in the presence of the various controlling ligands (no ligands, SAM, Co^{2+} or Trp). (**B**) Representative binding curves fitting the Hill equation to average steady state SPR responses. Curves show binding to single site tracks containing individual cognate sites (brown – *metJ*; magenta – *dtxR*; cyan – *trpR*) for both TW (circles) and individual repressor dimers (triangles) in the presence of the controlling ligand. (**C**) Mass photometry data comparing the distribution of estimated molecular masses for the complex of TW and a duplex DNA containing the *metJ* and *trpR* sites (TW:*metJ-trpR* complex) in the absence of ligands (gray) and in the presence of SAM and Trp (orange). The estimated molecular mass of 128 ± 2 kDa in the absence of ligands, and 183 ± 3 kDa in the presence of SAM and Trp are in excellent agreement with the predicted masses of 133 and 183 kDa, respectively. (D) Apparent half-lives of different TW:DNA complexes as calculated from fitted k_{off} values determined by SPR experiments in which TW was bound to DNA in the presence (filled circles) or after removal (open circles) of ligands. Dissociation was triggered by adding an excess of free competing specific DNA in the presence or absence of ligands. Controlling ligands correspond to the respective DNA tracks, as indicated. Circles represent the individual values from replicate experiments.

Fig. 4 TW stepping on a DNA track observed by single molecule FRET. (A) Four-site (*trpR-dtxR-metJ-trpR* bottom to top) DNA track attached via biotin-streptavidin linker to a biotinylated PLL-g-PEG passivated surface. The *trpR*-sites at the bottom and top are labelled with different FRET acceptors (the yellow emitter Atto565 and the red emitter Atto647N, respectively). The TrpR foot is labelled with the FRET donor AlexaFluor488, a green emitter. **(B-D)** Expected

TW:DNA complexes formed in presence of pairs of ligands as indicated. (Trp + Co^{2+} in (**B**), SAM + Co^{2+} in (**C**), SAM + Trp in (**D**)). Each of the three complexes is designed to yield a distinct fluorescence signal on excitation with a 488 nm laser: (**B**) FRET to Atto565 (yellow emission) when the TrpR foot is bound to the bottom-*trpR* site, (**C**) AlexaFluor488 (green emission) from the free TrpR foot, and (**D**) FRET to Atto647N (red emission) when TrpR is bound to the top-*trpR* site. (**E**) Microfluidic device used to rapidly change solution conditions in the imaging volume. (**F-I**) **Experiment 1**: two-colour imaging with emission split at 543 nm to simultaneously record AlexaFlour488 emission (shown as green) and ATTO565 plus ATTO647N emission (shown in red). (**F**) Schematic of expected TW position and false colour representation of expected fluorescence signal given solution changes (dashed lines). (**G**) Normalized example single-molecule trace. (**H**) Kymographs of all detected colocalizing single molecules sorted by time of last identified fluorescence. (**I**) Sum of all single-molecule traces in the respective experiment. (**J-M**) **Experiment 2**: same as in (**F-I**), but using two-colour imaging with emission split at 633 nm to simultaneously record ATTO565 emission (FRET emission from the bottom *trpR* site, shown in cyan) and ATTO647N emission (FRET emission from the top *trpR* site, shown in magenta).

Supplementary Materials for

Tumbleweed: an artificial motor protein that walks along a DNA track

Authors: Patrik Nilsson[1,2†], Neil O. Robertson[3,4†], Nils Gustafsson[1,2†], Roberta B. Davies[3†], Chu Wai Liew[3,4], Aaron Lyons[5], Ralf Eichhorn[6], Cassandra S. Niman[1,2], Gerhard A. Blab[5‡], Elizabeth H.C. Bromley[7], Andrew E. Whitten[8], Anthony P. Duff[8], Jason P. Beech[1,2], Peter Jönsson[1,9], Till Böcking[10,11], Birte Höcker[12], Derek N. Woolfson[13,14,15], Nancy R. Forde[5], Heiner Linke[1,2*], Paul M. G. Curmi[3,4*]

Corresponding author: heiner.linke@ftf.lth.se, p.curmi@unsw.edu.au

The PDF file includes:

    Materials and Methods
    Supplementary Text S1 to S3
    Figs. S1 to S20
    Tables S1 to S8
    References

Other Supplementary Materials for this manuscript include the following:

Materials and Methods

Construct design and cloning

Protein constructs were based on the following proteins: DtxR (UniProt Acc. No. P0DJL7), Geminin (UniProt Acc. No. E2QRF9), TrpR (UniProt Acc. No. P0A881), MetJ (UniProt Acc. No. P0A8U6) and the SpyTag:SpyCatcher system (*29*). For the repressor and Tumbleweed (TW) constructs DtxR refers to residues 2—121 with a C102D mutation (based on PDB: 1F5T (*30*)), Geminin$_{Coiled-Coil}$ refers to residues 110—145 with an additional N-terminal threonine and C-terminal glutamine (based on PDB: 1T6F (*31*)), TrpR refers to residues 2—108, and MetJ refers to residues 2—105 with a Q45K mutation (based on PDB: 1MJM (*32, 33*)).

Both MetJ and TrpR were expressed from pET28a vectors as fusion proteins with a N-terminal 8 x His-tag. DtxR was expressed from a pET19b vector as a fusion protein with a C-terminal 6 x His-tag.

TW was produced from two separate constructs, TW1 and TW2 (Fig. S13). TW1 consists of DtxR—Geminin$_{Coiled-Coil}$—SpyCatcher. TW2 consists of TrpR—Geminin$_{Coiled-Coil}$—SpyTag—Geminin$_{CoiledCoil}$—MetJ. For both TW1 and TW2 the protein domains were connected using flexible glycine-serine-rich linkers. Both TW1 and TW2 constructs were synthesized by Genescript within a pET15b vector allowing for the expression of protein with an N-terminal 8 x His-tag and TEV protease cleavage site.

Protein expression and purification

All proteins were expressed in *E. coli*. Individual repressors and TW1 were expressed using BL21 cells in 1–2 L of LB media. TW2 was poorly expressed under these conditions. After screening expression in different cell and media types TW2 was eventually expressed using C41 cells in 6–12 L of 2xTY media. Cells were grown with agitation at 37 °C until the culture reached $OD_{600} \sim 0.6$, at which point the incubation temperature was reduced to 24 °C and protein expression was induced by the addition of IPTG to a final concentration of 0.4 mM. Cultures were then incubated with agitation for approximately 20 h, after which cells were harvested by centrifugation and pellets stored at –80 °C until further processing.

Cell pellets were resuspended at 4 °C with stirring in 20 mM Tris pH 8, 150 mM NaCl, 20 mM Imidazole, 10% glycerol, 2.5 mM $MgCl_2$, 0.5 mM $CaCl_2$, 1 mM NaN_3, 0.1 mg/mL lysozyme, 10 µg/mL DNase I and 1 x protease tablet (Sigma-Aldrich). Approximately 25 mL of lysis buffer per liter of cell culture was used to resuspend the cell pellets. The resuspended cell slurry was then lysed using a cell disruptor at 20 kPSI pressure. The soluble fraction was harvested by centrifugation by 50 000 x g and then loaded onto a 5 mL HisTrap^{TM} HP nickel affinity chromatography column (Cytiva). The column was then subjected to a series of washes using a combination of buffer A (20 mM Tris pH 8, 400 mM NaCl, 10% glycerol, 1 mM NaN_3) and buffer B (buffer A + 1 M imidazole); first with 2% buffer B, then 5% buffer B, and then finally an elution step with 30% buffer B.

TW1 and TW2 were subjected to TEV cleavage and reverse His-Trap purification to increase the purity of the final product. After the initial His-trap purification, the eluted proteins were concentrated and diluted to a final concentration of 3% buffer B. This sample was then subjected to an overnight TEV protease cleavage (1:50 mg/mg TEV:protein), with DTT added to the sample to a final concentration of 1 mM. The TEV-cleaved sample was then flowed over the HisTrap^{TM} column to remove contaminants, collecting the flowthrough.

Finally, all proteins were subjected to size-exclusion chromatography (SEC) using either a HiLoad Superdex 75 26/60 column for individual repressors, or a HiLoad 200 16/600 column for TW1 and TW2 (Cytiva). SEC was performed in 20 mM Tris pH 8, 400 mM NaCl, 10% glycerol, 1 mM NaN$_3$ and 1 mM EDTA. Protein yields per cell culture volume were typically ~ 1 mg/L for individual repressors, ~4 mg/L for TW1 and ~0.3 mg/L for TW2. Purified protein was then frozen using liquid nitrogen and stored at –80 °C until further processing.

Assembly and purification of Tumbleweed

Purified TW1 and TW2 were mixed with a molar ratio of 1.2:1 (with TW2 at a final concentration of ~20–40 µM) in 20 mM Tris pH 8, 400 mM NaCl, 10% glycerol, 1 mM NaN$_3$ and 1 mM EDTA at 4 °C for 16 h. The assembly mixture was then subjected to SEC using a Superdex 200 Increase 10/300 column (Cytiva), using the same buffer as the assembly. Purified assembled Tumbleweed was then frozen in liquid nitrogen and stored at –80 °C until further processing.

Fluorescent labelling of TW

Fluorescent TW was produced by thiol conjugation of a maleimide dye to a TW containing a cysteine mutation. A TW2 construct containing TrpR$_{S107C}$ (henceforth referred to as TW2$_{S107C}$) was synthesized by Genscript. The protein was expressed and purified as described above, with 1 mM DTT included in purification buffers to maintain the protein in a reduced state.

Alexa Fluor™ 488 C$_5$ Maleimide (AF488; Thermofisher Scientific) was resuspended to 10 mM in DMSO. Purified TW2$_{S107C}$ was reduced with 1 mM TCEP pH 7 prior to labelling at a final concentration of 30–60 µM with 5–10x molar excess of maleimide dye. Reactions were performed in 50 mM Tris pH 8, 400 mM NaCl, 10% (v/v) glycerol either for 2 h at room temperature or overnight at 4°C. Reactions were quenched with 50 mM DTT for 30 min at room temperature. TW1 was added to the mixture such that the final molar ratio of TW1:TW2 was 1.2:1, and left to assemble TW as described above. The assembly mixture was purified from excess dye by SEC using a Superdex 200 Increase 10/300 column (Cytiva), in the same buffer as the assembly.

Following removal of excess dye, the degree of protein labelling was measured using the UV-Vis function of an ND-1000 Nanodrop Spectrophotometer (ThermoFisher Scientific). The absorbance of the protein:dye conjugate was measured at 280 nm (protein) and at 495 nm (AF488) (See Table S4). The degree of labelling was calculated according to the equation (*34*):

$$\text{Degree of Labelling} = \frac{\epsilon_{\text{Protein}} \cdot A_{495}}{(A_{280} - CF_{280} \cdot A_{495}) \cdot \epsilon_{\text{Dye}}}. \qquad 1$$

Here, $\epsilon_{\text{Protein}}$ is the extinction coefficient of the protein at 280 nm (55350 M^{-1} cm^{-1}), ϵ_{Dye} is the extinction coefficient of AF488 at 495 nm (73000 M^{-1} cm^{-1}) and CF_{280} is the correction factor for the absorption of light at 280 nm by AF488 (0.11). TW was assessed to be ~100% labelled (two labels per dimeric TrpR foot), and frozen with liquid nitrogen and stored at -80°C.

SAXS data collection

Small-angle X-ray scattering (SAXS) data were collected on the small/wide angle X-ray scattering beamline at the Australian Synchrotron (*35*), part of ANSTO in Melbourne, Australia.

The camera length used was 2680 mm corresponding to a q-range of 0.005 – 0.50 Å^{-1}. Proteins were auto-loaded from a 96-well plate and 50 mL of ~10 mg/mL of sample injected onto a Superdex 200 Increase 5/150 GL equilibrated in 10 mM HEPES pH 7.4, 150 mM NaCl, 5mM MgCl$_2$, connected in line with a coflow cell (*36*) through which X-rays were passed at 8 x 10^{12} photons per second at 11.5 keV. The data was collected on a Pilatus 2M detector (Table S5).

SAXS data processing
 SAXS data were processed using software from the ATSAS suite (*37, 38*) (Table S5). Protein elution profiles were generated from scattering intensities using custom written scripts. The elution profiles were used to determine the signals corresponding to protein and buffer frames. Firstly, 21 buffer frames were selected and averaged for buffer subtraction. The radius of gyration (R_g) was then determined for each frame across the protein peak (with buffer subtraction performed for each individual frame). Guinier analysis to determine R_g was performed in PRIMUS (*37*) to monitor data quality. Distance distribution ($P(r)$) curves were generated using GNOM (*37*) to estimate the maximum dimension (D_{max}) of the protein.

Multi-state modelling with MultiFoXS
 Multi-state modelling using MultiFoXS (*16*) was used to determine the population-weighted conformational states of TW that could be contributing to the experimental SAXS profile. The inputs used for MultiFoXS included a protein model, SAXS profile, flexible hinge residues and rigid body connections. To obtain protein models to use as templates for MultiFoXS, models of TW1 and TW2 were generated using ColabFold v1.4: AlphaFold2 using MMseqs2 (*39*). These Alphafold-generated models were subsequently used to construct a complete model for TW assembly using UCSF Chimera (*40*). For MultiFoXS, TW was divided into rigid elements and hinge regions. The rigid elements comprise: the three repressor dimers, the three coiled-coils and the SpyTag/SpyCatcher hub. Six hinges were placed in the Gly-Ser rich linkers. To simplify the computation by MultiFoXS, the polypeptide chain was cut in one of the two Gly-Ser rich linkers joining each rigid element. Thus, the six hinges were defined as: TW1: Gly125-Gly126 and Gly170-Gly171, between DtxR, the coiled-coil and SpyCatcher, respectively; and TW2: Ser114-Gly115, Ser155-Ser156, Gly183-Ser184, Gly228-Gly229, between TrpR, the coiled-coil, SpyTag, coiled-coil and MetJ, respectively (see Fig. S13).
 Preliminary tests were initially conducted with conformational sampling of 100 conformations and final calculations were performed to sample 10,000 conformations. For each sampled conformation, a SAXS profile is calculated followed by scoring of multi-state models to generate population-weighted ensembles with associated R_g. The best multi-state models that fit the experimental SAXS data were based on the χ^2 values and residual plots.

Design and assembly of DNA tracks
 The *metJ* cognate sequence (GAGACGTCTC) was taken directly from the consensus Met box sequence from the MetJ:DNA crystal structure (PDB: 1MJM (*32*)). The *trpR* cognate sequence (GTACTCGCTAGCGAGTAC) was based on a *trpR^S* sequence previously shown to bind TrpR dimer with a 1:1 stoichiometry (*41*). The *dtxR* cognate sequence (TTAGGTTAACCTAA) was based on a DtxR:DNA crystal structure (PDB: 1F5T (*30*)), with the sequence altered to produce a palindrome that is capable of binding only one DtxR dimer.

 Flanking sequences were designed to create blocks that could be rearranged within a multisite track that maintains consistent sequences either side of the cognate sites. The *metJ* flanking sequences were designed based on the native *E. coli metC* protomer sequence (GenBank

Ref. Seq. NC_000913.3.). This promoter sequence was chosen on the basis that it contains a single Met box (*32*). The *dtxR* flanking sequences were conserved from the DtxR:DNA crystal structure (PDB: 1F5T (*30*)). The *trpR* flanking sequences were designed by a genetic algorithm to minimize the presence of cryptic repressor binding sites (*18*). Double, triple and quadruple site tracks were designed by arranging the cognate sites and their flanking sequences. Models of the tracks were produced using cgNA+web (*42, 43*) and analyzed using PyMOL (*44*) to ensure the cognate sites were appropriately distanced from one another and remained in the same phase (i.e. on the same side of the track).

All tracks were ordered as single strand oligonucleotides from Integrated DNA Technologies (IDT). Oligonucleotides were resuspended to 100 µM in Milli-Q water and stored at –20 °C. Complementary oligonucleotides were annealed at concentrations of up to 20 µM in 25 mM HEPES pH 7.4, 200 mM KCl, 5 mM $MgCl_2$ by first heating the sample to 95 °C for 2 min, and then subjecting to a 95–20 °C gradient at 1°C/min. Annealed tracks were stored at -20 °C until further processing. Table S6 lists all DNA tracks used in SPR and mass photometry experiments.

Atomic force microscopy (AFM)

E. coli containing the pK8 plasmid (*18*) were cultured in LB media overnight in the presence of 25 mg/mL kanamycin. They were pelleted by centrifugation for 3 min at 12000 rpm. The plasmid was purified using a MiniPrep kit (Qiagen). The DNA was linearized by digestion with EagI (nine parts purified plasmid to one part NEBuffer 3, with 1 unit of EagI restriction endonuclease for every 150 ng of DNA) for 2 h at 25ºC, after which the enzyme was heat inactivated by incubation at 65 ºC for 25 min. The product was purified using a PCR Cleanup kit (Qiagen). Completion of restriction digest was confirmed by gel electrophoresis.

TW (0.26 mg/ml) and 10X HEPES buffer (40 mM HEPES, 100 mM NaCl, 20 mM $MgCl_2$, pH 7.4) were both diluted five-fold in Milli-Q water (1:1:3 TW:10X HEPES:H_2O). 2.5 µL of this stock was placed into a pre-lubricated microcentrifuge tube, to which was added 1.25 µL of linearized DNA (69.3 ng/mL) and 1.25 µL of 40 mM tryptophan. This mixture was incubated at room temperature on a spinner for 1 hour. After this time, 45 µL of 1X HEPES Buffer was added to the incubated solution, mixed with a pipette, and plated immediately onto freshly cleaved mica, where it was allowed to sit for 4 minutes. Following this, the excess solution was shaken off, and the plate was rinsed five times with 1 mL of Milli-Q water, then dried gently under a stream of filtered compressed air to remove any residual liquid.

AFM images were recorded using an Asylum Research MFP-3D SPM, using Mikromasch HQ:NSC15/AL BS probes with a spring constant of 40 N/m and resonance frequency of 325 kHz.

The chain-tracing software SmarTrace (*45*) was used to determine the contour lengths of DNA flanking the bound TW proteins. For this analysis, *N*=5 images were chosen that included seven or eight TW bound to clearly define the inserted track cassette. The cassette is located asymmetrically with respect to the ends of the linearized plasmid, and thus the long and short contour lengths, and their ratio of lengths, were used to confirm that the TW were bound in the expected region of the DNA.

Surface Plasmon Resonance (SPR)

SPR experiments were performed on an Biacore™ S200 instrument (Cytiva). All experiments were performed in running buffer (25 mM HEPES, pH 7.4, 200 mM KCl, 5 mM MgCl$_2$, 0.2 mM EDTA, 0.05% (v/v) Tween-20), which was passed through a 0.22 µm filter and degassed by vacuum for at least 30 min prior to use. Data was collected at 40 Hz in multi detection mode, with the instrument temperature set to 25 °C. All sensorgrams were double referenced.

DNA coating of SPR chips was prepared as previously described (*46*). Series S CM5 sensor chips (Cytiva) were initially prepared by depositing ~5000 RU of streptavidin on to the chip surface via amine coupling. A biotinylated single-stranded oligonucleotide (henceforth referred to as the biotin anchor) was added to the chip for a response of ~50–70 RU. Finally, annealed single or double cognate site DNA containing a single-stranded overhang complementary to the biotin anchor was flowed over the non-reference channels for a response of ~10 RU. Chips were regenerated using injections of 10 mM glycine pH 2.5, which removed the annealed DNA and made the biotin-anchor re-available to bind new DNA.

Individual repressors and TW were titrated onto single and double site DNA at 100 µL/min. When required, ligands were used with the following concentrations: 1 mM SAM, 0.4 mM CoCl$_2$ and 0.5 mM Trp. All titrations were performed using the A-B-A injection method, with a 30 s initial ligand injection, a contact injection containing protein for 15–30 s, and a final ligand injection of 40–60 s to trigger dissociation. A 60 s injection of buffer containing 0.5 M KCl was used to remove any residual protein from the surface before the next titration point. The average steady state positions were determined by the Biacore Evaluation software (Cytiva) and fit to a Hill equation using Prism 10 (Graphpad):

$$RU = \frac{RU_{Max} * [Protein]^h}{K_d{}^h + [Protein]^h}. \qquad\qquad 2$$

Here, K_d is the apparent affinity, h is the Hill coefficient and RU_{Max} is the fitted maximum binding response.

We were unable to fit the above titrations to a simple 1:1 kinetic model. Increasing the amount of DNA on the chip surface lead to sensorgrams with slower apparent binding kinetics, indicating the system suffered from mass transfer effects (*47*). Protein:DNA interactions are particularly prone to mass transfer in SPR as electrostatically steered interactions lead to high association rates (*48*). A consequence of mass transfer is that during the dissociation phase of an SPR experiment, the protein can release and rebind to the surface before it enters the bulk solution, resulting in a slower apparent dissociation rate (*48, 49*).

To avoid complications arising from TW rebinding DNA on the SPR surface, we decided to study TW:DNA dissociation kinetics using a SPR-based displacement assay which was based on classic tracer experiments (*50, 51*). This involved dissociating TW in the presence of free DNA in solution, such that TW would preferentially bind to the free DNA and not rebind to the surface. A dual injection method was used to first load TW onto the surface DNA with an immediate second injection used to trigger dissociation. The assay was performed using a 100 µL/min flow rate. For single-site DNA experiments, a 5 nM TW solution was used to bind TW to DNA on the SPR chip in the relevant ligand for 25 s, before dissociation of the TW:DNA

complex was triggered with an immediate 60s injection of a solution containing 1 µM free DNA ± ligand. The free DNA contained the relevant cognate binding site but lacked the single stranded anchor overhang and was thus unable to bind the sensor chip. For experiments involving double-site DNA, 1 nM TW was loaded onto the DNA in the presence of two ligands for 60 s before dissociation was triggered with a 60 s injection of a solution containing both ligands and 1 µM of each of the relevant free DNA sequences. The resulting dissociation curves were fit to a single exponential decay function using Prism (Graphpad):

$$RU = (RU_{Initial} - RU_{NS}) * e^{-k_{off}*t} + RU_{NS}. \qquad\qquad 3$$

Here, $RU_{Initial}$ is the initial response value, RU_{NS} is the non-specific response value, and k_{off} is the apparent dissociation rate constant. For double-site DNA, double-ligand experiments, RU_{NS} was set to 0.

Mass photometry

Mass photometry experiments were performed as described (52). Experiments were performed on an Refeyn TwoMP instrument using 24 x 50 mm high precision 1.5H coverslips (Marienfeld) and CultureWell™ gaskets (Grace Biolabs). Coverslips were first prepared by sonicating in a 50% (w/w) isopropanol bath for 3 min, followed by another 3 min sonication in MilliQ-Water. These sonication rounds were repeated for a total of 3 rounds before the coverslips were dried with nitrogen gas and stored at room temperature. Gaskets were affixed to the coverslip immediately prior to an experiment.

Buffers (25 mM HEPES, pH 7.4, 200 mM KCl, 5 mM MgCl_2, and 0.2 mM EDTA ± 0.5 mM SAM, 0.4 mM CoCl_2 and/or 0.5 mM Trp) were passed through a 0.22 µm filter prior to use in mass photometer experiments. Data were collected over a 60 s timeframe using the normal measurement mode and regular image size as set by the Refeyn AcquireMP software. Mass calibrations were performed using a combination of purified bovine serum albumin (Sigma-Aldrich) and NativeMark™ unstained protein standard ladder (ThermoFisher Scientific), with the 66, 132 and 480 kDa peaks providing a calibration curve with $R^2 > 0.99$. Manually picked mass distributions within the relevant mass ranges were selected and fit with a Gaussian function to determine an average molecular mass using the Refeyn DiscoverMP software. SAM buffer was found to produce an interference with the mass photometry signal. Additionally, DNA transiently interacted with the glass surface when in the presence of CoCl_2. Both conditions result in an additional mass peak centered around 60–80 kDa.

TW1, TW2 and TW samples were added to slides to a final concentration of 10 nM. TW and 4–10-fold molar excess DNA were preincubated for ~5 min before analysis on the mass photometer. Final concentrations of 25–30 nM TW and 250–300 nM DNA were used for single-site DNA experiments, whereas double-site DNA experiments used 5–25 nM TW and 50–250 nM DNA.

DNA tracks for single molecule observation

Single strand oligonucleotides used to generate DNA tracks (Table S7) for single molecule experiments were ordered from IDT. Oligonucleotides were resuspended to 100 µM in Milli-Q water and stored at –20 °C. All oligonucleotides required for the track were annealed at concentrations of 11.4 µM of each component oligo in 70 µL 1.25X PNK (T4 polynucleotide

kinase reaction) buffer (New England BioLabs, Cat. No. B0201S) by first heating the sample to 95 °C for 2 min, followed by a 95—20 °C gradient at 1°C/min. The annealed oligonucleotides were then ligated by adding 8 µL ATP (New England BioLabs, Cat. No. P0756S) to a final concentration of 1 mM and 2 µL T4 DNA ligase (New England BioLabs, Cat. No. M0202S) to a final concentration of 10,000 units/ml. The resulting 80 µL ligation reaction mixture in 1X PNK was kept at 16 °C for 16 h before heat inactivation at 65 °C for 10 min. The DNA was purified using bind-wash-elute to a commercial spin column (QIAquick PCR Purification Kit, Qiagen) as per manufacturer's instructions. Purified tracks were stored at 4 °C until use. Models of the tracks were produced using cgNA+web (*42, 43*) and analyzed using PyMOL (*44*) to ensure the cognate sites were appropriately distanced from one another and remained in the same phase.

Microfluidics for single molecule observation

To switch the TW running solutions, a previously published microfluidic device, developed by Niman *et al.* (*21*), was used (Fig. 4E; Fig. S14). This device is designed to allow switching between three solutions (three combinations of ligands) in arbitrary time order, with sub-second time resolution, and without stopping the flow. It consists of three inlet channels of equal length that intersect at the entrance to a primary observation channel where the TW stepping takes place. Each inlet has an "escape" channel of equal length, microfluidic resistance and pressure originating directly before the intersection and terminating at a common waste outlet (circle, Fig. S14A far left). These escape channels provide routes away from the observation channel. Increasing the flow rate in any one of the three inlet channels propagates that solution alone along the observation channel, while pushing the other two solutions away from the observation channel to their respective escape channels. This design allows for the solution in the observation channel to be changed rather than stopped, allowing for faster and more complete switching, as well as for switching the three fluids in arbitrary time order. Flow rates were controlled using Fluigent lineup pumps, flow unit and OxyGEN software (Fluigent). Flow rates of 7 µL/min in the "on" channel and 1 µL/min in the "off" channels provided good switching and arbitrary sequences of ligand solutions were achieved by the application of 7 µl/min to any of the inlet channels (and 1 µl/min to the others) (Fig. S14C).

The time resolution for switching the ligand concentration in immediate proximity to the channel surface where TW is located (the chemical switching time) is limited by the diffusion time of ligands in and out of the non-slip boundary layer. The chemical switching time inside the boundary layer depends on the fluid speed and the location along the microfluidic channel (*21*). For the experiments reported here, observations were conducted in the main observation channel, approximately 1 mm away from the intersection with the inlet lines. A 3D finite element simulation (incompressible, laminar flow) of the volumetric flow rates in the device was performed using COMSOL Multiphysics software with the fluid set as water, with a density of $1000 kg/m^3$ and a dynamic viscosity of 0.001 Pa s. For inlet flow rates of 7, 1 and 1 µl/min, the simulated volumetric flow rate in the observation channel of 2.285 µl/min which corresponds to a fluid speed in the observation channel of 7.6 mm/s. Based on the results in (*21*) (Fig. S14) we expect that the chemical switching time in the experiments reported here was about 0.2 s or less, much faster than the 7 s during which each solution was held constant.

Microfluidic master mold fabrication protocol

Microfluidic devices were produced by fabricating a mold in which polydimethylsiloxane (PDMS) could be cured and subsequently bonded to a glass surface. To fabricate the mold, a photomask was fabricated using an MLA150 maskless lithography system (Heidelberg

Instruments GmbH). A 4" silicon wafer was dried at 180°C for 5 min. A 50 µm thick dry film resist (SUEX®, K50, DJ Microlaminates) was applied to the wafer using a laminating machine (Catena 35, Acco UK Ltd) at 65 °C. A pre-exposure bake was performed at 85 °C on a hotplate (Model 1000-1 Precision Hot Plate, Electronic Micro Systems Ltd) to remove air bubbles and to relax the film. SUEX film was exposed at 365 nm in a contact mask aligner (Karl Suss MJB4 soft UV) for 27 s at a lamp power of 30 mW/cm^2. Post exposure baking was done at 85 °C for 5 min. Development was done in Mr DEV 600 (Micro Resist Technology GmbH) for 15 min plus 5 min in fresh developer followed by rinsing in flowing isopropyl alcohol and drying with nitrogen. A final bake was done in a convection oven at 200 °C for 15 min. To reduce adhesion of PDMS to the master, a layer of aluminum oxide (~1 nm) followed by a monolayer of perfluordecyltriklorosilan (FDTS) was deposited in an atomic layer deposition system (Fiji – Plasma Enhanced ALD, Veeco).

Microfluidic device preparation and single molecule observation
Base imaging buffer (25 mM HEPES, pH7, 200 mM NaCl, 5 mM MgCl$_2$) was passed through 0.22 µm filter and stored at 4 °C for no longer than 1 month.

Coverslips (40 mm x 20 mm, #1.5 thickness, VWR) were prepared by sonicating in isopropyl alcohol (Merck) for 10 min and blow dried with pressurized nitrogen. These were then cleaned with acid piranha solution of 3 parts sulfuric acid (Merck) and 1 part hydrogen peroxide (Merck) for 30 min at 80 °C after which they were rinsed with MilliQ water (Millipore MQ) and blow dried with pressurized nitrogen. Final cleaning was performed in oxygen-nitrogen mixed plasma (Zepto Plasma Cleaner, Diener Electronic) for 3 min at 40 kHz, 100W.

PDMS microfluidic devices were prepared by thoroughly mixing 2-component Sylgard 184 silicone elastomer and curing agent (G A Lindberg) in a 10:1 ratio and degassing to remove air bubbles. The mixture was poured onto the device master mold and allowed to cure at 80 °C for > 1h. After curing, using a 1 mm biopsy tool, inlets were punched at the three points indicated by inward pointing arrows (Fig. S14A). Using a 3 mm biopsy tool, outlets were punched at the circles (Fig. S14A) indicating the common waste outlet and the 10 mm mark along the primary observation channel. The molded PDMS was washed with isopropyl alcohol. Immediately after coverslip cleaning, the molded PDMS and coverslip contact surfaces were both etched in nitrogen plasma for 10 s and then bonded together by placing the molded PDMS onto the cleaned coverslip. Immediately, the device was wetted and then incubated for a minimum of 1h in imaging buffer with 1 mg/ml of 1:10000 PLL-g-PEG-Biotin (3.4 kDa PEG):PLL-g-PEG (2 kDa PEG). The device was visually inspected via white light transmission microscopy at 20X magnification for blockages, bonding defects, and bubbles. Devices were stored at RT submerged in imaging buffer.

Before use, the devices were washed with imaging buffer by aspirating from the common waste outlet while filling the observation channel outlet with buffer under 1 atmosphere pressure. The device channels were then incubated in imaging buffer containing 20 nM streptavidin (Merck) for at least 10 min. Glass slides were cut to size and glued to the ends of the coverslip using two component dental cement (Abberior Instruments) (Fig. S14B). Immediately before imaging, 0.2 mg/ml TROLOX (Merck) was added to the imaging buffer which was bubbled with nitrogen for 15 min to remove dissolved oxygen.

The microfluidic system was prepared by washing all microfluidic lines and flow meters with chlorine, MilliQ water, isopropyl alcohol (Merck) and finally with MilliQ water. The cleaned microfluidic system was loaded with three 2 ml reservoirs containing the ligand-pair solutions; imaging buffer, 2 mg/mL TROLOX (Merck) and

1) 0.5 mM Trp, 0.2 mM $CoCl_2$
2) 0.2 mM $CoCl_2$, 1mM SAM
3) 1mM SAM, 0.5 mM L-Trp.

For each control experiment, the microfluidic system was loaded with two reservoirs containing the same ligand-pair solution, either 1, 2, or 3, and the third reservoir containing imaging buffer, 2 mg/mL TROLOX (Merck) and no ligands.

Finally, the device was connected to the microfluidic system via the three 1 mm inlets and rinsed with running flows set to 7 µl/min for the solution desired in the observation channel and 1 µl/min for the two alternate solutions. After rinsing for a minimum of 1 min the device was mounted and secured to the microscope.

Single molecule fluorescence imaging was performed using a Nikon-TI2 inverted optical microscope in TIRF mode. A $100 \times$ TIRF objective (Plan-APOCHROMAT 100×1.45 NA Oil, Nikon) was used to collect fluorescence onto an sCMOS camera (Prime 95B, Photometrics), yielding a pixel size of 110 nm. Alternating laser excitation was provided by a laser combiner (LightHUB Ultra, Omicron) equipped with an acousto-optic tunable filter triggered by the camera digital trigger out via a multifunction I/O device (PCIe-6323, National Instruments). For excitation of AlexaFluor 488, ATTO565 and ATTO647N laser excitation was alternated between 488 nm, 561 nm and 640 nm at intensities on the order of 100 W cm^{-2}, 50ms exposure time per excitation wavelength producing 3 camera frames with no time delay between wavelengths. This sequence was repeated every 700 ms to generate a time sequence. Simultaneous imaging of two emission wavelength bands on two halves of the camera sensor was achieved using an image splitter (Optosplit II, Cairn). Two sets of filter combinations (Chroma) were used. For simultaneous imaging of AlexaFluor 488 emission and combined ATTO565 and ATTO647N emission, a ZT543rdc dichroic with ET570LP and ET525/50m emission filters. For simultaneous imaging of ATTO565 emission and ATTO647N emission a ZT633rdc dichroic with ET600/50m and ET700/75m emission filters.

In situ on the microscope, both outlets were emptied by aspirating. A 200 µl of a mixture of 1 nM TW and 200 pM DNA track was prepared at room temperature and immediately pipetted into the observation channel outlet at 1 atmosphere. All flows where set to 0.5 µl/min and a rolled kimwipe tissue paper (Kimtech) inserted into the common waste outlet to wick away waste buffer. The surface density of reagents was observed in the observation channel approximately 1 mm away from the intersection. Once single molecule binding on the surface began to reach 1 $µm^{-2}$, as judged by visual inspection, the flows were set to operating values of 7 µl/min (desired solution) and 1 µl/min (two alternative solutions), and the tissue was removed from the waste outlet. An unexposed field of view was selected, and the experiment was started. During the experiment, the solution present in the main observation channel was altered by changing the driving (7 µl/min) buffer every 7 s following the patterns indicated in Figs 4F, 4J, S7A, S8A and S9A. This was achieved in an automated fashion using Fluigent lineup pumps, flow units and OxyGEN software (Fluigent).

Single molecule FRET analysis

The pixel coordinates of bright spots (Fig. S15), independent of time point and color channel, were found using the sci-kit image package (*53*) as follows. For each color channel a maximum intensity, *I*, projection over time was calculated. For each of the three color channels this projection was normalized using the following equation:

$$I_{norm} = \frac{I - min\,(I)}{max\,(I - min(I))}. \qquad\qquad 4$$

From this normalized maximum intensity projection image, a scale-space volume is generated by computing the Laplacian of Gaussian filtered image with successively increasing standard deviation and generating a stack of images from the result. Single particles were detected using the local maxima in this scale-space volume (*54*). Detections close to the edges (35 pixels) of the field of view were removed.

The list of detected coordinate centers was filtered to include only detections that exhibited colocalization between the donor and FRET acceptor channels indicating the presence of both DNA and TW (Fig. S15). This was achieved by first identifying nearest neighbor pairs with one member in the donor channel and one in the FRET channel. Then this list of pairs of detections was filtered to only include nearest neighbors closer than 330 nm (3 pixels) of each other. The 330 nm distance threshold was chosen to allow for localization uncertainty, alignment of the optosplit and chromatic aberration.

For each colocalization that was identified between the FRET acceptor and donor channels a background subtracted intensity trace over time was extracted from the full unnormalized image sequence. The background subtracted intensity of the molecule was calculated for each frame by taking the average intensity of the pixels within a radius of 3 pixels (Fig. S16A) minus the median of the intensities in the pixels a radius of 4 pixels away (Fig. S16B). This was done for each frame and channel, resulting in the traces shown in Fig. S17. These traces were matched to the ligand solution in the observation channel at the time of the image acquisition by comparing exact start times and elapsed times from the pump log and metadata of the image. Any trace where the intensity of the first frame was either ~2.5 times greater than, or ~0.2 times less than, the average single fluorophore intensity was considered a spurious detection (for example aggregates or unidentified debris) and removed from the analysis. This only resulted in the removal of approximately 10% of colocalizing traces indicating successful preparation of a monodisperse 1:1 ratio of single molecule DNA:TW complexes fixed to the coverslip.

Traces were sorted based on the last detected falling edge in either color channel. Edges (Fig. S18A) were determined by Gaussian filtering with a standard deviation of 1.2 (Fig. S18B) and taking the first derivative (Fig. S18C). Peaks and troughs were identified using sci-pys peakfinder (*55*) using a threshold of 2.5 (Fig. S18C). Each trace was then normalized between its max value as 1 and the average value of the final 15 frames of the trace as 0. By mapping this normalized data to colors, sorted traces are shown in the kymographs in Figs 4H and L. In Experiment 1, 24-bit RGB color mapping was used with acceptor intensity represented by red and donor intensity by green. In Experiment 2, CMYK color space mapping was used with Atto647N (red) acceptor intensity represented by magenta and Atto565 (yellow) acceptor intensity represented by cyan.

Thresholding of the normalized intensity traces was used to determine the time which TW spent either bound to the intended (adjacent) *trpR* site or bound to the non-adjacent *trpR* site farthest from the bound foot. Due to the differing FRET efficiencies of the two fluorophores, independently selected absolute thresholds were used, specifically 0.8 for red FRET and 0.5 for yellow FRET. Frames in the first buffer condition for which only the red FRET intensity exceeded the specified threshold were counted as intended binding. Frames in the third buffer condition for which only the yellow FRET intensity exceeded the specified threshold were also counted as intended binding. Otherwise, frames were counted as non-adjacent binding including frames in which neither or both FRET intensities exceeded the threshold. To include an equal amount of data from each ligand solution, only the first three solution conditions were included in the analysis and only traces where the protein remained colocalized with the DNA for the full duration of this period were included. The fraction of time spent in the intended binding sites was calculated as $t_{intended}/(t_{intended} + t_{unintended})$ for each trace resulting in a mean of 65% ± 7.7% while the fraction of time spent in the unintended binding sites was calculated as $t_{unintended}/(t_{intended} + t_{unintended})$ for each trace resulting in a mean of 35% ± 6.3% where limits are 95% confidence intervals.

Supplementary Text 1

Single molecule FRET: Control Experiments

The effect of the fluid exchange in the absence of ligand changes was tested by performing the same biochemical preparation, imaging and mechanical flow switching as in the stepping experiments, but instead with Co^{2+}+SAM buffer in each of the three microfluidic reservoirs. Fig. S7 shows traces obtained using a single-molecule FRET analysis. In this experiment the TW is never expected to bind to a site labelled with a FRET acceptor, so the analyses for colocalization was relaxed by removing the requirement for colocalization with FRET intensity. Instead, relying only on colocalization with detections in the acceptor excitation channel. Keeping the FRET requirement removes all molecules from the analysis as expected. Otherwise, analysis was performed as in the above Methods section. Solutions were changed between identical Co^{2+}+SAM buffer 3 times and then changed to a buffer containing no ligands at all as indicated by Fig. S7A. Fig. S7B shows an example of these traces where TW dissociates from the DNA after the removal of ligands. All traces are shown as overlayed kymographs in Fig. S7C where green is TW donor signal and red is *trpR* site FRET. The kymograph shows no FRET nor any reaction to the changes in flow at all. At the point at which all ligands are removed (Fig. S7A) a slower than anticipated rate of signal loss is observed (Fig. S7B). This is due to the relaxed colocalization criteria allowing non-specifically bound nearby TW and DNA and potentially non-functional TW:DNA complexes to be included in the analysis. This is particular to this control experiment where we do not have a FRET signal as an analysis criterion.

Similar detachment experiments were performed with Trp+Co^{2+} (Fig. S8) and SAM+Trp, (Fig. S9), with a single solution change to the ligand-free solution. When changing to a ligand-free solution, the rate of TW detachment increases sharply for both of these ligand-pair conditions.

Supplementary Text 2

Simulation of TW Stepping

The single-molecule FRET data in Fig. 4 clearly demonstrate that TW can step bi-directionally on a short track consisting of four binding sites, but does not allow us to directly and experimentally evaluate performance parameters such as processivity and average speed on a long track. However, experimentally measured binding and stepping characteristics allow us to predict this performance using a model.

To analyze potential stepping behavior of TW along an extended track we devised a coarse-grained model based on Master equations governing the binding and unbinding kinetics of the three TW feet and the track. The track is represented by the periodic, unlimited sequence of binding sites …-*dtxR-metJ-trpR-dtxR-metJ-trpR-dtxR-metJ*-… (Fig. S19), mimicking the experimental setup used in Fig. 4. The kinetics of the three individual feet are described by three independent rate equations. The rate equation for each foot describes the transitions between an unbound state and bound states, in which the foot is bound specifically to its matching binding site on the track, e.g., the DtxR foot can only bind to *dtxR* etc., while non-specific binding is neglected, supported by the SPR results in Fig. 3A. Since these equations govern transitions between states (not concentrations), the corresponding rates have units [1/time] and are the inverse of the average waiting time between transitions. The values of these rates depend on the ligands in solution (see Table S8), which are cycled periodically through the combinations Trp + Co^{2+}, Co^{2+} + SAM, SAM + Trp, with a period of 21 seconds (i.e. the solution is switched every 7 seconds). Since there are always two ligands present at any given time, the rate equations for the individual feet become effectively coupled.

As an example, the set of possible states and the considered transitions between them are shown in Fig. S19 for the ligand combination Co^{2+} + SAM. In particular, Fig. S19 illustrates that we set up the rate model for each foot to distinguish between two distinct bound states, representing binding of a foot to its intended site (adjacent to the other bound foot) and unintended binding to the non-adjacent binding site (overstepping). For the other ligand combinations, Trp + Co^{2+} and SAM + Trp, analogous state combinations and transitions are realized by the rate equations.

The dynamics of each foot is thus driven by four rates for the transitions between its unbound state and two distinct bound states, which all depend on the foot-specific ligand being present in the solution or not. Fixing the explicit values for these rates in the model is guided by the experimental findings and is described below.

The rate processes for the three TW feet, their dependence on the ligand combination present in solution, and the 21-second cycle of the solution through three ligand combinations were implemented in a custom-built C++ code. A simulation run started with two feet bound on the track in the presence of the corresponding ligand combination, and ran until the TW detached from the track, *i.e.*, until none of the feet was in a bound state. The sequence of binding and unbinding events recorded during such a run for all three feet was translated into the corresponding stepping motion of the TW along the track. This resulted in a trajectory of the TW showing the position of its center (in units of track periods) as a function of time (in seconds). Examples of such trajectories are shown in Fig. S11.

Repeating simulations for individual trajectories many times gives access to the distribution of observables with which the expected stepping performance of the TW along an extended track can be characterized. We focus on the stepping distance as a function of time (giving the stepping velocity, Fig. S10A), the variance of the trajectories as a function of time (giving the effective diffusion coefficient, Fig. S10B), and the distance covered before the TW detaches from the track (Fig. S12).

The specific values for the various rates of the individual TW feet in our model were chosen as follows. We first focus on the rates describing (un-)binding between the TW repressor feet and the intended binding site (adjacent to the other bound foot): The off-rates for each repressor in the presence and absence of its ligand were adapted from the SPR experiments, see Table S8 (compare to Table S3), *i.e.*, the off-rates are characteristic for the type of repressor (TrpR, DtxR, MetJ). To choose the on-rates, we assumed that (re-)binding to the track is predominantly determined by the geometrical arrangement of the track and the binding sites, and thus can be assumed to be independent of the repressor type. For this reason, we took on-rates to be identical for all feet. Moreover, neglecting non-specific binding, the on-rates are all zero in the absence of the foot's ligand (see Table S8). In the presence of ligands, we adjusted the value of the on-rates such that the three *double-ligand* off-rates measured in the SPR dissociation experiments for the ligand combinations Trp + Co^{2+}, SAM + Co^{2+}, and SAM + Trp (Table S3) were best reproduced. To do this, we ran 10^6 simulations for each pair of feet and corresponding ligands, with the assumed value of the on-rates, until neither of the feet was bound . From the exponential distribution of the so obtained "survival times" we extract the effective dissociation rate; the results are listed in Table S8 (compare to Table S3 for the experimental values).

For unintended binding to the non-adjacent site (overstepping), we assumed the off-rates to be identical to the ones for unbinding from the intended binding site, because the off-rates are characteristic for the type of repressor, no matter where on the track it is bound. In the spirit of our assumption that the binding process is predominantly geometry-determined, the on-rates are again all the same for the different feet, but in general different from the on-rates to the intended binding site (different geometry). We encoded the difference in a factor α that multiplies the on-rates for intended binding:

$$\text{(on-rates for overstepping)} = \alpha * \text{(on-rates for intended binding).} \qquad 8$$

For $\alpha=0$, there is no overstepping, $0<\alpha<1$ implies that overstepping is less likely than intended binding, while $\alpha>1$ indicates that overstepping is more likely; for $\alpha=1$ intended binding and overstepping are equally likely. The relative weight of overstepping is calculated by comparing its rate to the total rate of binding, $\alpha/(1+\alpha)$. Hence, for $\alpha=0.5$ we find $1/3=33\%$, reproducing the experimentally observed frequency of overstepping of about 35%.

Supplementary Text 3

Can DNA bend sufficiently to facilitate overstepping?
    The inherent flexibility of the DNA track may be responsible for the observed overstepping of TW. To test this possibility, we used a simple physical model to determine the energetic cost of bending the DNA sufficiently to allow overstepping, *i.e.*, for two repressor feet to bind sites separated by an intervening empty site.

    If DNA is uniformly flexible, then it can follow a circular arc described by a radius of curvature, R_c, an arc length, s, and the angle subtended by the arc, θ, where

$$R_c = \frac{s}{\theta}. \qquad\qquad 5$$

    The largest possible reach of two TW feet occurs when the two legs (each of length d_l) are straight and each repressor foot (additional length d_f) binds to DNA perpendicularly (thus aligning the foot's axis with the local radius of curvature) (Fig. S20A,B; similar to Model 2 in Fig. 2C). For an angle θ subtended by the feet, this configuration of TW can be described by:

$$sin\left(\frac{\theta}{2}\right) = \frac{d_l}{R_c - d_f}. \qquad\qquad 6$$

    From our model of TW, we approximate: $d_l = 100\text{Å}$ and $d_f = 35\text{Å}$. From our model of the DNA track, the arc length between two non-consecutive repressor binding sites is approximately $s = 330\text{Å}$. Using these parameters, we can plot equations 5 and 6 for radius of curvature, R_c, as a function of the angle θ subtended – the equation 5 describing the DNA arc and equation 6 describing TW. The intersection of these two curves represents the solution of these two geometric equations (Fig. S20C), where $R_c \cong 145\text{Å}$ and $\theta \cong 130°$.

    By modelling DNA as a rigid rod of length s (*56*), we can estimate the energy, ΔU, required to bend DNA sufficiently to satisfy this geometric solution:

$$\Delta U = \frac{1}{2}\frac{BLs}{R_c^2}, \qquad\qquad 7$$

where B is the bending stiffness, $B = 50nm.\,k_BT$ (*56*). This gives $\Delta U = 3.9k_BT$, where k_B is the Boltzmann constant. Thus, it would not require a large thermal fluctuation to achieve the required bending of the DNA that would facilitate overstepping (along with flexibility in the TW linker regions). This bending could be reduced by supporting the DNA track in a straight configuration.

Fig. S1. P(r) plot calculated from the SEC-SAXS data. The figure shows the pair-distance distribution function, $P(r)$, derived from the experimental data (solid black line). Overlaid are the $P(r)$ functions for the representative compact (dotted blue line) and extended model structures (dotted red line) and their weighted sum representing the ensemble average (black dotted line).

Fig. S2. DNA binding by MetJ, DtxR and TrpR is orthogonal. MetJ, DtxR and TrpR were titrated onto each of their respective cognate sites via SPR in the presence of (**A**) SAM, (**B**) Co^{2+} and (**C**) Trp. The DNA sites used in these experiments are named above the figures, with sequences reported in Table S6. The steady state responses were fit to the Hill equation, with the fitted binding parameters for each specific interaction reported in Table S2.

Fig. S3. TW binding double-site DNA in the presence of single ligands, by SPR. TW was titrated onto DNA containing two repressor cognate sites **(A)** in the absence of ligand, and in the presence of **(B)** SAM, **(C)** Co²⁺ or **(D)** Trp. The DNA sites used in these experiments are named above the figures, with sequences reported in Table S6. **(E)** Representative steady state SPR binding isotherms of TW binding to double-site DNA were fitted to the Hill equation. The ligands and DNA used in these experiments are as per the labels within the figure panels. The fitted parameters for TW binding to the double site DNA are reported in Table S2.

Fig. S4. Dissociation kinetics of TW, by SPR. Representative dissociation experiments in which TW was loaded with the relevant ligand onto DNA containing either one or two repressor cognate sites. Dissociation was triggered by injection of a solution containing 1 µM free DNA containing the same cognate site as the DNA bound to the SPR chip surface. Free DNA sequences lacked an SPR anchor overhang and therefore could not bind to the SPR surface. The DNA sites used in these experiments are named above the figures, with sequences reported in Table S6. Ligands used in each experiment are labelled within the figure. Curves were fit to a single exponential decay function, with the apparent off-rates reported in Table S3. **(A)** Dissociation in the presence of DNA but absence of ligand. **(B)** Dissociation in the presence of both DNA and single ligands. **(C)** Dissociation in the presence of double ligands and two free oligonucleotides, each containing a single relevant repressor cognate site.

Fig. S5. TW forms monodisperse 1:1 complexes with both single- and double-site DNA, by mass photometry. TW:DNA mixtures were subjected to mass photometry. The resulting main mass distributions were fit to a Gaussian function, with molecular masses for each complex reported in Table S1. Signal interference by SAM and the binding of DNA to the glass slide in the presence of Co^{2+} resulted in additional mass peaks centered around 60–80 kDa. The DNA sites used in these experiments are named above the figures, with sequences reported in Table S6. The ligands used in each experiment are as per the labels within the figure panels. **(A)** TW binding to single-site DNA in the absence and presence of single ligands. The observed molecular masses of the TW:single-site complexes (150–154 kDa) were in good agreement with the theoretical molecular masses (155–161 kDa). **(B)** TW binding to double-site DNA in the absence and presence of double ligands. The observed molecular masses of the TW:double-site complexes (183–187 kDa) were in excellent agreement with the theoretical molecular masses (182–168 kDa).

Fig. S6. Normalized examples of single-molecule traces using two-colour imaging with emission split at 633 nm to simultaneously record ATTO565 emission (FRET emission from the bottom *trpR* site, shown in cyan) and ATTO647N emission (FRET emission from the top *trpR* site, shown in magenta). At the top of each column, a schematic of expected TW position and false colour representation of expected fluorescence signal given solution changes (dashed lines).

Fig. S7. **TW remained static on DNA track when ligand combination (Co²⁺+SAM) remained the same after microfluidic solution changes, as observed by single-molecule FRET.** **(A)** Schematic of expected TW position and false-color representation of expected fluorescence signal given solution changes (dashed lines). **(B)** Normalized example single-molecule trace. **(C)** Kymographs of all detected colocalizing single molecules sorted by time of last identified fluorescence. **(D)** Sum of all single-molecule traces in the experiment.

Fig. S8. **TW rapidly detached from DNA track after removal of ligands (SAM+Trp), as observed by single-molecule FRET.** **(A)** Schematic of expected TW position and false-color representation of expected fluorescence signal given solution changes (dashed lines). **(B)** Normalized example single-molecule trace. **(C)** Kymographs of all detected colocalizing single molecules sorted by time of last identified fluorescence. **(D)** Sum of all single-molecule traces in the experiment.

Fig. S9. TW rapidly detached from DNA track after removal of ligands (Trp+Co^{2+}), as observed by single-molecule FRET. (A) Schematic of expected TW position and false-color representation of expected fluorescence signal given solution changes (dashed lines). **(B)** Normalized example single-molecule trace. **(C)** Kymographs of all detected colocalizing single molecules sorted by time of last identified fluorescence. **(D)** Sum of all single-molecule traces in the experiment.

Fig. S10. Dynamical characteristics of the stepping motion along the track as a function of the parameter α which controls the relative frequency of overstepping. For α=0 there is no overstepping, α=0.5 corresponds to the situation observed in experiments where 65% of the time TW correctly steps while 35% oversteps (vertical dashed lines), for α=1 correct stepping and overstepping are equally likely, and for α>1 overstepping dominates. **(A)** Average stepping velocity obtained from 500,000 independent realizations of the stepping simulation. For the units of velocity, a track period consists of three binding sites and a ligand cycle is considered a full cycle of three ligand solution changes, so that a velocity of 1 corresponds to ideal stepping. **(B)** Effective diffusion coefficient of the same set of 500,000 trajectories.

'

Fig. S11. Trajectories of TW stepping along the DNA track from simulations. In each panel 10 independent simulation runs are shown in different colors. The vertical dashed lines indicate the time points at which the ligand combination in the solution is switched. **(A)** Trajectories with a probability of overstepping that reproduces the experimentally observed 65% correct stepping and 35% overstepping. **(B)** Trajectories without overstepping (only correct binding to the track is included in the model). Substeps visible in the traces correspond to binding/unbinding events within the current spatial period on the DNA track.

Fig. S12: The distances the simulated TW steps along the track before detaching are exponentially distributed. From a fit to the exponential curves, we obtain the average distance covered before detaching from the track (in units of [track periods]). **(A)** (left panel): α=0.5 (experimentally observed frequency of overstepping); 1/e stepping distance is 2.1 track periods ≈ 103 nm. **(B)** (right panel): α=0 (no overstepping); 1/e stepping distance is 1.6 track periods ≈ 78 nm.

```
>TW1
GKDLVDTTEM YLRTIYELEE EGVTPLRARI AERLEQSGPT VSQTVARMER  50   DtxR
DGLVVVASDR SLQMTPTGRT LATAVMRKHR LAERLLTDII GLDINKVHDE 100   Geminin_Coiled-Coil
ADRWEHVMSD EVERRLVKVL KGSSGGSGTL YEALKENEKL HKEIEQKDNE 150   SpyCatcher
IARLKKENKE LAEVAQGSSG GSGSSGDSAT HIKFSKRDED GKELAGATME 200   TrpR
LRDSSGKTIS TWISDGQVKD FYLYPGKYTF VETAAPDGYE VATAITFTVN 250   SpyTag
EQGQVTVNGK ATKGDAHI                                          MetJ

>TW2
GAQQSPYSAA MAEQRHQEWL RFVDLLKNAY QNDLHLPLLN LMLTPDEREA LGTRVRIVEE LLRGEMSQRE  70
LKNELGAGIA TITRGSNSLK AAPVELRQWL EEVLLKSDGS SGGSGTLYEA LKENEKLHKE IEQKDNEIAR 140
LKKENKELAE VAQGSSGGSG SSGAHIVMVD AYKPTKGSSG GSGSSGTLYE ALKENEKLHK EIEQKDNEIA 210
RLKKENKELA EVAQGSSGGS GGAEWSGEYI SPYAEHGKKS EQVKKITVSI PLKVLKILTD ERTRRKVNNL 280
RHATNSELLC EAFLHAFTGQ PLPDDADLRK ERSDEIPEAA KEIMREMGIN PETWEY
```

Fig. S13. Protein sequences of the TW constructs. Protein domains are colored per the top right inset. Glycine-serine rich linkers are colored grey. TW is formed via the formation of an isopeptide bond between the side chains of K186 in the SpyCatcher domain of TW1 and D170 in the SpyTag part of TW2. The S107C mutation was introduced into the TrpR domain of TW2 to enable labelling of the protein with AF488 via maleimide chemistry.

Fig. S14. Design of microfluidic device for single molecule TW stepping experiments. (A) Locations where the three inlets are punched are marked with arrows pointing inwards. Each inlet is to be connected to a reservoir containing the running buffer with a pair of ligands. Each inlet is connected via an equal length escape channel to a common outlet (circle, far left). Directly following the intersection of the three inlet channels, the main observation channel can be seen labelled with distance in mm from the intersection. The observation channel terminates in another outlet (circle, far right) placed 10 mm from the intersection. **(B)** Photo of assembled microfluidic device bonded to a coverslip including glued glass slide fragments for support and mounting. **(C)** 2D finite element simulation (incompressible, laminar flow) of flow rates in the microfluidic device for three fluids (red, green and blue) injected into the device inlets, one at 7 µL/min ("on" channel) the other two at 1 µL/min ("off" channel). The fluid injected at 7 µL/min (Top: red, Middle: green, Bottom: blue) propagates in the observation channel at 2.285 µL/min while the other two fluids are directed to their respective "escape" channel.

Fig. S15. Example of detected bright spots. A small field of view from a normalized maximum intensity projection of the data from **(A)** the FRET acceptor channel and **(B)** the donor channel from Experiment 1. All bright spots detected by the algorithm are outlined with white squares or circles. Squares denote those detected spots that were identified as not colocalizing in Donor and FRET channels. Circles denote those detected spots that were identified as colocalizing in Donor and FRET channels.

Fig. S16. Regions of interest created for single-molecule analysis. **(A)** shows the pixels used to determine intensity of a spot, and **(B)** shows the surrounding pixels whose intensity is used for background removal. The final background subtracted intensity of the molecular fluorescence signal in each frame is calculated by averaging the intensity of all pixels in the region of interest in **(A)** and subtracting the median in **(B)**.

Fig. S17. The resulting traces after background subtraction and normalization. Top: Experiment 1. Red is the FRET acceptor signal, green the donor. Black dashed lines indicate times of buffer changes. Bottom: Experiment 2. Magenta is the FRET acceptor signal from the red acceptor (top-*trpR* site) and cyan is the FRET acceptor signal from the yellow acceptor (bottom-*trpR* site). Black dashed lines indicate buffer switches.

Fig. S18. Outline of the edge detection algorithm. **(A)** shows the raw traces, **(B)** the raw traces after Gaussian filtering with standard deviation 1.2 and **(C)** shows the derivative of **(B)**. The threshold for peak detection is indicated by a yellow dashed line and the threshold for valley detection is indicated by a blue dashed line. In all panels, the detected rising edges are marked with a yellow dot and detected falling edges with a blue dot.

Fig. S19 Illustration of states and transitions modelled by the rate equations. Lower panel: The DNA track, extending ad infinitum, is sketched by a horizontal black line with colored dots, which label the different cognate binding sites on the track (compare Fig. 1B in the main text); two track periods are marked as n and $n+1$. The TW molecule is depicted in a star-like manner with three colored dots representing the repressor proteins TrpR (cyan), DtxR (magenta) and MetJ (brown), see Fig. 1 in the main text. Main panel: The five different possible states in the presence of the ligand combination Co^{2+} + SAM, including intended binding events and unintended overstepping events. The states are labelled by the repressor(s) bound to the track and the track period they are bound to, e.g., $(DtxR)_n(MetJ)_{n+1}$ indicates that DtxR is bound to track period n and MetJ to track period $n+1$ (this is the state in the middle); an asterisk at the repressor label indicates an overstepped configuration. The arrows indicate transitions between these different states. For the chosen ligand combination, there are no states in which TrpR is bound, or transitions towards such states, because we assume that TrpR can only bind in the presence of its ligand Trp. The sequence of events from the state $(DtxR)_n$ (upper left corner) to the state $(MetJ)_{n+1}$ (lower right corner) via the state $(DtxR)_n(MetJ)_{n+1}$ in the middle corresponds to the desired forward stepping behavior from one track period into the next. The transitions from the states $(MetJ^*)_n(DtxR)_n$ and $(MetJ)_{n+1}(DtxR^*)_{n+1}$ marked "TW oversteps" end up in the states $(MetJ)_n$ and $(DtxR)_{n+1}$ (not shown), which are equivalent to $(MetJ)_{n+1}$ and $(DtxR)_n$ but shifted by one track period backward and forward, respectively.

Fig. S20. Model for TW overstepping due to DNA flexibility. **(A)** The geometry of DNA bending into a uniform circular arc. TW is represented by two stretched legs, total length $2d_l$ that connect to DNA via two repressor feet, each of length, d_f, that bind to DNA in a perpendicular fashion, each tracing a radius of curvature. **(B)** An overlay of a model of fully stretched TW on the geometric figure from (**A**). (**C**) Plot of the equation for the radius of curvature describing the DNA geometry (blue) and the TW geometry (orange) as a function of angle. The intersection of these two curves marks a simultaneous solution to the two equations.

Protein	Ligand	DNA	Theoretical Mass (kDa)	Observed Mass (kDa)	n
TW1	-	-	59	56 ± 2	4
TW2	-	-	74	72 ± 3	4
TW	-	-	133	133 ± 1	4
TW	-	*metJ*	133	133 ± 3	3
TW	SAM	*metJ*	155	150 ± 1	3
TW	-	*dtxR*	133	133 ± 2	3
TW	Co^{2+}	*dtxR*	159	154 ± 1	3
TW	-	*trpR*	133	132 ± 1	3
TW	L-Trp	*trpR*	161	150 ± 2	4
TW	-	*trpR-dtxR*	133	129 ± 2	3
TW	$L\text{-Trp} + Co^{2+}$	*trpR-dtxR*	188	187 ± 2	4
TW	-	*dtxR-metJ*	133	130 ± 3	3
TW	$Co^{2+} + SAM$	*dtxR-metJ*	182	184 ± 2	5
TW	-	*metJ-trpR*	133	128 ± 2	3
TW	SAM + L-Trp	*metJ-trpR*	183	183 ± 3	4

Table S1. TW construct and complex molecular masses. Protein and protein:DNA molecular masses were determined using mass photometry, as shown in Fig. 3C & Fig. S5. Theoretical masses were calculated assuming TW and TW constructs act as obligate dimers. Observed masses represent average ± SEM, calculated from replicate experiments with sample size of n.

DNA	Protein	Ligand	Apparent K_d (nM)	h	n
metJ	MetJ	SAM	4 ± 0.4	1.22 ± 0.02	4
metJ	TW	SAM	18 ± 5	0.96 ± 0.07	4
dtxR	DtxR	Co^{2+}	5.5 ± 0.2	0.99 ± 0.03	5
dtxR	TW	Co^{2+}	8 ± 1	0.92 ± 0.10	4
trpR	TrpR	L-Trp	48 ± 2	1.61 ± 0.02	4
trpR	TW	L-Trp	27 ± 4	1.07 ± 0.03	4
trpR-dtxR	TW	Co^{2+}	3 ± 2	1.0 ± 0.2	2
trpR-dtxR	TW	L-Trp	15 ± 4	1.27 ± 0.09	4
dtxR-metJ	TW	SAM	14 ± 2	1.2 ± 0.2	3
dtxR-metJ	TW	Co^{2+}	5 ± 2	1.2 ± 0.1	3
dtxR-metJ	TW	L-Trp	50 ± 20	1.72 ± 0.03	3
metJ-trpR	TW	SAM	18 ± 2	1.3 ± 0.2	3
metJ-trpR	TW	L-Trp	13 ± 4	1.22 ± 0.09	4

Table S2. TW:DNA affinities in the presence of a single ligand by SPR. Affinities were determined by titrating TW onto a surface containing DNA. The average steady state response units in the contact phase were fitted to the Hill equation to determine the apparent K_d and Hill coefficient (h). Values represent average $\pm$ SEM, calculated from replicate experiments with sample size of n.

DNA	Ligand	Apparent k_{off} (s^{-1})	Apparent Half-life (s)	n
metJ	No ligand	3.7 ± 0.3	0.19 ± 0.02	3
metJ	SAM	1.14 ± 0.06	0.61 ± 0.03	3
dtxR	No ligand	0.45 ± 0.02	1.56 ± 0.07	3
dtxR	Co^{2+}	0.164 ± 0.009	4.2 ± 0.4	3
trpR	No ligand	0.99 ± 0.02	0.70 ± 0.02	3
trpR	L-Trp	0.36 ± 0.04	2.0 ± 0.2	3
trpR-dtxR	L-Trp + Co^{2+}	0.0239 ± 0.0008	29 ± 0.6	3
dtxR-metJ	SAM + Co^{2+}	0.0145 ± 0.0005	48 ± 1	2
metJ-trpR	SAM + L-Trp	0.0233 ± 0.002	29.7 ± 0.2	3

Table S3. TW:DNA dissociation kinetics by SPR. TW was loaded onto a DNA surface in the presence of the relevant ligands. Dissociation was triggered by injecting a solution containing no TW, the relevant ligands, and 1 µM DNA. 1 µM of each of the relevant single site DNA oligonucleotides were used for double ligand dissociation experiments. Dissociation curves were fitted to a single exponential decay function to determine apparent k_{off}. Half-lives were calculated from the apparent k_{off}'s. Values represent average ± SEM, calculated from replicate experiments with sample size of n.

Dye	A_{max} (nm)	Extinction Coefficient ($M^{-1}cm^{-1}$)	Correction Factor	Molecular Weight (g/mol)
Alexa Fluor™ 488 C_5 Maleimide	495	73000	0.11	720.66
Alexa Fluor™ 647 C_2 Maleimide	650	265000	0.03	1250
Atto 647 Maleimide	647	120000	0.04	829

Table S4. Properties of dye conjugates used for protein labelling. Data from Thermofisher Scientific and Sigma Aldrich.

(a) Sample details	
Organism	Synthetic construct
Source	Recombinant expression of components TW1 and TW2 in E. coli.
Scattering protein composition	TW protein composed of constructs TW1 and TW2 linked covalently via SpyTag/SpyCatcher
Sample environment/configuration	
Solvent composition	10 mM HEPES pH 7.4, 150 mM NaCl, 5mM $MgCl_2$
Sample temperature (°C)	Sample stored at ~10°C in a 96-well plate, but measured at ~20°C
In-beam sample cell	Sheath flow configuration
SEC-SAXS measurements	
Initial sample concentration (mg/mL)	~10
Column type	Superdex 200 Increase 5/150 GL
Volume injected (μL)	50
Flow rate (mL/min)	0.2

(b) SAS data collection	
Data acquisition / reduction software	SAXS/WAXS beamline control software, *ScatterBrain* (v 2.30), *CHROMIX* (*38*)
Source/instrument description	SAXS/WAXS beamline at the Australian Synchrotron, with Pilatus 2M detector at a SDD of 2680mm, with an incident intensity of ~8 x 10^{12} ph/s
Measured q-range (q_{min}–q_{max}) (Å^{-1})	0.005 – 0.5
Method for scaling intensities	Absolute scaling (cm^{-1}) referenced to water
Exposure time(s), No. of exposures	1 s x 21 frames for buffer/background 1 x 10 frames for sample
Additional relevant details	Buffer scattering frames taken immediately before the void fraction. Sample scattering frames taken at the elution peak over a region of stable R_g

(c) SAS-derived structural parameters	*PRIMUS/qt*, *AUTORG* and *GNOM* (*38*)
Guinier analysis	
$I(0) \pm \sigma$ (cm^{-1})	0.06175 ± 0.00022
$R_g \pm \sigma$ (Å)	67.7 ± 0.6
qR_g range (datapoint range)	0.39 – 1.00 (points 4-30)
Linear fit assessment (*AUTORG* fidelity)	0.47
PDDF/$P(r)$ analysis	
$I(0) \pm \sigma$ (cm^{-1})	0.06177 ± 0.00015
$R_g \pm \sigma$ (Å)	69.1 ± 0.4
d_{max} (Å)	270
q-range (Å^{-1})	0.006 – 0.492

$P(r)$ reciprocal-space fit: χ^2, CorMap P-value	1.63, 0.106

(*d*) Scattering particle size.	
Method(s)/software	*MoW* from *PRIMUS/qt* (*38*)
Volume estimates ($\mathring{A}^3$)	
Porod volume V_P	190
Molecular mass M estimates (kDa)	
Molecular mass M estimates	156
From chemical composition	133

(*e*) Modelling.	
Methods/software	
Atomistic modelling/software	*Multi-FoXS* (*16*)
q-range for fit ($\mathring{A}^{-1}$)	0.006 – 0.492
Symmetry/anisotropy assumptions	*N/A*
No. of individual model reconstructions	2 structures used to represent the scattering data taken from a pool of 100 conformations
χ^2, CorMap P-value	1.56, 0.156

(*f*) Data and model deposition.	
SASBDB ID	SASDWM8

Table S5. SEC-SAXS data collection, reduction and analysis.

Oligo name	Oligo type	Oligo sequence
NR019_Biotin-Anchor	Anchor	ACCTTAGGGCGTGG-Biotin
NR139_metJ_top	Single site	CCACGCCCTAAGGTATGCTAGTTT**GAGACGTCTC**AAAACAGGA
NR140_metJ_bottom	Single site	TCCTGTTTT**GAGACGTCTC**AAACTAGCAT
NR141_dtxR_top	Single site	CCACGCCCTAAGGTCATGCAAGGC**TTAGGTTAACCTAA**CTTGCATGTT
NR142_dtxR_bottom	Single site	AACATGCAAG**TTAGGTTAACCTAA**GCCTTGCATG
NR206_trpR_top	Single site	CCACGCCCTAAGGTGAGCATAGTG**GTACTCGCTAGCGAGTAC**CTCTGACTTC
NR207_trpR_bottom	Single site	GAAGTCAGAG**GTACTCGCTAGCGAGTAC**CACTATGCTC
NR349_metJ_top	Competitive DNA	GTTT**GAGACGTCTC**AAAA
NR350_metJ_bottom	Competitive DNA	TTTT**GAGACGTCTC**AAAC
NR351_dtxR_top	Competitive DNA	AGGC**TTAGGTTAACCTAA**CTTG
NR352_dtxR_bottom	Competitive DNA	CAAG**TTAGGTTAACCTAA**GCCT
NR353_trpR_top	Competitive DNA	AGTG**GTACTCGCTAGCGAGTAC**CTCT
NR354_trpR_bottom	Competitive DNA	AGAG**GTACTCGCTAGCGAGTAC**CACT
NR420_trpR-dtxR_top	Double site	CCACGCCCTAAGGTGAGCATAGTG**GTACTCGCTAGCGAGTAC**CTCTGACTTCCATGACGGGTGTCACATCATGCAAGGC**TTAGGTTAACCTAA**CTTGCATGTT
NR421_trpR-dtxR_bottom	Double site	AACATGCAAG**TTAGGTTAACCTAA**GCCTTGCATGATGTGACACCCGTCATGGAAGTCAGAG**GTACTCGCTAGCGAGTAC**CACTATGCTC
NR423_dtxR-metJ_top	Double site	CCACGCCCTAAGGTCATGCAAGGC**TTAGGTTAACCTAA**CTTGCATGTTGATCATATAGCCAATATGCTAGTTT**GAGACGTCTC**AAAACAGGA
NR424_dtxR-metJ_bottom	Double site	TCCTGTTTT**GAGACGTCTC**AAACTAGCATATTGGCTATATGATCAACATGCAAG**TTAGGTTAACCTAA**GCCTTGCATG
NR426_metJ-trpR_top	Double site	CCACGCCCTAAGGTATGCTAGTTT**GAGACGTCTC**AAAACAGGAATGGCACGCGTCCTGAGCATAGTG**GTACTCGCTAGCGAGTAC**CTCTGACTTC
NR427_metJ-trpR_bottom	Double site	GAAGTCAGAG**GTACTCGCTAGCGAGTAC**CACTATGCTCAGGACGCGTGCCATTCCTGTTTT**GAGACGTCTC**AAACTAGCAT

Table S6. Oligonucleotide sequences used for SPR and mass photometry. Underlined
sequences represent complements to Biotin-Anchor strand used to anneal DNA surfaces in SPR

experiments. Single- and double-site oligos refer to sequences that contain one and two repressor cognate sites, respectively. Competing DNA refers to sequences containing a single cognate binding site that was used to trigger dissociation in SPR experiments. Bold sequences represent the cognate binding sequences for the relevant repressor. Top and bottom sequences were annealed to form double-stranded DNA for use in experiments.

Oligo name	Description	Sequence
Up-strand 1	Anchor + bottom-*trpR* site + *dtxR* site	CCACGCCCTAAGGTGAGCATAGTG**GTACTCGCTAGCGAGTAC**CTCTGACT TCCATACGGGTGTCACATCATGCAAGGC**TTAGGTTAACCTAA**C
Up-strand 2	*metJ* site + top-*trpR* site	Phosphate-TTGCATGTTGATCATATAGCCAATATGCTAGTTT**GAGACGTCTC**AAAACA GGAATGGCACGCGTCCTGAGCATAGTG**GTACTCGCTAGCGAGTAC**CTCT
Down-strand 1	Atto647N + top-*trpR* site 2	ATTO647N–AGAG**GTACTCGCTAGCGAGTAC**CACTATGCTCAGGACGCGTGCCATT
Down-strand 2	*metJ* site + *dtxR* site	Phosphate-CCTGTTTT**GAGACGTCTC**AAACTAGCATATTGGCTATATGATCAACATGC AAG**TTAGGTTAACCTAA**GCCTTGCATG
Down-strand 3	bottom-*trpR* site + Atto565	Phosphate-ATGTGACACCCGTATGGAAGTCAGAG**GTACTCGCTAGCGAGTAC**CAC–ATTO565
Down-strand 4	Anchor	TATGCTCACCTTAGGGCGTGG-Biotin

Table S7. Oligonucleotide sequences used to construct the track for single molecule FRET. The four-site track was annealed, ligated, and purified as described in Materials and Methods from the six oligonucleotides listed here. Up-strands refer to strands designed with 5' to 3' direction away from the coverslip. Down-strands refer to strands designed with 5' to 3' direction towards the coverslip. Bold sequences represent the cognate binding sequences for the relevant repressor.

DNA	Ligand	Off rate (s^{-1})	On rate (s^{-1})
metJ	SAM	1.14	10
	none	3.7	0
dtxR	Co^{2+}	0.164	10
	none	0.45	0
trpR	Trp	0.36	10
	none	0.99	0
trpR-dtxR	Trp + Co^{2+}	0.011	n/a
dtxR-metJ	SAM + Co^{2+}	0.031	n/a
metJ-trpR	SAM + Trp	0.067	n/a

Table S8: Single- and no-ligand rates used in the simulations for the transition between the unbound and the intended bound state. Single- and no-ligand off-rates are identical to the rates measured in the SPR dissociation experiments (Table S3). The on-rates are assumed to be repressor-independent and are chosen to reproduce the measured double-ligand rates as well as possible; the double-ligand rates resulting from the model are all within a factor 3 from the measured rates, compare Table S3. They are extracted from the model by simulating the detachment of TW from the track in the presence of fixed ligand combinations Trp + Co^{2+}, Co^{2+} + SAM, and SAM + Trp, respectively. Transitions between unbound state and overstepped bound state use the same off-rates, but all on-rates are rescaled by a factor α.

1. Z. Wang, S. Raunser, Structural Biochemistry of Muscle Contraction. *Annu Rev Biochem* **92**, 411-433 (2023).

2. J. Robert-Paganin, O. Pylypenko, C. Kikuti, H. L. Sweeney, A. Houdusse, Force Generation by Myosin Motors: A Structural Perspective. *Chem Rev* **120**, 5-35 (2020).

3. A. Yildiz, Mechanism and regulation of kinesin motors. *Nature reviews. Molecular cell biology* **26**, 86-103 (2025).

4. S. L. Reck-Peterson, W. B. Redwine, R. D. Vale, A. P. Carter, The cytoplasmic dynein transport machinery and its many cargoes. *Nature reviews. Molecular cell biology* **19**, 382-398 (2018).

5. C. Bustamante, Y. R. Chemla, N. R. Forde, D. Izhaky, Mechanical processes in biochemistry. *Annu Rev Biochem* **73**, 705-748 (2004).

6. V. Ramakrishnan, The ribosome emerges from a black box. *Cell* **159**, 979-984 (2014).

7. N. V. DelRosso, N. D. Derr, Exploiting molecular motors as nanomachines: the mechanisms of de novo and re-engineered cytoskeletal motors. *Current opinion in biotechnology* **46**, 20-26 (2017).

8. C. S. Korosec *et al.*, Motility of an autonomous protein-based artificial motor that operates via a burnt-bridge principle. *Nat Commun* **15**, 1511 (2024).

9. M. Baroncini *et al.*, Making and Operating Molecular Machines: A Multidisciplinary Challenge. *ChemistryOpen* **7**, 169-179 (2018).

10. T. Zhang, H. Liu, The motive forces in DNA-enabled nanomachinery. *iScience* **27**, 109453 (2024).

11. P. F. Nilsson *et al.*, Walking by design: how to build artificial molecular motors made of proteins. *Nano Futures* **8**, 042501 (2024).

12. H. Linke, B. Höcker, K. Furuta, N. R. Forde, P. M. G. Curmi, Synthetic biology approaches to dissecting linear motor protein function: towards the design and synthesis of artificial autonomous protein walkers. *Biophys Rev* **12**, 1041-1054 (2020).

13. E. H. C. Bromley *et al.*, The Tumbleweed: towards a synthetic protein motor. *Hfsp Journal* **3**, 204-212 (2009).

14. B. Zakeri *et al.*, Peptide tag forming a rapid covalent bond to a protein, through engineering a bacterial adhesin. *Proceedings of the National Academy of Sciences of the United States of America* **109**, E690-697 (2012).

15. R. D. Astumian, Thermodynamics and kinetics of a Brownian motor. *Science* **276**, 917-922 (1997).

16. D. Schneidman-Duhovny, M. Hammel, J. A. Tainer, A. Sali, FoXS, FoXSDock and MultiFoXS: Single-state and multi-state structural modeling of proteins and their complexes based on SAXS profiles. *Nucleic Acids Res* **44**, W424-429 (2016).

17. J. Jumper *et al.*, Highly accurate protein structure prediction with AlphaFold. *Nature* **596**, 583-589 (2021).

18. S. Kovacic *et al.*, Design and Construction of a One-Dimensional DNA Track for an Artificial Molecular Motor. *J Nanomater* **2012**, 109238 (2012).

19. N. J. Kuwada *et al.*, Tuning the performance of an artificial protein motor. *Phys Rev E Stat Nonlin Soft Matter Phys* **84**, 031922 (2011).

20. N. J. Kuwada, G. A. Blab, H. Linke, A classical Master equation approach to modeling an artificial protein motor. *Chem Phys* **375**, 479-485 (2010).

21. C. S. Niman *et al.*, Controlled microfluidic switching in arbitrary time-sequences with low drag. *Lab Chip* **13**, 2389-2396 (2013).

22. S. Sudhakar *et al.*, Germanium nanospheres for ultraresolution picotensiometry of kinesin motors. *Science* **371**, (2021).

23. T. Yanagida, M. Iwaki, Y. Ishii, Single molecule measurements and molecular motors. *Philos Trans R Soc Lond B Biol Sci* **363**, 2123-2134 (2008).

24. S. L. Reck-Peterson *et al.*, Single-molecule analysis of dynein processivity and stepping behavior. *Cell* **126**, 335-348 (2006).

25. F. Mathieu *et al.*, Six-helix bundles designed from DNA. *Nano Lett* **5**, 661-665 (2005).

26. A. Spang *et al.*, Complex archaea that bridge the gap between prokaryotes and eukaryotes. *Nature* **521**, 173-179 (2015).

27. J. Vosseberg *et al.*, The emerging view on the origin and early evolution of eukaryotic cells. *Nature* **633**, 295-305 (2024).

28. F. J. Kull, R. D. Vale, R. J. Fletterick, The case for a common ancestor: kinesin and myosin motor proteins and G proteins. *Journal of muscle research and cell motility* **19**, 877-886 (1998).

29. L. Li, J. O. Fierer, T. A. Rapoport, M. Howarth, Structural analysis and optimization of the covalent association between SpyCatcher and a peptide Tag. *J Mol Biol* **426**, 309-317 (2014).

30. C. S. Chen, A. White, J. Love, J. R. Murphy, D. Ringe, Methyl groups of thymine bases are important for nucleic acid recognition by DtxR. *Biochemistry* **39**, 10397-10407 (2000).

31. M. Thepaut *et al.*, Crystal structure of the coiled-coil dimerization motif of geminin: structural and functional insights on DNA replication regulation. *J Mol Biol* **342**, 275-287 (2004).

32. C. W. Garvie, S. E. Phillips, Direct and indirect readout in mutant Met repressor-operator complexes. *Structure* **8**, 905-914 (2000).

33. Y. Y. He *et al.*, Structural and functional studies of an intermediate on the pathway to operator binding by Escherichia coli MetJ. *J Mol Biol* **320**, 39-53 (2002).

34. E. Tully, R. O'Kennedy, in *Encyclopedia of Microfluidics and Nanofluidics,* D. Li, Ed. (Springer US, Boston, MA, 2008), pp. 737-749.

35. N. M. Kirby *et al.*, A low-background-intensity focusing small-angle X-ray scattering undulator beamline. *Journal of Applied Crystallography* **46**, 1670-1680 (2013).

36. T. M. Ryan *et al.*, An optimized SEC-SAXS system enabling high X-ray dose for rapid SAXS assessment with correlated UV measurements for biomolecular structure analysis. *Journal of Applied Crystallography* **51**, 97-111 (2018).

37. K. Manalastas-Cantos *et al.*, ATSAS 3.0: expanded functionality and new tools for small-angle scattering data analysis. *J Appl Crystallogr* **54**, 343-355 (2021).

38. M. V. Petoukhov *et al.*, New developments in the ATSAS program package for small-angle scattering data analysis. *J Appl Crystallogr* **45**, 342-350 (2012).

39. M. Mirdita *et al.*, ColabFold: making protein folding accessible to all. *Nat Methods* **19**, 679-682 (2022).

40. E. F. Pettersen *et al.*, UCSF Chimera--a visualization system for exploratory research and analysis. *J Comput Chem* **25**, 1605-1612 (2004).

41. M. Jeeves, P. D. Evans, R. A. Parslow, M. Jaseja, E. I. Hyde, Studies of the Escherichia coli Trp repressor binding to its five operators and to variant operator sequences. *Eur J Biochem* **265**, 919-928 (1999).

42. R. Sharma, A. S. Patelli, L. De Bruin, J. H. Maddocks, cgNA+web : A Visual Interface to the cgNA+ Sequence-dependent Statistical Mechanics Model of Double-stranded Nucleic Acids. *J Mol Biol* **435**, 167978 (2023).

43.   L. De Bruin, J. H. Maddocks, cgDNAweb: a web interface to the cgDNA sequence-dependent coarse-grain model of double-stranded DNA. *Nucleic Acids Res* **46**, W5-w10 (2018).

44.   W. L. DeLano, *The PyMOL User's Manual*. (DeLano Scientific, San Carlos, CA, USA, 2002).

45.   N. Rezaei, A. Lyons, N. R. Forde, Environmentally Controlled Curvature of Single Collagen Proteins. *Biophysical journal* **115**, 1457-1469 (2018).

46.   S. Hertel *et al.*, The stability and number of nucleating interactions determine DNA hybridization rates in the absence of secondary structure. *Nucleic Acids Res* **50**, 7829-7841 (2022).

47.   P. Schuck, H. Zhao, The role of mass transport limitation and surface heterogeneity in the biophysical characterization of macromolecular binding processes by SPR biosensing. *Methods Mol Biol* **627**, 15-54 (2010).

48.   S. Wang, G. M. Poon, W. D. Wilson, Quantitative Investigation of Protein-Nucleic Acid Interactions by Biosensor Surface Plasmon Resonance. *Methods Mol Biol* **1334**, 313-332 (2015).

49.   P. G. Stockley, B. Persson, Surface plasmon resonance assays of DNA-protein interactions. *Methods Mol Biol* **543**, 653-669 (2009).

50.   T. D. Pollard, A guide to simple and informative binding assays. *Mol Biol Cell* **21**, 4061-4067 (2010).

51.   E. C. Hulme, M. A. Trevethick, Ligand binding assays at equilibrium: validation and interpretation. *Br J Pharmacol* **161**, 1219-1237 (2010).

52.   D. Wu, G. Piszczek, Standard protocol for mass photometry experiments. *Eur Biophys J* **50**, 403-409 (2021).

53.   S. van der Walt *et al.*, scikit-image: image processing in Python. *PeerJ* **2**, e453 (2014).

54.   T. Lindeberg, Scale Selection Properties of Generalized Scale-Space Interest Point Detectors. *Journal of Mathematical Imaging and Vision* **46**, 177-210 (2013).

55.   P. Virtanen *et al.*, SciPy 1.0: fundamental algorithms for scientific computing in Python. *Nat Methods* **17**, 261-272 (2020).

56.   J. F. Marko, S. Cocco, The micromechanics of DNA. *Physics World* **16**, 37 (2003).