1

The distinct role of human PIT in attention control

Siyuan Huang[1,2], Lan Wang[1], Sheng He[1,2].

1State Key Laboratory of Brain and Cognitive Science, Institute of Biophysics, Chinese Academy of
Sciences, Beijing 100101, China

2University of Chinese Academy of Sciences, Beijing 100049, China

* Sheng He.

Email: hes@ibp.ac.cn

Author Contributions: S. Huang, L. Wang, and S. He designed research; S. Huang performed research;
S. Huang analyzed data; and S. Huang, L. Wang, and S. He wrote the paper.

Competing Interest Statement: The authors declare no competing interest.

Classification: Psychological and Cognitive Sciences

Keywords: attention; cognition; fMRI; PITd; priority map

This Microsoft Word file includes:

16   Main Text
17   Figures 1 to 7

Abstract

Attentional distribution depends on both endogenous and exogenous processes, but how they interact in

attention allocation remains unclear. The attentional priority map, jointly determined by stimulus saliency

and task relevance, provides a framework for investigating their interplay. We propose that the human

posterior inferotemporal cortex (hPIT), located near object-processing cortical areas, serves as an

attentional priority map. Using fMRI with behavioral tasks, we show that hPIT shows stronger attentional

modulation than classical attention regions across motion, color, and shape tasks. hPIT shows lateralized

attentional enhancement even in the absence of visual input, with further elevated modulation in the

presence of stimuli, indicating its integrated role in priority control. Furthermore, its modulation is invariant

to stimulus category but sensitive to attentional demands, and the region is functionally connected to both

dorsal and ventral attentional networks. These findings highlight the hPIT as an integrator in attentional

control and provide critical insights into the brain's strategy for optimizing responses to the environment.

Main Text

Introduction

Attention serves as a pivotal mechanism within the brain, facilitating the allocation of its finite resource

towards the most pertinent stimuli in our environment (Desimone and Duncan 1995). Attention can be

systematically categorized according to its origin: bottom-up attention (exogenous), triggered by highly

salient stimuli, and top-down attention (endogenous), initiated through voluntary selection (Connor et al.

2004; Buschman and Miller 2007; Katsuki and Constantinidis 2014). The neural networks for bottom-up

and top-down attention have been extensively investigated, with ventral and dorsal attention networks

(VAN and DAN) specialized to support exogenous and endogenous attention respectively (Corbetta and

Shulman 2002; Corbetta et al. 2008; Vossel et al. 2014). How these two networks flexibly interact to

achieve integrated attentional selection, under situations that both salient stimulus and intentional goal

are present, remains inadequately understood (Noudoost et al. 2010).

Critical to the integration of exogenous and endogenous attention is the concept of "priority map",

defined as a map that reflects both the low-level salience and top-down influences (Treue 2003; Bisley

and Goldberg 2010; Bisley and Mirpour 2019). An important question arises concerning the location of

the priority map in the brain. For a brain region to qualify as the neural substrate supporting a priority

map, it must satisfy the following criteria (Ptak 2012; Stemmann and Freiwald 2019): 1) having spatially

restricted receptive fields, 2) exhibiting robust attentional modulation to predict attentional focus

regardless of stimulus attributes but sensitive to attentional load, 3) displaying little feature specificity and

high visual responsivity, and 4) exhibiting integration of bottom-up and top-down inputs. While the

saliency map - where objects compete for greater representation and attentional allocation (Fecteau and

Munoz 2006) across all feature dimensions - plays an important role in the mechanism of bottom-up

attention, the priority map (jointly determined by saliency and task guidance) is crucial for the cooperation

of bottom-up and top-down attention, making it a key aspect to understand for the implementation of

attention.

Previous research has indicated that the prefrontal cortex (PFC) and posterior parietal cortex (PPC) are

involved in both top-down and bottom-up attention, rendering them potential candidates for supporting a

priority map guiding attentional selection (Herrington and Assad 2009; Arcizet et al. 2011; Katsuki and

Constantinidis 2012; Ninomiya et al. 2012; Katsuki and Constantinidis 2014). In addition, the superior

colliculus (SC), especially its intermediate layers, receives signals from the frontal eye fields (FEF) and

lateral intraparietal area (LIP) / intraparietal sulcus (IPS) (Rushworth et al. 2006; Schall et al. 2011;

Benarroch 2023), could also support a priority representation (White et al. 2017). While visual areas,

driven by stimulus features, were considered unlikely to exhibit priority representation (Bisley and Mirpour

2019), a recent study in monkeys revealed that the posterior inferotemporal cortex (PITd) could encode

the locus of attention independent of features (Stemmann and Freiwald 2019). The structural connectivity

between PITd and parieto-frontal attentional areas (specifically LIP and FEF) further supports the idea

that PITd is part of attentional networks (Sani et al. 2019). In the context of human brains, recent results

have identified an area in the posterior inferotemporal cortex (hPIT) as a retinotopic and functional

homologue of the macaque PITd, which was proposed to be a putative node for the human endogenous

attentional control network (Sani et al. 2021). However, since only endogenous attention was investigated

in the previous experiments on monkey PITd as well as human PIT, it remains unclear whether hPIT is

important for the integration of endogenous and exogenous attention.

The current study investigated if attention can modulate activation in hPIT independent of stimulus

properties and cognitive demands. Instead of relying solely on the attentive motion task (Sani et al. 2021),

hPIT was localized and validated using three distinct spatial attention tasks. Furthermore, we aim to

discern the role of hPIT in both top-down and bottom-up attention, by manipulating the presence or

absence of visual stimuli. Distinct from nodes of classical endogenous attention network, hPIT was more

strongly modulated by attention in the presence of visual input. Furthermore, attention load, rather than

object category, significantly impacts the modulation in hPIT. Additionally, hPIT showed strong functional

connection with nodes in both ventral and dorsal attention network. Our results strongly suggest that

hPIT, different from its adjacent object-processing areas and other parietal and frontal nodes of attention

network, functions as an attention priority map.

Materials and Methods

1. Participants

Fifteen health volunteers (8 males and 7 females, age ranged from 22 to 28 years old) with normal or

corrected to normal vision participated in this study. None of the participants reported history of

neurological or psychiatric symptoms. Each participant completed three separate experimental sessions,

conducted on three different days. Written informed consent was obtained from all participants, and the

study protocol was approved by the Institutional Review Board of the Institute of Biophysics, Chinese

Academy of Sciences (#2020-IRBH-001).

2 Stimuli and procedures

Visual stimuli for the fMRI experiment were programmed using Matlab (MathWorks, Natick, MA, USA)

with Psychtoolbox (http://psychtoolbox.org/) and displayed via an MRI-compatible projector (1024 ×

768@60 Hz) onto a screen positioned at the rear of the MRI scanner. Participants viewed the stimuli

through a mirror attached to the head coil.

Because of the complexity of the task, each participant was trained on 2 or 3 separate days, before the

fMRI sessions. This training was designed to familiarize participants with the tasks to relieve some

possible tension during scanning and minimize the effects of learning.

2.1 Stimuli and procedures for experiment 1

In Experiment 1 (Day 1), participants completed three distinct spatial attention tasks across three

separate experimental blocks (Experiments 1a, 1b, 1c). Throughout each task, participants were

instructed to maintain fixation on a central point.

Experiment 1a (Fig.1A): Participants focused on a unilateral moving dot display and was required to

discriminate the direction of the coherent motion. As shown on Fig.1A, each trial began with an initial cue

indicating the target side, represented by a bar (0.6° visual angle) attached to the fixation point. Random

dot stimuli were then presented within a circular aperture (radius: 3° visual angle) centered 9° from the

fixation point. The dot display consisted of dots (size: 0.15° visual angle; density: 5 dots per degree of

visual angle) in motion for 2.5 seconds. During this sequence, a 0.5-second interval of coherent motion

(coherence: 50%, velocity: 6°/s) occurred randomly between 0.5 and 1.55 seconds after motion onset,

while the remaining duration involved random dot movement. Following the motion sequence, the circular

aperture disappeared, and an arrow appeared at the center of the screen for 1.5 seconds. Participants

were required to press a key to indicate whether the direction of coherent motion matched the direction

indicated by the arrow. Each block consisted of three trials with the same attended side, followed by a 6-

second rest period.

Experiment 1b (Fig.1B): Participants were instructed to attend to the proportion of red and green dots

presented within a unilateral circular aperture. The red and green dots were adjusted individually by each

participant to achieve iso-luminance, ensuring perceptual equality between the colors. The positioning of

the circular aperture was consistent with Experiment 1a, though the dot size was increased to 0.3° of

visual angle in this task. At the start of each trial, within the first second, dots were displayed in random

proportions of red (20%, 40%, 60%, 80%) with the remainder being green (pattern 1). Following this, the

position of the dots remained static during the 2nd second, but their color could potentially change

(pattern2). The proportion of red dots remain to be one of the pre-set ratios. In the final, third second of

each trial, the bilateral dots disappeared. Participants were then required to press a key to indicate

whether the color ratio of the dots had changed from pattern 1 to pattern 2. Each block in this experiment

consisted of four trials, with participants attending to the same side throughout, followed by a 6-second

rest period.

Experiment 1c (Fig.1C): In this experiment, participants were instructed to attend to a unilateral

geometrical pattern display, focusing on the proportion of different shapes. Each trial began with the first

pattern presented for 2 seconds, followed by a gradual transition in which the second pattern emerged

over the next 2 seconds and then remained visible. At the end of this sequence, participants were

required to press a key indicating whether the shape proportion between the two patterns had changed.

Each block consisted of two trials with attention directed to the same side, followed by a 6-second rest

period.

There were 16 blocks in one run, 8 attending left and 8 attending right for experiment 1a, 1b and 1c. On

the first day of scanning, participants completed 6 runs of Experiment 1a, 4 runs of Experiment 1b, and 4

runs of Experiment 1c. The average accuracy was 95% for Experiment 1a, 90% for Experiment 1b, and

90% for Experiment 1c.

2.2 Stimuli and procedures for experiment 2

In experiment 2 (day 2), participants first underwent a 488-second resting-state functional scan, during

which they passively viewed a grey screen. For the subsequent attention tasks (event-related design),

participants were instructed to maintain fixation on a central dot while directing their attention to one side.

The procedure (Fig.2) began with the central dot (10 pixels in diameter) flashing three times over 2

seconds, serving as an alert for the upcoming cue and target. Following this, a cue was presented for

0.8s. After a variable delay, randomly selected from intervals of 0.1, 0.15, 0.2, or 0.25 seconds, a target

dot (8 pixels in diameter, RGB: [0.7,0.7,0.7]) either appeared on the cued side (dot condition) or did not

appear (blank condition). In the dot condition, participants were required to report the location of the

target by pressing a key. In the blank condition, participants were instructed to refrain from any response.

Each run consisted of 16 trials, with individual trial durations of 14, 16, or 18 seconds. On this second

day of scanning, participants completed one run of the resting-state scan, followed by 8 task runs.

2.3 Stimuli and procedures for experiment 3

In experiment 3 (day 3), participants were asked to attend unilateral images presented within a circular

aperture, consistent with the method used in Experiment 1. The images were drawn from three distinct

categories:

Faces: Asian faces with an equal gender distribution (50% female).

Scenes: Equally divided between indoor and outdoor settings (50% outdoor).

Scrambles: Phase-scrambled images superimposed with a grating, with 50% of the gratings spanning

the 1st and 3rd quadrants.

To ensure visual consistency across images, they were standardized for luminance histograms and

spatial frequency using the SHINE toolbox (as referenced from Willenbockel et al., 2010).

At the onset of each run, participants were instructed to pay extra attention to a designated category

throughout that run. The procedure (Fig.3) began with the center dot flashing three times over two

seconds, signaling the forthcoming presentation of the cue and target. Following this, both the cue and

bilateral images from the same category were presented simultaneously. This images briefly shifted in

one direction before returning to their original position (0.25s+0.25s). During the next 1.5 seconds,

participants first pressed a key to indicate whether the direction of the images' motion matched that of the

arrow at the center. A green dot then appeared at the center, reminding participants to categorize the
image in their attended location. If the image matched the category to which they had been instructed to
pay attention, participants were required to identify specific content about the image (e.g., determining
whether a face was female or male) and responded by pressing either key '1' or '2'. If the image did not
belong to the designated category, participants simply pressed '3'.

Each run consisted of 32 trials: 50% of the trials featured images from the emphasized category, and
the remaining 50% were evenly distributed between the other two categories (25% each). Each trial
lasted either 8, 10, or 12 seconds. Across Experiment 3, there were a total of nine runs, with each
category being the focus of three runs. The accuracy of extra judgements is 82.9% for face, 81.8% for
scene and 84.3% for scramble [$F_{(1.373, 19.23)} = 1.096$, $P=0.3308$].

3 MRI Data Acquisition

MRI scanning was conducted using a 3T Siemens Prisma scanner at the Beijing MRI Center for Brain
Research (BMCBR), utilizing a standard 20-channel head coil. High-resolution T1-weighted anatomical
images were acquired at the start of each session (TR = 3000 ms; TE = 3.02 ms; 176 slices; slice
thickness = 1 mm; no inter-slice gap; field of view = 256 mm; flip angle = 8°; image matrix: 256×256).

Functional data were collected with gradient-echo EPI sequences (TR = 2000 ms; TE = 30.0 ms; 52
slices; slice thickness = 2 mm; no inter-slice gap; voxel resolution 2.0 x 2.0 x 2.0 mm, field of view = 192
mm; flip angle = 80°; image matrix: 96×96).

4 MRI Data Analysis

fMRI data were preprocessed and analyzed using FreeSurfer and AFNI and custom Python code
(primarily the mripy package, https://github.com/herrlich10/mripy). The preprocessing steps included de-
spiking, slice timing correction, EPI distortion correction (PE blip-up), rigid body motion correction, spatial
smoothing (4 mm FWHM Gaussian kernel, for the task runs), and per run scaling (as percent signal
change).

For task runs, general linear models were used to estimate BOLD signal change from baseline for each
stimulus condition. For each individual, bilateral FFA, OFA and PPA were defined based on the functional
contrast between faces and scenes from experiment 3. Bilateral V1 were defined using function contrast
[attend contra moving dot - baseline] (P < 0.001, uncorrected), and IPS, FEF and MT were defined using
the same contrast (P < 0.01, uncorrected) and referring the cerebral atlas (Glasser, 2016). Due to the lack
of standardized anatomical definitions, bilateral VFC and TPJ were identified using the same functional
contrast (P < 0.01, uncorrected) (Vossel et al. 2014). Bilateral hPIT was localized using data from all three
tasks in Experiment 1, by intersecting activation maps on the inferotemporal surface, while avoiding the
locations of FFA and PPA.

The resting-state data was preprocessed similarly to the functional data, with additional steps for
removing white matter and cerebrospinal fluid signals. Subsequently, confound regression, spatial
smoothing (4 mm), and bandpass filtering (0.01–0.1 Hz) were performed using AFNI's 3dTproject.
220
221   **Results**
222
Localization and Validation of hPIT from Task-Invariant Activation to Spatial Attention
To identify brain area(s) that encode the location of attention invariant to stimulus type and cognitive
demand (Stemmann and Freiwald 2019), we employed three different spatial attention tasks in Exp. 1. In
all three tasks, participants were instructed to fixate on a center dot and pay attention to circular aperture
on one side, as directed by the cue. First task was about motion discrimination, and the second and third
task required the discrimination of color proportion and shape proportion.
To isolate cortical areas modulated by endogenous spatial attention, contrasts between the conditions
[attend contralateral – attend ipsilateral] were calculated for each of the three task blocks. Significant
attention-modulated voxel maps were generated for individual participant (in most cases $p < 0.05$, cluster
size > 20, but for activated voxel number > 10000, using $p < 0.01$). Then we took the intersection of these
three maps and projected it to the cortex in every participant (example shown in Fig 4A), as our goal was
to locate hPIT in inferotemporal cortex of human brain which should exhibit the properties of priority map,
specifically in this context, insensitive to stimulus and cognition dimension. On the intersection of three
maps, we found bilateral brain areas significantly modulated by spatial attention in intra-parietal sulcus
(exclude the right hemisphere of S13) and inferotemporal cortex in all participants, and in most
participants also the middle temporal cortex (exclude S07). Only three participants (S03, S04, S11)
showed intersection area in the prefrontal cortex.
As illustrated in Fig 4B, the hPIT was manually identified bilaterally on the intersection maps within the
posterior and ventral part of mid-fusiform sulcus as a contiguous cluster of voxels that are non-
overlapping with fusiform face area (FFA) nor parahippocampal place area (PPA). In all participants, the
hPIT is ventromedial to the occipital face area (OFA) and posterior to the FFA. The locations of hPIT,
OFA and FFA identified in our study are shown in Fig 4C on the surface of standard brain
MNI152_2009c. The MNI coordinates delineating the center of mass (COM) for the ROIs are specified as
follows: hPIT (-34, -72, -14), (33, -73, -13); FFA (-43, -55, -19), (41, -54, -17); OFA (-46, -79, -3), (46, -74,
-8). The average cluster size of the hPIT is 210 voxels in left hemisphere and 191 voxels in right
hemisphere, with 2x2x2 mm^3 voxels. Detailed location and cluster size of hPIT in every participant's
cortical surface are reported in Supplementary Fig. 1.
Enhanced activation to attended than unattended moving dots in experiment 1, constrained by cerebral
atlas, allowed us to also locate cortical areas that are critical nodes of attention network (Corbetta and
Shulman 2002; Vossel et al. 2014) bilaterally, including medial temporal visual area (MT), IPS (classified

into IPS_P, the posterior part, and IPS_A, the anterior part), FEF, temporal parietal junction (TPJ), and

ventral frontal cortex (VFC) (see Methods). Additionally, V1 was included as a ROI, so that we can see

the consequences of attentional modulation in the primary visual cortex. Furthermore, the posterior part of

lateral–occipital cortex (LOp) and FFA) were localized using cortical parcellation and functional contrast

as controls, since they were adjacent to hPIT physically.

Attentional modulation of hPIT With and Without Bottom-up input

To investigate the modulation in hPIT and other brain areas applied by top-down attention (without

stimulus) and top-down combined with bottom-up attention (with stimulus), we manipulated the presence

or absence of visual stimuli in Exp. 2. Using event-related design, participants were required to direct their

attention to the left or right visual field based on central cues. A dot was then presented on the attended

side on half of the trials (50% probability), requiring participants to report, by pressing keys, its location

relative to two reference points that were constantly presented (Fig. 2).

ROI-based analysis was performed to BOLD activation signals from hPIT and other ROIs identified in

Exp. 1. Fig. 5A shows the averaged beta of contrast [attended - unattended (attend contralateral - attend

ipsilateral)] from bilateral ROIs for each participant, illustrating activation patterns of our ROIs under both

blank and dot conditions. A Two-Way ANOVA (n=15), considering blank/dot and ROI as factors, revealed

significant main effects for both: ROI [$F_{(2.246, 31.44)}$ =19.20, P<0.0001] and condition blank/dot [F

$_{(1.000, 14.00)}$ =20.41, P=0.0005]. A significant interaction between these two factors was also observed

[$F_{(4.061, 56.86)}$ = 10.98, P<0.0001].

The condition wherein participants were required to attend to one side in the absence of any stimulus

except the constant background, served to elucidate the impact of top-down attentional modulation. As

illustrated by the blue bars (Fig. 5A), top-down modulation was apparent in multiple ROIs (V1, hPIT, MT,

IPS_P, IPS_A, FEF, VFC), but it was strongest in hPIT. To establish quantitative metrics, a post-hoc

multiple comparison test (Dunnett, one-sided) was utilized to compare the strength of attentional

modulation between hPIT and other ROIs. The attentional modulation in hPIT was significantly stronger

than all the other ROIs : V1 [$q_{(14)}$ =2.648, adjusted P= 0.0451], MT [$q_{(14)}$ =2.856, adjusted P= 0.031],

IPS_P [$q_{(14)}$ =2.680, adjusted P= 0.0426], IPS_A [$q_{(14)}$ =3.883, adjusted P= 0.005], FEF [$q_{(14)}$

=3.823, adjusted P= 0.005], TPJ [$q_{(14)}$ =4.261, adjusted P= 0.002], VFC [$q_{(14)}$ =3.161, adjusted P=

0.018]. Critically, since higher-order regions may exhibit large receptive fields potentially reducing the

degree of contralateral bias, we specifically evaluated whether these frontal and parietal regions (FEF,

VFC, IPS_A, IPS_P) demonstrated engagement during top-down attention. One-sample t-tests confirmed

significantly greater activation during the attend-blank condition compared to baseline (all P < 0.05; see

Figure S2), providing direct evidence for their involvement in top-down attentional deployment.

With the introduction of a small dot as a target stimulus, we were able to examine the combined

influence of top-down and bottom-up attention. As indicated by the pink bars, hPIT again exhibited the

strongest attentional modulation effect, however, here the modulation effect reflects the combined effect

of top-down and bottom-up attention. Post-hoc comparison (Dunnett, one-sided) revealed again that hPIT

had significantly higher combined attentional modulation effect than all the other ROIs: V1[q (14) =5.717,

adjusted P< 0.001], MT [q (14) =5.443, adjusted P< 0.001], IPS_P [q (14) =6.742, adjusted P< 0.001],

IPS_A [q (14) =7.804, adjusted P< 0.001], FEF [q (14) =7.939, adjusted P< 0.001], TPJ [q (14) =7.944,

adjusted P< 0.001], and VFC [q (14) =7.554, adjusted P< 0.001]. Further, among all the ROIs, hPIT and

MT showed significantly stronger elevation in attentional modulation in the dot than the blank condition

[Bonferroni, hPIT: t (14) =5.321, adjusted P< 0.001; MT: t (14) =4.326, adjusted P =0.003].

Comparatively, the combined attentional effect was stronger in hPIT than in MT [paired t test, one-tailed, t

(14) =1.969, p=0.035]. Specific values representing attentional modulation for all ROIs (V1, MT, IPS, FEF,

TPJ, VFC, FFA, LOp) under both blank and dot conditions are provided in Supplementary Table S1.

As a supplement, comparison of attentional modulation across ROIs adjacent to hPIT was conducted

(Fig.5B). Results of Two-way ANOVA showed significant main effect of condition blank/dot [df=1, F

(1.000, 14.00) = 30.23, P<0.0001] and ROIs [df=2, F (1.921, 26.90) = 9.788, P=0.0007], as well as the

interaction [df=2, F (1.311, 18.36) = 5.691, P=0.0210]. Multiple comparisons (Tukey's) demonstrated that

the modulation in hPIT was significantly stronger than LOp and FFA in both conditions [Blank: q (14)

=3.822, P=0.043 for LOp, q (14) =4.581, P=0.015 for FFA; Dot: q (14) =7.054, P=0.001 for LOp, q (14)

=4.137, P=0.028 for FFA]. These findings highlight hPIT's distinctive pattern of attentional engagement

compared to the other two areas.

Complementary analysis revealed hemispheric differences in hPIT attentional modulation. A two-way

ANOVA (n=15, hemisphere x condition) was performed, results showed significant main effect of hPIT

hemisphere [df=1, F (1.000, 14.00) = 7.816, P=0.0143], significant main effect of condition [df=1, F

(1.000, 14.00) = 28.31, P=0.0001] and non-significant interaction [df=1, F (1.000, 14.00) = 0.9477,

P=0.3468].

Generally, while most ROIs showed significant modulation by attention in both the blank and dot

conditions (see supplementary Table 2), hPIT is unique in that it showed the strongest attentional

modulation in each condition as well as showing the largest elevation effect from blank to dot condition,

suggesting that hPIT is deeply engaged in both bottom-up attention and top-down attention.

Image Category-invariant response but load-sensitive attentional effect in hPIT

To effectively guide spatial attention, hPIT should be broadly responsive to different stimulus features

(i.e., lack of feature selectivity) but show high sensitivity to locations with salient and task-relevant

features. In addition, such spatial sensitivity should be modulated by attentional load. In order to explore

stimulus feature and attentional load sensitivity of hPIT, in experiment 3, images of three different

categories (face, scene, scramble) were presented, participants were required to pay attention to the

image on the left or right side according to the cue and report its moving direction. To modulate the

attentional load, in each run, participants were instructed to make more demanding judgements (higher
load) about the content of one pre-specified category of the attended images. For the pre-specified
category, the additional responses were female or male for faces, indoor or outdoor for scenes, overlay
grating tilted clockwise counterclockwise for the scrambled images (Fig. 3).

Fig.6A shows the response of hPIT to different categories of images in different conditions, indicated by
the beta values. To test if hPIT showed any effect of category preference, spatial attention and attention
load sensitivity, we first performed a three-way ANOVA (n=15, spatial attention x stimulus category x task
load). Results showed that the main effect of attention was significant [$F_{(0.7628, 10.68)} = 119.8$,
$P<0.0001$], the interaction between attention and category was significant [$F_{(2.000, 28.00)} = 4.068$,
$P=0.0281$], and the interaction between attention and task load was significant [$F_{(0.6964, 9.749)} = 8.307$,
$P=0.0230$]. Other main effects and interactions did not reach significance.

Since there is attention x category interaction, we further explored stimulus category sensitivity
separately in the attend and un-attend conditions. Bayesian repeated measures ANOVA were adopted to
compare the responses to face, scene and scramble stimuli under attend and un-attend conditions:
attend [$BF_{10}=0.93$], unattend [$BF_{10}=0.202$]. The small Bayesian Factors (<1) under both conditions
provide support that hPIT response was insensitive to these three categories, consistent with the criteria
for priority map.

In addition, as the main effect of attention is significant, to gain a better understanding of the attention
effect related to stimulus categories, we calculated the attentional modulation of hPIT for different image
categories and task load (Fig.6B). A Two-Way ANOVA (n=15, stimulus category x task load) was
performed. Significant main effects of category [$F_{(1.852, 25.93)} = 4.069$, $P= 0.0317$] and task load [F
$_{(1.000, 14.00)} = 8.307$, $P=0.0121$] were found while their interaction was non-significant [$F_{(1.527, 21.37)}$
$= 2.714$, $P=0.1004$]. Further investigating whether the observed attentional load effect (difference
between high load and low load condition) differed between hemispheres, we directly compared the load
modulation (averaged across stimulus categories) in left and right hPIT (Fig.6C). This comparison
revealed no significant difference between hemispheres [two-tailed paired t-test: $t_{(14)} = 0.4791$,
$P=0.6393$].

In general, unlike its adjacent object-processing areas, the response of hPIT is not sensitive to stimulus
category. There is evidence that attentional modulation in hPIT is slightly stronger to faces than scenes
and scrambled images, possibly reflecting the intrinsic saliency difference between these image
categories. The observation that the attentional modulation in hPIT is sensitive to task load further
supports hPIT's role as an attentional priority map.

Functional connectivity of hPIT to attentional networks
To unravel the functional connectivity of hPIT with the whole brain (especially the nodes of attention
network), resting-state data were analyzed, with ROIs defined by task-based data (experiment 1 and 3,

Fig. 7A) and cortical parcellation atlas (Glasser et al. 2016). Firstly, to visualize the patterns of functional

connectivity, we calculated the average time course of hPIT, FFA and LOp using small sphere-shaped

ROIs (2 mm radius) to avoid signal contamination for better visualization, and obtained the correlation

coefficient between the seeds and all other voxels of brain to generate functional connectivity maps from

the three spatially restricted seeds (Fig.7B). It is apparent that hPIT had strong functional connections

with nodes of dorsal (FEF, IPS, MT) and ventral (VFC, TPJ) attention networks. Then the functional

correlation using whole ROIs (hPIT, FFA, and LOp, example shown in Fig.7C) with bilateral attention

network nodes were calculated on each participant. The connection strength of bilateral hPIT, FFA, and

LOp with DAN and VAN were compared using Two-Way ANOVA [n=15, attention networks x seed ROIs],

results showed significant main effects of seed ROIs [$F_{(1.732, 24.25)} = 8.653$, P=0.0021] and attentional

networks [$F_{(1.000, 14.00)} = 60.56$, P<0.0001] (Fig.7D). Specifically, the correlation coefficients of hPIT

with attention networks were stronger than the other two seed areas (Dunnett, one-sided): PIT vs. FFA [q

(14) = 2.135, adjusted P= 0.0452], PIT vs. LOp [q (14) = 4.725, adjusted P=0.0003]. As for hemispheric

differences, a Two-Way ANOVA (Hemisphere [Left hPIT, Right hPIT] × Attention Network [Left DAN,

Right DAN, Left VAN, Right VAN]) revealed a significant main effect of Attention Network [$F_{(2.120, 29.67)}$

= 37.38, P < 0.0001], but no significant main effect of hPIT hemisphere was observed [$F_{(1.000, 14.00)}$ =

3.129, P = 0.0987] or the interaction [$F_{(1.266, 17.72)} = 0.3339$, P = 0.6223].

We also performed psychophysiological interaction (PPI) using data from experiment 2, which revealed

task-modulated right hPIT connectivity. During attended dot conditions (contrast [attend contralateral dot

– attend ipsilateral dot]), right hPIT showed significantly enhanced functional coupling with FFA [t (14) =

2.15, P = 0.048] and LOp [t (14) = 2.50, P = 0.029], with trending enhancement to TPJ [t (14) = 1.85, P =

0.088]. This suggests hPIT may allocate attentional resources to object-processing regions following

priority map generation.

The stronger connection of hPIT with each attentional nodes could be visualized in a circular

connectivity plot (Fig.7E). The strong functional connections of hPIT with attention networks, especially

the dorsal attention network, supports the important role it plays in attention control.

Discussion

It is traditionally accepted that the networks of frontal and parietal cortex play important roles in the

control and engagement of endogenous and exogenous attention, while regions of occipito-temporal

cortex specialize in the processing of colors and shapes, leading to the representation of scenes and

objects, such as faces, bodies, words (Ungerleider 2000; McCandliss et al. 2003; Moore et al. 2003;

Peelen and Downing 2005; Lafer-Sousa and Conway 2013; Baldauf and Desimone 2014; Grill-Spector

and Weiner 2014; Moore and Zirnsak 2017; Stemmann and Freiwald 2019). Our results provide new

distinct insights about the role of the human posterior inferotemporal cortex in the control and
implementation of attention. We identified a specific area within the posterior inferotemporal cortex, hPIT,
where its activation exhibited little category selectivity, but was strongly modulated by attention across
tasks, and reflected a combined effect of top-down attention and bottom-up attention. Further, the
attentional modulation in hPIT was significantly affected by attention load as well as image category,
consistent with the nature of attention. In addition, functional connectivity analysis revealed that hPIT,
compared to its neighbor areas FFA and Lop, exhibited stronger connectivity with nodes of the attention
network. Taken together, these results suggest that hPIT fulfills the key criteria of an attentional priority
map. It shows spatially specific responses that are strongly modulated by attention, relatively invariant to
stimulus features, and functionally integrated with both dorsal and ventral attention networks. Although
classical regions such as LIP and FEF are often associated with attentional control, they are less
consistent with our more stringent priority-map criteria in the present study, as their modulation was
mainly driven by top-down attention. Regarding input integration, hPIT is situated closer to high-level
visual areas, making it well positioned to integrate bottom-up saliency information with top-down signals.
The hPIT is in close proximity to other object-processing regions in the inferotemporal cortex, including
the FFA, the visual word form area (VWFA), and lateral occipital complex (LOC). The hPIT ((-34, -72, -
14), (33, -73, -13)) is posterior to the FFA (40, -55, -10) (Kanwisher et al. 1997) and VWFA (−43, −56,
−16) (Rauschecker et al. 2012; Zhang et al. 2018). LOC is localized by more activation when viewing
objects than scramble, consisting of two subregions: LO (in our paper, LOp) and Loa/pFs (Grill-Spector et
al. 2001). The hPIT is inferior to the LOp [bound by (-41, -77, 3) and (-36, -71, -13)], posterior to the
Loa/pFs (−38, −50, −17) (Grill-Spector et al. 1999).
Why does a brain region in the inferotemporal cortex exhibit the properties of an attention priority map,
integrating the top-down and bottom-up attention? Previous studies have suggested that LIP/IPS, FEF,
and SC are closely linked to attention guidance and eye movement and controls: LIP as a representation
of attentional priority that remaps across saccades, FEF as an eye movement controller receiving LIP
responses, and SC as reflecting the final saccade (Andersen et al. 1992; Thompson et al. 1996; McPeek
and Keller 2002; Shen and Paré 2007; Buschman and Miller 2009; Arcizet et al. 2011; Zhou and
Desimone 2011; Foley et al. 2017; Bisley and Mirpour 2019). On the other hand, both top-down and
bottom-up attention are frequently associated with objects, ideally an area serving as attentional priority
map should be positioned where it is easy to assess object information and also has broad connections
with other regions of the attentional networks. The hPIT's location in the inferotemporal cortex facilitates
the integration of key information from adjacent areas specialized in object-processing. As shown in our
functional connectivity results as well as in previous studies (Bogadhi et al. 2019; Sani et al. 2019; Sani et
al. 2021; Boshra and Kastner 2022), hPIT is connected with key areas of attention and eye movement
controls, such as LIP/IPS, FEF, and SC. The placement of an area serving attentional priority map in the

inferotemporal cortex, i.e., hPIT, has strategic advantages for both bottom-up information transmission

and top-down attentional modulation.

In addition to the strong connection between hPIT and the dorsal attention network, consistent with a

recent study (Meng et al. 2024), our functional connectivity analysis also revealed positive connections

with TPJ and VFC, as shown in individual connectivity maps, though these connections were weaker

compared to those with dorsal attention nodes. The connectivity with both the dorsal and ventral attention

networks further supports hPIT's unique role in bridging the bottom-up and top-down attention.

Our study is limited in that the effects of feature-based attention were not specifically examined in our

experiments. Besides, although faces used in the task had neutral expressions and the scene pictures

were also neutral, we cannot exclusively eliminate the possibility that potential semantic familiarity or

emotional salience may contribute to the subtle category-related effects in the results of experiment 3. In

addition, our current study lacks the temporal dynamic information of processing in hPIT and its

downstream and upstream brain areas due to the use of fMRI measurements. Future studies would

benefit from utilizing imaging methods with higher time resolution, with designs that address both spatial

and feature-based attention.

In summary, the hPIT showed strong attentional modulation across stimuli and tasks, with sensitivity to

attentional load, and more robust attentional modulation in the presence compared to the absence of

visual input. The hPIT also demonstrated function connectivity to both dorsal and ventral attention

networks. Together, our findings demonstrate the distinct role of hPIT in attention control, namely as an

attentional priority map that integrates endogenous and exogenous attentional processes.

Acknowledgments

This work was supported by STI2030-Major Projects (Grant Nos. 2021ZD0204200 and 2021ZD0203800);

and Key Research Program of Frontier Sciences, Chinese Academy of Science (Grant No. KJZD-SW-

L08).

References

Andersen RA, Brotchie PR, Mazzoni P. 1992. Evidence for the lateral intraparietal area as the parietal
eye field. Current Opinion in Neurobiology. 2(6):840–846. doi:10.1016/0959-4388(92)90143-9.

Arcizet F, Mirpour K, Bisley JW. 2011. A Pure Salience Response in Posterior Parietal Cortex. Cerebral
Cortex. 21(11):2498–2506. doi:10.1093/cercor/bhr035.

Baldauf D, Desimone R. 2014. Neural Mechanisms of Object-Based Attention. Science. 344(6182):424–
427. doi:10.1126/science.1247003.

Benarroch E. 2023. What Are the Functions of the Superior Colliculus and Its Involvement in Neurologic
Disorders? Neurology. 100(16):784–790. doi:10.1212/WNL.0000000000207254.

Bisley JW, Goldberg ME. 2010. Attention, Intention, and Priority in the Parietal Lobe. Annu Rev Neurosci.
33(1):1–21. doi:10.1146/annurev-neuro-060909-152823.

Bisley JW, Mirpour K. 2019. The neural instantiation of a priority map. Current Opinion in Psychology.
29:108–112. doi:10.1016/j.copsyc.2019.01.002.

Bogadhi AR, Bollimunta A, Leopold DA, Krauzlis RJ. 2019. Spatial Attention Deficits Are Causally Linked
to an Area in Macaque Temporal Cortex. Current Biology. 29(5):726-736.e4.
doi:10.1016/j.cub.2019.01.028.

Boshra R, Kastner S. 2022. Attention control in the primate brain. Current Opinion in Neurobiology.
76:102605. doi:10.1016/j.conb.2022.102605.

Buschman TJ, Miller EK. 2007. Top-Down Versus Bottom-Up Control of Attention in the Prefrontal and
Posterior Parietal Cortices. Science. 315(5820):1860–1862. doi:10.1126/science.1138071.

Buschman TJ, Miller EK. 2009. Serial, Covert Shifts of Attention during Visual Search Are Reflected by
the Frontal Eye Fields and Correlated with Population Oscillations. Neuron. 63(3):386–396.
doi:10.1016/j.neuron.2009.06.020.

Connor CE, Egeth HE, Yantis S. 2004. Visual attention: bottom-up versus top-down. Current Biology.
14(19):R850–R852. doi:10.1016/j.cub.2004.09.041.

Corbetta M, Patel G, Shulman GL. 2008. The Reorienting System of the Human Brain: From Environment
to Theory of Mind. Neuron. 58(3):306–324. doi:10.1016/j.neuron.2008.04.017.

Corbetta M, Shulman GL. 2002. Control of goal-directed and stimulus-driven attention in the brain. Nat
Rev Neurosci. 3(3):201–215. doi:10.1038/nrn755.

Desimone R, Duncan J. 1995. Neural Mechanisms of Selective Visual Attention. Annu Rev Neurosci.
18(1):193–222. doi:10.1146/annurev.ne.18.030195.001205.

Fecteau J, Munoz D. 2006. Salience, relevance, and firing: a priority map for target selection. Trends in
Cognitive Sciences. 10(8):382–390. doi:10.1016/j.tics.2006.06.011.

Foley NC, Kelly SP, Mhatre H, Lopes M, Gottlieb J. 2017. Parietal neurons encode expected gains in
instrumental information. Proc Natl Acad Sci USA. 114(16). doi:10.1073/pnas.1613844114. [accessed
2024 Aug 26]. https://pnas.org/doi/full/10.1073/pnas.1613844114.

Glasser MF, Coalson TS, Robinson EC, Hacker CD, Harwell J, Yacoub E, Ugurbil K, Andersson J,
Beckmann CF, Jenkinson M, et al. 2016. A multi-modal parcellation of human cerebral cortex. Nature.
536(7615):171–178. doi:10.1038/nature18933.

Grill-Spector K, Kourtzi Z, Kanwisher N. 2001. The lateral occipital complex and its role in object
recognition. Vision Research. 41(10–11):1409–1422. doi:10.1016/S0042-6989(01)00073-6.

Grill-Spector K, Kushnir T, Edelman S, Avidan G, Itzchak Y, Malach R. 1999. Differential Processing of
Objects under Various Viewing Conditions in the Human Lateral Occipital Complex. Neuron. 24(1):187–
203. doi:10.1016/S0896-6273(00)80832-6.

Grill-Spector K, Weiner KS. 2014. The functional architecture of the ventral temporal cortex and its role in
categorization. Nat Rev Neurosci. 15(8):536–548. doi:10.1038/nrn3747.

Herrington TM, Assad JA. 2009. Neural Activity in the Middle Temporal Area and Lateral Intraparietal
Area during Endogenously Cued Shifts of Attention. J Neurosci. 29(45):14160–14176.
doi:10.1523/JNEUROSCI.1916-09.2009.

Kanwisher N, McDermott J, Chun MM. 1997. The Fusiform Face Area: A Module in Human Extrastriate
Cortex Specialized for Face Perception. J Neurosci. 17(11):4302–4311. doi:10.1523/JNEUROSCI.17-11-
04302.1997.

Katsuki F, Constantinidis C. 2012. Early involvement of prefrontal cortex in visual bottom-up attention. Nat
Neurosci. 15(8):1160–1166. doi:10.1038/nn.3164.

Katsuki F, Constantinidis C. 2014. Bottom-Up and Top-Down Attention: Different Processes and
Overlapping Neural Systems. Neuroscientist. 20(5):509–521. doi:10.1177/1073858413514136.

Lafer-Sousa R, Conway BR. 2013. Parallel, multi-stage processing of colors, faces and shapes in
macaque inferior temporal cortex. Nat Neurosci. 16(12):1870–1878. doi:10.1038/nn.3555.

McCandliss BD, Cohen L, Dehaene S. 2003. The visual word form area: expertise for reading in the
fusiform gyrus. Trends in Cognitive Sciences. 7(7):293–299. doi:10.1016/S1364-6613(03)00134-7.

McPeek RM, Keller EL. 2002. Saccade Target Selection in the Superior Colliculus During a Visual Search
Task. Journal of Neurophysiology. 88(4):2019–2034. doi:10.1152/jn.2002.88.4.2019.

Meng Z, Huang Y, Wang W, Zhou L, Zhou K. 2024. Orienting role of the putative human posterior infero-
temporal area in visual attention. Cortex. 175:54–65. doi:10.1016/j.cortex.2024.04.006.

Moore T, Armstrong KM, Fallah M. 2003. Visuomotor Origins of Covert Spatial Attention. Neuron.
40(4):671–683. doi:10.1016/S0896-6273(03)00716-5.

Moore T, Zirnsak M. 2017. Neural Mechanisms of Selective Visual Attention. Annu Rev Psychol.
68(1):47–72. doi:10.1146/annurev-psych-122414-033400.

Ninomiya T, Sawamura H, Inoue K, Takada M. 2012. Segregated Pathways Carrying Frontally Derived
Top-Down Signals to Visual Areas MT and V4 in Macaques. J Neurosci. 32(20):6851–6858.
doi:10.1523/JNEUROSCI.6295-11.2012.

Noudoost B, Chang MH, Steinmetz NA, Moore T. 2010. Top-down control of visual attention. Current
Opinion in Neurobiology. 20(2):183–190. doi:10.1016/j.conb.2010.02.003.

Peelen MV, Downing PE. 2005. Selectivity for the Human Body in the Fusiform Gyrus. Journal of
Neurophysiology. 93(1):603–608. doi:10.1152/jn.00513.2004.

Ptak R. 2012. The Frontoparietal Attention Network of the Human Brain: Action, Saliency, and a Priority
Map of the Environment. Neuroscientist. 18(5):502–515. doi:10.1177/1073858411409051.

Rauschecker AM, Bowen RF, Parvizi J, Wandell BA. 2012. Position sensitivity in the visual word form
area. Proc Natl Acad Sci USA. 109(24). doi:10.1073/pnas.1121304109. [accessed 2024 Sep 6].
https://pnas.org/doi/full/10.1073/pnas.1121304109.

Rushworth MFS, Behrens TEJ, Johansen-Berg H. 2006. Connection Patterns Distinguish 3 Regions of
Human Parietal Cortex. Cerebral Cortex. 16(10):1418–1430. doi:10.1093/cercor/bhj079.

Sani I, McPherson BC, Stemmann H, Pestilli F, Freiwald WA. 2019. Functionally defined white matter of
the macaque monkey brain reveals a dorso-ventral attention network. eLife. 8:e40520.
doi:10.7554/eLife.40520.

Sani I, Stemmann H, Caron B, Bullock D, Stemmler T, Fahle M, Pestilli F, Freiwald WA. 2021. The human
endogenous attentional control network includes a ventro-temporal cortical node. Nat Commun.
12(1):360. doi:10.1038/s41467-020-20583-5.

Schall JD, Purcell BA, Heitz RP, Logan GD, Palmeri TJ. 2011. Neural mechanisms of saccade target
selection: gated accumulator model of the visual–motor cascade. Eur J of Neuroscience. 33(11):1991–
2002. doi:10.1111/j.1460-9568.2011.07715.x.

Shen K, Paré M. 2007. Neuronal activity in superior colliculus signals both stimulus identity and saccade
goals during visual conjunction search. Journal of Vision. 7(5):15. doi:10.1167/7.5.15.

Stemmann H, Freiwald WA. 2019. Evidence for an attentional priority map in inferotemporal cortex. Proc
Natl Acad Sci USA. 116(47):23797–23805. doi:10.1073/pnas.1821866116.

Thompson KG, Hanes DP, Bichot NP, Schall JD. 1996. Perceptual and motor processing stages
identified in the activity of macaque frontal eye field neurons during visual search. Journal of
Neurophysiology. 76(6):4040–4055. doi:10.1152/jn.1996.76.6.4040.

Treue S. 2003. Visual attention: the where, what, how and why of saliency. Current Opinion in
Neurobiology. 13(4):428–432. doi:10.1016/S0959-4388(03)00105-3.

Ungerleider SKALG. 2000. Mechanisms of Visual Attention in the Human Cortex. Annu Rev Neurosci.
23(1):315–341. doi:10.1146/annurev.neuro.23.1.315.

Vossel S, Geng JJ, Fink GR. 2014. Dorsal and Ventral Attention Systems: Distinct Neural Circuits but
Collaborative Roles. Neuroscientist. 20(2):150–159. doi:10.1177/1073858413494269.

White BJ, Berg DJ, Kan JY, Marino RA, Itti L, Munoz DP. 2017. Superior colliculus neurons encode a
visual saliency map during free viewing of natural dynamic video. Nat Commun. 8(1):14263.
doi:10.1038/ncomms14263.

Zhang B, He S, Weng X. 2018. Localization and Functional Characterization of an Occipital Visual Word
form Sensitive Area. Sci Rep. 8(1):6723. doi:10.1038/s41598-018-25029-z.

Zhou H, Desimone R. 2011. Feature-Based Attention in the Frontal Eye Field and Area V4 during Visual
Search. Neuron. 70(6):1205–1217. doi:10.1016/j.neuron.2011.04.032.

Figure 1. Schematic representation of the experimental paradigm for the Experiment 1. (A): The task was to report whether the direction of coherent motion on the attended side matched that of the white arrow. Note: The black arrow, representing one potential direction for the coherent dot movement, is used for illustrative purposes and was not actually presented during the experiment. (B): Pattern 1 and pattern 2, consisting of iso-luminant red and green dots, were presented sequentially. The task was to compare the color ratios of these patterns on the attended side and respond accordingly if any changes in the color ratio were detected. (C): Pattern 1 and pattern 2, consisting of equal number of small shapes (circles and squares) were presented sequentially. The task was to compare the shape ratios of these patterns on the attended side and respond accordingly if any changes in the shape ratio were detected.

Figure 2. Schematic depiction of the experimental design for Experiment 2. Following the flashing of central point, a cue is presented. In the dot condition, a dot appears on the attended side, prompting participants to report its relative position. In the blank condition, participants are instructed not to press any keys. (Light grey dots: serve as indicators, showing participants the potential target locations during the experiment.).

Figure 3. Schematic overview of the experimental paradigm for the experiment 3. At the start of the "face run", participants were instructed to pay heightened attention to the face images throughout that specific run. Following the central point's flashing sequence, a cue and bilateral images was presented. Participants first judged if the movement direction of the attended image aligned with the central arrow. Upon the central point changing to green, participants then responded based on the content of the attended image.

Figure 4. Functional localization of hPIT. (A): The intersection of three maps of three block tasks on one typical participant (S02). The cortical areas which are attached in red has shown significant activation in all three attention tasks. (B): The exact position of hPIT. Left: Positions of FFA and hPIT in statistical parametric maps of the contrast [attend face – attend scene] (p=0.05) (top) and hPIT in intersection map on the right hemisphere of S02. Right: The location of hPIT on both hemispheres of S02 circled by white line. The cortical areas which are attached in red has shown significant activation in all three attention tasks. (C): The positions of hPIT, OFA and FFA of 15 participants overlapped on surface of MNI152_2009c, with larger numerical values (manifested as deeper colors) indicating a higher degree of overlap among participants. The color scale ranging from grey to red delineates the spatial distribution of the hPIT, while the scale from grey to blue represents the FFA, and grey to yellow signifies the OFA.

Figure 5. Attention modulation in different brain regions. (A): The modulation pattern of V1, hPIT, MT, IPS, FEF, TPJ and VFC in condition blank (blue bar) and condition dot (pink bar), using beta of contrast: [attended -unattended (attend contralateral - attend ipsilateral)]. The modulation difference of attention between condition blank and condition dot reached significant level in PITd and MT. (B): The modulation pattern of hPIT, LOp, and FFA in condition blank (blue bar) and condition dot (pink bar). Error bars indicate 95% confidence interval. ∗∗∗ indicates the paired t test with significance of P < 0.001. ∗∗ indicates the paired t test with significance of P < 0.01. ∗ indicates the paired t test with significance of P < 0.05.

Figure 6. Response and attention modulation of hPIT to images of different categories. (A): The activation level of hPIT when attending to or not attending to images of different categories. Red bars indicate condition with attention in the receptive field (attended), while blue bars indicate condition with attention in the other side (unattended). Bars with higher chroma represent high load condition, lower chroma represent low load condition. (B): The modulation pattern of hPIT when presenting different categories of images with different attention load. White bars represent condition with low-load attention; Grey bars represent condition high-load attention. (C): The attention load effect (high load – low load) of bilateral hPIT, averaged across three categories. Error bars indicate 95% confidence interval.

Figure 7. Functional connectivity analysis of hPIT and its neighboring areas. (A): Activation map showing beta of contrast [attend contra moving dot - baseline] (p < 0.01, uncorrected) and the location of critical nodes in dorsal and ventral attention network on one typical subject (S03). (B): Thresholded map showing functional connectivity of seed sphere right-hemi hPIT, right-hemi FFA, and right-hemi LOp (Spearman's rank correlation coefficient > 0.2, p<0.05, uncorrected), averaged across subjects and projected onto the surface of standard brain MNI152_2009c. Color bar attached indicate the intensity of activation (A) and correlation (B). (C): The relative location of LOp, hPIT and FFA on the inflated cortical surface of parcellation map (Glasser's atlas). (D): Strength of functional connectivity of seed hPIT, FFA and LOp with DAN and VAN. Error bars indicate 95% confidence interval. ∗∗∗ indicates the significance of P < 0.001. ∗ indicates the significance of P < 0.05. (E): Circular plot for functional connectivity of seed hPIT, FFA and LOp with nodes of attention network of right hemisphere, with pink lines indicating connection with nodes of DAN, blue lines indicating nodes of VAN. Connections to left hemisphere nodes show similar but weaker trends. Opacity of each line connecting seed and nodes represents the rank of its connectivity strength (the strongest 100%, the middle 44%, the weakest 11%). Width of each line is scaled to its cubic numerical intensities of connectivity.