Orexin neurons couple pupil-linked arousal to Agrp hunger neurons to shape adaptive eating and food value perception

Authors: Seher Kosar[1], Eva Bracey[1], Yihui Du[1], Daria Peleg-Raibstein[1], Denis Burdakov[1*]

Affiliations: 1, ETH Zürich: Department of Health Sciences and Technology (D-HEST); Neuroscience Center Zürich (ZNZ); Institute of Food, Nutrition and Health; Institute for Neuroscience. Schwerzenbach, Switzerland

* correspondence: denis.burdakov@hest.ethz.ch

Abstract:

Arousal and hunger are basic essentials for survival, and their modulation is of general interest in daily life and across highly prevalent human conditions[1-15]. Here we show that their rapid fluctuations[16-23] are linked, identify responsible neurons, and demonstrate roles of this link in behavior, perception, and disease. First, using pupillometry as a direct metric of arousal[18], we found that self-paced pupil dilations coincide with inhibitions of hunger-causing Agrp neurons[20,24-28]. Neuron-specific ablation and optostimulation revealed that pupil-tracking hypocretin/orexin (H/O) neurons[19] are responsible for these inhibitions. Next, we compared Agrp neuron responses to different arousal-increasing sensory stimuli in mice with intact vs ablated H/O neurons. Arousing non-food stimuli distracted hungry mice from eating; this involved Agrp neuron inhibition, and required H/O neurons. With familiar food stimuli, we found that the feedforward inhibition of Agrp neurons by palatable food perception[20-23] was also disrupted by H/O neuron ablation, producing Agrp hyperactivity. Finally, we used EEG-EMG recordings to track Agrp neurons during normal sleep and pathological sleep paralysis (cataplexy), which - together with unexplained overeating[10,11,29] - is a key symptom of H/O neuron loss in humans[30-33]. In H/O-deficient mice, we found that Agrp hyperactivation is time-locked to cataplexy and REM sleep. Together, these results reveal an unanticipated[34,35] inverse link between hunger and volitionally-controllable[36] arousal, which relies on neurons that are prone to become defective during ageing and in many disorders[30,31,37-41].

Introduction

Eating and sleep-wake disorders are highly prevalent in modern humans[2-5], leading to extensive investigations into interplay of arousal, sleep, and hunger states[6-13]. Being more aroused and awake increases hunger and eating in the long term [13], because arousal generally consumes more energy than being asleep, and energy depletion is an established cause of hunger. On the neural level, hunger is caused by agouti-related peptide/neuropeptide Y co-expressing neurons (AGRP neurons) of the hypothalamic arcuate nucleus[20,24-28]. The AGRP neurons are slowly activated by energy depletion[20,24-28], and inhibited by multiple aspects of feeding: rapid "feedforward/predictive"[20] inhibition by food cues[21-23], slower inhibition by gut nutrients[42,43], and even slower "feedback/reactive" inhibition upon restoration of energy balance[20,42]. Interestingly, the arousal-promoting hypocretin/orexin-producing neurons[19,44-47] (H/O neurons) directly innervate AGRP neurons[48], and the H/O neurotransmitters rapidly (secs-mins) activate AGRP neurons *in vitro* [34,35], leading to models[49-51] implying that the hunger signals are generated not only slowly after energy depletion, but also rapidly by neural arousal signals (Fig. 1a).

However, such models fail to explain phenomena where feeding and arousal appear to change in opposite directions. For example, the human disorder type 1 narcolepsy involves a loss of H/O neurons but increased eating[10,11] (up to 4 times of normal[29]). In turn, during ageing, there are reports of both an overall decline in food intake ("the anorexia of ageing"[1]), and of difficulties of falling and staying asleep[14,15], recently linked to ageing-associated hyperexcitation of H/O neurons[41]. Furthermore, in anorexia, there are reports of increased wakefulness and arousal[6,12].

Therefore, here we re-examined the functional relationship between rapid fluctuations in arousal, H/O neurons and AGRP neurons, focusing on time-resolved fluctuations of arousal and consciousness states, H/O neuron hyperexcitation, and AGRP dynamics in a model of type 1 (i.e. H/O-deficient[32]) human narcolepsy. We present evidence for a rapidly-acting arousal brake on hunger signals, and propose a new model for hunger-arousal interplay.

82 **RESULTS**
83
Momentary pupil dilations correlate with reductions in AGRP neuron activity
We first tested how AGRP neuron activity is related to pupil size, a widely-used
indicator of momentary arousal that combines multiple sources of neurocognitive
information into a one-dimensional metric (reviewed in [18]). Pupil-linked arousal
fluctuates rapidly and closely matches the natural activity dynamics of H/O neurons
[19]. Contrary to the mainstream model that predicts positive co-modulation of H/O-
linked arousal and AGRP activity (Fig. 1a), we found pupil size to be negatively
correlated with AGRP activity (Fig. 1b-d). As expected from activation of AGRP
neurons by energy depletion[20,24], the dilation-associated AGRP inhibitions were
clearly seen in fasted mice (Fig. 1c-f), but less so in fed mice where AGRP neurons
are expected to already be inhibited (Supplementary Fig. 1).
95
96 Since pupil dilations correspond to H/O neuron activations[19], we therefore next
97 explored a new hypothesis: that H/O neurons exert a non-canonical inhibitory
98 influence on AGRP neurons, which contributes to dilation-associated AGRP neuron
99 inhibition. This hypothesis predicts that H/O neuron activation elicits AGRP neuron
100 inhibition, and that H/O neuron loss elevates AGRP neuron activity.
101
H/O neurons are capable of rapid inhibition of AGRP neurons
To test the first prediction, we selectively optostimulated H/O neurons by expressing
the excitatory optoactuator ChrimsonR under the control of orexin promoter[52] (Fig.
2a). We first analyzed responses of individual arcuate AGRP neurons *in vitro*, in
acute hypothalamic slices. We found that, among AGRP neurons that responded to
the optostimulation, the responses were mostly inhibitory (Supplementary Fig. 2b-c).
We reasoned that this inhibition may came from arcuate-projecting inhibitory neurons
activated by the H/O neuron optostimulation, since AGRP neurons do not directly
respond to the only inhibitory transmitter made by H/O neurons, dynorphin[53-56].
Arcuate-projecting VGAT neurons that inhibit AGRP neurons have been described in
the mediodorsal hypothalamus[57]. We therefore anatomically mapped the responses
to H/O neuron optostimulation of ARC-projecting (GCaMP retro-labelled) VGAT-cre
neurons around mediodorsal hypothalamic area (Supplementary Fig. 2d). This
confirmed strong optostimulation-induced excitation in a subset of arcuate-projecting
VGAT neurons (Supplementary Fig. 2e,f).
117
To resolve how the H/O neuron stimulation influences AGRP neurons *in vivo*, we
then recorded AGRP-GCaMP population activity using fiber photometry in freely-
moving mice (Fig. 2b,c). We found that AGRP neuron activity was inhibited upon H/O
neuron optostimulation in a stimulation-frequency dependent manner (Fig. 2b,c).
These results thus suggest that – in agreement with our hypothesis – H/O neuron
activity can inhibit AGRP neurons *in vivo*.
124
AGRP neuron inhibition during spontaneous pupil dilations requires H/O
neurons
To test the second prediction of our hypothesis, we compared pupil-associated
AGRP neuron activity in mice with selectively ablated H/O neurons[58] vs control mice
with intact H/O neurons (Fig. 2d). We observed a significantly weakened negative
correlation between pupil size and AGRP neuron activity in H/O neuron ablated mice
(Fig. 2e,f). To visualize effects more directly, we peak-aligned individual pupil

dilations in H/O ablated and control mice, and examined associated AGRP neuron
dynamics (Fig. 2e,f). This revealed significantly reduced dilation-associated AGRP
neuron inhibition in the H/O neuron ablated mice (Fig. 2g,h). Overall, the data in Fig.
2 thus identify H/O neurons as important mechanistic mediators of inhibitory coupling
between pupil dilation and AGRP neuron activity.
137
AGRP neuron hyperactivity during cataplexy in a mouse model of type 1
narcolepsy
Type 1 narcolepsy is a human neurological disorder involving abnormal sleep-wake
fragmentation; this disorder is caused by loss of H/O neurons [30,32]. H/O neurons
normally display brief (sec-min) increases in activity during both wakefulness[59-61] and
sleep[59,62-64]. Based on our data, we hypothesized that these H/O neuron activations
may limit AGRP neuron activity, and thus, in brains without H/O neurons, AGRP
neurons may be hyperactive during certain brain states.
146
When we analyzed spontaneous phasic activations of AGRP neurons, we indeed
found their amplitudes to be higher in H/O neuron ablated than in control mice (Fig.
3a-e). To define brain states corresponding to these aberrant AGRP
hyperactivations, we used electroencephalogram-electromyography (EEG-EMG). Of
particular clinical interest in type 1 narcolepsy are symptoms known as cataplexy,
defined by cessation of motor activity, postural collapse, and entry into a theta
oscillation-rich, "REM sleep-like" state directly from wakefulness [32,65]. It remains
unclear whether cataplexy episodes relate to hunger signals, which are presumably
increased in narcolepsy patients since they overeat[10,11,29]. Our orexin-DTR mouse
model of H/O neuron loss exhibits both overeating[58,66] and cataplexy-like episodes
(Fig. 3g,h). When we used established EEG-EMG-video criteria to identify cataplexy
episodes, we noticed that large spikes in AGRP neuron activity coincide with the
episodes (Fig. 3g,i). At a higher temporal resolution, we observed that AGRP neuron
activity sharply increased during episode onsets, and fell back shortly after their
offset (Fig. 3h,i).
162
We also systematically examined AGRP neuron activity corresponding to normal
brain states (wakefulness, REM sleep, NREM sleep) (Fig. 3 and Supplementary Fig.
3). We identified pronounced increases in AGRP neuron activity during REM sleep
(Fig. 3j-l). These REM-associated AGRP neuron activations were present in both
control and H/O neuron ablated mice, but – as expected from removal of an input
from H/O neurons [59,63,64]- were increased in amplitude in the H/O neuron ablated
mice (Fig3k,l).
170
171 Together, these data reveal that, as predicted by our hypothesis that H/O neurons
172 exert an inhibitory influence on AGRP neurons, there is temporally-defined AGRP
173 neuron hyperactivity in type 1 narcolepsy, that marks cataplexy episodes, as well as
174 REM sleep.
175
Role for H/O neurons in suppression of hunger signals by external stimuli
So far, we showed that H/O neurons shape AGRP neuron activity during internal
events such as spontaneous arousals, REM sleep, and cataplexy. Do H/O neurons
also shape AGRP neuron responses to external occurrences that may be
evolutionarily-important for the brain to process and respond to, such as unexpected
events, or proximity of familiar food? We hypothesized that external stimuli that

rapidly activate H/O neurons would inhibit AGRP neuron output in an H/O neuron-
dependent manner, and designed two assays to investigate this hypothesis.
184
First, we noted that among H/O neuron-activating stimuli are unexpected loud noises
[59,67]. We designed a feeding assay during which such a noise was briefly played
(Fig. 4a; we confirmed that this noise activated H/O neurons in Supplementary Fig.
4a). In agreement with our hypothesis, we found that the noise inhibited AGRP
neurons, and reduced spontaneous food intake if played while the mouse was eating
(Fig. 4b-d, control mice). These AGRP neuron and behavioral responses were
significantly reduced in H/O neuron ablated mice (Fig. 4b-d).
192
Second, we noted that AGRP neurons are known to be rapidly inhibited by food
proximity. This response - termed feedforward inhibition [20] - occurs before the food is
touched or digested, based on food sensory cues [21-23]. Because H/O neurons are
also activated by sensory cues related to food[68,69] (Supplementary Fig. 4b), and
contribute to AGRP neuron inhibition (Figs. 1-3), we tested whether they contribute
to the feedforward AGRP neuron inhibition by food. Indeed, we found that the
amplitude of AGRP neuron inhibition by presentation of a highly palatable food
(milkshake) was greatly reduced in H/O neuron ablated mice (Fig. 4f). The
feedforward inhibition of AGRP neurons by food is thought to reflect food value
sensing, since its amplitude is proportional to caloric value of the food, with larger
inhibition evoked by calorie-dense foods such as peanut butter than by chow [23]. We
found that H/O neurons also displayed such responses, but in the opposite direction:
they were strongly activated by peanut butter and less so by chow (Supplementary
Fig. 4b). We hypothesized that these H/O responses may affect AGRP neurons and
thus contribute to the AGRP neuron sensing of food value. Consistent with this, in
mice lacking H/O neurons, the inhibitory responses of AGRP neurons to food
presentation no longer discriminated between peanut butter and chow (Fig. 4g,h). In
contrast, AGRP neuron modulation by internal feedback signals ("hunger hormone"
ghrelin, nutrient depletion [23]) persisted in mice without H/O neurons (Supplementary
Fig. S4c,d).
213
214 Together, these experiments demonstrate that H/O neurons are required for normal
215 Agrp neuron modulation by important external stimuli. Without H/O neurons, Agrp
216 neurons of hungry mice remain hyperactive in the presence of palatable food, and
217 the mice over-prioritize eating.
218
219
220

DISCUSSION

Based on our findings, we propose the following model of rapid interactions between arousal and hunger. External stimuli rapidly activate H/O neurons[59-61,70,71]. This increases multiple general aspects of arousal[19,44,72], enabling the organism to optimally address incoming dangers or opportunities (reviewed in [49,73,74]). During caloric deficiency states when AGRP neurons are most active[20,24], these momentary arousals rapidly translate into inhibition of hunger-causing AGRP neurons. In the case of stimuli whose meaning is not yet clear, this H/O→AGRP inhibition may enable a rapid shift from feeding-focused state driven by AGRP neurons[20] to a general arousal state required to assess a new situation[75]. In the case of food-related stimuli, the H/O→AGRP neuron inhibition would contribute to the rapid sensing of food value by AGRP neurons, which is thought to be important for optimal regulation of ingestion[20,76]. If it holds, this model implies an inhibitory interplay between momentary arousal and hunger states, which – possibly together with the slow modulation of AGRP neurons by circadian rhythms [77] - may explain eating and body weight consequences of H/O neuron loss, including in humans with type 1 narcolepsy [10,11,29]. Our findings in the mouse model of type 1 narcolepsy also document previously-missed neural dynamics during key symptoms: the AGRP neuron hyperactivation during cataplexy. Together, these observations thus open new avenues for understanding the full scope of this chronic neurological disorder, as well as other human disorders associated with loss of H/O neurons[40].

249   **METHODS**
250
Animal procedures
All procedures followed Swiss Federal Food Safety and Veterinary Office Welfare
Ordinance (TSchV 455.1, approved by the Zurich Cantonal Veterinary Office), or the
UK Home Office regulations. Adult (older than 22 weeks) female and male C57BL6
mice were used, sex is noted in Supplementary Table 1. Animals were housed in a
12-hour reversed light-dark cycle at 22°C with 55% humidity, and had ad libitum
access to standard chow (3430 Kliba Nafag, Kaiseraugst, Switzerland) and water
unless stated otherwise (Supplementary Table 1). AGRP-IRES-cre mice and VGAT-
IRES-cre mice were from Jackson Lab (Jax no. 012899 and 028862, respectively).
To study mice without H/O neurons, we selectively ablated H/O neurons using the
previously validated hOrx-DTR mouse model[58] . To study AGRP neuron activity in
H/O neuron ablated mice, AGRP-IRES-cre and orexin-DTR mice were crossed
(AGRP-IRES-Cre x hOrx-DTR mice). Experiments were perfromed during the active
(dark) phase of the light cycle, except for studies of sleep which were done during
the inactive (light) phase (Fig. 3j-l and associated supplementary data).
266
Stereotaxic surgeries and viral vectors
Mice were anesthetized with 5% isoflurane, then transferred to the stereotaxic setup
(Kopf Instruments, Tujunga, USA) and maintained on 1.5–2% isoflurane. They were
given analgesic 30 mins before anaesthesia, and for three days post surgery.
Lidocaine was applied to the scalp before an incision was made to access the
cranium. Craniotomies were performed using a stereotax-mounted drill (RWD Life
Sciences). Viral vectors and implants were positioned at specific coordinates (given
below, as mm from Bregma unless stated otherwise). A Nanoject III injector was
used to inject the viral vectors at a speed of $1\,nl\,s^{-1}$.

For AGRP neuron recordings, AAV9.CAG.Flex.GCaMP6s.WPRE.SV40 (titer ≥ 1.9 ×
10^{13} vg/mL, from Addgene, MA, USA, 100842-AAV9) was injected unilaterally into
the arcuate nucleus of AgRP-IRES-cre, or AgRP-IRES-cre x hOrx-DTR mice, at the
following coordinates: anteroposterior -1.75, mediolateral -0.28, dorsoventral -6.1, -
6.0 to -5.9 (100nl per site). Optic fibers (400 µm, 0.39 NA, 1.25 mm metal ferrule,
Doric) were implanted at dorsoventral -5.85.

For H/O neuron recordings, AAV1.hORX.GCaMP6s.hGH ($2.5\,\times\,10^{12}$ GC ml^{-1},
Vigene Biosciences), previously demonstrated to selectively target H/O neurons[58],
was injected at coordinates: anteroposterior -1.35, mediolateral -/+0.9, dorsoventral -
5.7, 5.4 and 5.1 (50nl per site). A GRIN lens (ProView™ Integrated Lens 0.6mm x
7.3mm, Inscopix) was then implanted at dorsoventral -5.1.

For H/O neuron optostimulation, injection of AAV9-hORX-ChrimsonR-mCherry
($2\,\times\,10^{12}$ GC ml^{-1}, UZH Viral Vector Facility) previously demonstrated to
selectively target H/O neurons[52], or control injection (PBS), was performed bilaterally
at: anteroposterior -1.45, mediolateral ±2.8, dorsoventral -4.7 and -4.8 (150nl per
site). For *in vivo* optostimulation of H/O neurons, optic fibers (200-µm diameter, 0.39
NA, 1.25-mm ceramic ferrule; Thorlabs), angled at a 20-degree angle from vertical,
were placed 100 um above injection sites. For *in vitro* optogenetic-assisted probing
of H/O→AGRP neuron connectivity, AAV1-Syn-FLEX-GCaMP7f (Addgene, titer 1 ×
10^{13}, 104492) was also injected into the arcuate nucleus (coordinates as above) of
AgRP-IRES-cre mice.

For *in vitro* optogenetic-assisted calcium imaging probing functional connectivity from
H/O neuron to arcuate-projecting VGAT neurons, we injected AAV9-hORX-
ChrimsonR-mCherry into the lateral hypothalamus (coordinates as above) and
AAVretro-Syn-FLEX-GCaMP7f (Addgene, at titer ≥v7×10^{12} vg/mL, catalog# 104492)
into the arcuate nucleus (coordinates as above) of Vgat-IRES-cre mice.

For pupillometry, a custom-made headplate (Protolabs) was cemented to the skull.

For the electroencephalogram (EEG) and electromyogram (EMG) surgeries, a 3-
channel EEG/EMG headmount (Pinnacle Technology, model 8235: 6-Pin
Headmount, USA) was cemented to the skull and connected via stainless steel or
copper wire to four stainless steel screws (Fine Science Tools) serving as cortical
surface electrodes. The four screws were positioned at: (1) anteroposterior +1,
mediolateral +1; (2) anteroposterior +1, mediolateral -1; (3) anterioposterior -4.5 and
lateral +2, (4) from lambda: anteroposterior -1, mediolateral 0 (see Fig. 3). Screw (1)
functioned as the ground connection. A pair of stainless steel/copper wires was
implanted into the posterior neck muscle to capture EMG signals.

All implanted probes and headmounts were secured to the skull with three-part
dental cement (Superbond, Sun Medical). Mice were allowed to recover from surgery
for at least two weeks before experiments.
315
In vivo recordings of neural activity
Standard, dual-excitation-wavelength fiber photometry recordings and analyses
(described in detail in previous work [71,78,79]) were used to obtain AGRP-GCaMP6s
signals. H/O-GCaMP6s signals (Supplementary Fig. 4a,b) were obtained by whole-
field imaging using an INSCOPIX head-mounted microscope, as described in detail
in [71]. Fluorescence outputs were converted to normalised photometry signal % ΔF/F
as described in [79-81]. Following guidelines in [82], for between-subjects visualisations,
the % ΔF/F data were then converted to z scores using the function $z = (F_n − \mu)/\sigma$.
Here, F_n is the normalized photometry signal, and μ and σ are the mean and
standard deviation (respectively) of F_n of the whole recording session (Fig.1c, 3a,
3g, Supplementary Fig. 1a), or of a local baseline period around an event or a
stimulus presentation (Fig.1e, 2g, 3h-k, 4a-h; Supplementary Figs except
Supplementary Fig.1a). The local baseline period was -10 to -5 sec in 1e, 2g,
Supplementary 1c; 20s of awake state in 3h,j; 25 sec before transition onset in 3k,l
and Supplementary 3; and the pre-stimulus baselines depicted in remaining
supplementary figures.
332
Combined pupillometry and neural activity recordings
Pupil dynamics in head-fixed mice allowed to run freely on a rotating wheel were
recorded and analysed at constant level of light (to avoid effects of light-induced
pupil constriction) as previously described[19], using an infrared camera (Blackfly, FLIR
Systems, Imaging software Spinnaker SDK, frame rate 30 Hz). Pupil boundaries
were tracked using DeepLabCut to identify eight edge points in each video frame. A
circle-fitting algorithm in MATLAB calculated pupil radius, enabling precise pupillary
measurement across experimental conditions. Photometry and pupil size data were
aligned and downsampled to 10 Hz using the function resample, and z-scored
across the whole trace. To visualise AGRP-GCaMP6s signals corresponding to
similar pupil dilations in different mouse groups, dilation peaks were detected using
findpeaks in Matlab (MinPeakProm=1SD) and events in which running was less than
(0.05cm/sec) were removed. To analyse temporal correlations between AGRP-

GCaMP6s signals and pupil size, AGRP events were detected using findpeaks in
Matlab (min peak prominence 5% DF/F), time windows of 10s made around the peak
of the event were taken, and cross-correlated with corresponding pupil data using
the Matlab function xcorr with 'coeff' normalisation.
350
***In vivo* optogenetics**
Optogenetic stimulation of H/O neurons was performed as described in detail in our
previous work [19]. In Fig. 2, 30 sec trains of 5 ms laser pulses at each of the stated
frequences were applied at 3 minute intervals. Laser stimulation is repeated 3 times
per indicated frequency, in Fig. 2C every line shows the average of responses to
these repetitions for each frequency per mouse.
357
Brain slice recordings and optogenetics
300 μm hypothalamic coronal slices were acutely dissected and studied as desribed
in detail in our previous work [83]. The slices were placed in a recording chamber of an
Olympus microscope, continuously perfused with ACSF[83] at 21°C, gassed with 95%
O_2 and 5% CO_2. To stimulate ChrimsonR expressed in H/O neurons and axons,
HEKA Patchmaster software was used to deliver 20 Hz, 30 sec trains of 5 ms pulses
of a red laser (635 nm, Laserglow Technologies, 10mW at the fiber tip) through a
fiberoptic tip (200 um diameter) placed above mediobasal or mediodorsal
hypothalamic areas (defined in Supplementary Fig. 2). Calcium signals were
recorded using a a UMPlanF l10x objective, Olympus (0.3 NA,10.10mm working
distance) and Retiga ELECTRO camera (Imaging). Excitation light for calcium
imaging was provided by a 1 Hz 465 nm Sutter Lambda 4DG light source, controlled
by HEKA Patchmaster software. Images were acquired using HEKA SmartLUX and
HEKA Patchmaster software. Regions of interest (ROIs) were manually labeled
using ImageJ, and data were analyzed using MATLAB. Single-cell GCaMP signal
was converted to z scores using mean and standard deviation of its baseline (30 sec
before laser stimulation). If the z-score exceeded 1.5 SD for at least 5 seconds, the
response was categorized as excitatory. Conversely, if the z-score was below -1.5
SD for at least 5 seconds, the response was categorized as inhibitory.
377
EEG/EMG recordings
EEG/EMG data were recorded using the Pinnacle tethered system and software
(Pinnacle Technology, Lawrence, KS, USA). Signals were filtered by the preamplifier
(high pass above 0.5 Hz for EEG and 10 Hz for EMG). Vigilance/sleep states
(NREM, REM, and wake) were then automatically classified using an OSD4 script in
Spike2 (CED, https://ced.co.uk/downloads/scriptspkanal), with 5-second epochs.
After automatic detection, manual checking was also performed to ensure that the
following criteria for EEG/EMG brain states were met: wakefulness was defined as
desynchronized low-amplitude EEG and tonic EMG activity with phasic bursts;
NREM sleep as synchronized, high-amplitude, low-frequency (1-4 Hz) EEG and
reduced EMG activity compared with wakefulness with no phasic bursts; REM sleep
as EEG with prominent theta power (4-9 Hz) and a flat EMG. For identification of
cataplexy, videos of behavioral state were also recorded from above and beside the
transparent recording cage using 2 cameras (Basler acA1300-200um, Chromos
Industrial). Video and EEG data were combined to identify cataplexy, as outlined in [84]
(in brief: the cataplexy episodes were defined as an entry from wakefulness into a
REM-like state with behavioral paralysis lasting a minimum of 10 sec).
395

Lickometer analysis

Palatable food consumption was measured using a custom-built HPF dispenser (cat. no. 161K011, NResearch) with a capacitor-based lick sensor (cat. no. AT42QT-1010, SparkFun Electronics); data recorded at 500 Hz using custom Python scripts and a digital I/O device (NI-DAQmx, National Instruments). As palatable food, we used milkshake (energy milk strawberry flavor, 0.76 kcal ml^{-1}, Emmi AG) because there is abundant evidence that it is highly attractive for both mice and humans [85-87]. 6 µl of HPF was dispensed for every 10 detected licks. In Fig. 4c, lick data was z scored to the local baseline period 5 sec before the sound stimulus.

Sudden noise experiments

Hungry mice (24 h fasted) were placed in a cage containing the lickometer dispensing milkshake. Mice were previously habituated to both lickometer and milkshake. After mice licked continuously for 30 sec, a white noise was played through a loudspeaker placed next to lickometer (1 sec duration, 120 dB), while licking and AGRP neuron responses were recorded (4b-d). To quantify the AGRP reponse to the noise without the confound of licking and moving around the case, AGRP responses were additionally recorded in the head-fixed setup (the same as used for pupillometry) without food (Fig. 4d).

Food presentation experiments

Animals were habituated to peanut butter (Migros Classic Crunchy, 6.04 kcal g^{-1}) and milkshake (energy milk strawberry flavor, 0.76 kcal ml^{-1}, Emmi AG) in home cages overnight and to experimental cages for a minimum of two days before experiments started. On experimental days, animals were placed in the experimental cage for 15 min and then different foods (peanut butter, milkshake, or their standard chow [Maintenace Standard diet]) were placed into the case as previously described [23].

Histology

In order to validate the locations of AAV virus expression and the fiber optic, we prepared 50 µm brain sections as described in [78], and aquired images using a fluorescence microscope (Eclipse Ti2, Nikon).

Data analysis and statistics

Data processing and statistical evaluations were conducted using MATLAB (Mathworks, USA). All animals and data points were included in the analyses, except for those with no signal in fiber photometry at the beginning of experiments. The results of all statistical tests are summarised in figure legends, and are fully specified in Supplementary Table 2. T-tests were two-tailed, except where alternative hypotheses were directional in which case one-tailed tests were used (as specified in Supplementary Table 2). P values less than 0.05 were deemed statistically significant. Results and figures are displayed as data mean ± SEM, unless stated otherwise.

Acknowledgements:

This work was funded by ETH Zürich. We thank Mathilde Guillaumin and Han-Tao Li for their guidance on EEG/EMG surgery; Mathilde Guillaumin, Julia Harris, Bianca Viberti, and Micaela Borsa for their guidance on EEG/EMG data analysis for sleep

446 scoring and narcolepsy episode identification; and Nikola Grujic for setting up and
447 advising on pupillometry. We also thank Beate Laube for her help with mouse
448 genotyping. We are grateful to Antoine Adamantidis and Markus Schmid, as well as
449 Alexander Tesmer and Nikola Grujic, for helpful discussions and suggestions.
450
451 **Supplementary Materials:**
452 Tables 1 and 2
453 Supp. Figures 1-4
454
455

FIGURES

458
Fig. 1. Momentary pupil dilations correlate with reductions in AGRP neuron activity.
A. Previously-proposed relationship.
B. Method for direct experimental test of (A), including histology a coronal section showing
example optical fiber and injection site.
C. Example recording of pupil radius and AGRP-GCaMP6s activity (representative trace of n = 5
mice).
D. Cross-correlation analysis of pupil radius and AGRP-GCaMP6s signal (mean of n = 5 mice).
Right, quantification of correlation minima (n = 5 mice, ** p = 0.0079).
E. Pupil dilation events aligned with corresponding AGRP-GCaMP6s signals (152 events from n
= 5 mice).
F. Quantification of data in (E), ** p = 0.0019 (pupil) and ** p = 0.0011 (AGRP). Throughout the
figure: shaded areas or vertical bars show s.e.m.

480
Supplemetnary Fig. 1. Pupil-AGRP neuron relations in ad lib fed mice.
A.   Example recording of pupil and AGRP-GCaMP6s activity (representative trace of n = 5 ad
libitum fed mice).
B. Cross correlation analysis of pupil radius and associated AGRP-GCaMP6s signals (n = 5
mice). Right, quantification of correlation minima (n = 5 mice,** P = 0.02).
C. Pupil dilation events aligned with corresponding AGRP-GCaMP6s signals (170 events from n
= 5 mice).
D. Quantification of data in (C), ** P = 0.005 (pupil) and ** P = 0.006 (AGRP), n = 5 mice.
489
490
491
492

493
494
Fig. 2. Causal role for H/O neurons in rapid inhibition of AGRP neurons.
A. Left, targeting schematic for simultaneous H/O neuron optostimulation and AGRP neuron
recording. Right, coronal section showing example optical fiber and injection sites.
B. AGRP-GCaMP6s responses to H/O neuron optostimulation (laser bar) at different frequencies
(means of n = 5 mice in each dataset).
C. Quantification of (B) (frequency *P = 0.03; Chrimson vs. control *P = 0.02; n = 5 mice in each
group).
D. Schematic for AGRP neuron recording in mice ablated of H/O neurons (DTR+) and control
mice (DTR-).
E. Cross-correlation analysis of pupil radius and AGRP-GCaMP6s signals (n = 5 mice per group)
in H/O neuron ablated (DTR+) and control (DTR-) mice.
F. Quantification of correlation minima from (E) (n = 5 mice in each group, ** P = 0.0086 .
G. Pupil dilation events (top) aligned with corresponding AGRP-GCaMP6s signals (bottom) in
H/O neuron ablated (DTR+) and control (DTR-) mice (196 events from n = 5 DTR- mice, 172
events from n = 5 DTR+ mice). N.s. = P = 0.59; ***P<0.001 (comparison of means of -5 to +5
sec epochs).
H. Quantification of data in (G), top: n.s. = P= 0.166; bottom: *P=0.032 (n = 5 mice per group).
Throughout the figure: shaded areas or vertical bars show s.e.m.

Supplementary Fig. 2. Single-cell analysis of responses to H/O neuron optostimulation of arcuate AGRP neurons and arctuate-projecting VGAT neurons.

- A. Experimental design for single cell calcium imaging of AGRP-GCaMP7f and stimulation of H/O Chrimson neurons.
- B. Heatmap shows single-cell AGRP responses to red laser stimulation of H/O neurons. Red bar indicates stimulation time (n=241 cells from 3 mice).
- C. Characterisation of cells identified in (B). Cell activity was divided into three groups based on responses to H/O stimulation: Responses were either above (Excited cells), or below (inhibited cells) a threshold of 1.5 SD from the baseline for at least 5 seconds during laser stimulation. Cells with other responses were counted as "No response".
- D. Top, Experimental design for single cell calcium imaging of AGRP-GCaMP7f and stimulation of H/O Chrimson axons in various nuclei. Bottom, Histology shows an example slice where the posterior hypothalamic nucleus (PH) responded to H/O neuron optical stimulation.
- E. Top, Heatmaps displaying responses of GCaMP7f-expressing AGRP neurons to H/O Chrimson-expressing axon stimulation in different nuclei. Onset and offset of 635 nm laser stimulation is indicated by red lines.
  Coordinates for different nuclei were determined using The Mouse Brain in Stereotaxic Coordinates, Franklin & Paxinos, 1997, are listed in mm from bregma (AP, ML, DV) as follows:
  ARC, arcuate nucleus, -1.60 – 1.95, +/- 0.3, -5.9
  DMH - dorsomedial hypothalamic nucleus, -1.70-1.95, +/- 0.5, ~ -5.5
  PH - posterior hypothalamic nucleus, -1.85-1.95, -0.4, -4.75
  PVH - paraventricular hypothalamic nucleus -0.70-1, +/- 0.20, - 4.75
  VMH - ventromedial hypothalamic nucleus -1.60-1.95, +/- 0.5, -5.8
  Below, Average neuronal responses from heatmaps (n=3 mice).
- F. Quanfication of population shown in (E, n = 214 cells, ***P < 0.0001).

556
Fig. 3. Loss of H/O neurons causes AGRP hyperactivity during specific brain states.
A.   Example recordings of spontaneous AGRP neuron activity in control (DTR-) and H/O neuron
ablated (DTR+) mice (representative examples of n = 5 mice in each group).
B.   Amplitude distributions of AGRP neuron activity spikes, showing a shift to larger amplitudes in
DTR+ mice (***P= 0.000089, n = 5 mice in each group).
C-E. Qualification of the AGRP neuron spikes: peak amplitude (C), width (D), and
number (E) (n=5 per each group, *P = 0.02, ns= P=0.39, 0.46, respectively).
F.   Top, schematic of EEG/EMG recordigs and AGRP neuron recordings in H/O neuron ablated
(DTR+) and control (DTR-) mice.

G. Examples of AGRP neuron activity during cataplexy episodes (cat., gray boxes) from a mouse lacking H/O neurons (DTR+); typical example of n = 5 mice.

H. A higher temporal resolution example of a cataplexy episode (cat.) showing temporal relations between (from top to bottom) EEG frequency power spectrum, raw EEG signal, raw EMG signal, and AGRP-GCaMP6s activity.

I. Quantification of AGRP neuron activity before/during/after cataplexy (5 seconds pre., 10 seconds post), n = 31 episodes from 5 H/O neuron ablated mice. ***$P<0.0001$.

J. Examples of a REM sleep event in H/O neuron ablated (DTR+) and control (DTR-) mouse, showing temporal relations between (from top to bottom) EEG frequency power spectrum, raw EEG signal, raw EMG signal, AGRP-GCaMP6s activity, and brain state hypnogram derived from EEG (REM=rapid eye movement sleep; NREM=non-rapid eye movement sleep).

K. Group data from experiment in J, showing AGRP-GCaMP6s signals aligned to transition into REM. Data show means and s.e.m of n = 7 DTR-, n= 7 DTR+ mice.

L. Quantification of AGRP-GCaMP6s activity associated with specific brain states (mean activity during 5-7 sec after transition) in H/O neuron ablated (DTR+) and control (DTR-) mice. *$P=0.016$, ns=$P>0.05$ (post-hoc tests following RM ANOVA).

588
Supplementary Fig. 3. Additional data relating to Figure 3.
A. Top, AGRP activity during transitions from wake to NREM. Bottom, corresponding average
traces of transition heatmaps.
B. As in A, for NREM to REM state (REM durations > 10 sec, NREM > 30 sec). Bottom, same
trace as in main Fig. 3 (shown here again for consistency).
C. As in A, for REM to Wake transitions.
Data are means +/- SEM, n = 7 mice for each group across all panels.
596
597

599
600
Fig. 4. Effect of H/O neuron ablation on rapid AGRP neuron responses to food and non-food stimuli.
A. Schematic of the "sudden noise" experiment used to obtain data shown in B-D.
B. Example effects on eating (top) and AGRP-GCaMP6s (bottom) of the noise (representative
trace of DTR- n = 5 and DTR+ n = 4 mice).
C. Left: Group data for eating response show in B, mean of DTR- n = 5, DTR+ n = 4 mice. Right:
Quantification, *P = 0.002).
D. Left: Group data for AGRP-GCaMP6s response, data are mean and s.e.m of n = 5 mice per
group. Right: Quantification, *P = 0.01.
E. Schematic of the "food detection" experiment used to obtain data shown in f-h.
F. Left: AGRP neuron responses to presentation of a highly palatable food in H/O neuron
ablated (DTR+, n = 9) and control (DTR-, n = 11) mice. Right: Quantification, *P=0.04.
G. AGRP neuron responses to presentation of peanut butter (PB, darker colours) vs chow
(lighter colours) in H/O neuron ablated (DTR+) and control (DTR-) mice (data are mean and
sem of n = 6 DTR-, n = 7 DTR+ mice).
H. Quantification of data in G. DTR-: chow vs palatable food *P=0.04; DTR+: chow vs palatable
food P=0.30.

624
Supplementary Fig. 4. Additional data relating to main Fig. 4.
A. H/O neuron GCaMP responses to the sound stimulus used in Fig. 4 (n= 6 mice, **P = 0.003).
B. Left, H/O neuron responses to different food cues (PB = peanut butter). Right, Quantification of
H/O neurons responses (n =6 mice, ***P = 0.0018).
C. Left, Comparison of AGRP neuron responses to ghrelin i.p. injection in DTR- vs. DTR+ mice. Right,
quantification of responses to ghrelin (n = 4 DTR+, n = 6 DTR-; DTR+ vs. DTR-, ns = 0.125).
D. Left, Average AgRP neuron responses to chow in hungry (24h fasted) or ad lib fed H/O-ablated
mice.
E. Left, Quantification of data in (D), *P = 0.03, **P = 0.01.
634
635
636

Figure	Experiment	n (female mice)	n(male mice)	internal state
Fig1	AGRP-pupil	2	3	24 hr food deprivation
Fig2A-C	AGRP-chrimson vs heating control	2 and 3	2 and 3	Ab lib Fed
Fig2D-H	AGRP-pupil DTR- vs DTR+	2 and 1	3 and 4	24 hr food deprivation
Fig3A-E	AGRP-peaks DTR- vs DTR+	2 and 2	3 and 3	Ab lib Fed
Fig3F-H	AGRP-cataplexy DTR+	2	3	Ab lib Fed
Fig3J-L	AGRP-sleep DTR - vs DTR+	5 and 3	2 and 4	Ab lib Fed
Fig4B-D	AGRP-noise DTR- vs DTR+	2 and 1	3 and 4	24 hr food deprivation
Fig4F	AGRP-milkshake access DTR- vs DTR+	5 and 3	6 and 6	24 hr food deprivation
Fig4G-H	AGRP-H/O DTR- chow vs peanut butter	3	3	24 hr food deprivation
Fig4G-H	AGRP-H/O DTR+ chow vs peanut butter	1	6	24 hr food deprivation
S.Fig1	AGRP-pupil	2	3	Ab lib Fed
S.Fig.2A-C	AGRP- *in vitro* ca imaging		3	Ab lib Fed
S.Fig.2D-F	VGAT- *in vitro* ca imaging	2	3	Ab lib Fed
S Fig. 3	AGRP-sleep DTR- vs DTR+	5 and 3	2 and 4	Ab lib Fed
S Fig. 4a	H/O -noise		6	24 hr food deprivation
S Fig. 4b	H/O - chow vs peanut butter		6	24 hr food deprivation
S Fig. 4c	AGRP-ghrelin DTR- vs DTR+	2 and 1	4 and 5	Ab lib Fed
S Fig.4d,e	AGRP-chow ad lib vs chow fasted DTR+	1	6	Ab lib Fed or 24 hr food deprivation
S Fig.4e	AGRP-chow ad lib vs chow fasted DTR-	3	3	Ab lib Fed or 24 hr food deprivation

Supplementary Table 1. Summary of mouse sexes and nutritive states used in the study.
Mice were either ad lib fed or fasted as indicated in the table, for the following resons. H/O neuron optostimulation (Fig. 2) was done in fed mice, to avoid ceiling effects due to fasting-induced activation of H/O neurons. In sleep/wake studies (Fig. 3 and S3) mice were fed to avoid effects of fasting on sleep states. In Fig. 4A-F mice were fasted to induce simular continous eating across mice. In Fig. 4G-K, mice were fasted to maximise AGRP neuron responses to food.

Figure	Descriptors	n	Test used	Statistics	p-value
Fig. 1D	correlation coefficient	5 mice	One sample t-test	t(4) = -4.9270	0.0079
Fig. 1E	pupil radius AgRP-GCaMP	5 mice 5 mice	One sample t-test	t(4) = 7.2063 t(4) = -8.3445	0.0019 0.0011
Fig. 2C	Response of AgRP-GCaMP H/O Chrimson vs heating control 1Hz vs 40Hz	H/O Chrimson : 5 mice heating control : 5 mice	Repeated-measures ANOVA Post-hoc analysis	$F_{(1,8)}$ = 8.0941 $t_{(8)}$ = 3.372	0.0216 0.039
Fig. 2F	H/O DTR- vs H/O DTR+	H/O DTR- : 5 mice H/O DTR+ : 5 mice	one-tailed t-test	t(8) = -2.9950	0.0086
Fig. 2G	pupil radius AgRP-GCaMP H/O DTR- vs H/O DTR+	H/O DTR- : 5 mice, 196 events H/O DTR+ : 5 mice , 172 events	one-tailed t-test	t(366) = -0.2269 t(366) = -3.4453	0.5897 3.1828e-04
Fig. 2H	pupil radius AgRP-GCaMP H/O DTR- vs H/O DTR+	H/O DTR- : 5 mice H/O DTR+ : 5 mice	one-tailed t-test	t(8) = 1.0290 t(8) = -2.1377	0.1668 0.0325
Fig. 3B	Spontanenous AgRP-GCaMP H/O DTR- vs H/O DTR+ : Probability Density(%)	H/O DTR- : 5 mice H/O DTR+ : 5 mice	one-sided Kolmogorov-Smirnov test	D= 0.3693	8.96E-05
Fig. 3B	Spontanenous AgRP-GCaMP H/O DTR- vs H/O DTR+ : Peak amplitude	H/O DTR- : 5 mice H/O DTR+ : 5 mice	t-test	$t_{(8)}$ = 3.01	0.016823
Fig. 3D	Spontanenous AgRP-GCaMP H/O DTR- vs H/O DTR+ : Peak width	H/O DTR- : 5 mice H/O DTR+ : 5 mice	t-test	$t_{(8)}$ = 0.90	0.393373
Fig. 3E	Spontanenous AgRP-GCaMP H/O DTR- vs H/O DTR+ : Number of events	H/O DTR- : 5 mice H/O DTR+ : 5 mice	t-test	$t_{(8)}$ = -0.77	0.462538
Fig. 3I	AgRP-GCaMP in cataplexy epidoses pre cat. vs cat. vs post cat.	31 episodes from 5 mice	Repeated-measures ANOVA post-hoc pre vs cat. post-hoc cat. vs post	F(2, 60) =56.28	<0.0001 <0.0001 <0.0001
Fig. 3L	AgRP-GCaMP in brain states H/O DTR- vs H/O DTR+ NREM REM Wake	H/O DTR- : 7 mice H/O DTR+ : 7 mice	Repeated-measures ANOVA post-hocs DTR+ vs DTR-	F(2, 24) =4.7	0.0189 >0.9999 (NREM) 0.0116 (REM) >0.9999 (WAKE)
Fig. 4C	Licking H/O DTR- vs H/O DTR+	H/O DTR- : 5 mice H/O DTR+ : 4 mice	t-test	$t_{(7)}$ = 4.93	0.001697
Fig 4D	AgRP-GCaMP response to noise H/O DTR- vs H/O DTR+	H/O DTR- : 5 mice H/O DTR+ : 5mice	one-tailed t-test	t=(8) = -2.2362	0.0279
Fig. 4F	AgRP-GCaMP response to milkshake access H/O DTR- vs H/O DTR+	H/O DTR- : 11 mice H/O DTR+ : 9 mice	t-test	$t_{(18)}$ = 2.17	0.043693
Fig. 4H	AgRP-GCaMP response to food value Chow vs PB H/O DTR- H/O DTR+	H/O DTR- : 6 mice H/O DTR+ : 7 mice	paired t-test	$t_{(5)}$ = 2.72 $t_{(6)}$ = -1.13	0.041916 0.303224
Supp. Fig. 1B	correlation coefficient	5 mice	One sample t-test	t(4) = -3.77	0.0196
Supp. Fig. 1D	pupil radius AgRP-GCaMP	5 mice 5 mice	One sample t-test	t(4) = 5.5229 t(4)= -5.1634	0.0052 0.0067
Supp. Fig. 2F	Response of AgRP-GCaMP ARC DMH PH PVH VMH	28 cells from 5 mice 691 cells from 5 mice 214 cells from 5 mice 45 cells from 5 mice 71 cells from 4 mice	paired t-test	$t_{(27)}$ = 0.07, $t_{(690)}$ = 0.71 $t_{(213)}$ = -17.34 $t_{(44)}$ = -1.54 $t_{(70)}$ = -0.86	0.948068 0.475319 0.000000 0.131615 0.394080
Supp. Fig. 4A	H/O - GCaMP response to noise	6 mice	t-test	$t_{(5)}$ = 4.57	0.003001
Supp. Fig. 4B	H/O-GCaMP response to food value chow vs Peanut Butter	6 mice	paired t-test	$t_{(5)}$ = 5.17	0.001784
Supp. Fig. 4C	Ghrelin	H/O DTR- :7 mice H/O DTR+ : 6 mice	t-test	$t_{(11)}$ = 1.66	0.125901
Supp. Fig. 4E	Fasted vs fed	H/O DTR- : 6 mice H/O DTR+ : 7 mice	paired t-test	$t_{(5)}$ = -4.06 $t_{(6)}$ = -2.76	0.009747 0.032947

647
648
649 **Supplementary Table 2. Statistical information.**
655 **References:**

656
657
658
1   Malafarina, V., Uriz-Otano, F., Gil-Guerrero, L. & Iniesta, R. The anorexia of
ageing: physiopathology, prevalence, associated comorbidity and mortality. A
systematic   review.   *Maturitas*   **74**,   293-302   (2013).
https://doi.org:10.1016/j.maturitas.2013.01.016
2   Galmiche, M., Dechelotte, P., Lambert, G. & Tavolacci, M. P. Prevalence of
eating disorders over the 2000-2018 period: a systematic literature review. *Am*
J Clin Nutr **109**, 1402-1413 (2019). https://doi.org:10.1093/ajcn/nqy342
3   Hauri, P. J. in *Handbook of Psychology and Health, Volume 1*   211-260
(Routledge, 2021).
4   Ram, S., Seirawan, H., Kumar, S. K. & Clark, G. T. Prevalence and impact of
sleep disorders and sleep habits in the United States. *Sleep Breath* **14**, 63-70
(2010). https://doi.org:10.1007/s11325-009-0281-3
5   Thorpy, M. J. Classification of sleep disorders. *Neurotherapeutics* **9**, 687-701
(2012). https://doi.org:10.1007/s13311-012-0145-6
6   Malcolm, A., Toh, W. L., Crocker, K. & Phillipou, A. The contribution of sleep to
anorexia nervosa severity. *Eat Weight Disord* **27**, 1563-1568 (2022).
https://doi.org:10.1007/s40519-021-01286-2
7   Allison, K. C., Spaeth, A. & Hopkins, C. M. Sleep and Eating Disorders. *Curr*
Psychiatry Rep **18**, 92 (2016). https://doi.org:10.1007/s11920-016-0728-8
8   Olguin, P. *et al.* Medical comorbidity of binge eating disorder. *Eat Weight*
Disord **22**, 13-26 (2017). https://doi.org:10.1007/s40519-016-0313-5
9   da Luz, F. Q. *et al.* A systematic review with meta-analyses of the relationship
between recurrent binge eating and sleep parameters. *Int J Obes (Lond)* **47**,
145-164 (2023). https://doi.org:10.1038/s41366-022-01250-9
10   Baldini, V. *et al.* Eating disorders in narcolepsy type 1: Evidence from a cross-
sectional   Italian   study.   *J   Sleep   Res*   **33**,   e14150   (2024).
https://doi.org:10.1111/jsr.14150
11   Fortuyn, H. A. *et al.* High prevalence of eating disorders in narcolepsy with
cataplexy: a case-control study. *Sleep* **31**, 335-341 (2008).
12   Burger, P., Bos, R. W., Maas, J., Simeunovic-Ostojic, M. & Gemke, R. Sleep
disturbances in anorexia nervosa. *Eur Eat Disord Rev* **33**, 318-342 (2025).
https://doi.org:10.1002/erv.3148
13   Antza, C., Kostopoulos, G., Mostafa, S., Nirantharakumar, K. & Tahrani, A.
The links between sleep duration, obesity and type 2 diabetes mellitus. *J*
Endocrinol **252**, 125-141 (2021). https://doi.org:10.1530/JOE-21-0155
14   Mander, B. A., Winer, J. R. & Walker, M. P. Sleep and Human Aging. *Neuron*
94, 19-36 (2017). https://doi.org:10.1016/j.neuron.2017.02.004
15   Gulia, K. K. & Kumar, V. M. Sleep disorders in the elderly: a growing
challenge.   *Psychogeriatrics*   **18**,   155-165   (2018).
https://doi.org:10.1111/psyg.12319
16   Reimer, J. *et al.* Pupil fluctuations track fast switching of cortical states during
quiet   wakefulness.   *Neuron*   **84**,   355-362   (2014).
https://doi.org:10.1016/j.neuron.2014.09.033
17   Reimer, J. *et al.* Pupil fluctuations track rapid changes in adrenergic and
cholinergic   activity   in   cortex.   *Nature   Communications*   **7**   (2016).
https://doi.org:ARTN 13289
10.1038/ncomms13289

18   Grujic, N., Polania, R. & Burdakov, D. Neurobehavioral meaning of pupil size.
Neuron (2024). https://doi.org:10.1016/j.neuron.2024.05.029
19   Grujic, N., Tesmer, A., Bracey, E., Peleg-Raibstein, D. & Burdakov, D. Control
and coding of pupil size by hypothalamic orexin neurons. *Nat Neurosci* **26**,
1160-1164 (2023). https://doi.org:10.1038/s41593-023-01365-w
20   Andermann, M. L. & Lowell, B. B. Toward a Wiring Diagram Understanding of
Appetite   Control.   *Neuron*   **95**,   757-778   (2017).
https://doi.org:10.1016/j.neuron.2017.06.014
21   Mandelblat-Cerf, Y. *et al.* Arcuate hypothalamic AgRP and putative POMC
neurons show opposite changes in spiking across multiple timescales. *Elife* **4**
(2015). https://doi.org:10.7554/eLife.07122
22   Betley, J. N. *et al.* in *Nature* Vol. 14;521   180-185 (Nature Research, 2015).
23   Chen, Y., Lin, Y. C., Kuo, T. W. & Knight, Z. A. Sensory detection of food
rapidly   modulates   arcuate   feeding   circuits.   *Cell* **160**,   829-841   (2015).
https://doi.org:10.1016/j.cell.2015.01.033
24   Hahn, T. M., Breininger, J. F., Baskin, D. G. & Schwartz, M. W. Coexpression
of Agrp and NPY in fasting-activated hypothalamic neurons. *Nat Neurosci* **1**,
271-272 (1998).
25   Aponte, Y., Atasoy, D. & Sternson, S. M. in *Nature Neuroscience* Vol. 14   351-
355 (2011).
26   Gropp, E. *et al.* Agouti-related peptide-expressing neurons are mandatory for
feeding. *Nature Neuroscience* **8**, 1289-1291 (2005).
27   Krashes, M. J. *et al.* Rapid, reversible activation of AgRP neurons drives
feeding   behavior   in   mice.   *J Clin Invest* **121**,   1424-1428   (2011).
https://doi.org:10.1172/JCI46229
28   Luquet, S., Perez, F. A., Hnasko, T. S. & Palmiter, R. D. NPY/AgRP neurons
are essential for feeding in adult mice but can be ablated in neonates.
Science **310**, 683-685 (2005).
29   van Holst, R. J. *et al.* Aberrant Food Choices after Satiation in Human Orexin-
Deficient   Narcolepsy   Type   1.   *Sleep* **39**,   1951-1959   (2016).
https://doi.org:10.5665/sleep.6222
30   Nishino, S., Ripley, B., Overeem, S., Lammers, G. J. & Mignot, E. Hypocretin
(orexin) deficiency in human narcolepsy. *Lancet* **355**, 39-40 (2000).
31   Thannickal, T. C. *et al.* Reduced number of hypocretin neurons in human
narcolepsy. *Neuron* **27**, 469-474 (2000).
32   Bassetti, C. L. A. *et al.* Narcolepsy - clinical spectrum, aetiopathophysiology,
diagnosis   and   treatment.   *Nat Rev Neurol* **15**,   519-539   (2019).
https://doi.org:10.1038/s41582-019-0226-9
33   Dauvilliers, Y., Arnulf, I. & Mignot, E. Narcolepsy with cataplexy. *Lancet* **369**,
499--511 (2007).
34   van den Top, M., Lee, K., Whyment, A. D., Blanks, A. M. & Spanswick, D.
Orexigen-sensitive NPY/AgRP pacemaker neurons in the hypothalamic
arcuate   nucleus.   *Nature neuroscience* **7**,   493-494   (2004).
https://doi.org:10.1038/nn1226
35   Muroya, S. *et al.* Orexins (hypocretins) directly interact with neuropeptide Y,
POMC and glucose-responsive neurons to regulate Ca 2+ signaling in a
reciprocal manner to leptin: orexigenic neuronal pathways in the mediobasal
hypothalamus. *The European journal of neuroscience* **19**, 1524-1534 (2004).
https://doi.org:10.1111/j.1460-9568.2004.03255.x

36 Meissner, S. N. *et al.* Self-regulating arousal via pupil-based biofeedback. *Nat*
Hum Behav **8**, 43-62 (2024). https://doi.org:10.1038/s41562-023-01729-z
37 Thannickal, T. C., Lai, Y. Y. & Siegel, J. M. Hypocretin (orexin) cell loss in
Parkinson's   disease.   *Brain*   **130**,   1586-1595   (2007).
https://doi.org:10.1093/brain/awm097
38 Fronczek, R. *et al.* Hypocretin (orexin) loss in Alzheimer's disease. *Neurobiol*
Aging   **33**,   1642-1650   (2012).
https://doi.org:10.1016/j.neurobiolaging.2011.03.014
39 Peyron, C. *et al.* A mutation in a case of early onset narcolepsy and a
generalized absence of hypocretin peptides in human narcoleptic brains. *Nat*
Med **6**, 991-997 (2000).
40 Wang, C. *et al.* The Orexin/Receptor System: Molecular Mechanism and
Therapeutic Potential for Neurological Diseases. *Front Mol Neurosci* **11**, 220
(2018). https://doi.org:10.3389/fnmol.2018.00220
41 Li, S. B. *et al.* Hyperexcitable arousal circuits drive sleep instability during
aging.   *Science*   **375**,   eabh3021   (2022).
https://doi.org:10.1126/science.abh3021
42 Beutler, L. R. *et al.* in *Neuron* Vol. 96   461-475.e465 (Elsevier, 2017).
43 Su, Z., Alhadeff, A. L. & Betley, J. N. Nutritive, Post-ingestive Signals Are the
Primary Regulators of AgRP Neuron Activity. *Cell Rep* **21**, 2724-2736 (2017).
https://doi.org:10.1016/j.celrep.2017.11.036
44 Adamantidis, A. R., Zhang, F., Aravanis, A. M., Deisseroth, K. & de Lecea, L.
Neural substrates of awakening probed with optogenetic control of hypocretin
neurons. *Nature* **450**, 420-424 (2007).
45 Sakurai, T. *et al.* Orexins and orexin receptors: A family of hypothalamic
neuropeptides and G protein-coupled receptors that regulate feeding
behavior.   *Cell*   **92**,   573-585   (1998).   https://doi.org:10.1016/S0092-
8674(00)80949-6
46 de Lecea, L. *et al.* The hypocretins: hypothalamus-specific peptides with
neuroexcitatory activity. *Proc Natl Acad Sci U S A* **95**, 322-327 (1998).
47 De Luca, R. *et al.* Orexin neurons inhibit sleep to promote arousal. *Nat*
Commun **13**, 4163 (2022). https://doi.org:10.1038/s41467-022-31591-y
48 Horvath, T. L., Diano, S. & van den Pol, A. N. Synaptic interaction between
hypocretin (orexin) and neuropeptide Y cells in the rodent and primate
hypothalamus: a novel circuit implicated in metabolic and endocrine
regulations. *J Neurosci* **19**, 1072-1087 (1999).
49 Sakurai, T. The role of orexin in motivated behaviours. *Nat Rev Neurosci* **15**,
719-731 (2014). https://doi.org:10.1038/nrn3837
50 Milbank, E. & Lopez, M. Orexins/Hypocretins: Key Regulators of Energy
Homeostasis.   *Front   Endocrinol   (Lausanne)*   **10**,   830   (2019).
https://doi.org:10.3389/fendo.2019.00830
51 Tsujino, N. & Sakurai, T. Role of orexin in modulating arousal, feeding, and
motivation.   *Front   Behav   Neurosci*   **7**,   28   (2013).
https://doi.org:10.3389/fnbeh.2013.00028
52 Duffet, L. *et al.* A genetically encoded sensor for in vivo imaging of orexin
neuropeptides.   *Nat   Methods*   **19**,   231-241   (2022).
https://doi.org:10.1038/s41592-021-01390-2
53 Li, Y. & van den Pol, A. N. Differential target-dependent actions of
coexpressed   inhibitory   dynorphin   and   excitatory   hypocretin/orexin
neuropeptides. *The Journal of neuroscience : the official journal of the Society*

805   *for   Neuroscience* **26**,   13037-13047   (2006).
806   https://doi.org:10.1523/jneurosci.3380-06.2006

54   Chou, T. C. *et al.* Orexin (hypocretin) neurons contain dynorphin. *J Neurosci*
21, RC168 (2001). https://doi.org:20015644 [pii]

55   Schöne, C., Apergis-Schoute, J., Sakurai, T., Adamantidis, A. & Burdakov, D.
Coreleased orexin and glutamate evoke nonredundant spike outputs and
computations in histamine neurons. *Cell Rep* **7**, 697-704 (2014).
https://doi.org:10.1016/j.celrep.2014.03.055

56   Thomas, C. S. *et al.* Optogenetic stimulation of lateral hypothalamic
orexin/dynorphin inputs in the ventral tegmental area potentiates mesolimbic
dopamine neurotransmission and promotes reward-seeking behaviours.
Neuropsychopharmacology **47**,   728-740   (2022).
https://doi.org:10.1038/s41386-021-01196-y

57   Garfield, A. S. *et al.* Dynamic GABAergic afferent modulation of AgRP
neurons. *Nat Neurosci* **19**, 1628-1635 (2016). https://doi.org:10.1038/nn.4392

58   Gonzalez, J. A. *et al.* Inhibitory Interplay between Orexin Neurons and Eating.
Current   Biology **26**,   2486-2491   (2016).
https://doi.org:10.1016/j.cub.2016.07.013

59   Mileykovskiy, B. Y., Kiyashchenko, L. I. & Siegel, J. M. Behavioral correlates
of activity in identified hypocretin/orexin neurons. *Neuron* **46**, 787-798 (2005).
https://doi.org:10.1016/j.neuron.2005.04.035

60   Gonzalez, J. A., Iordanidou, P., Strom, M., Adamantidis, A. & Burdakov, D.
Awake dynamics and brain-wide direct inputs of hypothalamic MCH and
orexin   networks. *Nat   Commun* **7**,   11395   (2016).
https://doi.org:10.1038/ncomms11395

61   Lee, M. G., Hassani, O. K. & Jones, B. E. Discharge of identified
orexin/hypocretin neurons across the sleep-waking cycle. *J Neurosci* **25**,
6716-6720 (2005).

62   Ito, H. *et al.* Deficiency of orexin signaling during sleep is involved in abnormal
REM sleep architecture in narcolepsy. *Proc Natl Acad Sci U S A* **120**,
e2301951120 (2023). https://doi.org:10.1073/pnas.2301951120

63   Broglia, G. *et al.* Hypothalamic control of noradrenergic neurons stabilizes
sleep.   *bioRxiv*   (2024).
https://doi.org:https://doi.org/10.1101/2023.10.22.563502

64   Feng, H. *et al.* Orexin signaling modulates synchronized excitation in the
sublaterodorsal tegmental nucleus to stabilize REM sleep. *Nat Commun* **11**,
3661 (2020). https://doi.org:10.1038/s41467-020-17401-3

65   Hara, J. *et al.* Genetic ablation of orexin neurons in mice results in narcolepsy,
hypophagia,   and   obesity. *Neuron* **30**,   345-354   (2001).
https://doi.org:10.1016/S0896-6273(01)00293-8

66   Viskaitis, P. *et al.* Ingested non-essential amino acids recruit brain orexin cells
to suppress eating in mice. *Curr Biol* **32**, 1812-1821 e1814 (2022).
https://doi.org:10.1016/j.cub.2022.02.067

67   Takahashi, K., Lin, J. S. & Sakai, K. Neuronal activity of orexin and non-orexin
waking-active neurons during wake-sleep states in the mouse. *Neuroscience*
153, 860-870 (2008). https://doi.org:10.1016/j.neuroscience.2008.02.058

68   Petrovich, G. D., Hobin, M. P. & Reppucci, C. J. Selective Fos induction in
hypothalamic orexin/hypocretin, but not melanin-concentrating hormone
neurons, by a learned food-cue that stimulates feeding in sated rats.

Neuroscience **224**, 70-80 (2012). https://doi.org:10.1016/j.neuroscience.2012.08.036

69   Hassani, O. K., Krause, M. R., Mainville, L., Cordova, C. A. & Jones, B. E. in *The Journal of Neuroscience* Vol. 36 1747 (Society for Neuroscience, 2016).

70   Giardino, W. J. *et al.* Parallel circuits from the bed nuclei of stria terminalis to the lateral hypothalamus drive opposing emotional states. *Nat Neurosci* **21**, 1084-1095 (2018). https://doi.org:10.1038/s41593-018-0198-x

71   Karnani, M. M. *et al.* Role of spontaneous and sensory orexin network dynamics in rapid locomotion initiation. *Prog Neurobiol* **187**, 101771 (2020). https://doi.org:10.1016/j.pneurobio.2020.101771

72   Yamashita, A. *et al.* Aversive emotion rapidly activates orexin neurons and increases heart rate in freely moving mice. *Mol Brain* **14**, 104 (2021). https://doi.org:10.1186/s13041-021-00818-2

73   Li, J., Hu, Z. & de Lecea, L. The hypocretins/orexins: integrators of multiple physiological functions. *Br J Pharmacol* **171**, 332-350 (2014). https://doi.org:10.1111/bph.12415

74   Mahler, S. V., Moorman, D. E., Smith, R. J., James, M. H. & Aston-Jones, G. Motivational activation: a unifying hypothesis of orexin/hypocretin function. *Nat Neurosci* **17**, 1298-1303 (2014). https://doi.org:10.1038/nn.3810

75   de Araujo Salgado, I. *et al.* Toggling between food-seeking and self-preservation behaviors via hypothalamic response networks. *Neuron* (2023). https://doi.org:10.1016/j.neuron.2023.06.006

76   Chen, Y. & Knight, Z. A. Making sense of the sensory regulation of hunger neurons. *Bioessays* **38**, 316-324 (2016). https://doi.org:10.1002/bies.201500167

77   Sayar-Atasoy, N. *et al.* AgRP neurons encode circadian feeding time. *Nat Neurosci* **27**, 102-115 (2024). https://doi.org:10.1038/s41593-023-01482-6

78   Tesmer, A. L. *et al.* Orexin neurons mediate temptation-resistant voluntary exercise. *Nat Neurosci* **27**, 1774-1782 (2024). https://doi.org:10.1038/s41593-024-01696-2

79   Lerner, T. N. *et al.* Intact-Brain Analyses Reveal Distinct Information Carried by SNc Dopamine Subcircuits. *Cell* **162**, 635-647 (2015). https://doi.org:10.1016/j.cell.2015.07.014

80   Kim, C. K. *et al.* Simultaneous fast measurement of circuit dynamics at multiple sites across the mammalian brain. *Nat Methods* **13**, 325-328 (2016). https://doi.org:10.1038/nmeth.3770

81   Concetti, C., Bracey, E. F., Peleg-Raibstein, D. & Burdakov, D. Control of fear extinction by hypothalamic melanin-concentrating hormone-expressing neurons. *Proc Natl Acad Sci U S A* **117**, 22514-22521 (2020). https://doi.org:10.1073/pnas.2007993117

82   Sherathiya, V. N., Schaid, M. D., Seiler, J. L., Lopez, G. C. & Lerner, T. N. GuPPy, a Python toolbox for the analysis of fiber photometry data. *Sci Rep* **11**, 24212 (2021). https://doi.org:10.1038/s41598-021-03626-9

83   Burdakov, D. & Karnani, M. M. Ultra-sparse Connectivity within the Lateral Hypothalamus. *Curr Biol* **30**, 4063-4070 e4062 (2020). https://doi.org:10.1016/j.cub.2020.07.061

84   Willie, J. T. *et al.* Distinct narcolepsy syndromes in Orexin receptor-2 and Orexin null mice: molecular genetic dissection of Non-REM and REM sleep regulatory processes. *Neuron* **38**, 715-730 (2003).

85 Bussey, T. J. *et al.* New translational assays for preclinical modelling of cognition in schizophrenia: the touchscreen testing method for mice and rats. *Neuropharmacology* **62**, 1191-1203 (2012). https://doi.org:10.1016/j.neuropharm.2011.04.011

86 Kim, E. W. *et al.* Optimizing reproducibility of operant testing through reinforcer standardization: identification of key nutritional constituents determining reward strength in touchscreens. *Mol Brain* **10**, 31 (2017). https://doi.org:10.1186/s13041-017-0312-0

87 Edwin Thanarajah, S. *et al.* Habitual daily intake of a sweet and fatty snack modulates reward processing in humans. *Cell Metab* **35**, 571-584 e576 (2023). https://doi.org:10.1016/j.cmet.2023.02.015