Biodiversity modulates the cross-community scaling relationship in changing environments

Vojsava Gjoni[*,1,2,3]**, Florian Altermatt**[1,4]**, Aurélie Garnier**[1,5]** Gian Marco Palamara**[1,6]**, Mathew
Seymour**[7]**, Mikael Pontarp**[1,8]**, and Frank Pennekamp**[1]

[1] *Department of Evolutionary Biology and Environmental Studies University of Zurich, Zurich,*
Switzerland

[2] *Department of Biology, University of South Dakota, Vermillion, South Dakota, USA*

[3] *current address: Institute for Biological Resources and Marine Biotechnologies (IRBIM), National*
Research Council (CNR), Mazzara del Valo, Italy

[4] *Department of Aquatic Ecology, EAWAG: Swiss Federal Institute of Aquatic Science and Technology,*
Dübendorf

[5] *ECOBIO UMR CNRS 6553, University of Rennes, France*

[6] *current address: Italian National Research Council, Institute of Marine Sciences (CNR-ISMAR),*
00133 Rome, Italy;

[7] *School of Biological Sciences, The University of Hong Kong, Hong Kong SAR, China*

[8] *Department of Biology, Lund University, Lund, Sweden*

Correspondence E-mail: vojsavagjoni@cnr.it

Running Title: *Cross-community scaling relationships across richness and temperature*

Article Type: *Letter*

Words in abstract*: 149*

Words in main text*: 4995*

Number of references*:*

Number of figures*: 4*

Number of tables*: 1*

Number of textboxes*: 0*

Author contributions:

VG and FP conceived of the study. FA, AG, GP, MS, MP and FP designed the experiment and collected
the data. VG and FP conducted formal data analysis. VG and FP wrote the first draft of the manuscript.
All co-authors contributed to revisions of the manuscript.

34
ABSTRACT

Organismal abundance tends to decline with increasing body size. Metabolic theory links this size
structure with energy use and productivity, postulating a size–abundance slope of –0.75 that is invariant
across environments. We tested the robustness of this relationship across gradients of protist species
richness (1–6 species), temperature (15–25 °C), and time. Using replicated microcosms, we provide an
empirical test of how temperature and biodiversity jointly shape the cross-community scaling
relationship (CCSR). While our results support the expected of slope –0.75, we also found interactive
effects showing the relationship is not invariant. Warming altered abundance scaling with size
depending on richness; in high-richness communities, temperature favoured small protists, steepening
the CCSR slope. These context-dependent responses emerged over time, suggesting a role of size-
dependent species interactions in shaping responses to environmental change. Our findings demonstrate
that cross-community size scaling is not fixed but shifts dynamically with ecological context.

KEYWORDS: body size, protists, species richness, size-abundance relationship, temperature

INTRODUCTION

Across ecological communities, smaller species tend to occur in higher numbers than larger ones. This consistent inverse relationship between body size and abundance is a well-documented empirical pattern (Damuth 1991; White *et al.* 2007)*,* with a typical slope of approximately –0.75 on a log-log scale (Brown *et al.* 2004; Nee *et al.* 1991)*.* The widespread occurrence of this pattern has made it central to community ecology, offering insights into how body size structures populations and communities (Arim *et al.* 2011; Enquist *et al.* 1998; Giometto *et al.* 2013; Jacquet *et al.* 2020; Li 2002; Long & Morin 2005; Meehan *et al.* 2004; Morán *et al.* 2010; White *et al.* 2004, 2007)*.*

Beyond its descriptive power, the size–abundance relationship can link individual-level traits to ecosystem-level processes, serving as a tool for predicting how communities respond to environmental change (Gjoni *et al.* 2023; Gjoni & Glazier 2020). A central theoretical explanation for this pattern is the "energy equivalence", which posits that total population energy use remains constant across body sizes (Brown *et al.* 2004; Damuth 1981). This is based on the observation that population abundance declines with body mass ($\propto mass^{-0.75}$), while individual metabolic rate increases with mass ($\propto mass^{0.75}$), resulting in a size-invariant energy flux per unit area. This assumption is grounded in the idea that population abundance is limited by resource availability, resulting in a trade-off: species with greater per capita energy demands must occur at lower densities. An increase in resource input in-turn could lead to a proportional increase in population abundance or sustain the same number of individuals but with increased energy demand (Isaac *et al.* 2011).

While theoretical models predict consistent size-abundance relationships, mounting empirical evidence suggests empirical patterns may be more variable and context-dependent. Body size and abundance are responsive to biotic and abiotic factors, which may lead to changes in the size-abundance scaling when either the abundance of small or large organisms is particularly affected (Gjoni *et al.* 2023). Indeed, deviations from the expected -0.75 slope for the size-abundance have been observed (Arim *et al.* 2011; Gjoni *et al.* 2017, 2023; Gjoni & Basset 2018; Gjoni & Glazier 2020; Long & Morin 2005; Meehan *et al.* 2004), suggesting that energy used may not be the same across all organism sizes (Nee *et al.* 1991) and size-abundance scaling is not as invariant as assumed.

Temperature represents one of the most pervasive environmental factors that could disrupt these scaling relationships, yet its effects remain poorly understood. Warming is predicted to accelerate metabolic rates, increasing the energy required to sustain individuals following the metabolic theory of ecology (MTE) (Brown *et al.* 2004; Gillooly *et al.* 2001)*.* As a result, under constant energy availability, warming reduces carrying capacity by limiting the number of individuals that can be supported (Allen *et al.* 2002). However, because MTE assumes that temperature affects organisms similarly regardless of body size, it predicts that the slope of the size–abundance relationship should remain unchanged with warming (Fig. 1a). Contrary to this prediction, empirical studies often report deviations from the

expected –0.75 slope in size–abundance relationships under warmer conditions (Gjoni *et al.* 2023;
Morán *et al.* 2010; Pomeranz *et al.* 2022; Saito *et al.* 2021). One potential contributing factor is the
temperature–size rule, which posits that organisms tend to attain smaller body sizes at higher
temperatures (Atkinson 1995; Daufresne *et al.* 2009; Finkel *et al.* 2010; Gardner *et al.* 2011; Sheridan
& Bickford 2011). However, the temperature–size rule alone does not necessarily explain changes in
the slope of size–abundance relationships. If all organisms shrink proportionally, both small and large
taxa could increase in abundance equally, maintaining the original scaling. A change in the slope would
require a differential effect of temperature across size classes—where, for instance, smaller organisms
benefit more from warming than larger ones, or vice versa. In addition, ecological processes not
accounted for by metabolic theory, such as species interactions, may also contribute to observed
deviations. These interactions can be sensitive to both temperature (Burnside *et al.* 2014) and body size
(Freckleton & Watkinson 2001), further complicating predictions based solely on metabolic scaling
principles.

Biodiversity can play a critical role in shaping the size-abundance relationship due to species
interactions that determine community resource use and structure. While predation is a major focus of
studies investigating the size structure of communities, competitive communities have been less
frequently studied in the context of size-abundance scaling *(but see Long et al. 2006; Long & Morin*
2005). Research on biodiversity-ecosystem functioning has, for example, revealed that increased
species richness consistently leads to higher community abundance (Hooper *et al.* 2005; Tilman *et al.*
2001). Higher community abundance can be driven by highly productive species being more likely to
occur in species-rich communities (i.e., selection effects), or resource partitioning (i.e.,
complementarity) among species, reducing competition (Blackburn & Gaston 1996; Tilman *et al.*
2001). How community size structure is related to biodiversity effects is less well understood. Size
variation in competitive communities is often small (Tilman *et al.* 2004) and size structure is rarely
linked to complementarity or selection effects (but see Norkko *et al.* 2013). This suggests size-
abundance scaling is unaffected by species richness per se (Figure 1a).

Temperature and diversity can also interactively shape the size-abundance relationship, if they
disproportionally affect the interactions between larger and smaller organisms (Fig. 1b). Higher
temperatures tend to reduce the number of larger individuals, due to their higher metabolic demands,
creating a competitive advantage for smaller individuals due to their lower resource requirements
(Petchey et al. 1999). Temperature may also counteract some of the positive biodiversity effects
observed at higher richness due to direct negative physiological effects on biomass, leading to shallower
or even negative biodiversity-abundance relationships (Parain *et al.* 2019; Pennekamp *et al.* 2018). In
addition to these metabolic constraints, increased species richness can lead to reductions in population-
level density due to interference competition, niche overlap, or antagonistic interactions among co-
occurring species (DeLong 2021; Finke & Denno 2005; Turchin 2003). These negative diversity–

density relationships may also alter community size structure, especially if the strength or symmetry of competition varies with body size. Larger organisms are often competitively superior, as increased richness intensifies asymmetric resource competition (Lawton & Hassell 1981; Gómez & González-Megías 2002). Yet, whether larger or smaller species are favoured depends on the ecological and thermal context. Notably, in the absence of an interaction between activation energy and body size, the metabolic effects of temperature scale proportionally across sizes—meaning temperature accelerates the metabolism of many small organisms and a few large organisms to the same relative degree, with the proportional effect governed by $-E/kT$ (Brown *et al.* 2004). Therefore, temperature alone does not necessarily alter the slope of density–mass scaling unless body-size-dependent differences in thermal sensitivity are present. In species-rich communities, such differential sensitivities—amplified by complex interactions—may shift the competitive balance, causing richness to drive either steeper or shallower size-abundance slopes depending on which size classes gain a relative advantage (Fig. 1c). More generally, temperature and biodiversity can affect body size distributions and abundance through a suite of interconnected mechanisms. These include: shifts in metabolic rates and energy demand with temperature; differential thermal sensitivities among species, which alter interaction strength and performance; resource partitioning or competition intensity in diverse communities; and the tendency of larger organisms to dominate in systems with strong interference effects. By altering these underlying processes, temperature and diversity may interactively drive changes in both the intercept and slope of the size–abundance relationship, particularly when interactions between these factors amplify size-structured outcomes.

Effects of biotic and abiotic factors on the size-abundance scaling may further vary through time, since ecological communities are complex, adaptive systems which exhibit temporal dynamics in response to both biotic and abiotic factors (McGrady-Steed *et al.* 1997). Over time, changes in temperature and biodiversity may alter the relative proportions of organisms of different sizes within a community. Due to the equilibrium assumptions of the energy equivalence (Brown *et al.* 2004; Nee *et al.* 1991), we would expect the slope to converge to -0.75 in the long run. Indeed, experimental studies have shown that closed communities with controlled species richness (e.g., low vs. high richness) converge to a stable size-abundance relationship (Fig. 1d; Long & Morin 2005). In contrast, natural communities can exhibit fluctuating size-abundance relationships over time, driven by dynamic species interactions and environmental variability (Fig. 1e; White *et al.* 2004) that deviate from the expected slope.

Here, we investigated the size–abundance relationship across temperature and biodiversity (i.e., initial species richness) in controlled microcosms over time (Pennekamp et al. 2017, 2018). All communities were initially started from the same set of species, thus changes in size and abundance were integrating both phenotypic change in size and abundance. Given evidence that temperature can differentially affect organisms of different sizes, and that species richness may mediate these effects

through competitive interactions, our hypotheses explicitly test whether these factors alter size–abundance scaling. Based on a null model informed by the MTE, we hypothesize the following: (1) the size–abundance relationship initially follows a slope of –0.75, as predicted by MTE, (2) this slope may vary with temperature and species richness due to differential effects on small and large species, rather than remaining invariant, and (3) the slope may converge toward –0.75 over time as communities stabilize.

While ecological theory has traditionally emphasized within-community size spectra, which describes the distribution of individual body sizes and abundances within a single community at a given time, our study takes a broader perspective. We examine cross-community scaling relationships (CCSRs), which link the average body size of individuals to the total community abundance across different communities (White *et al.* 2007). While size spectra offer a useful summary of internal community structure, they do not retain the direct relationship between size and abundance necessary to assess ecological scaling patterns. In contrast, CCSRs explicitly link body size to total abundance at the community level and are therefore well suited for testing predictions based on energy equivalence and metabolic scaling theory (Isaac *et al.* 2011). Our experimental design included 53 protist communities per temperature treatment, each varying in initial species richness and composition. By comparing CCSRs across temperature and richness gradients, we were able to examine how warming alters the scaling of abundance with size—particularly whether these relationships deviate from the theoretical –0.75 slope expected under the metabolic theory of ecology. To further understand the role of species interactions on the CCSR, we created polycultures without interactions by pooling monoculture data with the same richness and composition (i.e., communities with no interaction expectation) as the real polycultures that we studied experimentally. Unlike previous observational studies, this controlled approach allowed us to directly test how temperature and diversity jointly shape cross-community scaling.

Figure 1 Conceptual overview of cross-community scaling relationships across temperature and diversity. a)
Temperature and diversity may affect abundance and size regardless of the size of individuals, which may lead to
changes in the intercept, but not the scaling. In contrast, smaller or larger individuals may be favoured at b) higher
(T1) or lower temperature (T2), or c) lower or higher species richness, leading to changes in the cross-community
scaling. Over time, the cross-community scaling relationship may be stable or converge to the expected slope of
-0.75 (d) or show more variable trajectories with endpoints that differ from the expected slope (e).

MATERIALS & METHODS

We used microcosms to test the effect of temperature and species richness on the CCSR. To this end,
we factorially manipulated temperature (15, 17, 19, 21, 23, and 25 °C) and species richness (1 to 6
species) in communities of six bacterivorous ciliate species differing in average cell volume (ordered
from the smallest to the largest): *Dexiostoma campylum (0.0000109 millicubicmeter), Tetrahymena*
thermophila (0.0000173 millicubicmeter), Loxocephalus sp. *(0.0000542 millicubicmeter), Colpidium*
striatum (0.000111 millicubicmeter), Paramecium caudatum (0.000180 millicubicmeter), and
Spirostomum teres (0.000488 millicubicmeter). The species were cultured in organic protist pellet
medium on which the freshwater bacterium *Serratia fonticola* was inoculated as the sole protist food
source. Due to the non-sterile handling of cultures, additional bacteria may have colonized the
microcosms throughout the experiment. The organic medium consisted of protist pellets (Carolina
Biological Supplies, Burlington, NC) at a concentration of 0.55 g per L of Chalkley's medium, and two

wheat seeds for slow nutrient release (for details on cultivation and experimental procedures, see also

Altermatt *et al.* (2015)). Stocks of protist cultures (monocultures) were established before the start of

the experiment to inoculate the microcosms. Single species cultures (richness = 1) were started at a

density of three individuals per mL in 100 mL medium kept in 250 mL Duran bottles. Multispecies

communities with species richness ranging from 2 to 6 species were initiated as a fixed fraction of the

species-specific carrying capacity. For example, 2-species communities were started with 20% of their

respective carrying capacity, while 4-species communities were started with 10% each. While having

the same fraction of carrying capacity, smaller and more abundant species started with a higher number

of individuals than larger and less abundant species. Since each species occurred the same number of

times in each richness level, there was no bias in terms of the number of large or small species across

richness levels. Since communities were closed to dispersal, richness can only decrease, and species

may be lost due to extinctions. We replicated each level of species richness and composition at least

twice for all levels including an additional replicate for the lowest and additional three replicates for the

highest levels of species richness resulting in 115 experimental units per temperature. Communities

were randomly assigned to their temperature level and placed in climate-controlled incubators (Pol-Eko

Aparatura, Wodzislaw, Poland) set to 15, 17, 19, 21, 23, or 25 °C. We used two independent incubators

for each temperature level. For an overview of the experimental design, refer to table S1.

To measure abundance and size of individual cells, we sampled each experimental unit every

day for the first 7 days, then 3 times per week for the following 50 days, and a final sampling 7 days

later. We used an established automated video species classification analysis (Pennekamp *et al.* 2015,

2017) to calculate the average size (i.e., volume) and abundance of each species. Since monocultures

and polycultures had different starting densities, we aligned the time series to make the dynamics

directly comparable. Please refer to Supplementary material section S1 for details. We then calculated

the cross-community scaling relationship relating total community abundance (sum of the abundance

of all species in the community) and the weighted mean size of an individual in the community (White

et al. 2007).

We used a regression approach to quantify the relationship between abundance (response) and

size, richness and temperature in our controlled microcosm communities. We scaled size, temperature

and richness treatments (by subtracting the respective mean from all values) to facilitate the

interpretation of model coefficients (Schielzeth 2010). The random structure for all models consisted

of a random intercept for community composition and a random intercept for incubator. We then

perform four analyses: 1) we estimated the overall cross-community scaling relationship across protist

communities, regressing total abundance on mean individual size (White *et al.* 2007). 2) We regressed

total abundance on mean individual size, temperature, and richness as well as all interactions between

main effects averaged over time. We included all possible interactions of the 3-way interactions models

(cell volume x temperature x richness). 3) We fitted the previous model (3 main effects and all possible

interactions, same random effects) to each time point to understand how estimates change through time.

4) To understand the role of species interactions on the CCSR, we generated communities with no

interaction expectation by pooling monoculture data with the same richness and composition as the real

polycultures that we studied experimentally. We then used the same model as 3) to compare the effects

of size, richness, and temperature, as well as their interactions between the real communities and

communities with no interaction expectation. Deviations in coefficient estimates would be putatively

driven by species' interactions. All analyses were performed in the statistical computing environment R

(version 3.6.1).

RESULTS

Interacting temperature and richness effects on CCSR

Averaging over time, we found a strong negative CCSR (b = -0.71, CI = -0.777 to -0.644) across

community richness and temperature treatments, undistinguishable from the expected value of b = -

0.75 (Fig. 2A). Although deviations from the linear fit are visible, they disappear when data was

stratified by richness (Fig. 2B).

Including temperature and richness in the model (Table 1), the CCSR slope (b = -0.749 CI = -

0.813 to -0.685) perfectly matched the expected value. However, temperature and richness jointly

changed the effect of size on abundance (volume:temperature coefficient = -0.015, CI = -0.024 to -

0.005 and volume:richness coefficient = -0.055, CI = -0.100 to -0.010). The effect of size on abundance

was furthermore mediated by a three-way interaction (volume:temperature:richness coefficient = -

0.012, CI = -0.018 to -0.007). In other words, at low richness, the abundance of small and large protists

was not affected by temperature (similar slope), whereas in more species rich communities, temperature

increased the abundance of smaller protists relative to larger ones, resulting in a steeper CCSR (Fig. 3,

Figure S1, Table 1). Importantly, these changes were not driven by extinctions of the larger species (see

sections S4). Using only two replicates for each richness level (leading to a balanced design) yielded

quantitatively and qualitatively similar results.

Figure 2 The relationship between centred average log10 cell volume ($\mu m \, \mu L^{-1}$) and average $\log_{10}$ total abundance

of protist communities, approximated with a simple linear fit (A). Deviations are visible at the lowest (< -0.5 log

cell volume) and largest average cell sizes (> 0.5 log cell volume) but disappear when stratified by richness (B).

Table 1 Temperature and species richness had an interactive effect on CCSR across protist communities. A

confidence interval (CI) not overlapping with zero indicates a significant effect. All significant terms including

the effect of volume on abundance are shown in bold. All predictors were mean centred for the analysis.

Term	Estimate [95% CI]
Intercept	4.564 [4.524, 4.603]
volume	**-0.749 [-0.813, -0.685]**
temperature	-0.019 [-0.023, -0.014]
richness	0.080 [0.051, 0.109]
volume:temperature	**-0.015 [-0.024, -0.005]**
volume:richness	**-0.055 [-0.100, -0.010]**
richness:temperature	-0.003 [-0.004, -0.001]
volume:temperature:richness	**-0.012 [-0.018, -0.007]**

Figure 3 Temperature and richness interact to alter protist community CCSR. The dashed line indicates the expected size-abundance scaling of -0.75. The figure shows the fitted size-abundance scaling based on the model presented in Table 1. A slope steeper (shallower) than -0.75 indicates that total abundance declines faster (more slowly) than expected with increasing average cell volume of the community.

CCSR through time

At the beginning of the experiment (Fig. 4A), CCSR was steeper (b = -0.9) than expected but stabilized on day 20 with a slope of b=0.75, which was expected. Afterwards the CCSR slope became shallower than expected (b = -0.35). Similar temporal patterns were observed when we used community evenness accounting for species extinctions and changes in relative abundance (see Section S3). Initially, communities with smaller average individual sizes tended to have higher abundance compared to those with larger individuals. Over time, communities with intermediate species richness (e.g., 5–7 species) showed relatively stable or slightly increasing abundance trends. By the end of the experiment, communities with larger average individual sizes had higher abundance than would be expected based on early patterns.

Figure 4: Temporal dynamics of effect sizes (and 95% confidence intervals) of a linear mixed-effects model where abundance is modelled as a function of size (V = volume), temperature (= T) and richness (= R) and their interactions for the observed and communities with no interaction expectation (in blue and red, respectively). The x-axis shows the day. A) Effect of cell size (V) on abundance, B–C) effect of two-way interactions between size and temperature (V x T), size and richness (V x R) on abundance. D) Three-way interaction between size, temperature and richness (V x T x R) on abundance. The dashed lines indicate a slope of -0.75 in panel A and zero in panels B–D (a deviation from zero would indicate a slope modification). A slope steeper (shallower) than -0.75 indicates that total abundance declines faster (more slowly) than expected with increasing average cell volume of the community.

Temporal analysis of the CCSR revealed dynamic patterns: temperature (V x T) initially
favored larger organisms but shifted to favor smaller ones after day 17, becoming non-significant by
the experiment's end. Species richness (V x R) initially reduced cell size effects but reversed after day
20 to promote larger organisms. The temperature-richness interaction (V x T x R) alternated between
negative and positive effects throughout the experiment, resulting in high richness communities being
more strongly affected by temperature than low richness communities.

Comparing real polycultures with communities having no interaction expectation revealed
contrasting temporal dynamics (Fig. 4). Initially, no-interaction communities showed steeper size-
abundance slopes than expected, indicating higher relative abundance of smaller organisms. After day
14, these slopes became shallower (approaching -0.40 by day 20) while real communities maintained -
0.75 until day 20. By day 30, no-interaction communities returned to expected values while real
communities became increasingly shallow. Unlike observed protist communities, the combined
temperature-richness effect showed no significant influence, indicating that temperature's impact on
size-abundance slopes was not richness-dependent in the no-interaction scenario.

DISCUSSION

Our study revealed that the cross-community scaling relationship of protist communities is mediated by
temperature, biodiversity (i.e., initial species richness), and time, challenging the widely held
assumption that CCSRs remain stable within a trophic level across time and space (Damuth 1981;
Enquist & Niklas 2001; Niklas & Enquist 2001). We found a negative CCSR (b ~ -0.75) across richness,
and temperature variation, in line with the expected slope of the energy equivalence (Damuth 1991;
Nee *et al.* 1991), confirming previous studies using the CCSR across phytoplankton (Li 2002),
macroinvertebrates, amphibians, and fish (Arim *et al.* 2011), rodent (White *et al.* 2004) and tree
communities (Enquist *et al.* 1998).

A CCSR is close to -0.75 is sometimes mirrored by a metabolic rate with a 0.75 scaling
exponent with mass (Brown *et al.* 2004), for instance in phytoplankton communities (Malerba *et al.*
2017; Perkins *et al.* 2019). Despite this support of the –0.75 slope, deviations are frequently reported in
within-community or population-level studies, where scaling relationships may vary depending on the
level of organization, sampling method, or ecological context (DeLong *et al.* 2010; Hatton *et al.* 2019;
White *et al.* 2007). An experimental study using protist communities initially divided into low and high
species richness showed no change over time with a cross-community size scaling of -1, which
converged over time (Long & Morin 2005). As we lack direct measurements of metabolic rates for our
focal protist species, we cannot confirm whether the classic metabolic scaling holds in this system.

Our experiment revealed a CCSR close to -0.75, suggesting that the scaling of metabolic rate
with body size may be a possible explanation. However, when we explicitly considered the abiotic and
biotic context of communities, we found that the CCSR of protist communities is mediated by

temperature, biodiversity and their interaction. Interacting effects between CCSR and temperature as
well as diversity, are inconsistent with energetic equivalence. In our high-richness communities, small
organisms were favored (more abundant than expected) with high temperature, whereas large organisms
were favored (more abundant than expected) when the temperature was low. Smaller organisms have
lower absolute resource requirements (Gillooly *et al.* 2001; Reuman *et al.* 2014), allowing for a higher
number of smaller organisms to be sustained (Li *et al.* 2006; Morán *et al.* 2010). Temperature may lead
to smaller organisms on average according to the temperature-size rule (Atkinson *et al.* 2003), however,
this physiological effect will affect organisms regardless of their size. Therefore, this cannot explain the
observed changes in CCSR, requiring additional explanations. Species richness increased abundance,
which is in line with experimental evidence that increasing the number of species (i.e., in this case,
competing for the same resources, such as bacteria) increases the total acquisition of resources due to
complementarity (Gómez & González-Megías 2002b). However, richness did not interact with size,
suggesting that complementarity and selection effects are not size-dependent in our communities. We
propose that temperature-dependent changes in species interactions provide the key explanation.

The interactions of the CCSR with temperature and richness may hence be driven by
temperature-dependent size asymmetries in resource competition (Cotgreave 1993; Nee *et al.* 1991;
Russo *et al.* 2003). One possible explanation is that temperature alters the competitive asymmetry
among organisms of different body sizes by differentially affecting their metabolic rates, feeding
efficiencies, and growth responses—traits that govern resource acquisition and competitive ability. At
higher temperatures, smaller organisms, with inherently faster metabolic rates, may outcompete larger
ones by exploiting resources more quickly. Conversely, at lower temperatures, the metabolic and
growth advantages of small size diminish, potentially allowing larger-bodied organisms to dominate
through superior resource access or longer generation times. These patterns are consistent with size–
temperature relationships described in the context of Bergmann's rule (Blackburn et al. 1999).

Another plausible mechanism for the increasing effect of temperature with rising species
richness is that temperature intensifies the strength and asymmetry of species interactions, particularly
in more complex communities. In species-rich systems, where interaction networks are more densely
connected, warming can amplify small differences in species' thermal sensitivities, resulting in
cascading effects on competition and resource partitioning (Dell et al. 2014; Gilbert 2014; O'Connor et
al. 2011). These dynamics may shift dominance patterns and energy flow, thereby altering the CCSR
more strongly under high richness. For instance, stronger interaction networks under warming can lead
to greater resource monopolization by thermally favoured species, magnifying size asymmetries (Sentis
et al. 2017; Uszko et al. 2017). Thus, species richness not only increases the potential for interaction-
driven community structuring but also makes the system more sensitive to temperature-induced change.

Previous analyses on our protist communities have shown that species interactions become
more asymmetric when temperatures rise. Tabi et al. (2020) showed that this leads to lower coexistence
of protist communities and a decoupling between abundance in isolation and abundance in

communities. Comparing the temporal dynamics of size-abundance scaling in communities with no
interaction expectation and real communities provides indirect evidence for this: in observed
polycultures, the initial slope aligned with the expected slope of -0.75, and over time, there was an
interactive effect of size, temperature and richness. In contrast, this pattern did not occur in communities
with no interaction expectation, as they lack the influence of species interactions. Maynard et al. (2020)
estimated coefficients that describe temperature effects on species interactions, showing mostly
negative effects of temperature and decreasing coefficients across the temperature gradient. Jointly, this
suggests that the observed steepening of the size-abundance scaling in high richness communities at
high temperature is driven by species interactions that are both temperature-dependent and size
asymmetric.

Our results show that size-abundance scaling can shift dynamically over time. Contrary to
expectations, the slope remained relatively stable in the first half of the experiment but deviated from
the -0.75 benchmark by the end, indicating a greater proportion of larger individuals in the community.
While this pattern diverges from the predictions of the Metabolic Theory of Ecology (MTE)—which
suggests that energy use becomes size-independent over time (Brown et al. 2004)—it highlights that
energy distribution in communities may become more complex, especially over extended timeframes
than suggested by theory. Specifically, larger individuals may acquire more energy than smaller ones
as communities mature, aligning with prior research suggesting resource monopolization by larger
protists under resource scarcity (Malerba & Marshall 2019; Tabi et al. 2019). Their capacity to store
resources may give them a survival advantage during periods of limited availability (Fenchel 1980).
This idea is further supported by hypotheses proposing that larger organisms endure fasting conditions
better due to their storage capacity (Millar & Hickling 1990), and experimental work showing that larger
microbial cells maintain efficiency under shifting nutrient levels (Malerba et al. 2018; Marshall et al.
2022). Similar trends have been observed in field studies, where temporal changes in CCSRs were
linked to shifts in species size and composition (White et al. 2004). However, given the arbitrary
endpoint of the experimental timeline, we caution against overinterpreting the final size-abundance
patterns as indicative of long-term outcomes.

Our study system differs from natural systems, which should be considered when interpreting
the results. First, since we did not allow for colonization by warm-adapted species/genotypes, we cannot
assess community turnover as a mechanism affecting community size structure under warming (Khaliq
et al. 2024). Second, microcosms are closed systems that do not allow for emigration or metacommunity
dynamics, which can buffer or balance ecological processes across space (Seymour *et al.* 2015).
Although all microcosms initially received the same amount of abiotic resources and energy inputs were
periodically replenished, the overall energy supply is not constant over time. Additionally, we did not
quantify the bacterial communities and their dynamics, which may further influence ciliate community
structure and interactions (Saleem *et al.* 2013). Microcosms can be colonized by air-borne bacteria; the
chance of such colonization events increases through time and may have been affected by our time

alignment. Replicate microcosms showed very repeatable dynamics, therefore we judge such chance
events have negligible effects on our results. Therefore, temporal effects need to be judged cautiously
and do not provide strong evidence against energetic equivalence. Third, we do not know whether the
observed changes in cross-community scaling are due to phenotypic plasticity, or also include
evolutionary responses. Despite these differences, our approach provides important complementary
insights to analyses of cross-community size scaling in natural communities.

CONCLUSION

Our experiments suggest that temperature and species richness play an important role in shaping the
size structure of ecological communities through time. We hence caution against assuming that energy
flux is size-independent in dynamic ecosystems, as changes in the biotic and abiotic environment may
change community size structure and hence the flow of energy. Testing these assumptions with
experimental communities provided new insights into the mechanisms driving community size
responses to environmental change, improving our ability to predict how biodiversity may influence the
size structure of communities in the face of global environmental change.

ACKNOWLEDGEMENTS
VG was funded by a Swiss National Science Foundation Scientific exchanges fellowship
(IZSEZ0_203498). We thank Roman Alther, Yves Choffat, Emanuel Fronhofer, Pravin
Ganesanandamoorthy, Suzanne Greene, Katherine Horgan, Elvira Mächler, Thomas M. Massie and
Owen Petchey for help with the data collection.

REFERENCES

Allen, A.P., Brown, J.H. & Gillooly, J.F. (2002). Global Biodiversity, Biochemical Kinetics, and the Energetic-Equivalence Rule. *Science*, 297, 1545–1548.

Altermatt, F., Fronhofer, E.A., Garnier, A., Giometto, A., Hammes, F., Klecka, J., *et al.* (2015). Big answers from small worlds: a user's guide for protist microcosms as a model system in ecology and evolution. *Methods Ecol Evol*, 6, 218–231.

Arim, M., Berazategui, M., Barreneche, J.M., Ziegler, L., Zarucki, M. & Abades, S.R. (2011). Determinants of Density–Body Size Scaling Within Food Webs and Tools for Their Detection. In: *Advances in Ecological Research*, The Role of Body Size in Multispecies Systems (ed. Belgrano, A.). Academic Press, pp. 1–39.

Atkinson, D. (1995). Effects of temperature on the size of aquatic ectotherms: Exceptions to the general rule. *Journal of Thermal Biology*, Effects of Rising Temperature on the Ecology and Physiology of Aquatic Organisms, 20, 61–74.

Atkinson, D., Ciotti, B.J. & Montagnes, D.J.S. (2003). Protists decrease in size linearly with temperature: ca. 2.5% per degree Celsius. *Proceedings of the Royal Society of London B: Biological Sciences*, 270, 2605–2611.

Blackburn, T.M., Gaston, K.J. & Loder, N. (1999). Geographic gradients in body size: a clarification of Bergmann's rule. *Diversity and Distributions*, 5, 165–174.

Brown, J.H., Gillooly, J.F., Allen, A.P., Savage, V.M. & West, G.B. (2004). Toward a Metabolic Theory of Ecology. *Ecology*, 85, 1771–1789.

Burnside, W.R., Erhardt, E.B., Hammond, S.T. & Brown, J.H. (2014). Rates of biotic interactions scale predictably with temperature despite variation. *Oikos*, 123, 1449–1456.

Cotgreave, P. (1993). The relationship between body size and population abundance in animals. *Trends in Ecology & Evolution*, 8, 244–248.

Damuth, J. (1981). Population density and body size in mammals. *Nature*, 290, 699–700.

Damuth, J. (1991). Of size and abundance. *Nature*, 351, 268–269.

Daufresne, M., Lengfellner, K. & Sommer, U. (2009). Global warming benefits the small in aquatic ecosystems. *Proceedings of the National Academy of Sciences*, 106, 12788–12793.

Dell, A.I., Pawar, S. & Savage, V.M. (2014). Temperature dependence of trophic interactions are driven by asymmetry of species responses and foraging strategy. *J Anim Ecol*, 83, 70–84.

DeLong, J.P. (2021). *Predator Ecology: Evolutionary Ecology of the Functional Response*. Oxford University Press.

DeLong, J.P., Okie, J.G., Moses, M.E., Sibly, R.M. & Brown, J.H. (2010). Shifts in metabolic scaling, production, and efficiency across major evolutionary transitions of life. *PNAS*, 107, 12941–12945.

Enquist, B.J., Brown, J.H. & West, G.B. (1998). Allometric scaling of plant energetics and population density. *Nature*, 395, 163–165.

Enquist, B.J. & Niklas, K.J. (2001). Invariant scaling relations across tree-dominated communities. *Nature*, 410, 655–660.

Fenchel, T. (1980). Relation between particle size selection and clearance in suspension-feeding ciliates. *Limnology and Oceanography*, 25, 733–738.

Finke, D.L. & Denno, R.F. (2005). Predator diversity and the functioning of ecosystems: the role of intraguild predation in dampening trophic cascades. *Ecology Letters*, 8, 1299–1306.

Finkel, Z.V., Beardall, J., Flynn, K.J., Quigg, A., Rees, T.A.V. & Raven, J.A. (2010).
Phytoplankton in a changing world: cell size and elemental stoichiometry. *Journal of*
Plankton Research, 32, 119–137.

Freckleton, R.P. & Watkinson, A.R. (2001). Asymmetric competition between plant species.
Functional Ecology, 15, 615–623.

Gardner, J.L., Peters, A., Kearney, M.R., Joseph, L. & Heinsohn, R. (2011). Declining body
size: a third universal response to warming? *Trends in Ecology & Evolution*, 26, 285–291.

Gilbert, P. (2014). The origins and nature of compassion focused therapy. *British Journal of*
Clinical Psychology, 53, 6–41.

Gillooly, J.F., Brown, J.H., West, G.B., Savage, V.M. & Charnov, E.L. (2001). Effects of
Size and Temperature on Metabolic Rate. *Science*, 293, 2248–2251.

Giometto, A., Altermatt, F., Carrara, F., Maritan, A. & Rinaldo, A. (2013). Scaling body size
fluctuations. *PNAS*, 110, 4646–4650.

Gjoni, V. & Basset, A. (2018). A Cross-community Approach to Energy Pathways Across
Lagoon Macroinvertebrate Guilds. *Estuaries and Coasts*, 41, 2433–2446.

Gjoni, V., Cozzoli, F., Rosati, I. & Basset, A. (2017). Size–Density Relationships: a Cross-
Community Approach to Benthic Macroinvertebrates in Mediterranean and Black Sea
Lagoons. *Estuaries and Coasts*, 40, 1142–1158.

Gjoni, V. & Glazier, D.S. (2020). A Perspective on Body Size and Abundance Relationships
across Ecological Communities. *Biology*, 9, 42.

Gjoni, V., Glazier, D.S., Wesner, J.S., Ibelings, B.W. & Thomas, M.K. (2023). Temperature,
resources and predation interact to shape phytoplankton size–abundance relationships at a
continental scale. *Global Ecology and Biogeography*, 32, 2006–2016.

Gómez, J.M. & González-Megías, A. (2002a). Asymmetrical Interactions Between Ungulates
and Phytophagous Insects: Being Different Matters. *Ecology*, 83, 203–211.

Gómez, J.M. & González-Megías, A. (2002b). Asymmetrical Interactions Between Ungulates
and Phytophagous Insects: Being Different Matters. *Ecology*, 83, 203–211.

Hatton, I.A., Dobson, A.P., Storch, D., Galbraith, E.D. & Loreau, M. (2019). Linking scaling
laws across eukaryotes. *Proceedings of the National Academy of Sciences*, 116, 21616–
21622.

Hooper, D.U., Chapin III, F.S., Ewel, J.J., Hector, A., Inchausti, P., Lavorel, S., *et al.* (2005).
Effects of Biodiversity on Ecosystem Functioning: A Consensus of Current Knowledge.
Ecological Monographs, 75, 3–35.

Isaac, N.J.B., Storch, D. & Carbone, C. (2011). Taxonomic variation in size–density
relationships challenges the notion of energy equivalence. *Biology Letters*, 7, 615–618.

Jacquet, C., Gounand, I. & Altermatt, F. (2020). How pulse disturbances shape size-
abundance pyramids. *Ecology Letters*, 23, 1014–1023.

Khaliq, I., Rixen, C., Zellweger, F., Graham, C.H., Gossner, M.M., McFadden, I.R., *et al.*
(2024). Warming underpins community turnover in temperate freshwater and terrestrial
communities. *Nat Commun*, 15, 1921.

Laakso, J., Löytynoja, K. & Kaitala, V. (2003). Environmental noise and population
dynamics of the ciliated protozoa Tetrahymena thermophila in aquatic microcosms. *Oikos*,
102, 663–671.

Lawton, J.H. & Hassell, M.P. (1981). Asymmetrical competition in insects. *Nature*, 289, 793–795.

Li, W.K.W. (2002). Macroecological patterns of phytoplankton in the northwestern North Atlantic Ocean. *Nature*, 419, 154–157.

Li, W.K.W., Glen Harrison, W. & Head, E.J.H. (2006). Coherent assembly of phytoplankton communities in diverse temperate ocean ecosystems. *Proceedings of the Royal Society B: Biological Sciences*, 273, 1953–1960.

Long, Z.T. & Morin, P.J. (2005). Effects of organism size and community composition on ecosystem functioning. *Ecology Letters*, 8, 1271–1282.

Long, Z.T., Steiner, C.F., Krumins, J.A. & Morin, P.J. (2006). Species Richness and Allometric Scaling Jointly Determine Biomass in Model Aquatic Food Webs. *Journal of Animal Ecology*, 75, 1014–1023.

Malerba, M.E. & Marshall, D.J. (2019). Size-abundance rules? Evolution changes scaling relationships between size, metabolism and demography. *Ecology Letters*, 22, 1407–1416.

Malerba, M.E., Palacios, M.M. & Marshall, D.J. (2018). Do larger individuals cope with resource fluctuations better? An artificial selection approach. *Proceedings of the Royal Society B: Biological Sciences*, 285, 20181347.

Malerba, M.E., White, C.R. & Marshall, D.J. (2017). Phytoplankton size-scaling of net-energy flux across light and biomass gradients. *Ecology*, 98, 3106–3115.

Marshall, D.J., Malerba, M., Lines, T., Sezmis, A.L., Hasan, C.M., Lenski, R.E., *et al.* (2022). Long-term experimental evolution decouples size and production costs in Escherichia coli. *Proceedings of the National Academy of Sciences*, 119, e2200713119.

Maynard, D.S., Miller, Z.R. & Allesina, S. (2020). Predicting coexistence in experimental ecological communities. *Nature Ecology & Evolution*, 4, 91–100.

McGrady-Steed, J., Harris, P.M. & Morin, P.J. (1997). Biodiversity regulates ecosystem predictability. *Nature*, 390, 162–165.

Meehan, T.D., Jetz, W. & Brown, J.H. (2004). Energetic determinants of abundance in winter landbird communities. *Ecology Letters*, 7, 532–537.

Morán, X.A.G., López-Urrutia, Á., Calvo-Díaz, A. & Li, W.K.W. (2010). Increasing importance of small phytoplankton in a warmer ocean. *Global Change Biology*, 16, 1137–1144.

Nee, S., Read, A.F., Greenwood, J.J.D. & Harvey, P.H. (1991). The relationship between abundance and body size in British birds. *Nature*, 351, 312–313.

Niklas, K.J. & Enquist, B.J. (2001). Invariant scaling relationships for interspecific plant biomass production rates and body size. *Proc Natl Acad Sci U S A*, 98, 2922–2927.

Norkko, A., Villnäs, A., Norkko, J., Valanko, S. & Pilditch, C. (2013). Size matters: implications of the loss of large individuals for ecosystem function. *Sci Rep*, 3, 2646.

O'Connor, M.I., Gilbert, B. & Brown, C.J. (2011). Theoretical Predictions for How Temperature Affects the Dynamics of Interacting Herbivores and Plants. *The American Naturalist*, 178, 626–638.

Parain, E.C., Rohr, R.P., Gray, S.M. & Bersier, L.-F. (2019). Increased Temperature Disrupts the Biodiversity–Ecosystem Functioning Relationship. *The American Naturalist*, 193, 227–239.

Pennekamp, F., Griffiths, J.I., Fronhofer, E.A., Garnier, A., Seymour, M., Altermatt, F., *et al.* (2017). Dynamic species classification of microorganisms across time, abiotic and biotic environments-A sliding window approach. *PLoS One*, 12, e0176682.

Pennekamp, F., Pontarp, M., Tabi, A., Altermatt, F., Alther, R., Choffat, Y., *et al.* (2018). Biodiversity increases and decreases ecosystem stability. *Nature*, 563, 109–112.

Pennekamp, F., Schtickzelle, N. & Petchey, O.L. (2015). BEMOVI, software for extracting behavior and morphology from videos, illustrated with analyses of microbes. *Ecology and Evolution*, 5, 2584–2595.

Perkins, D.M., Perna, A., Adrian, R., Cermeño, P., Gaedke, U., Huete-Ortega, M., *et al.* (2019). Energetic equivalence underpins the size structure of tree and phytoplankton communities. *Nat Commun*, 10, 255.

Petchey, O.L., McPhearson, P.T., Casey, T.M. & Morin, P.J. (1999). Environmental warming alters food-web structure and ecosystem function. *Nature*, 402, 69–72.

Pomeranz, J.P.F., Junker, J.R. & Wesner, J.S. (2022). Individual size distributions across North American streams vary with local temperature. *Global Change Biology*, 28, 848–858.

Reuman, D.C., Holt, R.D. & Yvon-Durocher, G. (2014). A metabolic perspective on competition and body size reductions with warming. *J Anim Ecol*, 83, 59–69.

Russo, S.E., Robinson, S.K. & Terborgh, J. (2003). Size-Abundance Relationships in an Amazonian Bird Community: Implications for the Energetic Equivalence Rule. *The American Naturalist*, 161, 267–283.

Saito, V.S., Perkins, D.M. & Kratina, P. (2021). A Metabolic Perspective of Stochastic Community Assembly. *Trends in Ecology & Evolution*, 36, 280–283.

Saleem, M., Fetzer, I., Harms, H. & Chatzinotas, A. (2013). Diversity of protists and bacteria determines predation performance and stability. *ISME J*, 7, 1912–1921.

Schielzeth, H. (2010). Simple means to improve the interpretability of regression coefficients. *Methods in Ecology and Evolution*, 1, 103–113.

Sentis, A., Binzer, A. & Boukal, D.S. (2017). Temperature-size responses alter food chain persistence across environmental gradients. *Ecology Letters*, 20, 852–862.

Seymour, M., Fronhofer, E.A. & Altermatt, F. (2015). Dendritic network structure and dispersal affect temporal dynamics of diversity and species persistence. *Oikos*, 124, 908–916.

Sheridan, J.A. & Bickford, D. (2011). Shrinking body size as an ecological response to climate change. *Nature Clim Change*, 1, 401–406.

Tabi, A., Pennekamp, F., Altermatt, F., Alther, R., Fronhofer, E.A., Horgan, K., *et al.* (2020). Species multidimensional effects explain idiosyncratic responses of communities to environmental change. *Nature Ecology & Evolution*, 4, 1036–1043.

Tabi, A., Petchey, O.L. & Pennekamp, F. (2019). Warming reduces the effects of enrichment on stability and functioning across levels of organisation in an aquatic microbial ecosystem. *Ecology Letters*, 22, 1061–1071.

Tilman, D., HilleRisLambers, J., Harpole, S., Dybzinski, R., Fargione, J., Clark, C., *et al.* (2004). Does Metabolic Theory Apply to Community Ecology? It's a Matter of Scale. *Ecology*, 85, 1797–1799.

Tilman, D., Reich, P.B., Knops, J., Wedin, D., Mielke, T. & Lehman, C. (2001). Diversity and productivity in a long-term grassland experiment. *Science*, 294, 843–845.

Turchin, P. (2003). *Complex Population Dynamics: A Theoretical/Empirical Synthesis*.

Princeton University Press, Princeton, N.J.

Uszko, W., Diehl, S., Englund, G. & Amarasekare, P. (2017). Effects of warming on

predator–prey interactions – a resource-based approach and a theoretical synthesis. *Ecology*

Letters, 20, 513–523.

White, E.P., Ernest, S.K.M., Kerkhoff, A.J. & Enquist, B.J. (2007). Relationships between

body size and abundance in ecology. *Trends in Ecology & Evolution*, 22, 323–330.

White, E.P., Ernest, S.K.M. & Thibault, K.M. (2004). Trade-offs in Community Properties

through Time in a Desert Rodent Community. *The American Naturalist*, 164, 670–676.

612