Optimizing crop clustering to minimize pathogen invasion in agriculture

Yevhen F. Suprunenko[1],* and Christopher A. Gilligan[1],‡

Department of Plant Sciences, University of Cambridge, Downing Street, Cambridge, CB2 3EA, UK
*,‡Corresponding authors: * ys526@cam.ac.uk; ‡ cag1@cam.ac.uk
Yevhen F Suprunenko (0000-0001-5927-7571) - ORCID
Christopher Gilligan (0000-0002-6845-0003) - ORCID

Acknowledgements
The authors are grateful to Dr Stephen Cornell for useful discussions at early stages of this work, and to Dr Alison Scott-Brown for helpful comments on the manuscript.

Funding
C.A.G. acknowledges financial support from the Gates Foundation (INV-010472).

Abstract

Context The initial rate of pathogen invasion in crops is influenced by the spatial clustering of susceptible crops and the characteristics of pathogen dispersal. Previous studies have shown that various degrees of crop clustering can effectively reduce this invasion rate. However, the optimal degrees of clustering that minimize pathogen invasion have not previously been identified.

Objectives This study aims to determine analytically the range of crop clustering that minimizes the initial rate of pathogen invasion.

Methods We studied artificial agricultural landscapes with crop areas arranged in identical square clusters on a regular square lattice. For pathogen dispersal, we used several common dispersal kernels, including Gaussian, negative exponential, and power-law. The optimal degree of clustering, defined by cluster size and separation distance, was calculated using a new analytical approximation for the pathogen invasion rate, which showed strong agreement with computer simulations. Additionally, we analysed a realistic cassava landscape at risk of invasion by cassava brown streak virus.

Results We identified a range of optimal cluster sizes and corresponding separation distances that minimize pathogen invasion rates for various dispersal kernels and landscapes with clusters of crop fields arranged on a regular square lattice. The methods can be extended to other geometrical configurations, such as long narrow fields. Using a cassava landscape as an example, we show how optimal crop clustering strategies can be derived to mitigate the potential invasion of cassava brown streak virus.

Conclusion The methods provides analytical insights that can help farmers and agricultural planners to optimize the spatial structure of agricultural landscapes to minimize initial pathogen invasion rates.

1. Introduction

One of the key factors that determine how quickly a crop pathogen (or pest) invades and spreads across an agricultural landscape is the relationship between the pathogen's dispersal and the spatial distribution of susceptible crops [1–13]. The rate of pathogen invasion can be slowed by adjusting the spatial configuration of host distribution relative to pathogen dispersal [1,5,8,9,11,12,14,15]. This can be achieved by reducing host aggregation [16–21], growing resistant varieties [22], or applying fungicides [7,23].

Two simple measures of pathogen invasion are the infection rate, r, which denotes the early exponential growth rate of the number of infected fields at the onset of an epidemic, and the basic reproduction number R_0. The latter is defined as $R_0 = r/\mu$, where $1/\mu$ represents the average infectious period of infected hosts.

Insights from theoretical [1,5,11,12] and experimental [8,9] studies indicate that pathogen invasion slows, and both r and R_0 decline, as the host landscape becomes more dispersed

(holding all other factors constant). While this might suggest that the minimal values of r and

R_0 would occur when fields of susceptible crops are evenly dispersed, recent theoretical work

[24] has shown that r and R_0 can be minimized in agricultural landscapes with non-uniform

spatial configurations. In these configurations, fields with susceptible crops can be aggregated

into clusters (i.e. clustered) up to a threshold size.

Suprunenko et al. (2025) [24] illustrated this process using artificial landscapes in which a fixed

area of susceptible crops was distributed in clusters on a regular square lattice (figure 1a). As

the total crop area remained constant, the number of clusters ($N_{clusters}$) decreased as cluster

size increased. The highest degree of clustering (i.e. the smallest $N_{clusters}$) associated with

the minimum r can be characterised by a threshold cluster size of width, L_H^*, and separation

distance between clusters, Δ_H^* (cf figure 1b).

Understanding these threshold values, L_H^* and Δ_H^*, would be invaluable for agricultural

planners in minimizing the risk of pathogen (or pest) invasion, given known dispersal

characteristics.

Fig. 1 (a) An illustration of a crop area, A_H (green), aggregated into one cluster

($N_{clusters} = 1$) or nine clusters ($N_{clusters} = 3 \times 3$), placed on a regular square grid

within an agricultural landscape, A, where arbitrarily $A_H A^{-1} = 0.15$. The degree of

clustering (aggregation) is characterised by the number of clusters, $N_{clusters}$, which

determines the size (L_H) of a square cluster and the separation distance (Δ_H) between

clusters. **(b)** Open circles indicate the infection rate r (the initial growth rate of the

number of infected fields) estimated from simulations of an individual-based epidemic

model initiated by a single randomly infected field within the landscapes of the type

shown in Figure 1a ([24]; Supplementary Note 1). Reconfiguring the crop area A_H into
clusters of size L_H smaller than the threshold size L_H^* ($L_H < L_H^*$) reduces the infection
rate from its maximum value (corresponding to $N_{clusters} = 1$) to its minimum value
(corresponding to $N_{clusters} \geq N_{clusters}^*$, where $N_{clusters}^*$ is the threshold number of
clusters). Therefore, the infection rate of susceptible agricultural crop fields at the
start of an epidemic can be minimized through various spatial reconfigurations of the
landscape. The illustration is a modified version of figure 1 in Ref. [24]. Computer
codes for all figures in the current paper are available from Figshare [25].

Computer simulation of epidemic models on simulated landscapes can, in principle, be used
to estimate the threshold size L_H^* and separation distance Δ_H^* for a given pathogen dispersal
kernel [15,20,21,26,27]. This involves analysing epidemics on host landscapes with many
different spatial configurations to identify the degrees of host clustering that minimize r (or
R_0). However, constructing realistic epidemic spread models for each new pest or pathogen
can be challenging, and considering numerous spatial configurations is computationally
demanding and potentially inefficient.

Here, we explore an alternative approach to identify landscapes that minimise the initial rates
of pathogen invasion using an analytical approximation for the initial infection rate, r. We test
this analytical approximation against empirical infection rates derived from computer
simulations of an individual-based model (IBM) for pathogen invasion and spread through a
host landscape. Using the analytical approximation for r, we derive values for the threshold
size L_H^* and separation distance Δ_H^* for a range of generic host landscapes and commonly-
used dispersal kernels, including Gaussian, negative exponential, and power-law kernels,
among others. We illustrate a potential application of this approach, motivated by the spread
of cassava brown streak virus (CBSV), which threatens cassava production in sub-Saharan
Africa [28–30]. Using a dispersal kernel for CBSV estimated by Godding *et al.* [31] and a section
of a cassava landscape provided by Szyniszewska (2020) [32], we present examples of maps
showing estimates of local threshold sizes and separation distances for clusters of cassava
fields that minimise infection rate.

2. Methods

2.1 Mathematical approximation for infection rate

We consider a susceptible-infected (SI) epidemiological compartmental individual-based
model (IBM) for the invasion and spread of a pathogen through an agricultural landscape
partially occupied by susceptible host crops. An analytical expression for the infection rate, r,
for this IBM has been variously derived in several studies, including those by Bolker (1999)[1],
North and Godfray (2017)[12], van den Bosch et al. (2024)[33] and others [11,24,34,35]. For
example, consider the approximation from Suprunenko et al. (2025)[24]:

$$r = \beta \times \left(n_S + \frac{n_S}{n^2} \int_0^{\frac{\sqrt{A}}{2}} 2\pi x b(x) g(x) \mathrm{d}x \right), \qquad (2.1)$$

where $b(x)$ is the pathogen dispersal kernel as a function of distance x, and β is the infection rate per contact density (i.e. the spatial density of susceptible hosts that an initial infected field can reach through the dispersal kernel $b(x)$). The host landscape is assumed to be a square plot of land with area A partially occupied by land containing a susceptible crop consisting of N sufficiently small, identical individual fields with area A_0. Each field is treated as an individual 'host', described as a point-like object located at the centre of a field. The initial densities $n_S = (N-1)A^{-1}$ and $n = NA^{-1}$ are the densities of susceptible hosts and all hosts respectively. The total area occupied by hosts is A_H, i.e. $A_H = A_0 N$, where $A_H \leq A$. The function, $g(x)$, denotes the second order spatial cumulant (also known as a truncated correlation function or autocovariance, see e.g. [36,37], that characterises the spatial structure of a host landscape.

The expression inside the brackets in equation (2.1) represents the spatial density of contacts with susceptible hosts (i.e. fields) from a randomly located initial infected field. Similarly, van den Bosch et al. (2024)[33] described an equivalent product, $b(x)g(x)$ (cf equation 2.1), as one that "describes how a pathogen 'sees' the host population around an infected host individual" [33]. Based on these insights, we construct an explicit expression for the spatial density of contacts to replace the expression in brackets in equation (2.1).

Our approach differs from conventional IBMs, which consider individual hosts as point-like entities and calculate the infection rate (2.1) based on their spatial distribution. In contrast, we explicitly consider individual hosts (i.e. fields) as square plots of cropped land with area A_0, and estimate the infection rate using the spatial distribution of these land polygons. We denote the spatial density of fields (measured as the number of fields per area) at location $\boldsymbol{x}$ as $n(\boldsymbol{x})$, defining $n(\boldsymbol{x}) = A_0^{-1}$ when a point $\boldsymbol{x}$ lies within a field, and $n(\boldsymbol{x}) = 0$ when $\boldsymbol{x}$ is outside a field. A pathogen at location $\boldsymbol{x}$ can infect directly the density of fields $n_b(\boldsymbol{x})$, where

$$n_b(\boldsymbol{x}) = \int b(|\boldsymbol{x_0} - \boldsymbol{x}|) n(\boldsymbol{x_0}) \mathrm{d}\boldsymbol{x_0},$$

and the integration is over the entire landscape. Averaging $n_b(\boldsymbol{x})$ over all possible locations, $\boldsymbol{x}$, within the landscape leads to the quantity $\bar{n}$,

$$\bar{n} = \left(\int n_b(\boldsymbol{x}) n(\boldsymbol{x}) d\boldsymbol{x} \right) / \left(\int n(\boldsymbol{y}) \mathrm{d}\boldsymbol{y} \right),$$

that can be interpreted as the spatially averaged density of contacts (with susceptible fields) that can result in infection from an initial randomly infected field. Therefore, the expression in brackets in equation (2.1) can be estimated by the density $\bar{n}$, i.e.

$$r = \beta \times \bar{n}. \qquad (2.2)$$

Further to this, we make the following two simplifying approximations.

Approximation 1: Here, we develop an approximation for equation (2.2) when there is a high
degree of aggregation of the target crop within a landscape (i.e. when $N_{clusters}$ is small): $r \approx$
$\beta \times \bar{n}_1(L_H)$, where $\bar{n}_1(L_H)$ is derived as follows. We assume that the crop area is distributed
into identical $L_H \times L_H$ square clusters on a regular square lattice, as illustrated in figure 1a.
The total number of clusters is $N_{clusters}$. We denote the spatial density of crops within a single
cluster as $n_1(x)$, where $n_1(x) = A_0^{-1}$ if $0 \le x_1 \le L_H$ and $0 \le x_2 \le L_H$ (where x_1 and x_2 are
the coordinates in a two-dimensional Cartesian coordinate system), and $n_1(x) = 0$
otherwise. Given a high degree of clustering, i.e. when all hosts are aggregated into a small
number of large clusters, we estimate r on a single cluster and use $n_1(x)$ instead of $n(x)$ in
the expression (2.2), obtaining $\bar{n}_1$ instead of $\bar{n}$, where

155
$$\bar{n}_1 = \left(\int n_b(x) n_1(x) dx \right) \bigg/ \left(\int n_1(y) dy \right).$$

153 Using straightforward algebraic transformations, shown in the electronic supplementary
154 material (Supplementary Note 2), the expression for $\bar{n}_1$ is simplified further:

$$\bar{n}_1(L_H) = 4 L_H^2 A_0^{-1} \int_0^1 \int_0^1 dx dy \, b(L_H \sqrt{x^2 + y^2})(1 - x)(1 - y). \tag{2.3}$$

The quantity $\bar{n}_1$ denotes the averaged density of fields within a single cluster that a randomly
introduced infected field within that cluster can infect directly via a dispersal kernel $b(x)$. The
upper bound of the quantity $\bar{n}_1$ is A_0^{-1}, i.e. the density of crops within a single cluster.

Approximation 2: Here, we develop an approximation for equation (2.2) when there is a low
degree of clustering in the host landscape: $r \approx \beta \times n$. We assume that the pathogen dispersal
kernel, $b(x)$, is sufficiently long-ranged relative to an individual field of area A_0, i.e. the
characteristic dispersal length, a, should be larger than the size of an individual field ($a >$
$\sqrt{A_0}$). Consequently, the minimum value of the infection rate can be approximated as $r_{min} =$
$\beta \times n$, where n is the spatial density of all N fields of a host crop within the landscape with
area A, $n = NA^{-1}$. For a low degree of clustering, we therefore use $r \approx r_{min}$.

Finally, we combine expressions for the infection rate within a single cluster, $\beta \times \bar{n}_1(L_H)$
(approximation 1), and its minimal value, $\beta \times n$ (approximation 2) noting that the larger
estimate should be used between approximations 1 and 2. Indeed, when the crop is highly
aggregated within a landscape, the estimate of r is given by $\beta \bar{n}_1(L_H)$ where $\beta \bar{n}_1(L_H) > \beta n$;
while for a low degree of crop aggregation the estimate of r is given by βn when $\beta n >$
$\beta \bar{n}_1(L_H)$. Therefore, using the maximum of two estimates, we use the following
approximation for r, denoted as $r_{approximate}$:

$$r_{approximate} = \beta \times \max[\bar{n}_1(L_H), n]. \tag{2.4}$$

173 It is important to stress that this approximation is applicable to dispersal kernels, $b(x)$, that
174 are decreasing functions of distance x. For dispersal kernels that do not decrease
175 monotonically with distance, such as inverse Gaussian (Wald) or lognormal dispersal kernels
176 (see [38]) the infection rate does not necessarily decrease as the number of clusters
177 ($N_{clusters}$) increases.

2.2 Computer simulations of an individual-based model

To test the approximation derived above, we compare the performance of equation (2.4) with
empirical rates for r at the start of an epidemic, originally derived from computer simulations
of an individual-based model of pathogen invasion in an agricultural landscape [24]. For
convenience, details of the individual-based model are summarised in Supplementary Note
1. Exploratory testing indicated that the transition of r from its maximum to its minimum
value captured by $r_{approximate}$ (equation 2.4) sufficiently well (see caption of figure 2);
therefore, we proceed to use this approximation.

In figure 2 we used the data [39] from computer simulations of individual-based model of
pathogen invasion and spread in agricultural landscapes that were taken from [24]. Here we
test a different analytical approximation from [24] for identifying optimal clustering to reduce
initial invasion rate, inspired by the original more computationally intensive method in [24]
that uses spatial cumulants to assess the impact of general landscape structures.

Fig. 2 The behaviour of the analytically approximated infection rate, $r_{approximate}$,
(lines corresponding to equation 2.4) as a function of the degree of clustering of
susceptible crops is compared with estimates of infection rate r (symbols) inferred
from computer simulations of an individual-based model (see Methods). Results are
shown for different landscapes and for two different dispersal kernels given by a
Gaussian function with standard deviation: **(a)** $\sigma = 10$ km, and **(b)** $\sigma = 1$ km. The
quantity $A_H A^{-1}$ denotes the fraction of the landscape occupied by susceptible fields;
$n = N A^{-1}$ is the spatial density of all N crop fields in the landscape with area A. Here,
$\beta = 1$, and the area of a unit field is $A_0 = 0.04$ km²; the upper bound of the rate r is
given by $\beta A_0^{-1} = 25$. The transition of r from its maximum to minimum is captured
by $r_{approximate}$ reasonably well, as quantified by the following measures. In both

panels, the maximum deviation $\max|r - r_{approximate}|$ is 0.6, 0.9, 1.3 and 2.0 for

$A_H/A = 1, 15, 30$ and 60, respectively. These deviations correspond to 3.7%, 4.5%,

8.2% and 22% of the total range $\max(r) - \min(r)$ for the same values of A_H/A.

2.3 Mathematical approximation for threshold size and separation distance

207 We define the threshold size L_H^* as the value of the argument of the function $\bar{n}_1(L_H)$ in

208 equation (2.4) such that

$$\bar{n}_1(L_H^*) = n. \qquad (2.5)$$

209 Using the definition $n = NA^{-1} = A_H(AA_0)^{-1}$, the condition (2.5) can be re-written as

$$\bar{n}_1(L_H^*)A_0 = A_H A^{-1}. \qquad (2.6)$$

The right-hand side of equation (2.6) is a dimensionless parameter $A_H A^{-1}$ denoting the

fraction of the landscape occupied by susceptible fields. For the most typical dispersal kernels,

$b(x)$, characterized by a scale parameter a (see Supplementary Note 3, Table S1), the left-

hand side of equation (2.6) depends on the dimensionless parameter $L_H^* a^{-1}$ explicitly (see

electronic supplementary material, Supplementary Note 4 for details). Therefore, for a

landscape characterized by $A_H A^{-1}$, the equation (2.6) determines L_H^* in units of the scale

parameter a that characterises the pathogen dispersal kernel $b(x)$. The threshold separation

distance Δ_H^* is then determined (on a regular square lattice, figure 1a) by L_H^* and $A_H A^{-1}$:

$$\Delta_H^* = L_H^* \times \left(\sqrt{A/A_H} - 1\right). \qquad (2.7)$$

218 Parameters and functions used in the current paper are summarized in Table 1.

219 **Table 1** Parameters used in the model, together with characteristic lengths of pathogen

220 dispersal and host landscape and their threshold values.

Symbol	Description
Model parameters	
A	Entire area of the landscape, a square plot of land.
A_0	Area occupied by a single field of host crop (a square plot of cropped land).
N	Number of host crop fields within the landscape.
A_H	Total area occupied by fields (host crop), $A_H = A_0 \times N; \ A_H \leq A$.
n	Density of all host crop fields within a landscape, $n = N/A$.
n_S	Density of susceptible fields within a landscape, assuming a single initial infected field, $n_S = (N - 1)/A$.
$b(x)$	Pathogen dispersal kernel.
σ	Standard deviation of a Gaussian function.
β	Infection rate per contact density.
r	Rate of infection of susceptible fields at the start of an epidemic, equation (2.1).
Spatial scales and degree of aggregation of a host landscape	

a, b_0	Scale parameter (a) and a shape parameter (b_0) of a pathogen dispersal kernel, see Supplementary Material, Supplementary Note 3, Table S1.
L_H, Δ_H	Cluster size (L_H) and separation distance (Δ_H) characterising a host crop landscape, figure 1a.
$N_{clusters}$	Number of clusters of cropped fields within the landscape, $N_{clusters} = A_H/L_H^2 = A/(L_H + \Delta_H)^2$. It can be seen as a measure of the degree of aggregation of a host landscape.
Threshold values	
L_H^*, Δ_H^* $N_{clusters}^*$	Threshold values of cluster size (L_H^*) and separation distance (Δ_H^*), figure 1c, expressed in units of the scale parameter a and calculated from equations (2.6) and (2.7). $N_{clusters}^*$ is the corresponding threshold value of the number of clusters.

2.4 Cassava landscape and CBSV

We illustrate the analytical approach with a real-life example: the invasion of CBSV in a
cassava landscape. Godding et al. (2023)[31] developed and parameterised an epidemic
spread model for CBSV with a pathogen dispersal kernel defined on a raster with a 1 km spatial
resolution, imposed by the spatial resolution of available host distribution data [32]. The CBSV
dispersal, determined in [31] by β and a function $K(d_{ij})$ of distance d_{ij} between centroids of
raster cells, accounts for the combined effect of short-distance CBSV spread via insect vectors
and long-distance spread via movement of virus-infected planting materials [31]. Specifically,
$K(d_{ij} = 0) = p$ and $K(d_{ij} > 0) = Cd_{ij}^{-\alpha}$, where p is the proportion of inoculum remaining
in the source cell, C is a normalization constant determined by $\sum_j K(d_{ij}) = 1$, and α is the
kernel exponent. By fitting this model to surveillance data, Godding and co-authors [31]
derived posterior distributions for the parameters β, p and α. Here, we use a radially
symmetric, staircase-like version of the raster-based dispersal kernel from [31]. Using their
results, we selected a power-law dispersal kernel $b(x)$, characterised by two parameters: the
exponent $\alpha = 3.75$ and $p = 0.12$ taken from a region of relatively high probability density in
posterior distributions estimated by Godding et al. (2023) [31].

In common with [31], we use CassavaMap [32] as a source of mapped landscape data for
cassava. The landscape data for cassava are derived [31] from production statistics [32] and
resolved to 1 km^2 raster cells across sub-Saharan countries that cultivate cassava. The
analytical estimates (2.6)-(2.7) depend on the ratio of host crop area A_H to the total area A.
We express the landscape in terms of hectares of cassava fields per raster cell, noting that,
according to [32], no more than half of the area of each raster cell (approximately 50 hectares)
can be allocated to cassava production. Therefore, the maximum value of $A_H A^{-1}$ in the
cassava landscape data cannot exceed 0.5, i.e. here $A_H A^{-1} \leq 0.5$.

Finally, using the approach described above, we calculated L_H^* and Δ_H^* (figure 4a-b, black
curves) for a cassava landscape at risk of CBSV invasion.

3. Results

Using the approximation for the rate r (equation 2.4, figure 2), we identified a range of values of cluster size L_H for which the infection rate, r, is minimized. Within this range, the largest cluster size, L_H, is referred to as a threshold cluster size L_H^*. Equation (2.6) determines the ratio L_H^*/a of the threshold cluster size L_H^* to the pathogen dispersal scale parameter a for a given pathogen dispersal kernel $b(x)$ and the fraction $A_H A^{-1}$ of the landscape occupied by susceptible crops. Correspondingly, the threshold separation distance is calculated using equation (2.7).

3.1 Artificial landscapes and commonly-used dispersal kernels

Applying this approach to different dispersal kernels and landscapes, we first considered the Gaussian, negative exponential and power-law kernels – commonly used in epidemic spread studies [13,40,41] – alongside several additional kernels detailed in Supplementary Note 3, Table S1. The results for the exponential and power-law kernels are summarised in figure 3, while the results for the other kernels from Table S1 are presented in the electronic supplementary material, Supplementary Note 4.

In addition to the numerical solution for L_H^* from equation (2.6), we derived an explicit analytical estimate for L_H^* when the value of $A_H A^{-1}$ is small. In this case, the value of L_H^* is also small. Therefore, using the approximation $b\left(L_H^*\sqrt{x^2 + y^2}\right) \approx b(0)$ in equation (2.6) we derived:

$$L_H^* \approx \sqrt{\frac{A_H}{b(0)A}}, \qquad \text{when} \quad \frac{A_H}{A} \ll 1. \tag{3.1}$$

The explicit analytical estimate (equation 3.1) depends only on the dispersal kernel at zero distance, $b(0)$, and ignores its distance-dependent component, $b(x > 0)$. Consequently, the explicit analytical estimate of L_H^* aligns better with the numerical solution for L_H^* from equations (2.6–2.7) when short-range dispersal dominates, as with Gaussian kernels (figure 3a), but less so for long-tailed kernels like the power-law (figure 3b). This is reflected in the closer agreement between the thin black (for the numerical solution, equations 2.6–2.7) and the thick gray (explicit analytical estimate, equation 3.1) lines in figure 3a than in figure 3b. As the ratio A_H/A increases, clusters move closer together, amplifying long-distance dispersal effects and reducing the accuracy of explicit analytical estimate (3.1), as seen in both panels.

Fig. 3 Threshold cluster size L_H^* and threshold separation distance Δ_H^* identified for two types of dispersal kernel. **(a)** Results are shown for exponential power kernels, that correspond to the Gaussian kernel when $b_0 = 2$ and the negative exponential kernel when $b_0 = 1$, as shown in the inset of the panel. **(b)** Results are shown for inverse power-law kernels with exponents $b_0 = 3$ and $b_0 = 4$, as illustrated in the inset of the panel. For the exact analytical expressions for the dispersal kernels used here see Table S1 in Supplementary Note 3. In both panels, black curves represent numerical estimates from equations (2.6) and (2.7), while grey curves at $A_H A^{-1} \leq 0.1$ represent explicit analytical estimates from equations (3.1) and (2.7). The R and Mathematica codes used to calculate these results are available from Figshare [25] and can be easily applied to other dispersal kernels.

We present an example of how results shown in figure 3 could inform stakeholders, including farmers and other agricultural planners, about the spatial structure of a host landscape. To keep the disease from spreading quickly at the start of an epidemic, it is important than no cluster of host crop fields is too large. Specifically, these clusters should not exceed a certain critical size (equal to $L_H^* \times L_H^*$). In addition, each cluster should be surrounded by a buffer zone – a strip of land of specific width (equal to $\Delta_H^*/2$) without susceptible host crops. Smaller clusters help limit disease spread within them, while buffer zones reduce the chance of transmission between clusters.

The same principle applies to clusters smaller than the critical size: any $L_H \times L_H$ clusters (where $L_H < L_H^*$) should be surrounded by a host-free buffer zone of width $\Delta_H/2$, where $\Delta_H = L_H\left(\sqrt{A/A_H} - 1\right)$ (cf equation (2.7)). The buffer zones around different clusters should not overlap. If there are two clusters within the landscape, first cluster $L_H^{(1)} \times L_H^{(1)}$ and second cluster $L_H^{(2)} \times L_H^{(2)}$, where both sizes are below the threshold size, i.e. $L_H^{(1),(2)} \leq L_H^*$, then the

minimal suggested separation distance between these clusters equals the sum of the two
corresponding buffer zones, i.e. $(\Delta_H^{(1)} + \Delta_H^{(2)})/2$.

Although we considered the spatial distribution of the cropped area into square clusters on a
regular square lattice, other geometrical configurations could be considered. For example, if
the cropped area is distributed among long narrow fields (e.g. see [8]), we present the
derivation of an explicit analytical estimate for L_H^* in Supplementary Note 5, analogous to
equation (3.1).

3.2 Cassava landscape and an example of CBSV dispersal kernel

To demonstrate the application of the analytical approach to CBSV invasion in a cassava
landscape, we used an area $A = 24 \times 24$ km^2 in an arbitrarily selected region on the border
between Cameroon and the Central African Republic (figure 4c) derived from [32]. We
calculated that cassava fields occupy 1.1% of this area (i.e. $A_H A^{-1} = 0.011$). According to
figures 4a and 4b, the corresponding threshold cluster size and separation distances are $L_H^* =$
0.27 km and $\Delta_H^* = 2.29$ km. Therefore, if it were logistically possible, our analysis suggests
that organizing the crop area from figure 4c into clusters of 0.27 km by 0.27 km ($L_H^* \times L_H^*$),
separated by 2.29 km (Δ_H^*), would result in the minimal infection rate, $r_{min} = \beta n$.

Fig. 4 Threshold cluster size L_H^* and threshold separation distance Δ_H^* identified for an example of CBSV dispersal kernel and a cassava landscape. **(a)-(b)** Threshold cluster size L_H^* and separation distance Δ_H^* that minimise the initial infection rate r as the fraction A_H/A of the landscape occupied by the host crop increases. Black curves represent numerical estimates obtained from equations (2.6) and (2.7), thick grey curves represent explicit analytical estimates from equations (3.1) and (2.7). **(c)** The cassava landscape, given at 1 km spatial resolution. **(d)** The same cassava landscape at 4 km spatial resolution, shown in terms of the fraction A_H/A of area occupied by the host crop. **(e)-(f)** Maps of L_H^* and Δ_H^* calculated using information from panels (a), (b) and (d), showing how the crop area in each cell can be maximally clustered for the infection rate to remain minimal ($r_{min} = \beta n$, with the value of n determined separately for each cell).

However, major redistributions of cropping patterns are unlikely to be practically feasible. Therefore, we considered a more practical approach by focusing on smaller 4 km by 4 km

local areas where spatial reconfigurations of cropping patterns might potentially be possible.
However, we emphasize that this is a hypothetical case to support the concept, as, in practice,
there are many other major factors to be considered. We aggregated the original landscape
to a 4 km spatial resolution (figure 4d). Using the fraction of area occupied by the host crop
in each 4 km by 4 km cell, we assigned each cell a corresponding value of the threshold cluster
size L_H^* (figure 4e) and separation distance Δ_H^* (figure 4f).

For illustration, consider the cell in figure 4d-f located between 4 km and 8 km on the x axis,
and between 12 km and 16 km on the y axis. Our analysis indicates that the infection rate
would remain minimal if cassava fields were aggregated into a cluster approximately 1 km by
1 km in size, surrounded by a cassava-free buffer zone with a width of $\Delta_H^*/2 \approx 900$ m. In
another cell, located between 16 km and 20 km on the x axis and between 12 km and 16 m
on the y axis, the infection rate would remain minimal if cassava fields were aggregated into
a cluster of 500 m by 500 m, surrounded by a cassava-free buffer zone with a width of $\Delta_H^*/2 \approx$
1 km.

Maps from figures 4e-f could in principle be used to inform decisions on where to plant new
cassava fields without increasing the risk of early CBSV epidemic spread. If a cluster of size
$L_H^* \times L_H^*$ already exists within a given 4 km by 4 km cell, new fields should not be placed within
the recommended cassava-free buffer zone (with a width of $\Delta_H^*/2$) around the cluster to avoid
increasing the infection rate. New fields can be planted near smaller clusters, but each smaller
cluster of size $L_H \leq L_H^*$ should be surrounded by a cassava-free buffer zone with a width of
$\Delta_H/2$, where $\Delta_H = L_H\left(\sqrt{A/A_H} - 1\right)$.

4. Discussion

To identify the range of optimal clustering of host crops that minimizes the pathogen invasion
rate, the analytical approach presented in this paper integrates the intrinsic spatial
characteristics of pathogen dispersal with the distribution of host crops. Considering a specific
geometrical configuration of host landscapes, where susceptible crops are distributed in
square clusters on a regular square lattice, the resulting threshold characteristics, cluster size
L_H^* and separation distance Δ_H^*, are derived in units of the scale parameter, a, of a given
pathogen dispersal kernel and depend on the fraction A_H/A of the landscape occupied by
susceptible crops. Consequently, this analytical approach is effective for a wide range of
dispersal kernels and host landscapes.

We used two main approximations in deriving the analytical results. The first approximation
was applied to landscapes where hosts were aggregated into a *small number* of clusters (i.e.
where r is a decreasing function of the number of clusters; see e.g. figure 2). In those cases,
we estimated r from a single cluster and the corresponding density $\bar{n}_1$ (that is the spatially
averaged density of contacts with susceptible crops within a single cluster that can result in
infection from an initial randomly infected host), $r \approx \beta \bar{n}_1$, (approximation 1, equation (2.3)).

In other landscapes, where crops were aggregated into a *large number* of clusters (i.e. where
r depends weakly on the number of clusters, see e.g. figure 2), we used a second

approximation and estimated r assuming a constant density, n (that is the density of crops
within the entire landscape and therefore independent of the number of clusters), $r \approx \beta n$,
(approximation 2).

Approximation 1 ($r \approx \beta \bar{n}_1$) applies when clusters are sufficiently large and well-isolated, such
that inter-cluster contacts are minimal. In this case, approximation 1 effectively considers
each single cluster in isolation, neglecting inter-cluster contacts and therefore
underestimating the actual value of r. In contrast, approximation 2 ($r \approx \beta n$) applies when
numerous sufficiently small clusters are located close to each other. In this case, contacts may
occur across many clusters, except the most distant ones. However, because approximation
2 assumes contacts with all hosts across the entire landscape, it overestimates the actual
value of r. The intersection of the two approximations determines the threshold cluster size
L_H^*, i.e. $\beta \bar{n}_1(L_H^*) = \beta n$ (equation 2.5), which is solved numerically to provide an exact value
for L_H^*. The underestimation (due to approximation 1) of the left-hand side of equation 2.5
and the overestimation on the right-hand side (due to approximation 2) lead to an
underestimation at the intersection point for the thresholds number of clusters, $N_{clusters}^*$
(figure 1c), and consequently, to an overestimation of L_H^* due to the relationship $L_H^* = $
$\sqrt{A_H/N_{clusters}^*}$. The resulting overestimation of L_H^* provides a practical upper bound to the
size of a cluster of crop fields that minimizes the risk of rapid spread during the initial stages
of epidemics.

By using methods based on r (or R_0 given the link in SIR models as $R_0 = r/\mu$ [24]), our analysis
focuses on short-term epidemiological dynamics. The analysis is therefore useful for
predicting outcomes during the initial invasion but does not necessarily provide an accurate
assessment of the long-term dynamics of an epidemic. However, in some cases, medium-term
dynamics can still be characterised by r. For example, in our earlier work [24] we
demonstrated that "the analytical estimates of the infection rate correctly identified the
relative ranking of the number of infected fields" [24] for different landscapes in sub-Saharan
Africa, using a six-month spread period as a proxy for medium-term dynamics. In general, the
nonlinear dynamics, typical of many epidemics, mean that initial fast or slow rates of epidemic
progress are often not sustained throughout the epidemic. For example, Brown and Bolker
[5] and Benincà et al. [42] have shown how initially high values of r or R_0 can result in a slow
spread of pathogens at later stages of an epidemic, where there are trade-offs between local
and long-range dispersal. Therefore, to generate robust predictions for long-term
epidemiological dynamics in realistic agricultural landscapes, it is necessary to use computer
simulations of epidemic spread models. For instance, Godding et al. [43] modelled the 25-year
spread of CBSV across sub-Saharan Africa, providing insights that go beyond what short-term
analytical methods can offer.

The results of the current paper improve our understanding of the relationship between
spatial scales of host distribution and pathogen dispersal, and their role in epidemic dynamics
[7,44]. Specifically, the analytical results in figure 3 provide some general insights into the
roles of the spatial scale parameter a and the shape parameter b_0 (cf Table S1 in
Supplementary Note 3) of a dispersal kernel in epidemics. These insights can be used to

quickly compare how different dispersal kernels affect the invasion speed of pathogens in the
same host landscape. This opens the door to future work on identifying optimal host
distributions under invasion by multiple pathogens with different dispersal kernels. In
addition, dispersal kernels are often poorly known, which makes the analytical approach
presented here particularly valuable. It enables rapid exploration of a broad family of
potential kernels – even before detailed empirical estimates are available.

For example, from equation (2.6) and its dependence on the dimensionless parameter L_H^*/a,
it follows that increasing the spatial scale parameter a (i.e., increasing dispersal length)
reduces the infection rate at the start of an epidemic towards its minimal value βn (see
Supplementary Note 6 for the details). Considering the role of the parameter b_0, figure 3a
shows that the values of L_H^* for the negative exponential kernel ($b_0 = 1$) are larger than for
Gaussian kernel ($b_0 = 2$). This result indicates that, in the same host landscape, the infection
rate r at the start of an epidemic will be higher for the Gaussian kernel than for the negative
exponential kernel, assuming β is constant.

For ease of presentation, we have focused on pathogens, but the method can equally be
applied to pest invasions where dispersal kernels are known or can be estimated. Ultimately,
the analytical approach of this work has the potential to inform agronomic planning decisions
[45,46] to optimise the size and spacing between clusters of fields, thereby minimizing the
initial infection rate of an invading pathogen. The results of this paper provide valuable
insights into selecting methods to decelerate the initial spread of an invading pathogen across
a landscape of susceptible crops. Future work could further develop this analytical approach
to encompass the role of characteristic spatial scales associated with control measures [7,23]
on initial epidemic dynamics.

Data availability

436 The datasets analysed during the current study were sourced from the previously published
437 work by Suprunenko *et al.* [39] and are available in the Figshare repository,
438 https://doi.org/10.6084/m9.figshare.25804702

References

1. Bolker, B. Analytic models for the patchy spread of plant disease. *Bulletin of Mathematical*

Biology **61**, 849–874 (1999).

2. Park, A. W., Gubbins, S. & Gilligan, C. A. Invasion and persistence of plant parasites in a spatially

structured host population. *Oikos* **94**, 162–174 (2001).

3. Gilligan, C. A. An Epidemiological Framework for Disease Management. *Advances in Botanical*

Research **38**, 1–64 (2002).

4. Stacey, A. J., Truscott, J. E., Asher, M. J. C. & Gilligan, C. A. A Model for the Invasion and Spread

of Rhizomania in the United Kingdom: Implications for Disease Control Strategies.

Phytopathology **94**, 209–215 (2004).

5. Brown, D. & Bolker, B. M. The effects of disease dispersal and host clustering on the epidemic

threshold in plants. *Bulletin of Mathematical Biology* **66**, 341–371 (2004).

6. Plantegenest, M., Le May, C. & Fabre, F. Landscape epidemiology of plant diseases. *J. R. Soc.*

Interface **4**, 963–972 (2007).

7. Gilligan, C. A., Truscott, J. E. & Stacey, A. J. Impact of scale on the effectiveness of disease

control strategies for epidemics with cryptic infection in a dynamical landscape: an example for

a crop disease. *J. R. Soc. Interface.* **4**, 925–934 (2007).

8. Sackett, K. E. & Mundt, C. C. Effect of plot geometry on epidemic velocity of wheat yellow rust.

Plant Pathology **58**, 370–377 (2009).

9. Mundt, C. C., Sackett, K. E. & Wallace, L. D. Landscape heterogeneity and disease spread:

experimental approaches with a plant pathogen. *Ecological Applications* **21**, 321–328 (2011).

10. Cunniffe, N. J. *et al.* Thirteen challenges in modelling plant diseases. *Epidemics* **10**, 6–10 (2015).

11. Mikaberidze, A., Mundt, C. C. & Bonhoeffer, S. Invasiveness of plant pathogens depends on the

spatial scale of host distribution. *Ecological Applications* **26**, 1238–1248 (2016).

12. North, A. R. & Godfray, H. C. J. The dynamics of disease in a metapopulation: The role of

dispersal range. *Journal of Theoretical Biology* **418**, 57–65 (2017).

13. Xing, Y. *et al.* Global Cropland Connectivity: A Risk Factor for Invasion and Saturation by

Emerging Pathogens and Pests. *BioScience* **70**, 744–758 (2020).

14. Lustig, A., Stouffer, D. B., Doscher, C. & Worner, S. P. Landscape metrics as a framework to

measure the effect of landscape structure on the spread of invasive insect species. *Landscape*

Ecol **32**, 2311–2325 (2017).

15. Etienne, L., Franck, P., Rusch, A. & Lavigne, C. Apple pest and pathogen reduction in landscapes

with large patch size and small area of orchards: a national-scale analysis. *Landsc Ecol* **39**, (2024).

16. Mundt, C. C. Effect of Host Genotype Unit Area on Development of Focal Epidemics of Bean Rust

and Common Maize Rust in Mixtures of Resistant and Susceptible Plants. *Phytopathology* **76**,

895–900 (1986).

17. Mundt, C. C. & Leonard, K. J. Analysis of factors affecting disease increase and spread in

mixtures of immune and susceptible plants in computer-simulated epidemics. *Phytopathology*

76, 832–840 (1986).

18. Skelsey, P., Rossing, W. A. H., Kessel, G. J. T. & Van Der Werf, W. Invasion of *Phytophthora*

infestans at the Landscape Level: How Do Spatial Scale and Weather Modulate the

Consequences of Spatial Heterogeneity in Host Resistance? *Phytopathology®* **100**, 1146–1161

(2010).

19. Papaïx, J., Touzeau, S., Monod, H. & Lannou, C. Can epidemic control be achieved by altering

landscape connectivity in agricultural systems? *Ecological Modelling* **284**, 35–47 (2014).

20. Beasley, E. M., Aristizábal, N., Bueno, E. M. & White, E. R. Spatially explicit models predict coffee

rust spread in fragmented landscapes. *Landsc Ecol* **37**, 2165–2178 (2022).

21. Précigout, P.-A., Renard, D., Sanner, J., Claessen, D. & Robert, C. Crop mixtures outperform

rotations and landscape mosaics in regulation of two fungal wheat pathogens: a simulation

study. *Landsc Ecol* **38**, 77–97 (2023).

22. Rimbaud, L. *et al.* Models of plant resistance deployment. *Annu. Rev. Phytopathol.* **59**, 125–152

(2021).

23. Meentemeyer, R. K., Haas, S. E. & Václavík, T. Landscape Epidemiology of Emerging Infectious

Diseases in Natural and Human-Altered Ecosystems. *Annu. Rev. Phytopathol.* **50**, 379–402

(2012).

24. Suprunenko, Y. F., Cornell, S. J. & Gilligan, C. A. Predicting the effect of landscape structure on

epidemic invasion using an analytical estimate for infection rate. *R. Soc. Open Sci.* **12**, 240763

(2025).

25. Suprunenko, Y. F. & Gilligan, C. A. Computer code for: Optimizing crop clustering to minimize

pathogen invasion in agriculture. Figshare https://doi.org/10.6084/m9.figshare.26038006

(2025).

26. Cunniffe, N. J., Cobb, R. C., Meentemeyer, R. K., Rizzo, D. M. & Gilligan, C. A. Modeling when,

where, and how to manage a forest epidemic, motivated by sudden oak death in California.

Proc. Natl. Acad. Sci. U.S.A. **113**, 5640–5645 (2016).

27. Nguyen, V.-A., Bartels, D. W. & Gilligan, C. A. Modelling the spread and mitigation of an

emerging vector-borne pathogen: Citrus greening in the U.S. *PLoS Comput Biol* **19**, e1010156

(2023).

28. Nweke, F. I., Lynam, J. K. & Spencer, D. S. C. *The Cassava Transformation: Africa's Best-Kept*

Secret. (Michigan State University Press, 2002).

29. Alicai, T. *et al.* Re-emergence of cassava brown streak disease in Uganda. *Plant Disease* **91**, 24–

29 (2007).

30. Muhindo, H. *et al.* Optimum time for harvesting cassava tubers to reduce losses due to cassava

brown streak disease in northeastern DRC. *JAS* **12**, 70 (2020).

31. Godding, D. *et al.* Developing a predictive model for an emerging epidemic on cassava in sub-

Saharan Africa. *Sci Rep* **13**, 12603 (2023).

32. Szyniszewska, A. M. CassavaMap, a fine-resolution disaggregation of cassava production and

harvested area in Africa in 2014. *Sci Data* **7**, 159 (2020).

33. van den Bosch, F., Helps, J. & Cunniffe, N. J. The basic-reproduction number of infectious

diseases in spatially structured host populations. *Oikos* e10616 (2024) doi:10.1111/oik.10616.

34. Suprunenko, Y. F., Cornell, S. J. & Gilligan, C. A. Analytical approximation for invasion and

endemic thresholds, and the optimal control of epidemics in spatially explicit individual-based

models. *J. R. Soc. Interface.* **18**, rsif.2020.0966, 20200966 (2021).

35. Wadkin, L. E. *et al.* Estimating the reproduction number, R_0, from individual-based models of

tree disease spread. *Ecological Modelling* **489**, 110630 (2024).

36. Ovaskainen, O. *et al.* A general mathematical framework for the analysis of spatiotemporal point

processes. *Theor Ecol* **7**, 101–113 (2014).

37. Cornell, S. J., Suprunenko, Y. F., Finkelshtein, D., Somervuo, P. & Ovaskainen, O. A unified

framework for analysis of individual-based models in ecology and beyond. *Nat Commun* **10**,

4716 (2019).

38. Nathan, R., Klein, E., Robledo-Arnuncio, J. J. & Revilla, E. Dispersal kernels: review. in *Dispersal*

ecology and evolution vol. chapter 15 187–210 (OXFORD UNIV PRESS, 2012).

39. Suprunenko, Y. F., Cornell, S. J. & Gilligan, C. A. Data from: Predicting the effect of landscape

structure on epidemic invasion using an analytical estimate for infection rate. Figshare

https://doi.org/10.6084/m9.figshare.25804702 (2024).

40. Fitt, B. D. L., Gregory, P. H., Todd, A. D., McCartney, H. A. & Macdonald, O. C. Spore Dispersal

and Plant Disease Gradients; a Comparison between two Empirical Models. *Journal of*

Phytopathology **118**, 227–242 (1987).

41. Mundt, C. C., Sackett, K. E., Wallace, L. D., Cowger, C. & Dudley, J. P. Aerial Dispersal and

Multiple-Scale Spread of Epidemic Disease. *EcoHealth* **6**, 546–552 (2009).

42. Benincà, E., Hagenaars, T., Boender, G. J., Van De Kassteele, J. & Van Boven, M. Trade-off

between local transmission and long-range dispersal drives infectious disease outbreak size in

spatially structured populations. *PLoS Comput Biol* **16**, e1008009 (2020).

43. Godding, D. *et al.* Predicting the cross-continental spread of the cassava brown streak disease

epidemic in sub-Saharan Africa. *bioRxiv* 2025.10.10.681618 (2025)

doi:10.1101/2025.10.10.681618.

44. Gilligan, C. A. Sustainable agriculture and plant diseases: an epidemiological perspective. *Phil.*

Trans. R. Soc. B **363**, 741–759 (2008).

45. Jeanneret, Ph. *et al.* Agroecology landscapes. *Landscape Ecol* **36**, 2235–2257 (2021).

46. Vega, D., Ibarra, S., Varela Pardo, R. A. & Poggio, S. L. Agroecological management of crop

diseases: a review. *Agroecology and Sustainable Food Systems* **47**, 919–949 (2023).