Title: Shared PRAME Epitopes are T-Cell Targets in NUT Carcinoma

Running Title: PRAME in NUT Carcinoma

Authors: Jeffrey L. Jensen[††1,2], Sara K. Peterson[††2,3], Maria Sambade[2], Jessica R.

Alley[2], Shawn Yu[4], Tomoaki Kinjo[2,4], Sarah N. Bennett[2,5], Steven P. Vensko II[2], Mitra

Shabrang[6], Johnathan D. DeBetta[7], Julie K. Geyer[2,8], Brandon A. Price[9], Kwangok P.

Nickel[10], Randall J. Kimple[10,11], Rishi S. Kotecha[12,13,14], Laura E. Herring[2], Ian J.

Davis[2,15], Jeremy R. Wang[2,8], Christopher A. French[16], Brian Kuhlman[2,4], Jared

Weiss[xx1,2], Alex Rubinsteyn[xx2,3], Benjamin G. Vincent[*xx5,17]

[††] = co-first authorship

[xx] = co-last authorship

[*] = corresponding author

Corresponding Author

Benjamin G. Vincent, MD

Associate Professor of Medicine, Division of Hematology

Joint Faculty, Department of Microbiology and Immunology

Curriculum in Bioinformatics and Computational Biology

Computational Medicine Program

Lineberger Comprehensive Cancer Center

University of North Carolina at Chapel Hill

5206 Marsico Hall, Chapel Hill, NC 27599

Benjamin_Vincent@med.unc.edu, @BenjaminGVincen

1: Division of Medical Oncology, Department of Medicine, The University of North

Carolina at Chapel Hill, Chapel Hill, North Carolina

2: Lineberger Comprehensive Cancer Center, Chapel Hill, North Carolina

3: Department of Genetics, Curriculum in Bioinformatics and Computational Biology,

The University of North Carolina at Chapel Hill, Chapel Hill, North Carolina

4: Department of Biochemistry and Biophysics, University of North Carolina at Chapel

Hill, Chapel Hill, North Carolina

5: Department of Microbiology and Immunology, The University of North Carolina at

Chapel Hill, Chapel Hill, North Carolina

6: Curriculum in Toxicology and Environmental Medicine, School of Medicine, University

of North Carolina at Chapel Hill, Chapel Hill, North Carolina

7: Department of Biochemistry and Molecular Biology, John Hopkins Bloomberg School

of Public Health, Baltimore, Maryland

8: Department of Pathology and Laboratory Medicine, University of North Carolina at

Chapel Hill, Chapel Hill, North Carolina

9: Tempus AI, Inc., Chicago Illinois

10: Department of Human Oncology, University of Wisconsin School of Medicine and

Public Health, University of Wisconsin, Madison, WI

11: University of Wisconsin Carbone Cancer Center, University of Wisconsin School of

Medicine and Public Health, University of Wisconsin, Madison, WI

12: Department of Clinical Haematology, Oncology, Blood and Marrow Transplantation,

Perth Children's Hospital, Perth, Australia

13: Leukaemia Translational Research Laboratory, WA Kids Cancer Centre, The Kids

Research Institute Australia, University of Western Australia, Perth, Australia

14: Curtin Medical School, Curtin University, Perth, Australia

15: Division of Hematology-Oncology, Department of Pediatrics, The University of North

Carolina at Chapel Hill, Chapel Hill, North Carolina

16: Department of Pathology, Brigham and Women's Hospital, Harvard Medical School,

Boston, Massachusetts

17: Division of Hematology, Department of Medicine, The University of North Carolina at

Chapel Hill, Chapel Hill, North Carolina

Keywords: NUT carcinoma, PRAME, T-Cell Receptor Bispecifics, Cancer-specific

Antigens, brenetafusp, IMC-F106C, BRD4::NUTM1, BRD3::NUTM1, NSD3::NUTM1,

anzutresgene autoleucel

Declarations:

Ethics approval and consent to participate: All genetic engineering experiments involving lentivirus or siRNAs were approved by the UNC Institutional Biosafety Committee. All animal experiments were approved by the UNC Institutional Animal Care and Use Committee. All patient samples analyzed in the Tempus AI Sequencing Database were deidentified, and patients consented for their deidentified data to be used in research.

Consent for publication: All authors consent that contingent on the acceptance thereof, for this research to be published in JITC with all appropriate requirements of copyright and access as this journal demands.

Availability of data and material: All scripts and non-genomic, non-mass spectrometry raw data used to generate the figures are available at GitHub (https://github.com/pirl-unc/PRAME_in_NUT_Carcinoma). RNA sequencing data from the five NUT cell lines and the patient derived xenograft will be available on Gene Expression Omnibus. A CSV file is available in supplementary data which includes gene expression data for cancer/testis genes in the Tempus AI patient cohort. All other sequencing data for this cohort is the property of Tempus AI and will be shared at their discretion. Any other raw data (including mass spectrometry, TMA PRAME staining images) not owned by Tempus AI are available upon request.

Competing interests: The authors declare no competing interests.

Funding: JLJ was supported by a Conquer Cancer Young Investigator Award, paid for

by the New Rhein Foundation. SY was also supported by a Conquer Cancer Young

Investigator Award, paid for by the Fred J. Ansfield, MD Endowed Young Investigator

Award. Any opinions, findings, and conclusions expressed in this material are those of

the author(s) and do not necessarily reflect those of the American Society of Clinical

Oncology®, Conquer Cancer®, Fred J. Ansfield or the New Rhein Foundation. JLJ was

supported additionally by a Lung Cancer Initiative Fellowship Award, the UNC

Department of Medicine Core Facilities CFAC Award, as well as the IM-TAG T32

(T32CA285257). SY was additionally supported by the American Cancer Society

Postdoctoral Fellowship, the UNC School of Medicine Translational Team Science

Award, and the UNC-Duke Collaborative Clinical Pharmacology T32 under NIH Grant

T32GM086330. RJK was supported by P50CA278595, R37 CA255330, and P30

CA014520. This work was also supported by The Max Vincze Foundation (AR and

BGV), The Victor Family Foundation (AR and BGV), philanthropic support from the Ben

Brown Family (JW), and the NIH grant R35GM131923. IJD is supported by the V

Foundation for Cancer Research and the NIH (R01-CA276663).

Authors' contributions:

JLJ – Generated hypotheses, performed experimentation for all figures, wrote the first

draft of the manuscript, revised the manuscript, and obtained funding

SKP – Generated hypotheses, performed sequencing analyses, produced the figures,

wrote the first draft of the manuscript, and revised the manuscript

MaS – Produced western blot samples, produced western blots, and prepared samples

for immunopeptidomics

JRA – Prepared samples for immunopeptidomics, performed bispecific and TCR T-cell

cytotoxicity assays, and edited the manuscript

SY – Designed and produced $\alpha PRAME_{425}$ x SP34 $\alpha CD3$ bispecific, designed and

produced the $\alpha HTLV$ bispecific, performed testing in T2 cells and NSCLC cells, and

revised the manuscript

TK - Designed and produced $\alpha PRAME_{425}$ x SP34 $\alpha CD3$ bispecific, designed and

produced the $\alpha HTLV$ bispecific, performed testing in T2 cells and NSCLC cells, and

revised the manuscript

SNB – Designed and produced $\alpha PRAME$ TCR T-cells and performed flow cytometry

staining to quantify transduction efficiency

SV – Predicted antigens using LENS, produced the corresponding figures, and also

wrote the methods section for LENS

MiS – Analyzed flow cytometry data

JDD – Prepared samples for western blotting

JKG – Analyzed long read DNA sequencing

BAP – Curated the *NUTM1* fusion cohort in the Tempus AI Sequencing Database and

performed sequencing analysis

KPN – Developed and provided the NUT carcinoma patient-derived xenograft

RJK - Developed and provided the NUT carcinoma patient-derived xenograft

RSK – Developed and provided the NUT carcinoma cell line JCM1

LEH - Analyzed immunopeptidomics data, generated mass spectrum figures, and

revised the mass spectrometry methods section.

IJD – Mentored SKP, assisted in sequencing analysis, and revised the manuscript

JRW – Mentored JKG and analyzed long read DNA sequencing

CAF – Provided NUT carcinoma cell lines, produced NUT carcinoma tissue microarray,

stained stained the TMA for PRAME, and revised the manuscript

BK – Mentored SY and TK, designed and produced $\alpha PRAME_{425}$ x SP34 $\alpha CD3$

bispecific, designed and produced the $\alpha HTLV$ bispecific, and revised the manuscript

JW – Conceived the project, mentored JLJ, generated hypotheses, revised the

manuscript, and obtained funding

AR – Conceived the project, mentored SKP, generated hypotheses, designed

experiments, analyzed sequencing data, revised the manuscript, and obtained funding

BGV – Conceived the project, supervised the project, mentored JLJ, generated

hypotheses, designed experiments, revised the manuscript, and obtained funding

Acknowledgments: We would like to acknowledge the work and contributions from

multiple core facilities at UNC including the Immune Monitoring and Genomics Facility,

the High Throughput Sequencing Facility, the Pathology Services Core Facility, the

Peptide Synthesis Core Facility, and the lentiviral/shRNA core facility.

List of Abbreviations:

Landscape of Neoantigen Software (LENS)

Non-small cell lung cancer (NSCLC)

T-cell receptor (TCR)

Cancer testis antigen (CTA)

Bispecific T-cell engager (BiTE)

Immune mobilizing monoclonal T-cell receptors against cancer (ImmTAC)

Dulbecco's modified eagle medium (DMEM)

Patient-derived xenograft (PDX)

Transcripts per million (TPM)

Relative centrifugal force (RCF)

Investigational new drug (IND)

Immunohistochemistry (IHC)

Data-dependent acquisition liquid chromatography tandem mass spectrometry (DDA

LC-MS/MS)

Data-independent acquisition (DIA)

Tissue microarray (TMA)

Abstract (Words=341):

Background: NUT carcinoma is a rare but highly lethal solid tumor without an effective

standard of care. NUT carcinoma is caused by bromodomain-containing *NUTM1* fusion

oncogenes, most commonly *BRD4::NUTM1*. BRD4::NUTM1 recruits p300 to acetylate

H3K27 forming expansive stretches of hyperacetylated chromatin called

"megadomains" with the overexpression of corresponding oncogenes, including *MYC*.

We hypothesized that transcriptional dysregulation caused by BRD4::NUTM1 would

lead to the generation of cancer-specific antigens that could be therapeutically

actionable.

Methods: We integrated genomics, computational antigen prediction software, targeted

immunopeptidomics using single- and double-labeled peptide standards, and gain/loss-

of-function genetic experiments on a panel of cell lines (N=5), a patient derived

xenograft, a tissue microarray (N=77), and patient samples from the Tempus AI

Sequencing Database harboring evidence of *NUTM1* fusions (N=165). We created an

$\alpha PRAME_{425}$ T-cell receptor x SP34 $\alpha CD3$ bispecific molecule modeled after

brenetafusp, an $\alpha PRAME_{425}$ T-cell receptor bispecific T-cell engager, as well as

$\alpha PRAME_{425}$ TCR T-cells based on anzutresgene autoleucel and we applied these

products to NUT carcinoma cells *in vitro*.

Results: We identified *PRAME* as the most commonly expressed cancer/testis antigen

in patient samples harboring the three canonical NUT carcinoma fusions

(*BRD4::NUTM1*, *BRD3::NUTM1*, and *NSD3::NUTM1*). Additionally, 56% (43/77) of NUT

carcinoma tissue microarray samples stained positive for PRAME. *BRD4::NUTM1*

expression in HEK 293T cells enhanced PRAME levels and *BRD4::NUTM1* knockout in

NUT carcinoma cells reduced PRAME levels. Immunopeptidomics detected more

PRAME-derived HLA ligands (N=9) than all other cancer/testis antigens combined

(N=5). Targeted mass spectrometry detected the HLA-A*02:01/SLLQHLIGL (PRAME$_{425}$)

epitope in 100% (4/4) of HLA-A*02+, PRAME+ NUT carcinoma samples at higher levels

(>0.01 fM) than HLA-A*02:01/RLDQLLRHV (PRAME$_{312}$) or HLA-A*02:01/YLHARLREL

(PRAME$_{462}$). The αPRAME$_{425}$ T-cell receptor x SP34 αCD3 bispecific molecule and

αPRAME$_{425}$ TCR T-cells each exhibited potent, T-cell mediated cytotoxicity against

PRAME+ NUT carcinoma cells.

Conclusions: PRAME is highly and frequently expressed in NUT carcinoma and the

most common oncoprotein causing NUT carcinoma, BRD4::NUTM1, contributes to

these high PRAME levels. PRAME epitopes presented by HLA Class I are a previously

unrecognized therapeutic vulnerability for NUT carcinoma that warrant clinical trials

testing PRAME targeted immunotherapies in this neglected patient population.

<u>What is already known on this topic:</u> NUT carcinoma is a devastating malignancy that is

recalcitrant to cytotoxic chemotherapy, T-cell checkpoint blockade, and targeted

therapies in the form of bromodomain inhibitors.

<u>What this study adds:</u> NUT carcinoma tumors are high in the cancer/testis gene

PRAME. The oncogene most commonly causing NUT carcinoma, *BRD4::NUTM1*,

contributes to these high levels. NUT carcinoma cells present PRAME epitopes on HLA

Class I molecules and are susceptible to PRAME-directed, T-cell mediated cytotoxicity.

<u>How this study might affect research, practice or policy:</u> Our results argue for phase I/II

clinical trials testing PRAME immunotherapies like brenetafusp or anzutresgene

autoleucel in *PRAME+* NUT carcinoma patients.

Graphical Abstract:

Background (Words = 684):

NUT carcinoma is a rare but underdiagnosed and devastating malignancy defined by

ectopic expression of the Nuclear Protein in Testis gene ("*NUT*" or "*NUTM1*"), most

commonly through an in-frame translocation to the bromodomain-containing *BRD4* with

formation of the *BRD4::NUTM1* fusion gene (75% of cases) (1). BRD4::NUTM1 is a

potent oncoprotein that initiates NUT carcinoma by binding to acetylated H3K27

("H3K27ac"), recruiting p300 and catalyzing the hyperacetylation of adjacent histones

resulting in extended hyperacetylated domains ("megadomains") with the corresponding

overexpression of corresponding oncogenes, including *MYC* and *SOX2* (2, 3). NUT

carcinoma was formerly labeled as NUT Midline Carcinoma owing to a predilection for

afflicting midline structures and is now known to occur most commonly in the lungs and

sinonasal passages (4). NUT carcinoma is not associated with smoking or any known

exposures and afflicts individuals of all ages including infants, children, and young

adults (5-7). Clinical outcomes in adults are dismal with median survival after diagnosis

of 6.7 months and similar outcomes in children due to rapid tumor growth, high

metastatic potential, and lack of sensitivity to systemic therapies (4, 8). Treatment for

localized disease is focused on aggressive attempts at local control with wide local

resection and adjuvant radiation and/or chemotherapy. However, even among NUT

carcinoma patients treated with complete surgical resection of the initial tumor,

outcomes are poor. In one cohort, patients treated with surgical resection experienced a

median survival time of 415 days post-diagnosis with no patient surviving for more than

3 years post diagnosis (4, 9).

Targeted therapeutic strategies to date have focused on blocking the binding of BRD4 to

H3K27ac using bromodomain inhibitors, but in phase I/II clinical trials these inhibitors

have yielded progression free survival times of only 1-2 months (10-13). At present, all

clinical trials testing new therapeutics for NUT carcinoma patients in the United States

are   testing   bromodomain   inhibitors   (NCT05019716,   NCT05372640,   and

NCT05488548). NUT carcinoma tumors are PD-L1 low, tumor mutational burden low,

and immune "cold" tumors that are also highly refractory to cytotoxic chemotherapy and

T-cell checkpoint blockade (4). New treatment modalities are clearly needed.

Interest in treating challenging solid tumors with HLA-restricted TCR-based therapies is

growing. The first phase III clinical trial of a TCR x CD3 activator bispecific (also known

as an ImmTAC or BiTE) showed overall survival benefit in the frontline setting for HLA-

A*02+ uveal melanoma targeting the gp100 peptide (14). Recently, the SPEARHEAD-1

trial showed durable responses in patients with heavily pre-treated synovial sarcoma or

myxoid round cell liposarcoma with afamitresgene autoleucel, a TCR T-cell therapy

expressing a TCR targeting an HLA-A*02 bound epitope from the cancer/testis gene

MAGE-A4 (15). This resulted in the first TCR T-cell product approval for afamitresgene

autoleucel and was quickly followed up by the IGNYTE-ESO trial which similarly

showed durable responses in the same disease types with letetresgene autoleucel,

another TCR T-cell product targeting epitopes from the cancer/testis gene *NY-ESO-1*

(HLA-A*02:01, -A*02:05, or -A*02:06) (16).

Like NUT carcinoma, synovial sarcoma and myxoid/round cell liposarcoma are

uncommon fusion protein-driven malignancies. >90% of these sarcomas harbor the

pathognomonic t(X;18) or t(12;16)(q13;p11), respectively (17, 18). Whereas MAGE-A4

and NY-ESO-1 antigens are common in these sarcomas, the antigenic landscape of

NUT carcinoma and the susceptibility of NUT carcinoma to T-cell mediated therapeutics

are unknown. We hypothesized that if BRD4::NUTM1 causes dysregulated gene

transcription through reliable epigenetic changes, then BRD4::NUTM1 could also

upregulate genes encoding cancer specific antigens that would serve as actionable

TCR targets. Here we show that BRD4::NUTM1 causes transcription of the

cancer/testis gene *PRAME*, and that *PRAME* is the predominant cancer/testis gene

expressed within a clinical patient cohort of *NUTM1* fusion genes and especially in

patients with canonical NUT carcinoma fusions (*BRD4::NUTM1*, *BRD3::NUTM1*,

NSD3::NUTM1) (19). We also show that NUT carcinoma cells present PRAME epitopes

with HLA Class I molecules and are susceptible to T-cell mediated killing mediated by

an affinity-matured PRAME epitope TCR x CD3 activating bispecific molecule or by

αPRAME TCR T-cells, with both modalities generated from TCRs currently in clinical

testing for common cancers. Our work identifies PRAME as a promising target for

cancer vaccines, TCR-based therapies, and TCR-mimetics in NUT carcinoma phase I/II

clinical trials.

Methods (Words = 2,255):

Reagents

Cell lines were maintained in DMEM with 10% FBS and 1% penicillin/streptomycin. The

PDX donor signed consent version 12 of the "Consent Form for Use of Tissue, Blood,

and Other Biological Materials in Research and Authorization for Use of Identifiable

Health Information in Biobanking" for the University of Wisconsin Carbone Cancer

Center Translational Science BioCore which was IRB approved on 10/21/2015. The

patient's tumor and a deidentified pathology report were provided to our investigators.

The tumor was chopped and minced into small pieces, dissolved in media containing

FBS (10%), penicillin/streptomycin (1%), amphotericin B (1%), and matrigel (1:1 mixture

with tissue prep) and injected through a 21-gauge needle into the flanks of female NSG

mice. The tumor was then propagated in the flanks of NSG mice (using the technique

above) and purified for analyses using tumor dissociation and mouse cell depletion kits

(Miltenyi Biotec). RNA was isolated using RNEasy columns (Qiagen). Western blots

used PVDF and blocking in 5% milk. Antibodies are described in **Supplementary Table**

1. siRNA transfections were performed with Lipofectamine RNAiMAX and 10 µM pools

of four siRNAs synthesized by IDT with [dT][dT] overhang, desalted purification, and no

3' modifications (**Supplementary Table 2**). Tet-on, blasticidin selection lentiviral

pInducer20 vectors containing *BRD4::NUTM1* or *GFP* have been described (20). VSV-G

pseudotyped lentiviral particles were packaged in 293T cells and purified by filtration.

TMA immunohistochemistry was done by BWH Dana-Farber-Harvard Medical School

Specialized Histopathology Services-Longwood (SHL) Core using αPRAME (Biocare

Medical ACI3252B, 1:100).

Immunopeptidome Predictions

Antigen prediction used LENS v1.5.1 (21). Sample-level FASTQs were trimmed and

filtered for quality using fastp 0.23.1 (22). The resulting FASTQ pairs were aligned to a

modified hg38 reference genome (excluding the Epstein-Barr Virus (EBV) contig) using

STAR 2.7.0f with parameters *--quantMode TranscriptomeSAM --outSAMtype BAM*

SortedByCoordinate --twopassMode Basic –outSAMunmappedWithin (23). Transcript-

level quantifications were generated using Salmon 1.1.0 with a GENCODE v37 GTF,

augmented with human retroviral-like sequences from gEVE (24). CTAs were defined

using a manually curated list of genes with testis and placenta-specific expression

(**Supplementary Table 3**). The initial list of CTAs was pulled from CTDatabase and

manually filtered using Human Protein Atlas.

Splice variants were detected using SNAF 0.7.0 (25). Expressed viruses were detecting

using the VirDetect workflow. HLA typing for class I alleles was performed using

seq2HLA 2.2 (26). We processed protein sequences from CTAs in the 50th expression

percentile for downstream peptide prediction. Gene fusions were predicted using

STARfusion 1.10.1 with recommended parameters and trimmed RNA FASTQs.

Relevant protein sequences derived from CTAs, fusion, splice variants, viruses, and

ERVs were filtered for potential peptides of interest using NetMHCpan 4.1b (27).

RNA Sequencing

Library preparation used Illumina's Stranded mRNA Prep Ligation kit, starting with 125

ng of RNA. The average fragment length for these libraries was between 300-400 bp.

Paired-end sequencing was performed using the NovaSeq 6000 version 1.5 (2X100),

obtaining at least 45M reads per sample. Bulk RNA-seq samples were processed using

selected components of the nf-core RNA-seq pipeline (v3.14), an open-source workflow

managed with Nextflow. Reads were demultiplexed, and quality was assessed using

FastQC, with results summarized in MultiQC (28). Adapter trimming was performed with

cutadapt, ribosomal RNA was removed with SortMeRNA, followed by alignment to the

GRCh38 reference genome using STAR (29). Transcript-level quantification was

conducted with Salmon (24). Counts were summarized to the gene level and

normalized to TPM to generate a gene x sample matrix. All downstream visualization

and analysis were performed in R. All scripts used for this project are publicly available

on GitHub (https://github.com/pirl-unc/PRAME_in_NUT_Carcinoma).

Immunopeptidomics and Targeted Mass Spectrometry

Snap frozen cell pellets were submitted to Biognosys for immunopeptidomics. Samples

and a processing quality control sample (JY cells) were lysed using Biognosys' Lysis

Buffer for 1 hour at 4^{o}C. After removal of cell debris, the lysates were incubated with

bead coupled antibodies (clone W6/32) to capture HLA class I complexes with

associated peptides. Bound protein complexes were washed and peptides eluted using

Biognosys' optimized immunopeptidomics protocol. Clean up for mass spectrometry

was carried out using an Oasis HLB µElution Plate (30 µm). Immunopeptides were

dried down using a SpeedVac system and resuspended in 10 µl of 1% acetonitrile and

0.1% formic acid in water. Samples were pooled and fractionated into 5 fractions using

high pH reversed phase fractionation (HPRP) for spectral library generation.

For DDA LC-MS/MS measurements of fractionated samples, 8 µl of immunopeptides

per sample were injected to a Biognosys in-house packed reversed-phase column on a

NeoVanquish UHPLC nano liquid chromatography system connected to a Orbitrap

Exploris 480 mass spectrometer equipped with a Nanospray Flex ion source and a

FAIMS Pro ion mobility device. LC solvents were A: water with 0.1% FA, B: 80%

acetonitrile. The nonlinear LC gradient was 1 - 40% solvent B in 120 minutes followed

by a column washing step in 90% B for 11 minutes, and a final equilibration step of 1%

B for 5 minutes at 64^{o}C with a flow rate set to 250 nl /min. MS1 precursor scans were

acquired between 330-1250 m/z at 120,000 resolution with data dependent MS2 scans

at 60,000 resolution. MS2 scans were acquired in cycle time of 1.6 s and 1.4 s per

applied FAIMS compensation voltage. Only precursors with charge state 1-5 were

isolated for dependent MS2 scans.

For DIA LC-MS/MS measurements of study samples, 8 µl of immunopeptides per

sample were injected to a Biognosys in house packed reversed phase column on a

NeoVanquish UHPLC nano liquid chromatography system connected to a Orbitrap

Exploris 480 mass spectrometer equipped with a Nanospray Flex ion source and a

FAIMS Pro ion mobility device (Thermo Scientific). The system was used in trap and

elute configuration using a 300 µm x 5 mm PepMap Neo C 18 Trap 3 PK cartridge pre

column (Thermo Scientific 174502). LC solvents were A: water with 0.1% FA, B: 80%

acetonitrile, 0.1% FA in water. The nonlinear LC gradient was 1-40% solvent B in 120

minutes followed by a column washing step in 90% B for 11 minutes, and a final

equilibration step of 1% B for 5 minutes at 64^o C with a flow rate set to 250 nl /min. The

FAIMS DIA method consisted of one full range MS1 scan and 20 DIA segments as

adopted from Bruderer et al (30) and Tognetti et al (31).

The shotgun and HRM mass spectrometric data was analyzed using Spectronaut v18,

the false discovery rate on peptide and protein level was set to 1 %. A human UniProt

.fasta database (Homo sapiens, 2025-01-01) appended with *NUTM1* fusion proteins

was used for the search engine, also supplemented with mutated *NUTM1* and *PRAME*

isoform fastas, allowing for non-specifically digested peptides (no trypsin cleaved

termini) with a length between 7 and 12 amino acids, max. 2 variable modifications (N-

term acetylation, carbamidomethylation [C], pSTY) and requiring carbamidomethylation

methionine oxidation as fixed modification. The identification results were then exported

as a hybrid spectral library. Raw HRM mass spectrometric data were analyzed using

Spectronaut v18 with Qvalue filtering and no background signal integration or

imputation. The hybrid spectral library generated in this study was used. Peptide level

false discovery rate control was 1 % and cross-run normalization using global

normalization to the median. MS2 spectra were visualized and annotated using the

Interactive Peptide Spectral Annotator. Fragment ion tolerance was set to 5ppm for a-,

b-, and y-ions.

For targeted mass spectrometry, cells were homogenized using a Precellys

homogenization device (Bertin Technologies) and then lysed as above along with a

quality control sample pellet of 10 million Raji cells. Protein concentrations were

measured with a BCA assay (Pierce, Thermo Scientific) then HLA Class I-bound

peptides were isolated, cleaned up, dried down and resuspended as above. Four stable

isotope labeled reference peptides were spiked into the final peptide samples at known

concentrations (manufactured by SciTide with quality grade +/-10% quantification

precision, >95% purity). For LC-PRM measurements, 50% of total immunopeptide yield

per sample (1 μg total peptide anticipated per injection) was injected with 5 fmol SIS

peptides on custom-length Aurora 3 Freontier-type reversed phase column (IonOpticks)

on a ThermoScientific NeoVanquishUHPLC nano-liquid chromatography system

connected to a ThermoScientific Orbitrap Exploris 480 mass spectrometer equipped

with a Nanospray Flex ion source and a FAIMS Pro ion mobility device

(ThermoScientific). LC solvents were A: water with 0.1% FA; 80% acetonitrile, 0.1% FA

in water. The LC gradient was 0-50% solvent B in 55.8 min followed by 51-90% B in 10

sec, 90% B for 5.9 min to wash the column and a 1.2 min step at 1% B to equilibrate the

column. The total gradient length was 63.3 min. An unscheduled run in PRM mode was

performed before data acquisition for retention time calibration as previously described

(32). Signal processing and data analysis was carried out using SpectroDive 12.4 and

Q-value filter of 1%.

Bispecific Protein Production

Coding sequences for PRAME soluble TCR specific for the HLA-A*02:01-restricted

SLLQHLIGL epitope were derived from sequences previously described for the PRAME

Vα and Vβ domains and from the Protein Data Bank ID 6U07 crystal structure for the

computationally thermostabilized Cα and Cβ domains (33, 34). Chimeric SP-34 αCD3

heavy and light chain sequences were published previously (35). One primary fusion

construct and two auxiliary constructs were generated as clonal genes in mammalian

expression vector plasmids (Twist Bioscience). The primary fusion constructs consisted

of the SP-34 αCD3 heavy chain, a $(Gly_4Ser)_4$ linker, PRAME TCR Vα/Cα, and an 8xHis-

Tag. The first auxiliary coding sequence encoded for SP-34 αCD3 light chain; the

second for PRAME TCR Vβ/Cβ (**Supplementary Table 4)**.

All three plasmid constructs were co-transfected for transient expression in Expi293F

cells cultured with Expi293 Expression Medium at 50 mL scale using ExpiFectamine

293 Transfection Kit, similarly to previously described with the following modifications

(36). Plasmids for primary fusion construct, first auxiliary construct, and second auxiliary

construct were mixed in a 3:1:1 mass ratio, totaling 1 µg plasmid mixture per milliliter of

cell culture. At 18 hours after transfection, enhancer 1 (300 µL at 50 mL scale) and

enhancer 2 (3 mL at 50 mL scale) and 6 mL of 3.5% (w/v) Hyclone Cell Boost 1 were

added to the cells. Cell culture supernatant was harvested at 96 hours after transfection

by pelleting cells at 3428 RCF at 4°C for 15 min and further clarified using a 0.22 µm

polyvinylidene difluoride membrane filter (Millipore). Assembled PRAME bispecific

constructs were affinity purified by batch-binding clarified medium with pre-equilibrated

penta-Ni resin (Marvelgent) for 1 hour at 4°C and then transferred into a column to

collect the resin. A three-step wash was performed by gravity at room temperature (RT)

to eliminate protein contaminants. The first wash was performed with 6 CV (column

volume) of high-salt Tris buffer [50 mM Tris (pH 8.0) and 1 M sodium chloride with 10

mM imidazole]; the second was performed with 2 CV wash with 1× PBS (pH 7.4) with

20 mM imidazole; and the third was performed with 2 CV wash with 1× PBS (pH 7.4)

with 40 mM imidazole. Bound proteins were eluted in one step using 5 CV of 1× PBS

(pH 7.4) and 500 mM imidazole to collect the protein and then sequential concentration

at 4°C using a 30K MWCO Amicon-4 followed by an Amicon-0.5 centrifugal filter at

maximum allowed RCF for each filter, serially exchanged into 1× PBS (pH 7.4) to

ensure at least 1:1000 dilution of imidazole. Purity was assessed by SDS-

polyacrylamide gel electrophoresis.

TCR T-Cells

TCR T-cells were generated using a previously described approach to knock-out

endogenous α and β chains, followed by retroviral transduction to express the

anzutresgene autoleucel αPRAME TCR (37-39). CD8+ T-cells were isolated from

healthy donor PBMCs using an EasySep Human CD8+ T-Cell Isolation Kit (StemCell

Technologies) and stimulated for 72 hours with plate-bound αCD3 (1 μg/mL,

BioLegend) and αCD28 (1 μg/mL, BioLegend) with Il-2 (300 IU/mL, StemCell

Technologies), IL-7 (5 ng/mL, StemCell Technologies), and IL-15 (5 ng/mL, StemCell

Technologies). Following activation, T-cells were electroporated with TCR α and β

chain-targeting ribonucleoproteins using a 4D Nucleofector X (Lonza) and cultured for

48-72 hours with 180 IU/mL IL-2. To generate retroviral supernatants containing the

TCR of interest, HEK292T cells were transfected with retroviral plasmids, including SPR

plasmid containing the αPRAME TCR, using GeneJuice Transfection Reagent (Sigma-

Aldrich) and harvested at 48 and 72 hours as previously described (39). TCR null CD8+

T-cells were transduced in Retro-Nectin-coated plates (Takara Bio) according to the

manufacturer's protocol and cultured for a minimum of 72 hours prior to screening. To

confirm TCR expression and specificity, TCR T-cells were stained on ice with peptide-

loaded tetramers (Flex-T A02 monomers, PE, BioLegend), αCD8 (BV421, Clone: Hit8a,

BioLegend) and fixable live/dead stain (NIR, ThermoFisher). Cells were fixed and run

on a BD Symphony A3 flow cytometer (BD Biosciences); analysis was performed with

FlowJo (v.10.10.0).

Cytotoxicity Assays

Whole blood was obtained from Gulf Coast Regional Blood Center. PBMCs were

isolated using Ammonium-Chloride-Potassium Lysing Buffer and Density Gradient

Medium (Stemcell Technologies). T-cells were isolated using a negative selection,

magnetic bead kit (Stemcell Technologies) and then incubated with RPMI and 10% FBS

for 5 days in an αCD3 coated (5 µg/mL, BD Biosciences) flask with carrier-free IL-2 (2

ng/mL, 18.2 IU/mL, R&D Systems) and αCD28 (2.5 µg/mL, BD Biosciences). Expanded

T-cells were co-cultured in RPMI media with 10% FBS. Luminescence output was either

read using CellTiter-Glo reagent for luciferase-free cell lines or by adding luciferin (250

µg/mL, Promega). Peptide pulses used 1 µg/mL of SLLQHLIGL peptide, synthesized by

the UNC High-Throughput Peptide Synthesis and Array Facility over three hours (T2

peptide pulse) or 10 µg/mL of SLLQHLIGL (GenScript) added at the beginning of the

experiment (TCR T-cell cytotoxicity).

Results (Words=2,121):

RNA sequencing and computational predictions suggest PRAME and NUTM1 are

sources of HLA Class I peptide ligands in NUT carcinoma

To test the hypothesis that transcriptional dysregulation due to NUTM1 fusion

oncoproteins would lead to the production of cancer-specific antigens, we obtained five

NUT carcinoma cell lines (HLA-A*02 positive:TC-797, 10-15, 14169, PER403. HLA-

A*02 negative: JCM1) and one HLA-A*02+ PDX described in **Table 1**. We then

performed RNA sequencing on these samples and applied the LENS computational

workflow (**Figure 1A**). LENS is an open-source method that predicts tumor-specific and

tumor-associated antigens from genomic data (21). LENS predicted HLA Class I binding

for peptides derived from several CTAs but only PRAME and NUTM1 had predicted

epitopes across multiple samples (**Figure 1B-C**). Interestingly, LENS predicted the

translationally relevant SLLQHLIGL (PRAME$_{425}$) epitope in every sample except for TC-

797 in which *PRAME* was transcriptionally silent (40, 41). Whole genome long-read

DNA sequencing failed to identify a genetic cause for the absence of PRAME

expression in TC-797 (data not shown). In screening cancer/testis genes using our

manually curated list of CTAs (**Supplementary Methods Table 3**), out of 164 different

cancer/testis genes only *PRAME* and *NUTM1* were consistently expressed across the

tested samples (**Figure 1D**). To evaluate protein-level heterogeneity of PRAME we

stained the samples with αPRAME and confirmed TC-797 to be PRAME- with PER403,

14169, and JCM1 showing high proportions of PRAME+ cells (**Figure 1E**). We

concluded that in this N=6 set of NUT carcinomas, PRAME and NUTM1 are the most

promising sources of cancer-specific antigens for therapeutic action.

RNA sequencing and immunohistochemistry identify high frequency and levels of

PRAME RNA and PRAME protein in NUT carcinoma patient samples

To determine if the predominance of *PRAME* and *NUTM1* expression among

cancer/testis genes was reflected in patient samples, we searched the de-identified

Tempus multimodal database for patients with detected *NUTM1* fusion genes (identified

via Tempus xT [solid-tumor DNA] (42), Tempus xR [solid-tumor RNA] (43), or Tempus xF

[circulating tumor DNA] assays) (44). Research filters below clinical reporting were

applied to obtain patients with evidence of a *NUTM1* fusion, defined as containing at

least 10 total supporting reads from DNA or at least 5 total supporting reads from RNA.

Subject to these thresholds, 165 patients had evidence of *NUTM1* fusions, with a total

of 33 distinct fusion partners (**Table 2**). Consistent with their role as canonical NUT

carcinoma fusion partners, *BRD4::NUTM1, NSD3::NUTM1,* and *BRD3::NUTM1*

accounted for the majority of cases (35.8%, 17.6%, and 10.9% respectively) with the

remaining samples harboring one of the 30 other fusion partners (**Supplementary**

Figure 1A). As expected, *BRD4::NUTM1*, *BRD3::NUTM1*, and *NSD3::NUTM1* fusions

primarily occurred in the lung and head/neck primary sites – typical anatomic locations

for NUT carcinoma (9). Analysis of gene expression from 162 of the patients with

available RNA data revealed that out of 157 CTA genes covered in the Tempus xR

assay, *PRAME* was identified as having the highest average expression after *NUTM1*

(**Figure 2A**). We observed a bimodal distribution of *PRAME* expression in comparison

to *NUTM1* across the cohort, designating *PRAME*^{high} and *PRAME*^{low} groups using a

threshold of 3.1 $\log_2$(TPM+1) (**Figure 2B**). We found the *BRD4::NUTM1* group to be

significantly enriched in *PRAME*^{high} samples (fisher exact, *p*<0.001) with *BRD3::NUTM1*

and *NSD3::NUTM1* also having more *PRAME*^{high} samples than *PRAME*^{low} (**Figure 2C**).

Of the remaining non-canonical fusion partners, 14 partners had at least one instance of

PRAME^{high} with 3 fusion partners having more than one: *MGA::NUTM1*,

SLC12A6::NUTM1, and *CIC::NUTM1* (**Supplementary Figure 1B**). Notably, only 1 of

13 *YAP1::NUTM1* samples was classified as *PRAME*^{high}. Using the *PRAME*^{high} and

PRAME^{low} groups, we identified several primary sites with the highest percentage of

PRAME^{high} samples (**Figure 2D**). These frequently *PRAME*^{high} sites included typical

NUT carcinoma sites of origin including sinuses, head/face/neck, the nasal cavity, and

lung.

To assess for intratumoral heterogeneity of protein level PRAME expression in NUT

carcinoma patient samples, we stained a previously described NUT carcinoma tissue

microarray for PRAME and quantified the percentage of PRAME positive cells within the

tumor (**Figure 2E, 2F**) (45). We found high frequency of PRAME expression in 28/77

(36%) of patient samples with ≥75% of tumor cells staining positive. Overall 43/77

(56%) of patient samples exhibited some PRAME staining.

BRD4::NUTM1 drives PRAME expression

Due to the high levels and frequency of *PRAME* observed in cell lines (N=5), a PDX

(N=1), the Tempus AI sequencing database (N=165), and the NUT carcinoma tissue

microarray (N=77), and because PRAME expression is regulated by epigenetic changes

including H3K27 acetylation in non-malignant human physiology (46), we hypothesized

that *NUTM1* fusion oncoproteins might be contributing to transcriptional dysregulation

and causing the ectopic expression of PRAME. To test this hypothesis, we transduced

HEK 293T cells with tet-on *BRD4::NUTM1* or *eGFP* lentiviral particles and observed an

increase in PRAME protein levels with *BRD4::NUTM1* expression but not with GFP

(**Figure 3A**). We then used siRNAs targeting *NUTM1* to knockdown BRD4::NUTM1 in

the five NUT carcinoma cell lines and immunoblotted for PRAME (**Figure 3B**). We found

that in all cell lines except for the *PRAME* negative TC-797 cell line, BRD4::NUTM1 loss

was accompanied by a decrease in PRAME protein levels. The decrease in PRAME

with BRD4::NUTM1 knockdown was marginal in 14169 compared to other NUT

carcinoma cell lines, so we performed a western blot time course in 14169 cells which

revealed a greater effect on PRAME levels at a later timepoint, 4 days instead of 3

days, after *NUTM1* siRNA transfection (**Figure 3C**). We considered whether these

effects might be mediated through the corresponding loss of MYC after BRD4::NUTM1

knockout, but in our review of the literature describing MYC-target genes we could find

no evidence to suggest that *PRAME* is a MYC-target gene (47). We concluded that

BRD4::NUTM1 is contributing to the expression of PRAME in *BRD4::NUTM1* NUT

carcinomas, which may be independent of MYC.

Immunopeptidomics on NUT carcinoma samples identifies more HLA Class I PRAME

ligands than all other analyzed CTAs combined

We hypothesized that HLA Class I-bound PRAME epitopes would be detectable using

immunopeptidome profiling with mass spectrometry. Shotgun (DIA) mass spectrometry

was performed with the five cell lines and with the PDX to identify and characterize

NUTM1, PRAME, and other CTA epitopes being presented by HLA Class I molecules

and to guide therapeutic TCR selection. We identified a total of 33,644 unique peptides

among the six samples **(Supplementary Figure 2)**. Focusing on the presented CTAs,

we identified 14 unique peptides from 6 proteins out of all CTAs with detection in at least

1 sample. Of these 14 peptides, 9 corresponded to PRAME epitopes and 1

corresponded to a novel, previously undescribed NUTM1 epitope (**Figure 4A**). The

remaining four CTA peptides corresponded to *MAGE-A1*, *CT83*, *CTAG2* and *ACTL8*. As

expected, no PRAME epitopes were discovered in the *PRAME* negative TC-797 cell

line. The well described and therapeutically actionable HLA-A*02:01/SLLQHLIGL

(PRAME$_{425}$) epitope was detected in the HLA-A*02+ PER403 and PDX samples but not

10-15 or 14169 (48-50). To detect fusion junctional peptides, long read RNA sequencing

was performed and used to generate consensus fusion sequences (**Supplementary**

Figure 3). A reference human proteome augmented with these fusion sequences failed

to identify junctional fusion peptides.

Due to differential fragmentation of amino acids, quantification of peptide epitopes with

untargeted immunopeptidomics is inherently semi-quantitative. Because SLLQHLIGL

(PRAME$_{425}$) was only detected in 2/4 (50%) of PRAME+, HLA-A*02+ samples and

because the SLLQHLIGL (PRAME$_{425}$) epitope with the most clinical developmental

interest had lower semi-quantitative signal intensities than YLHARLREL (PRAME$_{462}$),

we developed targeted mass spectrometry assays for three PRAME epitopes and one

novel NUTM1 epitope using single- and double-labeled peptide standards to quantify

these epitopes across our six NUT carcinoma samples. Single- or double-labeled

standards were used to enhance our confidence in the quantitative results across two

independent sets of experiments. The labeled peptide standards used for these assays

are described in **Table 3**. We found similar results and trends using single- and double-

labeled peptide standards, and were able to detect SLLQHLIGL (PRAME$_{425}$) in 4/4

(100%) of PRAME+, A02+ samples at abundances higher than PRAME$_{312}$ or

PRAME$_{462}$. Interestingly, DVYENFRQW (NUTM1$_{237}$) (**Figure 4B-D**) exhibited the

highest absolute abundance of all epitopes and samples in the JCM1 cell line in the

single-labeled assays and the second highest absolute abundance in the JCM1 cell line

in the double-labeled assays but was very low abundance or non-detectable in the other

5/6 samples, likely due to DVYENFRQW being a HLA-A*25:01 ligand (MHCflurry

predicts this as the only JCM1 HLA class I molecule with binding specificity for

DVYENFRQW).

Based on the increased prevalence of PRAME epitopes in our samples relative to

NUTM1, the rarity of HLA-A*25:01 compared to HLA-A*02:01, and based on the

availability of existing PRAME therapeutics for testing, we focused on therapeutic

targeting of PRAME epitopes in NUT carcinoma.

PRAME+ NUT carcinoma cells are susceptible to T-cell killing mediated by an

αPRAME$_{425}$ x SP34 αCD3 BiTE or by αPRAME$_{425}$ TCR T-cells

The susceptibility of NUT carcinoma to TCR-mediated therapeutics like TCR bispecifics

or TCR T-cell therapies is unknown. To explore this issue and to evaluate the

translational relevance of PRAME cancer-specific antigens, we created an αPRAME$_{425}$

TCR x SP34 αCD3 BiTE molecule with the same variable regions as the affinity-

matured TCR BiTE brenetafusp (**Figure 5A**). Brenetafusp has completed phase I/II

testing and is currently being evaluated as first line therapy in a randomized phase 3

trial (NCT06112314) for melanoma patients (51). Because our αPRAME$_{425}$ TCR x SP34

αCD3 BiTE has significant homology to brenetafusp including perfect homology to the

variable regions of the TCR, we anticipated our bispecific molecule to have specificity

for the same SLLQHLIGL (PRAME$_{425}$) epitope brenetafusp targets. We used

SLLQHLIGL peptide pulsing and TAP deficient T2 cells in co-culture with allogeneic T-

cells to demonstrate enhancement of bispecific-mediated cytotoxicity with peptide pulse.

TAP deficient T2 cells have empty HLA molecules on the cell surface and are loaded by

incubating the cells with an HLA peptide ligand. Killing of peptide-pulsed T2 cells thus

demonstrated the specificity of the bispecific molecule for the SLLQHLIGL epitope

(**Figure 5B**). We then titrated the concentration of the bispecific in the co-cultures of a

PRAME+, HLA A02+ NSCLC cell line in the presence of allogeneic T-cells and found a

dose-dependent effect (**Figure 5C**). The *PRAME* negative TC-797 and the *PRAME*

positive 10-15 and 14169 cell lines were tested alongside a haplo-incompatible NSCLC

cell line (H460, HLA A*24:02, *68:01) for T-cell mediated cytotoxicity in co-culture

experiments using 10^{-8} M of bispecific molecule. We observed no cytotoxic effect of the

bispecific in TC-797 or H460 at either of the tested effector:target ratios but we did

observe potent cytotoxicity that was enhanced by a higher effector:target ratio in both

PRAME positive NUT carcinoma cell lines (**Supplementary Figure 4A-B**).

This first cytotoxicity experiment was done using luminescent cell lines, however,

PER403 and JCM1 luminescent cell lines were unavailable. Thus, we designed a

second experiment to include PER403 and JCM1 in which T-cells were washed off of

the adherent tumor cells and then tumor cell viability was assessed using CellTiter-Glo.

To further demonstrate specificity of the αPRAME$_{425}$ x SP34 αCD3 bispecific for HLA

Class I-mediated cytotoxicity we also created a control, αHTLV-1 Tax x SP34 αCD3

bispecific molecule identical to the PRAME bispecific except for changes in the variable

regions (**Figure 5A**). Again, we observed potent T-cell mediated cytotoxicity in

PRAME+, A02+ NUT carcinoma cells with the αPRAME$_{425}$ bispecific molecule but not

with the αHTLV-1 Tax bispecific or in no bispecific controls (**Figure 5E, Supplementary**

Figure 4C).

Lastly, to demonstrate that NUT carcinoma is likely susceptible to PRAME TCR

immunotherapies as a class we also generated off-the-shelf αPRAME$_{425}$ TCR T-cells

using the TCR from anzutresgene autoleucel: a PRAME TCR T-cell product in early

phase clinical testing for patients with melanoma and other advanced solid tumors

(NCT03686124, NCT06743126). The native TCR was genetically ablated using

transfection with Cas9 protein and TCR sgRNA guides (1 α chain sgRNA, 1 β chain

sgRNA), and then TCR null T-cells were transduced with αPRAME$_{425}$ TCR encoding

lentiviral particles or with vehicle control (**Figure 5A**). Flow cytometry with αTCR

immunoglobulin and SLLQHLIGL tetramer staining demonstrated successful ablation of

the naïve TCR and lentiviral TCR transduction efficiencies of 66% or 41%, respectively

by staining method (**Figure 5D, Supplementary Figure 5**). We then co-cultured TCR

null T-cells and αPRAME$_{425}$ TCR T-cells with the five NUT carcinoma cell lines, with and

without SLLQHLIGL peptide pulse and found complete or nearly complete ablation of

target tumor cells at the lowest E:T ratio tested (1:1) in all 3 PRAME+, HLA-A*02+ NUT

carcinoma cell lines treated with αPRAME$_{425}$ TCR T-cells but not with αHTLV-1 Tax TCR

T-cells (**Figure 5F, Supplementary Figure 4D**). Importantly, in HLA-A*02:01 but

PRAME- TC-797, peptide pulse with SLLQHLIGL peptide rescued the cytotoxicity

effects of αPRAME$_{425}$ TCR T-cells consistent with antigen-directed killing. We

concluded that PRAME epitopes are a previously unrecognized therapeutic vulnerability

in NUT carcinoma.

Discussion/Conclusion (Words=1,102):

Recent advances in antigen-directed immunotherapy have resulted in a new generation

of therapeutics that leverage cell surface protein or TCR/peptide/HLA binding

interactions for tumor targeting and specificity. Approved agents in solid tumors target

either HLA-presented peptide epitopes such as gp100 in uveal melanoma, MAGE-

A4/NY-ESO-1 epitopes in synovial sarcoma/myxoid round cell liposarcoma, or surface

targets such as DLL3 in small cell lung cancer (14, 15). To identify promising targeted

immunotherapies for NUT carcinoma, a highly lethal and treatment-refractory

malignancy, we performed RNA sequencing on a selection of NUT carcinoma samples

and applied the results to the antigen prediction software LENS which predicted

PRAME as a plentiful source of cancer-specific antigens in NUT carcinoma (21). We

found that *PRAME* was highly expressed in cell lines and the PDX model, consistent

with observations in a large cohort of *NUTM1* fusion patient samples (especially among

canonical *NUTM1* fusion patient samples) as well as in a NUT carcinoma tissue

microarray.

Using genetic approaches, we showed that elevated PRAME levels in *BRD4::NUTM1*

NUT carcinoma are enhanced or at least maintained by BRD4::NUTM1. By applying

immunopeptidomics to our panel of NUT carcinoma samples we also detected 9

PRAME epitopes, including the actionable SLLQHLIGL (PRAME$_{425}$) epitope that is

targeted by brenetafusp and anzutresgene autoleucel. NUTM1 epitopes were nearly

absent from our immunopeptidomics results, likely due to immunoselection for non-

antigenic fusion proteins, limitations of detection by shotgun immunopeptidomics, and/or

strong nuclear localization of NUTM1 fusion proteins (13). However, we did identify a

single novel NUTM1 epitope, likely binding HLA-A*25:01, confirmed through targeted

mass spectrometry to be absent or at very low levels of abundance in all of the HLA-

A*25:01 negative samples. Future identification of other NUTM1 epitopes for NUTM1-

specific TCR therapeutic development will require more sensitive immunopeptidomic

methods, or perhaps targeted mass spectrometry using computationally predicted

epitopes.

PRAME epitopes are appealing candidates for peptide-HLA-directed therapeutics

because of the strong localization of PRAME to the testis in healthy human physiology

and because they arise as a direct result of the initiating genetic lesion in NUT

carcinoma (*BRD4::NUTM1*). We created an αPRAME$_{425}$ TCR x SP34 αCD3 molecule

modeled after brenetafusp that then showed potent, T-cell-mediated cytotoxicity against

PRAME+ NUT carcinoma cells but not *PRAME-* NUT carcinoma or haploincompatible

NSCLC cells. We also showed NUT carcinoma cells to be susceptible to αPRAME$_{425}$

TCR T-cell antigen-directed killing using the TCR from anzutresgene autoleucel,

building our confidence that PRAME immunotherapies as a class might perform well

against NUT carcinomas in clinical trials.

There are multiple limitations of this work. We demonstrated PRAME antigen-specific

cytotoxicity *in vitro* have not presented *in vivo* treatment models here. This is in part

because brenetafusp and anzutresgene autoleucel are already in clinical trials, making

unexpected toxicities or dosing issues in mice less relevant. Secondly, mouse models

for human antigen-directed therapeutics are contrived as they require human T-cell

injections into immunocompromised mice and the human T-cells have alloreactivity to

mouse tissues. We also encountered unexpected results in our cytotoxicity

experiments. Specifically, the αPRAME bispecific enhanced proliferation in PRAME-

TC-797 at E:T=5:1 in two independent experiments. The αHTLV-1 Tax bispecific also

exhibited this effect. These data suggest that CD3 stimulation enhances TC-797

proliferation in the absence of antigen-binding, perhaps through cytokine-mediated

effects. There was also unexpected cytotoxicity against HLA-A*02- JCM1 cells with both

the αPRAME BiTE and αPRAME TCR T-cells even though SLLQHLIGL (PRAME$_{425}$) is

broadly recognized as an HLA A02 ligand. This may be due to reactivity against the

SLLQHLIGL (PRAME$_{425}$) peptide presented by HLA-B*08:01, as computational

predictions suggest SLLQHLIGL (PRAME$_{425}$) has high binding affinity for this JCM1

HLA-B allele (52).

In conclusion, we report here that the most common oncogene initiating NUT

carcinoma, *BRD4::NUTM1*, enhances PRAME expression, that PRAME is a frequently

and highly expressed cancer/testis gene in *BRD4::NUTM1, NSD3::NUTM1* and

BRD3::NUTM1 NUT carcinomas, and that the resulting PRAME epitopes are actionable

targets for TCR therapeutics in NUT carcinoma. Our work adds NUT carcinoma to the

growing cadre of malignancies for which there is immense interest in targeting PRAME.

Specifically, PRAME is prevalent in genetically heterogeneous cancers including

melanoma, acute myeloid leukemia, NSCLC, ovarian cancer, and others (53). In healthy

human physiology, *PRAME* is turned on at the earliest stages of spermatogenesis and

is silenced by histone modifications upstream of the translation initiation site (46). In

malignancies, PRAME expression has been associated with worse clinical outcomes,

DNA double strand breaks, aneuploidy, and genetic instability (46, 54-56). Oncogenic

selection pressures in NUT carcinoma likely favor overriding the natural protections

against ectopic PRAME production via BRD4::NUTM1-p300 acetylation of H3K27 (2,

45, 57). EZH2 is active in NUT carcinoma and promotes inhibitory trimethylation of

H3K27 which is mutually exclusive with H3K27ac domains in NUT carcinoma (45).

Thus, H3K27 trimethylation is a plausible mechanism by which TC-797 and other

BRD4::NUTM1 patient samples could restrict *PRAME* expression. Ultimately, further

assays of histone states, such as ChIP-Seq or CUT&RUN, could explore this possibility.

The mechanisms promoting transcription of *PRAME* are likely conserved between

BRD4::NUTM1, BRD3::NUTM1, and NSD3::NUTM1 because BRD4 and BRD3 are

highly homologous (>80%) and because NSD3::NUTM1 recruits BRD4, thus forming a

protein complex with the same constituents as BRD4::NUTM1 (19, 58).

Our work suggests that clinical trials of PRAME targeting TCR therapeutics including

TCR T-cell therapies, endogenous T-cell therapy, TCR BiTEs/ImmTACs, tumor

vaccines, and/or immunoglobulins specific for HLA-bound PRAME epitopes ("TCR

mimetics") are warranted in NUT carcinoma. Potential agents to test include the

aforementioned brenetafusp and the TCR T-cell therapy anzutresgene autoleucel (40,

41). A PRAME-specific TCR is included in the presently recruiting trial NCT05973487: a

personalized TCR T-cell trial for patients with locally advanced or metastatic solid

tumors. An RNA-loaded dendritic cell vaccine using *PRAME* has completed phase I

testing for allogeneic stem cell transplant-ineligible acute myeloid leukemia and other

PRAME vaccines are in development (59, 60). Pr20, a PRAME-specific TCR mimetic

has been developed and showed activity against human leukemias in xenografts (61).

Brenetafusp has been tested in PRAME+ and PRAME- tumors (as determined by IHC)

with PRAME+ tumors showing better responses (40). Our clinical cohort showed

common NUT carcinoma anatomical origins to be associated with very high or universal

PRAME expression, suggesting that either IHC and/or anatomical origin could be

reasonable inclusion criteria for PRAME immunotherapy trials in this population.

Further, the growing set of PRAME therapeutics may have greater efficacy in NUT

carcinoma than in melanoma, NSCLC or other common malignancies without unifying

genetic etiologies because of a mechanistic link between the central genetic

rearrangements in NUT carcinoma and *PRAME* expression. Mechanistic studies linking

bromodomain fusion proteins to PRAME and clinical trials testing PRAME therapeutics

in NUT carcinoma are logical next steps to discovering efficacious treatments for this

rare, orphan disease.

References:

1.   Eagen KP, French CA. Supercharging BRD4 with NUT in carcinoma. Oncogene.
2021;40(8):1396–408.
2.   Alekseyenko AA, Walsh EM, Wang X, Grayson AR, Hsi PT, Kharchenko PV, et al.
The oncogenic BRD4-NUT chromatin regulator drives aberrant transcription within large
topological domains. Genes & Development. 2015;29(14):1507–23.
3.   Wang R, Liu W, Helfer CM, Bradner JE, Hornick JL, Janicki SM, et al. Activation
of SOX2 expression by BRD4-NUT oncogenic fusion drives neoplastic transformation in
NUT midline carcinoma. Cancer research. 2014;74(12):3332–43.
4.   Kloker LD, Sidiras M, Flaadt T, Brecht IB, Deinzer CKW, Groß T, et al. Clinical
management of NUT carcinoma (NC) in Germany: Analysis of survival, therapy
response, tumor markers and tumor genome sequencing in 35 adult patients. Lung
Cancer. 2024;189:107496.
5.   Flaadt T, Wild H, Abele M, Frühwald M, Dirksen U, Classen CF, et al. NUT
carcinoma in pediatric patients: Characteristics, therapeutic regimens, and outcomes of
11 cases registered with the German Registry for Rare Pediatric Tumors (STEP).
Pediatric Blood & Cancer. 2024;71(3):e30821.
6.   Lemelle L, Moya-Plana A, Dumont B, Fresneau B, Laprie A, Claude L, et al. NUT
carcinoma in children, adolescents and young adults. Bulletin du Cancer.
2022;109(4):491–504.
7.   French CA. Pathogenesis of NUT Midline Carcinoma. Annual Review of
Pathology: Mechanisms of Disease. 2012;7(1):247–65.
8.   Bauer DE, Mitchell CM, Strait KM, Lathan CS, Stelow EB, Lüer SC, et al.
Clinicopathologic Features and Long-term Outcomes of NUT Midline Carcinoma.
Clinical Cancer Research. 2012;18(20):5773–9.
9.   French CA, Cheng ML, Hanna GJ, DuBois SG, Chau NG, Hann CL, et al. Report
of the first international symposium on NUT carcinoma. Clinical Cancer Research.
2022;28(12):2493–505.
10.   Lewin J, Soria JC, Stathis A, Delord JP, Peters S, Awada A, et al. Phase Ib Trial
With Birabresib, a Small-Molecule Inhibitor of Bromodomain and Extraterminal Proteins,
in Patients With Selected Advanced Solid Tumors. J Clin Oncol. 2018;36(30):3007–14.
11.   Piha-Paul SA, Hann CL, French CA, Cousin S, Braña I, Cassier PA, et al. Phase
1 study of molibresib (GSK525762), a bromodomain and extra-terminal domain protein
inhibitor, in NUT carcinoma and other solid tumors. JNCI Cancer Spectrum.
2020;4(2):pkz093.
12.   Shapiro GI, LoRusso P, Dowlati A, T. Do K, Jacobson CA, Vaishampayan U, et al.
A Phase 1 study of RO6870810, a novel bromodomain and extra-terminal protein
inhibitor, in patients with NUT carcinoma, other solid tumours, or diffuse large B-cell
lymphoma. British journal of cancer. 2021;124(4):744–53.
13.   French CA, Ramirez CL, Kolmakova J, Hickman TT, Cameron MJ, Thyne ME, et
al. BRD–NUT oncoproteins: a family of closely related nuclear proteins that block
epithelial differentiation and maintain the growth of carcinoma cells. Oncogene.
2008;27(15):2237–42.

14.   Nathan P, Hassel JC, Rutkowski P, Baurain J-F, Butler MO, Schlaak M, et al.
Overall Survival Benefit with Tebentafusp in Metastatic Uveal Melanoma. New England
Journal of Medicine. 2021;385(13):1196–206.
15.   D'Angelo SP, Araujo DM, Abdul Razak AR, Agulnik M, Attia S, Blay J-Y, et al.
Afamitresgene autoleucel for advanced synovial sarcoma and myxoid round cell
liposarcoma (SPEARHEAD-1): an international, open-label, phase 2 trial. The Lancet.
2024;403(10435):1460–71.
16.   D'Angelo SP, Furness AJS, Thistlethwaite F, Burgess MA, Riedel RF, Haanen J,
et al. Lete-cel in patients with synovial sarcoma or myxoid/round cell liposarcoma:
Planned interim analysis of the pivotal IGNYTE-ESO trial. American Society of Clinical
Oncology; 2024.
17.   Gazendam AM, Popovic S, Munir S, Parasu N, Wilson D, Ghert M. Synovial
Sarcoma: A Clinical Review. Current Oncology. 2021;28(3):1909–20.
18.   Knight JC, Renwick PJ, Dal Cin P, Van Den Berghe H, Fletcher CDM.
Translocation t(12;16)(q13;p11) in Myxoid Liposarcoma and Round Cell Liposarcoma:
Molecular and Cytogenetic Analysis1. Cancer Research. 1995;55(1):24–7.
19.   Luo J, Bishop JA, DuBois SG, Hanna GJ, Sholl LM, Stelow EB, et al. Hiding in
plain sight: NUT carcinoma is an unrecognized subtype of squamous cell carcinoma of
the lungs and head and neck. Nature Reviews Clinical Oncology. 2025.
20.   Durall RT, Huang J, Wojenski L, Huang Y, Gokhale PC, Leeper BA, et al. The
BRD4–NUT Fusion Alone Drives Malignant Transformation of NUT Carcinoma. Cancer
Research. 2023;83(23):3846–60.
21.   Vensko SP, Olsen K, Bortone D, Smith CC, Chai S, Beckabir W, et al. LENS:
Landscape of Effective Neoantigens Software. Bioinformatics. 2023;39(6):btad322.
22.   Chen S, Zhou Y, Chen Y, Gu J. fastp: an ultra-fast all-in-one FASTQ
preprocessor. Bioinformatics. 2018;34(17):i884–i90.
23.   Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, et al. STAR:
ultrafast universal RNA-seq aligner. Bioinformatics. 2013;29(1):15–21.
24.   Patro R, Duggal G, Love M, Irizarry R, Kingsford C. Salmon: fast and bias-aware
quantification of transcript expression using dual-phase inference. Salmon: fast and
bias-aware quantification of transcript expression using dual-phase inference.10.
25.   Li G, Mahajan S, Ma S, Jeffery ED, Zhang X, Bhattacharjee A, et al. Splicing
neoantigen discovery with SNAF reveals shared targets for cancer immunotherapy.
Science translational medicine. 2024;16(730):eade2886.
26.   Boegel S, Scholtalbers J, Löwer M, Sahin U, Castle JC. In silico HLA typing using
standard RNA-Seq sequence reads. Molecular Typing of Blood Cell Antigens: Springer;
2015. p. 247–58.
27.   Reynisson B, Alvarez B, Paul S, Peters B, Nielsen M. NetMHCpan-4.1 and
NetMHCIIpan-4.0: improved predictions of MHC antigen presentation by concurrent
motif deconvolution and integration of MS MHC eluted ligand data. Nucleic acids
research. 2020;48(W1):W449–W54.
28.   Ewels P, Magnusson M, Lundin S, Käller M. MultiQC: summarize analysis results
for multiple tools and samples in a single report. Bioinformatics. 2016;32(19):3047–8.
29.   Kopylova E, Noé L, Touzet H. SortMeRNA: fast and accurate filtering of
ribosomal RNAs in metatranscriptomic data. Bioinformatics. 2012;28(24):3211–7.

30.   Bruderer R, Bernhardt OM, Gandhi T, Xuan Y, Sondermann J, Schmidt M, et al.
Optimization of Experimental Parameters in Data-Independent Mass Spectrometry
Significantly Increases Depth and Reproducibility of Results *. Molecular & Cellular
Proteomics. 2017;16(12):2296–309.
31.   Tognetti M, Sklodowski K, Müller S, Kamber D, Muntel J, Bruderer R, et al.
Biomarker Candidates for Tumors Identified from Deep-Profiled Plasma Stem
Predominantly from the Low Abundant Area. Journal of Proteome Research.
2022;21(7):1718–35.
32.   Escher C, Reiter L, MacLean B, Ossola R, Herzog F, Chilton J, et al. Using iRT, a
normalized retention time for more targeted measurement of peptides. PROTEOMICS.
2012;12(8):1111–21.
33.   Addis Philip W, Bedke Nicole JOY, Bouard L, Harper S, Liddy N, Mahon T, et al.,
inventors; IMMUNOCORE LTD, assignee. T CELL RECEPTORS. WO patent WO
2018/234319 A1. 2018 2018/06/19.
34.   Froning K, Maguire J, Sereno A, Huang F, Chang S, Weichert K, et al.
Computational stabilization of T cell receptors allows pairing with antibodies to form
bispecifics. Nature Communications. 2020;11(1):2330.
35.   Wu X, Sereno AJ, Huang F, Lewis SM, Lieu RL, Weldon C, et al., editors. Fab-
based bispecific antibody formats with robust biophysical properties and biological
activity. MAbs; 2015: Taylor & Francis.
36.   Kudlacek ST, Metz S, Thiono D, Payne AM, Phan TT, Tian S, et al. Designed,
highly expressing, thermostable dengue virus 2 envelope protein dimers elicit
quaternary epitope antibodies. Science advances. 2021;7(42):eabg4084.
37.   Leonie Alten DM, Sebastian Bunk, Claudia Wagner, Mathias Ferber, inventorT
cell receptors and immune therapy using the same against PRAME positive cancers.
United States of America2018.
38.   Schober K, Müller TR, Gökmen F, Grassmann S, Effenberger M, Poltorak M, et
al. Orthotopic replacement of T-cell receptor α-and β-chains with preservation of near-
physiological T-cell function. Nature biomedical engineering. 2019;3(12):974–84.
39.   Kim SH, Yu SS, Park JS, Robbins PD, An CS, Kim S. Construction of retroviral
vectors with improved safety, gene expression, and versatility. Journal of virology.
1998;72(2):994–1004.
40.   Davar D, Williams A, Lopez J, Olson D, Sato T, Shaw H, et al. 694 Phase 1
safety and efficacy of brenetafusp, a PRAME × CD3 ImmTAC bispecific, in post-
checkpoint cutaneous melanoma (CM). Journal for ImmunoTherapy of Cancer.
2024;12(Suppl 2):A796–A.
41.   Friedman C, De Burgh Williams A, Lopez JS, Ouali K, Middleton MR,
Thistlethwaite F, et al. 750P Phase I safety and efficacy of brenetafusp, a PRAME
× CD3 ImmTAC T cell engager, in platinum resistant ovarian cancer (PROC).
Annals of Oncology. 2024;35:S569–S70.
42.   Beaubier N, Tell R, Lau D, Parsons JR, Bush S, Perera J, et al. Clinical validation
of the tempus xT next-generation targeted oncology sequencing assay. Oncotarget.
2019;10(24):2384–96.
43.   Michuda J, Breschi A, Kapilivsky J, Manghnani K, McCarter C, Hockenberry AJ,
et al. Validation of a Transcriptome-Based Assay for Classifying Cancers of Unknown
Primary Origin. Mol Diagn Ther. 2023;27(4):499–511.

44.   Finkle JD, Boulos H, Driessen TM, Lo C, Blidner RA, Hafez A, et al. Validation of
a liquid biopsy assay with molecular and clinical profiling of circulating tumor DNA. NPJ
Precis Oncol. 2021;5(1):63.
45.   Huang Y, Durall RT, Luong NM, Hertzler HJ, Huang J, Gokhale PC, et al. EZH2
cooperates with BRD4-NUT to drive NUT carcinoma growth by silencing key tumor
suppressor genes. Cancer research. 2023;83(23):3956–73.
46.   Kurtenbach S, Sanchez MI, Kuznetsoff J, Rodriguez DA, Weich N, Dollar JJ, et
al. PRAME induces genomic instability in uveal melanoma. Oncogene. 2024;43(8):555–
65.
47.   Menssen A, Hermeking H. Characterization of the c-MYC-regulated
transcriptome by SAGE: identification and analysis of c-MYC target genes. Proceedings
of the National Academy of Sciences. 2002;99(9):6274–9.
48.   Spel L, Boelens JJ, van der Steen DM, Blokland NJ, van Noesel MM, Molenaar
JJ, et al. Natural killer cells facilitate PRAME-specific T-cell reactivity against
neuroblastoma. Oncotarget. 2015;6(34):35770–81.
49.   Rosskopf S, Leitner J, Paster W, Morton LT, Hagedoorn RS, Steinberger P, et al.
A Jurkat 76 based triple parameter reporter system to evaluate TCR functions and
adoptive T cell strategies. Oncotarget. 2018;9(25):17608–19.
50.   Pentcheva-Hoang T, Torres D, Rodriguez T, Korngold A, Lu A, Crisostomo J, et
al. Inducible MyD88/CD40 (iMC) Enhances Proliferation and Survival of Tumor-Specific
TCR-Modified T Cells and Improves Anti-Tumor Efficacy in Myeloma. Blood.
2016;128(22):4550.
51.   Hamid O, Sato T, Davar D, Callahan MK, Thistlethwaite F, Aljumaily R, et al.
728O Results from phase I dose escalation of IMC-F106C, the first PRAME ×
CD3 ImmTAC bispecific protein in solid tumors. Annals of Oncology. 2022;33:S875.
52.   Bassani-Sternberg M, Bräunlein E, Klar R, Engleitner T, Sinitcyn P, Audehm S, et
al. Direct identification of clinically relevant neoepitopes presented on native human
melanoma tissue by mass spectrometry. Nature Communications. 2016;7(1):13404.
53.   Al-Khadairi G, Decock J. Cancer Testis Antigens and Immunotherapy: Where Do
We Stand in the Targeting of PRAME? Cancers. 2019;11(7):984.
54.   Doolan P, Clynes M, Kennedy S, Mehta JP, Crown J, O'Driscoll L. Prevalence
and prognostic and predictive relevance of PRAME in breast cancer. Breast Cancer
Research and Treatment. 2008;109(2):359–65.
55.   Epping MT, Hart AAM, Glas AM, Krijgsman O, Bernards R. PRAME expression
and clinical outcome of breast cancer. British Journal of Cancer. 2008;99(3):398–403.
56.   Takata K, Chong LC, Ennishi D, Aoki T, Li MY, Thakur A, et al. Tumor-associated
antigen PRAME exhibits dualistic functions that are targetable in diffuse large B cell
lymphoma. The Journal of Clinical Investigation. 2022;132(10).
57.   Alekseyenko AA, Walsh EM, Zee BM, Pakozdi T, Hsi P, Lemieux ME, et al.
Ectopic protein interactions within BRD4–chromatin complexes drive oncogenic
megadomain formation in NUT midline carcinoma. Proceedings of the National
Academy of Sciences. 2017;114(21):E4184–E92.
58.   Rahman S, Sowa ME, Ottinger M, Smith JA, Shi Y, Harper JW, et al. The Brd4
extraterminal domain confers transcription activation independent of pTEFb by
recruiting multiple proteins, including NSD3. Mol Cell Biol. 2011;31(13):2641–52.

59. Fløisand Y, Remberger M, Bigalke I, Josefsen D, Vålerhaugen H, Inderberg EM,
et al. WT1 and PRAME RNA-loaded dendritic cell vaccine as maintenance therapy in de
novo AML after intensive induction chemotherapy. Leukemia. 2023;37(9):1842–9.
60. Bayat Z, Majidi Z, Taherkhani A, Samie L. Engineering a Novel T-Cell
Epitope-Based Vaccine Targeting PRAME in Cancer Immunotherapy. Peptide Science.
2025;117(1):e24393.
61. Chang AY, Dao T, Gejman RS, Jarvis CA, Scott A, Dubrovsky L, et al. A
therapeutic T cell receptor mimic antibody targets tumor-associated PRAME
peptide/HLA-I antigens. The Journal of Clinical Investigation. 2017;127(7):2705–18.