Transcriptomic Analysis of CAD Cell Differentiation

*Carlos A. Cevallos[1], *Anna Leigh White[1], *Brooke A. Fazio[1], Lillian S. Wendt[1], Jasmine W. Feng[1], Dora Posfai[2], April L. Horton[1,3], John M. Warrick[1], Omar A. Quintero-Carmona[1]

[1]Department of Biology, University of Richmond, VA, 23173
[2]Department of Molecular Genetics and Microbiology, Duke University, Durham, NC, 27708
[3]Department of Biology, Bates College, Lewiston, ME, 04240

*Authors contributed equally

ORCID

Cevallos	
White	0009-0004-9081-1268
Fazio	
Wendt	0009-0005-3939-1832
Feng	0009-0009-1778-7915
Posfai	0000-0003-3965-6456
Horton	0000-0001-6810-7435
Warrick	
Quintero-Carmona	0000-0002-9314-1704

Abstract

CAD cells were derived from Cath.a cells, a mouse central nervous system catecholaminergic cell line. Serum-starved CAD cells undergo morphological changes and resemble isolated neurons when observed by microscopy. We carried out an RNAseq transcriptomic analysis to examine differentiated CAD cells for expression signatures related to neuronal functions, identifying ~1900 transcripts whose expression changed with differentiation. Pathview analysis identified ~80 KEGG pathway gene sets that were differentially expressed, including upregulation of at least 13 neuron-related pathways. This dataset can be explored more deeply, allowing further investigation into expression changes relevant to studying neuronal functions in this easy-to-culture model system.

Introduction

Neuron differentiation is difficult to study with primary neuronal cultures due to their delicacy and dependence on supporting glial cells. Most immortalized nerve cell lines also present challenges as proliferating neurons do not differentiate, making them unsuitable as neuronal models. Cath.a-differentiated (CAD) cells, derived from the CNS catecholaminergic cell line Cath.a (Suri, Fung, Tischler, andChikaraishi, 1993), present a possible solution to this issue (Qi, Wang, McMillian, andChikaraishi, 1997). While CAD cells proliferate in serum-containing media, they differentiate, growing neurite-like processes when switched to serum-free media. CAD cells can be maintained for at least 6 weeks in differentiating media, making them an attractive model for studying neuronal processes and differentiation, especially for researchers where primary neurons are not easily accessible, as is the case for researchers at primarily undergraduate institutions.

The Cath.a cell line is derived from a transgenic mouse brain tumor carrying the SV40 T antigen (Tag) oncogene (Suri et al., 1993). Tag stimulates cell proliferation and likely blocks differentiation by keeping Cath.a cells in cycle (Fanning and Knippers, 1992). Like Cath.a, CAD cells express tyrosine hydroxylase (*Th*) and are generally accepted to be catecholaminergic. Unlike Cath.a, the CAD cell line produces no dopamine or norepinephrine despite expressing *Th* and instead exhibits large buildups of L-dihydroxyphenylalanine (L-DOPA), the product of TH enzymatic activity on tyrosine and the precursor to dopamine and norepinephrine, suggesting issues with DOPA decarboxylase activity. Additionally, CAD cells have lost the Tag gene and have gained the ability to differentiate under appropriate conditions (Qi et al., 1997).

Initial characterization by Qi and colleagues demonstrated that CAD expresses neuron-specific proteins such as class III β-tubulin, an isotype of tubulin expressed only in neurons (Sullivan, Havercroft, Machlin, andCleveland, 1986), GAP-43, a cortically associated protein found at high concentration in axons and growth cones (Skene et al., 1986), SNAP-25, a presynaptic protein necessary for vesicle endocytosis (Oyler, Polli, Wilson, andBillingsley, 1991), and synaptotagmin, a synaptic vesicle protein (Matthew, Tsavaler, andReichardt, 1981). Additionally, Qi et al. demonstrated that CAD cells do not express glial fibrillary acidic protein (GFAP), indicative of its neuron-like nature as many immortalized neuron lines express both glial and neuronal proteins (Schubert et al., 1974). Structurally, Qi et al. identified that CAD cells processes contain many of the same elements as neurons, including cytoskeletal elements, the presence of mitochondria, the absence of other organelles, dense-core and clear vesicles, and varicosities. Previous works have looked at changes in expression for subsets of genes related to neuronal functions (Bisig et al., 2009; Lazaroff, Dunlap, andChikaraishi, 1996; Suri et al., 1993), but a comprehensive assessment of gene expression in response to differentiation has not existed for CAD cells.

Results and Discussion

To better understand the neuron-like nature of differentiated CAD cells, we conducted global examination of changes in gene expression resulting from CAD cell differentiation. In addition to examining gene expression changes for individual genes, we also investigated expression changes in gene systems as described by the Kyoto Encyclopedia of Genes and Genomes (KEGG) (Kanehisa and Goto, 2000). We

differentiated CAD cells by growing them in serum-free media, and after 5 days the CAD cells produced long projections that ultrastructurally resembled neurites when visualized by fluorescence microscopy (Figure 1A) or scanning electron microscopy (Figure 1B). These processes contained mitochondria and terminated in structures resembling actin-rich growth cones.

We examined gene expression patterns of differentiated and undifferentiated cells by isolating RNA from cultures of cells and then completing an RNAseq analysis. Principal component analysis (PCA) revealed robust biological replication with undifferentiated and differentiated samples clustering together. The PCA also suggested up to 89% of variation between the transcriptome of samples can be attributed to the differentiated or undifferentiated state of the cells (Figure 1C). Upon differential gene analysis, 1445 genes were found to be significantly upregulated > 2-fold and 451 downregulated > 2-fold in differentiated cells compared to undifferentiated cells (p_{adj}<0.1). Of the 1445 upregulated genes, 1050 had a baseMean value greater than 20. The baseMean values for *Actb* (100,911)*, GAP34* (7203), *Tubb3* (16,487), *Snap25* (5,365), and *Myo10* (267) demonstrate a wide range of expression levels for genes relevant to neuronal function. The same was observed within gene families, such as the synaptotagmins, where baseMean values ranged from 30 (*Synt9*) to ~7000 (*Synt1*). The full gene expression data are included in the extended data.

RNAseq analysis revealed that many of the upregulated genes are implicated in all aspects of neuronal activities. For example, action potential firing has been observed in CAD cells (Wang and Oxford, 2000), and differentiated CAD cells displayed increased expression of a subset of the subunits for voltage-gated sodium channels

such as *Scn1b* (baseMean=46, 6.3-fold increase, $p_{adj}<1.1x10^{-11}$) or *Scn3a* (baseMean=1344, 4.3-fold increase, $p_{adj}<1.2x10^{-31}$), potassium channels such as *Kcnab1* (baseMean=1838, 2-fold increase, $p_{adj}<4.5x10^{-20}$) or *Kcnc1* (baseMean=923, 2.8-fold increase, $p_{adj}<0.001$), calcium channels such as *Cacna1b* (baseMean=1590, 2.2-fold increase, $p_{adj}<3.3x10^{-5}$) or *Cacna2d2* (baseMean=571, 2.7-fold increase, $p_{adj}<0.012$), and sodium/potassium pumps such as *Atp1b2* (baseMean=491, 2.7-fold increase, $p_{adj}<5.6x10^{-10}$) (Bean, 2007; Bezanilla, 2008; Burke and Bender, 2019; Rama, Zbili, andDebanne, 2018).

Expression of tyrosine hydroxylase (*Th*) does not change significantly upon differentiation (baseMean=91, 1.3-fold expression change, $p_{adj}<0.3$). Although, dopa decarboxylase (*Ddc*) expression increases approximately 2-fold (baseMean=60, 2-fold expression increase, $p_{adj}<0.03$) upon differentiation, previous data indicate that this change is insufficient to result in appreciable levels of dopamine expression (Qi et al., 1997). This could be due to posttranscriptional contexts that limit dopamine synthesis, including translational regulation, cellular conditions insufficient for the DDC enzyme to be sufficiently active for dopamine synthesis. *Ddc* is also the second enzyme in the serotonin synthesis pathway, while the first enzyme in the pathway, tryptophan hydroxylase, showed a decrease in expression (*Tph2*, baseMean=146, ~40% decrease in expression, $p_{adj}<4.8x10^{-4}$). Following neurotransmitter synthesis, these small molecules are loaded into synaptic vesicles by membrane transporters such as *Slc17a7/Vglut1* (baseMean=47, 3.8-fold expression increase, $p_{adj}<2.4x10^{-7}$). This glutamate transporter is found in synaptic vesicle membranes (Sanchez-Mendoza et al., 2017), is one of the most highly expressed glutamate transporters in the adult brain

(Wojcik et al., 2004) and is preferentially expressed in neuroblasts over glial cells (Sanchez-Mendoza et al., 2017).

Differentiation also resulted in a significant increase in receptors involved in synaptic function, such as dopamine receptor D1 (*Drd1,* baseMean=55, 4-fold increase, $p_{adj}<3.7\times10^{-8}$), in agreement with previous observations (Pasuit, Li, andKuzhikandathil, 2004). Proteins involved in synapse establishment (Craig and Kang, 2007) were also upregulated, such as the pre-synaptic cell adhesion molecule neurexin 1 (*Nrxn1*, baseMean=289, 2.5-fold increase, $p_{adj}<1.8\times10^{-11}$) the post-synaptic protein neuroligin 2 (*Nlgn2*, baseMean=1848, 1.8-fold increase, $p_{adj}<1.6\times10^{-4}$). Neurotransmitter reuptake attenuates neuronal signaling. *Slc6a17*, is a glutamate transporter present at synaptic junctions in both pre- and post-synaptic sites, particularly in glutamatergic terminals (Iqbal et al., 2015), and was upregulated approximately 8-fold in in differentiated CAD cells (baseMean=18, $p_{adj}<3.4\times10^{-6}$).

In addition to these specific genes involved in neuronal activity, Pathview analysis (Luo and Brouwer, 2013) revealed upregulation for entire pathways as well. Gene enrichment analysis and functional classification by KEGG revealed 83 gene sets that are significantly upregulated and 28 that are downregulated in differentiated cells (q<0.1). The pattern of upregulation for neurotransmitter biosynthetic enzymes, neurotransmitter receptors, and other related machinery varied across the different pathways. The full Pathview analysis is included in the extended data.

Thirteen of the upregulated gene sets were related to neuronal function (Figure 1F). Genes in the glutaminergic (mmu04724, $q=4.0\times10^{-2}$), cholinergic (mmu04725, $q=4.9\times10^{-5}$), GABAergic (mmu04727, $q=1.6\times10^{-3}$), and dopaminergic (mmu04728,

q=2.1×10^{-2}) synapse pathways were all upregulated, as well as the synaptic vesicle

cycle pathway (mmu04721, q=1.1×10^{-2}), suggesting that differentiated CAD cells

possess a subset of the machinery required for some forms of neuronal communication.

Other KEGG pathways related to extracellular signaling relevant to neuronal function

were also upregulated, including neuroactive ligand-receptor interaction (mmu0480,

q=2.3×10^{-2}), retrograde endocannabinoid signaling (mmu04723, q=5.1×10^{-2}), and Notch

signaling pathway (mu04330, q=3.3×10^{-2}). Additional upregulated pathways included

systems related to migration, such as cell adhesion molecules (mmu04514, q=4.9×10^{-5}),

ECM-receptor interaction (mmu04512, q=1.4×10^{-4}), focal adhesion (mmu04510,

q=3.1×10^{-3}) and axon guidance (mmu4360, q=2.3×10^{-3}).

Pathview analysis also revealed the downregulation of pathways less prevalent in

mature neurons, including DNA replication (mmu03030, q=2.1×10^{-25}) and cell cycle

(mmu04110, q=1.5×10^{-23}). Additionally downregulated were several pathways related to

DNA repair including homologous recombination (mmu3440, q=6.2×10^{-14}), mismatch

repair (mmu03430, q=2.7×10^{-9}), nucleotide excision repair (mmu03420, 4.0×10^{-9}), and

non-homologous end-joining (mmu03450, q=4.7×10^{-2}). These findings align with Qi et

al.'s observation that initiating differentiation slowed CAD cell proliferation (Qi et al.,

1997). Taken together, the downregulation of these pathways is indicative of a

population of cells no longer actively dividing, as would be the case for neurons and

other terminally differentiated cells (Anda et al., 2016). However, transcriptionally

active, long-lived cells such as neurons require repair mechanisms to maintain genomic

integrity (Li et al., 2022), so the downregulation of nucleotide excision repair

(mmu03420, q=$4.0x10^{-9}$) and base excision repair (mmu03410, q=$1.3x10^{-5}$) pathways is surprising.

A variety of model systems exist for studying neuronal function at the scale of individual cells, such as isolated primary neurons (Jiang et al., 2006; Mains and Patterson, 1973), induced pluripotent stem cells (Compagnucci, Nizzardo, Corti, Zanni, andBertini, 2014; Dolmetsch and Geschwind, 2011; Inoue, 2010; Swistowski et al., 2010), and a variety of cultured cell lines (Biedler, Helson, andSpengler, 1973; Jones-Villeneuve, McBurney, Rogers, andKalnins, 1982; Pleasure, Page, andLee, 1992; Schubert et al., 1974; Schubert, Heinemann, andKidokoro, 1977). Many of these systems are expensive to maintain and challenging to grow in culture. Additionally, some of the cell lines are not derived purely from the CNS (Wang and Oxford, 2000). The CAD cell transcriptome suggests that there are gene-specific, pathway-specific and global similarities in gene expression between differentiated CAD cells and mature neurons. In conclusion, CAD cells provide an interesting and viable option for researchers to employ a relatively easy-to-culture model system with many of the structural features of neurons. This transcriptomic dataset allows researchers to determine if their particular research question could be investigated using CAD cells, based on examination of expression changes in relevant genes and pathways.

Acknowledgements

We thank the Duke University School of Medicine for the use of the Sequencing and Genomic Technologies Shared Resource, which provided library generation, RNA sequencing services, and the guidance to make initiation of this project possible. Thank you to the other members of the Q-lab for their enthusiasm and support in pursuit of new knowledge. The University of Richmond School of Arts & Sciences funded undergraduate summer research for Anna Leigh White, Brooke Fazio, Lillie Wendt, and Jasmine Feng through "The Richmond Guarantee." This work was also funded by NIH: R15GM119077 to Omar Quintero-Carmona and R15GM119077-S1 (Research Supplements to Promote Diversity in Health-Related Research) which supported Carlos Cevallos' work as a post-baccalaureate researcher.

Author contributions

Carlos A. Cevallos: Funding acquisition, Formal analysis, Investigation, Visualization, Writing – original draft

Anna Leigh White: Formal analysis, Investigation, Visualization, Writing – original draft, Writing – review & editing

Brooke A. Fazio: Formal analysis, Investigation, Writing – original draft

Lillian S. Wendt: Visualization, Writing – original draft, Writing – review & editing

Jasmine W. Feng: Formal analysis, Visualization, Writing – original draft, Writing – review & editing

Dora Posfai: Conceptualization, Data curation, Formal analysis, Investigation, Methodology, Resources, Software, Validation, Visualization, Writing – original draft, Writing – review & editing

April L. Horton: Funding acquisition, Conceptualization, Supervision, Writing – original draft, Writing – review & editing

John M. Warrick: Supervision, Writing – original draft, Writing – review & editing

Omar A. Quintero-Carmona: Conceptualization, Data curation, Formal analysis, Funding acquisition, Investigation, Methodology, Project administration, Resources, Supervision, Validation, Visualization, Writing – original draft, Writing – review & editing

Materials and Methods

CAD cell culturing

Cath.a-differentiated (CAD) cells were a gift from Dona Chikaraishi (Duke University) (Qi et al., 1997), and were maintained in growth media (DMEM/F12 media supplemented with 10% fetal bovine serum, penicillin, and streptomycin) in a humidified atmosphere containing 5% CO_2 at 37°C. To induce differentiation, cells were transitioned to differentiating media (DMEM/F12 media supplemented with penicillin, streptomycin, and ITS supplement (Gemini) containing insulin, transferrin, and selenium).

Scanning electron microscopy

Acid-washed coverslips were coated with $10\mu g/ml$ mouse laminin in PBS for 1 hour and washed with PBS to remove unbound laminin. Cells were plated on coverslips in growth media at a density appropriate for imaging individual cells and allowed to adhere overnight. Samples were fixed in PBS with 2.5% glutaraldehyde for a minimum of 20 minutes. The coverslips were then washed in PBS followed by a series of washes in increasing concentrations of ethanol for 5 minutes at each step (30%, 50%, 70%, 90%, and 100%). The samples were then critical-point dried and sputter-coated as described previously (Applewhite, Davis, Griffis, andQuintero, 2016). Cells were imaged using a JEOL 6360LV scanning electron microscope. For differentiated samples, cells were transitioned to differentiating media for five days prior to processing for imaging.

Fluorescence microscopy and immunostaining sample preparation

Cells were plated on acid-washed, 22mm square #1.5 coverslips which had been previously coated with 10μg/ml mouse laminin in PBS for 1 hour. Coverslips were seeded at a cell density appropriate for imaging and allowed to adhere overnight in growth media. Differentiated samples were transitioned to differentiating media for five days prior to processing for imaging. For simple staining for DNA, and F-actin, samples were fixed in 37°C PBS containing 4% paraformaldehyde for 10 minutes, and then permeabilized in PBS containing 0.5% Triton X-100 for 5 minutes. Samples were then stained in PBS containing 6.6nM ALEXA488 phalloidin and 100nM DAPI for 20 minutes. The coverslips were then washed 4 times in PBS for 5 minutes and mounted on slides in PBS containing 80% glycerol and 0.5% N-propyl gallate, and sealed with clear Sally Hansen Xtreme Wear nail polish.

Fluorescence microscopy image acquisition

Images were acquired using an Olympus IX-83 microscope outfitted with a PLAN APON 60x/1.42NA DIC objective, or a UPlanFL N 40x/1.3NA DIC objective. Fluorescence imaging required an EXFO mixed gas light source, Sutter filter wheels and shutters, and a Sedat Quad filterset (Chroma). Images were acquired using a Hamamatsu ORCA-Flash 4.0 V2 sCMOS camera, and Metamorph imaging software. Images were prepared for publication using Metamorph and FIJI/ImageJ (Schindelin et al., 2012), which included brightness adjustment, contrast adjustment, and pseudocoloration.

RNA isolation

RNA was isolated from T25-sized cultures of CAD cells using the RNeasy Plus kit (Qiagen) according the supplied protocol for purification of RNA from animal cells, with the addition of the Qiashredder step prior to loading the lysate on the column as well as the addition of an on-column DNAse digestion prior to washing and eluting the RNA from the column. RNA was then frozen and sent to the Duke Center for Genomic and Computational Biology for further processing and sequencing.

Library-generation and RNA sequencing

The Duke Sequencing and Genome Technologies Core built stranded mRNA-seq libraries from RNA using Kapa Stranded RNA HyperPrep Kits (Roche). Libraries were pooled in equimolar concentration and were sequenced on the Illumina HiSeq 4000 producing 50 bp single-end reads.

RNA-seq informatic analyses

Briefly, RNA-seq analysis was conducted through the following methods. Quality of FASTQ files from RNA sequencing were assessed using FastQC v.0.11.5 (Andrews, 2010). Adaptor sequences and low-quality reads were trimmed using Trimmomatic v.0.36 (Bolger, Lohse, andUsadel, 2014). Next, alignment was performed using the STAR v.2.5.1a aligner (Dobin et al., 2013) to map reads to the mouse reference transcriptome, GRCm38.p6, to create a count matrix. We used DESeq2 (Love, Huber,

andAnders, 2014) to normalize the read count matrix, and to perform differential

analysis. All statistical analyses and plots were generated using R v3.4.3. Pathview

v.1.20.0 package was used for gene set enrichment analysis utilizing the KEGG

database (Luo and Brouwer, 2013). KEGG pathways were classified manually using

the BRITE functional hierarchies (Aoki-Kinoshita and Kanehisa, 2007) as an initial frame

of reference, and final categorization was based on a consensus among three authors.

Figure 1: CAD cells differentiate into neuron-like cells with serum deprivation.

The cell morphology of CAD cells changes to become more neuron-like after 5 days of growth in serum-free media. Undifferentiated cells form lamellae and sit flat or in clumps, while differentiated cells have long projections that form networks. (A) Phalloidin staining for actin is shown in yellow and DAPI staining for DNA is shown in magenta as visualized by fluorescence microscopy. (B) Representative scanning electron microscopy images. Displayed images are representative of commonly observed phenotypes. All scale bars are 20 μm. (C) Principal component analysis (PCA) indicated that 89% of the variation between RNAseq samples of differentiated and undifferentiated CAD cells could be attributed to differentiation state of the cells. (D) After 5 days of differentiation, 1445 genes were upregulated and 451 genes were downregulated with a more than 2-fold change in gene expression. Red markers outside of the shaded area indicate expression-level changes greater than 2-fold in magnitude with a $p_{adj} \leq 0.1$. (E) Pathview analysis revealed differential expression for a subset of KEGG pathway gene sets compared to undifferentiated CAD cells, including upregulation of 13 gene pathways related to neuronal function (q ≤ 0.1).

References:

Anda, F. C., Madabhushi, R., Rei, D., Meng, J., Graff, J., Durak, O., . . . Tsai, L. H. (2016). Cortical neurons gradually attain a post-mitotic state. *Cell Res, 26*(9), 1033-1047. doi:10.1038/cr.2016.76

Andrews, S. (2010). FastQC: a quality control tool for high throughput sequence data. Retrieved from https://www.bioinformatics.babraham.ac.uk/projects/fastqc/

Aoki-Kinoshita, K. F., & Kanehisa, M. (2007). Gene annotation and pathway mapping in KEGG. *Methods Mol Biol, 396*, 71-91. doi:10.1007/978-1-59745-515-2_6

Applewhite, D. A., Davis, C. A., Griffis, E. R., & Quintero, O. A. (2016). Imaging of the Cytoskeleton Using Live and Fixed Drosophila Tissue Culture Cells. *Methods Mol Biol, 1365*, 83-97. doi:10.1007/978-1-4939-3124-8_4

Bean, B. P. (2007). The action potential in mammalian central neurons. *Nat Rev Neurosci, 8*(6), 451-465. doi:10.1038/nrn2148

Bezanilla, F. (2008). How membrane proteins sense voltage. *Nat Rev Mol Cell Biol, 9*(4), 323-332. doi:10.1038/nrm2376

Biedler, J. L., Helson, L., & Spengler, B. A. (1973). Morphology and growth, tumorigenicity, and cytogenetics of human neuroblastoma cells in continuous culture. *Cancer Res, 33*(11), 2643-2652. Retrieved from https://www.ncbi.nlm.nih.gov/pubmed/4748425

Bisig, C. G., Chesta, M. E., Zampar, G. G., Purro, S. A., Santander, V. S., & Arce, C. A. (2009). Lack of stabilized microtubules as a result of the absence of major maps in CAD cells does not preclude neurite formation. *FEBS J, 276*(23), 7110-7123. doi:10.1111/j.1742-4658.2009.07422.x

Bolger, A. M., Lohse, M., & Usadel, B. (2014). Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics, 30*(15), 2114-2120. doi:10.1093/bioinformatics/btu170

Burke, K. J., Jr., & Bender, K. J. (2019). Modulation of Ion Channels in the Axon: Mechanisms and Function. *Front Cell Neurosci, 13*, 221. doi:10.3389/fncel.2019.00221

Compagnucci, C., Nizzardo, M., Corti, S., Zanni, G., & Bertini, E. (2014). In vitro neurogenesis: development and functional implications of iPSC technology. *Cell Mol Life Sci, 71*(9), 1623-1639. doi:10.1007/s00018-013-1511-1

Craig, A. M., & Kang, Y. (2007). Neurexin–neuroligin signaling in synapse development. *Current opinion in neurobiology, 17*(1), 43. doi:10.1016/j.conb.2007.01.011

Dobin, A., Davis, C. A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., . . . Gingeras, T. R. (2013). STAR: ultrafast universal RNA-seq aligner. *Bioinformatics, 29*(1), 15-21. doi:10.1093/bioinformatics/bts635

Dolmetsch, R., & Geschwind, D. H. (2011). The human brain in a dish: the promise of iPSC-derived neurons. *Cell, 145*(6), 831-834. doi:10.1016/j.cell.2011.05.034

Fanning, E., & Knippers, R. (1992). Structure and function of simian virus 40 large tumor antigen. *Annu Rev Biochem, 61*, 55-85. doi:10.1146/annurev.bi.61.070192.000415

Inoue, H. (2010). Neurodegenerative disease-specific induced pluripotent stem cell research. *Exp Cell Res, 316*(16), 2560-2564. doi:10.1016/j.yexcr.2010.04.022

Iqbal, Z., Willemsen, M. H., Papon, M. A., Musante, L., Benevento, M., Hu, H., . . . van Bokhoven, H. (2015). Homozygous SLC6A17 mutations cause autosomal-recessive intellectual disability with progressive tremor, speech impairment, and behavioral problems. *Am J Hum Genet, 96*(3), 386-396. doi:10.1016/j.ajhg.2015.01.010

Jiang, X. Y., Fu, S. L., Nie, B. M., Li, Y., Lin, L., Yin, L., . . . Xu, X. M. (2006). Methods for isolating highly-enriched embryonic spinal cord neurons: a comparison between enzymatic and mechanical dissociations. *J Neurosci Methods, 158*(1), 13-18. doi:10.1016/j.jneumeth.2006.05.014

Jones-Villeneuve, E. M., McBurney, M. W., Rogers, K. A., & Kalnins, V. I. (1982). Retinoic acid induces embryonal carcinoma cells to differentiate into neurons and glial cells. *J Cell Biol, 94*(2), 253-262. doi:10.1083/jcb.94.2.253

Kanehisa, M., & Goto, S. (2000). KEGG: kyoto encyclopedia of genes and genomes. *Nucleic Acids Res, 28*(1), 27-30. doi:10.1093/nar/28.1.27

Lazaroff, M., Dunlap, K., & Chikaraishi, D. M. (1996). A CNS catecholaminergic cell line expresses voltage-gated currents. *J Membr Biol, 151*(3), 279-291. doi:10.1007/s002329900078

Li, X., Cao, G., Liu, X., Tang, T. S., Guo, C., & Liu, H. (2022). Polymerases and DNA Repair in Neurons: Implications in Neuronal Survival and Neurodegenerative Diseases. *Front Cell Neurosci, 16*, 852002. doi:10.3389/fncel.2022.852002

Love, M. I., Huber, W., & Anders, S. (2014). Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol, 15*(12), 550. doi:10.1186/s13059-014-0550-8

Luo, W., & Brouwer, C. (2013). Pathview: an R/Bioconductor package for pathway-based data integration and visualization. *Bioinformatics, 29*(14), 1830-1831. doi:10.1093/bioinformatics/btt285

Mains, R. E., & Patterson, P. H. (1973). Primary cultures of dissociated sympathetic neurons. I. Establishment of long-term growth in culture and studies of differentiated properties. *J Cell Biol, 59*(2 Pt 1), 329-345. doi:10.1083/jcb.59.2.329

Matthew, W. D., Tsavaler, L., & Reichardt, L. F. (1981). Identification of a synaptic vesicle-specific membrane protein with a wide distribution in neuronal and neurosecretory tissue. *J Cell Biol, 91*(1), 257-269. doi:10.1083/jcb.91.1.257

Oyler, G. A., Polli, J. W., Wilson, M. C., & Billingsley, M. L. (1991). Developmental expression of the 25-kDa synaptosomal-associated protein (SNAP-25) in rat brain. *Proc Natl Acad Sci U S A, 88*(12), 5247-5251. doi:10.1073/pnas.88.12.5247

Pasuit, J. B., Li, Z., & Kuzhikandathil, E. V. (2004). Multi-modal regulation of endogenous D1 dopamine receptor expression and function in the CAD catecholaminergic cell line. *J Neurochem, 89*(6), 1508-1519. doi:10.1111/j.1471-4159.2004.02450.x

Pleasure, S. J., Page, C., & Lee, V. M. (1992). Pure, postmitotic, polarized human neurons derived from NTera 2 cells provide a system for expressing exogenous proteins in terminally differentiated neurons. *J Neurosci, 12*(5), 1802-1815. doi:10.1523/JNEUROSCI.12-05-01802.1992

Qi, Y., Wang, J. K., McMillian, M., & Chikaraishi, D. M. (1997). Characterization of a CNS cell line, CAD, in which morphological differentiation is initiated by serum deprivation. *J Neurosci, 17*(4), 1217-1225. Retrieved from http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=9006967

Rama, S., Zbili, M., & Debanne, D. (2018). Signal propagation along the axon. *Curr Opin Neurobiol, 51*, 37-44. doi:10.1016/j.conb.2018.02.017

Sanchez-Mendoza, E. H., Bellver-Landete, V., Arce, C., Doeppner, T. R., Hermann, D. M., & Oset-Gasque, M. J. (2017). Vesicular glutamate transporters play a role in neuronal differentiation of cultured SVZ-derived neural precursor cells. *PLoS One, 12*(5), e0177069. doi:10.1371/journal.pone.0177069

Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., . . . Cardona, A. (2012). Fiji: an open-source platform for biological-image analysis. *Nat Methods, 9*(7), 676-682. doi:10.1038/nmeth.2019

Schubert, D., Heinemann, S., Carlisle, W., Tarikas, H., Kimes, B., Patrick, J., . . . Brandt, B. L. (1974). Clonal cell lines from the rat central nervous system. *Nature, 249*(454), 224-227. doi:10.1038/249224a0

Schubert, D., Heinemann, S., & Kidokoro, Y. (1977). Cholinergic metabolism and synapse formation by a rat nerve cell line. *Proc Natl Acad Sci U S A, 74*(6), 2579-2583. doi:10.1073/pnas.74.6.2579

Skene, J. H., Jacobson, R. D., Snipes, G. J., McGuire, C. B., Norden, J. J., & Freeman, J. A. (1986). A protein induced during nerve growth (GAP-43) is a major component of growth-cone membranes. *Science, 233*(4765), 783-786. doi:10.1126/science.3738509

Sullivan, K. F., Havercroft, J. C., Machlin, P. S., & Cleveland, D. W. (1986). Sequence and expression of the chicken beta 5- and beta 4-tubulin genes define a pair of divergent beta-tubulins with complementary patterns of expression. *Mol Cell Biol, 6*(12), 4409-4418. doi:10.1128/mcb.6.12.4409-4418.1986

Suri, C., Fung, B. P., Tischler, A. S., & Chikaraishi, D. M. (1993). Catecholaminergic cell lines from the brain and adrenal glands of tyrosine hydroxylase-SV40 T antigen transgenic mice. *J Neurosci, 13*(3), 1280-1291. doi:10.1523/JNEUROSCI.13-03-01280.1993

Swistowski, A., Peng, J., Liu, Q., Mali, P., Rao, M. S., Cheng, L., & Zeng, X. (2010). Efficient generation of functional dopaminergic neurons from human induced pluripotent stem cells under defined conditions. *Stem Cells, 28*(10), 1893-1904. doi:10.1002/stem.499

Wang, H., & Oxford, G. S. (2000). Voltage-dependent ion channels in CAD cells: A catecholaminergic neuronal line that exhibits inducible differentiation. *J Neurophysiol, 84*(6), 2888-2895. doi:10.1152/jn.2000.84.6.2888

Wojcik, S. M., Rhee, J. S., Herzog, E., Sigler, A., Jahn, R., Takamori, S., . . . Rosenmund, C. (2004). An essential role for vesicular glutamate transporter 1 (VGLUT1) in postnatal development and control of quantal size. *Proc Natl Acad Sci U S A, 101*(18), 7158-7163. doi:10.1073/pnas.0401764101