Revealing Multidimensional Associations in Life History Traits Across Four Animal Taxa

Yang Zhang[1], Yunshu Zhang[1], Chunhui Hao[1,2*]

[1] School of Life Sciences, Nanjing University, Nanjing 210023, China

[2] School of Ecological and Environmental Sciences, East China Normal University, Shanghai 200241, China

[*]Corresponding author:

Chunhui Hao

School of Life Sciences, Nanjing University

163 Xianlin Avenue, Nanjing 210023, China

Email: haochunhui1996@gmail.com

Abstract

Trade-offs—negative associations among life history traits—are fundamental to understanding life history evolution. While traditional studies have focused mainly on pairwise trait relationships, organisms allocate finite energy across multiple processes simultaneously, necessitating a multivariate perspective. Here, by integrating both bivariate and multivariate phylogenetic frameworks, we investigated the complex relationships among six key life history traits (adult body size, offspring size, development time, adult lifespan, clutch size, and clutch frequency) across species of mammals, birds, reptiles, and insects. We demonstrated that multivariate frameworks significantly alter inferred trait relationships across species. For instance, a direct relationship between offspring size and adult lifespan becomes indirect when additional traits are considered. Moreover, while some multivariate patterns are consistent across taxa, others are taxon-specific. Our findings suggest that life history variations emerge from the interplay of multiple traits, emphasizing the need to view life history evolution through the lens of interconnected ecological and evolutionary processes.

Keywords: life history, trade-off, multivariate correlation, phylogenetic comparative method, phylogenetic path analysis

Introduction

Life history traits are measurable characteristics of an organism that determine its survival, growth, and reproductive success, such as body size, developmental duration and overall lifespan. These traits vary substantially across taxa, reflecting the diverse strategies used by different taxa in response to varying evolutionary pressures[1–4]. For example, mammals and birds typically exhibit lower reproductive rates and higher levels of parental investment, enhancing offspring survival in relatively stable environments. In contrast, fish and amphibians tend to have higher reproductive rates and shorter growth periods, strategies that maximize reproductive output and population survival in unstable environments[1–4].

Researchers have developed life history theory as a conceptual framework to explain the evolution of distinct life history strategies adopted by different organisms. Central to this framework is the idea of life history trade-offs, which arise because organisms have limited energy, time and nutrient that must be allocated among competing functions such as growth, reproduction, and maintenance[5–7]. Consequently, increased investment in one function necessarily reduces the resources available for another, leading to trade-offs among life history traits[5–7]. For instance, the trade-off between survival and reproduction is evident in organisms like annual herbs and insects, which allocate more resources to reproduction at the cost of their long-time survival, ensuring reproductive success in environments with high mortality risks[4,8,9].

Although numerous empirical studies have provided evidence for theoretical trade-offs among life history traits, examples have also been identified that do not conform to theoretical expectations. For example, the expected trade-off between body size and clutch size is sometimes absent or even reversed in amphibians or reptiles, where larger individuals may achieve higher reproductive outputs simultaneously[3,10]. Moreover, the relationship between specific life history traits can vary across taxa. For example, a negative correlation between development and reproduction has been consistently

observed in mammals and birds[2,3], yet this relationship was not always found in certain plants and insects[9,11].

Why was the evidence on the life history trade-offs mixed? One potential reason is that most of the current studies have only examined bivariate correlations between pairs of life history traits, while overlooking the possible multivariate relationships among these traits[12,13]. For instance, in *Drosophila*, the challenge of ephemeral food resources can simultaneously enhance stress resistance, metabolic reserves, and somatic maintenance at the expense of reduced fecundity. Additionally, abundant feeding and breeding opportunities can favour increased larval competitive ability and high fecundity at the cost of reduced size and maintenance[14,15]. Such complex multidimensional trade-offs could arise because organisms have a finite energy budget that must be divided among multiple life history processes simultaneously, rather than just two traits at a time[16–18]. Consequently, correlations between pairwise traits can be obscured by additional traits, leading to discrepancies between theoretical predictions and empirical findings[12,13,19].

To address these discrepancies, we carried out phylogenetic comparative analyses to investigate the relationships among six key life history traits, including adult body size (BS), offspring size (CS), development time (DT), adult lifespan (AL), clutch size (CS) and clutch frequency (CF) across a diverse range of animal species. Our work focused on two central questions: (1) Do multivariate relationships among life history traits alter the bivariate relationships observed between pairs of traits? (2) Are these multivariate relationships consistent across major animal taxa, including mammals, birds, reptiles, and insects? To answer these questions, we first used phylogenetic generalized least squares (PGLS) to investigate bivariate relationships between pairs of traits across species while accounting for phylogenetic relatedness. Next, we applied phylogenetic path analysis (PPA) to test multiple hypotheses regarding the multivariate relationships among these traits and identify the best-supported model (see Methods, Figure 1). By

88   comparing PGLS and PPA results, we differentiated direct pairwise associations from

89   indirect associations mediated by other traits, providing deeper insights into the

90   evolution of life history strategies in animals.

Methods

Data collection

(1) Life history traits

We collected species-level life history traits data from two key literature sources[4,20],

which aggregated data from numerous studies on four major animal taxa:

mammals[2,3,21], birds[2,3,22,23], reptiles[3,10], and insects[e.g., 24–26]. To enable

cross-taxon comparisons, we focused on six key traits that are consistently available for

all four animal taxa: (1) Adult body size: measured as adult body mass for vertebrate

species, and body length for insects; (2) Offspring size: defined as offspring body mass

for vertebrate species and body length for insect species; (3) Development time:

measured as the time from birth to sexual maturity for vertebrates, and from hatching

to adult eclosion for insects; (4) Adult lifespan: defined as the time from sexual maturity

to death for vertebrates, and from adult eclosion to death for insects; (5) Clutch size:

defined as the number of eggs per clutch or litter across all species, and (6) Clutch

frequency: measured as the number of clutches or litters per year for vertebrate species,

and per day for insect species, reflecting their typically shorter lifespan. Due to the

absence of direct clutch frequency data for insect species, we estimated it indirectly via

dividing the reported lifetime fecundity by clutch size, ensuring consistency across

datasets.

For vertebrate taxa (i.e., mammals, birds and reptiles), we included only species with

complete data for all six life history traits, resulting in a final dataset comprising 335

mammalian, 148 avian, and 417 reptilian species (Table S1). In contrast, for insect,

restricting the dataset to only include insect species with complete data for all six traits

would result in a small, taxonomically unrepresentative sample. To address this, we

applied a phylogenetic imputation method implemented in the R package *Rphylopars* (version 0.3.10) to impute the missing data while accounting for the phylogeny[27]. Specifically, we only included insect species with real data available for at least four of the six life history traits, and then imputed the remaining missing values. This approach resulted in a final dataset of 205 insect species, ensuring a complete and taxonomically representative insect dataset for subsequent analyses. Furthermore, all measurements of life history traits were natural log-transformed to meet the assumptions of statistical methods. The complete dataset used in this study is provided in supplementary material (Table S1).

(2) Phylogeny

We pruned published phylogenetic trees for each animal taxon to create phylogenies that reflect the shared evolutionary histories of the species included in our dataset. For mammals, we obtained the phylogenetic tree from Bakewell et al. [4], which was originally derived from Fritz et al. [28], and pruned it using "*keep.tip*" function from the R package "*ape*" (version 5.8) [29] to include only the species present in our dataset. For reptiles, Bakewell et al. [4] merged three existing molecular phylogenies[30–32], which we similarly pruned. The avian phylogeny used in this study was pruned based on a most recent avian molecular phylogeny inferring from genomic data[33]. For insects, the phylogeny used in this study was based on a most comprehensive insect molecular phylogeny available, which is constructed hierarchically using transcriptome, mitogenome data and species-rich DNA data[34]. The resulting phylogenetic trees for each taxon are provided as dendrograms in the supplementary material (Figure S1).

Statistical analyses

(1) Phylogenetic generalized least squares

We used phylogenetic generalized least squares (PGLS) [35] to examine the pairwise relationships between every pair of life history traits across species within each taxon. PGLS is a statistic approach that fits a linear model while accounting for the

phylogenetic non-independence among species, and can incorporate different

evolutionary models (e.g., Pagel's λ, Brownian motion or early burst) to define the

distribution of the residuals from the linear model[36]. We used Pagel's λ model in our

main analyses. Pagel's λ, ranging from 0 to 1, measures the phylogenetic signal in the

data: λ = 0 implies that the traits are independent of the phylogeny (i.e., there is no

phylogenetic signal), whereas λ = 1 suggests that the traits perfectly follow the

phylogenetic relationships as expected under a Brownian motion model of

evolution[37]. We conducted PGLS using R package "*caper*" (version 1.0.3) [38],

which estimates λ by automatically determining the value that maximizes the likelihood

of the fit of the data to the phylogenetic tree and regression model, ensuring optimal

branch length transformations given the dataset. For each PGLS model, we derived the

regression coefficients and associated statistical outputs, including t-values, *P*-values,

and R-squared (R^2) statistics to assess the significance, explanatory power of the

regression and the overall fit of the models to the data (Table S2). We initiated our

analysis to test all pairwise relationships (Figure 1) between life history traits in

mammals. Then, we extended our investigation by conducting separate PGLS models

to examine the correlations between life history traits across species within other taxa,

including birds, reptiles and insects.

(2) Phylogenetic path analyses

We employed phylogenetic path analysis (PPA) [39,40] to examine the multivariate

relationships, including both direct and indirect associations, among diverse life history

traits across species. PPA is an extension of structural equation modelling (SEM)

approach that can test the goodness-of-fit of the data to different hypotheses regarding

the associations among multiple variables while incorporating phylogenetic regression

methods, such as PGLS, to account for phylogenetic non-independence among species

[41,42].

We carried out the PPA by testing previously proposed hypotheses. Previous studies

have documented several robust associations, including: positive associations between
adult body size and offspring size or lifespan[2,8,43], and between adult body size,
offspring size and development time[14,44,45]. Additionally, a trade-off between clutch
size and offspring size or development time has been consistently reported[46–48]. In
contrast, some associations remain contentious, with inconsistent findings in literature,
such as those between body size and clutch size, and between development time and
lifespan[2,4,49]. To simplify hypothesis testing, we first incorporated the well-
established associations into all candidate models as fixed components (Figure 1, Table
S3). Next, for the associations that remain debated or unresolved, we systematically
included or excluded them across various candidate models, generating a list of distinct
hypotheses for evaluation (Figure 1; Table S3).

Figure 1. Conceptual diagram illustrating the hypotheses of our study, highlighting the hypothetical
relationships among life history traits. Blue arrows indicate hypothesized positive direct
associations between life history traits, while red arrows denote negative correlations. Solid lines
represent well-established associations consistently supported by previous studies, and were
therefore included in every candidate models in phylogenetic path analysis. In contrast, dotted lines
denote associations that were not consistently observed, reflecting mixed support across different
studies. References for the well-established associations are provided alongside the corresponding
arrows.

We initially conducted PPA on mammalian life history traits utilizing the R package "*phylopath*" (version 1.3.0) [50] while assuming Pagel's λ model for the trait evolution. We tested the goodness-of-fit for a list of 511 candidate models, each represented as a formula in Tables S3. Model fitting was assessed via the Fisher's C statistic, with a proposed model deemed a good fit if the C statistic was non-significant ($P > 0.05$) [41,42]. Moreover, to rank the candidate models and identify the best-supported model, we calculated the C statistic information criterion (CICc) for each model[39,40], analogous to the Akaike information criterion (AIC). The best-supported model was the one with a non-significant C statistic ($P > 0.05$) and the lowest CICc value, when the difference in CICc (ΔCICc) between this model and the next best model did not exceed 2[42]. We then conducted PPA for the other animal taxa following the same protocol, enabling comparisons with the patterns observed in mammals.

Visualization

Phylogenetic relationships among species were visualized using dendrograms generated with the *ggtree* package (version 3.13.0.1) [51]. We mapped the life history trait values for each species onto the tips of the phylogenetic trees, with color gradients representing trait value variation. In addition, we created graphical representations of the relationships among multiple life history traits when reporting the results of PGLS models and PPA. All visualizations were produced in R (version 4.4.1) [52] using the *ggplot2* package (version 3.5.1) [53].

Sensitivity Analyses

We used different evolutionary models in phylogenetic imputation method, PGLS, and PPA to check whether our findings are robust to alternative evolutionary models. The alternative evolutionary models included Brownian motion (BM) and early burst (EB). The results of these sensitivity analyses are presented in supplementary materials (Figures S2-6 and Tables S2 and S5).

Results

PGLS revealed bivariate associations among life history traits in mammals.

We formulated hypotheses regarding the relationships among life history traits, drawing from previous studies (see in Methods, Figure 1). Using PGLS while assuming a Pagel's λ model, we tested these hypotheses by examining the bivariate associations between pairs of life history traits across species while accounting for phylogenetic non-independence (Figure 1). Our main analysis focused on mammals, compiling data on six key life history traits across 335 mammalian species: adult body size (BS), offspring size (OS), development time (DT), adult lifespan (AL), clutch size (CS), and clutch frequency (CF) (Figure 2a).

Figure 2. Phylogenetic correlations between life history traits in mammalian species. (a) Phylogenetic tree of 335 mammalian species, with heatmap rings displaying life history trait values.

Ring colours range from yellow (low value) to dark purple (high value). All measurements of life history traits were natural log-transformed and standardized. The outermost rings denote the major mammalian orders. Strong covariation is observed among life history traits across species, for example, between body size, offspring size, development time and lifespan. (b) The correlations among pairs of life history traits across 335 mammalian species derived from PGLS models. Each dot represents a species, with its coordinates corresponding to the values of two life history traits. The best-fit regression line represents the overall trend.

First, we validated all previously well-established associations (Figure 1). Specifically, we confirmed the positive correlations between adult body size and both offspring size and lifespan (PGLS, OS ~ BS, $\beta \pm SE = 0.87 \pm 0.014$, $P < 0.001$; AL ~ BS, $\beta \pm SE = 0.158 \pm 0.015$, $P < 0.001$; Figure 2b; Tables 1 and S2). Additionally, we observed a positive correlation between adult size or offspring size and development time, while offspring size was negatively correlated with clutch size (PGLS, DT ~ BS, $\beta \pm SE = 0.186 \pm 0.022$, $P < 0.001$; DT ~ OS, $\beta \pm SE = 0.207 \pm 0.024$, $P < 0.001$; CS ~ OS, $\beta \pm SE = -0.089 \pm 0.014$, $P < 0.001$; Figure 2b; Tables 1 and S2). We also identified a significant negative association between clutch size and development time (PGLS, DT ~ CS, $\beta \pm SE = -0.448 \pm 0.107$, $P < 0.001$; Figure 2b; Tables 1 and S2). These findings are consistent with previously reported trends[2,8,14,45,46,48], reinforcing the robustness of these correlations in mammalian species (Figure 1).

Next, we examined associations that have received mixed support in previous studies (Figure 1). Our analysis confirmed some widely recognized associations among life history traits in mammals. Specifically, clutch frequency showed significant negative associations with adult body size, offspring size, development time, and lifespan (PGLS, CF ~ BS, $\beta \pm SE = -0.103 \pm 0.014$, $P < 0.001$; CF ~ OS, $\beta \pm SE = -0.112 \pm 0.016$, $P < 0.001$; CF ~ DT, $\beta \pm SE = -0.222 \pm 0.031$, $P < 0.001$; AL ~ CF, $\beta \pm SE = -0.202 \pm 0.065$, $P = 0.002$; Figure 2b; Tables 1 and S2). Additionally, we found lifespan had a positive correlation with offspring size or development time, as well as a negative association

with clutch size (PGLS, AL ~ OS, $\beta \pm$ SE = 0.174 ± 0.016, P < 0.001; AL ~ DT, $\beta \pm$

SE = 0.183 ± 0.038, P < 0.001; AL ~ CS, $\beta \pm$ SE = -0.333 ± 0.077, P < 0.001; Figure

2b; Tables 1 and S2). In contrast, we also revealed some correlations that deviated from

existing hypothetical associations. While some studies have reported a positive

relationship between adult body size and clutch size[3,10,54], our results indicated a

trade-off between the two traits in mammals (PGLS, CS ~ BS, $\beta \pm$ SE = -0.063 ± 0.013,

P < 0.001; Figure 2b; Tables 1 and S2). Finally, we found no significant correlation

between clutch size and clutch frequency (PGLS, P = 0.156; Figure 2b; Tables 1 and

S2).

Discrepancies between PGLS and PPA indicated indirect associations among life

history traits.

We observed inconsistencies between the results of PGLS and the hypothetical

associations or previous findings, which we attribute to a key limitation in earlier

studies: Prior studies primarily focused only on pairwise relationships between life

history traits, often overlooking the complex multivariate associations among multiple

traits. In our dataset, we observed that life history traits such as body size, offspring

size, development time and lifespan might strongly covary, suggesting that some

observed pairwise correlations may result from confounding effects of additional

variables (Figure 2a). Therefore, to examine whether multivariate associations would

change the bivariate correlations, we used phylogenetic path analysis (PPA) to study

life history evolution across 335 mammalian species, a method that simultaneously tests

both direct and indirect relationships among traits while accounting for phylogenetic

relationships[39,40].

We formulated 511 models to explore potential relationships among six life history

traits, and conducted PPA to select the best-support model (Table S3, see Method for

details). In the best-support model, we found that some direct correlations between pairs

of traits were consistently observed in both PGLS and PPA models. Specifically, adult

body size showed positive direct correlations with offspring size, development time, and lifespan (PPA, OS ~ BS: β ± SE = 0.942 ± 0.015; DT ~ BS: β ± SE = 0.303 ± 0.191; AL ~ BS: β ± SE = 0.506 ± 0.062; Figure 3a; Table 1). Moreover, a significant positive direct correlation was also identified between between development time and both offspring size and lifespan (PPA, DT ~ OS: β ± SE = 0.169 ± 0.2; AL ~ DT: β ± SE = 0.08 ± 0.052; Figure 3a; Table 1). In contrast, a negative direct correlation was found between offspring size and clutch size (PPA, CS ~ OS: β ± SE = -0.721 ± 0.158; Figure 3a; Table 1). Both adult body size and development time were negatively associated with clutch frequency (PPA, CF ~ BS: β ± SE = -0.349 ± 0.068; CF ~ DT: β ± SE = -0.28 ± 0.055; Figure 3a; Table 1), Additionally, clutch size was negatively correlated to both development time and lifespan (PPA, DT ~ CS: β ± SE = -0.128 ± 0.058; AL ~ CS: β ± SE = -0.113 ± 0.052; Figure 3a; Table 1). We found no direct correlations between clutch size and clutch frequency in both PGLS models and PPA models (PPA, β = 0; Figure 3a; Table 1).

Figure 3. Best-supported models in phylogenetic path analysis using Pagel's λ model for (a) mammals, (b) birds, (c) reptiles and (d) insects. Blue arrows represent positive correlations, while

red arrows represent negative correlations, with their width corresponding to the magnitude of the

standardized regression coefficients. Exact coefficient values are provided alongside the arrows.

Notably, some correlations identified in PGLS were not retained as direct associations in PPA, but were instead mediated by third variables, making them indirect associations in PPA models (Figure 3a; Table 1). Specifically, the negative association between offspring size and clutch frequency identified in PGLS was mediated by adult body size and/or development time in PPA (Figure 3a; Table 1). Similarly, the negative correlation between clutch frequency and lifespan was mediated through adult body size and/or development time, and the positive association between offspring size and lifespan was mediated by adult body size, development time and/or clutch size (Figure 3a; Table 1).

Table 1. Comparison of PGLS and PPA results for pairs of life history traits in mammals. The first column lists the trait pairs, and shaded rows indicate associations that were found discrepancies between the two models.

Model	PGLS (Coef ± SE)	PPA (Coef ± SE)
Offspring Size ~ Body Size	$0.87 \pm 0.014^*$	0.942 ± 0.015
Development Time ~ Body Size	$0.186 \pm 0.022^*$	0.303 ± 0.191
Adult Lifespan ~ Body Size	$0.158 \pm 0.015^*$	0.506 ± 0.062
Clutch Size ~ Body Size	$-0.063 \pm 0.013^*$	0.337 ± 0.156
Clutch Frequency ~ Body Size	$-0.103 \pm 0.014^*$	-0.349 ± 0.068
Development Time ~ Offspring Size	$0.207 \pm 0.024^*$	0.169 ± 0.2
Adult Lifespan ~ Offspring Size	$0.174 \pm 0.016^*$	0
Clutch Size ~ Offspring Size	$-0.089 \pm 0.014^*$	-0.721 ± 0.158
Clutch Frequency ~ Offspring Size	$-0.112 \pm 0.016^*$	0
Adult Lifespan ~ Development Time	$0.183 \pm 0.038^*$	0.08 ± 0.052
Clutch Frequency ~ Development Time	$-0.222 \pm 0.031^*$	-0.28 ± 0.055
Development Time ~ Clutch Size	$-0.448 \pm 0.107^*$	-0.128 ± 0.058
Adult Lifespan ~ Clutch Size	$-0.333 \pm 0.077^*$	-0.113 ± 0.052
Clutch Frequency ~ Clutch Size	0.095 ± 0.067	0
Adult Lifespan ~ Clutch Frequency	$-0.202 \pm 0.065^*$	0

Lastly, PPA revealed a reversal in the direction of one correlation: the association

between adult body size and clutch size, which was negative in PGLS, became positive

in PPA (PPA, CS ~ BS, $\beta \pm SE = 0.337 \pm 0.156$; Figure 3a; Table 1).

Patterns found in mammals were partially consistent in other animal taxa.

We then extended the analyses to other animal taxa, including reptiles, birds and insects,

to understand whether the patterns observed in mammals were consistent across these

groups.

In PGLS models, many associations were consistent across animal taxa, while some

notable exceptions (Figures S2-4; Table S2). For example, adult body size was

negatively correlated with clutch size in mammals (PGLS, CS ~ BS, $\beta \pm SE = -0.063 \pm$

0.013, $P < 0.001$; Figure 2b; Tables 1 and S2). In contrast, reptiles and insects exhibited

a significant positive association between these traits, while no relationship was

detected in birds (PGLS, reptiles: $\beta \pm SE = 0.242 \pm 0.018$, $P < 0.001$; Figure S3b; insects:

$\beta \pm SE = 0.374 \pm 0.128$, $P = 0.004$; Figure S4b; birds: $P = 0.521$; Figure S2b; Table S2).

In PPA models, we first identified several consistent direct correlations between pairs

of life history traits across different animal taxa (Figure 3). Specifically, adult body size

was positively correlated with both offspring size and development time across all four

groups, as well as between offspring size and development time (Figure 3; Table S5).

Despite these common trends, we observed taxon-specific variations. In mammals,

birds, and reptiles, we consistently observed positive direct associations between body

size and lifespan (Figures 3a-c; Table S5). However, insects exhibited the opposite

trends, with negative correlations (Figure 3d; Table S5). Similarly, while both clutch

size and clutch frequency were negatively correlated with development time in

mammals, birds and reptiles (Figures 3a-c; Table S5), insects displayed a positive

correlation between clutch size and development time, and no significant direct

correlation between clutch frequency and development time (Figure 3d; Table S5).

In addition, while positive direct correlations between adult size and clutch size were evident in mammals, reptiles and insects (Figures 3a, 3c and 3d; Table S5), no significant direct correlation was found in birds (Figure 3b; Table S5). Similarly, except birds (Figure 3b; Table S5), a negative correlation was consistently identified between offspring size and clutch size across most taxa (Figures 3a, 3c and 3d; Table S5). Additionally, no direct association was identified between clutch size and clutch frequency in mammals and birds, nor between clutch frequency and lifespan in mammals and reptiles (Figure 3; Table S5).

Notably, some associations observed in mammals reversed in other taxa (Figure 3; Tables S5). For example, the correlation between clutch size and lifespan was negative in mammals and reptiles (Figures 3a and 3c; Table S5), but became positive in birds and insects (Figures 3b and 3d; Table S5).

Sensitive analyses

We evaluated the robustness of the PGLS and PPA models across four taxa using two alternative evolutionary models, including Brownian motion (BM) and early burst (EB). The results derived from different evolutionary models displayed only minimal differences (Figures S5 and S6; Tables S2 and S5). For example, in mammals, while model with Pagel's λ did not detect a significant relationship between clutch size and clutch frequency (PGLS, $P = 0.156$; Figure 2b; Table S2), a negative correlation was demonstrated in the PGLS models with both BM and EB (PGLS, CF ~ CS, $\beta \pm SE = 0.132 \pm 0.06$, $P = 0.029$; Table S2). Furthermore, in the PPA models, the Pagel's λ model showed positive direct associations between body size and clutch size (PPA, CS ~BS, $\beta \pm SE = 0.337 \pm 0.156$; Figure 3a; Table S5), as well as between development time and lifespan (PPA, AL ~ DT, $\beta \pm SE = 0.08 \pm 0.052$; Figure 3a; Table S5). However, no direct relationships were observed between these traits in the PPA models with both BM and EB (PPA, $\beta = 0$; Figures S5a and S6a; Tables S5).

Discussion

In this study, we investigated the species-wise relationships among six key life history traits (e.g., adult body size, development time, and clutch size) across mammals, birds, reptiles, and insects. Using phylogenetic generalized least squares (PGLS) and phylogenetic path analysis (PPA), we examined both bivariate and multivariate relationships among these traits, uncovered patterns that traditional bivariate approaches could not capture alone, providing deeper insights into the evolution of animal life histories.

Traditional studies typically rely on bivariate analyses to assess relationships among life history traits[5,14,55,56]. While informative, such analyses may overlook confounding effects from additional variables[12,13,19]. Although multivariate methods like principal component analysis (PCA) have been used to reduce dimensionality[57–60], they do not allow for hypotheses testing or the disentanglement of more nuanced relationships among traits[13]. By employing PPA, we extended beyond these limitations to disentangle both direct and indirect associations among traits under a hypothesis-testing framework. This allowed us to reveal patterns that bivariate analyses alone cannot capture. For instance, although previous work supported a trade-off between adult body size and clutch size in mammals[2–4], our PPA revealed a direct positive correlation between these traits (Figure 3a; Table 1). Similarly, while earlier studies proposed a negative association between offspring size and clutch frequency[2–4], we found no direct relationship between these traits (Figure 3a; Table 1).

More importantly, these findings provide novel insights into the complex nature of life history evolution, offering implications that extend beyond previous theories. Traditional theories have framed organisms along a continuum from "fast" to "slow" life histories[6,7,56]. "Fast" life histories, characterized by rapid growth, early reproduction, and short lifespans, are typically associated with unstable environments.

In contrast, "slow" life histories involve slower growth, delayed reproduction, and longer lifespans, often reflecting adaptation to more stable conditions[61,62]. While this framework has been instrumental in explaining life history variation among organisms, accumulating evidence suggested that life history strategies are more complex than the traditional "fast-slow" model implies[12,13,19]. As Stott et al. [13] argue, this framework oversimplifies the intricate dynamics of life history evolution, and fails to account for the full diversity of strategies that exist beyond this continuum. Our findings support this perspective by demonstrating that life history evolution is shaped by complex, multidimensional relationships rather than simple bivariate correlations. We showed that some associations, such as between adult body size and clutch size in mammals, was actually mediated by other traits, such as development time, underscoring the importance of accounting for additional mediating variables when examining life history evolution. (Figure 3a).

Additionally, studies have also uncovered the genetic architecture underlying the multidimensional nature of life history trade-offs[19]. Genes with pleiotropic effects influence multiple traits simultaneously by regulating interconnected gene networks. Often acting as key transcription factors or components of signaling pathways, these genes coordinate multiple processes like survival, growth, and fecundity[19]. For instance, In *Glanville fritillary* butterflies, the pleiotropic gene *Sdhd* modulates critical metabolic pathways that enhances adult flight performance and dispersal capacity while concurrently imposing costs such as reduced metabolic rate and delayed lavral growth[63].

Furthermore, we observed taxon-specific life history strategies across different animal taxa. For example, in vertebrates, there is a positive relationship between adult body size and lifespan, and a negative relationship between clutch size and development time, but insects exhibit the opposite patterns on both associations (Figure 3). Such differences likely arise from the distinctions in the environments in which these groups

inhabit. Mammals and birds, which often inhabit relative stable and predictable environments, tend to invest in fewer offspring with extensive parental care and longer developmental periods[5,6,64,65]. Larger organisms in these groups generally exhibit slower metabolic rates, reduced reproductive rates, and prolonged parental investment, all of which contribute to longer lifespans and more gradual life histories[64–66]. This strategy allows for greater resource allocation per offspring, enhancing survival rates in stable habitats[61,62,67]. In contrast , insects often inhabit unpredictable environments with fluctuating resource availability[66,68]. To cope with these conditions, they adopt a high-fecundity strategy, producing large clutch sizes at the expense of individual offspring quality[69–71]. This leads to a positive correlation between clutch size and development time, as larger clutches intensify larval competition, prolonging maturation[72,73]. Additionally, a trade-off emerges between adult body size and lifespan: while larger body size enhances resource acquisition, it also increases predation risk, resulting in a negative correlation between body size and lifespan[74,75].

Variations in life history trait patterns across animal taxa indicate that the associations among these traits may differ depending on the taxonomic scale[76]. For example, recent studies have also revealed a greater diversity of strategies adopted by different individuals within populations. This study also revealed that individual variation does not always align neatly along a slow–fast continuum[60]. When more individual-level data are available, future studies should investigate the multivariate relationships among traits within species and compare with patterns observed across species to better understand life history evolution across scales.

Conflict of interest declaration. The authors declare no competing interests.

Acknowledgements. We deeply thank Shucun Sun and Xinqiang Xi for their valuable suggestions; Xintong Zhong, Zhimin Qin and Quanquan Jin for the daily help. This research was supported by "Yuxiu Young Scholars Program" of Nanjing University.

References

1. Rochet M, Cornillon P, Sabatier R, Pontier D. 2000 Comparative analysis of phylogenetic and fishing effects in life history patterns of teleost fishes. *Oikos* **91**, 255–270. (doi:10.1034/j.1600-0706.2000.910206.x)

2. Jeschke JM, Kokko H. 2009 The roles of body size and phylogeny in fast and slow life histories. *Evol. Ecol.* **23**, 867–878. (doi:10.1007/s10682-008-9276-y)

3. Myhrvold NP, Baldridge E, Chan B, Sivam D, Freeman DL, Ernest SKM. 2015 An amniote life-history database to perform comparative analyses with birds, mammals, and reptiles: ecological archives E096-269. *Ecology* **96**, 3109–3109. (doi:10.1890/15-0846R.1)

4. Bakewell AT, Davis KE, Freckleton RP, Isaac NJB, Mayhew PJ. 2020 Comparing life histories across taxonomic groups in multiple dimensions: how mammal-like are insects? *Am. Nat.* **195**, 70–81. (doi:10.1086/706195)

5. Stearns SC. 1998 *The evolution of life histories*. Oxford University Press. See https://doi.org/10.1093/oso/9780198577416.001.0001.

6. Reznick D, Bryant MJ, Bashey F. 2002 r - and K -selection revisited: the role of population regulation in life-history evolution. *Ecology* **83**, 1509–1520. (doi:10.1890/0012-9658(2002)083[1509:RAKSRT]2.0.CO;2)

7. Stearns SC. 1989 Trade-offs in life-history evolution. *Funct. Ecol.* **3**, 259–268. (doi:10.2307/2389364)

8. Blackburn TM. 1991 Evidence for a `fast-slow' continuum of life-history traits among parasitoid hymenoptera. *Funct. Ecol.* **5**, 65. (doi:10.2307/2389556)

9. Salguero-Gómez R, Jones OR, Jongejans E, Blomberg SP, Hodgson DJ, Mbeau-Ache C, Zuidema PA, De Kroon H, Buckley YM. 2016 Fast–slow continuum and reproductive strategies structure plant life-history variation worldwide. *Proc. Natl. Acad. Sci.* **113**, 230–235. (doi:10.1073/pnas.1506215112)

10. Allen WL, Street SE, Capellini I. 2017 Fast life history traits promote invasion success in amphibians and reptiles. *Ecol. Lett.* **20**, 222–230. (doi:10.1111/ele.12728)

11. Mayhew PJ. 2016 Comparing parasitoid life histories. *Entomol. Exp. Appl.* **159**, 147–162. (doi:10.1111/eea.12411)

12. Thorson JT *et al.* 2023 Identifying direct and indirect associations among traits by merging phylogenetic comparative methods and structural equation models. *Methods Ecol. Evol.* **14**, 1259–1275. (doi:10.1111/2041-210X.14076)

13. Stott I *et al.* 2024 Life histories are not just fast or slow. *Trends Ecol. Evol.* **39**, 830–

840. (doi:10.1016/j.tree.2024.06.001)

14. Roff DA. 1993 *Evolution Of Life Histories : Theory and Analysis*. 1st edn. New
York,NY: Springer.

15. Flatt T. 2020 Life-history evolution and the genetics of fitness components in
drosophila melanogaster. *Genetics* **214**, 3–48. (doi:10.1534/genetics.119.300160)

16. Pease CM, Bull JJ. 1988 A critique of methods for measuring life history trade-offs.
J. Evol. Biol. **1**, 293–303. (doi:10.1046/j.1420-9101.1988.1040293.x)

17. De Jong G. 1993 Covariances between traits deriving from successive allocations
of a resource. *Funct. Ecol.* **7**, 75. (doi:10.2307/2389869)

18. Cressler CE, Bengtson S, Nelson WA. 2017 Unexpected nongenetic individual
heterogeneity and trait covariance in daphnia and its consequences for ecological
and evolutionary dynamics. *Am. Nat.* **190**, E13–E27. (doi:10.1086/691779)

19. Chang C, Moiron M, Sánchez-Tójar A, Niemelä PT, Laskowski KL. 2024 What is
the meta-analytic evidence for life-history trade-offs at the genetic level? *Ecol. Lett.*
27, e14354. (doi:10.1111/ele.14354)

20. Bakewell AT.. 2020 The diversity, intercorrelation, and macroevolutionary
consequences of insect life histories. PhD thesis, University of York.

21. Capellini I, Baker J, Allen WL, Street SE, Venditti C. 2015 The role of life history
traits in mammalian invasion success. *Ecol. Lett.* **18**, 1099–1107.
(doi:10.1111/ele.12493)

22. Lislevand T, Figuerola J, Székely T. 2007 Avian body sizes in relation to fecundity,
mating system, display behavior, and resource sharing. *Ecology* **88**, 1605–1605.
(doi:10.1890/06-2054)

23. De Magalhães JP, Costa J. 2009 A database of vertebrate longevity records and their
relation to other life-history traits. *J. Evol. Biol.* **22**, 1770–1774.
(doi:10.1111/j.1420-9101.2009.01783.x)

24. Fountain MT, Hopkin SP. 2005 Folsomia candida (collembola): a 'standard' soil
arthropod. *Annu. Rev. Entomol.* **50**, 201–222.
(doi:10.1146/annurev.ento.50.071803.130331)

25. Corbet PS, Suhling F, Soendgerath D. 2006 Voltinism of odonata: a review. *Int. J.*
Odonatol. **9**, 1–44. (doi:10.1080/13887890.2006.9748261)

26. Brooks SJ. 2015 A field guide to the dragonflies and damselflies of britain and
Ireland. In *Descriptive Taxonomy: The Foundation of Biodiversity Research* (eds
MF Watson, CHC Lyal, CA Pendry), pp. 113–126. Cambridge: Cambridge

University Press.

27. Goolsby EW, Bruggeman J, Ané C. 2017 Rphylopars: fast multivariate phylogenetic comparative methods for missing data and within-species variation. *Methods Ecol. Evol.* **8**, 22–27. (doi:10.1111/2041-210X.12612)

28. Fritz SA, Bininda-Emonds ORP, Purvis A. 2009 Geographical variation in predictors of mammalian extinction risk: big is bad, but only in the tropics. *Ecol. Lett.* **12**, 538–549. (doi:10.1111/j.1461-0248.2009.01307.x)

29. Paradis E, Claude J, Strimmer K. 2004 APE: analyses of phylogenetics and evolution in R language. *Bioinformatics* **20**, 289–290. (doi:10.1093/bioinformatics/btg412)

30. Jaffe AL, Slater GJ, Alfaro ME. 2011 The evolution of island gigantism and body size variation in tortoises and turtles. *Biol. Lett.* **7**, 558–561. (doi:10.1098/rsbl.2010.1084)

31. Oaks JR. 2011 A time-calibrated species tree of crocodylia reveals a recent radiation of the true crocodiles: recent radiation of crocodiles. *Evolution* **65**, 3285–3297. (doi:10.1111/j.1558-5646.2011.01373.x)

32. Pyron R, Burbrink FT, Wiens JJ. 2013 A phylogeny and revised classification of squamata, including 4161 species of lizards and snakes. *BMC Evol. Biol.* **13**, 93. (doi:10.1186/1471-2148-13-93)

33. Feng S *et al.* 2020 Dense sampling of bird diversity increases power of comparative genomics. *Nature* **587**, 252–257. (doi:10.1038/s41586-020-2873-9)

34. Chesters D. 2020 The phylogeny of insects in the data-driven era. *Syst. Entomol.* **45**, 540–551. (doi:10.1111/syen.12414)

35. Martins EP, Hansen TF. 1997 Phylogenies and the comparative method: a general approach to incorporating phylogenetic information into the analysis of interspecific data. *Am. Nat.* **149**, 646–667. (doi:10.1086/286013)

36. Freckleton RP, Harvey PH, Pagel M. 2002 Phylogenetic analysis and comparative data: a test and review of evidence. *Am. Nat.* **160**, 712–726. (doi:10.1086/343873)

37. Pagel M. 1999 Inferring the historical patterns of biological evolution. *Nature* **401**, 877–884. (doi:10.1038/44766)

38. Orme CD L *et al*. 2013. caper: comparative analyses of phylogenetics and evolution in R. R package version 1.0.3. https://CRAN.R-project.org/package=pcaper.

39. Hardenberg AV, Gonzalez-Voyer A. 2013 Disentangling evolutionary cause-effect relationships with phylogenetic confirmatory path analysis: phylogenetic path

analysis. *Evolution* **67**, 378–387. (doi:10.1111/j.1558-5646.2012.01790.x)

40. Gonzalez-Voyer A, Von Hardenberg A. 2014 An introduction to phylogenetic path analysis. In *Modern Phylogenetic Comparative Methods and Their Application in Evolutionary Biology* (ed LZ Garamszegi), pp. 201–229. Berlin, Heidelberg: Springer Berlin Heidelberg.

41. Shipley B. 2000 *Cause and correlation in biology: a user's guide to path analysis, structural equations and causal inference*. Cambridge, U.K: Cambridge University Press.

42. Shipley B. 2013 The AIC model selection method applied to path analytic models compared using a d-separation test. *Ecology* **94**, 560–564. (doi:10.1890/12-0976.1)

43. Jiménez-Ortega D, Kolm N, Immler S, Maklakov AA, Gonzalez-Voyer A. 2020 Long life evolves in large-brained bird lineages*. *Evolution* **74**, 2617–2628. (doi:10.1111/evo.14087)

44. Deeming DC, Birchard GF, Crafer R, Eady PE. 2006 Egg mass and incubation period allometry in birds and reptiles: effects of phylogeny. *J. Zool.* **270**, 209–218. (doi:10.1111/j.1469-7998.2006.00131.x)

45. Maino JL, Pirtle EI, Kearney MR. 2017 The effect of egg size on hatch time and metabolic rate: theoretical and empirical insights on developing insect embryos. *Funct. Ecol.* **31**, 227–234. (doi:10.1111/1365-2435.12702)

46. Siikamaki P, Hovi M, Ratti O. 1994 A trade-off between current reproduction and moult in the pied flycatcher- an experiment. *Funct. Ecol.* **8**, 587. (doi:10.2307/2389919)

47. Czesak ME, Fox CW. 2003 Evolutionary ecology of egg size and number in a seed beetle: genetic trade-off differs between environments. *Evolution* **57**, 1121–1132. (doi:10.1111/j.0014-3820.2003.tb00321.x)

48. Charnov EL, Ernest SKM. 2006 The offspring-size/clutch-size trade-off in mammals. *Am. Nat.* **167**, 578–582. (doi:10.1086/501141)

49. Eijs IEM, Van Alphen JJM. 1999 Life history correlations: why are hymenopteran parasitoids an exception? *Ecol. Lett.* **2**, 27–35. (doi:10.1046/j.1461-0248.1999.21045.x)

50. Van Der Bijl W. 2018 phylopath : easy phylogenetic path analysis in R. *PeerJ* **6**, e4718. (doi:10.7717/peerj.4718)

51. Yu G, Smith DK, Zhu H, Guan Y, Lam TT. 2017 ggtree : an r package for visualization and annotation of phylogenetic trees with their covariates and other associated data. *Methods Ecol. Evol.* **8**, 28–36. (doi:10.1111/2041-210X.12628)

52. R Core Team. 2024 R: A language and environment for statistical computing.
https://www.R-project.org/

53. Hadley Wickham. 2016 *ggplot2*. Second Edition. New York, NY: Springer Cham.
See https://ggplot2.tidyverse.org.

54. Wright J, Solbu EB, Engen S. 2020 Contrasting patterns of density-dependent
selection at different life stages can create more than one fast–slow axis of life-
history variation. *Ecol. Evol.* **10**, 3068–3078. (doi:10.1002/ece3.6122)

55. Stearns SC. 1976 Life-history tactics: a review of the ideas. *Q. Rev. Biol.* **51**, 3–47.
(doi:10.1086/409052)

56. Sæther BE. 1987 The influence of body weight on the covariation between
reproductive traits in european birds. *Oikos* **48**, 79. (doi:10.2307/3565691)

57. Healy K, Ezard THG, Jones OR, Salguero-Gómez R, Buckley YM. 2019 Animal
life history is shaped by the pace of life and the distribution of age-specific mortality
and reproduction. *Nat. Ecol. Evol.* **3**, 1217–1224. (doi:10.1038/s41559-019-0938-
7)

58. Oli MK, Dobson FS. 2003 The relative importance of life-history variables to
population growth rate in mammals: cole's prediction revisited. *Am. Nat.* **161**, 422–
440. (doi:10.1086/367591)

59. Salguero-Gómez R *et al.* 2016 COMADRE : a global data base of animal
demography. *J. Anim. Ecol.* **85**, 371–384. (doi:10.1111/1365-2656.12482)

60. Van De Walle J *et al.* 2023 Individual life histories: neither slow nor fast, just
diverse. *Proc. R. Soc. B Biol. Sci.* **290**, 20230511. (doi:10.1098/rspb.2023.0511)

61. John D. Reynolds. 2003 Life histories and extinction risk. In *Macroecology:*
Concepts and Consequences (eds Tim M Blackburn, Kevin J Gaston), pp. 195–217.
Oxford: Wiley-Blackwell.

62. Braendle C, Heyland A, Flatt T. 2011 Integrating mechanistic and evolutionary
analysis of life history variation. In *Mechanisms of Life History Evolution: The*
Genetics and Physiology of Life History Traits and Trade-Offs (eds T Flatt, A
Heyland), pp. 3–10. Oxford University Press.

63. Marden JH, Langford EA, Robertson MA, Fescemyer HW. 2021 Alleles in
metabolic and oxygen-sensing genes are associated with antagonistic pleiotropic
effects on life history traits and population fitness in an ecological model insect*.
Evolution **75**, 116–129. (doi:10.1111/evo.14095)

64. Russell EM, Yom-Tov Y, Geffen E. 2004 Extended parental care and delayed
dispersal: northern, tropical, and southern passerines compared. *Behav. Ecol.* **15**,

647   831–838. (doi:10.1093/beheco/arh088)

65. Stearns SC. 1983 The influence of size and phylogeny on patterns of covariation among life-history traits in the mammals. *Oikos* **41**, 173–187. (doi:10.2307/3544261)

66. Zera AJ, Harshman LG. 2001 The Physiology of Life History Trade-Offs in Animals. *Annu. Rev. Ecol. Syst.* **32**, 95–126. (doi:10.1146/annurev.ecolsys.32.081501.114006)

67. Flatt T, Heyland A, editors. 2011 *Mechanisms of life history evolution: the genetics and physiology of life history traits and trade-offs*. Oxford (G. B.): Oxford university press. See https://doi.org/10.1093/acprof:oso/9780199568765.001.0001.

68. Berger D, Walters R, Gotthard K. 2008 What limits insect fecundity? Body size- and temperature-dependent egg maturation and oviposition in a butterfly. *Funct. Ecol.* **22**, 523–529. (doi:10.1111/j.1365-2435.2008.01392.x)

69. Tammaru T, Haukioja E. 1996 Capital breeders and income breeders among lepidoptera: consequences to population dynamics. *Oikos* **77**, 561–564. (doi:10.2307/3545946)

70. Branson DH. 2008 Influence of individual body size on reproductive traits in melanopline grasshoppers (orthoptera: acrididae). *J. Orthoptera Res.* **17**, 259–263. (doi:10.1665/1082-6467-17.2.259)

71. Boggs CL. 2009 Understanding insect life histories and senescence through a resource allocation lens. *Funct. Ecol.* **23**, 27–37. (doi:10.1111/j.1365-2435.2009.01527.x)

72. Moreau J, Desouhant E, Louâpre P, Goubault M, Rajon E, Jarrige A, Menu F, Thiéry D. 2017 Chapter eight - how host plant and fluctuating environments affect insect reproductive strategies? In *Advances in Botanical Research* (eds N Sauvion, D Thiéry, P-A Calatayud), pp. 259–287. Academic Press.

73. Haave-Audet E, Besson AA, Nakagawa S, Mathot KJ. 2022 Differences in resource acquisition, not allocation, mediate the relationship between behaviour and fitness: a systematic review and meta-analysis. *Biol. Rev.* **97**, 708–731. (doi:10.1111/brv.12819)

74. Peacor SD, Werner EE. 2001 The contribution of trait-mediated indirect effects to the net effects of a predator. *Proc. Natl. Acad. Sci.* **98**, 3904–3908. (doi:10.1073/pnas.071061998)

75. Kuang JJ, Chesson P. 2008 Predation-competition interactions for seasonally recruiting species. *Am. Nat.* **171**, E119–E133. (doi:10.1086/527484)

682   76. Agrawal AA. 2020 A scale-dependent framework for trade-offs, syndromes, and
683   specialization in organismal biology. *Ecology* **101**, e02924. (doi:10.1002/ecy.2924)