Title

Hatching with Numbers: Pre-natal Light Exposure Affects Number Sense and the Mental Number Line in young domestic chicks

Short Title

Prenatal stimulation enhances numerical cognition

Authors

Rosa Rugani[1*], Matteo Macchinizzi[1], Yujia Zhang[2], Lucia Regolin[1]

[1] Department of General Psychology, University of Padua, Padua, Italy.
[2] Department of Psychology, The Ohio State University, Columbus, OH 43210, USA.

*Correspondence concerning this article should be addressed to Rosa Rugani
Email: rosa.rugani@unipd.it

Abstract

Humans order numerosity along a left-to-right Mental Number Line (MNL), traditionally considered culturally rooted. Yet, some species at birth show Spatial-Numerical Associations, SNA, suggesting neural origins. Various accounts link SNA to brain lateralization, but lack evidence.

We investigated brain lateralization effects on numerical spatialization in 100 newborn domestic chicks. In-ovo light exposure yielded strongly-lateralized brains in half the chicks and weakly-lateralized in the other half. Chicks learned to select the 4^{th} item in a sagittal array. At test, the array was rotated 90°, with left and right 4^{th} items correct. Strongly-lateralized chicks outperformed weakly-lateralized ones when ordinal and spatial cues were reliable (experiment 1), but not with unreliable spatial cues (experiment 2). Moreover, only strongly-lateralized chicks showed left-to-right directionality, suggesting the right hemisphere's key role in integrating spatial and numerical cues. We demonstrate that brain lateralization is fundamental for developing a left-to-right oriented SNA.

Keywords

Number Sense, Brain lateralization, Prenatal stimulation, Mental Number Line, Spatial-numerical association, Comparative cognition

Introduction

Humans share with other animals basic numerical capacities (Cantlon and Brannon, 2007; Cordes et al., 2001; Cordes and Brannon, 2009; Rugani, 2018; Vallortigara, 2018); this evolutionarily ancient number sense serves as a building block for our unique mathematical abilities (Dehaene, 2011). A distinctive feature of the human numerical system is the Mental Number Line, MNL: an association of small numbers with the left and large ones with the right. Numeral visualization in various spatial configurations (right-to-left, circular, and vertical) was first reported by Galton (1880) (see also Bertillon, 1880; Seron et al., 1992; Tang et al., 2008). The first experimental demonstration of left-to-right numerical spatialization was observed in a parity judgment task, where participants pressed a left or right key for odd or even numerals. Faster responses to small numbers on the left and to large numbers on the right have been described as the SNARC effect (Spatial-Numerical Association of Response Codes): a spatial congruency between the response side (left or right) and the relative position of the represented numerical magnitude on an oriented MNL (left-space/small numbers and right-space/large numbers) (Dehaene et al., 1993). Left-to-right numerical spatialization has been replicated across various experiments (e.g. Bächtold et al., 1998; Dehaene, 2011); however, evidence remains inconsistent across paradigms (Karolis et al., 2011) and reading direction (Shaki et al., 2009; Zebian, 2005), emphasizing the role of task-dependent factors and culture in shaping numerical spatialization. Moreover, spatial biases in numerical processing that demonstrated Spatial Numerical Association, SNA, have been reported in young infants (Bulf et al., 2016; de Hevia, 2021; de Hevia and Spelke, 2010; Rugani et al., 2022; West and McCrink, 2021), newborns (de Hevia et al., 2017, 2014; Di Giorgio et al., 2019; McCrink et al., 2020) and non-humans (Drucker and Brannon, 2014; Giurfa et al., 2022; Rugani et al., 2024, 2020b, 2015a, 2010, 2007), challenging the predominant role of culture in determining the left-to-right orientation. Evidence in non-human animals indicates that nervous systems across various species, despite their different levels of complexity, are prewired in how they relate numbers to space. Together, conflicting evidence of an oriented representation of numerical magnitude, as described in the MNL in humans, and broader spatial biases in numerical processing, as reported in the SNA in non-verbal or preverbal subjects, suggests that numerical spatialization likely originates from neural bases, yet it remains flexible and subject to modulation by experience and contextual factors (Rugani and de Hevia, 2017).

The first evidence of numerical spatialization in non-human subjects was observed in day-old domestic chicks (Rugani et al., 2007, 2010) and adult Clark's nutcrackers (Rugani et al., 2010). Birds learned to identify a target item, e.g., the 4[th], in a sagittally oriented array of identical items. They were then tested with the array rotated by 90°, thus laying on a fronto-parallel plane with the items oriented left-to-right. Animals selected the 4[th] left item more often than the 4[th] right one, suggesting that number is intrinsically represented from left to right (Rugani et al., 2007). Using this paradigm, the same left-to-right proto-counting tendency has been observed also in adult rhesus monkeys (Drucker and Brannon, 2014), and preschoolers (West and McCrink, 2021).

Behavioral observations supporting the SNA have increased (Adachi, 2014; Bulf et al., 2016; de Hevia et al., 2017; de Hevia and Spelke, 2010; Di Giorgio et al., 2019; Drucker and Brannon, 2014; Giurfa et al., 2022; McCrink et al., 2020; Rugani et al., 2024, 2015a, 2010; West and McCrink, 2021); (but see for null results Beran et al., 2019; Triki and Bshary, 2018). Concurrently, specific neurons responding to numerosities have been identified in human (Pearson et al., 2009; Piazza et al., 2004; Piazza and Eger, 2016), monkey (Nieder, 2002), crow (Ditz and Nieder, 2015), and chick (Kobylkov et al., 2022) brains. Nevertheless, the biological mechanisms and neural underpinnings of SNA are still largely unknown.

The right hemisphere is specialized for analyzing spatial information (Rashid and Andrew, 1989; Regolin et al., 2005) and for processing numerical information (Piazza and Eger, 2016; Rugani and Regolin, 2020). This functional overlap (Harvey et al., 2013; Zorzi et al., 2002) has been proposed as the basis for the SNA that prompts animals to start counting from left to right.

This will be referred to as the right-hemisphere dominance model (Rugani et al., 2016, 2015a). Two other explicatory models have been put forward: the Emotional-Valence (Vallortigara, 2018) and the Brain Asymmetric Frequency Tuning, BAFT (Felisatti et al., 2020).

The Emotional-Valence model is grounded in evidence demonstrating that the right hemisphere processes negative emotions and the left hemisphere handles positive ones (Davidson, 2004; Vallortigara, 2018). This model posits that small numerosities are associated with a negative valence, and activate the right hemisphere that directs movement toward the left; large numerosities would be associated with a positive valence, activating the left hemisphere that leads movement toward the right (Vallortigara, 2018).

BAFT links spatial frequency processing to higher cognitive functions. The right hemisphere specializes in processing low spatial frequencies, while the left hemisphere high spatial frequencies (Gazzaniga, 2000). Because smaller numerosities correspond to low frequencies, and large numerosities to high frequencies, such hemispheric specialization may explain SNA (Felisatti et al., 2020).

Although all models, whether mutually exclusive or not, acknowledge the role of hemispheric specialization in SNA, no research has directly investigated its effect on number spatialization. Direct manipulation of brain lateralization is essential for determining its impact on SNA. While a stronger bias is expected in highly lateralized chicks, the right-hemisphere model predicts a left bias, the Emotional Valence model predicts a right bias, and BAFT predicts no bias.

The aim of the present study is to assess whether prenatal light stimulation influences SNA. Specifically, the study has two main objectives: (i) to assess the effect of hemispheric lateralization on number spatialization by varying its degree; (ii) to disentangle the role of the two cerebral hemispheres using monocular occlusion.

As for the first aim, lateralization levels in domestic chicks can be easily manipulated by controlling exposure to light in the final period of incubation, between embryonic days 18 and 21 (Rogers and Bolden, 1991; Rogers and Sink, 1988). During development, the embryo rotates such that the right eye faces outward toward the translucent eggshell and any available environmental light. In contrast, the left eye is oriented toward the body, receiving little to no light. The chick embryo's visual system, specifically the thalamofugal pathway, undergoes differentiation in-ovo. Egg exposure to light leads to an asymmetrical stimulation of the two eyes such that there is an increase in forebrain projections from the left side of the thalamus (fed by the light-stimulated right eye) compared with the right side (Deng and Rogers, 1998). Such asymmetries extend to the strength of visual projections from the thalamus to the visual Wulst (a laminated bulge in the dorsal telencephalon, functionally analogous to mammalian visual cortex (Clark and Colombo, 2020; Rogers and Sink, 1988)). Specifically, the right visual Wulst receives more bilateral information from the two eyes than the left one (Costalunga et al., 2022). Evidence shows that as little as two-hour light exposure prior to hatching is sufficient to induce these brain asymmetries (Deng and Rogers, 1997; Rogers, 1982). As a consequence of light-enhanced hemispheric differentiation, light-incubated chicks, Li-chicks, are strongly lateralized, as demonstrated by cognitive and behavioral biases (Daisley et al., 2009). For the present study, we expect that Li-chicks will exhibit a clearer left-to-right oriented numerical spatialization compared to dark-incubated chicks, Di-chicks, whose brain lateralization is mainly prevented (Rogers and Bolden, 1991).

Additionally, higher levels of lateralization enhance cognitive performance (Daisley et al., 2009). Strongly lateralized chicks perform better in dual tasks requiring simultaneous foraging and predator monitoring (Dharmaretnam and Rogers, 2005), in discrimination and categorization tasks such as distinguishing pebbles from food grains (Deng and Rogers, 1997), and in transitive inference tasks where they learn hierarchical stimulus relationships (A > B > C > D > E) that then apply to novel pairings (i.e., AE and BD) (Daisley et al., 2010). Brain lateralization seems to improve also numerical ability: strongly lateralized fish outperform non-lateralized individuals in quantity discrimination tasks. In a shoal-choice task, non-lateralized fish failed to distinguish

between 4 vs. 6 conspecifics, while lateralized fish reliably selected the larger group. Similarly, in an abstract dot discrimination task, lateralized fish successfully distinguished between 3 vs. 4 dots, whereas non-lateralized fish did not (Dadda et al., 2015). Although lateralization was inferred indirectly through the mirror test, similar results in shoal-choice tasks emerged using a detour test (Gatto et al., 2019), supporting the hypothesis that brain lateralization enhances cognitive efficiency and was a key selective force in evolution (Dadda et al., 2015). In this study, rather than inferring brain lateralization from behavioral tests, we directly examined the impact of prenatal light stimulation, which is known to influence lateralization in domestic chickens (Costalunga et al., 2022; Rogers, 1982; Rogers and Bolden, 1991). This approach provides a more structural understanding of its potential effects on numerical abilities and the relationship between environmental factors and number sense. Based on previous findings, we hypothesized that numerical performance would be enhanced in Li-chicks compared to Di-chicks.

As for the second aim, to investigate hemispheric dominance, we used temporary monocular occlusion by applying a removable eye patch to cover one eye of the chicks. Due to the complete decussation of fibers at the optic chiasm (Weidner et al., 1985) and lack of a structure homologous to the corpus callosum (even though other smaller tracts allow interhemispheric communication (Robert and Cuénod, 1969; Theiss et al., 2003)), information received by each eye is mainly elaborated by the contralateral hemisphere (Deng and Rogers, 1998; Rogers and Vallortigara, 2021). Restriction of the visual input by monocular occlusion will allow us to disentangle how the two hemispheres elaborate ordinal information, and to test whether one hemisphere is dominant.

Results

Here, we tested 100 male domestic chicks (*Gallus gallus*) of the Aviagen ROSS 308 line (experiment 1, $n = 48$; experiment 2, $n = 52$). Dark-incubated chicks (Di-chicks; $n=24$ in experiment 1, $n =26$ in experiment 2) were obtained from eggs incubated in darkness, while Light-incubated chicks (Li-chicks; $n =24$ in experiment 1, $n =26$ in experiment 2) were obtained from eggs exposed to light (LED 4.8W lightbulb) from day 18 to 21 of incubation.

All chicks were trained to peck for food reward at the 4^{th} item in an array of 10 identical, equally-spaced, and sagittally aligned items (plastic caps); Fig. 1A. Each chick underwent four tests. In the first test, the array was oriented as during training, i.e., sagittally with respect to the chick's starting point; Sagittal test, Fig. 1A. This test was run in Binocular condition of vision.

Subsequently, each chick underwent tests in which the array was rotated by 90° so as to lay fronto-parallel with respect to the chick's starting position; Fronto-Parallel tests; Fig. 1B. Fronto-Parallel tests were administered to the chick in three conditions of vision: Binocular, Fig. 1C, Monocular-Right (left eye patched and right eye in use; Fig. 1D), and Monocular-Left (right eye patched and left eye in use; Fig. 1E).

The Fronto-Parallel test presented the array as oriented from left to right. This orientation produces two possible correct options, both equidistant from the subject: the 4^{th} left and the 4^{th} right item; Fig. 1B. The critical difference between experiment 1 and 2 was that, during testing, spatial information was available in experiment 1, but was unreliable in experiment 2. Specifically, in experiment 1, the inter-item distance and the total length of the array was constant and identical to the training array, thus chicks could identify the items on the basis of either ordinal or spatial cues. In experiment 2, the inter-item distance was uniform within each trial, but varied systematically between trials (1.44 cm, 2.55 cm, 3.11 cm, and 3.66 cm resulting in total array lengths of respectively 43.0, 53.0, 58.0, and 63.0 cm) thus the spatial information was unreliable.

These experimental manipulations allow us to test and compare the three models proposed to explain the origin of the Mental Number Line (Table 1).

193   i)   According to the right-hemisphere dominance model (Rugani et al., 2016, 2015a),
194 which highlights the role of the right hemisphere in processing spatial and numerical cues, a left

Fig. 1. Illustration of the experimental apparatus and of the three conditions of vision. (**A**) Array configuration used at training and Sagittal test. (**B**) Array configuration used for the Fronto-Parallel tests. S.P. indicates the chick's starting position. (**C**) Binocular condition of vision: both eyes, thus both hemispheres (left in red and right in blue), in use. (**D**) Right monocular condition with the right eye and the left hemisphere in use, in red. (**E**) Left monocular condition with the left eye and the right hemisphere in use, in blue. In monocular conditions, the eye not receiving light and its contralateral hemisphere are depicted in gray.

bias is expected whenever both eyes and hemispheres are in use, particularly in individuals with
greater interhemispheric differences, i.e., Li-chicks. Moreover, the left bias should be more
pronounced when only the right hemisphere processes information, as in the monocular-left eye
condition.
ii)   Following the emotional-valence model (Vallortigara, 2018), which suggests that
the left hemisphere processes positive emotions, considering that food itself is associated with
positive emotions, a right bias might be expected in bilateral processing during food search. This
bias should be more pronounced in Li-chicks due to their enhanced hemispheric differentiation.

iii)   As for the BAFT model (Felisatti et al., 2020), the left/right symmetry in item
disposition within the array would lead to an absence of bias with no difference between strong and
weak lateralized animals.
In each trial, chicks were allowed a single peck. We recorded the selected item to calculate
the percentage of responses at each position and averaged them separately for each group and test.
We analyzed the group percentage for choosing the 4th item above chance (10%), using Bonferroni
correction for multiple comparisons (data and significant results are reported in Table 2; Additional
analyses on the selection of each item are reported in Table S1 and Table S2). To assess side bias
in the Fronto-Parallel tests, we compared correct choices on the left (4L) *versus* the right (4R) using
a paired *t* test, with Cohen's *d* as the effect size, and Bonferroni correction, see Table 2. Moreover,
we tested whether brain lateralization influenced accuracy by comparing the percentage of correct
choices (i.e., the selection of the 4th item in the Sagittal test; the 4L or 4R items in the Fronto-
Parallel tests) between Li-chicks and Di-chicks using a two-sample *t* test, with Cohen's *d* as the
effect size, and Bonferroni correction. We conducted both frequentist and Bayesian statistics to
ensure reliable interpretations of our results.
218

Table 1. Models and hypotheses explaining SNA. For each of the three main models, is
reported a brief explanation of the hypotheses regarding SNA mechanisms, and the predicted
outcomes for experiment 1 and 2.

Model	Hypothesis	Predictions for experiment 1	Predictions for experiment 2
Right-Hemisphere Dominance	Right hemisphere dominance in visuospatial attention.	Right-hemisphere dominance leads to left-to-right scanning. The left bias should be more pronounced in Light-incubated chicks.	Absence of spatial cues results in diminished involvement of the right hemisphere, and the left bias is predicted to be less pronounced.
Emotional-Valence	Left hemisphere dominance in processing positive emotions; right hemisphere dominance in processing negative emotions.	Left hemisphere processing of positive rewards (e.g., food) leads to a rightward bias. This should be more pronounced in Light-incubated chicks.	Lack of spatial cues is not expected to cause differences compared to the predictions of Experiment 1.
Brain Asymmetric Frequency Tuning (BAFT)	Left hemisphere dominance in high-frequency processing; right hemisphere dominance in low-frequency processing.	No bias expected due to the symmetrical array configuration. No difference is expected between Light- and Dark-incubated chicks.	Lack of spatial cues is expected to have no significant impact on the predicted outcomes of Experiment 1.

219
220
221   **Results in the tests allowing utilization of reliable ordinal and spatial cues (experiment 1)**
222
223   **Sagittal test conducted under binocular vision condition.**

In the Sagittal test (Fig. 2A), Li-chicks selected the 2^{nd} and 4^{th} items above chance; Di-chicks selected the 1^{st}, 2^{nd} and 4^{th} items above chance (Table 2 and Table S1). Moreover, Li-chicks showed higher accuracy in selecting the 4^{th} item than Di-chicks (Li-chicks: $n = 24$, mean = 36.38, SE = 1.666; Di-chicks: $n = 24$, mean = 28.44, SE = 1.913; $t(45.1) = 3.132$, $P = 0.003$, $d = 0.904$; BF = 12.588); Fig. 2A.

Fronto-Parallel transfer test conducted under binocular vision condition.
In the Fronto-Parallel Binocular test Fig. 2B), Li-chicks successfully transferred what was learned to the rotated series: they selected only the 4^{th} left item above chance (Table 2). Li-chicks showed a left bias, pecking the 4^{th} left item more than the 4^{th} right one ($n = 24$, $t(23) = 4.791$, $P < 0.001$, $d = 1.635$; BF = 337.124). Di-chicks pecked both 4^{th} left and 4^{th} right items above chance (Table 2) without a difference ($n = 24$, $t(23) = 0.218$, $P = 1.000$ $d = 0.076$; BF = 0.219). Remarkably, only Li-chicks showed a left bias, indicating a tendency to proto-count from left to right.
Comparing accuracy between the two groups, Li-chicks selected the 4^{th} left item more than Di-chicks did (Li-chicks: $n = 24$, mean = 28.85, SE = 2.572; Di-chicks: $n = 24$, mean = 18.96, SE = 2.149 ; $t(44.6) = 2.952$, $P = 0.010$, $d = 0.852$; BF = 8.456); while Li-chicks choose the 4^{th} right item less than Di-chicks (Li-chicks: $n = 24$, mean = 11.89, SE = 1.532; Di-chicks: $n = 24$, mean = 19.79, SE = 2.321; $t(39.8) = -2.840$, $P = 0.014$, $d = 0.820$; BF = 6.657).

Fronto-Parallel transfer test conducted under left-monocular vision condition.
In the Fronto-Parallel Monocular-Left test (Fig. 2C), Li-chicks were able to transfer learning to a differently oriented series, correctly selecting the 4^{th} left item above chance, even if they also pecked at the 1^{st} left item (Table 2; Table S1). Moreover, they selected the 4^{th} left more than the 4^{th} right item ($n = 24$, $t(23) = 6.056$, $P < 0.001$, $d = 2.088$; BF = 5598.452,). Di-chicks failed: they only selected the 1^{st} and 2^{nd} left items above chance (Table S1).
As for the difference between the two groups, Li-chicks selected the 4^{th} left item more than Di-chicks (Li-chicks, $n = 24$, mean = 22.59, SE = 1.783; Di-chicks, $n = 24$, mean = 12.78, SE = 1.696; $t(45.9) = 3.988$, $P < 0.001$, $d = 1.151$; BF = 104.830).
These results suggest that whenever the right hemisphere is processing the information, light exposure affects the left bias and numerical performance. This evidence, on one side, supports the relevance of the right hemisphere in directing SNA directionality (Rugani et al., 2015a; Rugani and de Hevia, 2017). On the other side, it shows how significant experiences that stimulate brain development, although limited to a few hours of exposure to moderate ambient light, can boost cognitive performance.

Fronto-Parallel transfer test conducted under right-monocular vision condition.
In the Fronto-parallel Monocular-Right test (Fig. 2D), Li-chicks succeeded correctly in selecting the 4^{th} right item, which was pecked more than the 4^{th} left item ($n = 24$, $t(23) = -6.151$, $P < 0.001$, $d = 1.844$; BF = 6887.511), even if they also selected the 1^{st} left item above chance (Table 2 and Table S1). Di-chicks failed: they did not select the 4^{th} left nor right item above chance; the only item selected above chance was the 1^{st} right one (Table S1).
As for the accuracy differences between the two groups, Li-chicks selected the 4^{th} right item more than Di-chicks (Li-chicks: $n = 24$, mean = 21.48, SE = 2.191; Di-chicks: $n = 24$, mean = 13.42, SE = 1.792; $t(44.3) = 2.850$, $P = 0.013$, $d = 0.823$; BF = 6.799).
Again, only Li-chicks succeeded while Di-chicks failed, corroborating evidence on the importance of light stimulation in favoring the development of both hemispheres (Costalunga et al., 2022) and boosting cognitive performance.
Overall, these data showed that prenatal light experience can stimulate brain development and hemispheric specialization, which emphasizes the Spatial Numerical Association and enhances

performance in a spatial/numerical task. This provides novel evidence of the role of brain lateralization in determining SNA and in boosting proto-numerical counting.

Results in the tests allowing utilization of reliable ordinal and unreliable spatial cues (experiment 2)

Sagittal test conducted under binocular vision condition.
In the Sagittal test (Fig. 3A), both groups succeeded: Li-chicks exclusively selected the correct 4^{th} item above chance (Table 2), while Di-chicks selected the 4^{th} item, but also mistakenly pecked the 2^{nd} item (Table 2 and Table S2). Yet, accuracy in pecking the 4^{th} item did not differ in the two groups (Li-chicks: $n = 26$, mean $= 32.27$, SE $= 2.203$; Di-chicks: $n = 26$, mean $= 31.49$, SE $= 3.008$; $t(45.8) = 0.211$, $P = 1.000$, $d = 0.059$; BF $= 0.283$).

Fronto-Parallel transfer test conducted under binocular vision condition.
In the Fronto-Parallel Binocular test (Fig. 3B), Li-chicks and Di-chicks selected both 4^{th} left and right items above chance (Table 2). For each group, no differences emerged in selecting the 4^{th} left and 4^{th} right items, indicating a lack of side bias in both groups (Li-chicks: $n = 26$, $t(25) = 1.910$, $P = 0.135$, $d = 0.624$; BF $= 0.997$; Di-chicks: $n = 26$, $t(25) = -0.218$, $P = 1.000$, $d = 0.098$; BF $= 0.219$). Li-chicks did not differ from Di-chicks in responses on the 4^{th} left (Li-chicks: $n = 26$, mean $= 21.77$, SE $= 1.939$; Di-chicks: $n = 26$, mean $= 18.73$, SE $= 1.813$; $t(49.8) = 1.147$, $P = 0.513$, $d = 0.318$; BF $= 0.478$) or 4^{th} right item (Li-chicks: $n = 26$, mean $= 15.80$, SE $= 1.812$; Di-chicks: $n = 26$, mean $= 19.67$, SE $= 1.966$; $t(49.7) = -1.446$, $P = 0.309$, $d = 0.401$; BF $= 0.655$). This confirms that whenever the spatial information is unavailable at the test, the left bias disappears (Rugani et al., 2011), highlighting the role of the right hemisphere in processing spatial information and determining the left-to-right orientation of the Spatial-Numerical Association.

Fronto-Parallel transfer test conducted under left-monocular vision condition.
In the Fronto-Parallel Left-Monocular test (Fig. 3C), Li-and Di-chicks failed: Li-chicks selected the 1^{st} left item (Table S2); Di-chicks selected the 1^{st} and the 2^{nd} left item above chance (Table S2). As for the accuracy, the two groups equally selected the 4^{th} left (Li-chicks: $n = 26$, mean $= 15.01$, SE $= 1.832$; Di-chicks: $n = 26$, mean $= 14.84$, SE $= 1.937$; $t(49.8) = 0.066$, $P = 1.000$, $d = 0.018$; BF $= 0.279$) and the 4^{th} right items (Li-chicks: $n = 26$, mean $= 7.94$, SE $= 1.193$; Di-chicks: $n = 26$, mean $= 8.33$, SE $= 1.356$; $t(49.2) = -0.218$ $P = 1.000$, $d = 0.060$; BF $= 0.284$).
Whenever spatial information is unavailable, the right hemisphere fails to transfer tasks. This indicates that independent of the hemisphere's development, unilateral right hemispheric processing is insufficient in dealing with an ordinal task; thus, ordinality does not appear to be lateralized to the right hemisphere.

Fronto-Parallel transfer test conducted under right-monocular vision condition.
In the Fronto-Parallel Monocular-Right test (Fig. 3D), only Li-chicks succeeded and selected the 4^{th} right item; even if they also pecked the 1^{st} and the 2^{nd} right items (Table 2 and Table S2), moreover they selected the 4^{th} right more than the 4^{th} left item ($n = 26$, $t(25) = -4.946$, $P < 0.001$, $d = -1.447$; BF $= 573.520$). Di-chicks failed to select the 4^{th} left or right item; instead they selected only the 1^{st} right item above chance (Table S2). As for the difference in accuracy between the two groups, Li-chicks did not select the 4^{th} right item more than Di-chicks (Li-chicks: $n = 26$, mean $= 17.98$, SE $= 1.889$; Di-chicks: $n = 26$, mean $= 12.46$, SE $= 1.815$; $t(49.9) = 2.110$, $P = 0.080$, $d = 0.585$; BF $= 1.678$). This suggests that lateralization influences numerical cognition even in the absence of spatial information, and that the left hemisphere plays a significant role in processing ordinal information.

Fig. 2. Results of experiment 1. The average percentage of chicks' choices (y-axis) as a function of item positions (x-axis), light exposure modulating brain lateralization, and tests. Error bars indicate standard errors. The gray dashed line indicates chance level (10%). (**A**) Results of the Sagittal test. (**B**) Results of the Fronto-Parallel Binocular test. (**C**) Results of the Fronto-Parallel Monocular Left test. (**D**) Results of Fronto-Parallel Monocular Right test. ***$P < 0.001$, **$P < 0.01$, and *$P < 0.05$.

Fig. 3. Results of experiment 2. The average percentage of chicks' choices (y-axis) as a function of item positions (x-axis), light exposure modulating brain lateralization, and tests. Error bars indicate standard errors. The gray dashed line indicates chance level (10%). (**A**) Results of the Sagittal test. (**B**) Results of the Fronto-Parallel Binocular test. (**C**) Results of the Fronto-Parallel Monocular Left test. (**D**) Results of Fronto-Parallel Monocular Right test. ***$P < 0.001$, **$P < 0.01$, and *$P < 0.05$

Discussion
General
Our main findings are that prenatal exposure that can modulate brain lateralization in domestic
chicks impacts the left-to-right oriented numerical spatialization and numerical performance.
In experiment 1, which allowed chicks to reliably use both ordinal and spatial cues in
identifying the 4th item, chicks exhibited different behaviors despite identical learning experiences
and tasks, either showing or not showing a left bias, depending on their prenatal light exposure. In
the Fronto-Parallel Binocular test, when both eyes and hemispheres processed the information,
chicks hatched from light-incubated eggs, Li-chicks, selected only the 4th left item; while chicks
hatched from dark-incubated eggs, Di-chicks, equally selected the 4th left and the 4th right item.
Only Li-chicks (i.e., more strongly lateralized) demonstrated left-to-right proto-counting, indicating
that brain lateralization influences Mental Number Line directionality. When the two hemispheres
engaged in differential processing, as observed in Li-chicks, a unidirectional left-to-right oriented
numerical spatialization emerged. Conversely, when prenatal stimulation did not enhance
hemispheric specialization, resulting in more homogeneous hemispheric processing, animals (Di-
chicks) showed no directional bias. These pioneering findings corroborate all models positing
hemispheric specialization as the neural basis for Spatial Numerical Association (Felisatti et al.,
2020; Rugani et al., 2016, 2015a; Vallortigara, 2018), while establishing lateralization essential

prerequisite for numerical spatialization. However, pre-hatching light stimulation did not affect chicks' performance when spatial information was unavailable (experiment 2). This finding substantiates the relevance of spatial information and highlights that its integration with numerical processing within shared cortical regions is fundamental to the neurobiological underpinning of number spatialization. This integration is coherent with the fact that in the chick's brain the right hemisphere is dominant in processing spatial information (Rashid and Andrew, 1989; Regolin et al., 2005), but can also process numerical information (Rugani et al., 2011; Rugani and Regolin, 2020) (consistently with primates and human literature (Piazza and Eger, 2016)). The present data align with previous research that used monocular occlusion to disentangle the engagement of the two hemispheres with spatial or ordinal cues (Rugani and Regolin, 2020). In prior research, day-old chicks learned to select the 4th item in an array of 10 identical sagittal-aligned items maintained in fixed positions, so that both spatial and ordinal cues were available during learning. At test, chicks faced a left-to-right-oriented series where the inter-item distance was manipulated so that the 3rd item was at the same distance from the beginning of the series as the 4th item had been at training. This forced chicks to choose either spatial or ordinal cues. Chicks tested binocularly selected both the 4th left and right items above chance expectation, confirming that a coherent use of numerical and spatial information is essential in limiting birds' responses toward the left (Rugani et al., 2011). Chicks tested monocularly chose the 3rd and 4th items on the seeing side, suggesting that birds relied on spatial or ordinal cues to a similar extent in different trials and that each hemisphere can process both cues (Rugani and Regolin, 2020).

Here, in monocular conditions, Li-chicks succeeded in Fronto-Parallel tests. Even if they directed their pecks to the visible side (Rugani et al., 2016; Rugani and Regolin, 2021), they selected the 4th item above chance expectation. This was the case in experiment 1 when ordinal and spatial cues were available at test. Nevertheless, in experiment 2, when spatial information was available at training but unreliable at test, chicks succeeded in the right-eye/left-hemisphere, but not in the left-eye/right-hemisphere condition. This corroborates the hypothesis of the right hemisphere specificity in the analysis of spatial cues and suggests that the left hemisphere is more specialized in processing ordinal information. The left-to-right spatialization of numerosity appears to be based on preferential processing by the right hemisphere when spatial information is available and hemispheric specialization is favored by environmental stimulation.

Ecological Implications and Adaptive Variability

Experimental contexts allow for selective manipulation of the environment, enabling changes that are highly improbable in nature. In experiment 2, the item arrangement, that had been experienced as a stable context that provided coherent and reliable numerical and spatial information during training, was manipulated to eliminate spatial information at test. This manipulation resulted in the disappearance of left-to-right oriented directionality in the binocular condition and in a failure in the left-eye/right-hemisphere condition. We can speculate that in some naturalistic contexts, establishing an anchor point (potentially based on reliable landmarks or beacons) and predetermining a privileged starting position might be advantageous in facilitating processing and reducing conflicts between incompatible responses. Throughout evolution, this may have maximized right hemisphere engagement, specialized for spatial information processing, and triggered an imbalance favoring left space. This left-biased space would serve as an anchor point from which to initiate environmental scanning, avoiding the delay presumably implied if there was not a hemisphere taking control of processing and guiding behavior (Rogers et al., 2013). Such an intrinsic, left-oriented bias might be advantageous for other ecological situations requiring number processing, such as quantifying conspecifics or food items. For example, when foraging, a consistent left-to-right scanning pattern could help animals to efficiently locate and quantify food sources without overlooking areas. The tendency to scan items from the preferred left side (Diekamp et al., 2005), might be evolved as an adaptive behavior to maximize fitness. This

tendency could then have been assimilated by other cognitive processes that share neural substrates, including numerical cognition. Numerosities relevant for animals (such as the number of conspecifics, food items, or predators) are inherently distributed in space. The right hemisphere specialized in spatial processing incorporated some rudimentary forms of enumeration. This resulted in a right hemispheric dominance for both space and number. This might explain the observed left-to-right bias in numerical cognition tasks. The absence of this bias in conditions where spatial cues are artificially eliminated, as in experiment 2, underscores the spatial nature of numerical processing of objects in the environment. Remarkably, the left-to-right directionality is not reported in weakly lateralized, Di-chicks, in both experiments, irrespective of spatial cues availability. It should be noted that in some situations like predator detection, even if a systematic approach to surveying the surroundings could be beneficial for prompt threats detection, it could also lead to more predictability. This could favor predators with a complementary approach directionality. From an evolutionary perspective, lateralization variability within a species can be viewed as an adaptive strategy. This variability may represent a form of evolutionary bet-hedging (Simons, 2011), where different degrees of lateralization confer different adaptive advantages in fluctuating ecological contexts. Bet-hedging strategies maintain population fitness by promoting phenotypic diversity, optimizing adaptation in unpredictable environments (Philippi and Seger, 1989). Lateralization variability might also contribute to the species' behavioral unpredictability, offering an advantage in predator-prey dynamics (Güntürkün et al., 2020). The persistence of both lateralized and non-lateralized individuals within a population may be an evolutionarily stable strategy (Ghirlanda and Vallortigara, 2004), conferring differential advantages to different individuals, overall maintaining high fitness and making the overall population less predictable (Rogers, 2021).

Monocular test outcomes support that light-induced lateralization enhances spatial-numerical performance

Our investigation produced a second major result and showed that prenatal exposure significantly affected performance. In the Sagittal test, Li-chicks outperformed Di-chicks, demonstrating that a higher degree of lateralization led to greater accuracy when both spatial and numerical cues were available (experiment 1).

The results of the monocular Fronto-Parallel tests further support the effect of embryonic stimulation in enhancing performance, as only Li-chicks succeeded in both the left and right monocular conditions. Remarkably, in the Fronto-Parallel test of experiment 1 allowing a coherent use of spatial and numerical information, Li-chicks tested in left-monocular condition of vision showed a bias alike to Li-chicks tested in binocular condition of vision. When a hemisphere is dominant for a function, behavior under its sole control often matches that observed in normal conditions of vision when both hemispheres are active (Rogers et al., 2013). In the present scenario, similarities between chicks tested in Binocular and Left Monocular conditions of vision suggest that when both hemispheres are processing the information, the availability of spatial cues triggers an overactivation of the right hemisphere resulting in a leftward bias (Rugani et al., 2016).

Whenever processing was confined to a single hemisphere, either one, only strongly lateralized Li-chicks succeeded while weakly lateralized Di-chicks failed, corroborating evidence on the importance of light stimulation in favoring the development and specialization of both hemispheres and in boosting cognitive performance. This finding aligns with previous anatomical studies that demonstrated the presence of light-dependent lateralization in bilaterally responsive units of the right visual Wulst (Costalunga et al., 2022; Rogers and Deng, 1999). Nevertheless, in the specific case of Li-chicks tested with the left-eye/right-hemisphere in use, subjects failed the Fronto-Parallel test when the use of spatial cues was prevented (experiment 2). This highlights the reliance of the right hemisphere on spatial information.

Interpreting results through proposed models for the origin of the Mental Number Line

The present results reveal that brain lateralization influences performance in ordinal tasks involving both spatial and numerical cues, suggesting a joint contribution of hemispheric specialization and environmental stimulation to the spatial organization of numbers. These results allow us to reconsider the models proposed to explain the SNA.

1. The BAFT model would not predict any asymmetry as it refers number spatialization to differences in spatial frequencies (Felisatti et al., 2020), but in the present study the spatial distribution of the stimuli is symmetrical. The predictions based on this model fit with the results of the experiment 2, where chicks did not show any spatial bias in the Fronto-Parallel Binocular test. Yet the model fails in predicting the left bias found in the Fronto-Parallel Binocular and Left Monocular tests of experiment 1. While this model can still be valuable in explaining spatial-numerical association in other contexts, it fails to account for the left-to-right proto-counting observed in this experimental setting (Drucker and Brannon, 2014; Rugani et al., 2016, 2011, 2010, 2007; West and McCrink, 2021).

2. According to the emotional valence model chicks should exhibit a rightward bias in food-seeking behavior, since food rewards trigger positive emotions, particularly when lateralization is pronounced. The assumption that a larger numerosity can be associated with more positive emotion and consequent preferential processing by the left hemisphere would explain the right bias (Vallortigara, 2018). Conversely, the assumption that a smaller numerosity is associated with a negative emotion would activate the right hemisphere, driving animals toward the left (Vallortigara, 2018). Nevertheless, in the present study, chicks showed a left bias when they could rely on consistent spatial and numerical information (experiment 1) and no bias when spatial information was unreliable (experiment 2). Thus, the results obtained in the present study do not fit the emotional model.

3. The left bias meets the predictions based on the right-hemisphere dominance model, which links the origin of the Mental Number Line to the right hemisphere's specialization in visual/numerical processing —resulting in a predisposed left-to-right scanning tendency (Rugani et al., 2015a). Remarkably, this model has been put forward to explain outcomes in both the previously-mentioned paradigms (Giurfa et al., 2022; Rugani et al., 2016, 2015a; Rugani and de Hevia, 2017), thus explaining both the left-to-right oriented searching found in ordinal tasks, as well as the association of smaller numerosity with the left and larger numerosity with the right. Such association likely results from brain asymmetry driven by right-hemisphere dominance in visuospatial attention. Although in principle arbitrary, the left-to-right mapping direction during evolution may have been imposed by brain asymmetry: a common and ancient trait that occurs in a wide range of vertebrates (Rogers and Andrew, 2002; Vallortigara and Rogers, 2005) and invertebrates (Anfora et al., 2011; Baracchi et al., 2018; Frasnelli et al., 2014, 2012), which possibly optimizes simultaneous processing of different kinds of information (Rogers et al., 2013). In natural environments, relevant numerosity (e.g., predators, food items, conspecifics) are intrinsically linked to their spatial arrangements, and their enumeration might be facilitated whenever a clear scanning directionality is present. Consequently, numerical estimation may have evolved in conjunction with spatial processing biases, leading to a left-anchored, rightward-directed environmental scanning strategy. This hypothesis suggests that numerical cognition evolved incorporating spatial processing biases and reflecting the spatial nature of ecologically relevant numerosity.

It should be noted that BAFT and the emotional valence models have been elaborated to explain the performance in a different task (Di Giorgio et al., 2019; Giurfa et al., 2022; Rugani et al., 2015a, 2020b). This required animals first to learn to associate a food reward with a central numerosity, e.g., an array depicting 5 dots. At test, when presented with new but identical numerosity, placed one on the left and one on the right, animals chose the left option when the test numerosity was smaller than the one experienced during learning, e.g., 2, and the right option when the numerosity was larger, e.g., 8 (Di Giorgio et al., 2019; Giurfa et al., 2022; Rugani et al., 2015a,

2020b). Such shift in the bias from left to right depending on numerosity cannot be simply associated with the lateralization of feeding responses typically guided by the left hemisphere in chicks (Deng and Rogers, 2002, 1997; Dharmaretnam and Rogers, 2005). Nevertheless, numerosities could correlate with spatial frequencies (Felisatti et al., 2020) or with emotional valence (Vallortigara, 2018). Remarkably, the right hemisphere dominance model provides a more comprehensive explanation. In fact, it accounts for the above mentioned task (Di Giorgio et al., 2019; Giurfa et al., 2022; Rugani et al., 2015a; Rugani and de Hevia, 2017) as well as for the ordinal task (Rugani et al., 2016; Rugani and Regolin, 2020).

Although all models (Felisatti et al., 2020; Rugani et al., 2016, 2015a; Vallortigara, 2018) identify the key role of hemispheric specialization in determining numerical spatialization, the Right-Hemisphere Dominance model can explain animal behavior across multiple contexts, providing a more parsimonious and generalizable explanation of cognitive processes, and potentially revealing fundamental connections between seemingly distinct cognitive domains.

Materials and Methods
Subject
We tested 100 male domestic chicks (*Gallus gallus*) of the Aviagen ROSS 308 line (experiment 1, $n = 48$; experiment 2, $n = 52$). We chose male chicks because of their superior response to food reinforcement compared to females (Regolin et al., 2005; Vallortigara et al., 1990) and significantly greater degree of lateralization in the thalamofugal pathway (Rajendra and Rogers, 1993). The fertilized eggs were obtained weekly from two local hatcheries (Agricola Berica, Montegalda, Vicenza, Italy, or Società Agricola La Pellegrina Spa, San Pietro in Gù, Padova, Italy). Eggs on the seventh or fourteenth day of incubation were delivered to the lab and placed in a FIEM incubator MG 70/100 ($45 \times 58 \times 43$ cm) at a controlled temperature of 36-38 °C and 60% humidity. On the eighteenth day of incubation, eggs were moved to a hatching machine ($60 \times 32 \times 40$ cm) at controlled temperature and humidity (36-38 °C; 60%) until the 21^{st} day of incubation (hatching day). Eggs were incubated under two conditions: in darkness to obtain Dark-incubated (weakly-lateralized) chicks (Di-chicks, $n = 24$ in experiment 1, $n = 26$ in experiment 2), and under light exposure using an LED 4.8W lightbulb to obtain Light-incubated (strongly-lateralized) chicks (Li-chicks, $n = 24$ in experiment 1, $n = 26$ in experiment 2). Only the chicks that completed all four tests were included in the final sample size. A few hours after hatching, chicks were feather-sexed and caged in pairs or triplets in standard metal cages ($28 \times 32 \times 40$ cm) with the floor covered with absorbent paper. The rearing room was maintained at a temperature of 28-31°C and humidity of about 60%. The cages were illuminated by neon lights (36 W) placed about 15 centimeters above each cage, with a standard 24-hour light-dark rearing cycle. Food (chick crumbles) and water were available in transparent glass jars (5×5 cm) *ad libitum*. Daily, chicks were familiarized and fed with some mealworms (*Tenebrio molitor larvae*) that were used as reinforcement during training. These rearing conditions were maintained until the initiation of the experimental protocol on Wednesday (8 a.m.), their third day of life, when food jars were removed from cages and chicks were isolated one per cage. Chicks underwent two hours of food deprivation before the start of each experimental session (shaping, training, and tests). Following the last testing session, chicks were rehoused in social groups with water and food *ad libitum*. On Friday afternoon, their fifth day of life, chicks were donated to local farmers (see Fig. 4 for a graphical illustration of the experimental schedule).

Apparatus
The experimental apparatus was located in the experimental room, near the rearing room, and maintained at constant temperature and humidity (28 °C; 70%). The apparatus consisted of a square arena constructed from green polypropylene ($100 \times 100 \times 40$ cm) with the floor covered with wood shavings (Fig. 1). Inside the arena were ten identical red plastic caps (3 cm in diameter, 0.9 cm in

height), each filled with wood shavings. To minimize the potential use of external cues, the entire setup was elevated and rotated randomly between trials or sessions (Rugani et al., 2007, 2010). Bottle caps were frequently shuffled to ensure that the choices made by the chicks did not depend on some unique characteristics of the caps (Rugani et al., 2016, 2011; Rugani and Regolin, 2021). The apparatus comprised two mirrored starting positions ($15 \times 15 \times 10$ cm boxes) outside opposing walls, one of which serves as a starting position (labeled as "S.P." in Fig. 1). The starting boxes were designed to allow consistent visual input of the inner apparatus to the chicks. Access to the arena was provided through an entrance door made of green polypropylene (10×17 cm), which could be lifted by a nylon thread.

Shaping

In both experiments, the shaping, pre-training, and training procedures were conducted in the same setting. Shaping started on Wednesday morning after 2 hours of food deprivation (8:00-10:00 am). This was essential in motivating foraging behavior during training and testing. During Shaping, the array made of 10 items was centrally aligned along the median sagittal line, thus sagittally oriented, with respect to either starting point (Fig. 1A), with the first cap positioned 28.5 cm from each entrance door. Each cap was positioned 1.44 cm apart from the subsequent one, with the overall array length being 43 cm. The array was situated 48.5 cm from either side wall.

The experimenter first introduced the chick into the arena for habituation, which lasted for about two minutes, allowing the bird to explore until it showed no signs of distress. Subsequently, the experimenter placed the chick into the starting point and the Shaping began. A piece of mealworm was placed and remained visible on the 4th cap to reinforce pecking behavior at that ordinal position. After the chick entered the arena and first pecked at any item, the trial was over, and the chick was immediately placed back into the starting box. If the chick did not peck at any item within 30 seconds, the experimenter used a metal stick to direct the chick to the fourth cap. After the chick had successfully pecked the 4th cap in 10 trials (whether consecutive or not), in the subsequent trials, the food was gradually covered with wood shavings until it was completely hidden. The shaping lasted for 10-15 minutes, followed by 30-40 minutes of rest back in the rearing cage with access to water but not to food.

Pre-Training

After the resting period, the Pre-Training began. From the Pre-Training all items looked identical as the food reward in the correct (4th) cap was completely buried in wood shavings. The chick had to complete three consecutive correct trials to reach the learning criterion and pass the re-training phase (Rugani et al., 2011, 2016; Rugani and Regolin, 2020, 2021). This usually took about 5-10 minutes. If the chick did not reach the learning criterion, the Pre-Training was repeated after 30 minutes of rest. If the chick again did not reach the criterion, it was excluded from the study.

Training

Immediately after completion of the Pre-Training, chicks underwent Training, comprising 20 trials, using the identical sequence employed in the Pre-Training phase. In each trial, only one choice was allowed, and the trial was terminated as soon as the chick pecked any item, with its choice being recorded. If the chick did not peck any item in 180 seconds, the trial was considered null and terminated. A choice was considered correct if the chick pecked at the 4th item. The learning criterion for passing the training phase was eight correct trials out of 20 trials (Rugani et al., 2011, 2016; Rugani and Regolin, 2020, 2021). If a subject did not achieve the learning criterion, after 40 minutes of rest it underwent another Pre-Training and, if this was successful the Training began. Each chick had three chances to pass the training criterion. During this phase, five chicks were excluded in Experiment 1 and nine in Experiment 2 due to motivational or health problems, an exclusion rate that aligns with those of previous studies involving similar procedures (Rugani et al.,

594 2020a, 2015b). These subjects were subsequently replaced to maintain the predetermined sample
595 size.
596
597 **Re-Training**
598 Re-Training was conducted prior to every test. The Re-Training procedure was the same as Pre-
599 Training and ended with three consecutive correct trials. Test sessions started immediately after
600 Re-Training was completed.
601
Test sessions
All chicks participated in all four tests: first the Sagittal Test, then the Fronto-Parallel Binocular
Test. Thereafter, monocular tests were administered in counter-balanced order. The Fronto-Parallel
Monocular Right Test was conducted prior to the Fronto-Parallel Monocular Left Test for $n = 24$
chicks in experiment 1 and $n = 26$ in experiment 2. The remaining chicks ($n = 24$ in experiment 1,
$n = 26$ in experiment 2) underwent the Fronto-Parallel Monocular Tests in the reversed order.
608
Sagittal Test
The procedure for the Sagittal Test (Fig. 1A) was the same as the Training. The Sagittal
Test consisted of 20 trials, and the time limit for each trial was 60 seconds. During Testing, food
reinforcement was available only in pre-established trials to prevent the extinction of responses
over multiple unrewarded test trials (reinforced trials: 4, 5, 7, 10, 13, 14, 16 and 19 (Rugani et al.,
2016, 2011; Rugani and Regolin, 2021, 2020)). Thereafter, subjects rested for at least 60 minutes
before entering the Fronto-Parallel Tests.
In experiment 1 ordinal and spatial cues were available to identify the 4th correct item, in
fact the array was arranged as during Training, with the length of the series being kept constant
throughout the trials.
In experiment 2, to eliminate spatial cues to locate the 4th item, the inter-item distance varied
between test trials (1.44 cm, 2.55 cm, 3.11 cm, and 3.66 cm), while remaining equally-spaced within
each trial, resulting in total array lengths of 43.0, 53.0, 58.0, and 63.0 cm, respectively. The first
cap was set at 28.5 cm from the starting position.
623
Fronto-parallel Tests
In both experiments, the Fronto-Parallel Test was conducted on each subject in three different
conditions of vision (Fig. 1). The Binocular Fronto-Parallel Test was always administered first.
Then half of the chicks underwent the Left Monocular Fronto-Parallel Test and finally the Right
Monocular Fronto-Parallel Test, while the other half underwent the monocular tests in reverse
order.
In the Monocular Fronto-Parallel Test, a temporary eye patch was carefully applied to
restrict visual input to one of the chicks' eyes. The patch, made of removable paper tape, did not
obstruct eyelid movements and enabled smooth removal post testing without harming the subject.
Before the actual test, chicks were habituated to wear the eye patch for about 15 minutes, during
which they were closely monitored; any signs of distress or excessive scratching prompted
immediate intervention to ensure the animals' well-being (Rogers and Vallortigara, 2017; Rugani
et al., 2016; Rugani and Regolin, 2021).
In the three Fronto-Parallel Tests, the array was rotated 90°, fronto-parallel with respect to
the Starting Point (Fig. 1B). Inter-item distances matched those in the Sagittal Test for each
experiment. In this rotated array, both the 4th item from left (4L) and right (4R) were considered
correct and rewarded during the pre-established trials (as described for the Sagittal Test). Each
Fronto-Parallel Test comprised 20 trials, with two-hour rest periods between tests.
642
Statistical analyses

In each trial, chicks were allowed a single peck. We recorded the selected item to calculate the
percentage of responses at each position as [(number of pecks to a given item ÷ total number valid
trials) × 100] and averaged them separately for each group and test. We employed both frequentist
and Bayesian statistical approaches, conducting corresponding Bayesian analyses for each
frequentist test. We analyzed the group percentage for choosing each item above chance (10%),
using Wilcoxon one-sample signed-rank tests with Bonferroni correction for multiple comparisons
(data and significant results are reported in Table S1 and Table S2), and one-sample Bayesian t-
tests. To assess side bias in the Fronto-Parallel tests, we compared correct choices on the left (4L)
versus the right (4R) using paired *t* tests, with Cohen's *d* as the effect size, and Bonferroni as the
correction method; moreover, we conducted two-sample Bayesian *t* tests. Additionally, we tested
whether brain lateralization influenced accuracy by comparing the percentage of correct choices
(i.e., the selection of the 4th item in the Sagittal test, and of the 4L or 4R items in the Fronto-Parallel
tests) between Li-chicks and Di-chicks using two-sample *t* tests, with Cohen's *d* as the effect size,
and Bonferroni as the correction method, additionally we conducted two-sample Bayesian t-tests.
Bayesian factors were computed using BayesFactor package (Morey and Rouder, 2012).
The analyses were conducted using R (version 4.3.1; R Core Team, 2022). We used the
classification by Lee & Wagenmakers (2014) (Lee and Wagenmakers, 2014) to interpret BFs.
661
662
663
664

Fig. 4 Time schedule of the experiments. During the last three days of incubation, eggs were incubated either in darkness (weakly lateralized chicks) or under light exposure (strongly lateralized chicks). When chicks were three days old, the shaping procedure started, followed by Pre-Training and Training. In the following days, chicks were re-trained and then underwent all four tests.

Table 2. Descriptive statistics. For each test in the two experiments, the accuracy of selecting the 4th item in the Sagittal test and the 4L or 4R items in the Fronto-Parallel (FP) tests is reported.

Experiment 1: Ordinal and Spatial cue									
Test	**Hatch Condition**	**Choice**	**mean**	**SD**	**SE**	***n***	***r***	***P***	**BF**
Sagittal	**Di-chicks**	4	28.436	9.374	1.913	24	0.878	< 0.001	>10000
	Li-chicks	4	36.382	8.164	1.666	24	0.88	< 0.001	>10000
FP Binocular	**Di-chicks**	4L	18.958	10.527	2.149	24	0.727	0.005	169
		4R	19.792	11.371	2.321	24	0.724	0.003	188
	Li-chicks	4L	28.851	12.598	2.572	24	0.859	< 0.001	>10000
		4R	11.894	7.503	1.532	24	0.337	0.798	0.741
FP Monocular Left	**Di-chicks**	4L	12.780	8.309	1.696	24	0.376	0.532	1.287
		4R	3.990	6.154	1.256	24	-0.708	1.000	0.05
	Li-chicks	4L	22.595	8.736	1.783	24	0.876	< 0.001	>10000
		4R	6.297	6.747	1.377	24	-0.465	1.000	0.068
FP Monocular Right	**Di-chicks**	4L	3.746	3.668	0.749	24	-0.798	1.000	0.013
		4R	13.417	8.781	1.792	24	0.406	0.398	1.936
	Li-chicks	4L	5.219	6.356	1.297	24	-0.594	1.000	0.057
		4R	21.484	10.734	2.191	24	0.807	0.001	1843
Experiment 2: Ordinal cue only									
Test	**Hatch Condition**	**Choice**	**mean**	**SD**	**SE**	***n***	***r***	***P***	**BF**
Sagittal	**Di-chicks**	4	31.487	15.337	3.008	26	0.863	< 0.001	>10000
	Li-chicks	4	32.275	11.232	2.203	26	0.876	< 0.001	>10000
FP Binocular	**Di-chicks**	4L	18.725	9.246	1.813	26	0.74	0.001	836.246
		4R	19.666	10.024	1.966	26	0.786	0.002	1070.73
	Li-chicks	4L	21.771	9.888	1.939	26	0.819	< 0.001	>10000
		4R	15.800	9.239	1.812	26	0.603	0.024	21.919
FP Monocular Left	**Di-chicks**	4L	14.835	9.878	1.937	26	0.473	0.122	5.337
		4R	8.333	6.912	1.356	26	-0.316	1.000	0.102
	Li-chicks	4L	15.011	9.344	1.832	26	0.518	0.050	8.449
		4R	7.939	6.082	1.193	26	-0.302	1.000	0.084
FP Monocular Right	**Di-chicks**	4L	8.290	7.659	1.502	26	-0.269	1.000	0.102
		4R	12.457	9.255	1.815	26	0.29	0.850	0.839
	Li-chicks	4L	6.288	6.151	1.206	26	-0.515	1.000	0.06
		4R	17.984	9.634	1.889	26	0.78	0.001	214

Supplementary Materials:
The PDF file includes:
Tables S1 and S2

References

Adachi I. 2014. Spontaneous Spatial Mapping of Learned Sequence in Chimpanzees: Evidence for a SNARC-Like
    Effect. *PLoS ONE* **9**:e90373. doi:10.1371/journal.pone.0090373
Anfora G, Rigosi E, Frasnelli E, Ruga V, Trona F, Vallortigara G. 2011. Lateralization in the Invertebrate Brain:
    Left-Right Asymmetry of Olfaction in Bumble Bee, Bombus terrestris. *PLoS ONE* **6**:e18903.
    doi:10.1371/journal.pone.0018903
Bächtold D, Baumüller M, Brugger P. 1998. Stimulus-response compatibility in representational space.
    Neuropsychologia **36**:731–735. doi:10.1016/s0028-3932(98)00002-5
Baracchi D, Rigosi E, De Brito Sanchez G, Giurfa M. 2018. Lateralization of Sucrose Responsiveness and Non-
    associative Learning in Honeybees. *Front Psychol* **9**:425. doi:10.3389/fpsyg.2018.00425
Beran MJ, French K, Smith TR, Parrish AE. 2019. Limited evidence of number–space mapping in rhesus monkeys
    (*Macaca mulatta*) and capuchin monkeys (*Sapajus apella*). *Journal of Comparative Psychology* **133**:281–
    293. doi:10.1037/com0000177
Bertillon J. 1880. Sur la vision de la série des nombres. *La Nature* **378**:196–98. doi:10.3406/bmsap.1880.3301
Bulf H, de Hevia MD, Macchi Cassia V. 2016. Small on the left, large on the right: numbers orient visual attention
    onto space in preverbal infants. *Dev Sci* **19**:394–401. doi:10.1111/desc.12315
Cantlon JF, Brannon EM. 2007. Basic Math in Monkeys and College Students. *PLoS Biol* **5**:e328.
    doi:10.1371/journal.pbio.0050328
Clark WJ, Colombo M. 2020. The functional architecture, receptive field characteristics, and representation of objects
    in the visual network of the pigeon brain. *Progress in Neurobiology* **195**:101781.
    doi:10.1016/j.pneurobio.2020.101781
Cordes S, Brannon EM. 2009. Crossing the divide: Infants discriminate small from large numerosities.
    Developmental Psychology **45**:1583–1594. doi:10.1037/a0015666
Cordes S, Gelman R, Gallistel CR, Whalen J. 2001. Variability signatures distinguish verbal from nonverbal counting
    for both large and small numbers. *Psychonomic Bulletin & Review* **8**:698–707. doi:10.3758/BF03196206
Costalunga G, Kobylkov D, Rosa-Salva O, Vallortigara G, Mayer U. 2022. Light-incubation effects on lateralisation
    of single unit responses in the visual Wulst of domestic chicks. *Brain Struct Funct* **227**:497–513.
    doi:10.1007/s00429-021-02259-y
Dadda M, Agrillo C, Bisazza A, Brown C. 2015. Laterality enhances numerical skills in the guppy, Poecilia
    reticulata. *Front Behav Neurosci* **9**:285. doi:10.3389/fnbeh.2015.00285
Daisley JN, Mascalzoni E, Rosa-Salva O, Rugani R, Regolin L. 2009. Lateralization of social cognition in the
    domestic chicken (*Gallus gallus*). *Phil Trans R Soc B* **364**:965–981. doi:10.1098/rstb.2008.0229
Daisley JN, Vallortigara G, Regolin L. 2010. Logic in an asymmetrical (social) brain: Transitive inference in the
    young domestic chick. *Social Neuroscience* **5**:309–319. doi:10.1080/17470910903529795
Davidson RJ. 2004. Well–being and affective style: neural substrates and biobehavioural correlates. *Phil Trans R Soc
    Lond B* **359**:1395–1411. doi:10.1098/rstb.2004.1510
de Hevia MD. 2021. How the Human Mind Grounds Numerical Quantities on Space. *Child Dev Perspect* **15**:44–50.
    doi:10.1111/cdep.12398
de Hevia MD, Izard V, Coubart A, Spelke ES, Streri A. 2014. Representations of space, time, and number in
    neonates. *Proc Natl Acad Sci USA* **111**:4809–4813. doi:10.1073/pnas.1323628111
de Hevia MD, Spelke ES. 2010. Number-Space Mapping in Human Infants. *Psychol Sci* **21**:653–660.
    doi:10.1177/0956797610366091
de Hevia MD, Veggiotti L, Streri A, Bonn CD. 2017. At Birth, Humans Associate "Few" with Left and "Many" with
    Right. *Current Biology* **27**:3879-3884.e2. doi:10.1016/j.cub.2017.11.024
Dehaene S. 2011. The number sense: how the mind creates mathematics, Rev. and updated ed. ed. New York: Oxford
    University Press.
Dehaene S, Bossini S, Giraux P. 1993. The mental representation of parity and number magnitude. *Journal of
    Experimental Psychology: General* **122**:371–396. doi:10.1037/0096-3445.122.3.371
Deng C, Rogers LJ. 2002. Prehatching visual experience and lateralization in the visual Wulst of the chick.
    Behavioural Brain Research **134**:375–385. doi:10.1016/S0166-4328(02)00050-5

Deng C, Rogers LJ. 1998. Bilaterally projecting neurons in the two visual pathways of chicks. *Brain Research*
794:281–290. doi:10.1016/S0006-8993(98)00237-6
Deng C, Rogers LJ. 1997. Differential contributions of the two visual pathways to functional lateralization in chicks.
Behavioural Brain Research **87**:173–182. doi:10.1016/S0166-4328(97)02276-6
Dharmaretnam M, Rogers LJ. 2005. Hemispheric specialization and dual processing in strongly versus weakly
lateralized chicks. *Behavioural Brain Research* **162**:62–70. doi:10.1016/j.bbr.2005.03.012
Di Giorgio E, Lunghi M, Rugani R, Regolin L, Dalla Barba B, Vallortigara G, Simion F. 2019. A mental number line
in human newborns. *Dev Sci* **22**. doi:10.1111/desc.12801
Diekamp B, Regolin L, Güntürkün O, Vallortigara G. 2005. A left-sided visuospatial bias in birds. *Current Biology*
15:R372–R373. doi:10.1016/j.cub.2005.05.017
Ditz HM, Nieder A. 2015. Neurons selective to the number of visual items in the corvid songbird endbrain. *Proc Natl*
Acad Sci USA **112**:7827–7832. doi:10.1073/pnas.1504245112
Drucker CB, Brannon EM. 2014. Rhesus monkeys (*Macaca mulatta*) map number onto space. *Cognition* **132**:57–67.
doi:10.1016/j.cognition.2014.03.011
Felisatti A, Laubrock J, Shaki S, Fischer MH. 2020. A biological foundation for spatial–numerical associations: the
brain's asymmetric frequency tuning. *Annals of the New York Academy of Sciences* **1477**:44–53.
doi:10.1111/nyas.14418
Frasnelli E, Haase A, Rigosi E, Anfora G, Rogers L, Vallortigara G. 2014. The Bee as a Model to Investigate Brain
and Behavioural Asymmetries. *Insects* **5**:120–138. doi:10.3390/insects5010120
Frasnelli E, Vallortigara G, Rogers LJ. 2012. Left–right asymmetries of behaviour and nervous system in
invertebrates. *Neuroscience & Biobehavioral Reviews* **36**:1273–1291. doi:10.1016/j.neubiorev.2012.02.006
Galton F. 1880. Visualised Numerals. *Nature* **21**:252–256. doi:10.1038/021252a0
Gatto E, Agrillo C, Brown C, Dadda M. 2019. Individual differences in numerical skills are influenced by brain
lateralization in guppies (*Poecilia reticulata*). *Intelligence*, Intelligence in non human animals **74**:12–17.
doi:10.1016/j.intell.2019.04.004
Gazzaniga MS. 2000. Cerebral specialization and interhemispheric communication: Does the corpus callosum enable
the human condition? *Brain* **123**:1293–1326. doi:10.1093/brain/123.7.1293
Ghirlanda S, Vallortigara G. 2004. The evolution of brain lateralization: a game-theoretical analysis of population
structure. *Proc R Soc Lond B* **271**:853–857. doi:10.1098/rspb.2003.2669
Giurfa M, Marcout C, Hilpert P, Thevenot C, Rugani R. 2022. An insect brain organizes numbers on a left-to-right
mental number line. *Proc Natl Acad Sci USA* **119**:e2203584119. doi:10.1073/pnas.2203584119
Güntürkün O, Ströckens F, Ocklenburg S. 2020. Brain Lateralization: A Comparative Perspective. *Physiological*
Reviews **100**:1019–1063. doi:10.1152/physrev.00006.2019
Harvey BM, Klein BP, Petridou N, Dumoulin SO. 2013. Topographic Representation of Numerosity in the Human
Parietal Cortex. *Science* **341**:1123–1126. doi:10.1126/science.1239052
Karolis V, Iuculano T, Butterworth B. 2011. Mapping numerical magnitudes along the right lines: differentiating
between scale and bias. *J Exp Psychol Gen* **140**:693–706. doi:10.1037/a0024255
Kobylkov D, Mayer U, Zanon M, Vallortigara G. 2022. Number neurons in the nidopallium of young domestic
chicks. *Proc Natl Acad Sci USA* **119**:e2201039119. doi:10.1073/pnas.2201039119
Lee MD, Wagenmakers E-J. 2014. Bayesian Cognitive Modeling: A Practical Course, 1st ed. Cambridge University
Press. doi:10.1017/CBO9781139087759
McCrink K, Veggiotti L, Hevia MD. 2020. A left visual advantage for quantity processing in neonates. *Ann NY Acad*
Sci **1477**:71–78. doi:10.1111/nyas.14457
Morey RD, Rouder JN. 2012. BayesFactor: Computation of Bayes Factors for Common Designs.
doi:10.32614/CRAN.package.BayesFactor
Nieder A. 2002. Representation of the Quantity of Visual Items in the Primate Prefrontal Cortex. *Science* **297**:1708–
1711. doi:10.1126/science.1072493
Pearson JM, Hayden BY, Raghavachari S, Platt ML. 2009. Neurons in Posterior Cingulate Cortex Signal Exploratory
Decisions in a Dynamic Multioption Choice Task. *Current Biology* **19**:1532–1537.
doi:10.1016/j.cub.2009.07.048
Philippi T, Seger J. 1989. Hedging one's evolutionary bets, revisited. *Trends in Ecology & Evolution* **4**:41–44.
doi:10.1016/0169-5347(89)90138-9
Piazza M, Eger E. 2016. Neural foundations and functional specificity of number representations. *Neuropsychologia*
83:257–273. doi:10.1016/j.neuropsychologia.2015.09.025
Piazza M, Izard V, Pinel P, Le Bihan D, Dehaene S. 2004. Tuning Curves for Approximate Numerosity in the Human
Intraparietal Sulcus. *Neuron* **44**:547–555. doi:10.1016/j.neuron.2004.10.014
Rajendra S, Rogers LesleyJ. 1993. Asymmetry is present in the thalamofugal visual projections of female chicks. *Exp*
Brain Res **92**. doi:10.1007/BF00229044
Rashid N, Andrew RJ. 1989. Right hemisphere advantage for topographical orientation in the domestic chick.
Neuropsychologia **27**:937–948. doi:10.1016/0028-3932(89)90069-9

Regolin L, Garzotto B, Rugani R, Pagni P, Vallortigara G. 2005. Working memory in the chick: parallel and
lateralized mechanisms for encoding of object- and position-specific information. *Behavioural Brain*
Research **157**:1–9. doi:10.1016/j.bbr.2004.06.012
Robert F, Cuénod M M. 1969. Electrophysiology of the intertectal commissures in the pigeon I. Analysis of the
pathways. *Exp Brain Res* **9**. doi:10.1007/BF00238325
Rogers LJ. 2021. Brain Lateralization and Cognitive Capacity. *Animals* **11**:1996. doi:10.3390/ani11071996
Rogers LJ. 1982. Light experience and asymmetry of brain function in chickens. *Nature* **297**:223–225.
doi:10.1038/297223a0
Rogers LJ, Andrew R, editors. 2002. Comparative Vertebrate Lateralization, 1st ed. Cambridge University Press.
doi:10.1017/CBO9780511546372
Rogers LJ, Bolden SW. 1991. Light-dependent development and asymmetry of visual projections. *Neuroscience*
Letters **121**:63–67. doi:10.1016/0304-3940(91)90650-I
Rogers LJ, Deng C. 1999. Light experience and lateralization of the two visual pathways in the chick. *Behavioural*
Brain Research **98**:277–287. doi:10.1016/S0166-4328(98)00094-1
Rogers LJ, Sink HS. 1988. Transient asymmetry in the projections of the rostral thalamus to the visual hyperstriatum
of the chicken, and reversal of its direction by light exposure. *Exp Brain Res* **70**. doi:10.1007/BF00248362
Rogers LJ, Vallortigara G. 2021. Brain and behavioural asymmetries in non-human species. *Laterality* **26**:v–vii.
doi:10.1080/1357650X.2021.1891675
Rogers LJ, Vallortigara G, editors. 2017. Lateralized Brain Functions: Methods in Human and Non-Human Species,
Neuromethods. New York, NY: Springer New York. doi:10.1007/978-1-4939-6725-4
Rogers LJ, Vallortigara G, Andrew RJ. 2013. Divided Brains: The Biology and Behaviour of Brain Asymmetries, 1st
ed. Cambridge University Press. doi:10.1017/CBO9780511793899
Rugani R. 2018. Towards numerical cognition's origin: insights from day-old domestic chicks. *Phil Trans R Soc B*
373:20160509. doi:10.1098/rstb.2016.0509
Rugani R, de Hevia M-D. 2017. Number-space associations without language: Evidence from preverbal human
infants and non-human animal species. *Psychon Bull Rev* **24**:352–369. doi:10.3758/s13423-016-1126-2
Rugani R, Kelly DM, Szelest I, Regolin L, Vallortigara G. 2010. Is it only humans that count from left to right? *Biol*
Lett **6**:290–292. doi:10.1098/rsbl.2009.0960
Rugani R, Loconsole M, Simion F, Regolin L. 2020a. Individually distinctive features facilitate numerical
discrimination of sets of objects in domestic chicks. *Sci Rep* **10**:16408. doi:10.1038/s41598-020-73431-3
Rugani R, Platt ML, Zhang Y, Brannon EM. 2024. Magnitude shifts spatial attention from left to right in rhesus
monkeys as in the human mental number line. *iScience* **27**:108866. doi:10.1016/j.isci.2024.108866
Rugani R, Regolin L. 2021. Approach direction and accuracy, but not response times, show spatial-numerical
association in chicks. *PLoS ONE* **16**:e0257764. doi:10.1371/journal.pone.0257764
Rugani R, Regolin L. 2020. Hemispheric specialization in spatial versus ordinal processing in the day-old domestic
chick (*Gallus gallus*). *Ann NY Acad Sci* **1477**:34–43. doi:10.1111/nyas.14345
Rugani R, Regolin L, Vallortigara G. 2007. Rudimental numerical competence in 5-day-old domestic chicks (*Gallus*
gallus): Identification of ordinal position. *Journal of Experimental Psychology: Animal Behavior Processes*
33:21–31. doi:10.1037/0097-7403.33.1.21
Rugani R, Vallortigara G, Priftis K, Regolin L. 2020b. Numerical magnitude, rather than individual bias, explains
spatial numerical association in newborn chicks. *eLife* **9**:e54662. doi:10.7554/eLife.54662
Rugani R, Vallortigara G, Priftis K, Regolin L. 2015a. Number-space mapping in the newborn chick resembles
humans' mental number line. *Science* **347**:534–536. doi:10.1126/science.aaa1379
Rugani R, Vallortigara G, Regolin L. 2016. Mapping number to space in the two hemispheres of the avian brain.
Neurobiology of Learning and Memory **133**:13–18. doi:10.1016/j.nlm.2016.05.010
Rugani R, Vallortigara G, Regolin L. 2015b. The use of proportion by young domestic chicks (*Gallus gallus*). *Anim*
Cogn **18**:605–616. doi:10.1007/s10071-014-0829-x
Rugani R, Vallortigara G, Vallini B, Regolin L. 2011. Asymmetrical number-space mapping in the avian brain.
Neurobiology of Learning and Memory **95**:231–238. doi:10.1016/j.nlm.2010.11.012
Rugani R, Zhang Y, Ahmed N, Brannon E. 2022. Children perform better on left than right targets in an ordinal task.
Acta Psychologica **226**:103560. doi:10.1016/j.actpsy.2022.103560
Seron X, Pesenti M, Noël M-P, Deloche G, Cornet J-A. 1992. Images of numbers: or "When 98 is upper left and 6
sky blue." *Cognition* **44**:159–196. doi:10.1016/0010-0277(92)90053-K
Shaki S, Fischer MH, Petrusic WM. 2009. Reading habits for both words and numbers contribute to the SNARC
effect. *Psychonomic Bulletin & Review* **16**:328–331. doi:10.3758/PBR.16.2.328
Simons AM. 2011. Modes of response to environmental change and the elusive empirical evidence for bet hedging.
Proc R Soc B **278**:1601–1609. doi:10.1098/rspb.2011.0176
Tang J, Ward J, Butterworth B. 2008. Number Forms in the Brain. *Journal of Cognitive Neuroscience* **20**:1547–1556.
doi:10.1162/jocn.2008.20120
Theiss MPH, Hellmann B, Güntürkün O. 2003. The architecture of an inhibitory sidepath within the avian tectofugal
system: *NeuroReport* **14**:879–882. doi:10.1097/00001756-200305060-00021

Triki Z, Bshary R. 2018. Cleaner fish Labroides dimidiatus discriminate numbers but fail a mental number line test.
Anim Cogn **21**:99–107. doi:10.1007/s10071-017-1143-1
Vallortigara G. 2018. Comparative cognition of number and space: the case of geometry and of the mental number
line. *Phil Trans R Soc B* **373**:20170120. doi:10.1098/rstb.2017.0120
Vallortigara G, Cailotto M, Zanforlin M. 1990. Sex differences in social reinstatement motivation of the domestic
chick (*Gallus gallus*) revealed by runway tests with social and nonsocial reinforcement. *Journal of
Comparative Psychology* **104**:361–367. doi:10.1037/0735-7036.104.4.361
Vallortigara G, Rogers LJ. 2005. Survival with an asymmetrical brain: advantages and disadvantages of cerebral
lateralization. *Behav Brain Sci* **28**:575–589. doi:10.1017/S0140525X05000105
Weidner C, Repérant J, Miceli D, Haby M, Rio JP. 1985. An anatomical study of ipsilateral retinal projections in the
quail using radioautographic, horseradish peroxidase, fluorescence and degeneration techniques. *Brain
Research* **340**:99–108. doi:10.1016/0006-8993(85)90778-4
West E, McCrink K. 2021. Eye Tracking Lateralized Spatial Associations in Early Childhood. *Journal of Cognition
and Development* **22**:678–694. doi:10.1080/15248372.2021.1926254
Zebian S. 2005. Linkages between Number Concepts, Spatial Thinking, and Directionality of Writing: The SNARC
Effect and the REVERSE SNARC Effect in English and Arabic Monoliterates, Biliterates, and Illiterate
Arabic Speakers. *J Cogn Cult* **5**:165–190. doi:10.1163/1568537054068660
Zorzi M, Priftis K, Umiltà C. 2002. Neglect disrupts the mental number line. *Nature* **417**:138–139.
doi:10.1038/417138a

Acknowledgments

Funding:
European's Union Horizon 2020 Research and Innovation program under the Marie Sklodowska-Curie Grant/Award Number: 795242 to R. R.; PRIN 2022, grant/award number: 202254RHRT to R. R.;
PRIN 2022 PNRR, grant/award number: P2022TKY7B, to R. R. and L. R.; PRIN 2017: grant/award number: 2017PSRHPZ_003, to L. R. and R. R..

Author contributions:
Conceptualization: RR, and LR.
Methodology: RR, LR., and MM
Investigation: RR, YZ, and MM
Data analysis: YZ, MM, and RR
Writing—original draft: RR
Writing—review and editing: LR, MM, and YZ

Competing interests: The authors declare that they have no competing interests.

Data and materials availability: All data are available in the main text or the supplementary materials. Additional material containing metadata, row data, script used for the analysis, output of the script, and a folder with the same materials available for macOS for this article is available at: Hatching with Numbers: How Pre-natal Experience Affects Chicks' Left-to-Right Mental Number Line - Research Data Unipd
URI: https://researchdata.cab.unipd.it/id/eprint/1424

Ethics: All experimental procedures employed were evaluated, approved, and conducted in strict adherence to the guidelines provided by the Committee for Animal Welfare of the University of Padua, the Ethical Committee of the University of Padova for Animal Experimentation and the Ministry of Health of the Italian Republic (Prot. N.9245, 17/01/2019). This comprehensive compliance addressed both national and European directives concerning animal research.

Table S1. Descriptive statistic experiment 1. Descriptive statistics of the accuracy in selecting each of the 10 items in the Sagittal test, and in selecting the 5 items on the left (L) and the 5 items on the right (R) in the Fronto-Parallel (FP) tests of Experiment 1.

Test	Hatch Condition	Choice	Mean	SD	SE	n	r	P	BF
Experiment 1: Ordinal and Spatial cue									
Sagittal	Di-chick	1	**17.152**	**10.579**	**2.159**	**24**	**0.651**	**0.019**	**26.604**
		2	**15.378**	**8.408**	**1.716**	**24**	**0.682**	**0.027**	**18.426**
		3	11.294	7.015	1.432	24	0.227	1.000	0.497
		4	**28.436**	**9.374**	**1.913**	**24**	**0.878**	**< 0.001**	**>10000**
		5	7.986	5.794	1.183	24	-0.345	1.000	0.089
		6	7.326	6.902	1.409	24	-0.395	1.000	0.083
		7	5.914	5.176	1.057	24	-0.732	1.000	0.055
		8	2.546	3.999	0.816	24	-0.917	1.000	0.041
		9	1.678	2.424	0.495	24	-0.91	1.000	0.319
		10	2.292	3.290	0.672	24	-0.912	1.000	0.040
	Li-chick	1	14.649	8.392	1.713	24	0.569	0.070	8.031
		2	**18.629**	**8.566**	**1.748**	**24**	**0.828**	**0.002**	**930.306**
		3	9.002	6.779	1.384	24	-0.0943	1.000	0.135
		4	**36.382**	**8.164**	**1.666**	**24**	**0.88**	**< 0.001**	**>10000**
		5	4.974	4.596	0.938	24	-0.783	1.000	0.048
		6	5.914	6.701	1.368	24	-0.57	1.000	0.064
		7	4.583	6.743	1.376	24	-0.704	1.000	0.055
		8	2.511	4.669	0.953	24	-0.822	1.000	0.043
		9	2.094	2.930	0.598	24	-0.909	1.000	0.042
		10	1.261	2.232	0.456	24	-0.923	1.000	4.647
FP Binocular	Di-chick	1L	6.458	8.272	1.689	24	-0.473	1.000	0.078
		1R	6.250	5.566	1.136	24	-0.591	1.000	0.060
		2L	9.583	7.790	1.590	24	-0.058	1.000	0.178
		2R	5.833	6.019	1.229	24	-0.626	1.000	0.059
		3L	7.292	5.103	1.042	24	-0.539	1.000	0.069
		3R	6.458	5.610	1.145	24	-0.626	1.000	0.062
		4L	**18.958**	**10.527**	**2.149**	**24**	**0.727**	**0.005**	**168.591**

		4R	**19.792**	**11.371**	**2.321**	**24**	**0.724**	**0.003**	**188.153**
		5L	9.792	6.164	1.258	24	-0.006	1.000	0.190
		5R	9.583	5.882	1.201	24	-0.127	1.000	0.169
	Li-chick	1L	5.095	5.598	1.143	24	-0.704	1.000	0.053
		1R	7.350	6.753	1.378	24	-0.452	1.000	0.082
		2L	8.023	7.186	1.467	24	-0.342	1.000	0.101
		2R	7.164	6.443	1.315	24	-0.365	1.000	0.077
		3L	6.576	5.782	1.180	24	-0.57	1.000	0.065
		3R	7.785	7.907	1.614	24	-0.277	1.000	0.100
		4L	**28.851**	**12.598**	**2.572**	**24**	**0.859**	**< 0.001**	**>10000**
		4R	11.894	7.503	1.532	24	0.337	0.798	0.741
		5L	10.180	7.009	1.431	24	0.026	1.000	0.237
		5R	7.081	6.262	1.278	24	-0.436	1.000	0.074
FP Monocular Left	Di-chick	**1L**	**31.591**	**18.397**	**3.755**	**24**	**0.826**	**< 0.001**	**5702.882**
		1R	5.716	9.515	1.942	24	-0.449	1.000	0.076
		2L	**17.327**	**9.620**	**1.964**	**24**	**0.683**	**0.009**	**64.802**
		2R	2.775	4.944	1.009	24	-0.813	1.000	0.044
		3L	11.907	9.773	1.995	24	0.189	1.000	0.527
		3R	2.348	3.652	0.745	24	-0.859	1.000	0.040
		4L	12.780	8.309	1.696	24	0.376	0.532	1.287
		4R	3.990	6.154	1.256	24	-0.708	1.000	0.050
		5L	6.625	7.226	1.475	24	-0.446	1.000	0.074
		5R	4.940	4.561	0.931	24	-0.803	1.000	0.048
	Li-chick	**1L**	**23.035**	**10.533**	**2.150**	**24**	**0.863**	**< 0.001**	**>10000**
		1R	6.256	6.615	1.350	24	-0.544	1.000	0.066
		2L	11.109	7.272	1.484	24	0.171	1.000	0.419
		2R	3.244	4.112	0.839	24	-0.851	1.000	0.042
		3L	12.500	7.108	1.451	24	0.357	0.455	1.457
		3R	3.643	5.890	1.202	24	-0.728	1.000	0.048
		4L	**22.595**	**8.736**	**1.783**	**24**	**0.876**	**< 0.001**	**>10000**
		4R	6.297	6.747	1.377	24	-0.465	1.000	0.068
		5L	7.578	7.194	1.468	24	-0.379	1.000	0.090

Test	Hatch Condition	Choice	Mean	SD	SE	n	r	P	BF
		5R	3.743	4.773	0.974	24	-0.803	1.000	0.045
FP Monocular Right	Di-chick	1L	3.991	4.710	0.961	24	-0.823	1.000	0.045
		1R	**33.744**	**16.545**	**3.377**	**24**	**0.861**	**< 0.001**	**90653.398**
		2L	2.763	4.737	0.967	24	-0.872	1.000	0.043
		2R	15.779	12.802	2.613	24	0.411	0.255	3.214
		3L	4.083	5.947	1.214	24	-0.729	1.000	0.050
		3R	9.537	8.191	1.672	24	-0.0384	1.000	0.176
		4L	3.746	3.668	0.749	24	-0.798	1.000	0.042
		4R	13.417	8.781	1.792	24	0.406	0.398	1.936
		5L	5.033	5.004	1.022	24	-0.716	1.000	0.050
		5R	7.908	6.329	1.292	24	-0.304	1.000	0.091
	Li-chick	1L	5.375	7.537	1.539	24	-0.613	1.000	0.063
		1R	**23.770**	**14.388**	**2.937**	**24**	**0.765**	**0.002**	**535.661**
		2L	4.345	5.663	1.156	24	-0.768	1.000	0.050
		2R	13.847	9.447	1.928	24	0.419	0.287	2.233
		3L	4.405	5.204	1.062	24	-0.759	1.000	0.048
		3R	11.261	5.861	1.196	24	0.293	0.984	0.591
		4L	5.219	6.356	1.297	24	-0.594	1.000	0.057
		4R	**21.484**	**10.734**	**2.191**	**24**	**0.807**	**0.001**	**1842.938**
		5L	5.994	4.638	0.947	24	-0.685	1.000	0.053
		5R	4.300	4.755	0.971	24	-0.772	1.000	0.046

896
897

Table S2. Descriptive statistic experiment 2. Descriptive statistics of the accuracy in selecting each of the 10 items in the Sagittal test, and in selecting the 5 items on the left (L) and the 5 items on the right (R) in the Fronto-Parallel (FP) tests of Experiment 2

Experiment 2: Ordinal cue only									
Test	Hatch Condition	Choice	Mean	SD	SE	n	r	P	BF
Sagittal	Di-chick	**1**	**17.152**	**10.579**	**2.159**	**24**	**0.651**	**0.019**	**26.604**
		2	**15.378**	**8.408**	**1.716**	**24**	**0.682**	**0.027**	**18.426**
		3	11.294	7.015	1.432	24	0.227	1.000	0.497
		4	**28.436**	**9.374**	**1.913**	**24**	**0.878**	**< 0.001**	**>10000**
		5	7.986	5.794	1.183	24	-0.345	1.000	0.089

		6	7.326	6.902	1.409	24	-0.395	1.000	0.083
		7	5.914	5.176	1.057	24	-0.732	1.000	0.055
		8	2.546	3.999	0.816	24	-0.917	1.000	0.041
		9	1.678	2.424	0.495	24	-0.91	1.000	0.319
		10	2.292	3.290	0.672	24	-0.912	1.000	0.040
	Li-chick	1	14.649	8.392	1.713	24	0.569	0.070	8.031
		2	**18.629**	**8.566**	**1.748**	**24**	**0.828**	**0.002**	**930.306**
		3	9.002	6.779	1.384	24	-0.0943	1.000	0.135
		4	**36.382**	**8.164**	**1.666**	**24**	**0.88**	**< 0.001**	**>10000**
		5	4.974	4.596	0.938	24	-0.783	1.000	0.048
		6	5.914	6.701	1.368	24	-0.57	1.000	0.064
		7	4.583	6.743	1.376	24	-0.704	1.000	0.055
		8	2.511	4.669	0.953	24	-0.822	1.000	0.043
		9	2.094	2.930	0.598	24	-0.909	1.000	0.042
		10	1.261	2.232	0.456	24	-0.923	1.000	4.647
FP Binocular	Di-chick	1L	6.458	8.272	1.689	24	-0.473	1.000	0.078
		1R	6.250	5.566	1.136	24	-0.591	1.000	0.060
		2L	9.583	7.790	1.590	24	-0.058	1.000	0.178
		2R	5.833	6.019	1.229	24	-0.626	1.000	0.059
		3L	7.292	5.103	1.042	24	-0.539	1.000	0.069
		3R	6.458	5.610	1.145	24	-0.626	1.000	0.062
		4L	**18.958**	**10.527**	**2.149**	**24**	**0.727**	**0.005**	**168.591**
		4R	**19.792**	**11.371**	**2.321**	**24**	**0.724**	**0.003**	**188.153**
		5L	9.792	6.164	1.258	24	-0.006	1.000	0.190
		5R	9.583	5.882	1.201	24	-0.127	1.000	0.169
	Li-chick	1L	5.095	5.598	1.143	24	-0.704	1.000	0.053
		1R	7.350	6.753	1.378	24	-0.452	1.000	0.082
		2L	8.023	7.186	1.467	24	-0.342	1.000	0.101
		2R	7.164	6.443	1.315	24	-0.365	1.000	0.077
		3L	6.576	5.782	1.180	24	-0.57	1.000	0.065
		3R	7.785	7.907	1.614	24	-0.277	1.000	0.100
		4L	**28.851**	**12.598**	**2.572**	**24**	**0.859**	**< 0.001**	**>10000**

		4R	11.894	7.503	1.532	24	0.337	0.798	0.741
		5L	10.180	7.009	1.431	24	0.026	1.000	0.237
		5R	7.081	6.262	1.278	24	-0.436	1.000	0.074
FP Monocular Left	Di-chick	**1L**	**31.591**	**18.397**	**3.755**	**24**	**0.826**	**< 0.001**	**5702.882**
		1R	5.716	9.515	1.942	24	-0.449	1.000	0.076
		2L	**17.327**	**9.620**	**1.964**	**24**	**0.683**	**0.009**	**64.802**
		2R	2.775	4.944	1.009	24	-0.813	1.000	0.044
		3L	11.907	9.773	1.995	24	0.189	1.000	0.527
		3R	2.348	3.652	0.745	24	-0.859	1.000	0.040
		4L	12.780	8.309	1.696	24	0.376	0.532	1.287
		4R	3.990	6.154	1.256	24	-0.708	1.000	0.050
		5L	6.625	7.226	1.475	24	-0.446	1.000	0.074
		5R	4.940	4.561	0.931	24	-0.803	1.000	0.048
	Li-chick	**1L**	**23.035**	**10.533**	**2.150**	**24**	**0.863**	**< 0.001**	**>10000**
		1R	6.256	6.615	1.350	24	-0.544	1.000	0.066
		2L	11.109	7.272	1.484	24	0.171	1.000	0.419
		2R	3.244	4.112	0.839	24	-0.851	1.000	0.042
		3L	12.500	7.108	1.451	24	0.357	0.455	1.457
		3R	3.643	5.890	1.202	24	-0.728	1.000	0.048
		4L	**22.595**	**8.736**	**1.783**	**24**	**0.876**	**< 0.001**	**>10000**
		4R	6.297	6.747	1.377	24	-0.465	1.000	0.068
		5L	7.578	7.194	1.468	24	-0.379	1.000	0.090
		5R	3.743	4.773	0.974	24	-0.803	1.000	0.045
	Di-chick	1L	3.991	4.710	0.961	24	-0.823	1.000	0.045
		1R	**33.744**	**16.545**	**3.377**	**24**	**0.861**	**< 0.001**	**90653.398**
FP Monocular Right		2L	2.763	4.737	0.967	24	-0.872	1.000	0.043
		2R	15.779	12.802	2.613	24	0.411	0.255	3.214
		3L	4.083	5.947	1.214	24	-0.729	1.000	0.050
		3R	9.537	8.191	1.672	24	-0.0384	1.000	0.176
		4L	3.746	3.668	0.749	24	-0.798	1.000	0.042
		4R	13.417	8.781	1.792	24	0.406	0.398	1.936

	5L	5.033	5.004	1.022	24	-0.716	1.000	0.050
	5R	7.908	6.329	1.292	24	-0.304	1.000	0.091
Li-chick	1L	5.375	7.537	1.539	24	-0.613	1.000	0.063
	1R	**23.770**	**14.388**	**2.937**	**24**	**0.765**	**0.002**	**535.661**
	2L	4.345	5.663	1.156	24	-0.768	1.000	0.050
	2R	13.847	9.447	1.928	24	0.419	0.287	2.233
	3L	4.405	5.204	1.062	24	-0.759	1.000	0.048
	3R	11.261	5.861	1.196	24	0.293	0.984	0.591
	4L	5.219	6.356	1.297	24	-0.594	1.000	0.057
	4R	**21.484**	**10.734**	**2.191**	**24**	**0.807**	**0.001**	**1842.938**
	5L	5.994	4.638	0.947	24	-0.685	1.000	0.053
	5R	4.300	4.755	0.971	24	-0.772	1.000	0.046

Task creation

Logic iteration

Shaping
Fine-tuning
Training

Reviewing
Testing