The retinoic acid receptor regulates development of a key evolutionary novelty - the

molluscan shell.

Keisuke Shimizu[1#], Kazuyoshi Endo[2], and Tetsuhiro Kudoh[3#]

1: JAMSTEC 2-15, Natsushima-cho, Yokosuka-city, Kanagawa, 237-0061, Japan

2: Department of Earth and Planetary Science, The University of Tokyo, 7-3-1 Hongo, Tokyo

113-0033, Japan.

3: Biosciences, University of Exeter, Stocker Road, Exeter EX4 4QD, UK.

Corresponding authors: k.shimizu.bio14@gmail.com, t.kudoh@exeter.ac.uk

Abstract

The shells of molluscs are iconic structures of invertebrate exoskeletons supporting and

protecting their soft body parts. The shell matrix molecules are synthesized and secreted

from a shell-producing tissue, the shell gland. Shell gland cells develop at the early

trochophore stage of the larvae. To date, the molecular signalling pathways by which the

shell gland forms and starts to secrete the shell remains elusive. Here we demonstrate in the

Pacific oyster *Crassostrea gigas* and the limpet *Nipponacmea fuscoviridis*, that the retinoic

acid receptor (RAR), is crucial for inducing shell gland formation. RAR is expressed in both

species in the shell gland at the late gastrula to early trochophore stages prior to the first

production of shell. Suppression of the RAR by chemical inhibitors or gene-knock-down lead

to a complete loss of the larval shell. Transcriptomic and *in situ* hybridisation analyses

revealed that the developmental regulatory genes that are normally expressed in the shell

gland, including *engrailed*, are down-regulated in the RAR-suppressed embryos. Using the

RAR functional assay carried out on zebrafish embryos, we also revealed that the oyster

RAR cannot transduce the RA signal in zebrafish, indicating that the molluscan RAR is

clearly different from vertebrate RARs in its binding capacity to the RA. Our finding

27   represents a key example of adaptive evolution of developmental "toolkit" genes for the

28   origin of a major novel trait, the molluscan shell, in animals.

Introduction

Fossil evidence, combined with phylogenetic relationships, suggests that skeletogenesis

evolved many times in the early Cambrian, accompanying the evolution of novel animal body

plans. At this time, molluscs evolved to have one or more mineralized shells. During the

process of spiral cleavage in their early development, cells fated to be the shell gland are

segregated (Lambert 2010; Perry 2015): by invagination at the gastrula stage, and then the

gland is evaginated to form the shell field (Reviewed in Kniprath 1981). Some transcription

factors are known to be expressed in the shell gland and shell field cells. Most notable is the

homeobox gene *engrailed*, which is expressed in those cells in a specific manner in a variety

of species representing major molluscan lineages (Jacobs et al., 2000; Wanninger &

Haszprunar 2001; Nederbragt et al., 2002; Hinman et al., 2003; Samadi et al., 2009; 2012;

Salamanca-Díaz et al., 2021), suggesting that it may act as a master gene, controlling initial

larval shell development. But direct evidence that *engrailed* is responsible for shell formation

is lacking, and the genes upstream of *engrailed* in this regulatory network are still unknown. It

has also been reported that the Dpp signalling pathway has a crucial role in shell

development by regulating shell growth from the early larval stage (Iijima et al., 2008; Kin et

al., 2009; Shimizu et al., 2011, 2013; Hashimoto et al., 2012,). In *Lymnaea stagnalis*,

asymmetrical expression of Dpp in the left and right side of the shell gland determines

asymmetric shell growth, leading to the formation of either dextral or sinistral coiling of the

shell, whereas equally distributed Dpp in the left and right sides of shell gland in the limpet

Nipponacmea fuscoviridis results in L-R symmetric shell growth (Shimizu et al., 2013).

Recently, it has been shown that the Wnt signalling pathway is involved in the coiling pattern

and morphogenesis of *L. stagnalis* shell (Ohta et al., 2024). Although these studies improved

our understanding of shell growth mechanisms, it is still unclear what signals and

mechanisms trigger initial shell formation, namely the specification of the lineage of the shell

gland and shell field cells.

Transcription factors and nuclear receptors (NRs), bind directly to DNA and regulate the

expression of key genes involved in early development, homeostasis, and metabolism. NRs

consist of two domains, DNA binding domain (DBD) and ligand binding domain (LBD). LBD

in a NR interacts with specific ligands that can activate the NR. Subsequently, the DBD of

NRs binds to a specific transcriptional regulatory element that exists in the promoter region of

the target genes and regulate their transcription (Germain et al., 2006). The retinoic acid

receptor (RAR) is one of the NRs, and its LBD binds to the ligand retinoic acid (RA) in

vertebrates. RAR signalling is one of the major signalling pathways that regulates the

expression of key developmental genes, including hox genes, during early embryonic

development and determines body patterning and tissue differentiation (Marshall et al., 1994,

Kudoh et al. 2002). RA is a metabolite of vitamin A (retinol) and mainly synthesized from

retinal aldehyde by aldehyde dehydrogenase 1a (Aldh1a) (Kudoh et al. 2002). RA is

degraded to oxidized retinoic acid (inactive form) by the RA-degrading enzyme cytochrome

P450 26 (Cyp26) to remove unwanted RA molecules (e.g. Kudoh et al. 2002). Hence, these

two enzymes, Aldh1a and Cyp26, control the spatio-temporal distribution of RA levels during

embryogenesis in chordates (e.g. Niederreither et al., 2002; Kudoh, 2002; Reijntjes et al.,

2005).

Here we report that *rar* is predominantly expressed in the shell gland of trochophore

larvae of the Pacific oyster *Crassostrea gigas* and the limpet *Nipponacmaea fuscoviridis*, and

the loss of function of RAR causes loss of shell gland and shell field, leading to the failure of

larval shell formation. Based on these findings, we discuss the master role of the *rar* gene in

molluscan shell development and its implications on the evolutionary origin of molluscan

shells.

80 **Results**

81 *Expression of the retinoic acid receptor gene in the shell field cells*

To explore the roles of RAR in molluscs, spatial expression patterns of the RAR gene (Cg-*rar*)

in the Pacific oyster *C. gigas* larvae were examined using wholemount *in situ* hybridisation.

Cg-*rar* is expressed in the dorsal invaginated region, namely the shell field invagination (SFI)

(Eyster and Morse, 1984) at late gastrula stage (8 hours post-fertilization [hpf]) (Fig. 1a). The

expression of Cg-*rar* in SFI is confirmed by comparison with the expression of two shell field

marker genes, Cg-*gata2/3* and *larval shell matrix protein 1* (Cg-*lsmp1*) in SFI. At late gastrula

stage (8 pfh), Cg-*gata2/3* is expressed around the SFI (Fig. 1b), but the larval shell matrix

protein Cg-*lsmp1* is not expressed yet (Fig. 1c). The expression of Cg-*lsmp1* was first detected

in the shell field cells that originated from SFI at early trochophore stage (10 hpf) and

overlapped with the Cg-*rar* expression (Fig. 1d, e). Cg*gata2/3* continued its expression around

SFI in 10 hpf larvae (Fig. 1f). After the shell field evagination (12 hpf), Cg-*rar* and Cg-*lsmp1*

are expressed in the shell field cells (Fig. 1g, h), and Cg-*gata2/3* expression is restricted to the

edge of the shell field cells (Fig. 1i). We also examined expression of *rar* (Nf-*rar*) and two shell

marker genes, *chitin synthase 1* (Nf-*CS1*) and *6D12,* that were identified as biomineralization

genes in the abalone *Haliotis asinina* (Jackson et al., 2006) in the trochophore larvae of the

limpet *N. fuscoviridis.* Nf-*rar* is expressed in the shell field cells (Fig. 1j), which are marked by

the shell marker gene *6D12* (Fig. 1k). The other shell marker Nf-*CS1* is expressed at the edge

of the shell field cells (Fig. l). These observations suggest that *rar* has a role in molluscan larval

shell gland development.

Figure 1. RAR is expressed in the shell gland and progenitor cells from gastrula to trochophore stages.

In situ hybridisation of the oyster *C. gigas* (a to i) and the limpet *N. fuscoviridis* (j to l). The embryos and larvae were stained with *in situ* probes for *Cg-rar* (a-c), *Cg-lsmp1* (d-f), *Cg-gata2/3* (g-i), *Nf-rar* (j), *Nf-6D12* (k) and *Nf-CS1* (l).

Retinoic acid receptor is essential for shell fate specification in molluscs

To investigate the function of RAR in molluscs, the embryos of the pacific oyster *C. gigas* and

the limpet *N. fuscoviridis* were treated with atRA and two RAR inhibitors (Ro-41-5253 and

AGN-193109) at the 1-2 cell stage onward. Although the 24 hpf larvae precipitated the

embryonic shell in normal development, the larvae that were treated with the RAR inhibitors

completely failed to form their shells both in *C. gigas* (Fig. 2c,d,k) and *N. fuscoviridis* (Fig.

2h,i,l). The 24 hpf larvae treated with at-RA showed abnormal morphology of larval shells with

reduction of areas covered by the shell (Fig. 2b,g). We then confirmed the crucial role of RAR

in the shell development using another loss-of-function approach: Nf*rar* was knocked down by

microinjection of a specific morpholino oligo (Nf*rar*-MO) in *N. fuscoviridis*. The Nf*rar*-MO

injected embryos indeed developed non-shelled veliger larvae (Fig. 2n), which are similar to

those treated with RAR inhibitors (Fig. 2X). These results suggest that RAR has a crucial role

in promoting shell development in molluscan larvae.

Figure 2 Suppression of RAR blocks shell formation in the oyster and limpet larvae.
Live *C. gigas* (a-e) and *N. fuscoviridis* (f-j) embryos were treated with atRA (b,g), the retinoic
acid inhibitors RO (c,h), AGN (d,i) and aldh1a inhibitor DEAB (e,j). For *C. gigas*, the live
larval shell was visualized by calcein with green fluorescence (a-e) indicating absence of the
shell in the Ro and AGN treated embryos (c,d). In the limpet, presence (black arrow head)
and absence (white arrow head) of the larval shell was visible with bright field microscope (f-
j). k,l. Ratios of normal shelled larvae, small shelled larvae, and abnormal larvae in three
biological replicate experiments using the oyster (**k**) and the limpet (**l**). Scale bar: 20μm (**a**)
and 50 μm (**b**). (m.n.) Gene knockdown of *Nf_rar* by morpholino injection. Consistent with the
results of Ro and AGN, Nf*rar*-MO suppressed larval shell formation in the limpet (n, white
arrow head).

RNAseq reveals that RAR regulates the shell field genes

In order to identify the target genes for the retinoic acid receptor (RAR) in molluscs, we treated

the oyster, *C. gigas* embryos/larvae with RAR inhibitor (Ro-41-5253) and atRA at the 1-2 cell

stage onward and analzyed the transcriptome in early trochophores (9.5 hpf). The results show

that the expression levels of 1289 and 426 genes (14.1 and 4.7 %) were significantly changed

by the Ro-41-5253 and atRA treatment, respectively (|logFC| > 1 and FDR < 0.05, Fig. 3a). If

Ro-41-5253 blocks the atRA signal, a common set of genes would be expected to show an

inverted response to these treatments. However, we found only one gene that was down- and

up-regulated by Ro-41-5253 and atRA respectively. This suggests that atRA may not act as

the ligand of RAR to activate the genes downstream of RAR. In contrast, 246 and 47 genes

were down- and up-regulated in both Ro-41-5253 and atRA treated larvae, respectively (Fig.

3a), suggesting there may be some level of similarity between atRA and Ro-41-5253 treated

embryos. Indeed, a LogFC plot of these two conditions shows a linear correlation pattern

through the down-regulated to up-regulated genes (Fig. 3b). It should also be noted that the

angle of the plot leans toward the Ro-41-5253 side, suggesting the level of gene expression

change is more prominent in the Ro-41-5253 compared with atRA treated samples (Fig. 3b).

Among the 846 genes that were down-regulated in Ro-41-5253 treated larvae (Fig. 4a), eight

key domains or families are significantly enriched in InterProScan (dynein heavy chain domain,

caveolin, EF-hand domain, homeobox, G protein-coupled receptor, FAD/NAD (P) binding

domain, and immunoglobulin) (Fig. 4c) ($p < 0.05$).

We then focused on the larval shell matrix protein (LSMPs) encoding genes in *C. gigas* (Zhao

et al., 2018). We found that a total of 13 and 9 of the 29 Cg-LSMPs encoding genes were

significantly down regulated in Ro-41-5253 and atRA treatment, respectively (Fig. 3b). These

genes had relatively high expression levels in the control samples (FPKM > 10) except for one

gene, CGI_10008969, in Ro-41-5253 (Fig. 3b). The data also confirmed that the effect of Ro-

41-5253 is stronger than that of atRA. The homeobox genes *engrailed*, *post2, evx, cux*, and

ssuh2, were down-regulated in both conditions (Fig. 3d), and other homeobox genes, *hox1,*

lox1, hhex, *nkx6.2*, *pou4*, *six1/2* and *rx,* were down-regulated only in Ro-41-5253 treatment,

suggesting these genes might be involved in shell gland and shell field cell differentiation and

function (Fig. 3d).

Figure 3. Transcriptomics analyses of *C. gigas* 24hpf larvae treated with RA or Ro.

a. Number of transcripts up or down regulated by RA and/or Ro (|logFC| > 1 and FDR < 0.05.

b. LogFC plot of expression levels of RA and Ro samples at Y and X axes, respectively. Red

highlights up- or down-regulated genes by RA. InterProScan enriched gene families in the

transcripts down-regulated by Ro identified 8 gene familes/domains. d. LogFC plot of

expression levels of RA and Ro samples focusing on homeobox genes.

RAR determines shell gland cell fate

To confirm that shell-related gene expression is regulated by the RAR signalling pathway,

Ro-41-5253 treated larvae were *in situ* stained with probes for shell-marker genes (Pacific

oyster: Cg*engrailed,* Cg-*lsmp1* and Cg-*tyr1*; limpet: Nf-*engrailed,* Nf-*CS1* and Nf-*6D12*) in *C.*

gigas (8 hpf trochophore larvae) and *N. fuscoviridis* (10 hpf trochophore larvae) (Fig. 3a, b, e,

f). Ro-41-5253-treated larvae showed suppressed expression of these shell-marker genes in

both species, strongly suggesting that RAR signalling has a crucial role in cell fate

specification of the shell gland. Besides these shell field-marker genes, the transcription

factor *engrailed* was also suppressed in the shell gland region of both species after Ro-41-

5253 treatment.

181
182

Figure 4 Gene expression patterns in shell gland and shell field are dependent on RAR signalling.

185   *In situ* hybridization of oyster (18hpf) (a-c) and limpet (f-h) (10h) larvae. Expression of *Cg-*

186   *lsmp-1* (a), *Cg-tyr1* (b), *Cg-en2* (c), *Nf-cs1* (f), *Nf-6D12* (g) and *Nf-en* (h) are all suppressed

187   by the RAR inhibitor (Ro-41-5253). Embryos treated with all trans Retionic Acid (atRA) did

188   not show strong up or down regulation of these marker genes. Scale bar: 20μm.

CgRAR disturbs the RA signaling in zebrafish embryo

The DNA binding domain (DBD) of oyster RAR (CgRAR_DBD) has a high similarity (81-86%) to that of chordate RAR. In contrast, the ligand binding domain (LBD) of oyster RAR (CgRAR_LBD) shows a low similarity (53-60%) to that of chordate RAR. It has been known that the 25 amino acid sequence around the LBD region, called the ligand binding pocket (LBP), is structurally important for interaction with RA in the human RAR. Five of these 25 amino acids in the LBP of CgRAR are different from human and zebrafish RAR suggesting that the ligand-binding ability and/or specificity might be different between CgRAR and vertebrate RARs. To test if Cg-RAR could respond to the RA *in vivo*, we used zebrafish (*Danio rerio*) embryos as an *in vivo* assay system. Firstly, zebrafish RARab (Dr*RARab*) was overexpressed in the zebrafish embryo by injecting mRNA at 1-cell stage. When uninjected zebrafish embryos were treated with 0.1 μM atRA, they showed mild reduction of the head and tail (Fig. 5Bb). However, when the Dr*RARab* was overexpressed, the head and tail became further reduced (Fig. 5Be), suggesting that overexpression of Dr*RARab* enhanced the action of atRA and induced the posteriorisation of the brain (Kudoh et al. 2002). However, when Cg*RAR* is overexpressed and the embryos subsequently treated with 0.1 μM atRA, the phenotype was similar to that of uninjected embryos treated with 0.1 μM atRA (Fig. 5Bk), suggesting that CgRAR did not

properly transduce the signal of atRA in the zebrafish embryo. To examine if the lack of

response is due to the different LBD of the CgRAR, we generated a chimeric construct of

DrRARab in which the LBD was replaced with that of CgRAR (DrRAR-CgLBD) (Fig. 5A).

Indeed, the embryos injected with mRNA of DrRAR-CgLBD did not show any enhancement of

the effect of 0.1 μM atRA (Fig.5Bh). The same result was observed with 9-cis-RA, indicating

that CgRAR cannot respond to either at-RA or 9-cis-RA as the DrRAR does (Fig. 5Bc,f,i). To

further examine brain-regional specific action of the three RAR constructs (DrRAR, CgRAR,

and DrRAR-CgLBD) to atRA, we conducted expression analysis using three positional marker

genes, Dr*pax2* (mid-hindbrain boundary [MHB]), Dr*hoxb3a* (tail to hindbrain rhombomere 5)

and Dr*hoxb5b* (tail to hindbrain rhombomere 7). Uninjected embryos treated with 0.1 μM atRA

showed loss of Dr*pax2* and expansion of Dr*hoxb3a* and Dr*hoxb5b* to the anterior end (Fig.

5Cb,f,j), suggesting the loss of the forebrain, midbrain and MHB. However injection of the Cg-

rar mRNA rescued the Dr-*pax2* expression in the MHB and moved the anterior border of the

Dr-*boxb3a* and Dr-*hoxb5b* expression domain backward in the presence of 0.1 μM at-RA

(Fig.5Cd.h.j), suggesting Cg-RAR acts as a dominant negative form against the Dr-RAR in the

zebrafish embryo. Consistently, the DrRAR-CgLBD showed a similar rescue phenotype

(Fig.5Cc,g,k), indicating the dominant negative activity is due to the diverse functions of the

LBDs in the oyster and zebrafish RARs. We then examined Dr-*hoxb1b* expression, one of the

direct target genes of RA signaling, using zebrafish embryos at late gastrula. DrRAR-CgLBD

or Cg*RAR* mRNA injection led to suppress Dr-*hoxb1b* expression (Fig. 5Db,c).

Finally, to identify the key amino acid that alters LBD activity between vertebrates and

molluscs, the amino acid Valine (V) at position 223, a residue that is only specific to molluscs,

was changed to Phenylalanin (F), which is conserved in vertebrates (DrRAR-CgLBD V223F),

its activity was examined. Although DrRAR-CgLBD showed a dominant negative activity, this

single amino acid modification DrRAR-CgLBD V223F was sufficient for the gene to lose the

dominant negative activity, and act instead similar to the zebrafish RAR, enhancing the RA

treatment in the 24hpf embryos by reducing head and tail (Fig. 5Bm) and by inducing *hoxb1b*

expression at late gastrula stage (Fig. 5Df). Altogether, we conclude that the molluscan RAR

has a property distinct from the vertebrate RAR in response to the RA, and this difference is

largely due to the single amino acid substitution at the position 223 with valine in molluscs and

phenylalanine in vertebrates respectively.

Figure 5. Comparative and functional domain analysis of *Cg-rar* and *Dr-rar* using the

zebrafish embryo *in vivo* model. A. zebrafish RAR (DrRAR) and Pacific oyster RAR

(CgRAR) show conserved domain structure with N-terminal DNA binding domain (DBD) and

C-terminal ligand binding domain (LBD). Chimeric construct of DrRAR_CgLDB (DrRAR

backbone replaced with the LBD domain of CgLBD), and DrRAR_CgLDB(V223F), a point

mutation of DrRAR_CgLDB that altered the amino acid V(valine) 223 to F(phenylalanine), B.

Fertilised zebrafish eggs were injected with mRNA encoding RAR constructs listed in A and

treated with or without atRA or 9cRA. Live images were taken at 24hpf. C. *In situ* hybridisation

of 24hpf embryos injected with RAR constructs, treated with atRA, and stained with *Dr_hoxb3a*

(a-d), *Dr_hoxb5b* (e-h) or *Dr_pax2a* (i-l) probes. D. *In situ* hybridisation staining of *Dr-hoxb1b*

in the injected and RA treated embryos. Embryos in b and c were injected with mRNAs at 2-

cell stage in one blastomere, showing effects in one side of the embryo only.

Discussion

We report here that the retinoic acid receptor (RAR) is essential trigger for shell gland

development and initial shell secretion in molluscs. The expression patterns of the homeobox-

containing regulatory gene *engrailed* are likely associated with the development of molluscan

shells, as *engrailed* is expressed in the ectoderm cells at the boundary of the shell forming

region of various taxa (Polyplacophora, Jacobs et al., 2000; Cephalopoda, Bratte et al., 2007;

Scaphopoda, Wanninger and Haszuprunar, 2001; Bivalvia, Kin et al., 2009, Gastropoda,

Moshel et al., 1998). Other transcription factors, *distal-less* (*dlx*)*, soxC, gata2/3, pax2/5/8, and*

lox 4 are also expressed inside or the boundary of the shell field cells in bivalve or gastropod

(Jackson and Degnan 2016; Liu et al., 2015, 2017, 2020). However, there has been no

functional evidence showing that these transcription factors are involved in the initial shell

development. In this study, we found that *engrailed* expression in the shell field was

significantly down-regulated in the larvae of both oyster and limpet that were treated by Ro-

41-5253 (Fig. 3c, h), and these larvae failed to develop their embryonic shells with down-

regulation of the expression of LSMP genes (Figs. 2-4). On the other hand, the expression of

other shell-related candidate genes (*dpp*, *dlx*, *gata2/3*, and *soxC*) was not affected under the

Ro-41-5253 treatment in oyster larvae (9.5 phf). These results suggest that RAR signaling

directly or indirectly regulates *engrailed* expression, and both *rar* and *engrailed* are involved in

the differentiation of shell secretary cells and crucial signals for their embryonic shell

development. In contrast, other shell related genes (*dpp*, *dlx*, *gata2/3*, and *soxC*) possibly

belong to other gene regulatory networks (GRN) and play roles in the shell growth or other

shell-related functions rather than the differentiation of shell secretory cells.

The signaling pathway encompassing the retinoic acid receptor has been well studied in

chordates. Its ligand, RA, is an important morphogen involved in body axis formation and

growth patterning corresponding to its gradient, as RA can diffuse over long distances, and its

spatio-temporal distribution regulates the expression of homeotic genes (Marshall et al. 1994;

Aulehla & Purquite, 2010). In contrast, RAR is not found in insect species such as *Drosophila,*

thus it has been suspected that the morphogenetic role of the RA/RAR signaling may have

been acquired along the branch leading to chordates, and therefore is linked to the origin of

their innovative body plans (Shimeld 1996; Manzanares et al. 2000; Schilling & Knight 2001;

Wada 2001; Holland 2005). However, a recent study using the genomic databases including

Ambulacraria and Protostome revealed that the components of RA signaling machinery, RA

synthetase (*aldh1a*), RA-degrading enzyme (*cyp26*) and receptors (*rar*), originated in the last

common ancestor of bilaterians, rather than in the last common ancestor of chordates (Albalat

& Canestro 2009). This suggests that RA signaling pathway may well have been involved in

the evolution and developmental diversity of not only Chordata but also Lophotrochozoa. The

effects of RA overdose have been studied in a variety of lophotrochozoans including

platyhelminthes, annelids, and molluscs (e.g. platyhelminthes: Romero & Bueno, 2001,

annelids, Handberg-Thorsager et al., 2018, molluscs: Creton et al., 1993; Dmetrichuk et al.,

2006; Carter et al., 2015; Vogeler et al., 2017; Johnson et al., 2019). These previous reports

suggest that an ancestral function of RAR signal was neuronal growth and cell survival

(Dmetrichuk et al., 2006; Campo-Paysaa et al., 2008; Handberg-Thorsager et al., 2018). In

chordates, an important role of RAR signaling is embryonic patterning along the anterior-

posterior axis via regulation of *hox* genes (Shimeld, 1996; Schilling and Knight, 2001, Kudoh

et al. 2002). For instance, treatment of the zebrafish embryo with atRA induces loss of the

anterior brain cell fates (Fig. 5, Kudoh et al. 2002). However, a loss of these anterior structures

has not been observed in invertebrate larvae treated with atRA. Our transcriptome results also

showed that atRA does not induce expression of *hox* genes (Fig. 3), albeit some hox and other

homeobox genes were down-regulated (*hox1*, *lox5*, *lox4*, and *post2*) by Ro-41-5253 (Fig. 3).

Interestingly, many homeobox genes that are downregulated by Ro-41-5253 showed an

overall linear relationship to the downregulated pattern seen in the treatment with atRA,

suggesting that atRA in this experiment may have acted as a weak inhibitor of RAR. If this is

the case, it might be possible that molluscan RAR may have a ligand other than atRA (or

9cisRA). Or otherwise, the role of RA in molluscs could primarily be to restrain the activity of

RAR. Within the 846 genes that were down regulated in the Ro-41-5253 treated larvae (Fig.

4a), eight key domains or families are significantly enriched in InterProScan (dynein heavy

chain domain, caveolin, EF-hand domain, homeobox, G protein-coupled receptor, FAD/NAD

(P) binding domain, and immunoglobulin) (Fig. 4c) ($p < 0.05$). One of the enriched domains,

EF-hand domain, is well known in calcium-binding proteins, which may be involved in the

secretion of calcium carbonates in the larval shell. Homeobox containing genes play key roles

in the gene regulation for fate specification and body patterning. These data suggest that RAR

is involved in the process of cell fate specifications including regulation of gene expression for

key transcription factors (such as *engrailed* and tissue specific genes including calcium

secretion proteins and LSMPs.In chordates, it is known that the RAR/RXR heterodimer binds

to the RA response elements (RAREs) and maintains the expression of target genes. RAREs

consist of two direct repeats of a core hexameric motif 5'-(A/G)G(G/T)(G/T)(G/C)A-3',

separated by 1, 2, or 5 bp long nucleotides (DR1, DR2, or DR5, respectively) (Balmer and

Blomhoff, 2002). In the mollusc *Nucella lapillus* and the annelid *Platynereis dumerilli*, the

RAR/RXR heterodimer is able to bind to RAREs (DR1, DR2, and DR5) like chordate

counterparts (Gutierrez-Mazariegos et al., 2014; Handberg-Thorsager et al., 2018). Indeed,

sequences of DBD in RAR including the three box regions (P-box, D-box, and T-box) are highly

conserved between vertebrates and lophotrochozoans. In the zebrafish embryo assay, CgRAR

acted like a dominant negative form and antagonized the signaling pathway with DrRAR. This

suggests that CgRAR cannot bind to atRA in the manner similar to chordate RAR but can still

bind to the RARE DNA sequence via DBD, therefore, can act as a dominant negative form in

zebrafish.

Although the DBD of RAR is well conserved among lophotrochozoans and chordates, LBD

sequences of RAR in lophotrochozoans are more divergent from those in chordates. In the

annelid *P. dumerilli,* a single amino acid difference at position 356 is observed among the

important residues of the LBP, at which position human and other vertebrates have

phenylalanine while the annelid has valine, and this mutation (V356F) is shown to decrease

the binding ability to atRA (Handberg-Thorsager et al., 2018). The valine at position 356 in

RAR of *P. dumerilli* is shared by the annelid *Capitella teleta*, pacific oyster *C. gigas*, and the

brachiopod *Lingula anatina*. In the zebrafish embryo assay, the phenotype and the gene

expression pattern under the overexpression of CgRAR and DrRAR-CgLBD with atRA

treatment rescue, indicated a behaviour as an antagonist to DrRAR (Fig. 5). On the other hand,

the zebrafish embryos with overexpression of DrRAR-CgLBD-V223F, which corresponds to

V356F mutant of *P. dumerili* RAR (Handberg-Thorsager et al., 2018), mimicking vertebrate

RAR, indicated results just like those of the overexpression of DrRAR (Fig. 5). Indeed, DrRAR-

CgLBD-V223F was able to induce *hoxb1b* expression, a direct target gene of RAR in zebrafish,

while DrRAR-CgLBD inhibited its expression (Fig. 5). These results suggest that the RAR of

molluscs and other lophotrochozoans such as annelids and brachiopods cannot activate

downstream gene expression using atRA as the ligand, and that the single amino acid

modification in the LBP critically affects the ligand binding and/or transcriptional regulation.

The marginal or differential role of atRA in molluscs is also supported by the experiment with

DEAB, an inhibitor of atRA synthesis, showing no clear inhibitory effect in the shell

development (Fig. 2).

Invertebrate exoskeletons constitute one of the key products of the Cambrian explosion,

and may have diversified by, and contributed to, the evolutionary arms race, because external

skeletons play important roles for survival (e.g. feeding, protection, and body support). Major

bilaterian lineages with exoskeletons have probably evolved their hard tissues independently

in each phylum within a very short time across the Ediacaran-Cambrian transition (Murdock

and Donoghue, 2011). However, the evolutionary origins of exoskeletons remain unclear.

Various homologous genes or proteins that are associated with skeletogenesis have been

reported from different groups of bilaterians (Ettensohn et al. 2003; Livingston et al., 2006;

Jackson et al., 2007), leading to a notion that a "biomineralization toolkit" might have evolved

in parallel among those taxa. The homeobox-containing transcription factor *engrailed* has been

thought to be one of the key biomineralization toolkit genes. Jacobs & Gates (2003) compared

expression patterns of *engrailed* among invertebrate phyla and pointed out the relationship

between the *engrailed* expression pattern in ectodermal cells and skeletal development among

protostomes; absence of *engrailed* expression in ectodermal cells corresponds with absence

of exoskeleton in onychophorans (Wedeen et al., 1997) and annelids (Wedeen & Weisblat,

1991; Lans et al., 1993; Seaver & Kanesige, 2006). In juveniles of brittle stars, *engrailed* is

expressed in the boundaries between newly forming spine ossicles (Lowe & Wray 1997). The

cells that form spine ossicles are called skeletogenic mesenchyme cells, which are derived

from mesoderm, but they rely in part on ectodermally derived cues (Malinda & Ettensohn,

1994). Thus, the patterns of *engrailed* expression in skeleton boundaries in echinoderms are

consistent with those in protostomes. Similar *engrailed* expression patterns have been

reported in Lophotorochozoa. In molluscs and brachiopods, *engrailed* is expressed around the

shell gland and the mantle lobes that are derived from ectodermal cells, respectively (Moshel

et al., 1998; Jacobs et al., 2000; Wanninger and Haszuprunar, 2001; Bratte et al., 2007; Kin et

al., 2009; Shimizu et al., 2017). We found that RAR signal is the key for inducing shell gland

development and thus subsequent shell formation, and regulates the expression of the

homeobox gene *engrailed*. Thus, previously existing gene modules including *engrailed* that

have been deployed in dorsal ectodermal cells might have contributed to molluscan shell

evolution, and this novel body plan is most likely to have been recruited by the RAR pathway.

Such an evolutionary process, in which a new gene function is added to an existing toolkit,

known as co-option, is not limited to the origin of the molluscan shell (Carrol et al., 1994; beetle

horn, Moczek & Rose, 2005). Qur new findings provide clues about the origin and evolution of

exoskeletons based on the shell gland cell development and involvement of divergent nuclear

receptor signalling pathway, and pose more general questions about how various novel body

plans evolved in animals.

Material and Methods

Animals

Individuals of *Nipponacmaea fuscoviridis* were collected from the intertidal rocky shores in

Manazuru, Kanagawa, Japan. Individuals of the oyster *Crassostrea gigas* were obtained

from Guernsey Sea Farms Ltd. (Guernsey, UK). Methods of egg collection followed previous

studies (oyster: Vogeler et al., 2017, limpet: Deguchi 2007). Embryos of the limpet and oyster

were cultured in filtered seawater at 24 ºC and 22 ºC, respectively.

Chemical treatment

Embryos of *C. gigas* and *N. fuscoviridis* were incubated in filtered natural seawater

containing 0.1% (v/v) DMSO and treated with 2μM of all-trans retinoic acid (R2625, Sigma-

Aldrich Japan, Tokyo, Japan) and 0.5-2 μM of RAR inhibitors Ro-41-5253 (SML0573, Sigma-

Aldrich Japan, Tokyo, Japan) and AGN-193109 (SML2034, Sigma-Aldrich) at the 1 or 2-cell

stages. As a negative control, embryos were also exposed to filtered natural seawater with

0.1% (v/v) DMSO. For the examination of the presence or absence of calcium carbonates in

the larval shells of oyster, we added 1/400 calcein stock solution (Sigma, 6g/L in DMSO) into

culture dishes from 4 to 24 hpf embryo.

Microinjection of MO and mRNA

The MO (Gene Tools LLC) NfuRAR-MO (GTCATACATTGTATTTGGCACATCA) was

designed against their start codons. NfuRAR-MO was adjusted to 2.0 mg mL^{-1} in 3 %

Rhodamine B dextran isothiocyanate-Dextran (R8881, Sigma). Full length of CgiRAR

(CGI_10028545) was amplified by PCR using specific primers (CgiLBD-F

TGTGGGATAAAGTGACAGAAC, CgiLBD-R CTAACCGTAAATGAAGGACTC,

ZfRAR_CgiLBD-F TTCATTTACGGTTAGAGATCCCAGGCTCCATG and ZfRAR-CgiLBD-R

CTGTCACTTTATCCCACA-GGTCGACATCCAG) and cloned into pCS2+ vector. The mRNA

was synthesized using the mMESSAGE SP6 kit (AM1340, Thermo Fisher Scientific) and

adjusted to 25 µg mL^{-1} in 5 % Phenol red (Sigma). NfuRAR-MO was injected into unfertilized

eggs of *N. fuscoviridis* using FemtoJet microinjector (Eppendorf). The cocktail of MOs

(DreRARaa and ab) and mRNA (CgiRAR) were injected into one-cell zebrafish embryos.

RNA extraction and transcriptome analysis

We collected 9 hpf trochophore larvae of oyster that are three conditions in triplicate (0.1%

DMSO, 1 µM of all-trans retinoic acid, and 1 µM of RAR inhibitor Ro-41-5253) and 10 hpf

trochophore larvae of limpet that are no treatment in triplicate. We isolated mRNA according

to the manufacturer's protocols for RNA extraction using TRIzol reagent (15596026, Thermo

Fisher Scientific) and RNeasy (Quiagen, 74104), and stored at -80 ºC until for complementary

DNA (cDNA) synthesis and/or transcriptome analysis. We prepared 100-bp DNA libraries from

the mRNA samples using ScriptSeq RNA-seq library preparation kit (ScriptSeq: V224-40412)

according to the manufacturer's protocols and then performed 100-base paired-end

sequencing by HiSeq2500 system (Illumina). We then performed de novo transcriptome

assembly using Trinity (Grabherr et al., 2011; Haas et al., 2013) release (r20131110) and

mapping these reads using Tophat (version 2.0.8b). We picked up transcripts of which FPKM

values are higher than 5 for our analyses. Differential expression analysis was evaluated using

edgeR in the bioconductor package of R (3.7.14). To examine the changes of expression levels

of larval shell matrix protein (LSMP) genes by Ro and RA, we selected candidate genes from

our *C. gigas* larval transcriptome database (Zhao et al., 2018). Among 110 annotated Cgi-

LSMPs genes in D-shape larvae of *C. gigas*, we selected 29 Cgi-LSMPs for analyses; we

removed 58 housekeeping-like genes that are constantly expressed in all developmental

stages (38 stages) and adult tissues (11 tissues) (RPKM > 1, Zhang et al., 2012), 13 transcripts

that are not predicted as gene models in the genome study (Zhang et al., 2012), and 11

transcripts that are quite low expression levels in our transcriptome result (FPKM < 1).

cDNA synthesis and gene subcloning

cDNA was synthesized according to the manufacturer's protocols for cDNA synthesis using

ReverTra Ace (TRT-101 ,Toyobo, Osaka, Japan). Nine gene sequences (*Cgi-rar*, *Cgi-cyp26,*

Cgi-engrailed, *Cgi-gata2/3*, *Cgi-tyr1*, *Cgi-lsmp1, Nfu-rar*, *Nfu-cyp26*, *Nfu-6D12*) were

amplified with PCR using specific primers and then purified using the SV Gel Extraction and

PCR Clean-Up system (A9281, Promega). Subcloning of these genes was performed using

pGEM-T easy Vector Systems (A1360, Promega) and *Escherichia coli* JM109 competent

cells (L1001, Promega).

Whole mount **in situ** ***hybridization***

Molluscan embryos were fixed with 4% paraformaldehyde (PFA) buffer (with 0.1M MOPS,

0.5M NaCl, and 2mM EGTA) for 1 h at room temperature. After fixation, samples were

washed with phosphate-buffered saline with 0.1 % Tewwn-20 (PBT) for 5 minutes five times,

followed by the dehydration with a gradual series of methanol/PBT (50/50, 80/20, 100/0) for

15 minutes each, and stored in 100 % methanol at -20ºC. Zebrafish embryos were fixed with

4% PFA in PBS at 4 ºC overnight, washed three times with PBT, and stored in 100 %

methanol at -20ºC. Probe synthesis and whole mount *in situ* hybridization was performed

following the protocol from a previous study on the brachiopod *Lingula anatina* (Shimizu et

al., 2017) and zebrafish *D. renio* (Kudoh et al. 2002) for molluscan embryos and zebrafish

embryos respectively. Color development was performed with NBT/BCIP (Sigma-Aldrich,

11681451001) or BM Purple (Sigma-Aldrich, 11442074001) in the dark. All color reactions

were stopped at the same time.

Ethical declaration

All methods are reported in accordance with ARRIVE guidelines. All experiments and

methods approved by the Animal Welfare and Ethical Review Board at the University of

Exeter and University of Tokyo.

Data Availability

The transcriptome data of *C. gigas* larvae treated with Ro41-5253 and atRA is desposited

and publicly available from NCBI GEO accession GSE291194.

Acknowledgements

This work was supported by the JSPS Overseas Long Term Fellowship to KS, the NC3Rs

project grant (NC/X001121/1) to TK, and the JSPS KAKENHI grants (23244101 and

18H01323) to KE. We thank the Aquatic Resource Facility staff of the University of Exeter for

maintenance and husbandry of animals. We also thank to Liz Williams and David

Salamanca-Diaz for constructive comments on our manuscript and data, and to Paul O'Neill

for bioinformatics support.

References

Albalat, R. The retinoic acid machinery in invertebrates: ancestral elements and vertebrate
innovations. *Mol. Cell. Endocrinol.* **313**, 23-35 (2009).

Albalat, R. & Cañestro, C. Identification of Aldh1a and Cyp26 and RAR orthologs in
protostomes pushes back the retinoic acid genetic machinery in evolutionary time to the
bilaterian ancestor. *Chem. Biol. Interact.* **178**, 188–196 (2009).

Anderson, D. T. Embryology and phylogeny in annelids and arthropods. Pergamon Press,
Oxford (1973).

Aulehla, A. & Pourquié, O. Signaling gradients during paraxial mesoderm development. *Harb.*
Perspect. Biol. **2**, a000869 (2010).

Baratte, S., Andouche, A. & Bonnaud, L. Engrailed in cephalopods: a key gene related to the
emergence of morphological novelties. *Dev. Genes Evol.* **217**, 353-62 (2007).

Campo-Paysaa, F., Marlétaz F, Laudet, V. & Schubert, M. Retinoic acid signaling in
development: tissue-specific functions and evolutionary origins. *Genesis* **46**, 640–656
(2008).

Cather, J. N. Cellular interactions in the development of the shell gland of the gastropod,
Ilyanassa. *J. Exp. Zool.* **166**, 205-23 (1967).

Conway-Morris, S. The cuticular structure of the 495-Myr-old type species of the fossil worm
Palaeoscolex, *P. piscatorum*. *Zoological Journal of the Linnean Society* **119**, 69–82 (1997).

Créton, R., Zwaan, G. & Dohmen, R. Specific developmental defects in molluscs after
treatment with retinoic acid during gastrulation. *Dev. Growth Differ.*, 357–364 (1993).

Deguchi, R. Fertilization causes a single Ca2 þ increase that fully depends on Ca2 þ influx in
oocytes of limpets (Phylum Mollusca, Class Gastropoda). *Dev. Biol.* **304**, 652–663. (2007).

Dmetrichuk, J.M., Carlone, R.L. & Spencer, G. E. Retinoic acid induces neurite outgrowth and
growth cone turning in invertebrate neurons. *Dev. Biol.* **294**, 39–49 (2006).

Dzik, J. Early metazoan evolution and the meaning of its fossil record. In M. K. Hecht, R. J.

MacIntyre, and M. T. Clegg (eds.), Evolutionary biology, pp. 339–386. New York (1993)

Germain P, Staels B, Dacquet C, Spedding M, Laudet V: Overview of nomenclature of

nuclear receptors. Pharmacol Rev 2006, 58:685–704.

Grabherr, M. G. *et al.* Full-length transcriptome assembly from RNA-Seq data without a

reference genome. *Nat. Biotechnol.* **29**, 644–652 (2011).

Hashimoto, N., Kurita, Y. & Wada, H. Developmental role of dpp in the gastropod shell plate

and co-option of the dpp signaling pathway in the evolution of the operculum. *Dev. Bio.*

366, 367–373 (2012).

Haas, B. J. *et al.* De novo transcript sequence reconstruction from RNA-seq using the Trinity

platform for reference generation and analysis. *Nat. Protoc.* **8**, 1494–1512 (2013).

Deployment of regulatory genes during gastrulation and germ layer specification in a model

spiralian mollusc Crepidula.

Perry KJ, Lyons DC, Truchado-Garcia M, Fischer AH, Helfrich LW, Johansson KB, Diamond

JC, Grande C, Henry JQ. *Dev Dyn*. **244**:1215-48. (2015)

Hinman VF, O'Brien EK, Richards GS, Degnan BM. Expression of anterior Hox genes during

larval development of the gastropod Haliotis asinina. *Evol Dev* **5** 508-21 (2003).

Holder, N., & Hill, I. Retinoic acid modifies development of the midbrain-hindbrain border and

affects cranial ganglion formation in zebrafish embryos. *Development* **113**, 1159-1170

(1991).

Holland, L. Z. Non-neural ectoderm is really neural: evolution of developmental patterning

mechanisms in the non-neural ectoderm of chordates and the problem of sensory cell

homologies. *J. Exp. Zool. B Mol. Dev. Evol.* **304**, 304–323 (2005).

Holland, L. Z., Kene, M. & Holland, N. D. Sequence and embryonic expression of the

amphioxus engrailed gene (AmphiEn): The metameric pattern of transcription resembles

that of its segment-polarity homolog in Drosophila. *Development* **124**, 1723–1732. (1997).

Iijima M, Takeuchi T, Sarashina I, Endo K. Expression patterns of engrailed and dpp in the

gastropod Lymnaea stagnalis. *Dev Genes Evol*. **218**, 237-51 (2008).

Jacobs, D. K. & Gates R. D. Developmental Genes and the Reconstruction of Metazoan

Evolution – Implications of Evolutionary Loss, Limits on Inference of Ancestry and Type 2

Errors. Integrative & Comparative Biology v. 43 pp. 619-646 (2003).

Jacobs, D. K., Wray, C. G., Wedeen, C. J., Kostiken, R., Desalle, R., Staton, J. L., Gates, R.

D. & Lindberg, D. R. Molluscan engrailed expression, serial organization, and shell

evolution. *Evol. Dev.* **2**, 340-347 (2000).

Kin, K., Kakoi, S. & Wada, H. Novel role for dpp in the shaping of bivalve shells revealed in a

conserved molluscan developmental program. *Dev. Biol.* **329**, 152–166 (2009).

Kniprath, E. Ontogeny of the molluscan shell field. *Zoologica Scripta* **10**, 61–79 (1981).

Kudoh, T., Wilson, S. W. & Dawid, I. B. Distinct roles for Fgf, Wnt and retinoic acid in
posteriorizing the neural ectoderm. *Development* **129**, 4335-4346 (2002).

Lans, D., Wedeen, C. J. & Weisblat, D. A. Cell lineage analysis of the expression of an
engrailed homolog in leech embryos. *Development* **117**, 857-871 (1993).

Lowe, C. J., Wray, G. A. Radical alterations in the roles of homeobox genes during echinoderm
evolution. *Nature* **16**, 718-721. (1997)

Lambert DJ. Developmental Patterns in Spiralian Embryos *Curr Biol.* 20, R72-R77 (2010).

Malinda, K. M. & Ettensohn, C. A. Primary mesenchyme cell migration in the sea urchin
embryo: distribution of directional cues. *Dev. Biol.* **164**, 562–578 (1994).

Manzanares, M., et al. Conservation and elaboration of Hox gene regulation during evolution
of the vertebrate head. *Nature* **408**, 854– 857 (2000).

Marshall, H., et al. A conserved retinoic acid response element required for early expression
of the homeobox gene Hoxb-1. *Nature* **370**, 567–571 (1994).

Moshel, S. M., Levine, M. & Collier, J. R. Shell differentiation and engrailed expression in the
Ilyanassa embryo. *Dev. Genes Evol.* **208**, 135-141 (1998).

Nederbragt, A. J., van Loon, A. E. & Dictus, W. J. Expression of Patella vulgata orthologs of
engrailed and dpp-BMP2/4 in adjacent domains during molluscan shell development
suggests aconserved compartment boundary mechanism. *Dev. Biol.* **246**, 341-55 (2002).

Nüsslein-Volhard, C. & E. Weischaus. Mutations affecting segment number and polarity in
Drosophila. *Nature* **287**, 795– 801 (1980).

Ohta S, Noshita K, Kimoto K, Ishikawa A, Sato H, Shimizu K, Endo K. Possible roles of Wnt in
the shell growth of the pond snail Lymnaea stagnalis. *Sci Rep.* 14:26488. (2024)Patel, N.
H., Kornberg, T. B. & Goodman, C. S. Expression of engrailed during segmentation in
grasshopper and crayfish. *Development* **107**, 201–212 (1989).

Perry KJ, Lyons DC, Truchado-Garcia M, Fischer AH, Helfrich LW, Johansson KB, Diamond
JC, Grande C, Henry JQ. Deployment of regulatory genes during gastrulation and germ
layer specification in a model spiralian mollusc Crepidula. *Dev Dyn.* **244,** 1215-48. (2015).

Ramskold, L. & Hou, X. New early Cambrian animal and onychophoran affinities of enigmatic
metazoans. Nature 351: 225–228 (1991).

Reijntjes, S., Blentic, A., Gale, E., & Maden, M. The control of morphogen signalling: regulation
of the synthesis and catabolism of retinoic acid in the developing embryo. *Dev. Biol.* **285**,
224–237 (2005).

Romero, R. & Bueno, D. Disto-proximal regional determination and intercalary regeneration in
planarians, revealed by retinoic acid induced disruption of regeneration. *Int. J. Dev. Biol*.

45, 669–673 (2001).

Schilling, T. F., & Knight, R. D. Origins of anteroposterior patterning and Hox gene regulation

during chordate evolution. *Philos. Trans. R. Soc. Lond. B. Biol. Sci.* **356** 1599–1613 (2001).

Seaver, E. C. & Kaneshige, L. M. Expression of 'segmentation' genes during larval and juvenile

development in the polychaetes Capitella sp. I and H. elegans. *Dev. Biol*. **289**, 179–194

(2006).

Shimeld, S. M. Retinoic acid, hox genes and the anterior-posterior axis in chordates.

BioEssays **18**, 613–616 (1996).

Shimizu, K., Sarashina, I., Kagi, H. & Endo, K. Possible functions of dpp in gastropod shell

formation and shell coiling. *Dev. Genes Evol.* **221**, 59–68 (2011).

Shimizu, K., Iijima, M., Setiamarga, D. H. E., Sarashina, I., Kudoh, T., Asami, T., Gittenberger,

E. & Endo, K. Left-right asymmetric expression of dpp in the mantle of gastropods

correlates with asymmetric shell coiling. *EvoDevo* **4**, 15. (2013).

Shimizu, K., Luo, Y-J., Satoh, N. & Endo, K. Possible co-option of engrailed during brachiopod

and   mollusc   shell   development.   *Biol.*   *Lett.*   **13**,   20170254.

http://dx.doi.org/10.1098/rsbl.2017.0254 (2017).

Vogeler, S., Galloway, T. S., Isupov, M. & Bean, T. P. Cloning retinoid and peroxisome

proliferator-activated nuclear receptors of the Pacific oyster and in silico binding to

environmental   chemicals.   *PLoS*   *ONE*   **12**,   e0176024.   https://doi.

org/10.1371/journal.pone.0176024 (2017).

Vachon, G., Cohen, B., Pfeifle, C., McGuffin, M. E., Botas, J. & Cohen, S. M. Homeotic genes

of the bithorax complex repress limb development in the abdomen of the Drosphila embryo

through the target gene *Distal-less*. *Cell* **71**, 437-450 (1992).

Wada, H. Origin and evolution of the neural crest: a hypothetical reconstruction of its

evolutionary history. *Dev. Growth Differ.* **43**, 509– 520 (2001).

Wanninger, A. & Haszprunar, G. The expression of an engrailed protein during embryonic shell

formation of the tusk-shell, *Antalis entalis* (Mollusca, Scaphopoda). *Evol. Dev.* **3**, 312-21

(2001).

Wedeen, C. J., & Weisblat, D. A. Segmental expression of an engrailed-class gene during early

development and neurogenesis in an annelid. *Development* **113**, 805–814 (1991).

Wedeen, C. J., Kostriken, R. G., Leach, D., & Whitington, P. Segmentally iterated expression

of an engrailed-class gene in the embryo of an Australian onychophoran. *Dev. Genes Evol.*

270, 282-286 (1997).

Yang, J., Martin, R. S., Lan, T., Hou, J. & Zhang, X. Articulated Wiwaxia from the Cambrian

Stage 3 Xiaoshiba Lagerstätte. *Scientific Reports* **4** (2014).

Zamora, S., Sumrall, C. D. & Vizcaïno, D. Morphology and ontogeny of the Cambrian

edrioasteroid echinoderm Cambraster cannati from western Gondwana. *Acta Palaeontol.*

Polonica (2013).

Zhao, F., Bottjer, D. J., Hu, S., Yin, Z. & Zhu, M. Complexity and diversity of eyes in early

Cambrian ecosystems. *Sci. Rep.* **3**, 2751 (2013).

Table 1 Number of gene expressed-larvae in the oyster and limpet.

C. gigas (18 hpf)

	Trial	*tyr*	*smp-L1*	*collagen*	*en*	*cyp26*
Control	1	36/38	35/40	39/42	47/51	
	2	50/55	32/35	40/45	39/40	
RA (2 µM)	1	34/40	35/39	30/37	32/40	
	2	48/55	42/46	28/38	40/48	
Ro (2 µM)	1	3/35	4/50	2/40	8/44	
	2	5/52	41/48	5/44	3/35	

N. fuscoviridis (10 hpf)

	Trial	*cs-1*	*6D-12*	*en*	*dpp*	*cyp26*
Control	1	32/32	30/30	33/33	39/40	24/24
	2	40/40	34/34	34/35	33/33	35/35
RA (2 µM)	1	35/40	40/40	28/38	25/25	32/32
	2	31/38	32/32	29/40	36/37	44/44
Ro (2 µM)	1	8/32	5/37	10/45	29/32	37/37
	2	12/45	8/40	17/50	36/38	52/52