Title: Environmental dynamics influence scale cortisol levels in Mexican cavefish

Authors: Marc B. Bauhus[1], Luise Wander[1], David Boy[1], Ana Santacruz[2,3], Ernesto Maldonado[4], Joachim Kurtz[1,5], Sylvia Kaiser[5,6] and Robert Peuß[1,2,5*]

Affiliations:
[1]: Institute for Evolution and Biodiversity, University of Münster, Germany
[2]: Institute for Integrative Cell Biology and Physiology, University of Münster, Germany
[3]: Stowers Institute for Medical Research, Kansas City, MO, USA
[4]: EvoDevo Research Group, Unidad Académica de Sistemas Arrecifales, Instituto de Ciencias del Mar y Limnología, Universidad Nacional Autónoma de México, Puerto Morelos, Quintana Roo, Mexico
[5]: Joint Institute for Individualisation in a Changing Environment, University of Münster and Bielefeld University, Münster, Bielefeld, Germany
[6]: Department of Behavioural Biology, University of Münster, Germany

* Corresponding author:
Robert Peuß: robert.peuss@uni-muenster.de

Keywords: Stress, Environment, Scale Cortisol, *Astyanax mexicanus*

Abstract:
Cortisol is a key hormone involved in stress responses, and its regulation can vary across populations experiencing different environmental pressures. While short term changes in certain environmental factors can result in immediate changes in cortisol levels, it remains unclear whether variable and fluctuating environments affect lifetime stress responses of individuals. In this study, we estimated lifetime-experienced stress in cave- and surface-dwelling populations of *Astyanax mexicanus*. The advantage of this model is that different populations are confronted with varied dynamics of environmental change, making it an ideal study system to investigate lifetime stress responses upon environmental fluctuations. To estimate lifetime-experienced stress, we measured cortisol levels in the scales of *A. mexicanus* for the first time using an ELISA assay. We studied six populations from six different habitats (four cave and two surface habitats) that vary in their dynamics of abiotic factors. We analysed scale cortisol levels across two years to examine temporal changes and individual variability within and between populations. Our results show that increased fluctuations in abiotic parameters are associated with an increased cortisol level in scales and, in addition, an increased inter-individual variability within a population irrespective of cave or surface habitat. These findings highlight the influence of varying ecological conditions on lifetime-experienced stress in *A. mexicanus*, providing insight into potential adaptive mechanisms underlying stress regulation in extreme environments.

Introduction

Stress is a response to environmental challenges or threats (stressors), triggering a complex physiological and behavioral reaction that enables individuals to cope with environmental changes (Schulte et al., 2014). Stress can be induced by changes in abiotic conditions such as temperature (Gates et al., 1992), oxygen levels (Van Raaij et al., 1996) and pH (Wang et al., 2009) or by biotic factors such as predators (Van Dievel et al., 2016), pathogens (Huffman et al., 2004), aggressive interactions among conspecifics (Sachser and Lick, 1991), and food availability (Pravosudov et al., 2001). During the vertebrate stress response, the adrenal glands (or interrenal glands in fish) produce glucocorticoids, for example cortisol or corticosterone, which has downstream effects on metabolism (Brillon et al., 1995; Christiansen et al., 2007; Vijayan et al., 1997) and immunity (Webster Marketon and Glaser, 2008). Populations and individuals can vary in their sensitivity to stressors and their stress responses, thereby adapting to highly variable environments (DeKoning et al., 2004; Schulte et al., 2014). However, it remains unclear whether highly variable and fluctuating environments impose greater lifetime stress on individuals compared to more stable environments.

To investigate how fluctuations in abiotic parameters affect the cortisol levels, we use the Mexican Tetra *Astyanax mexicanus*. This model exhibits an extreme example of local adaptation to different habitats and environmental conditions. *A. mexicanus* consists of two ecotypes: surface-dwelling (surface fish) and cave-dwelling (cavefish), both distributed in north-eastern Mexico (Mitchell et al., 1977, see Fig. 1A). Cave aquatic environments present stressors such as reduced and infrequent availability of nutritional resources due to the absence of primary energy production in total darkness (Poulson, 2001; Venarsky et al., 2014). However, they also provide more stable abiotic conditions (Krishnan et al., 2020; Poulson and White, 1969; Tabin et al., 2018) but with differences between caves and pools within caves depending on the hydrological regime (Legendre et al., 2025). Consequently, cavefish have evolved several adaptations that allow them to thrive in resource-limited conditions. These include metabolic adaptations such as elevated blood sugar levels and insulin resistance (Riddle et al., 2018; Shimobayashi et al., 2023), enhanced lipogenesis (Xiong et al., 2022) and increased fat accumulation (Xiong et al., 2018). Behavioral adaptations include binge eating (Aspiras et al., 2015), reduced sleep (Yoshizawa et al., 2015) and decreased aggression (Elipot et al., 2013). Additionally, cavefish must cope with hypoxia (Krishnan et al., 2020; Ornelas-García et al., 2018; Rohner et al., 2013) which has led to an increase in erythrocyte production (van der Weele and Jeffery, 2022) as well as larger erythrocytes with higher hemoglobin content per cell (Boggs et al., 2022).

Unlike cavefish, surface-dwelling *A. mexicanus* experience greater fluctuations in abiotic conditions, such as temperature variation (Tabin et al., 2018), and are exposed to more biotic stressors, including parasites (Peuß et al., 2020; Santacruz et al., 2023, 2020) predators (Chin et al., 2018; Mitchell et al., 1977; Yoshizawa and Library, 2015) and aggressive interactions among conspecifics (Elipot et al., 2013; Rodriguez-Morales et al., 2022).

Strong ecological differences between cave and surface habitats resulted in specific stress response adaptations of cave- and surface fish. For example, cavefish show lower plasma cortisol levels upon exposure to confined space (Gallo and Jeffery, 2012), lower whole-body

cortisol in total darkness (Bilandžija et al., 2020), and reduced stress behaviors (Chin et al.,
2018; Padmanaban et al., 2025) compared to surface fish.

While short-term elevations in blood-plasma cortisol occur in response to acute stress, scale
cortisol levels provide a measure of chronic or lifetime-experienced stress, as cortisol
accumulates in scales over time (Aerts et al., 2015). Consequently, scale cortisol levels have
been widely used to assess lifetime stress across various fish species (Britton et al., 2023;
Carbajal et al., 2019; Lebigre et al., 2022; Roque d'orbcastel et al., 2021; Vercauteren et al.,
2022)

To examine differences in lifetime-experienced stress between surface and cave-adapted *A.*
mexicanus, we measured cortisol levels in scales of wild surface and cave individuals from
different populations across two consecutive years. We sampled two distinct surface
populations: Rio Choy (a stream population) and Presa El Oyul (a lake population) – as well
as four cave populations: Pachón, Los Sabinos, Tinaja and Rio Subterráneo - which represent
an isolation gradient from surface habitats (Mitchell et al., 1977; Moran et al., 2023; Simon et
al., 2017; see Fig. 1A-D). Additionally, we measured several abiotic parameters in both years
to assess fluctuation in abiotic parameters across habitats. To disentangle environmental and
genetic effects on cortisol levels (Krishnan et al., 2020), we compared scale cortisol levels from
wild individuals to those of laboratory-reared individuals from the same population. We
hypothesized that scale cortisol levels, i.e. life-time experienced stress increases with a more
dynamic and fluctuating environment. Therefore, we predicted lower scale cortisol levels and
individual variation in cavefish compared to surface fish due to the less dynamic and
fluctuating cave environment.

Results

Scale cortisol levels

To compare lifetime-experienced stress across wild and laboratory populations of cave and
surface fish, we measured scale cortisol levels using an ELISA assay. We fitted a generalized
linear model (GLM) with scale cortisol level as response, population as the predictor and body
condition (Fulton's K-index), standard length (as a proxy for age), and sex as covariates. Our
results indicate that wild Pachón cavefish (Fig. 1H) in 2023 had the lowest scale cortisol levels
(EMM: 8.74 pg/mg, 95% CI: 5.65–13.5; Fig. 1E, see Supplemental Table 1 for standard errors),
followed by laboratory-raised Pachón cavefish (EMM: 9.24, 95% CI: 6.09–14.01; Fig. 1E) and
wild Los Sabinos cavefish (Fig. 1I) (EMM: 12.9, 95% CI: 10.11–16.46; Fig. 1E). Wild Tinaja
cavefish (Fig. 1J) exhibited intermediate cortisol levels in both 2023 (EMM: 22.51, 95% CI:
14.71–34.44; Fig. 1E) and 2024 (EMM: 23.87, 95% CI: 16.76–34.0; Fig. 1E). Interestingly,
Rio Choy surface fish (Fig. 1F) showed similar cortisol levels in both wild and laboratory
conditions: EMMs were 23.08 (95% CI: 17.98–29.62) in 2023, 25.35 (95% CI: 19.67–32.67)
in 2024, and 21.17 (95% CI: 14.95–29.98) in the lab (Fig. 1E). The highest cortisol levels were
observed in wild Presa El Oyul surface fish (Fig. 1G), with an EMM of 43.87 (95% CI: 35.24–
54.61) in 2023 and 30.15 (95% CI: 24.01–37.87) in 2024 (Fig. 1E). Rio Subterráneo cavefish
(Fig. 1K) exhibited intermediate cortisol levels in 2023 (EMM: 19.38, 95% CI: 15.27–24.6),
but levels increased in 2024 (EMM: 37.32, 95% CI: 28.4–49.05) (Fig. 1E).

126
Figure 1. Scale cortisol levels of wild and laboratory *A. mexicanus* populations. (A) Map of sampling area in the
Sierra de El Abra Tanchipa and Sierra Micos region in north-eastern Mexico with sampling locations of respective
populations. (B) Stream surface fish habitat (Nacimiento Rio Choy). (C) Lake surface fish habitat (Presa El Oyul).
(D) Cavefish habitat (Pachón) cave. (E) Scale cortisol levels in pg/mg scales per population and sampling year

with respective sample size (N) in brackets. The raw data points are plotted together with violins to show the
distribution per population. Large points with error bars show the estimated marginal means with 95% confidence
intervals under the fitted generalized linear model with scale cortisol as response, population as predictor and
body condition (K-index), mean-centred standard length and sex as covariates. Significant differences (adjusted
p-value <0.05) between populations are indicated by the compact letter design with two populations being
significantly different when no letter overlaps. Representative images of (F) Rio Choy and (G) Presa El Oyul
surface- and (H) Pachón, (I) Los Sabinos, (J) Tinaja and (K) Rio Subterráneo cavefish.

Pairwise comparisons between populations

To assess statistical differences between populations and across years within populations, we
conducted pairwise comparisons based on EMMs (see Supplemental Table 2 for all pairwise
comparisons). Wild and laboratory Pachón cavefish and wild Los Sabinos cavefish had
significantly lower scale cortisol levels than wild Rio Choy surface fish in both years (Fig. 1E,
Supplemental Table 2). The largest differences in cortisol levels were observed between wild
and lab Pachón and Los Sabinos cavefish and wild Presa El Oyul surface fish in both years
(Fig. 1E, Supplemental Table 2). Comparisons between wild Tinaja cavefish and both wild Rio
Choy and Presa El Oyul surface fish did not yield any significant differences (Fig. 1E,
Supplemental Table 2). However, wild Tinaja cavefish in 2024 exhibited higher cortisol levels
than wild Pachón cavefish and showed a tendency toward higher levels compared to laboratory
Pachón cavefish (Fig. 1E, Supplemental Table 2). A notable within-population difference was
found in Rio Subterráneo cavefish, where individuals caught in 2024 had significantly higher
cortisol levels than those caught in 2023 (Fig. 1E, Supplemental Table 2). Finally, Presa El
Oyul surface fish caught in 2023 had significantly higher cortisol levels than both laboratory
and wild Rio Choy surface fish caught in the same year and showed a trend toward higher
levels compared to Rio Choy surface fish in 2024 (Fig. 1E, Supplemental Table 2).

Covariate effects on scale cortisol levels

Overall, body condition (Fulton's k-index) was a significant predictor of cortisol levels
(β=0.509, SE=0.178, p=0.0047, see Supplemental Figure 3, Model Summary 1), indicating that
individuals with better body condition exhibited higher cortisol levels. The effect of mean-
centred standard length on cortisol was negative but not statistically significant ($\beta = -0.084$,
$SE = 0.048$, $p = 0.085$, see Supplemental Figure 4, Model Summary 1), suggesting a trend
where larger individuals might have lower cortisol levels. Similarly, sex did not significantly
influence cortisol levels ($\beta = -0.104$, $SE = 0.097$, $p = 0.288$, see Supplemental Information,
Model Summary 1).

Table 1. Abiotic water parameters in surface (Rio Choy, Presa El Oyul) and cave (Pachón, Los Sabinos, Rio Subterráneo, Tinaja) habitats measured in spring 2023 and 2024.

Site	Year	Temp. [C°]	pH	LDO [mg/L]	Cond. [µs/cm]	PO4 [mg/L]	CaCO3 [mg/L]	Fe [mg/L]	Cu [mg/L]	NO2 [mg/L]
Rio Choy	2023	25.9	7	7.36	1164.5	0.75	710.5	0	0	0.004
Rio Choy	2024	25.8	7.26	6.8	1123	0.06	560	0	0.02	0.005
Presa El Oyul	2023	26	8.47	7.59	428	1	214	0.05	0.04	0.002
Presa El Oyul	2024	30.5	9.5	9.4	316	0.6	64	0.13	0	1E-10
Pachón	2023	24.35	7.35	5.345	568	0.45	307	0.005	0.005	0.013
Pachón	2024	25.7	7.58	5.8	490	0.6	208	0	0.01	0.027
Los Sabinos (pool 1)	2023	22.8	7.43	1.97	484	1.3	251	0	0	0.065
Los Sabinos (pool 1)	2024	23.1	7.84	1.73	534	0.6	239	0	0.03	0.177
Tinaja (pool 2)	2024	23.3	8.17	6.29	667	2.4	346	0	0	0.009
Rio Subterráneo (pool 1)	2023	23.7	7.22	1.76	470	1.3	232	0	0	0.229
Rio Subterráneo (pool 2)	2023	23.8	7.33	5.77	481	1.4	228	0	0	0.048
Rio Subterráneo (pool 1)	2024	25	7.59	3.37	524	1.9	234	0	0	0.007
Rio Subterráneo (pool 2)	2024	25.1	7.25	3.38	519	4.7	212	0	0	0.063

Abiotic parameters

To explore the impact of abiotic factors on scale cortisol levels, we measured temperature (°C), pH, liquid dissolved oxygen (mg/L), conductivity (µs/cm), phosphate (mg/L), hardness (CaCO₃) (mg/L), iron (mg/L), copper (mg/L), nitrite (mg/L), and ammonium (mg/L) in each habitat (Table 1). A principal component analysis (PCA) based on these parameters revealed notable differences not only between cave and surface habitats but also between the stream (Rio Choy) and lake (Presa El Oyul) surface habitats (Fig. 2A). Additionally, Presa El Oyul exhibited significant differences in some parameters between 2023 and 2024, primarily reflected in principal component (PC) 1 (2023: 2.257, 2024: 5.416) (Fig. 2A). PCA loading factors indicated that the variation in PC 1 was primarily driven by temperature (0.455), pH (0.47), dissolved oxygen (0.396), and iron (0.498) (Fig. 2B). We correlated scale cortisol levels with PC 1 and fitted a simple GLM with cortisol as the response variable and abiotic PC 1 as the predictor. We found that abiotic PC 1 had a significantly positive effect on cortisol (β=0.095, SE=0.022, p<0.0001), indicating that higher PC 1 values correspond to elevated scale cortisol levels (Fig. 2C).

Figure 2. Dynamics of abiotic parameters influencing scale cortisol levels and individual variation in *A. mexicanus*. (A) Principal component analysis (PCA) plot based on parameters from Table 1 showing scores of principal components (PC) 1 (x-axis) and 2 (y-axis) per habitat and year. The percentage of explained variation per PC is displayed in brackets. (B) Heatmap showing PC loading factors per measured parameter. (C) Correlation of individual scale cortisol levels with PC 1 scores for each population and sampling year. The regression line represents the predicted regression with 95% confidence intervals (shaded area) of a generalized linear model with scale cortisol levels as response and PC 1 as predicting variables. (D) Individual variation in scale cortisol levels displayed by 95% confidence intervals (error bars) with means (points) based on bootstrap resampling (1000 iterations) per population and year.

Individual variation in scale cortisol levels

To examine individual variation in scale cortisol levels across populations, we performed bootstrap resampling with 1000 iterations and reported 95% confidence intervals (CIs). We found that Los Sabinos cavefish in 2023 exhibited the lowest degree of individual variation (95% CI: 12.56–48.83), followed by wild Rio Choy surface fish in 2023 (95% CI: 19.3–68.63), Tinaja cavefish in 2024 (95% CI: 21.12–75.14), wild Pachón cavefish in 2023 (95% CI: 3.14–59.77), Tinaja cavefish in 2023 (95% CI: 15.55–94.94), and laboratory Pachón cavefish (95% CI: 10.25–93.98) (Fig. 2D). More individual variation was observed in Rio Subterráneo cavefish in 2023 (95% CI: 55.95–282.39), wild Rio Choy surface fish in 2024 (95% CI: 16.28–271.85), laboratory Rio Choy surface fish (95% CI: 79.49–612.57), and Presa El Oyul surface fish in 2024 (95% CI: 104.87–954.53) (Fig. 2D). The highest degree of individual variation in scale cortisol levels was found in Presa El Oyul surface fish in 2023 (95% CI: 58.54–1018.52) and Rio Subterráneo cavefish in 2024 (95% CI: 267.53–908.98) (Fig. 2D). Significant differences in individual variation (non-overlap in CIs) were observed between laboratory Rio Choy surface fish and wild Pachón, Los Sabinos, and Tinaja cavefish in 2024 (Fig. 2D). Additionally, Rio Subterráneo cavefish in 2023 and Presa El Oyul surface fish in 2023 showed significantly higher individual variation compared to Los Sabinos cavefish (Fig. 2D). In 2024, both Presa El Oyul surface fish and Rio Subterráneo cavefish exhibited consistently higher individual variation compared to all other cave populations and wild Rio Choy surface fish in 2023 (Fig. 2D).

Discussion

Cave stressors

We found substantial evidence that life-time stress in *A. mexicanus* is connected to variability of specific abiotic factors. Our results show that increased fluctuations in abiotic parameters are associated with increased cortisol levels in both cave and surface habitats. Furthermore, low scale cortisol levels of Pachón and Los Sabinos cavefish suggest that those fish are adapted to extreme environmental conditions such as food scarcity. Specific metabolic and behavioural adaptations (Aspiras et al., 2015; Olsen et al., 2023; Riddle et al., 2018; Shimobayashi et al., 2023; Xiong et al., 2022, 2018; Yoshizawa et al., 2015) enable cavefish to thrive upon resource-limited conditions in the cave environment over long periods. However, cortisol showed an ambiguous role in response to starvation in other fish, being sometimes elevated (Arslan et al., 2016; Caruso et al., 2010) and sometimes reduced or unaffected (Poursaeid and Falahatkar, 2022; Sumpter et al., 1991; White and Fletcher, 1986).

Another stressor in caves is hypoxia (Krishnan et al., 2020; Legendre et al., 2025; Ornelas-García et al., 2018). Our measurements confirmed lower liquid dissolved oxygen levels in all cave habitats compared to surface habitats, with particularly low levels in Los Sabinos cave. Low scale cortisol levels in Pachón and Los Sabinos cavefish (Fig. 1E) may indicate that hypoxia adaptations (Boggs et al., 2022; van der Weele and Jeffery, 2022) contribute to a reduced life-time stress response. However, the effects of hypoxia on cortisol release in other fish species remain inconsistent (Léger et al., 2021; O'Connor et al., 2011).

Dynamics in abiotic parameters

While Tinaja and Rio Subterráneo cavefish experience similar hypoxic conditions and show similar starvation adaptations as Pachón and Los Sabinos fish (Aspiras et al., 2015; Boggs et al., 2022; Legendre et al., 2025; Ornelas-García et al., 2018; Riddle et al., 2018; Wilkens and Hüppop, 1986), their higher scale cortisol levels in 2024 suggest the presence of additional stressors. Hydrological regimes influence the dynamics of physico-chemical water parameters in cave habitats (Legendre et al., 2025). Unlike Los Sabinos, Tinaja, and Rio Subterráneo caves, Pachón cave is not affected by seasonal epigean water inflow due to its elevated entrance (210 m asl) (Mitchell et al., 1977). Our PCA analysis showed minimal annual variation in measured water parameters for Pachón, Los Sabinos, and Rio Subterráneo caves. However, long-term data revealed that Pachón cave exhibits the most stable conditions, whereas Los Sabinos and Rio Subterráneo experience seasonal fluctuations in temperature, pH, and conductivity due to epigean water inflow (Legendre et al., 2025; Tabin et al., 2018). Such fluctuations are known to elevate cortisol and stress responses in fish (Choi and An, 2008; Cockrem et al., 2019; Mota et al., 2017; Rohner et al., 2013; Wagner et al., 1997), potentially explaining higher scale cortisol levels in Tinaja and Rio Subterráneo cavefish in 2024. Higher scale cortisol levels in Rio Subterráneo in 2024 compared to 2023 suggest differences in experienced stress across years in this cave. We observed drastically lower water levels and even the disappearance of smaller pools in this cave in 2024 compared to 2023 (personal observation, Marc Bauhus), potentially due to lower precipitation in the area ("Servicio Meteorológico Nacional," n.d.). Alongside potential variations in abiotic dynamics, this could suggest heightened stress due to increased population density and resource competition, both of which can elevate cortisol levels (Bolasina et al., 2006; Ramsay et al., 2006; Ruane and Komen, 2003).

Rio Choy surface fish had similar cortisol levels to Tinaja and Rio Subterráneo cavefish, despite differing abiotic conditions. Our data showed no significant annual variation in abiotic parameters in Rio Choy, but seasonal fluctuations—undetected due to our single annual measurements—may still impact cortisol levels. Long-term monitoring of stream habitats is necessary to fully understand the environmental dynamics affecting surface fish stress levels.

In contrast to stream and cave habitats, we observed a much greater dynamic in abiotic parameters in the lake habitat Presa El Oyul. The difference in abiotic conditions in the lake between spring 2023 and 2024 was similarly high as the difference to other habitats such as Rio Choy and the caves, with parameters being the most extreme in 2024. This was mostly driven by higher temperature, pH, oxygen and iron levels. In addition to the beforementioned potential effects of temperature, pH and oxygen fluctuations on cortisol levels, increased iron concentrations were shown to affect cortisol and physiological stress responses in whitefish (Lappivaara et al., 1999; Lappivaara and Marttinen, 2005). We could show that scale cortisol levels were positively correlated with increased abiotic PC 1 values, suggesting an association of temperature, pH, oxygen and iron with lifetime stress levels across populations

Biotic stressors

Biotic stressors, such as increased predation risk in the lake habitat, could likewise explain increased scale cortisol levels and lifetime stress. During fieldwork, we observed piscivorous birds and predatory fish in Presa El Oyul Lake (personal observation, Marc Bauhus). Studies in other fish report mixed effects of predation risk on cortisol levels (Fischer et al., 2014;

Humphrey et al., 2020). In *A. mexicanus*, surface fish show more predation associated stress
behaviors, such as prolonged bottom-dwelling and freezing, compared to cavefish (Chin et al.,
2018; Padmanaban et al., 2025). Given that these behaviors are mediated by glucocorticoid
hormones (De Marco et al., 2013; Ziv et al., 2012) surface *A. mexicanus* likely experience
greater predation-induced cortisol responses. However, differences in scale cortisol levels
between Presa El Oyul and Rio Choy in 2023 could suggest that predation pressures vary
among surface populations. Further studies are needed to clarify this relationship.
Parasite infections can also trigger elevated glucocorticoid hormone responses in fish (Allan et
al., 2020; O'Dwyer et al., 2020). Therefore, variations in parasite abundance and diversity
between and among cave and surface populations may account for the higher stress levels
observed in Rio Choy and Presa El Oyul surface fish, as well as Tinaja and Rio Subterráneo
cavefish in 2024, compared to Pachón cavefish. This difference could be due to the reduced
parasite diversity in the Pachón cave population (Peuß et al., 2020; Santacruz et al., 2023,
2020).
Another potential stressor is intraspecific aggression. Cavefish display reduced aggressive
behaviours compared to surface fish (Elipot et al., 2013; Rodriguez-Morales et al., 2022).
Aggressive interactions among conspecifics cause stress in fish (Winberg and Lepage, 1998),
suggesting that reduced aggression in cavefish could contribute to their lower scale cortisol
levels.
A comparison of wild and lab populations of Pachón cavefish and Rio Choy surface fish,
respectively, revealed no significant differences in scale cortisol levels between lab-reared and
wild-caught fish. However, a trend toward higher cortisol levels in lab surface fish compared
to lab cavefish suggests a potential genetic basis for stress differences. Given that
environmental conditions were standardized in the lab, differences in aggression behaviour
could be a contributing factor. Genetic mapping of scale cortisol levels through crossbreeding
experiments could help identify underlying genetic determinants.
306
Individual variation in scale cortisol levels
We found evidence that individual variation in scale cortisol levels varies across populations.
Low individual variation in Los Sabinos, Pachón, and Tinaja cavefish suggests that these cave
habitats are characterised by low fluctuations in stressors, leading individuals to occupy similar
niches within the given habitat. Adaptations to strong and consistent selective pressures
(Poulson, 2001; Venarsky et al., 2014), such as limited food availability and stable, yet
sometimes extreme abiotic conditions (Krishnan et al., 2020; Legendre et al., 2025; Poulson
and White, 1969; Tabin et al., 2018), may contribute to reduced individual variation in stress
responses and lifetime stress levels. This reduced variability could, in turn, limit the ability to
conform to niche changes through stress-related mechanisms. Interestingly, Rio Choy surface
fish did not exhibit greater individual variation in scale cortisol levels than the mentioned cave
populations. This could be due to adaptations to different but equally stable selective pressures.
In contrast, Presa El Oyul Lake exhibits greater environmental heterogeneity and dynamic
stressors, including fluctuating abiotic parameters, and potentially, predation, and aggressive
interactions among conspecifics, which may explain the higher individual variation in scale
cortisol levels. In Rio Subterráneo, periodic flooding from nearby surface streams, along with
the introduction and hybridization of surface individuals (Garduño-Sánchez et al., 2023; Moran

et al., 2023) together with invasive parasites (Santacruz et al., 2023), could have contributed to greater individual variation in scale cortisol levels. This variation may stem from abiotic fluctuations or from gradients of cave-adapted traits across the sampled individuals leading to differences in stress responses.

Scale cortisol as a measure for lifetime stress

Surprisingly, we found a significant positive relationship between body condition (K-index) and scale cortisol levels, contrasting with previous studies in other fish species, where higher scale cortisol levels were either associated with reduced body condition (Brosset et al., 2023) or showed no correlation (Carbajal et al., 2019). One possible explanation is that cavefish exhibit increased fat accumulation (Xiong et al., 2022, 2018), which elevates Fulton's K-index through increased body mass. In humans, obesity and metabolic syndrome are linked to increased cortisol levels and hyperactivity of the hypothalamic-pituitary-adrenal axis (Anagnostis et al., 2009). A similar mechanism may influence cortisol regulation in *A. mexicanus*, warranting further research into interactions between cortisol, fat storage, and body condition in this species.

Additionally, we observed a tendency toward a negative relationship between standard length and scale cortisol levels (see Supplemental Information). Standard length was included as a proxy for age in the model, given its positive correlation with age despite differing growth rates between cave and surface fish (Simon et al., 2017). However, previous studies have shown that scale cortisol levels tend to increase with age (Lebigre et al., 2022), raising the possibility that standard length is an inadequate proxy for age in *A. mexicanus*. Alternatively, our results could reflect a selection bias, where individuals with chronically high cortisol levels experience higher mortality (Pickering and Pottinger, 1989), leading to the selective survival of individuals with lower cortisol levels in older (larger) age classes. In addition, exclusion of some small individuals because of insufficient scale material could have further exacerbated this potential source of error. Future studies should aim to integrate precise age estimates, such as counting annual growth increments on scales or otoliths (Simon et al., 2017), into scale cortisol analyses of wild populations and directly test whether cortisol accumulates in scales over time in *A. mexicanus* to support its suitability as a measure for lifetime-experienced stress. Furthermore, the actual responsiveness to cortisol needs to be further investigated in this system since there could be individual or population differences due to varying glucocorticoid receptor expression in the brain, as shown for example in arctic charr (Vindas et al., 2017).

Conclusion

Our findings suggest that lifetime stress, as indicated by scale cortisol levels, is lower in isolated cave populations of *A. mexicanus*, likely due to their stable, yet extreme environments, and associated adaptations. In contrast, surface fish experience greater environmental fluctuations and biotic stressors, leading to higher and more variable cortisol levels. Environmental dynamics, particularly hydrological fluctuations and associated changes in abiotic parameters, appear to shape life-time stress responses across populations, emphasizing the complex interplay between abiotic and biotic factors in shaping stress physiology. Future research should experimentally test the effects of specific stressors, such as hypoxia, starvation,

367 abiotic parameters, predation, parasites and aggression on cortisol regulation and explore the
368 genetic basis of stress-related differences between ecotypes. Our study highlights the
369 importance of scale cortisol as a biomarker for long-term stress and provides insights into the
370 role of environmental stability in shaping stress adaptations.
371

Material & methods

Field sample collection

We collected wild individuals (males and females) of *A. mexicanus* in 2023 and 2024 under
permit no. SPARN/DGVS/03371/23 granted by the Secretaría de Medio Ambiente y Recursos
Naturales (SEMARNAT) to Ernesto Maldonado. In 2023, we sampled fish from Rio
Subterráneo (Micos), Pachón and Los Sabinos caves as well as from Nacimiento Rio Choy
(stream) and Presa El Oyul (lake) between April 5th and April 15th. Additionally, we collected
fish from Tinaja cave between December 2nd and 3rd. In 2024, we resampled Rio Subterráneo,
Tinaja, Nacimiento Rio Choy and Presa El Oyul (only females) between March 9th and March
19th.
In cave environments, we used hand nets to capture fish, while in stream and lake habitats, we
employed improvised fishing poles equipped with size 16 hooks (Gamakatsu, Japan) baited
with tortilla or dried mango to catch surface fish. We euthanized all fish with an overdose of
MS-222 (0.5 g/L) and dissected them within 24h post capture. We weighed all fish to the
nearest mg and took standardized pictures to determine the standard length of each fish to the
nearest mm with ImageJ. Using a metal spatula, we carefully collected scales and transferred
them to pre-weighed 2 mL Eppendorf-tubes, which we immediately flash froze in liquid
nitrogen. We stored the samples at -20 °C until further processing.
Furthermore, we measured temperature [C°], pH (in 2023), liquid dissolved oxygen [mg/L],
conductivity [μS/cm], phosphate high range [mg/L], hardness ($CaCO_3$) high range [mg/L], iron
[mg/L], copper [mg/L], nitrite [mg/L] and ammonium [mg/L] with a Hach SL 1000
multiparametric device and the respective chemkeys and probes at each sampling except Tinaja
in 2023. Additionally, we measured these parameters in Pachón and Los Sabinos in 2024
despite not sampling fish. For pH measurements in 2024, we used an Akozon Smart Sensor
Pen Type PH Meter Portable Water Quality Tester Acidimeter PH818.
397

Laboratory fish husbandry and sample collection

To generate F1 offspring, we obtained parental Pachón cavefish from the Retaux laboratory
(Paris) and Rio Choy surface fish from the Wilkens laboratory (Hamburg). We grouped at least
four males and two females together for breeding, following the methods described by
(Baumann and Ingalls, 2022). We maintained F1 offspring from both populations at 22°C and
fed them daily with freshly hatched *Artemia* larvae until they reached approximately 10 mm in
size. Between 10 and 30 mm, we provided food five times per week using Gemma Micro. As
they grew between 30 and 45 mm, we adjusted the feeding schedule to four times per week
with Gemma Silk and once per week with frozen bloodworms. After reaching 45 mm, we fed
them twice per week with Gemma Silk and once per week with frozen bloodworms. We housed
adult fish (> 45 mm) in groups of at least six individuals in 14 L aquaria with recirculating tap
water, temperature control, and mechanical, UV, and biological filtration (Vewa Tech, Hamm,

Germany). To minimize aggression, we enriched Rio Choy surface fish tanks with hexagonal
plastic structures. We euthanized fish at 10 months (Rio Choy) and 14 months (Pachón) of age
using an overdose of MS-222 (0.5 g/L) in accordance with the permit T24.016UMS granted by
the University of Münster. We collected scales with a metal spatula and transferred them to
pre-weighed 2 mL Eppendorf tubes, storing them at -20°C until analysis.
415

Extraction of cortisol from scales and determination of corticosterone levels
Before processing, we thawed the scales and washed them three times by adding 1.2 mL of
isopropanol to each tube, shaking them at 20 Hz for 3 minutes in a ball mill (MM 400, Retsch
GmbH, Germany). We discarded the supernatant and allowed the scales to dry overnight in a
fume cupboard before recording their dry weights. Next, we placed two 5 mm steel balls into
each tube and pulverized the scales for 3 minutes at 30 Hz in the ball mill. We weighed up to
13 mg of powdered scales and added 1.3 mL of 100% methanol, maintaining a final
concentration of 10 mg powdered scales/mL methanol. We incubated the samples overnight in
methanol on an overhead shaker. The next day, we centrifuged the samples at $1,000 \times g$ for 1
minute, transferred the supernatant to a new tube, and centrifuged it again at $20,000 \times g$ for 10
minutes. We transferred 1 mL of the supernatant to a new vial and evaporated it in a vacuum
concentrator. We then resuspended the cortisol residue in 250 µL of ELISA buffer provided by
the assay kit (Neogen, product no. #402710, USA). We selected this assay kit based on previous
validations in other fish species (Britton et al., 2023; Carbajal et al., 2018; Ding et al., 2024)
and because it showed the best performance during our own validation trials. We stored the
samples at −20°C until performing the ELISA. Before measurement, we thawed and vortexed
the samples. We determined cortisol concentrations in duplicate following the manufacturer's
instructions. To minimize potential batch effects, we randomly loaded each ELISA plate with
samples from different populations.
435

Statistical analysis
After excluding some individuals due to non-sufficient scale material and a technical problem
in the laboratory, we obtained the following sample sizes per population for statistical analysis:
Rio Choy 2023: N=21, Rio Choy 2024: N=23, Rio Choy lab: N=10, Presa El Oyul 2023: N=23,
Presa El Oyul 2024: N=24, Pachón 2023: N=7, Pachon lab: N=9, Los Sabinos 2023: N=23,
Tinaja 2023: N=8, Tinaja 2024: N=10, Rio Subterráneo 2023: N=21, Rio Subterráneo 2024:
N=16. All statistical analyses were conducted in R (version 4.3.2). To test for differences in
scale cortisol levels between populations, we first assessed normality using Shapiro-Wilk tests
and visualized distributions with histograms and density plots. Since cortisol levels were not
normally distributed but strictly positive, and right-skewed, we fitted generalized linear models
(GLMs) with a Gamma distribution and log-link function. Model selection was based on
Akaike Information Criterion (AIC) values, model diagnostics (residual plots from the car
package) (see Supplemental Fig. 1), and the biological meaningfulness of fixed effects. The
final model included population, Fulton's k-index as a measure for body condition (Barton et
al., 2002), mean-centred standard length and sex as fixed effects. To evaluate the relationship
between cortisol and environmental variation, we performed a principal component analysis
(PCA) on abiotic parameters where we imputed zero values to 1E-10. We then incorporated
the first principal component (PC1) into an additional GLM to assess its effect on cortisol

levels. Model residuals were checked for normality and homoscedasticity. We conducted pairwise comparisons of estimated marginal means (EMMs) using the emmeans package and applied compact letter display (CLD) to summarize significant differences among populations. Additionally, we used bootstrap resampling (1,000 iterations) to compare individual cortisol variability across populations. All statistical tests were two-tailed, and significance was set at $p < 0.05$.

Data availability

Data and code will be made available upon successful publication in an online repository.

References

Aerts J, Metz JR, Ampe B, Decostere A, Flik G, De Saeger S. 2015. Scales Tell a Story on the Stress History of Fish. *PLoS One* 10:e0123411. doi:10.1371/JOURNAL.PONE.0123411

Allan BJM, Illing BR, Fakan EP, Narvaez P, Sgrutter A, Sikkel PC, McClure EC, Rummer JL, McCormick MI. 2020. Parasite infection directly impacts escape response and stress levels in fish. *Journal of Experimental Biology* 223. doi:10.1242/JEB.230904

Anagnostis P, Athyros VG, Tziomalos K, Karagiannis A, Mikhailidis DP. 2009. The Pathogenetic Role of Cortisol in the Metabolic Syndrome: A Hypothesis. *J Clin Endocrinol Metab* 94:2692–2701. doi:10.1210/JC.2009-0370

Arslan G, Sahin T, Hisar O, Hisar SA. 2016. Effects of low temperature and starvation on plasma cortisol, triiodothyronine, thyroxine, thyroid-stimulating hormone and prolactin levels of juvenile common carp (Cyprinus carpio). *Mar Sci Technol Bull* 4:5–9.

Aspiras AC, Rohner N, Martineau B, Borowsky RL, Tabin CJ. 2015. Melanocortin 4 receptor mutations contribute to the adaptation of cavefish to nutrient-poor conditions. *Proc Natl Acad Sci U S A* 112:9668–9673. doi:10.1073/PNAS.1510802112

Barton B, Morgan J, Vijayan M. 2002. Physiological and condition-related indicators of environmental stress in fish.

Baumann DP, Ingalls A. 2022. Mexican tetra (Astyanax mexicanus): biology, husbandry, and experimental protocols. *Laboratory Fish in Biomedical Research: Biology, Husbandry and Research Applications for Zebrafish, Medaka, Killifish, Cavefish, Stickleback, Goldfish and Danionella Translucida* 311–347. doi:10.1016/B978-0-12-821099-4.00003-1

Bilandžija H, Hollifield B, Steck M, Meng G, Ng M, Koch AD, Gračan R, Ćetković H, Porter ML, Renner KJ, Jeffery W. 2020. Phenotypic plasticity as a mechanism of cave colonization and adaptation. *Elife* 9. doi:10.7554/ELIFE.51830

Boggs TE, Friedman JS, Gross JB. 2022. Alterations to cavefish red blood cells provide evidence of adaptation to reduced subterranean oxygen. *Scientific Reports 2022 12:1* 12:1–10. doi:10.1038/s41598-022-07619-0

Brillon DJ, Zheng B, Campbell RG, Matthews DE. 1995. Effect of cortisol on energy expenditure and amino acid metabolism in humans. *https://doi.org/101152/ajpendo19952683E501*

Britton JR, Andreou D, Lopez-Bejar M, Carbajal A. 2023. Relationships of scale cortisol content suggest stress resilience in freshwater fish vulnerable to catch-and-release angling in recreational fisheries. *Fish Res* 266:106776. doi:10.1016/J.FISHRES.2023.106776

Brosset P, Averty A, Mathieu-Resuge M, Schull Q, Soudant P, Lebigre C. 2023. Fish morphometric body condition indices reflect energy reserves but other physiological processes matter. *Ecol Indic* 154:110860. doi:10.1016/J.ECOLIND.2023.110860

Carbajal A, Monclús L, Tallo-Parra O, Sabes-Alsina M, Vinyoles D, Lopez-Bejar M. 2018. Cortisol detection in fish scales by enzyme immunoassay: Biochemical and methodological validation. *Journal of Applied Ichthyology* 34:967–970. doi:10.1111/JAI.13674

Carbajal A, Tallo-Parra O, Monclús L, Vinyoles D, Solé M, Lacorte S, Lopez-Bejar M. 2019. Variation in scale cortisol concentrations of a wild freshwater fish: Habitat quality or seasonal influences? *Gen Comp Endocrinol* 275:44–50. doi:10.1016/J.YGCEN.2019.01.015

Caruso G, Maricchiolo G, Micale V, Genovese L, Caruso R, Denaro MG. 2010. Physiological responses to starvation in the European eel (Anguilla anguilla): Effects on haematological, biochemical, non-specific immune parameters and skin structures. *Fish Physiol Biochem* 36:71–83. doi:10.1007/S10695-008-9290-6

Chin JSR, Gassant CE, Amaral PM, Lloyd E, Stahl BA, Jaggard JB, Keene AC, Duboue ER. 2018. Convergence on reduced stress behavior in the Mexican blind cavefish. *Dev Biol* 441:319–327. doi:10.1016/J.YDBIO.2018.05.009

Choi CY, An KW. 2008. Cloning and expression of Na+/K+-ATPase and osmotic stress transcription factor 1 mRNA in black porgy, Acanthopagrus schlegeli during osmotic stress. *Comp Biochem Physiol B Biochem Mol Biol* 149:91–100. doi:10.1016/J.CBPB.2007.08.009

Christiansen JJ, Djurhuus CB, Gravholt CH, Iversen P, Christiansen JS, Schmitz O, Weeke J, Jørgensen JOL, Møller N. 2007. Effects of Cortisol on Carbohydrate, Lipid, and Protein Metabolism: Studies of Acute Cortisol Withdrawal in Adrenocortical Failure. *J Clin Endocrinol Metab* 92:3553–3559. doi:10.1210/JC.2007-0445

Cockrem JF, Bahry MA, Chowdhury VS. 2019. Cortisol responses of goldfish (Carassius auratus) to air exposure, chasing, and increased water temperature. *Gen Comp Endocrinol* 270:18–25. doi:10.1016/J.YGCEN.2018.09.017

De Marco RJ, Groneberg AH, Yeh CM, Castillo Ramírez LA, Ryu S. 2013. Optogenetic elevation of endogenous glucocorticoid level in larval zebrafish. *Front Neural Circuits* 7:46978. doi:10.3389/FNCIR.2013.00082/BIBTEX

DeKoning ABL, Picard DJ, Bond SR, Schulte PM. 2004. Stress and interpopulation variation in glycolytic enzyme activity and expression in a teleost fish Fundulus heteroclitus. *Physiological and Biochemical Zoology* 77:18–26. doi:10.1086/378914

Ding J, Fanjara E, Stene A, Gansel LC, Finstad B, Cao Y. 2024. Comparison and Validation of Commercial Elisa Kits for Quantification of Fecal Cortisol Metabolites in Atlantic Salmon (Salmon Salar). doi:10.2139/SSRN.5012212

Elipot Y, Hinaux H, Callebert J, Rétaux S. 2013. Evolutionary shift from fighting to foraging in blind cavefish through changes in the serotonin network. *Current Biology* 23:1–10. doi:10.1016/J.CUB.2012.10.044

Fischer EK, Harris RM, Hofmann HA, Hoke KL. 2014. Predator exposure alters stress physiology in guppies across timescales. *Horm Behav* 65:165–172. doi:10.1016/J.YHBEH.2013.12.010

Gallo ND, Jeffery WR. 2012. Evolution of Space Dependent Growth in the Teleost Astyanax mexicanus. *PLoS One* 7:e41443. doi:10.1371/JOURNAL.PONE.0041443

Garduño-Sánchez M, Hernández-Lozano J, Moran RL, Miranda-Gamboa R, Gross JB, Rohner N, Elliott WR, Miller J, Lozano-Vilano L, McGaugh SE, Ornelas-García CP. 2023. Phylogeographic relationships and morphological evolution between cave and surface Astyanax mexicanus populations (De Filippi 1853) (Actinopterygii, Characidae). *Mol Ecol* 32:5626–5644. doi:10.1111/MEC.17128

Gates RD, Baghdasarian G, Muscatine L. 1992. Temperature Stress Causes Host Cell Detachment in Symbiotic Cnidarians: Implications for Coral Bleaching. *https://doi.org/102307/1542252* 182:324–332. doi:10.2307/1542252

Huffman DL, Abrami L, Sasik R, Corbeil J, Van Der Goot FG, Aroian R V. 2004. Mitogen-activated protein kinase pathways defend against bacterial pore-forming toxins. *Proc Natl Acad Sci U S A* 101:10995–11000. doi:10.1073/PNAS.0404073101

Humphrey EA, Crespi E, Travis J. 2020. Under pressure: Short- and long-term response to predation varies in two populations of a live-bearing fish. *Ethology* 126:463–475. doi:10.1111/ETH.12996

Krishnan J, Persons JL, Peuß R, Hassan H, Kenzior A, Xiong S, Olsen L, Maldonado E, Kowalko JE, Rohner N. 2020. Comparative transcriptome analysis of wild and lab populations of Astyanax mexicanus uncovers differential effects of environment and morphotype on gene expression. *J Exp Zool B Mol Dev Evol* 334:530–539. doi:10.1002/JEZ.B.22933

Lappivaara J, Kiviniemi A, Oikari A. 1999. Bioaccumulation and subchronic physiological effects of waterborne iron overload on whitefish exposed in humic and nonhumic water. *Arch Environ Contam Toxicol* 37:196–204. doi:10.1007/S002449900506/METRICS

Lappivaara J, Marttinen S. 2005. Effects of waterborne iron overload and simulated winter conditions on acute physiological stress response of whitefish, Coregonus lavaretus. *Ecotoxicol Environ Saf* 60:157–168. doi:10.1016/J.ECOENV.2004.01.003

Lebigre C, Woillez M, Barone H, Mourot J, Drogou M, Le Goff R, Servili A, Hennebert J, Vanhomwegen M, Aerts J. 2022. Temporal variations in scale cortisol indicate consistent local-and broad-scale constraints in a wild marine teleost fish. *Mar Environ Res* 182:105783. doi:10.1016/J.MARENVRES.2022.105783

Legendre L, Père S, Rebaudo F, Espinasa L, Rétaux S. 2025. Water parameters and hydrodynamics in rivers and caves hosting Astyanax mexicanus populations reveal macro-, meso- and micro-habitat characteristics. *bioRxiv* 2025.02.06.635682. doi:10.1101/2025.02.06.635682

Léger JAD, Athanasio CG, Zhera A, Chauhan MF, Simmons DBD. 2021. Hypoxic responses in Oncorhynchus mykiss involve angiogenesis, lipid, and lactate metabolism, which may be triggered by the cortisol stress response and epigenetic methylation. *Comp Biochem Physiol Part D Genomics Proteomics* 39:100860. doi:10.1016/J.CBD.2021.100860

Mitchell R, Russell W, Elliott W. 1977. Mexican eyeless Characin fishes, genus Astyanax: environment, distribution, and evolution. *KIP Monographs*.

Moran RL, Richards EJ, Ornelas-García CP, Gross JB, Donny A, Wiese J, Keene AC, Kowalko JE, Rohner N, McGaugh SE. 2023. Selection-driven trait loss in independently evolved cavefish populations. *Nature Communications 2023 14:1* 14:1–19. doi:10.1038/s41467-023-37909-8

Mota VC, Martins CIM, Eding EH, Canário AVM, Verreth JAJ. 2017. Cortisol and testosterone accumulation in a low pH recirculating aquaculture system for rainbow trout (Oncorhynchus mykiss). *Aquac Res* 48:3579–3588. doi:10.1111/ARE.13184

O'Connor EA, Pottinger TG, Sneddon LU. 2011. The effects of acute and chronic hypoxia on cortisol, glucose and lactate concentrations in different populations of three-spined stickleback. *Fish Physiol Biochem* 37:461–469. doi:10.1007/S10695-010-9447-Y

O'Dwyer K, Dargent F, Forbes MR, Koprivnikar J. 2020. Parasite infection leads to widespread glucocorticoid hormone increases in vertebrate hosts: A meta-analysis. *Journal of Animal Ecology* 89:519–529. doi:10.1111/1365-2656.13123

Olsen L, Levy M, Medley JK, Hassan H, Miller B, Alexander R, Wilcock E, Yi K, Florens L, Weaver K, McKinney SA, Peuß R, Persons J, Kenzior A, Maldonado E, Delventhal K, Gluesenkamp A, Mager E, Coughlin D, Rohner N. 2023. Metabolic reprogramming

underlies cavefish muscular endurance despite loss of muscle mass and contractility. *Proc Natl Acad Sci U S A* 120:e2204427120. doi:10.1073/PNAS.2204427120

Ornelas-García P, Pajares S, Sosa-Jiménez VM, Rétaux S, Miranda-Gamboa RA. 2018. Microbiome differences between river-dwelling and cave-adapted populations of the fish astyanax mexicanus (De Filippi, 1853). *PeerJ* 2018:e5906. doi:10.7717/PEERJ.5906/SUPP-3

Padmanaban N, Ambosie R, Choy S, Marcus S, Nilsson SRO, Keene AC, Kowalko JE, Duboué ER. 2025. Automated behavioral profiling using neural networks reveals differences in stress-like behavior between cave and surface-dwelling Astyanax mexicanus. *bioRxiv* 2025.01.30.635725. doi:10.1101/2025.01.30.635725

Peuß R, Box AC, Chen S, Wang Y, Tsuchiya D, Persons JL, Kenzior A, Maldonado E, Krishnan J, Scharsack JP, Slaughter BD, Rohner N. 2020. Adaptation to low parasite abundance affects immune investment and immunopathological responses of cavefish. *Nat Ecol Evol*. doi:10.1038/s41559-020-1234-2

Pickering AD, Pottinger TG. 1989. Stress responses and disease resistance in salmonid fish: Effects of chronic elevation of plasma cortisol. *Fish Physiol Biochem* 7:253–258. doi:10.1007/BF00004714

Poulson TL. 2001. Adaptations of cave fishes with some comparisons to deep-sea fishes. *Environ Biol Fishes* 62:345–364. doi:10.1023/A:1011893916855/METRICS

Poulson TL, White WB. 1969. The cave environment. *Science (1979)* 165:971–981. doi:10.1126/SCIENCE.165.3897.971

Poursaeid S, Falahatkar B. 2022. Starvation alters growth, stress metabolites and physiological responses in juvenile great sturgeon (Huso huso). *Anim Feed Sci Technol* 294:115429. doi:10.1016/J.ANIFEEDSCI.2022.115429

Pravosudov V V., Kitaysky AS, Wingfield JC, Clayton NS. 2001. Long-Term Unpredictable Foraging Conditions and Physiological Stress Response in Mountain Chickadees (Poecile gambeli). *Gen Comp Endocrinol* 123:324–331. doi:10.1006/GCEN.2001.7684

Riddle MR, Aspiras AC, Gaudenz K, Peuß R, Sung JY, Martineau B, Peavey M, Box AC, Tabin JA, McGaugh S, Borowsky R, Tabin CJ, Rohner N. 2018. Insulin resistance in cavefish as an adaptation to a nutrient-limited environment. *Nature 2018 555:7698* 555:647–651. doi:10.1038/nature26136

Rodriguez-Morales R, Gonzalez-Lerma P, Yuiska A, Han JH, Guerra Y, Crisostomo L, Keene AC, Duboue ER, Kowalko JE. 2022. Convergence on reduced aggression through shared behavioral traits in multiple populations of Astyanax mexicanus. *BMC Ecol Evol* 22:1–15. doi:10.1186/S12862-022-02069-8

Rohner N, Jarosz DF, Kowalko JE, Yoshizawa M, Jeffery WR, Borowsky RL, Lindquist S, Tabin CJ. 2013. Cryptic variation in morphological evolution: HSP90 as a capacitor for loss of eyes in cavefish. *Science (1979)* 342:1372–1375. doi:10.1126/SCIENCE.1240276

Roque d'orbcastel E, Bettarel Y, Dellinger M, Sadoul B, Bouvier T, Amandé JM, Dagorn L, Geffroy B. 2021. Measuring cortisol in fish scales to study stress in wild tropical tuna. *Environ Biol Fishes* 104:725–732. doi:10.1007/S10641-021-01107-6

Sachser N, Lick C. 1991. Social experience, behavior, and stress in guinea pigs. *Physiol Behav* 50:83–90. doi:10.1016/0031-9384(91)90502-F

Santacruz A, Claudia |, Ornelas-García P, Pérez-Ponce De León G, Mérida U, Mérida M. 2020. Incipient genetic divergence or cryptic speciation? Procamallanus (Nematoda) in freshwater fishes (Astyanax). *Zool Scr* 49:768–778. doi:10.1111/ZSC.12443

Santacruz A, Hernández-Mena D, Miranda-Gamboa R, De León GPP, Ornelas-García CP. 2023. Host-parasite interactions in perpetual darkness: Macroparasite diversity in the

cavefish Astyanax mexicanus. *Zool Res* 44:782. doi:10.24272/J.ISSN.2095-8137.2022.376

Schulte PM, Davies SA, Dow JAT, Lukowiak K. 2014. What is environmental stress? Insights from fish living in a variable environment. *Journal of Experimental Biology* 217:23–34. doi:10.1242/JEB.089722

Servicio Meteorológico Nacional. n.d. https://smn.conagua.gob.mx/es/

Shimobayashi M, Thomas A, Shetty S, Frei IC, Wölnerhanssen BK, Weissenberger D, Vandekeere A, Planque M, Dietz N, Ritz D, Meyer-Gerspach AC, Maier T, Hay N, Peterli R, Fendt SM, Rohner N, Hall MN. 2023. Diet-induced loss of adipose hexokinase 2 correlates with hyperglycemia. *Elife* 12. doi:10.7554/ELIFE.85103

Simon V, Elleboode R, Mahé K, Legendre L, Ornelas-Garcia P, Espinasa L, Rétaux S. 2017. Comparing growth in surface and cave morphs of the species Astyanax mexicanus: insights from scales. *SpringerV Simon, R Elleboode, K Mahé, L Legendre, P Ornelas-Garcia, L Espinasa, S RétauxEvodevo, 2017•Springer* 8. doi:10.1186/s13227-017-0086-6

Sumpter JP, Le Bail PY, Pickering AD, Pottinger TG, Carragher JF. 1991. The effect of starvation on growth and plasma growth hormone concentrations of rainbow trout, Oncorhynchus mykiss. *Gen Comp Endocrinol* 83:94–102. doi:10.1016/0016-6480(91)90109-J

Tabin JA, Aspiras A, Martineau B, Riddle M, Kowalko J, Borowsky R, Rohner N, Tabin CJ. 2018. Temperature preference of cave and surface populations of Astyanax mexicanus. *Dev Biol* 441:338–344. doi:10.1016/J.YDBIO.2018.04.017

van der Weele CM, Jeffery WR. 2022. Cavefish cope with environmental hypoxia by developing more erythrocytes and overexpression of hypoxia-inducible genes. *Elife* 11. doi:10.7554/ELIFE.69109

Van Dievel M, Janssens L, Stoks R. 2016. Short- and long-term behavioural, physiological and stoichiometric responses to predation risk indicate chronic stress and compensatory mechanisms. *Oecologia* 181:347–357. doi:10.1007/S00442-015-3440-1

Van Raaij MTM, Pit DSS, Balm PHM, Steffens AB, Van Den Thillart GEEJM. 1996. Behavioral Strategy and the Physiological Stress Response in Rainbow Trout Exposed to Severe Hypoxia. *Horm Behav* 30:85–92. doi:10.1006/HBEH.1996.0012

Venarsky MP, Huntsman BM, Huryn AD, Benstead JP, Kuhajda BR. 2014. Quantitative food web analysis supports the energy-limitation hypothesis in cave stream ecosystems. *Oecologia* 176:859–869. doi:10.1007/S00442-014-3042-3/FIGURES/2

Vercauteren M, Ampe B, Devriese L, Moons CPH, Decostere A, Aerts J, Chiers K. 2022. Explorative study on scale cortisol accumulation in wild caught common dab (Limanda limanda). *BMC Vet Res* 18:1–12. doi:10.1186/S12917-022-03385-3

Vijayan MM, Pereira C, Grau EG, Iwama GK. 1997. Metabolic Responses Associated with Confinement Stress in Tilapia: The Role of Cortisol. *Comp Biochem Physiol C Pharmacol Toxicol Endocrinol* 116:89–95. doi:10.1016/S0742-8413(96)00124-7

Vindas MA, Magnhagen C, Brännäs E, Øverli Ø, Winberg S, Nilsson J, Backström T. 2017. Brain cortisol receptor expression differs in Arctic charr displaying opposite coping styles. *Physiol Behav* 177:161–168. doi:10.1016/J.PHYSBEH.2017.04.024

Wagner EJ, Bosakowski,' And T, Intelmann S. 1997. Combined Effects of Temperature and High pH on Mortality and the Stress Response of Rainbow Trout after Stocking; Combined Effects of Temperature and High pH on Mortality and the Stress Response of Rainbow Trout after Stocking. doi:10.1577/1548-8659(1997)126

Wang WN, Zhou J, Wang P, Tian TT, Zheng Y, Liu Y, Mai W jun, Wang AL. 2009. Oxidative stress, DNA damage and antioxidant enzyme gene expression in the Pacific white shrimp, Litopenaeus vannamei when exposed to acute pH stress. *Comparative*

Biochemistry and Physiology Part C: Toxicology & Pharmacology 150:428–435. doi:10.1016/J.CBPC.2009.06.010

Webster Marketon JI, Glaser R. 2008. Stress hormones and immune function. *Cell Immunol* 252:16–26. doi:10.1016/J.CELLIMM.2007.09.006

White A, Fletcher TC. 1986. Serum cortisol, glucose and lipids in plaice (Pleuronectes platessa L.) exposed to starvation and aquarium stress. *Comp Biochem Physiol A Physiol* 84:649–653. doi:10.1016/0300-9629(86)90380-4

Wilkens H, Hüppop K. 1986. Sympatric speciation in cave fishes? *Journal of Zoological Systematics and Evolutionary Research* 24:223–230. doi:10.1111/J.1439-0469.1986.TB00630.X

Winberg S, Lepage O. 1998. Elevation of brain 5-HT activity, POMC expression, and plasma cortisol in socially subordinate rainbow trout. *Am J Physiol Regul Integr Comp Physiol* 274. doi:10.1152/AJPREGU.1998.274.3.R645

Xiong S, Krishnan J, Peuß R, Rohner N. 2018. Early adipogenesis contributes to excess fat accumulation in cave populations of Astyanax mexicanus. *Dev Biol* 441:297–304. doi:10.1016/J.YDBIO.2018.06.003

Xiong S, Wang W, Kenzior A, Olsen L, Krishnan J, Persons J, Medley K, Peuß R, Wang Y, Chen S, Zhang N, Thomas N, Miles JM, Alvarado AS, Rohner N. 2022. Enhanced lipogenesis through Pparγ helps cavefish adapt to food scarcity. *Current Biology* 32:2272-2280.e6. doi:10.1016/J.CUB.2022.03.038

Yoshizawa M, Library WO. 2015. Behaviors of cavefish offer insight into developmental evolution. *Mol Reprod Dev* 82:268–280. doi:10.1002/MRD.22471

Yoshizawa M, Robinson BG, Duboué ER, Masek P, Jaggard JB, O'Quin KE, Borowsky RL, Jeffery WR, Keene AC. 2015. Distinct genetic architecture underlies the emergence of sleep loss and prey-seeking behavior in the Mexican cavefish. *BMC Biol* 13:1–12. doi:10.1186/S12915-015-0119-3

Ziv L, Muto A, Schoonheim PJ, Meijsing SH, Strasser D, Ingraham HA, Schaaf MJM, Yamamoto KR, Baier H. 2012. An affective disorder in zebrafish with mutation of the glucocorticoid receptor. *Molecular Psychiatry 2013 18:6* 18:681–691. doi:10.1038/mp.2012.64

Acknowledgements

We thank Michelle Borgers, Ulla Pebesma, Annika Potthoff, David Perez Guerra, Riley Kellermeyer and Nicolas Rohner for their help with fieldwork and fish dissections. We also acknowledge people from Ejido Los Sabinos and from the Comisión Nacional de Áreas Naturales Protegidas (CONANP) for their guidance in Los Sabinos and Tinaja caves. Furthermore, we thank Edith Ossendorf and Sabine Kruse for support on establishing and conducting the ELISA assay. Finally, we thank Jaime Anaya-Rojas for advice with statistical analysis. This work was funded by the German Research Foundation (DFG) as part of the CRC TRR 212 (NC3, 316099922), projects B06 (granted to R.P., Project number 471673683) and project S, Endocrinology platform (granted to S.K., Project number 396782989).

Author contributions

Marc Bauhus: investigation (lead); formal analysis (lead); project administration (lead); writing – original draft (lead); visualization (lead); conceptualization (support). **Luise Wander**: investigation (support); formal analysis (support); writing – review & editing (equal). **David Boy:** investigation (support); formal analysis (support); writing – review & editing (equal).

Ana Santacruz: investigation (support); writing – review & editing (equal). **Ernesto Maldonado:** investigation (support); writing – review & editing (equal). **Joachim Kurtz:** resources (support); supervision (support); writing – review & editing (equal). **Sylvia Kaiser:** methodology (lead); resources (support); funding acquisition (support); writing – review & editing (equal). **Robert Peuß:** conceptualization (lead); funding acquisition (lead); project administration (support); supervision (lead); resources (lead); visualization (support); writing – original draft (support); writing – review & editing (equal).

Competing interests

The authors declare no competing interests.

Ethics statement

All wild animals were collected under permit no. SPARN/DGVS/03371/23 granted by the Secretaría de Medio Ambiente y Recursos Naturales (SEMARNAT) to Ernesto Maldonado. All animals were treated in accordance with local veterinary and animal welfare authorities and their respective guidelines. Scales from laboratory fish were used under the permit T24.016UMS granted by the University of Münster, Germany in accordance with local veterinary and animal welfare authorities.