MANUSCRIPT TITLE

Stable interaction of NONO with DBHS family members upon etoposide-
induced DNA damage

AUTHORS

Barbara Trifault[1,2], Stephanie Lamer[3], Andreas Schlosser[3#], Kaspar Burger[1,2#]*

[1]Mildred Scheel Early Career Center for Cancer Research (Mildred-Scheel-
Nachwuchszentrum, MSNZ) Würzburg, University Hospital Würzburg, Josef-Schneider-Str. 2,
97080 Würzburg, Germany.

[2]Department of Biochemistry and Molecular Biology, Biocenter of the University of Würzburg,
Am Hubland, 97074 Würzburg, Germany.

[3]Rudolf-Virchow-Zentrum - Center for Integrative and Translational Bioimaging, University
of Würzburg, Josef-Schneider-Str. 2, 97080 Würzburg, Germany

[#]equal contribution

*To whom correspondence should be addressed: E-mail: kaspar.burger@uni-wuerzburg.de
(KB)

Short title

NONO interactome during DNA damage

ABSTRACT

The Non-POU domain containing octamer binding (NONO) protein is a member of the multifunctional *Drosophila* behavior/human splicing (DBHS) protein family and a core component of nuclear paraspeckles. NONO forms dimers with the other two DBHS members splicing factor proline and glutamine rich (SFPQ) protein or the paraspeckle component 1 (PSCP1) to modulate RNA metabolism and gene expression both at the transcriptional and post-transcriptional level. Increasing evidence suggests that NONO participates in genome maintenance by stimulating the DNA damage response (DDR) upon induction of DNA double-strand breaks (DSBs). However, the molecular principles that engage NONO in genome stability are poorly understood. We hypothesized that the induction of DSBs alters NONO protein-protein interactions and applied label-free mass spectrometry to test for changes in the interactome of NONO in human U2OS cells upon treatment with the topoisomerase II inhibitor etoposide. Surprisingly, our mass spectrometry data reveal that etoposide treatment does not induce major changes in NONO protein-protein interactions. We confirmed this finding by orthogonal co-immunoprecipitation assays and co-localization assays. Our data suggest that the bulk of interactions between NONO and SFPQ or PSPC1 are insensitive to etoposide treatment and that DBHS family members promote genome stability as stable dimers.

Keywords

DNA damage response; mass spectrometry; NONO; etoposide; DBHS proteins; nucleolus.

INTRODUCTION

RNA-binding proteins (RBPs) are an abundant protein family with critical roles in RNA metabolism. Utilizing a characteristic RNA recognition motif (RRM) and variable RNA-binding domains, RBPs recognize a diverse array of transcripts to participate in virtually all aspects of gene expression, ranging from the production and processing of nascent RNA in the nucleus to posttranscriptional roles in translation and RNA stability [1,2]. The Non-POU domain containing octamer binding (NONO) protein is a member of the *Drosophila* behavior/human splicing (DBHS) protein family and concentrates in paraspeckles. Paraspeckles are stress-responsive nuclear bodies that are built on the lncRNA NEAT1 and sequester 40 or so RBPs to form RNA metabolic hubs [3–7]. Members of the DBHS family are abundant and dynamic RBPs that mediate numerous protein-protein and protein-nucleic acid interactions to participate in various steps of pre-mRNA synthesis [8]. NONO self-dimerizes or forms heterodimers with the splicing factor proline and glutamine rich (SFPQ) protein or the paraspeckle component 1 (PSCP1) protein to promote the retention and editing of a subset of pre-mRNA transcripts within paraspeckles [9,10]. NONO also enriches on chromatin and associates with promoter-associated nascent transcripts to stimulate the formation of transcriptional condensates and augment RNA polymerase II activity [11–15]. First discovered as a splicing factor, NONO directly interacts with spliceosomal subunits and recognizes the 5' splice site, but is not essential for the splicing machinery in unperturbed cells [16,17]. Instead, NONO enhances pre-mRNA processing of super enhancer-associated transcripts upon oncogenic stress [18]. NONO is upregulated in many tumors and high levels of NONO foster the expression oncogenic transcriptional programs, which promotes proliferation and drug-resistance of cancer cells [19]. Thus, NONO is a multifunctional RNA metabolic factor.

Genome instability is a hallmark of cancer [20]. DNA double-strand breaks (DSBs) are highly toxic and threat genome stability [21,22]. To counteract genotoxic stress, the DNA

damage response (DDR) engages a myriad of proteins that sense and repair DSBs, including

RBPs [23–25]. NONO and other DBHS family members are also linked to DDR and modulates

the recognition and repair of DSBs [26]. Early studies suggest that NONO heterodimerizes with

SFPQ to promote DSB pathway choice [27]. The depletion of NONO indeed delays the

clearance of DSB foci and increases chromosomal aberrations [28,29]. NONO directly

accumulates at DSBs and interacts with a subset of DDR factors, which fosters DSB repair via

non-homologous end-joining (NHEJ) [30,31]. The accumulation of NONO at DSBs occurs in

a poly(ADP-ribose)-dependent manner and involves the activation of the DSB-sensing DNA-

dependent protein kinase (DNA-PK) [28,32]. More recently, novel mechanisms for the genome

protective role of NONO have been reported, which include the relocalization of NONO and

other DBHS family members to the nucleolus and nucleolus-associated condensates via

interaction with nucleolar transcripts [33,34]. These findings underscore the relevance of

NONO in promoting efficient DSB repair and genome stability.

Interestingly, NONO binding affinity to RNA substrates is regulated by the coactivator

associated arginine methyltransferase 1 (CARM1/PRMT4) [35]. Moreover, NONO undergoes

glycosylation and ubiquitinylation to efficiently accumulate at DSBs and UV-induced lesions,

respectively [36,37]. However, to what extent DNA damage impacts on NONO protein-protein

interactions is poorly understood. We hypothesized that the induction of DSBs alters the

interactions of NONO with other DBHS family proteins and employed label-free quantitation

by mass spectrometry to determine NONO protein-protein interactions in human U2OS cells in

the absence or presence of the topoisomerase II inhibitor etoposide. Surprisingly, we observed

no clear changes in the NONO interactome and persistent association of both SFPQ and PSPC1

to NONO. Persistent interaction of NONO with DBHS family members in response to

etoposide treatment was validated by orthogonal assays and suggests remarkable stability of

the bulk of *bona fide* NONO protein-protein interactions upon induction of DSB signaling.

MATERIALS AND METHODS

Tissue culture, transfections, inhibitors and antibodies

Human U2OS osteosarcoma cells were acquired from the German collection of microorganisms and cell cultures (DSMZ), cultured at 37°C and 5 % CO_2 in Dulbecco's modified eagle's medium (DMEM, Gibco) complemented with 10 % fetal bovine serum (FBS, Capricorn), 100 U/mL penicillin-streptomycin (Gibco) and 2 mM L-glutamine (Gibco), and routinely tested for mycoplasma infection. For ectopic expression, cells were transfected with 1.5 µg plasmid DNA encoding HA-tagged NONO (pcDNA3.1-HA-NONO, kind gift from Nicolas Manel). For knockdown experiments, cells were transfected with 100 nM siRNA that selectively targets either NONO (siNONO, Dharmacon, L-007756-01 smart pool) or represents non-targeting control (siControl, Dharmacon, D-001810-01 smart pool). Lipofectamine 2000 (Invitrogen) and Opti-MEM (Gibco) was used for transfections according to the manufacturer´s protocol. Incubation conditions with etoposide (Sigma) were for 2 h with 20 µM, unless stated differently. The following antibodies were used in this study. Primary antibodies were: IgG control (Proteintech, 30000-0-AP); NONO (Proteintech, 11058-1-AP); PSPC1 (Proteintech, 16714-1-AP); SFPQ (Abcam, ab177149); Ki-67 (Thermo, MA5-14520); HA-tag (Abcam, ab9110; BioLegend, 901502); Fibrillarin (Abcam, ab5821); Vinculin (Sigma, V9131); histone H2B (Abcam, ab1790); phospho-ser-139 histone H2A.X (Cell Signaling, 2577; Millipore, 05-636); 53BP1 (Novus, NB100-304). Secondary antibodies were: HRP-linked IgG (Cytiva, GEHENA934; Cytiva, NA931); Alexa Fluor 546-linked IgG (Thermo, A10036), Alexa Fluor 488-linked IgG (Thermo, A21206), Alexa Fluor 546-linked IgG (Thermo, A10040).

Immunoblotting and immunoprecipitation

Cells were lysed, boiled (5 min, 95°C) and sonicated (Branson, 10 %, 5 sec) in 4x sample buffer

(250 mM tris-HCl pH 6.8, 8 % sodium dodecyl sulphate (SDS), 40 % glycerol, 0.8 % β-

mercaptoethanol, 0.02 % bromophenol blue). Whole cell extracts were separated by

denaturating SDS polyacrylamide gel electrophoresis (PAGE) and transferred onto a

nitrocellulose membrane (Cytiva). Membranes were blocked (30 min, RT) in PBST (PBS/0.1

% triton x-100) with 5 % milk, probed with primary (overnight, 4°C) and secondary antibodies

(1 h, RT) diluted in PBST with or without milk and washed in between incubation steps with

PBST. Signals were visualized with an ECL kit (Cytiva) with an imaging station (Fusion FX

Vilber, LAS-4000 Fuji) and quantified with ImageJ (NIH). For immunoprecipitation, cells were

trypsinised, resuspended in DMEM, centrifuged (1200 rpm, 3 min), washed in cold PBS and

centrifuged again. The pellet was lysed for 10 min on ice in 5 volumes of IP buffer (200 mM

NaCl, 0.5 mM EDTA, 20 mM HEPES, 0.2 % NP-40, 10 % glycerol, 1x protease/phosphatase

inhibitor (Roche), 100 U of Ribolock RNase inhibitor (Thermo). Lysates were centrifuged

(12000 rpm, 12 min) and supernatants were collected in a new tube. 10 % of lysate was kept as

input (IN) and the remaining 90 % was incubated (3 h, 4°C) on a rotative wheel with 5 μg

primary antibody pre-conjugated with 30 μL G beads overnight. Crosslinking of pre-conjugated

antibodies was performed with bis(sulfosuccinimidyl)suberate (BS3, Thermo) or dimethyl

pimelimidate (DMP, Thermo) according to the manufacturer's protocol. Immunocomplexes

were captured on a magnet (Invitrogen), washed three times with 800 μL IP buffer and eluted

(5 min, 95°C) with 40 μL of 4x sample buffer (reducing condition). For native PAGE, samples

were eluted in 4x sample buffer without β-mercaptoethanol (non-reducing condition). Proteins

were visualized by using either a SilverQuest staining kit (Thermo) according to the

manufacturer´s protocol prior to transfer or by staining for 5 min at RT with Ponceau solution

(0.5 % Ponceau S/1 % acetic acid) after the transfer.

Confocal imaging

Cells grown on cover slips (Roth) were washed in PBS, fixed (10 min, RT) with 3 % paraformaldehyde (Sigma), permeabilized (10 min, RT) with PBST, washed (5 min, three times) in PBS and blocked (2 h, 4°C) with PBS/10 % FBS. For staining, cells were incubated in a humidified chamber with primary antibodies (overnight at 4°C) and secondary antibodies (2 h, RT) diluted in PBS/0.15 % FBS and secondary antibodies. Cells were washed with PBST (three times, 5 min) between incubations and sealed in 4,6-diamidino-2-phenylindole (DAPI)-containing mounting medium (Vectashield). Signals were imaged with a confocal microscope (CLSM-Leica-SP2, x63 objective, 1024x1024 resolution, airy = 1) and channels were acquired sequentially, between frames, with equal exposure times as single scans. Images were analyzed and co-localization was assessed by ImageJ and RGB profiler (NIH).

Nanoscale liquid chromatography coupled to tandem mass spectrometry (Nano LC-MS/MS)

For in-gel digestion, immunoselected proteins were eluted with 4x lithium dodecyl sulphate (LDS) sample buffer (NuPAGE, Thermo) according to the manufacturer's protocol and precipitated in a four-fold volume of acetone (overnight, -20°C). Precipitated proteins were washed three times with acetone at -20°C, dissolved in 4x LDS sample buffer, reduced (10 min, 70°C) with 50 mM DTT and alkylated (20 min, RT) with 120 mM iodoacetamide. Proteins were separated by Novex 4-12 % bis-tris gels with MOPS buffer (NuPAGE, Thermo) according to the manufacturer's protocol. Gels were washed (5 min, three times) with ultrapure water, stained (60 min, RT) with Coomassie (Thermo) and washed again for 1 h. All gel lanes were cut in 15 slices, destained with 30 % acetonitrile in 0.1 M NH_4HCO_3 pH 8.0, shrunk with 100 % acetonitrile, and dried in a vacuum concentrator. Each gel band was digested (overnight, 37°C) with 0.1 µg trypsin in 0.1 M NH_4HCO_3 pH 8.0, the supernatant was removed and the

peptides were extracted from the gel slices with 5 % formic acid. The extracted peptides were

pooled with the supernatant. For NanoLC-MS/MS analysis, an Orbitrap Fusion (Thermo)

equipped with a PicoView Ion Source and coupled to an EASY-nLC 1000 was used. Peptides

were loaded on capillary columns (PicoFrit, 30 cm x 150 μm ID, New Objective) self-packed

with ReproSil-Pur 120 C18-AQ, 1.9 μm (Dr. Maisch) and separated with a linear gradient from

3 % to 30 % acetonitrile and 0.1 % formic acid (30 min, flow rate 500 nL/min). Scans were

acquired with a resolution of 7500 for MS/MS scans and 60000 for MS scans in the Orbitrap

analyzer. HCD fragmentation with 35 % normalized collision energy was applied and a top

speed data-dependent MS/MS method with a fixed cycle time of 3 sec was used. Dynamic

exclusion was set with a repeat count of 1 and an 30sec exclusion duration and singly charged

precursors were excluded from selection. Minimum signal threshold for precursor selection was

set to 5x104 and predictive AGC was used with a AGC target value of 2x105 for MS scans and

5x104 for MS/MS scans. EASY-IC was used for internal calibration.

Mass spectrometry data analysis

Raw MS data files were analyzed with MaxQuant version 1.6.2.2 [38]. Database search was

performed with Andromeda, which is integrated in the utilized version of MaxQuant. The

search was performed against the UniProt Human Reference Proteome database (January 2021,

UP000005640, 75777 entries). Additionally, a database containing common contaminants was

used. The search was performed with tryptic cleavage specificity with 3 allowed miscleavages.

Protein identification was monitored by setting a false-discovery rate (FDR) of <1 % on protein

and peptide spectrum match level. In addition to MaxQuant default settings, the search was

performed against the following variable modifications: Protein N-terminal acetylation, Gln to

pyro-Glu formation (N-term. Gln) and oxidation (Met). Carbamidomethyl (Cys) was set as

fixed modification. Further data analysis was performed using R scripts developed in-house.

LFQ intensities were used for protein quantitation [39]. Proteins with less than two razor/unique

peptides were removed. Missing LFQ intensities were imputed with values close to the baseline.

Data imputation was performed with values from a standard normal distribution with a mean

of the 5 % quantile of the combined log10-transformed LFQ intensities and a standard deviation

of 0.1. For the identification of significantly enriched proteins, median log2 transformed protein

ratios were calculated from the two replicate experiments and boxplot outliers were identified

in intensity bins of at least 300 proteins. Log2 transformed protein ratios of sample versus

control with values outside a 1.5x (significance 1) or 3x (significance 2) interquartile range

(IQR), respectively, were considered as significantly enriched in the individual replicates.

Ethics statement

All data are derived from the U2OS cancer cell line (non-pathogenic, non-infectious, safety

level S1), which has been isolated in 1964. U2OS cells are an established cancer model cell line

for more than 50 years and working with U2OS cells does not require ethical approval.

RESULTS

Validation of NONO antibody from mass spectrometry

We wished to study the impact of DNA damage on the NONO interactome in U2OS cells. We

acquired a knockout-validated polyclonal NONO antibody and confirmed its selectivity upon

RNAi-mediated depletion of NONO. Using the NONO antibody for immunoblotting and

immunofluorescence microscopy, we observed prominent reduction in reactivity upon

transfection of NONO-selective siRNA (Fig. S1A-C). Next, we validated the induction of DNA

damage by confocal imaging of DSB markers in response to etoposide treatment. As expected,

we observed prominent formation of nuclear foci in the presence of etoposide that stained

positive for the *bona fide* DSB marker p53-binding protein 1 (53BP1) and partially overlapped

with signals for the DNA damage marker serine-139 phosphorylated histone H2A.X variant

(γH2A.X) (Fig. S1D and E). We also tested the capacity of the antibody to immunoselect

NONO and performed immunoprecipitation experiments (Fig. S1F). Indeed, the antibody

efficiently enriched NONO from U2OS whole cell lysates, irrespective of etoposide treatment

conditions. Thus, we established conditions to detect and immunoselect NONO in the absence

and presence of DSBs.

We noticed prominent immunoglobulins signals in our immunoprecipitation samples,

suggesting prominent coelution of bead-conjugated immunoglobulins. Excessive amounts of

immunoglobulins can perturb mass spectrometry data and prevent the detection of less abundant

peptides. Thus, aimed to covalently bind the antibody to the matrix and prevent coelution. We

elaborated on our immunoprecipitation protocol and introduced a crosslinking step using either

BS3 or DMP reagent (Fig. S1G). Using non-reducing conditions, we found that both BS3 and

DMP efficiently prevented coelution of immunoglobulins from the beads. However, the NONO

antibody only tolerated BS3, but not DMP crosslinking and the latter prevented the efficient

immunoselection of NONO from lysates. We repeated the immunoprecipitation experiment and

assessed NONO levels after either non-reducing or reducing separation conditions (Fig. S1H).

Immunoblotting for NONO confirmed that BS3 crosslinking both preserves the affinity of the

antibody toward NONO and prevents coelution of immunoglobulins from the beads. We

conclude that etoposide treatment and antibody crosslinking by BS3 are appropriate to assess

changes in the NONO interactome upon DNA damage.

Persistent interaction of NONO with DBHS family members upon DNA damage

Having established conditions to immunoselect NONO, we applied mass spectrometry to

determine the impact of DNA damage on NONO protein-protein interactions. For comparative

analysis of two biological replicates, we incubated cells in the absence or presence of DNA

damage and performed immunoprecipitation from whole cell lysates using either the NONO

antibody or an isotype control (Fig. S2A). Correlation analysis revealed a high similarity

between replicates, irrespective of the antibody used for immunoprecipitation or etoposide

treatment (Fig. S2B). Overall, we detected 1031 proteins that were reproducibly present upon

NONO immunoprecipitation, albeit many of them with low peptide counts or poor sequence

coverage. To increase stringency, we considered only those as hits that were detected by at least

five unique peptides in the NONO immunoprecipitation samples. Such filtering narrowed down

the list of proteins to 238 that were immunoselected by the NONO antibody in the absence or

presence of etoposide treatment (Fig. 1A). Reassuringly, NONO was found as top hit together

with SFPQ and PSPC1, and all three DBHS family members scored with high significance

(Table S1). However, visual inspection of mass spectrometry data suggested an overall high

similarity between control and DNA damage conditions. Indeed, a stable, etoposide-resistant

interaction between NONO and DBHS family members was confirmed by immunoblotting

(Fig. 1B and C). Also, 76 of the top100 mass spectrometry hits were found in both conditions

(Fig. 1D; Table S2). To directly assess for differential co-enrichment of proteins in the absence

or presence of DNA damage, we integrated both treatment conditions into one graph (Fig. 2A).

Linear regression analysis revealed that PSPC1, SFPQ and virtually all other hits were co-

immunoprecipitated with NONO irrespective of etoposide treatment. We considered a second

filter and scored only those factors as significant that were at least 4-fold enriched over the

isotype control. This analysis yielded 49 proteins, out of which 28 were reported as nuclear

proteins. With a few exceptions, the list of top30 proteins is strikingly similar and the bulk of

filtered hits in the control condition are also found upon etoposide treatment (Fig. 2B). Of note,

90% or so of hits scoring as nuclear proteins were RBPs (Fig. S2C) and these included several

previously reported components of paraspeckles and NEAT1 interactors, such as HNRNPUL1,

YBX1 or SYNCRIP [40,41]. We conclude that our NONO interactome data are valid, but may

suffer from a considerable number of false-positive hits and cytoplasmic contaminants.

Surprisingly, the comparative analysis of NONO interactomes revealed no significant changes

in response to DNA damage.

Co-localization of DBHS family members *in vivo* and concomitant nucleolar re-

localization upon DNA damage

We could previously show that NONO, SFPQ and PSPC1 undergo nucleolar re-localization

upon etoposide treatment [34,42]. To corroborate our biochemical data and further assess the

association of NONO with DBHS family members in response to DNA damage *in vivo*, we

investigated the subcellular localization of DBHS proteins in the absence and presence of

etoposide by confocal imaging. We aimed to co-stain endogenous SFPQ or PSPC1 with

endogenous NONO using the same NONO antibody as used for biochemical assays. To the

best of our knowledge, however, there are currently no suitable and validated SFPQ or PSPC1

antibodies available that would allow co-staining with this NONO antibody. Thus, we

ectopically expressed full length, HA-tagged NONO (HA-NONO), in U2OS. As expected, HA-

NONO localized to the nucleoplasm in unperturbed cells and displayed a more pan-nuclear

localization upon etoposide treatment (Fig. S3A). A fraction of HA-NONO also accumulated

in the nucleolus, as monitored by partial co-localization with the nucleolar marker fibrillarin

(Fig. S3B). Importantly, treatment with etoposide did not alter the expression levels of HA-

NONO (Fig. S3C). Next, we co-stained endogenous DBHS proteins with HA-NONO.

Reassuringly, HA antibody reactivity displayed strong co-localization with each of the three

DBHS family members irrespective of etoposide treatment (Fig. 3A and B). Moreover, pan-

nuclear localization could be observed for a fraction of both HA-NONO, and each of the three

endogenous DBHS proteins in response to etoposide treatment. We conclude that NONO

associates with SFPQ and PSCP1 *in vivo* and that the bulk of these associations is resistant to

etoposide treatment.

DISCUSSION

Our study compared NONO protein interactions in the absence or presence of DNA damage.

Our mass spectrometry data confirmed the interaction of NONO with DBHS family member,

which was further validated by co-immunoprecipitation assays and co-localization studies. The

interactome data further indicate that NONO associates with numerous other RBPs. However,

our approach did not reveal any clear change in NONO interactions upon induction of DSBs.

Albeit confirming stable, etoposide-resistant formation of dimers between NONO and other

DBHS family members, the interpretation of DNA damage-induced changes in NONO

interactions may suffer from limitations.

NONO shares more than 70 % of sequence homology with SFPQ and PSPC1. The

conserved DBHS region of NONO contains two RNA recognition motifs (RRM1 and RRM2),

a NONA/paraspeckle domain (NOPS), and a coiled-coil domain [8]. NONO also contains an

amino-terminal histidine proline glutamine (HPQ)-rich region and a carboxy-terminal

glutamine proline (GP)-rich region, which harbors a nuclear localization sequence (NLS). The

canonical RRM1 and non-canonical RRM2 facilitate the binding of NONO to various nucleic

acid substrates, whereas the dimerization of DBHS proteins relies on the interaction between

the RRM2 domain of one paralog with the NOPS and coiled-coil domains of the other paralog

[8,43]. Intriguingly, phosphoproteomic data suggest that the induction of DNA damage by

ionizing radiation alters phosphorylation at NONO at threonine residues T428 and T450, which

are located in the carboxy-terminal region of NONO [44]. It is tempting to speculate that NONO

also undergoes differential phosphorylation upon etoposide treatment and that DNA damage-

responsive post-translational modifications alter NONO protein-protein interactions. However,

NONO is a multifunctional RBPs and modulates numerous processes in the nucleus. Thus, the

fraction of NONO susceptible to DNA damage may be rather small and its analysis likely

requires the generation of more selective tools such as phospho-specific antibodies. Our

imaging data using a pan-NONO antibody indeed indicate that a minor fraction of the nuclear

NONO pool is responsive to etoposide treatment, at least for the conditions used in this study.

Typically, factors involved in the recognition and repair of DSBs accumulate at such lesions

and form foci that can be visualized by microscopy. However, we did not observe the formation

of NONO-positive foci. A selective phospho-specific NONO antibody may not only reveal the

formation of NONO-positive DSB foci, but also facilitate a refined analysis of the NONO

protein-protein interactome in response to DNA damage. The data set could further benefit from

improvements in the mass spectrometry protocol, such as immunoprecipation from subcellular

fractions and the usage of orthogonal assays to study interactomes, such as proximity labelling

approaches. The latter allows profiling of proteins that localize in distinct subcellular

compartments including the nucleolus [45]. This may define changes in NONO protein-protein

interactions in the future.

Our data indicate that a subset of NONO accumulates in the nucleolus upon etoposide

treatment. However, NONO relocalization is not quantitative, and therefore potential changes

in NONO interactions upon DNA damage are difficult to resolve by mass spectrometry.

Interestingly, a more quantitative relocalization of NONO and other DBHS to the nucleolus has

been observed upon ionizing radiation [46]. Thus, DNA damage-induced changes in the NONO

interactome may occur concomitant with relocalization, for instance upon harsher treatment

conditions that induce more persistent DSBs and likely trigger a more comprehensive

engagement of NONO in DDR.

Overall, our study provides insights in the interactome of NONO in response to

etoposide treatment and suggests remarkable stability of the bulk of NONO protein-protein

interactions upon induction of DNA damage. It will be important to determine if the genome-

protective role of NONO occurs mostly in association with other DBHS family members or if

distinct types of DNA damage selectively engage individual DBHS protein family members in

DDR.

DATA AVAILABILITY

Proteomics data are available at the ProteomeXchange consortium via the PRIDE partner

repository [47] with the identifier PXD061274 (reviewer token: aFmB1YlDmtL1) or upon

login to the PRIDE website via reviewer account (username: reviewer_pxd061274@ebi.ac.uk,

password: 3Uo7pqlSPyJP).

CONFLICT OF INTEREST

The authors declare no conflict of interest.

SUPPORTING INFORMATION

Figure S1. Validation of NONO antibody selectivity in U2OS cells. (A, B) Immunoblot

detecting NONO in whole cell extracts (A) and confocal imaging of NONO (B) upon transient

transfection of siRNA targeting NONO (siNONO) or non-targeting control siRNA (siControl).

Vinculin, loading control; broken circle, nucleus. Representative images are shown. (C)

Quantitation of (B) using RGB profiler line scan analysis. a.u., arbitrary units. (D) Confocal

imaging of 53BP1 and γH2A.X foci. Representative images are shown. (E) Quantitation of (D)

using RGB profiler line scan analysis. (F) Immunoblot detecting NONO upon

immunoprecipitation (IP) with NONO antibody from whole cell lysates (IN, input). IgG,

isotype control antibody. (G) Immunoblot detecting NONO upon immunoprecipitation (IP)

with NONO antibody from whole cell lysates and native PAGE. Silver stain, loading control;

exp., exposure. Full blots are shown. (H) as in (G) but comparing NONO signals upon native

PAGE (left) and reducing PAGE (right). Full blots are shown. Control, non-crosslinked

condition, BS3, bis(sulfosuccinimidyl)suberate; DMP, dimethyl pimelimidate.

Figure S2. Quality control for NONO mass spectrometry in U2OS cells. (A) Coomassie stain

upon immunoprecipitation (IP) with NONO antibody from whole cell lysates. IgG, isotype

control antibody. (B) Correlation analysis of mass spectrometry replicates. Both the x-axis and

the y-axis display intensities of peptides. r, correlation coefficient. (C) Stratification of top30

mass spectrometry hits.

Figure S3. Quality control for ectopic expression of HA-tagged NONO in U2OS cells. (A)

Confocal imaging of HA-NONO upon transient transfection of pcDNA3.1-HA-NONO

expression plasmid. FBL, Fibrillarin, nucleolar marker. Representative images are shown. (B)

Quantitation of (A) using RGB profiler line scan analysis. a.u., arbitrary units. (C) Immunoblot

detecting HA-NONO upon transient transfection of pcDNA3.1-HA-NONO expression plasmid

in whole cell extracts. Vinculin, loading control; mock, non-transfected control.

Figure S4. Raw data. Uncropped immunoblots are shown. Red box, cropped area depicted in

figure panels.

Table S1. Mass spectrometry data.

Table S2. List of top100 mass spectrometry hits.

ACKNOWLEDGMENTS

We thank all members of the department for support. We acknowledge Lea Boten and Teresa

Klugt for excellent technical assistance. This work was funded by a grant from the German

Cancer Aid (grant code: 8606100-NG1) awarded to KB.

Author contributions: Conceptualization, BT, AS and KB; Methodology, BT, SL, AS and

KB; Investigation, BT and KB; Formal Analysis, BT, SL, AS and KB; Writing – Original Draft,

BT and KB; Writing – Review & Editing, AS and KB; Funding Acquisition, KB; Supervision,

AS and KB.

REFERENCES

1. Gebauer F, Schwarzl T, Valcárcel J, Hentze MW. RNA-binding proteins in human genetic
disease. Nat Rev Genet. 2021 Mar;22(3):185-198.

2. Corley M, Burns MC, Yeo GW. How RNA-Binding Proteins Interact with RNA:
Molecules and Mechanisms. Mol Cell. 2020 Apr 2;78(1):9-29.

3. Fox AH, Nakagawa S, Hirose T, Bond CS. Paraspeckles: Where Long Noncoding RNA
Meets Phase Separation. Trends Biochem Sci. 2018 Feb;43(2):124-135.

4. Bond CS, Fox AH. Paraspeckles: nuclear bodies built on long noncoding RNA. J Cell Biol.
2009 Sep 7;186(5):637-44.

5. Clemson CM, Hutchinson JN, Sara SA, Ensminger AW, Fox AH, Chess A, et al. An
architectural role for a nuclear noncoding RNA: NEAT1 RNA is essential for the structure
of paraspeckles. Mol Cell. 2009 Mar 27;33(6):717-26.

6. Hirose T, Virnicchi G, Tanigawa A, Naganuma T, Li R, Kimura H, et al. NEAT1 long
noncoding RNA regulates transcription via protein sequestration within subnuclear bodies.
Mol Biol Cell. 2014 Jan;25(1):169-83.

7. Sasaki YT, Ideue T, Sano M, Mituyama T, Hirose T. MENepsilon/beta noncoding RNAs
are essential for structural integrity of nuclear paraspeckles. Proc Natl Acad Sci U S A.
2009 Feb 24;106(8):2525-30.

8. Knott GJ, Bond CS, Fox AH. The DBHS proteins SFPQ, NONO and PSPC1: a
multipurpose molecular scaffold. Nucleic Acids Res. 2016 May 19;44(9):3989-4004.

9. Zhang Z, Carmichael GG. The fate of dsRNA in the nucleus: a p54(nrb)-containing
complex mediates the nuclear retention of promiscuously A-to-I edited RNAs. Cell. 2001
Aug 24;106(4):465-75.

10.   Prasanth KV, Prasanth SG, Xuan Z, Hearn S, Freier SM, Bennett CF, et al. Regulating
gene expression through RNA nuclear retention. Cell. 2005 Oct 21;123(2):249-63.

11. Lewis BA, Das SK, Jha RK, Levens D. Self-assembly of promoter DNA and RNA Pol
II machinery into transcriptionally active biomolecular condensates. Sci Adv. 2023 Oct
20;9(42):eadi4565.

12. Wei Y, Luo H, Yee PP, Zhang L, Liu Z, Zheng H, et al. Paraspeckle Protein NONO
Promotes TAZ Phase Separation in the Nucleus to Drive the Oncogenic Transcriptional
Program. Adv Sci (Weinh). 2021 Dec;8(24):e2102653.

13. Emili A, Shales M, McCracken S, Xie W, Tucker PW, Kobayashi R, et al. Splicing
and transcription-associated proteins PSF and p54nrb/nonO bind to the RNA polymerase II
CTD. RNA. 2002 Sep;8(9):1102-11.

14. Amelio AL, Miraglia LJ, Conkright JJ, Mercer BA, Batalov S, Cavett V, et al. A
coactivator trap identifies NONO (p54nrb) as a component of the cAMP-signaling
pathway. Proc Natl Acad Sci U S A. 2007 Dec 18;104(51):20314-9.

15. Mamontova V, Trifault B, Burger K. Nono induces Gadd45b to mediate DNA repair.
Life Sci Alliance. 2024 Jun 6;7(8):e202302555.

16. Kameoka S, Duque P, Konarska MM. p54(nrb) associates with the 5' splice site within
large transcription/splicing complexes. EMBO J. 2004 Apr 21;23(8):1782-91.

17. Peng R, Hawkins I, Link AJ, Patton JG. The splicing factor PSF is part of a large
complex that assembles in the absence of pre-mRNA and contains all five snRNPs. RNA
Biol. 2006 Apr;3(2):69-76.

18. Zhang S, Cooper JA, Chong YS, Naveed A, Mayoh C, Jayatilleke N, et al. NONO
enhances mRNA processing of super-enhancer-associated GATA2 and HAND2 genes in
neuroblastoma. EMBO Rep. 2022 Nov 23:e54977.

19. Ronchetti D, Traini V, Silvestris I, Fabbiano G, Passamonti F, Bolli N, et al. The
pleiotropic nature of NONO, a master regulator of essential biological pathways in cancers.
Cancer Gene Ther. 2024 Jul;31(7):984-994.

20. Hanahan D. Hallmarks of Cancer: New Dimensions. Cancer Discov. 2022
Jan;12(1):31-46.

21. Jackson SP, Bartek J. The DNA-damage response in human biology and disease.
Nature. 2009 Oct 22;461(7267):1071-8.

22. Ciccia A, Elledge SJ. The DNA damage response: making it safe to play with knives.
Mol Cell. 2010 Oct 22;40(2):179-204.

23. Blackford AN, Jackson SP. ATM, ATR, and DNA-PK: The Trinity at the Heart of the
DNA Damage Response. Mol Cell. 2017 Jun 15;66(6):801-817.

24. Zong D, Oberdoerffer P, Batista PJ, Nussenzweig A. RNA: a double-edged sword in
genome maintenance. Nat Rev Genet. 2020 Nov;21(11):651-670.

25. Burger K, Ketley RF, Gullerova M. Beyond the Trinity of ATM, ATR, and DNA-PK:
Multiple Kinases Shape the DNA Damage Response in Concert With RNA Metabolism.
Front Mol Biosci. 2019 Aug 2;6:61.

26.   Feng P, Li L, Deng T, Liu Y, Ling N, Qiu S, et al. NONO and tumorigenesis: More
than splicing. J Cell Mol Med. 2020 Apr;24(8):4368-4376.

27.   Bladen CL, Udayakumar D, Takeda Y, Dynan WS. Identification of the
polypyrimidine tract binding protein-associated splicing factor.p54(nrb) complex as a
candidate DNA double-strand break rejoining factor. J Biol Chem. 2005 Feb
18;280(7):5205-10.

28.   Krietsch J, Caron MC, Gagné JP, Ethier C, Vignard J, Vincent M, et al. PARP
activation regulates the RNA-binding protein NONO in the DNA damage response to
DNA double-strand breaks. Nucleic Acids Res. 2012 Nov 1;40(20):10287-301.

29.   Li S, Kuhne WW, Kulharya A, Hudson FZ, Ha K, Cao Z, et al. Involvement of
p54(nrb), a PSF partner protein, in DNA double-strand break repair and radioresistance.
Nucleic Acids Res. 2009 Nov;37(20):6746-53.

30.   Kuhnert A, Schmidt U, Monajembashi S, Franke C, Schlott B, Grosse F, et al.
Proteomic identification of PSF and p54(nrb) as TopBP1-interacting proteins. J Cell
Biochem. 2012 May;113(5):1744-53.

31.   Salton M, Lerenthal Y, Wang SY, Chen DJ, Shiloh Y. Involvement of Matrin 3 and
SFPQ/NONO in the DNA damage response. Cell Cycle. 2010 Apr 15;9(8):1568-76.

32.   Fan XJ, Wang YL, Zhao WW, Bai SM, Ma Y, Yin XK, et al. NONO phase separation
enhances DNA damage repair by accelerating nuclear EGFR-induced DNA-PK activation.
Am J Cancer Res. 2021 Jun 15;11(6):2838-2852.

33.   Yasuhara T, Xing YH, Bauer NC, Lee L, Dong R, Yadav T, et al. Condensates
induced by transcription inhibition localize active chromatin to nucleoli. Mol Cell. 2022
Aug 4;82(15):2738-2753.e6.

34.   Trifault B, Mamontova V, Cossa G, Ganskih S, Wei Y, Hofstetter J, et al. Nucleolar
detention of NONO shields DNA double-strand breaks from aberrant transcripts. Nucleic
Acids Res. 2024 Apr 12;52(6):3050-3068.

35.   Hu SB, Xiang JF, Li X, Xu Y, Xue W, Huang M, et al. Protein arginine
methyltransferase CARM1 attenuates the paraspeckle-mediated nuclear retention of
mRNAs containing IRAlus. Genes Dev. 2015 Mar 15;29(6):630-45.

36.   Cui Y, Xie R, Zhang X, Liu Y, Hu Y, Li Y, et al. OGA is associated with
deglycosylation of NONO and the KU complex during DNA damage repair. Cell Death
Dis. 2021 Jun 16;12(7):622.

37.   Deshar R, Yoo W, Cho EB, Kim S, Yoon JB. RNF8 mediates NONO degradation
following UV-induced DNA damage to properly terminate ATR-CHK1 checkpoint
signaling. Nucleic Acids Res. 2019 Jan 25;47(2):762-778.

38.   Cox J, Mann M. MaxQuant enables high peptide identification rates, individualized
p.p.b.-range mass accuracies and proteome-wide protein quantification. Nat Biotechnol.
2008 Dec;26(12):1367-72.

39.   Cox J, Hein MY, Luber CA, Paron I, Nagaraj N, Mann M. Accurate proteome-wide
label-free quantification by delayed normalization and maximal peptide ratio extraction,
termed MaxLFQ. Mol Cell Proteomics. 2014 Sep;13(9):2513-26.

40.   Naganuma T, Nakagawa S, Tanigawa A, Sasaki YF, Goshima N, Hirose T.
Alternative 3'-end processing of long noncoding RNA initiates construction of nuclear
paraspeckles. EMBO J. 2012 Oct 17;31(20):4020-34.

41.   West JA, Davis CP, Sunwoo H, Simon MD, Sadreyev RI, Wang PI, et al. The long
noncoding RNAs NEAT1 and MALAT1 bind active chromatin sites. Mol Cell. 2014 Sep
4;55(5):791-802.

42.   Trifault B, Mamontova V, Burger K. In vivo Proximity Labeling of Nuclear and
Nucleolar Proteins by a Stably Expressed, DNA Damage-Responsive NONO-APEX2
Fusion Protein. Front Mol Biosci. 2022 Jun 6;9:914873.

43.   Knott GJ, Chong YS, Passon DM, Liang XH, Deplazes E, Conte MR, et al. Structural
basis of dimerization and nucleic acid binding of human DBHS proteins NONO and
PSPC1. Nucleic Acids Res. 2022 Jan 11;50(1):522-535.

44.   Beli P, Lukashchuk N, Wagner SA, Weinert BT, Olsen JV, Baskcomb L, et al.
Proteomic investigations reveal a role for RNA processing factor THRAP3 in the DNA
damage response. Mol Cell. 2012 Apr 27;46(2):212-25.

45.   Hung V, Udeshi ND, Lam SS, Loh KH, Cox KJ, Pedram K, et al. Spatially resolved
proteomic mapping in living cells with the engineered peroxidase APEX2. Nat Protoc.
2016 Mar;11(3):456-75.

46.   Moore HM, Bai B, Boisvert FM, Latonen L, Rantanen V, Simpson JC, et al.
Quantitative proteomics and dynamic imaging of the nucleolus reveal distinct responses to
UV and ionizing radiation. Mol Cell Proteomics. 2011 Oct;10(10):M111.009241.

47.   Vizcaíno JA, Côté RG, Csordas A, Dianes JA, Fabregat A, Foster JM, et al. The
PRoteomics IDEntifications (PRIDE) database and associated tools: status in 2013. Nucleic
Acids Res. 2013 Jan;41(Database issue):D1063-9.

FIGURE LEGENDS

Figure 1. Assessment of NONO interactomes in U2OS cells. (A) Intensity versus ratio plots

displaying NONO mass spectrometry hits. 238 proteins that were detected with at least five

unique peptides identified by mass spectrometry are shown. Data are normalized to IgG isotype

controls and represent two merged biological replicates. The x-axis/y-axis displays

ratios/intensities of peptides. The number of unique peptides correlates with the size of each

circle and the significance is color-coded. (B, C) Immunoblots detecting NONO, PSPC1 and

SFPQ in whole cell extracts (IN, input) and upon immunoprecipitation (IP) with NONO

antibody (B) or SFPQ antibody (C). IgG, isotype control; Ki-67, negative control. (D) Venn

diagram showing overlap of NONO mass spectrometry hits. The top100 proteins have been

assessed.

Figure 2. Comparative analysis of NONO mass spectrometry hits in U2OS cells. (A)

Integration of mass spectrometry conditions by ratio versus ratio depiction. 49 proteins that

were detected with at least five unique peptides identified by mass spectrometry and that were

at least 4-fold enriched over IgG isotype controls are shown. Data represent two merged

biological replicates. Both the x-axis and the y-axis display ratios of peptides, whilst the

diagonal axis displays intensities. The number of unique peptides correlates with the size of

each circle and the significance is color-coded. Green box, nuclear protein; grey box, non-

significant hits. (B) Table summarizing mass spectrometry data for top30 proteins in the

absence or presence of etoposide.

Figure 3. Assessment of DBHS protein localization in U2OS cells. (A) Confocal imaging of

either NONO (left), SFPQ (middle) or PSPC1 (right) in combination with HA-NONO upon

transient transfection of pcDNA3.1-HA-NONO expression plasmid. Representative images are

shown. (B) Quantitation of (A) using RGB profiler line scan analysis. a.u., arbitrary units.

Figure 1

A

NONO interactome (Control)

NONO interactome (Etoposide)

B

C

D

Figure 2

A

Comparative NONO interactome (≥5 unique peptides in IP NONO)

B

Control

Rank	Gene	Average number of unique peptides		Average sequence coverage in %		Fold-enrichment (IP NONO / IP IgG)
		IP IgG	IP NONO	IP IgG	IP NONO	
1	NONO	13.0	62.0	28.5	70.4	231.7
	PSPC1	1.5	20.5	3.2	34.8	221.9
	SFPQ	19.0	63.0	28.7	58.3	107.8
	MOV10	7.0	27.5	8.2	28.5	83.5
	SRGAP2	6.0	30.0	6.4	31.6	76.9
	RBM27	0.0	17.0	0.0	17.0	54.1
	PABPC4	7.5	18.5	21.9	35.5	28.6
	RPL3	5.0	16.5	12.4	34.3	27.9
	UPF1	11.0	28.5	11.9	29.1	25.2
10	YBX1	3.5	7.5	33.4	44.5	22.7
	FXR2	0.5	8.5	4.0	21.5	22.2
	FAM120A	3.5	14.5	4.3	18.5	21.4
	NUFIP2	1.5	8.0	2.7	14.3	20.3
	FXR1	4.5	17.5	12.7	30.3	19.3
	TRIM21	14.0	26.5	24.5	42.3	19.2
	LARP1	4.0	15.5	4.7	17.2	18.6
	RPL13	2.0	7.5	11.4	26.8	18.3
	LACTB	0.5	9.0	1.0	15.3	16.2
	PRRC2C	2.0	16.0	1.1	7.6	14.9
20	RPL4	3.5	6.5	7.4	13.9	12.5
	PRRC2A	1.5	14.5	1.3	10.6	9.8
	DHX29	2.0	10.5	2.3	9.2	8.4
	CDC5L	3.5	7.5	5.6	12.6	8.2
	RPL24	0.0	6.0	0.0	32.2	7.8
	RPLP0; RPLP0P6	6.5	6.5	23.7	23.2	5.9
	DDX27	4.0	7.5	6.0	10.7	5.9
	PABPC1	13.5	25.5	30.7	44.5	5.8
	ATXN2L	6.5	12.5	8.1	16.2	5.7
	HNRNPUL1	12.5	17.0	17.3	22.9	5.6
30	RBM14	7.5	13.0	15.3	21.7	5.4

Etoposide

Gene	Average number of unique peptides		Average sequence coverage in %		Fold-enrichment (IP NONO / IP IgG)
	IP IgG	IP NONO	IP IgG	IP NONO	
PSPC1	0.5	31.5	1.0	43.5	317.1
NONO	13.0	68.5	24.9	77.0	179.4
SFPQ	8.5	72.5	14.2	62.4	145.7
SRGAP2	2.5	46.5	2.4	46.1	132.5
YBX1	0.5	12.5	1.4	55.0	117.5
RBM27	0.5	23.0	0.5	22.2	105.2
UPF1	2.0	38.5	2.0	38.0	97.9
RPL3	1.5	23.5	3.9	45.6	78.7
MOV10	3.0	39.0	3.1	40.6	67.7
FAM120A	0.5	24.5	0.8	27.6	49.3
PABPC4	2.0	18.0	8.1	37.5	44.6
FXR2	0.0	14.0	1.2	31.6	42.9
NUFIP2	1.0	10.5	1.9	16.4	37.7
RPLP0; RPLP0P6	3.0	10.0	14.9	30.5	37.3
LACTB	0.5	12.5	1.0	20.1	31.9
LARP1	4.0	26.0	4.2	28.8	24.6
FXR1	3.0	20.0	6.6	33.3	24.3
PRRC2C	0.0	30.0	0.0	12.5	23.0
PRRC2A	0.0	19.5	0.0	13.9	21.4
RPL13	1.0	13.0	5.7	43.1	20.8
RPL5	3.0	21.0	11.6	49.4	20.1
ATXN2L	2.5	19.0	2.6	23.2	19.1
DHX29	0.5	19.5	0.5	18.4	17.8
RPL4	0.0	16.0	0.0	31.4	17.4
USP10	0.5	10.5	0.8	15.5	14.7
TRIM21	16.0	28.5	27.5	46.1	12.6
RPL24	0.5	8.0	2.9	47.5	13.0
ZC3HAV1	1.5	13.0	2.3	17.4	11.0
PABPC1	9.0	25.5	18.9	48.5	10.3
SYNCRIP	1.5	8.0	6.4	21.5	10.2

Figure 3