1 **Functional-space alignment resolves the eco-evolutionary landscape of**

2 **siderophore biosynthesis across bacteria**

3 Jiqi Shao[1,2], Yanzhao Wu[3], Shizheng Tian[4], Ruichen Xu[4], Haohua Luo[4], Yuanzhe Shao[4],
4 Linlong Yu[4], Ruolin He[1], Guanyue Xiong[1], Peng Guo[5], Nan Rong[5], Zhong Wei[6*], Shaohua
5 Gu[2*], Zhiyuan Li[1,4*]
6
[1]Quantitative Biology, Academy for Advanced Interdisciplinary Studies, Peking University,
Beijing 100871, China.
[2]College of Resources and Environmental Sciences, State Key Laboratory of Nutrient Use
and Management (SKL-NUM), National Academy of Agriculture Green Development,
China Agricultural University, Beijing 100193, China.
[3]College of Computer Science and Technology (College of Data Science), Taiyuan
University of Technology, Taiyuan, 030600, China.
[4]Peking-Tsinghua Center for Life Sciences, Academy for Advanced Interdisciplinary
Studies, Peking University, Beijing 100871, China.
[5]Institute of Zoology, Chinese Academy of Sciences, Beijing 100101, China.
[6]Jiangsu Provincial Key Lab for Organic Solid Waste Utilization, Jiangsu Collaborative
Innovation Center for Solid Organic Waste Resource Utilization, National Engineering
Research Center for Organic-based Fertilizers, Nanjing Agricultural University, Nanjing
210095, China
21
22
[*]Correspondence: zhiyuanli@pku.edu.cn, gsh2025@cau.edu.cn, weizhong@njau.edu.cn

Abstract

Siderophores are central mediators of microbial iron acquisition, competition, and
ecological adaptation, yet their biosynthetic diversity remains difficult to resolve across
species because existing sequence-based BGC comparison is strongly constrained by
phylogenetic background. Here we combine large-language-model-assisted literature
mining, functional-space comparison, and genome-scale analysis to resolve the global
organization of siderophore biosynthesis across bacteria. We first built SideroBank, a
manually curated cross-species benchmark of siderophore biosynthetic gene clusters
(BGCs), and used it to show that many identical products recur across distant taxa
whereas the corresponding BGCs often fail to cluster in sequence space. We then
developed BGC Block Aligner, which compares BGCs as ordered systems of functionally
meaningful blocks and thereby converts comparison from sequence space to functional
space. Applied to 97,432 bacterial genomes, this framework produced the Siderophore
Atlas, revealing that siderophore synthesis is a remarkably pervasive trait encoded by

over 60% of the analyzed genomes, with certain clusters being the most widely disseminated secondary metabolites across the bacterial domain. This global landscape suggests that the adoption of specific biosynthetic strategies is predominantly driven by ecological lifestyle rather than strict phylogenetic relatedness. Furthermore, a stark macro-evolutionary dichotomy **was** observed between the continuous structural diversification of NRPS pathways and the standardized, HGT-driven dissemination of NIS systems, linking functional-space genomics to the global ecology and evolution of siderophore biosynthesis..

Introduction

Microbial natural products represent a major reservoir of chemical diversity and play central roles in survival, competition, and environmental adaptation. Advances in genome sequencing have transformed natural product discovery from a purely chemistry-driven enterprise into a comparative genomic problem, revealing immense and previously hidden biosynthetic potential across cultivated and uncultivated microorganisms[1]. Among natural products, siderophores are especially informative because they connect a chemically tractable biosynthetic system to a universal ecological problem: the acquisition of bioavailable iron. Beyond iron scavenging, siderophores also shape microbial competition, community interactions, host association, and pathogenicity, and have attracted growing attention in biotechnology and therapeutic development[2-4]. As such, resolving how siderophore biosynthesis is distributed across bacteria, and how similar products are encoded across divergent lineages, is central to understanding both microbial ecology and natural product evolution.

The rapid accumulation of genome and metagenome data has been accompanied by a broad suite of tools for BGC detection, annotation, and large-scale organization. antiSMASH has made genome-wide identification of biosynthetic pathways routine[5], while DeepBGC and related machine-learning or deep-learning approaches have expanded BGC prediction and classification[6,7]. Curated resources such as MIBiG provide experimentally validated reference clusters for comparative analysis[8]. At the comparison stage, BiG-SCAPE and BiG-SLiCE have enabled large-scale clustering of BGCs by using domain content, domain order, and sequence similarity to define gene cluster families[9-11], whereas pathway-specific tools such as RiPPMiner and TransATor demonstrate the value of incorporating biosynthetic logic for specialized metabolite classes[12,13]. Together, these frameworks have greatly accelerated the scalable exploration of biosynthetic diversity, but they mainly fundamentally rooted in sequence-based BGC comparison.

For many modular biosynthetic systems, however, sequence proximity is only an imperfect proxy for functional relatedness. In pathways such as NRPS and NRPS-independent(NIS), product identity is determined less by global full-length similarity than by the substrate specificity, functional identity, and ordered organization of key

biosynthetic modules[14]. In NRPS systems, substrate choice is encoded in
adenylation-domain recognition logic, classically captured by specificity-conferring
residues and later extended by machine-learning-based predictors[15-18]. More broadly,
NRPS diversification is shaped by recombination, exchange, and reorganization of
modular units, emphasizing that local functional modules rather than full-length sequence
similarity often determine biosynthetic output. In siderophore pathways, similar
complications arise in NRPS-independent systems, where IucA/IucC-like synthetases and
their active-site logic define core product assembly[3,19]. As a result, proteins from distant
taxa may encode equivalent biosynthetic functions despite extensive sequence
divergence, whereas globally similar proteins in closely related organisms may encode
different substrate preferences. Functional differences and phylogenetic differences are
therefore entangled in sequence space.
92
A second obstacle is that the reference knowledge needed to interpret siderophore
diversity remains highly fragmented. Existing resources typically emphasize either
chemical structures or a relatively small number of curated BGCs, whereas critical links
among product identity, biosynthetic genes, and taxonomic distribution remain dispersed
across the literature[8,20]. This fragmentation is particularly limiting for siderophores, whose
biosynthesis has been investigated across many taxa, habitats, and research traditions
but rarely organized into a unified cross-species reference system. Meanwhile, recent
advances in protein structure prediction suggest that functionally constrained features
may provide a more stable basis for cross-species comparison than full-length sequence
similarity alone[21]. Together, these observations point to two unmet needs: a
literature-derived cross-species benchmark that connects product identity to BGC
architecture, and a comparison framework capable of recovering functional equivalence
beyond the limits imposed by phylogenetic background.
106
Here, we address these limitations through an integrated framework that links
literature-scale knowledge reconstruction to genome-scale functional comparison. We
first developed Sidero-Mining, a large-language-model-assisted workflow for extracting
siderophore biosynthetic knowledge from the literature, and used it to construct
SideroBank, a manually curated cross-species benchmark of siderophore BGCs. Using
this benchmark, we show that many siderophores recur across distant taxa whereas the
corresponding BGCs often fail to cluster under existing sequence-based methods,
revealing a major phylogenetic constraint in current comparison frameworks. We then
developed BGC Block Aligner (BBA), which represents BGCs as ordered systems of
functionally meaningful blocks and compares them through block-level alignment, thereby
converting BGC comparison from sequence space to functional space. This framework
enables functional typing of siderophore BGCs at a global scale, allowing us to build the
Siderophore Atlas and to resolve the global organization, taxonomic spread, and
contrasting macro-evolutionary regimes of NRPS and NIS siderophore pathways across
bacteria.

122 Result

123 LLM-mined SideroBank exposes the phylogenetic constraint of sequence-based

124 BGC comparison

High-quality reference datasets are essential for linking siderophore molecules to their
biosynthetic gene clusters (BGCs), yet such resources remain incomplete. Existing
databases either focus primarily on chemical structures or contain only limited numbers of
annotated siderophore BGCs, whereas the key links among product identity, biosynthetic
genes, and taxonomic distribution remain scattered across the literature. To address this
gap, we developed Sidero-Mining, a large-language-model-assisted workflow for
literature-scale extraction of siderophore biosynthetic information, and used it to construct
SideroBank, a manually curated cross-species siderophore BGC resource (Figure 1A,
Figure S1−S12). Through systematic screening of more than 10,000 siderophore-related
publications, we retained 1,823 high-confidence articles for structured mining. After
automated extraction, manual verification, genome retrieval, and BGC annotation,
SideroBank compiled 738 nonredundant siderophore BGCs together with 325 NRPS
adenylation-domain substrate annotations(Figure S13), substantially expanding the
currently available siderophore reference space.
139

Figure 1. LLM-mined SideroBank reveals the phylogenetic constraint of sequence-based comparison for cross-species siderophore BGCs

(A) Schematic overview of the Sidero-Mining workflow. This pipeline uses a large language model for large-scale screening and structured extraction of siderophore-related literature, followed by manual curation, genome retrieval, and BGC annotation, ultimately generating SideroBank, a benchmark dataset of cross-species siderophore biosynthetic gene clusters.

(B) Taxonomic distribution of the major cross-species siderophore products curated in SideroBank for NRPS and NIS pathways.

(C) UMAP projection of pairwise sequence similarity among siderophore BGCs calculated by BiG-SCAPE. Red tones represent NRPS, and blue tones represent NIS. Color intensity indicates the abundance of BGCs within each product type.

(D) Pairwise similarity heatmap and statistical analysis of cross-species same-product

BGCs in the NRPS pathway.

(E) Pairwise similarity heatmap and statistical analysis of cross-species same-product

BGCs in the NIS pathway.

Systematic curation through SideroBank revealed that many siderophores recur across

distantly related taxa, including across genera and higher taxonomic ranks, indicating that

the taxonomic distribution of the same product is far broader than previously appreciated

(Figure 1B). To further test whether existing comparison frameworks could reliably

recognize these cross-species same-product BGCs, we used BiG-SCAPE to calculate

pairwise similarity and visualize the resulting relationships in reduced dimensional space.

For both NRPS and NIS pathways, BGCs derived from different species but encoding the

same product often failed to cluster together in sequence similarity space (Figure 1C).

Further analysis showed that similarity among BGCs encoding the same product was

largely concentrated within genera, whereas pairwise similarity across genera was

significantly lower. The mean within-genus versus between-genus similarity difference

reached 0.46 for NRPS and 0.38 for NIS, indicating that substantial sequence divergence

can arise even among cross-species same-product BGCs.

Because BiG-SCAPE measures similarity primarily from sequence similarity between

matched domains, this sequence-space-based comparison is poorly suited for stably

recognizing functionally equivalent BGCs across species and may systematically split

them into separate groups. Overall, SideroBank not only revealed the widespread

cross-species distribution of siderophore products, but also showed that existing

sequence-based BGC comparison is insufficient to stably recover the correspondence

among cross-species same-product BGCs. More importantly, by integrating a large

number of manually curated cross-species same-product cases, SideroBank provides a

critical benchmark dataset for evaluating the limitations of existing frameworks and for

developing new methods capable of decoupling functional similarity from sequence

divergence imposed by phylogenetic background.

BGC Block Aligner converts BGC comparison from sequence space to functional

space through block-level alignment

The inability of sequence-based BGC comparison to stably recover cross-species

same-product BGCs indicates that global sequence similarity cannot serve as a reliable

proxy for biosynthetic equivalence. Just as the biological meaning of nucleic acid and

protein sequences is determined by the identity and order of their basic units, BGC

function is likewise defined primarily by the functional identity, substrate specificity, and

ordered organization of key biosynthetic modules. Based on this principle, we developed

BGC Block Aligner (BBA). BBA first decomposes each BGC into functionally meaningful

blocks, each representing a unit responsible for substrate recognition, scaffold
construction, or key tailoring steps. These blocks are then organized into ordered block
sequences according to their native biosynthetic organization and compared through
block-level alignment, analogously to biological sequence alignment. In this way, BBA
compares not raw sequences themselves, but function-defining units and their
organizational relationships, thereby converting BGC comparison from sequence space to
functional space(Figure 2A).
202
203
In BBA, block definition follows a function-first principle: core modules directly involved in
product biosynthesis are extracted, whereas background genes unrelated to the target
biosynthetic process are minimized; fragmented BGCs are merged using core domains as
anchors when appropriate. For siderophore pathways, the most critical blocks are the
biosynthetic modules that determine substrate recognition and core scaffold assembly,
such as the A domain in NRPS pathways and IucA/IucC-like enzymes in NIS pathways.
BBA further assigns block-level similarity matrices to different block types(Figure S14,
S15). For NRPS, these matrices are based primarily on substrate-recognition features of
A domains; for NIS, they incorporate structure-guided feature sequences derived from
active-site-related positions. In both NRPS and NIS pathways, features extracted from
substrate-recognition sites or structural constraints showed greater cross-species stability
and stronger functional discrimination than full-length sequences, indicating that the
signals most informative for cross-species comparison reside not in overall full-length
similarity, but in function-defining sites directly related to biosynthetic activity.

Figure 2. BGC Block Aligner converts siderophore BGC comparison from sequence space to functional space through block-level alignment

(A) Schematic overview of BGC Block Aligner (BBA). BBA first decomposes each BGC into functionally meaningful blocks, where each block corresponds to a functional unit involved in substrate recognition, scaffold construction, or key tailoring steps. For core blocks responsible for product scaffold formation, functional features are extracted to construct variable similarity matrices; for auxiliary backbone blocks, fixed scoring matrices are defined. These blocks are then organized into ordered block sequences according to

their native biosynthetic arrangement and compared through block-level alignment.
Similarity among nearby domains is further integrated to derive an overall functional
similarity, thereby converting BGC comparison from sequence space to functional space.
(B) Functional similarity matrix and clustering results for cross-species same-product
BGCs in the NRPS pathway calculated by BBA. Compared with sequence-based
methods, BBA more strongly organizes BGCs by product identity rather than by host
taxonomy.
(C) Functional similarity matrix and clustering results for cross-species same-product
BGCs in the NIS pathway calculated by BBA. Similarly, BBA more stably recovers
functional correspondence among cross-species same-product BGCs in the NIS pathway.
(D) Relationship between pairwise BGC distance and host genome distance for NRPS
same-product BGCs across species. The x-axis represents host genome distance and the
y-axis represents pairwise BGC distance. Red circles indicate BBA, and grey triangles
indicate BiG-SCAPE.
(E) Relationship between pairwise BGC distance and host genome distance for NIS
same-product BGCs across species. Blue circles indicate BBA, and grey triangles indicate
BiG-SCAPE.
(F) Relationship between BGC similarity and product chemical similarity in the NRPS
dataset. Product pairs were grouped into three bins according to chemical similarity:
Chem low (<0.3), Chem mid (0.3−0.7), and Chem high (>0.7).
(G) Relationship between BGC similarity and product chemical similarity in the NIS
dataset. As in NRPS, BBA-derived functional similarity shows stronger agreement with
product chemical similarity in the NIS pathway.
(H) Comparison of clustering results under different distance thresholds. Red and blue
indicate NRPS and NIS siderophores, respectively; circles indicate BBA, and triangles
indicate BiG-SCAPE.
253
We next evaluated BBA on the SideroBank benchmark dataset. Compared with
BiG-SCAPE, the similarity matrices generated by BBA clustered BGCs more clearly by
product identity rather than by host taxonomy, with the strong genus-level partitioning
seen in sequence-based comparison markedly reduced (Figure 2B, C). In pairwise
distance analyses, BBA produced substantially lower within-group distances for
same-product BGCs than BiG-SCAPE: the mean distance was 0.09 for NRPS under BBA
versus 0.50 under BiG-SCAPE, and 0.19 for NIS under BBA versus 0.80 under
BiG-SCAPE, with markedly tighter distributions. When genome-level species distance
was used as the x-axis to compare pairwise BGC distances across methods, BiG-SCAPE
distances rose rapidly with increasing phylogenetic separation, whereas BBA functional
distances increased only slightly and remained comparatively stable overall (Figure 2D, E),
indicating much lower sensitivity to phylogenetic background. Moreover, when BGC
similarity was stratified by product structural similarity, BBA showed substantially stronger
agreement with chemical similarity than sequence-based comparison. In the subset with
chemical similarity greater than 0.7, the mean functional similarity under BBA reached
0.78 for NRPS and 0.84 for NIS (Figure 2F, G). Under this metric, BBA also enabled more
stable grouping at smaller thresholds and substantially reduced the inflation of cluster

counts caused by phylogeny-driven sequence divergence (Figure 2H). Together, these
results show that BBA provides a functional-space metric for cross-species BGC
comparison that is more closely aligned with biosynthetic output than raw sequence
similarity.

Siderophore Atlas: high-resolution functional typing of siderophore BGCs in

functional space

Having established the functional-space distance defined by BBA, we next applied it to
large-scale functional typing of siderophore BGCs across bacterial genomes. Unlike
conventional typing strategies that rely on sequence similarity, BBA provides a
cross-species comparable measure of functional similarity, enabling a unified global view
of siderophore biosynthetic potential. Guided by the distance distributions of manually
curated same-product BGCs in SideroBank, we applied stringent criteria to perform
fine-grained functional typing of NRPS and NIS BGCs and used the resulting classification
to construct the Siderophore Atlas.
285
We applied BBA to 97,432 bacterial genomes and functionally typed 148,112 annotated
NRPS BGCs, generating a global NRPS functional tree and identifying 7,479 distinct
NRPS BGC functional types. Among these, 1,196 (15.1%) were assigned as dedicated
NRPS-type siderophore biosynthetic types. Notably, although these siderophore-related
functional types represent only a minority of all NRPS types, they were distributed across
48.3% of genomes, indicating striking ecological prevalence. These NRPS siderophore
types were not confined to a small local region of the functional tree, but instead occupied
multiple separated branches, and several types were broadly distributed across many
genera. This pattern suggests that siderophore biosynthesis in NRPS functional space is
not restricted to a single localized sector, but is supported by multiple differentiated
combinations of function-defining modules.
297
Similarly, we functionally typed 39,246 NIS BGCs and identified 358 distinct NIS
siderophore functional types. Together with the NRPS results, we ultimately defined 1,554
independent functionally defined BGC families for siderophore biosynthesis and named
this global function-space-based classification framework the Siderophore Atlas. Overall,
64.8% of bacterial genomes encoded at least one siderophore biosynthetic system (Table
S3), indicating that siderophore production is not restricted to a few specialized lineages
but is instead a widespread adaptive capacity across bacterial genomes. Importantly, the
Siderophore Atlas is not a catalog organized by conventional sequence clustering, but a
high-resolution atlas organized by functional equivalence, enabling comparison of
siderophore biosynthetic potential across phylogenetically diverse bacteria within a unified
functional space.
309

Figure 3. The Siderophore Atlas resolves the global functional organization of siderophore biosynthesis across bacterial genomes

(A) Global functional landscape of NRPS biosynthetic gene clusters across bacterial genomes. This map was generated by applying BGC Block Aligner to NRPS BGCs identified from 97,432 bacterial genomes. Each point represents a functionally typed NRPS BGC cluster observed in a given genus. Red tones indicate siderophore-associated NRPS clusters, and deeper color indicates broader cross-genus distribution.

(B) Functional tree of more than 1,000 NRPS siderophore types. Each point represents a functionally typed NRPS BGC cluster observed in a given genus. Distinct colors highlight

321 the top siderophore types showing the broadest cross-species distribution.

322 (C) Functional tree of NIS siderophore types. Each point represents a functionally typed

323 NIS BGC cluster observed in a given genus. Distinct colors highlight the top siderophore

324 types showing the broadest cross-species distribution.

Macro-evolutionary landscape and taxonomic distribution of siderophore

biosynthetic strategies

A global survey on siderophore prevalence across taxonomic levels reveals a paradoxical

pattern. While siderophore biosynthesis is present in approximately two-thirds of genomes,

its prevalence declines sharply at higher taxonomic ranks, dropping to ~34% at the order

level and ~27% at the class level (Table S3). This steep reduction suggests that

siderophore production is not universally conserved, but instead represents a highly

selected trait shaped by specific ecological lifestyles.

To further resolve these lifestyle-associated pressures, we grouped bacterial classes

based on their overall siderophore biosynthetic strategies (Figure 4A), revealing four

distinct ecological archetypes that are also observed in other taxonomic levels (Figure

S16):

1. The NRPS Generalists (archetype 1). Represented by Gammaproteobacteria and

Actinomycetes, this archetype is characterized by a high abundance of NRPS

siderophore BGCs and a frequent co-occurrence of NIS siderophores. These lineages

invest heavily in complex modular biosynthetic pathways, consistent with adaptation

to highly competitive and resource-limited microenvironments.

2. The NIS Specialists (archetype 2). Represented by classes such as Bacilli and

Flavobacteriia, this group predominantly synthesize NIS siderophores while largely

lacking canonical NRPS machinery. This streamlined strategy suggests specialization

toward energetically efficient iron acquisition.

Notably, although these two archetypes comprise less than half of all recognized

bacterial classes, they encompass the majority of commonly sequenced taxa relevant

to human health and agriculture, explaining their disproportionate representation at

the genome level. For example, *Gammaproteobacteria* harbor notorious human

pathogens and *Bacilli* have been widely utilized agricultural biocontrol agents, while

Alphaproteobacteria and *Actinomycetes* are foundational to crop symbioses and soil

biogeochemistry.

3. In contrast, the remaining classes reflect alternative survival strategies. The

Non-siderophore NRPS Producers (archetype 3; spanning Desulfobacteria to

Tissierellia) retain substantial NRPS capacity but appear to redirect these biosynthetic

resources toward other classes of secondary metabolites (e.g., toxins, antimicrobials,

or pigments) rather than siderophores. This reallocation is ecologically viable because

these organisms secure iron through alternative avenues, such as anaerobic

sulfate-reducers Desulfobacteria access soluble ferrous iron, and predatory lineages

Myxococcia recycle iron from lysed prey, freeing their biosynthetic potential for

chemical warfare and spatial competition.

4. The Minimalists (archetype 4; from Bacteroidia to Synergistia) lack both NRPS-
potential and siderophore biosynthetic pathways, consistent with a minimalists
lifestyles such as obligate anaerobiosis or host dependence, where iron acquisition
may rely on alternative mechanisms or iron-sparing strategies.

Figure 4. Macro-evolutionary landscape and taxonomic distribution of siderophore
biosynthetic strategies.
(A) Distribution of siderophore biosynthetic strategies across bacterial classes. The
heatmap quantifies the percentage of genomes within each class adopting one of four
mutually exclusive pathways: *Other NRPS* (possessing NRPS machinery but lacking
dedicated siderophore BGCs); *NI-Sid only*; *NRP-Sid only*; and *Both Sid* (co-occurrence of
both systems). Classes are vertically ordered by hierarchical clustering based on strategy
profile similarity.
(B) Compositional diversity of siderophore BGCs. The total length of each horizontal bar
represents the Shannon Index (in bits), quantifying the cumulative siderophore diversity
within each class. Bars are subdivided proportionally to reflect the relative abundance of
specific BGCs. Focal siderophores are color-coded according to the legend (NIS in cold
tones, NRPS in warm tones), while grey and white segments denote the minor
contributions of other annotated and unannotated siderophore BGCs, respectively.
(C) Biosignature landscape of siderophore potential across bacterial orders. A
dual-hierarchical clustering heatmap illustrating the occurrence of distinct siderophore
BGCs across varied orders. Color intensity denotes the Z-score normalized occurrence of
each BGC (red: high; dark blue: absent/low). Rows represent bacterial orders, with labels
color-coded by their parent class. Columns represent individual siderophore BGCs; labels
for focal BGCs are colored consistently with plot B, non-focal BGCs with annotation are
colored black, while unannotated BGCs are omitted for visual clarity. The top annotation
band delineates the overarching biosynthetic logic: NRPS-type (peach) versus NIS-type
(light blue). Clustering was performed using Ward's linkage based on the Euclidean
distance of prevalence profiles.

Focusing on bacterial classes with siderophore prevalence exceeding 10%, we observe
substantial heterogeneity in their internal biosynthetic repertoires (Figure 4B). To
characterize this variation, we tracked the proportional abundance of 20 focal
siderophores representing major chemical classes of clinical and ecological significance.
The resulting compositional profiles reveal a marked contrast in diversification strategies.
Certain clades within the NIS Specialist archetype, such as Bacilli, exhibit relatively low
internal diversity, dominated by a few highly successful focal siderophores. In contrast,
lineages such as Actinomycetes display substantially broader biosynthetic diversity,
driven largely by a long tail of non-focal and frequently unannotated BGCs. Accordingly,
internal chemical diversity does not strictly correlate with overall siderophore prevalence,
as evident in Cyanophyceae. These compositional differences warrant further
investigation at the order level.

To further resolve these lifestyle-associated pressures at higher resolution,
dual-hierarchical clustering of siderophore biosignatures (Figure 4C, Figure S17, Figure S
18) reveals that biosynthetic strategies across bacterial orders frequently group according
to shared functional profiles, rather than phylogenetic relatedness. For example, certain
orders within Gammaproteobacteria (e.g., Enterobacterales and Pseudomonadales)
cluster in close proximity to Betaproteobacteria (e.g., Burkholderiales), suggesting

convergence in siderophore repertoires. This pattern is consistent with adaptation to
similarly competitive environments, where complex NRPS or hybrid siderophore systems
may provide selective advantages. Conversely, marked divergence is observed even
among closely related taxa. Within Actinomycetes, Mycobacteriales separates distinctly in
the clustering space and is strongly dominated by mycobactin-like BGCs, whereas related
orders such as Kitasatosporales and Pseudonocardiales display alternative repertoires,
including NIS-type desferrioxamine and diverse NRPS pathways. This contrast highlights
the functional plasticity of siderophore biosynthesis within phylogenetically related groups.
421
Shifting the focus from host taxonomy to the metabolites themselves, the heatmap reveals
substantial variation in the distribution patterns of individual siderophores. Some BGCs,
such as enterobactin and desferrioxamine, show relatively broad occurrence across
multiple, phylogenetically distant orders, with recurring hotspots of high prevalence. This
pattern is consistent with high evolutionary mobility and potential horizontal transfer. In
contrast, other siderophores, including yersiniabactin and bacillibactin, exhibit more
restricted distributions, largely confined to specific lineages. These differences point to
distinct evolutionary trajectories, potentially reflecting variation in ecological specialization
and dissemination capacity.
431
432 Together, these observations support a framework in which siderophore diversity is
433 shaped not only by vertical inheritance but also by differential mobility and ecological
434 selection, motivating a conceptual decoupling of siderophore repertoires from host
435 taxonomy.
436

Taxonomic spread reveals contrasting evolutionary patterns of NRPS and NIS

pathways

The taxonomic distributions explored in the previous section expose a decoupling: a
BGC's prevalence at the genome level does not necessarily dictate its ability to permeate
higher taxonomic ranks. To quantify this cross-lineage mobility, we introduce the metric of

"taxonomic spread." We define this as the absolute number of distinct host clades (e.g.,

classes) in which a specific BGC is detected, regardless of its genome frequency within
those clades (Figure 5A). In essence, if a BGC appears in at least one genome of a given
class, that class contributes to its total spread score.
446
Visualizing the BGCs with the highest class-level spread (spread > 3; Figure 5A)
highlights two preliminary trends. First, NRPS siderophores appear disproportionately
represented among the most widely disseminated NRPS clusters. Despite comprising
only 15% of the total NRPS repertoire, they account for three of the top five most broadly
distributed NRPS BGCs, led by enterobactin (spread = 6). Second, the absolute highest
spread values are dominated by a select group of NIS siderophores. Although NIS

systems are present in a smaller fraction of global genomes (30%) compared to NRPS
siderophores (~48%), their top-ranking BGCs achieve exceptional cross-clade distribution.
Seven distinct NIS pathways exceed a class-level spread of 3, spearheaded by
schizokinen (spread = 8) and desferrioxamine (spread = 7).
457

458
Figure 5. Taxonomic spread reveals contrasting evolutionary patterns of NRPS and
NIS pathways.
(A) Taxonomic distribution of high-spread siderophore BGCs at the class level. The
top panel displays the specific spread value (number of distinct host classes) for each
biosynthetic gene cluster (BGC), highlighting those with a spread greater than three. The
lower bubble matrix illustrates the occurrence of these BGCs across host classes, where
bubble size corresponds to the genome prevalence of a BGC within each taxon. "Star" or
focal molecules are identified by their specific assigned colors (consistent with Figure 4),
while other NRPS and NIS clusters are shown in black (e.g., indigoidine) and grey. Host
taxa on the Y-axis are sorted by BGC richness, and BGCs on the X-axis are ordered by
descending spread value. A vertical dashed line separates NRPS-type clusters (left) from
NIS-type clusters (right).
(B) Comparative spread distribution of siderophore and non-siderophore NRPS
BGCs across taxonomic levels. The 3X2 panel illustrates the proportion of biosynthetic
gene clusters (BGCs) with specific spread values (2, 3, and >3) relative to the total
number of BGCs, Color coding distinguishes different spread values: light green

(Spread=2), yellow (Spread = 3), and salmon (Spread > 3). "Sid" denotes
siderophore-related NRPS clusters, and "Other" denotes non-siderophore NRPS clusters.
Each subplot corresponds to a distinct taxonomic level, from species to phylum.

(C). **Rank-abundance distribution of siderophore BGCs at the genus level**. The
log-log plots illustrate the spread of NRP-siderophores (left panel) and NIS-siderophores
(right panel) across host genera. Siderophores are ordered by their descending spread
value on the X-axis (Rank), with the Y-axis representing the number of distinct host
genera (Spread) for each BGC. The grey line indicates the Zipf (power-law) fit, with the
resulting Zipf coefficient (beta) and Kolmogorov-Smirnov (KS) statistic provided in each
panel. Focal molecules are highlighted with specific assigned colors, while other
annotated and unknown clusters are shown in black and grey, respectively.

Defined quantification confirmed that NRPS siderophores consistently exhibit a vastly
superior expansion capacity compared to non-siderophore NRPSs. Specifically, the
fraction of NRPS siderophore BGCs achieving broad cross-lineage distribution (i.e.,
successfully spanning >1, >2, or >3 distinct clades) substantially exceeds that of their
non-siderophore counterparts, at taxonomic scales from genus to phylum.
Complementary cumulative distribution function (CCDF) analysis confirms the statistical
robustness of NPRS siderophores's higher spread (Figure S19). Furthermore, from the
species level up to the order level, the CCDF reveals that the spread frequencies for both
NRPS categories follow a heavy-tailed distribution that tightly fits a power-law model, a
scaling frequently carring intriguring eco-evolutionary implications.

To untangle the macro-ecological dynamics driving these pathways, we visualized the
rank-abundance distributions of siderophore BGCs at the genus level (Figure 5C). Plotting
taxonomic spread against rank on a log-log scale exposes fundamentally divergent
evolutionary architectures between the two biosynthetic classes:

NRPS siderophores strictly adhere to a power-law distribution (KS = 0.07, $\beta = 0.73$).
Scale-free is a typical character of such distribution: While canonical BGCs like
enterobactin occupy the apex, the spread of subsequent clusters (such as vanchrobactin
and yersiniabactin) decays log-linearly. This scale-free geometry precludes any defined
threshold that scales the taxonomic spread, ultimately manifesting as a massive long tail
of rare, frequently unannotated BGCs restricted to single genera. Ecologically, such
power-law topologies can be classically explained by the Simon model, a generative
"duplication-innovation" framework. Under this model, the empirical scaling exponent
($\beta = 0.73$) translates to an estimated innovation rate of approximately $u = 0.27$. This
remarkably high rate implies continuous, localized invention, perfectly mirroring the
modular nature of NRPS assembly lines where domain swapping and recombination
readily generate novel products (Figure 5D).

In contrast, the NIS siderophore distribution displays a cliff-like pattern. This deviation
from power-law is confirmed statistically ($KS = 0.23$) and is visually apparent as a
pronounced downward curvature on the log-log scale, culminating in a substantially

steeper decay ($\beta = 1.05$) than that observed in NRPS. Unlike the continuous decay of
rank-abundance in NRPS, spread of NIS clearly separates into two tiers: The top 6 or 8
highly mobilized focal pathways, including desferrioxamine, schizokinen, and aerobactin,
attain massive taxonomic spread. This is then followed by a sharp decrease, where most
pathways exhibit very low spread values less than 3(Figure 5C, gray dots). This
mathematical profile, capped by a definitive diversity threshold, suggest an evolutionary
reliance on horizontal gene transfer (HGT) of these "top-tier" solutions over *de novo*
assembly-line invention.
527
528   Together, these contrasting mathematical profiles capture two fundamental evolutionary
529   solutions to iron scarcity: the relentless structural diversification of NRPS systems versus
530   the rapid, standardized dissemination of NIS pathways (Figure 5D).
531

Discussion

Natural product genomics has transformed the discovery of microbial biosynthetic
potential by shifting the field from culture-based compound isolation to large-scale
genome- and metagenome-enabled analysis. Tools such as antiSMASH, BiG-SCAPE,
and BiG-SLiCE have greatly improved BGC detection, annotation, and family-level
organization, making comparative natural product genomics a central component of
molecular discovery[5,9-11]. Yet our results indicate that, in the cross-species setting,
sequence-based BGC comparison still faces a deeper bottleneck: global sequence
similarity does not stably recover biosynthetic equivalence. This problem is particularly
evident in siderophores, a class of natural products with well-defined molecular function,
broad ecological importance, and strong clinical relevance. Siderophores are not only
central to microbial iron acquisition but also participate in host-associated interactions,
microbial competition, and therapeutic applications[2]. When cross-species same-product
BGCs cannot be reliably recovered, the consequences extend beyond cluster
classification to downstream functional annotation and ecological interpretation.
547
In this context, SideroBank is valuable not simply because it expands siderophore
reference coverage, but because it establishes a missing cross-species benchmark
dataset. Existing resources tend to emphasize either chemical structure, local cluster
annotation, or a limited set of curated exemplars, but rarely connect product identity, BGC
architecture, host distribution, and cross-species correspondence within one framework.
By integrating a large number of manually curated cross-species same-product cases,
SideroBank makes it possible to systematically ask whether BGCs associated with the
same product should still be recognized as corresponding systems when they occur in
distant lineages. Prior work has shown that siderophore pathways can be replaced, lost,
or horizontally disseminated across bacterial groups, indicating that their distribution does
not map trivially onto host phylogeny[22]. In this sense, SideroBank is not merely a resource
but a reference coordinate system for evaluating the boundary conditions of cross-species
BGC comparison.

Our study also highlights a deeper role for LLM-assisted knowledge extraction in natural
product genomics. Here, the LLM does not merely accelerate literature mining; rather, it
enables fragmented and heterogeneous biosynthetic knowledge to be formalized at scale.
Sidero-Mining therefore acts as a knowledge reconstruction layer, converting
literature-derived molecular knowledge into a structured reference space that can support
benchmark construction, method development, and global functional typing. In other
words, the LLM made knowledge reconstruction at scale possible, while manual curation
made the resulting benchmark reliable. This suggests that the value of LLMs in natural
product research may extend beyond annotation assistance to the construction of
machine-readable biosynthetic reference systems.

Against this benchmark, BGC Block Aligner (BBA) addresses the central comparison
problem at a more fundamental level by redefining what should be compared. Existing
BGC comparison frameworks have greatly advanced natural product genomics, but they
remain rooted largely in domain content, gene order, and full-length sequence similarity,
making them better suited to recovering sequence proximity than stable functional
equivalence. Prior work on NRPS and other modular assembly lines has shown that
product diversification is closely linked to module exchange, local recombination, and the
reorganization of substrate-selecting units[14]. Our results support the view that, for
complex natural product BGCs, the key determinants of product scaffolds and functional
divergence are not global full-length similarity, but the functional identity, substrate
specificity, and ordered organization of core biosynthetic modules. Rather than treating a
BGC as a contiguous stretch of raw sequence, BBA represents it as an ordered system of
functionally meaningful blocks and compares these through block-level alignment, thereby
moving BGC comparison from sequence space to functional space.

The significance of BBA extends beyond this conceptual shift. First, it provides a more
biosynthetically grounded basis for defining functionally defined BGC families, reducing
the tendency of phylogeny-driven sequence divergence to fragment closely related
biosynthetic solutions into separate groups. Second, it improves the transfer of
product-linked annotation into distant genomic backgrounds, making cross-species
functional annotation more stable. Third, because BBA operates on functionally
meaningful blocks, it provides a natural entry point for studying block-level replacement,
recombination, insertion, deletion, and convergence in BGC evolution. At the same time, it
helps distinguish true biosynthetic novelty from cases that appear novel only because
sequence divergence obscures underlying functional similarity. In siderophores, this
framework also provides an upstream foundation for linking biosynthesis to receptor
specificity, resource sharing, and microbial interaction networks. With the emergence of
structure prediction tools such as AlphaFold[21], structural constraints and active-site
features are increasingly becoming a more unified functional scale than full-length
sequence similarity alone, consistent with our results for NIS pathways. The broader
implication is that the signals most informative for cross-species comparison do not reside
primarily in full-length similarity, but in substrate-recognition sites, active-site-associated

features, and structurally constrained functional modules.

The resulting Siderophore Atlas extends this functional-space comparison to a global
scale. Its value lies not in listing siderophore BGCs per se, but in organizing siderophore
biosynthesis through functional typing rather than conventional sequence clustering. Our
results show that, although siderophore biosynthesis is broadly prevalent across bacterial
genomes, this prevalence does not imply simple conservation across higher taxonomic
ranks. Instead, bacterial groups differ strikingly in their biosynthetic strategies, internal
biosynthetic profiles, and taxonomic spread. This is consistent with broader views of
siderophores as components of iron competition, host-associated interactions, and
ecological networks, rather than merely as isolated iron chelators. Thus, the Siderophore
Atlas is not a catalog organized by sequence clustering, but a global framework organized
by functional equivalence, enabling siderophore biosynthetic potential to be compared
across phylogenetically diverse bacteria on a unified scale.

At the macro-evolutionary level, our results further suggest that NRPS siderophores and
NIS siderophores follow contrasting organizational regimes. NRPS siderophores exhibit a
long-tailed, innovation-rich architecture, consistent with continuous diversification driven
by modular recombination and local innovation. By contrast, NIS siderophores are
dominated by a relatively small number of highly disseminated pathways, suggesting a
dissemination-oriented architecture in which a few successful solutions account for much
of the cross-lineage spread. This contrast is consistent with prior observations of
recombination-driven NRPS diversification and siderophore pathway replacement or
dissemination across bacterial lineages. In light of our taxonomic spread analysis, we
interpret these patterns as evidence for at least two different biosynthetic
macro-evolutionary strategies for coping with iron limitation: one based on continued
expansion of chemical space through modular innovation, and another based on repeated
dissemination of a limited set of highly transferable pathways. In this sense, siderophore
diversity is not simply an accumulation of molecular variants, but a reflection of how
different biosynthetic systems balance innovation against dissemination.

Several limitations should also be acknowledged. First, both SideroBank and the
Siderophore Atlas remain constrained by currently sequenced genomes, available
references, and annotation quality, and therefore cannot capture the full extent of
siderophore diversity. Second, the functional equivalence recovered by BBA remains an
inference based on biosynthetic features rather than direct chemical or genetic validation
for every family. Third, BBA was developed and validated primarily in siderophore
pathways; although its principles should be extensible to PKS and other modular natural
product systems, the definition of blocks, extraction of functional features, and alignment
rules will need to be adapted for each biosynthetic class. Even so, this study points to two
broader methodological transitions that are now becoming feasible in natural product
genomics: Sidero-Mining, aided by an LLM, converts fragmented biosynthetic knowledge
in literature space into a computable reference space, whereas BBA further converts BGC
comparison from a phylogeny-constrained sequence space into a more biosynthetically

meaningful functional space. As structure prediction, substrate characterization, chemical
annotation, and ecological datasets continue to improve, such a functional-space
genomics framework may provide a more general route for linking gene clusters,
molecular products, and community-level function.
653
654
655

Methods

Literature mining and construction of the SideroBank dataset

To construct a high-confidence cross-species siderophore knowledgebase, we
first performed large-scale literature mining using the Semantic Scholar
Dataset. By querying the Semantic Scholar API with the keyword *"siderophore"*
and restricting publication years to post-1980 studies, we retrieved a total of
12,004 potentially relevant publications.

To evaluate the performance of different models in extracting
siderophore-related information, we manually constructed a benchmark
dataset consisting of 60 articles, including equal numbers of positive articles
containing siderophore-related information and negative articles without such
information. To reduce potential biases arising from publication period, specific
siderophore names, or recurring cue words, the benchmark set was designed
to include articles from different time periods and a diverse range of
siderophore systems. Ground-truth labels were manually curated from these
articles. Model outputs were then classified as true positives, false positives,
true negatives, or false negatives, and overall precision, recall, and F1-score
were calculated.

To improve the accuracy and consistency of literature extraction, we
developed the Sidero-Mining pipeline with carefully designed task-specific
prompts. Each prompt generally consisted of three components: (i) explicit
specification of the extraction task, such as extracting siderophore names or
producer species; (ii) a structured output template in Markdown format to
facilitate downstream parsing, verification, and conversion into more visually
organized formats such as XMind; and (iii) constraint instructions designed to
suppress hallucinations, for example by requiring the model to return "null" for
unavailable information and to avoid generating content outside the predefined
template.

For siderophore biosynthesis extraction, we further implemented a multi-step
reasoning workflow designed to mimic manual literature interpretation. In the
first step, the model extracted all genes related to the target siderophore and

summarized their putative biosynthetic functions. In the second step, these
genes and their functional descriptions were used to infer the biosynthetic
class of the siderophore. In the third step, the predicted biosynthetic type and
the previously extracted gene set were jointly used to identify genes directly
involved in siderophore synthesis. In the fourth step, the refined
synthesis-gene information was combined with the original article content to
extract substrate-level biosynthetic details. By decomposing a complex
synthesis-mining task into a sequence of connected subtasks, this workflow
enabled progressive reasoning and improved the accuracy of
biosynthesis-related information extraction.

For large-scale screening, literature filtering was performed in two stages. In
the first stage, titles and abstracts of the retrieved articles were input into
gpt-4o-mini (gpt-4o-mini-2024-07-18), and six predefined prompts were used
to classify articles into major topic categories, including siderophore chemical
structure, biosynthesis, ecology, pathology, environmental and agricultural
relevance, and metal-ion transport. For ease of downstream statistics, the
model was instructed to return only binary "Y" or "N" outputs.

In the second stage, articles retained after title-and-abstract screening were
further evaluated at the full-text level. Full texts were downloaded
automatically using the links provided in the Semantic Scholar Dataset
whenever possible, and the remaining articles were collected manually. Full
texts were then processed using **gpt-4o-mini (gpt-4o-mini-2024-07-18)** with
seven additional prompts to determine whether each article contained
information related to single-gene studies, genomic studies, siderophore
transport, iron-affinity measurements, siderophore receptors, chemical
structure characterization, or microbial interactions. Articles primarily focused
on siderophore synthesis and biosynthetic genes were retained as the final
literature set for siderophore information mining.

Genome retrieval and BGC annotation

After extracting siderophore-related information from the literature, we
retrieved the corresponding genome data based on the strain information
reported in the mined articles. To this end, we developed an automated
genome retrieval workflow using the NCBI API. By querying strain names
extracted from the literature, the program automatically downloaded the
corresponding genome files from the NCBI Dataset. For strains that could not
be directly matched due to renaming, taxonomic updates, or redundant
strain-name annotations in the original articles, the relevant genome
assemblies were collected manually from the NCBI website.

Once genome files were obtained, siderophore biosynthetic gene clusters
were annotated using a custom annotation workflow based on antiSMASH 7.0
with default parameters. These annotations served as the primary basis for
downstream BGC curation and comparison.

Dataset collection and preprocessing

Manually curated cross-species BGC dataset
We first collected literature-reported records of siderophore distribution across species
and assembled a cross-species reference dataset of siderophore biosynthetic gene
clusters (BGCs). For each reported strain, the corresponding genome data were further
examined and candidate BGCs were annotated. These annotations were then manually
validated according to antiSMASH product-class assignments and the internal
organizational structure of each BGC, to ensure consistency between the collected
clusters and their reported siderophore products.
738
Predicted protein structure dataset
To establish a structural reference dataset for NIS biosynthetic enzymes, we queried
InterPro using the Pfam entry corresponding to the IucA/IucC domain (IucA_IucC,
PF04183) as the key domain and retrieved approximately 14,000 protein entries
associated with AlphaFold2-predicted structures. Metadata for these entries, including the
identifiers of their corresponding AlphaFold2 models, were obtained through the InterPro
download API. The corresponding predicted protein structures were then downloaded
through the AlphaFold database API.
Based on this dataset, amino acid sequences were extracted and compiled into a
searchable DIAMOND sequence database, which was used in subsequent analyses to
match query NIS biosynthetic enzymes to reference sequences and structures.

BGC Block Aligner algorithm and parameter settings

Definition of core biosynthetic domains
For NRPS pathways, the set of domains used for BGC comparison was defined based on
the antiSMASH annotation framework for NRPS core components. This set included both
the major catalytic domain subtypes that constitute canonical NRPS modules and
additional domains involved in modification or accessory biosynthetic functions. These
domain definitions were used for downstream module decomposition, feature extraction,
and scoring-matrix construction.
For NIS pathways, the analysis focused on the key NIS biosynthetic enzymes responsible
for siderophore assembly, which served as the basic units for feature extraction and
similarity comparison.
761
Extraction features from NRPS A domains

Because adenylation (A) domains are the primary determinants of substrate specificity in
NRPS pathways, we extracted multiple types of sequence-derived features from A
domains for downstream similarity calculation. First, the 34-amino-acid (34AA) active-site
feature sequences were extracted using a modified implementation based on the source
code of AmenPredictor. In this procedure, protein sequences were aligned against a given
hidden Markov model using HMMER2, and residues corresponding to predefined
active-site positions in the template were extracted as feature residues.
770
In addition, we extracted the 27-amino-acid (27AA) specificity-conferring features defined
in DeepAden, enabling an alternative representation of the substrate-determining region
of each A domain. To further capture broader contextual information beyond manually
defined active-site residues, we also obtained embedding features for each A domain
using the DeepAden framework. These embeddings provide a continuous representation
of sequence-level functional properties and were used as an additional feature space for
pairwise comparison.
778
After feature extraction from all A domains in the BGCs to be compared, redundant feature
sequences were removed using DIAMOND or CD-HIT where applicable. Pairwise
similarity between A domains was then quantified on the basis of sequence identity
between the extracted 34AA or 27AA feature sequences, or by similarity measures
computed from the corresponding embedding representations. These similarity values
were subsequently used for scoring-matrix construction.
785

Feature extraction for NIS biosynthetic enzymes
For NIS pathways, we first constructed a DIAMOND sequence database from the
predicted protein structure dataset described above. Query NIS biosynthetic enzyme
sequences were searched against this database, and the highest-scoring hit passing the
predefined threshold was selected and mapped to its corresponding AlphaFold2-predicted
structure. When no related sequence could be identified under the preset threshold of 0.9,
the query structure was predicted using ESMFold.
After obtaining the predicted structures of the query NIS enzymes, pairwise structural
alignments were performed between query and reference proteins using gtalign. Based
on the active-site positions defined in the reference templates, the corresponding residues
in the aligned query sequences were extracted as active-site feature sequences for
downstream similarity calculation.
Different reference proteins were used for different classes of NIS biosynthetic enzymes,
including DesD (PDB: 7tgm), lucA (PDB: 5jm8), SbnC (PDB: 7cbb), and AsbB (PDB: 3to3).
For each reference, the active-site residue positions were defined according to the
corresponding structural template and used to extract the aligned residues from the query
proteins.
Construction of the scoring matrices
For NRPS pathways, a unified scoring matrix was constructed by combining A domains
with other biosynthetically relevant domains. For non-A domains, a match between
domains of the same family and the same subtype was assigned a score of 0.2, whereas

a match between domains of the same family but different subtypes was assigned a score
of 0.12. A mismatch between domains from different families was penalized by -0.1.
Because A domains carry the most important functional information, their match scores
were not assigned as fixed constants but were instead derived directly from the pairwise
similarity matrix of A-domain feature sequences, or from transformed values based on that
matrix, with scores constrained to the range of 0 to 1. To prevent biologically implausible
matches between A domains and non-A domains, such matches were penalized by -0.5.
The final NRPS scoring matrix therefore integrated both A-domain similarities and scores
for other biosynthetic domains.
816
For NIS pathways, the relative order of biosynthetic enzymes was not explicitly
considered. Instead, the analysis focused on similarity relationships among NIS enzymes
within each pair of BGCs. Active-site feature sequences extracted from each structural
template were used to generate separate similarity matrices, and for each enzyme pair,
the maximum similarity value across these matrices was retained as the final similarity
score. This final similarity matrix, with values ranging from 0 to 1, was then used directly,
or after optional transformation, as the scoring matrix for downstream optimization.
824
Generation of pathway arrangements and BGC comparison
For NRPS pathways, all NRPS modules were treated as the basic units for arrangement
and comparison. Under the exhaustive arrangement mode, when the number of module
units was below a predefined threshold, such as five, all possible permutations of the
modules were generated and evaluated. Considering the linear biosynthetic logic of
NRPS systems, adjacent modules located within the same operon and oriented in the
same transcriptional direction could be merged into a single unit, thereby reducing the
size of the permutation space and minimizing fragmentation of contiguous module blocks.
When the number of module units exceeded the threshold, a greedy strategy was used
instead. Specifically, the BGC containing the largest number of modules was selected as
the reference BGC, and its module units were fixed according to their genomic order. As in
the exhaustive mode, adjacent and co-directional modules within the same operon could
be merged into a single unit. For each query BGC, all module units were extracted and
locally scored against different regions of the reference arrangement using the predefined
scoring matrix. Candidate segments were then iteratively placed according to their local
scores, and penalties were applied when newly placed segments overlapped with regions
that had already been assigned in previous steps.
842
After the arrangement step, overall NRPS BGC comparison was performed on the
concatenated domain sequences using the Biopython pairwise aligner under the specified
scoring matrix. In the exhaustive arrangement mode, all possible arrangements were
evaluated and the highest-scoring alignment was retained as the final result. In the greedy
mode, the alignment score of the assembled arrangement was used as the final similarity
score for the BGC pair.
849

For NIS pathways, no direct global ordering of biosynthetic enzymes was imposed.
Instead, the method evaluated the quality of correspondence between the sets of NIS
enzymes in the two BGCs. Based on the pairwise similarity scoring matrix described
above, the Hungarian algorithm was used to identify the optimal enzyme matching
between the two BGCs. The resulting total matching score was then normalized by the
larger number of NIS enzymes present in either BGC, thereby reducing the effect of
copy-number differences on the final similarity estimate.
857
BiG-SCAPE parameter settings
To benchmark the results of BGC Aligner against an established BGC comparison
framework, the same set of siderophore BGCs was also analyzed using BiG-SCAPE.
BiG-SCAPE was run with Pfam version 34, using the --no_classify option and --mix mode.
Computations were performed with 32 CPU cores, and the cutoff was set to 1 to retain all
pairwise comparison results. All other parameters were kept at their default settings. The
raw similarity values output by BiG-SCAPE were extracted and further assembled into
complete pairwise similarity matrices for downstream analyses.
866
867

Author contributions

Jiqi Shao: Conceptualization; data curation; algorithm development; analysis;
visualization; writing - original draft. Yanzhao Wu: Data curation; visualization; coding and
analysis; writing - original draft. Shizhen Tian: Data curation. Haohua Luo: Data curation.
Yuanzhe Shao: Data curation. Linlong Yu: Data curation. Guanyue Xiong:Data curation.
Ruolin He: Data curation. Ruichen Xu: Data curation. Peng Guo: Data curation. Nan
Rong:Data curation. Shaohua Gu: Conceptualization; funding acquisition. Zhiyuan Li:
Conceptualization; funding acquisition; analysis; visualization; writing - original draft.
876
877

878 **Declaration of Interests**

879 The authors declare no competing interests.
880
881

Reference

1.   Medema, M.H., and Fischbach, M.A. (2015). Computational approaches to natural
product discovery. Nat Chem Biol *11*, 639-648. 10.1038/nchembio.1884.
2.   Kramer, J., Özkaya, Ö., and Kümmerli, R. (2020). Bacterial siderophores in community
and   host   interactions.   Nature   Reviews   Microbiology   *18*,   152-163.
10.1038/s41579-019-0284-4.
3.   Carroll, C.S., and Moore, M.M. (2018). Ironing out siderophore biosynthesis: a review
of non-ribosomal peptide synthetase (NRPS)-independent siderophore synthetases.
Crit Rev Biochem Mol Biol *53*, 356-381. 10.1080/10409238.2018.1476449.
4.   Wei, Z., Gu, S., Vollenweider, V., Zuo, Y., Li, Z., and Kümmerli, R. (2025). Microbial
siderophores   for   One   Health.   Trends   in   Microbiology   *33*,   1277-1285.
10.1016/j.tim.2025.05.002.
5.   Blin, K., Shaw, S., Augustijn, H.E., Reitz, Z.L., Biermann, F., Alanjary, M., Fetter, A.,
Terlouw, B.R., Metcalf, W.W., Helfrich, E.J.N., et al. (2023). antiSMASH 7.0: new and
improved predictions for detection, regulation, chemical structures and visualisation.
Nucleic Acids Research *51*, W46-W50. 10.1093/nar/gkad344.
6.   Hannigan, G.D., Prihoda, D., Palicka, A., Soukup, J., Klempir, O., Rampula, L., Durcak,
J., Wurst, M., Kotowski, J., Chang, D., et al. (2019). A deep learning genome-mining
strategy   for   biosynthetic   gene   cluster   prediction.   Nucleic   Acids   Research   *47*,
e110-e110. 10.1093/nar/gkz654.
7.   Lai, Q., Yao, S., Zha, Y., Zhang, H., Zhang, H., Ye, Y., Zhang, Y., Bai, H., and Ning, K.
(2025). Deciphering the biosynthetic potential of microbial genomes using a BGC
language processing neural network model. Nucleic Acids Research *53*, gkaf305.
10.1093/nar/gkaf305.
8.   Zdouc, Mitja M., Blin, K., Louwen, Nico L.L., Navarro, J., Loureiro, C., Bader,
Chantal D., Bailey, Constance B., Barra, L., Booth, Thomas J., Bozhüyük, Kenan A.J.,
et al. (2024). MIBiG 4.0: advancing biosynthetic gene cluster curation through global
collaboration. Nucleic Acids Research *53*, D678-D690. 10.1093/nar/gkae1115.
9.   Kautsar, S.A., van der Hooft, J.J.J., de Ridder, D., and Medema, M.H. (2021).
BiG-SLiCE: A highly scalable tool maps the diversity of 1.2 million biosynthetic gene
clusters. GigaScience *10*, giaa154. 10.1093/gigascience/giaa154.
10.   Navarro-Muñoz, J.C., Selem-Mojica, N., Mullowney, M.W., Kautsar, S.A., Tryon, J.H.,
Parkinson, E.I., De Los Santos, E.L.C., Yeong, M., Cruz-Morales, P., Abubucker, S., et
al. (2020). A computational framework to explore large-scale biosynthetic diversity.
Nature Chemical Biology *16*, 60-68. 10.1038/s41589-019-0400-9.
11.   Draisma, A., Loureiro, C., Louwen, N.L.L., Kautsar, S.A., Navarro-Muñoz, J.C.,
Doering, D.T., Mouncey, N.J., and Medema, M.H. (2026). BiG-SCAPE 2.0 and
BiG-SLiCE 2.0: scalable, accurate and interactive sequence clustering of metabolic
gene clusters. Nature Communications *17*, 2000. 10.1038/s41467-026-68733-5.
12.   Agrawal, P., Khater, S., Gupta, M., Sain, N., and Mohanty, D. (2017). RiPPMiner: a
bioinformatics resource for deciphering chemical structures of RiPPs based on
prediction of cleavage and cross-links. Nucleic Acids Research *45*, W80-W88.
10.1093/nar/gkx408.

13. Helfrich, E.J.N., Ueoka, R., Dolev, A., Rust, M., Meoded, R.A., Bhushan, A., Califano,
G., Costa, R., Gugger, M., Steinbeck, C., et al. (2019). Automated structure prediction
of trans-acyltransferase polyketide synthase products. Nat Chem Biol *15*, 813-821.
10.1038/s41589-019-0313-7.

14. Calcott, M.J., Owen, J.G., and Ackerley, D.F. (2020). Efficient rational modification of
non-ribosomal peptides by adenylation domain substitution. Nat Commun *11*, 4554.
10.1038/s41467-020-18365-0.

15. Stachelhaus, T., Mootz, H.D., and Marahiel, M.A. (1999). The specificity-conferring
code of adenylation domains in nonribosomal peptide synthetases. Chem Biol *6*,
493-505. 10.1016/s1074-5521(99)80082-9.

16. Röttig, M., Medema, M.H., Blin, K., Weber, T., Rausch, C., and Kohlbacher, O. (2011).
NRPSpredictor2—a web server for predicting NRPS adenylation domain specificity.
Nucleic Acids Research *39*, W362-W367. 10.1093/nar/gkr323.

17. Rausch, C., Weber, T., Kohlbacher, O., Wohlleben, W., and Huson, D.H. (2005).
Specificity prediction of adenylation domains in nonribosomal peptide synthetases
(NRPS) using transductive support vector machines (TSVMs). Nucleic Acids Res *33*,
5799-5808. 10.1093/nar/gki885.

18. Mongia, M., Baral, R., Adduri, A., Yan, D., Liu, Y., Bian, Y., Kim, P., Behsaz, B., and
Mohimani, H. (2023). AdenPredictor: accurate prediction of the adenylation domain
specificity of nonribosomal peptide biosynthetic gene clusters in microbial genomes.
Bioinformatics *39*, i40-i46. 10.1093/bioinformatics/btad235.

19. Yang, J., Banas, V.S., Patel, K.D., Rivera, G.S.M., Mydy, L.S., Gulick, A.M., and
Wencewicz, T.A. (2022). An acyl-adenylate mimic reveals the structural basis for
substrate recognition by the iterative siderophore synthetase DesD. Journal of
Biological Chemistry *298*. 10.1016/j.jbc.2022.102166.

20. He, R., Gu, S., Xu, J., Li, X., Chen, H., Shao, Z., Wang, F., Shao, J., Yin, W.-B., Qian, L.,
et al. (2024). SIDERITE: Unveiling hidden siderophore diversity in the chemical space
through digital exploration. iMeta *3*, e192. https://doi.org/10.1002/imt2.192.

21. Jumper, J., Evans, R., Pritzel, A., Green, T., Figurnov, M., Ronneberger, O.,
Tunyasuvunakool, K., Bates, R., Žídek, A., Potapenko, A., et al. (2021). Highly
accurate protein structure prediction with AlphaFold. Nature *596*, 583-589.
10.1038/s41586-021-03819-2.

22. Bruns, H., Crüsemann, M., Letzel, A.C., Alanjary, M., McInerney, J.O., Jensen, P.R.,
Schulz, S., Moore, B.S., and Ziemert, N. (2018). Function-related replacement of
bacterial siderophore pathways. Isme j *12*, 320-329. 10.1038/ismej.2017.137.

960