Cap-adjacent 2`-*O*-ribose methylation of RNA in *C. elegans* is required for

postembryonic growth and germline development in the presence of the

decapping exonuclease EOL-1

Eileen Clemens[1], Sarah Brivo[1,6], Mohammed Al-Khafaji[1], Peter Eijlers[1], Maheshika

Kurukulasuriya[1], Irmgard U. Haussmann[2], David MacLeod[1,7], Marius Wenzel[3],

Berndt Müller[1], Matthias Soller[4,5] and Jonathan Pettitt[1]*

1. School of Medicine, Medical Sciences and Nutrition, Institute of Medical Sciences,
University of Aberdeen, Aberdeen, AB25 2ZD, United Kingdom

2. College of Life Science, Birmingham City University, Birmingham, B15 3TN,
United Kingdom

3. School of Biological Sciences, University of Aberdeen, Aberdeen AB24 2TX,
United Kingdom

4. School of Biosciences, College of Life and Environmental Sciences, University
of Birmingham, Edgbaston, Birmingham, B15 2TT, United Kingdom

5. Division of Molecular and Cellular Function, School of Biological Sciences,
Faculty of Biology, Medicine and Health, University of Manchester, Manchester,
M13 9PT, United Kingdom

6. Current address: Department of Pathology and Immunology, University of
Geneva, CMU - Rue Michel-Servet 1, CH-1211 Geneva 4, Switzerland

7. Current address: Wellcome Trust Centre for Cell Biology, Michael Swann
Building, Max Born Crescent, Edinburgh EH9 3BF, United Kingdom

* Corresponding author: Tel +44 1224 437516; Email j.pettitt@abdn.ac.uk

ABSTRACT

Cap-adjacent 2`-*O*-ribose methylation (cOMe) of the first two transcribed nucleotides of RNA polymerase II transcripts is a conserved feature in many eukaryotes. In mammals, these modifications are key to a transcript surveillance system that regulates the interferon response, but the broader functions of cOMe remain poorly understood. To understand the role of cOMe in *C. elegans*, we functionally characterised the methyltransferases (CMTR-1 and CMTR-2) responsible for installing these modifications. These enzymes have distinct expression patterns, protein interaction partners, and loss of function phenotypes. Loss of CMTR-1 causes dramatic reductions in cOMe, impaired growth and sterility. In contrast, animals lacking CMTR-2 are superficially wild-type, though CMTR-2 loss enhances the severity of the *cmtr-1* mutant phenotype. Depletion of CMTR-1 causes downregulation of transcripts associated with germline sex determination and upregulation of those involved in the intracellular pathogen response (IPR). We show that absence of the decapping exonuclease, EOL-1, an IPR component, completely suppresses the sterility and growth defects caused of loss of CMTR-1, suggesting that EOL-1 degrades cellular transcripts lacking cOMe. Our work shows the physiological relevance of cOMe in protecting transcripts from decapping exonucleases, raising the possibility that cOMe plays a role in RNA-mediated immune surveillance beyond the vertebrates.

45 **GRAPHICAL ABSTRACT**

INTRODUCTION

Cap-adjacent 2`-*O*-ribose methylation (cOMe) of RNA was first discovered in mammalian cells several decades ago (Adams and Cory, 1975; Anreiter et al., 2023; Furuichi et al., 1975; Galloway and Cowling, 2019). Subsequent work showed that these modifications are found throughout the Metazoa, as well as in the viruses and protists that infect these organisms. In animals, the first two transcribed nucleotides can harbour cOMe, with methylation of the first transcribed nucleotide being designated "cap1", and methylation of both first and second nucleotides referred to as "cap2". Although these modifications are ubiquitous, their physiological significance remains incompletely understood.

The first defined and the best understood role for cOMe is in preventing activation of an RNA-based immune response (Daffis et al., 2010; Li et al., 2013; Schuberth-Wagner et al., 2015; Züst et al., 2011): an antiviral mechanism consisting of sensor proteins that recognise transcripts lacking cOMe, which brings about the activation of the interferon response. This has created strong selective pressure for viruses to evolve the means to ensure their transcripts are modified by cOMe (Hyde and Diamond, 2015). However, this function does not explain the presence of cOMe in animals that lack the interferon pathway. It also doesn't explain the full significance of cOMe in mammals: loss of either of the two highly conserved methyltransferases, CMTR1 and CMTR2, responsible for cOMe in mice results in embryonic lethality without inducing the interferon pathway (Dohnalkova et al., 2023); and there are multiple lines of evidence supporting diverse roles for cOMe, including the regulation of transcription, RNA splicing, and translation (Dönmez et al., 2004; Drazkowska et al., 2022; Haussmann et al., 2022; Kuge et al., 1998; Liang et al., 2022; Yu et al., 1998).

Studying cOMe in genetically tractable model organisms such as *Drosophila*

melanogaster and *C. elegans* should allow us to understand the functional

significance of cOMe from a whole animal perspective. Both organisms have

orthologues of mammalian CMTR1 and CMTR2 (Dix et al., 2022; Haussmann et al.,

2022). However, detailed functional genetic analysis in *Drosophila* has been

hampered by the relatively subtle phenotypic consequences of loss of cOMe.

Drosophila CMTr1/2 double mutants are viable, though they show specific

behavioural defects consistent with defects in the spatial regulation of translation in

neurons (Haussmann et al., 2022).

Previous studies have shown that subcellular mislocalisation of *C. elegans* CMTR-

1can affect the expression of specific genes (Meisel et al., 2024). Here we present a

comprehensive analysis of CMTR function, showing that loss of cOMe in *C. elegans*

leads to growth retardation and pronounced defects in germline development, with

cmtr-1 loss-of-function mutants conferring a fully penetrant sterile phenotype. Loss of

cOMe leads to activation of a set of genes that constitute the intracellular pathogen

response (IPR), which are induced by viruses and intracellular pathogens (Lažetić et

al., 2023), raising the possibility that as in mammals, there exists a surveillance

mechanism to detect transcripts lacking these modifications. From a forward genetic

screen for suppressors of *cmtr-1* loss-of-function we identified the decapping

exonuclease EOL-1, showing that this enzyme is a major component of the cellular

response to transcripts lacking cOMe. Our work suggests evolutionary continuity in

the role of cOMe in RNA immune surveillance and provides physiological

confirmation of the role of this modification in protecting transcripts from degradation

by the DXO/Rai decapping exonuclease family.

MATERIALS AND METHODS

Nematode Strains

C. elegans strains were grown using standard culture conditions (Stiernagle, 2006) at 20°C unless otherwise indicated. N2 (Bristol) was used as the wild type. The following previously described strains were used in this study: HML1012 (Hills-Muckey et al., 2022); CGC45; CGC66. Strains generated in the current study are described in Supplementary Table 1.

The *cmtr-1* deletion allele, *syb3613*, was generated by Fujian Sunybiotech Co. LTD, and obtained as strain PHX3613. The *cmtr-2* deletion allele, *tm4453*, was obtained from the Mitani Lab as strain FX4453. PE1000 was generated from FX4453 by outcrossing six times against N2. Fluorescent-tag/degron knock-in strains were created as described previously (DeMott et al., 2021; Dickinson et al., 2015; Hills-Muckey et al., 2022). The CMTR-1(K244A) mutation was generated through oligonucleotide-templated repair of CRISPR/Cas9 induced double-strand breaks, using the previously described *dpy-10* based co-conversion strategy (Arribere et al., 2014). CRISPR alleles of *eol-1* were made using the same *dpy-10*-based approach, but the injections were performed using PE1176 animals and the resultant F1 progeny grown on 5-Ph-IAA containing plates, selecting for suppression of the sterile phenotype. Alleles were confirmed by Sanger sequencing (Eurofins Genomics). Oligonucleotides used to verify the allelic status of genes, generate guide RNA constructs, and produce homologous repair templates are given in Supplementary Table 1.

Transgenic rescuing assay

The wild-type *cmtr-1* cDNA was amplified from total RNA using primers cmcDNAFo

and cmcDNARev (Supplementary Table 1). The plasmid backbone, which contains

the *rps-0* promoter and *unc-54* 2`-O- UTR, was amplified using primers prpo-u54Fwd

and prpo-u54Re (Supplementary Table 1), and the *cmtr-1* cDNA cloned between the

rps-0 promoter and *unc-54* 3`UTR using the NEBuilder HiFi DNA Assembly kit (NEB)

to create plasmid pSB2. The plasmid was verified by Sanger Sequencing (Eurofins

Genomics). PE1033 worms were co-injected with pSB2 and a plasmid containing

myo-2p::tdTomato (Eijlers et al., 2024). Transgenic lines were established as

previously described (Mello et al., 1991; Nance and Frøkjær-Jensen, 2019). The

rescue assay was performed by picking non-tdTomato offspring as L1/L2 larvae to

separate plates and scoring their ability to reach fertile adulthood.

Fluorescent microscopy

134 Worms were mounted in 5 µl M9 supplemented with 10 mM sodium azide on 5%

135 agar pads. Images were obtained using either a Zeiss Axioplan 2, equipped with a

136 Hamamatsu Orca ER camera, or a Zeiss Imager M2 upright microscope, equipped

137 with a Hamamatsu Flash 4 LT camera.

Western blot

Western blots were performed as described previously (Fasimoye et al., 2022) using

lysates from 300 – 500 adult/L4 larvae, or 5000 L1 larvae per well. To detect

embryonic expression of CMTR-1/-2 in embryos, 40 µg of total protein from

embryonic extracts was loaded per gel lane. For detecting larval expression, GFP-

tagged proteins were detected with mouse anti-GFP antibodies (Roche

11814460001), mNeonGreen tagged proteins with rabbit anti-mNeonGreen

antibodies (Proteintech, 29523-1-AP) and GAPDH using mouse anti-GAPDH

antibodies (Invitrogen AM4300). Secondary antibodies used were anti-mouse or anti-

rabbit HRP conjugated IgG antibody (Cell Signalling Technology 7074 and 7076). To

detect proteins in embryonic extracts, GFP tag Polyclonal antibody (Proteintech,

50430-2-AP) (1:1000 dilution) was used together with the above anti-rabbit

secondary (1:3000 dilution). Blots were also stained with amidoblack and quantified

using either Fiji (Schindelin et al., 2012)or ImageQuant (Cytiva) software (Fasimoye

et al., 2022).

Preparation of *C. elegans* embryo extracts

C. elegans embryo extracts were prepared essentially as described previously

(Eijlers et al., 2024)(see also Supplementary Methods). Extracts were treated with

RNase as described previously (Eijlers et al., 2024).

Cap-methylation status analysis of mRNA

Total RNA was extracted with Trizol (Sigma) according to the manufacturers

description using 20 µg glycogen (Roche) for precipitation. poly(A) mRNA was

prepared using the NEBNext® Poly(A) mRNA Magnetic Isolation Module (NEB) by

double oligo dT selection according to the manufacturer's instructions.

For the analysis of 5` cap structures, 5 µg total RNA was used for decapping by

yDcpS in 20 µl for 1 h at 37°C according to the manufacturer's instruction (NEB). The

RNA was extracted by phenol/CHCl$_3$ and ethanol precipitated in the presence of

glycogen. The RNA was then labeled in a total volume of 20 µl containing 2 µl

capping buffer (NEB), 1 µl SAM (2 mM), 0.25 µl ^{32}P-αGTP (3000 Ci/mmol, 6.6 µM;

Hartmann Analytics), 0.5 µl RNase Protector (Roche) and 0.5 µl capping enzyme

(NEB) by incubation for 1 h at 37°C. The volume was then increased to 50 µl with

water and poly(A) RNA selected as described above. The RNA was then digested on

the beads in 5 µl using 0.5 µl NEB buffer 3 and 0.5 µl RNase I for 2 h, and then 10 µl

gel loading buffer was added (98% deionized formamide,10 mM EDTA, 0.025%

xylene cyanol FF and 0.025% bromphenol blue); products were analysed on 22%

denaturing polyacrylamide gels (National Diagnostics) and pre-run for 2 h. Gels were

soaked in 20% PEG400, 10% acetic acid and 40% methanol for 10 min and then

dried on a Whatman 3MM paper. Dried gels were then exposed to a storage

phosphor screen (Bio-Rad) and scanned by a Molecular Imager FX in combination

with QuantityOne software (Bio-Rad). Markers were prepared as described

previously (Dix et al., 2022; Haussmann et al., 2022).

For the analysis of the first nucleotide in mRNA, 5 µg poly(A) mRNA was purified as

described above and 5 µl incubated with terminator nuclease (Epicenter), according

to the manufacturer's instructions, to remove rRNA followed by phenol/CHCl$_3$ and

ethanol precipitation in the presence of glycogen (Roche). The mRNA was then

decapped using RppH (NEB) and dephosphorylated by Antarctic phosphatase (NEB)

in NEB buffer 2 supplemented with 0.1 mM ZnCl$_2$ in 20 µl. Then the RNA was

extracted by phenol/CHCl$_3$ and precipitated in the presence of glycogen. The 5`-ends

of dephosphorylated mRNAs were then labelled using 10 units of T4 PNK (NEB) and

0.25 µl ^{32}P-γATP (6000 Ci/mmol, 12.5 µM; Hartmann Analytics). The labelled RNA

was precipitated, and resuspended in 10 µl of 50 mM sodium acetate buffer (pH 5.5)

and digested with nuclease P1 (SIGMA) for 2 h at 37°C. Two microliters of each

sample was loaded on cellulose F TLC plates (20x 20 cm; Merck) and run in a

solvent system of isobutyric acid:0.5 M NH_4OH (5:3, v/v, solvent A), as first

dimension, and isopropanol:HCl:water (70:15:15, v/v/v, solvent B) or sodium

phosphate, 0.1 M (pH 6.8), ammonium sulfate, n-propanol (100:60:2, v/w/v, solvent

C), as the second dimension. TLCs were repeated from biological replicates. The

identity of the nucleotide spots was determined as described (Keith, 1995; Kruse et

al., 2011). TLCs were exposed to a storage phosphorscreen (Bio-Rad) and scanned

by a Molecular Imager FX in combination with QuantityOne software (Bio-Rad).

Monitoring SL1 *trans*-splicing using quantitative PCR

RNA was isolated from embryonic extracts prepared from PE1176 and PE1220

embryos, treated for 4 h with either 50 μM 5-Ph-IAA-AM in DMSO, or control treated

in DMSO by extraction with TriZol, and then purified using the PureLink RNA Mini kit

(Life Technologies), with modifications for TRIzol treated samples and DNase

treatment, as described by the manufacturer. Reverse transcription and analysis by

qPCR were done as described (Eijlers et al, 2024, Philippe et al, 2017). Analyses

were done as three technical replicates. Shown are ΔC_T values derived for each

replicate (Schmittgen and Livak, 2008).

***In vitro* spliced leader *trans*-splicing assays.**

For extract preparation, PE1176 and PE1220 animals were grown in liquid culture

and embryos were isolated and treated with 50 μM 5-PH-IAA-AM or control-treated

with DMSO as described (Eijlers et al., 2024), except that treatment was for 4h.

Extracts were then prepared as described (Eijlers et al., 2024) and stored at -80°C.

The synthetic SL *trans*-splicing substrate, consisting of *rps-3* genomic DNA

containing 404 bp of 5` UTR sequence (including 382 bp of outron sequence

upstream of the known SL1 *trans*-splice site) and extending 266 base pairs

downstream from the ATG translation start site, was amplified from *C. elegans*

genomic DNA using primers *rps-3*genFwd and *rps-3*genRev (Supplementary Table

1). The resulting amplicon was inserted into pBlueScript KS(-) cleaved with XbaI

using NEB HIFI gene assembly to produce pBS-*rps-3*. The sequence was confirmed

by Sanger sequencing (Eurofins Genomics). pBS-*rps-3* DNA linearised with EcoRI

was transcribed with T7 RNA polymerase using the MEGAscript® kit (Invitrogen).

This produces a synthetic *rps-3* RNA with a 3`-end derived from pBlueScript KS(-)

that serves as a primer binding site in PCR. RNA was extracted using

phenol/chloroform, concentrated by ethanol precipitation, resuspended in RNase-

free water and purified using MicroSpin G-25 columns (Cytiva). RNA concentration

was determined by measuring absorption at 260 nm assuming 1 AU = 40 µg/ml.

In vitro SL1 *trans*-splicing reactions were done essentially as described (Lasda et al.,

2011). 15 µl reactions were done in 10 mM Tris-HCl (pH 8.0), 60 mM KCl, 4 mM

$MgCl_2$, 2 mM ATP, 20 mM creatine phosphate, 50 µg/ml creatine kinase, 2 mM DTT,

3 % PEG 8000, 0.25 mM EDTA, 5% glycerol, 0.5 mM PMSF, 2 U/µl RNaseOUT, 8

µg/µl embryonic extract and 2.5 ng/µl synthetic *rps-3* mRNA. Control reactions were

performed without addition of ATP, creatine phosphate and creatine kinase.

Incubation was at 15°C for 2 h.

For analysis by one-step RT-PCR, reactions were diluted 1:80 in RNase-free MilliQ

water and aliquots were then used for one-step RT-PCR using the Luna® Universal

One-Step RT-qPCR kit (NEB) according to the manufacturer's instructions; primers

are given in Supplementary Table 1. Reactions were done with 1 µl or 2 µl diluted

reactions and analysed by agarose gel electrophoresis, or quantified using a

LightCycler 480 (Roche), with software release 1.5.1.62 Sp3. qPCR data was

analysed using the ΔC_T method (Schmittgen and Livak, 2008). Products P1, P2 and

P3 were identified by cloning into pGEM-T Easy (Promega) followed by Sanger

sequencing (Eurofins Genomics).

Protein Immunoprecipitation

Immunoprecipitations of GFP-tagged proteins were performed in triplicates or

quadruplicates using anti-GFP nanobody coupled agarose beads and control

agarose beads (GFP-Trap and control agarose beads, Chromotek GmbH),

essentially as described previously (Eijlers et al., 2024).

Protein analysis by LC-MS/MS and differential protein expression analysis

Proteomic analysis by mass spectrometry was done at the Aberdeen Proteomics unit

as previously described (Eijlers et al., 2024; Fasimoye et al., 2022). MaxQuant

version 1.6.5.0 (Cox and Mann, 2008) was used to process the raw data files with

the *C. elegans* reference proteome UP000001940 downloaded on the 24[th] of August

2021 as described (Eijlers et al., 2024), except that the feature "match between runs"

was applied. The MaxQuant protein group files were further analysed using Perseus

version 1.6.5.0 as described (Eijlers et al., 2024). Graphs were drawn in GraphPad

Prism.

RNA extraction and mRNA-Seq

Prior to RNA extraction, the worms were treated with 5-Ph-IAA when they reached

the L3 larval stage. Therefore, synchronised PE1176 L1s were grown in liquid culture

for 22.5 h until they started moulting into larval stage 3. They were then divided into

two and plated on control plates or 5-Ph-IAA plates, respectively, and cultured for 16

h, at which point the animals had started moulting into the L4 larval stage. Worms

were washed off NGM plates using M9 and washed 3 times by centrifugation at 749g

for 2 min at 4°C. Subsequently, 1 ml of Trizol (TRI Reagent, T9424, Sigma Aldrich)

was added per 50 µl of worm pellet and samples frozen in liquid nitrogen. Samples

were thawed at 37°C and re-frozen four times. They were then vortexed for 30 s, left

to rest for 30 s and the vortex/rest process repeated for a further three times. Phase

separation by addition of chloroform and recovery of RNA was done as described by

the manufacturer. The aqueous fraction was transferred to a low DNA-binding-tube

and diluted 1:1 with 70% ethanol. RNA was purified using PureLink RNA Mini Kit

(Thermo Fisher Scientific), using the on-column DNase-treatment option. Three

libraries per treatment group were sequenced on an Illumina NovaSeq instrument,

generating 2x150bp paired-end reads (Novogene UK Ltd).

Differential gene expression analysis

The quality of the raw reads was inspected in FASTQC 0.11.9 (Andrews, 2010) and

MULTIQC 1.12 (Ewels et al., 2016), and bases with a Phred score below 20 were

trimmed using TRIMGALORE 0.6.6 (Krueger, 2015). The trimmed reads were then

aligned to the *C. elegans* WB235 reference genome using HISAT2 2.2.0 (Kim et al.,

2015). Alignments were processed using SAMTOOLS 1.14 (Li et al., 2009). Gene-

level read counts were obtained using FEATURECOUNTS 2.0.2 (Liao et al., 2014),

quantifying against exon annotations and assigning fractional counts to all alignment

locations of multi-mapping reads.

The read counts were analysed in DESeq2 1.42. (Love et al., 2014), identifying

differentially expressed genes with an FDR (Benjamini and Hochberg, 1995)

significance threshold of 0.05, and shrinking fold changes at low-count genes using

the *apeglm* method (Zhu et al., 2019). Significantly differentially expressed genes

(DEGs) were examined for overrepresentation GeneOntology terms (biological

process ontology) using CLUSTERPROFILER 4.10.0 (Yu et al., 2012) with the

org.Ce.eg.db 3.18.0 database (June 2024). The GO results were simplified by

removing terms at graph levels 1-3, clustering GO terms semantically with the Wang

method at a similarity cutoff of 0.7 (Yu et al., 2010), and retaining for each cluster the

single GO term with the smallest *P*-value.

Data analysis using RStudio

RStudio was used for data analysis and graph generation (http://www.rstudio.com/).

The following R packages were used: ggplot2 (Wickham, 2016), dplyr

(https://dplyr.tidyverse.org/), carData/car (10.32614/CRAN.package.car), multcomp

(10.32614/CRAN.package.multcomp), ggpubr (https://rpkgs.datanovia.com/ggpubr/),

rstatix (https://rpkgs.datanovia.com/rstatix/), tibble (https://tibble.tidyverse.org/),

RcolorBrewer (10.32614/CRAN.package.RColorBrewer).

Screen for suppressors of *cmtr-1* loss-of-function

PE1176 worms were subjected to a standard ethyl methylsulfonate (EMS)

mutagenesis. Thirty F1 cultures were established by subjecting the mutagenised

parental animals to alkaline hypochlorite treatment and plating 500 F1 animals per 9

cm plate. The mutagenised animals and their F1 progeny were propagated on

standard NGM agar plates. The F2 embryos from alkaline hypochlorite treated F1s

were plated onto 9 cm NGM plates supplemented with 1 µM 5-Ph-IAA. Suppressor

mutants, identified on the basis of presence of eggs in the uterus of adult F2s, were

picked to establish lines, taking care to only maintain one line for each F1 lineage.

Genetic analysis of suppressor mutants

We reasoned that our screen would generate two classes of fertile mutants: those

that are *bona fide* extragenic suppressors of mNG^AID::CMTR-1 knockdown; and

loss-of-function mutants in the *TIR1* transgene, which would lead to 5-Ph-IAA-

independent expression of mNG^AID::CMTR-1. We thus screened each suppressor

line for the presence of nuclear mNeonGreen fluorescence when grown on media

containing 5-Ph-IAA. Mutant lines that showed mNeonGreen fluorescence in the

presence 5-Ph-IAA were assumed to be *TIR1* mutants; the *TIR1* coding region for a

subset of these was sequenced to confirm this assumption. Amplicons generated

from single-worm PCRs using primer pairs TIR1Fwd+TIR1Rev1 and

TIR1fo1+TIR1re2 were sequenced using the same oligos to prime Sanger

sequencing reactions (Eurofins Genomics).

Mutants that lacked nuclear mNeonGreen fluorescence in the presence of 5-Ph-IAA

were assumed to be genuine epistatic suppressors and were backcrossed with non-

mutagenised PE1176. The resultant strains were crossed into the *cmtr-1(syb3613)*

background to determine the ability of the suppressor mutations to suppress the

sterile phenotype of the *cmtr-1* null allele.

Sib-Selection and DNA Preparation

Two suppressor strains, PE1237 (*fe152*) and PE1245 (*fe154*), were subject to a

sibling subtraction/whole genome sequencing method adapted from a previously

reported method (Joseph et al., 2017). Male PE1053 worms were crossed with

hermaphrodites from PE1237/1245. F1 progeny carrying the *feEx342* transgenic

array were identified on the basis of pharyngeal tdTomato expression, picked to

separate plates, and allowed to segregate F2 progeny. Two hundred *feEx342*

transgenic F2s were picked singly to separate 35 mm plates and allowed to

propagate. Homozygous suppressor lines (Sup) were identified on the basis of the

presence of fertile non-tdTomato F3/F4 worms, and the absence of non-tdTomato

"scrawny" F3/F4 worms, the latter being observed on plates whose frequency

suggested that these were broods of suppressor heterozygotes. Lines that failed to

inherit suppressor mutations (Nonsup) were identified on the basis of the complete

absence of fertile non-tdTomato F3/F4 worms. Contaminated plates were discarded.

Worms were washed off plates, and Sup/Nonsup cultures pooled as outlined

previously (Doitsidou et al., 2010). The number of 35 mm F2 plates used to generate

the Sup and Nonsup pools were as follows: *fe152* Sup = 16; *fe152* Nonsup = 19;

fe154 Sup = 19; *fe154* Nonsup = 19. Genomic DNA was prepared using the Qiagen

Puregene Tissue Kit (Qiagen, 158063) as described previously (Doitsidou et al.,

2010). One library per pool was sequenced on an Illumina NovaSeq X Plus

instrument, generating 2x150bp paired-end reads (Novogene UK Ltd).

Variant Detection

Quality control of the raw reads was carried out using FASTQC 0.11.9 (Andrews,

2010) and MULTIQC 1.12 (Ewels et al., 2016). Adaptor removal and trimming of

bases with a Phred score below 20 was performed using TRIMGALORE 0.6.6

(Krueger, 2015). The trimmed reads were then aligned to the *C. elegans* WB235

reference genome using BWA MEM 0.17.17 (Li and Durbin, 2010). Variant detection

was done for all libraries together in a single run of FREEBAYES 1.3.6 (Garrison and

Marth, 2012), and the resultant VCF files were filtered and manipulated using

BCFTOOLS 1.14 (Danecek et al., 2021). We considered biallelic SNPs with

experiment-wide genotyping quality of at least 20 and at least 10 reads in each

library, and retained SNPs fixed for the alternative allele (>90% of read depth) in Sup

libraries and fixed for the reference allele (>90% of read depth) in Nonsup libraries.

SNPEFF 4.3t (Cingolani et al., 2012) was then used to functionally annotate the

candidate SNPs identified by this filtering strategy.

RESULTS

CMTR-1 is required for oocyte cell fate

To determine the role of cOMe in *C. elegans,* we studied the phenotypes of animals

lacking one or both CMTR proteins. We obtained a deletion allele, *syb3613*, of *cmtr-*

1 (SunyBiotech), which removes the methyltransferase domain, and is thus a

predicted null allele. We also studied a deletion allele of *cmtr-2*, *tm4453*, that deletes

key residues in the methyltransferase domain (Dix et al., 2022). While *cmtr-*

1(syb3613) homozygotes were able to reach adulthood, they were 100% sterile

(Figure 1A,B) and were thinner than similarly staged *syb3613* heterozygotes.

We found that the sterility was due to defects in the formation of oocytes, which

ranged from a completely masculinised germline that produced only sperm, through

to animals that were able to produce only a few embryos that failed to develop

(Figure 1D,E,H). In all cases, *cmtr-1(syb3613)* homozygotes showed much smaller

germlines than wild-type worms, something also observed in mutants that cause

masculinised hermaphrodite germlines (Kimble and Crittenden, 2007).

We confirmed that the defects observed were due to loss of *cmtr-1* function by

demonstrating that an extrachromosomal transgene consisting of the *cmtr-1* cDNA

expressed under the control of the *rps-0* promoter rescued the germline sterility

defects (Figure 1F,G). As part of this analysis we found that offspring that failed to

inherit the extrachromosomal transgene generally showed a more severe phenotype

than the offspring of heterozygous mothers, displaying more severe growth defects

(Figure 1G). Since extrachromosomal transgenes are generally silenced in the

germline (Kelly et al., 1997), this likely indicates that there is a significant *cmtr-1*

maternal contribution, which is absent from the non-transgenic offspring of

transgenic mothers. The more severe phenotypes of these animals indicates that the

loss of *cmtr-1* function impacts other tissues beyond the germline.

In contrast to loss of *cmtr-1*, *cmtr-2* loss-of-function homozygotes are superficially

wild type (Dix et al., 2022). To investigate whether loss of *cmtr-2* function influenced

the *cmtr-1* loss-of-function phenotype, we examined the phenotype of *cmtr-1*; *cmtr-2*

double mutants. This analysis showed that loss of *cmtr-2* enhanced the phenotype of

cmtr-1 single mutants (Figure 1H). Thus, CMTR-2 contributes to the role played by

CMTR-1.

The failure of oocyte differentiation in *cmtr-1* mutant homozygotes suggested that

CMTR-1 function is required for some component(s) of the sperm-oocyte switch. To

determine when CMTR-1 protein is required during larval development for oocyte

cell fate, we generated an auxin-inducible degeneration (AID) allele of *cmtr-1*, which

expresses endogenous CMTR-1 tagged with the auxin-inducible degron (AID)

peptide and monomeric NeonGreen (mNG^AID::CMTr1). This strain (PE1176) also

expresses the *A. thaliana* TIR1(F79G) transgene necessary for auxin-dependent

protein depletion (Hills-Muckey et al., 2022). Growing this strain in the presence of 5-

Ph-IAA showed rapid depletion of mNG^AID::CMTr1 (Supplementary Figure 1), and

such worms were 100% sterile and showed reduced body size compared to

untreated controls (Figure I).

Using this strain, we performed a series of experiments designed to deplete

mNG^AID::CMTR-1 at specific times during postembryonic development. We

initiated synchronous cultures of 50 starved, first-stage larvae (L1s) on control plates

and transferred them to 5-Ph-IAA plates at defined times (Figure 1J). We also

performed the inverse experiment, culturing L1s on 5-Ph-IAA plates and transferring

them to control plates at the same time points. This "auxin-shift" experiment (named

by analogy to temperature-shift experiments performed with temperature-sensitive

mutants) revealed that 5-Ph-IAA depletion of mNG^AID::CMTR-1 resulted in sterility

only during a critical time period between approximately 24 and 40 hours following

initiation of starved L1 larval development. The critical period during which CMTR-1,

and thus cOMe, is required corresponds roughly to the L3 and L4 larval stages.

Given that the sperm-oocyte switch occurs during the L4 stage (Kimble and

Crittenden, 2007), the time-window during which CMTR-1 is required suggest that

this is when transcripts involved in germline sex determination need to be modified

by cOMe for normal germline cell fate.

Figure 1. Loss of *cmtr-1* function results in defects in germline development

and body size. Representative *cmtr-1(syb3613)/+* heterozygote (A), compared to

sibling *cmtr-1(syb3613)* homozygotes (B). Wild type young adult hermaphrodite (C),

showing posterior gonad arm (gold overlay), compared to the same regions of *cmtr-*

1(syb3613) homozygotes, one of which shows a single defective oocyte (D), and the

other has a fully masculinised germline that has only produced sperm (E). Position of

the vulva is indicated by "v". Scale bars represent 100 µm (A,B) and 20 µm (C,D,E).

(F) Transgenic expression of *cmtr-1* cDNA under the control of the *rps-0* promoter

(*Ex-cmtr-1(+)*) rescues the fertility defects of *cmtr-1(syb3613)* mutants (*cmtr-1(-)*). (G)

Comparison of the number of worms reaching adulthood 72 hours after hatching (n >

61 for each experiment). Mutant offspring (*cmtr-1(-)*; *cmtr-2(+)* or *cmtr-1(-)*; *cmtr-2(+)*)

of transgenic parents (transgene is indicated '*Ex-cmtr-1(+)*)are slower-growing

compared to genetically identical offspring of heterozygous parents. (H) Loss of *cmtr-*

2 increases the severity of the *cmtr-1* loss-of-function germline defects. Histograms

show the distribution of the three germline defect classifications caused by loss of

cmtr-1 function: production of a few arrested embryos (Emb, blue); abortive oocyte

production (Ooc, gold); masculinised germline (Mog, grey). (I) Worm diameter,

measured at the mid-body, for worms homozygous for the mNG^AID::*cmtr-1* allele

subject to control and 5-Ph-IAA treatment. Two-way ANOVA and a Tukey post hoc

test were used to determine p-values; * < 0.05, ** < 0.01, *** < 0.001, n = 75 animals

per group. (J) "Auxin-shift" experiment to determine when CMTR-1 function is

required for fertility. Starved L1 larvae (n=50) were placed on either 5-Ph-IAA or

control plates for the indicated time period, before being transferred to the alternative

plate type (i.e. those on 5-Ph-IAA plates were moved to control plates, and vice

versa). The worms were then left on the second plate, allowed to reach adulthood,

and scored for fertility.

The *C. elegans* CMTR proteins show distinct expression patterns and protein-protein interactions

Studies in cultured mammalian cells and *Drosophila* indicated that the two CMTR proteins have conserved, distinct subcellular distributions: in both cases, CMTR1 is primarily nuclear, whereas CMTR2 is both nuclear and cytoplasmic (Haline-Vaz et al., 2008; Haussmann et al., 2022; Werner et al., 2011). Using strains that express GFP-tagged alleles of the endogenous proteins, we found that *C. elegans* CMTR proteins are nuclear throughout postembryonic development (Figure 2A-D), being expressed in most, if not all cells, though CMTR-2 does not appear to be as strongly expressed in the nucleus as CMTR-1. While the overall levels of the two proteins are similar in embryos, as judged by Western blot (Figure 2H), the intracellular localisation of the two proteins is strikingly distinct: CMTR-1 is predominantly nuclear (Figure 2E), in contrast to CMTR-2, which is strongly cytoplasmic (Figure 2F, compare to Figure 2G).

To investigate the functional significance of the distinct localisation of the proteins during embryogenesis further, we performed immunoprecipitation analyses from embryonic lysates to assess their respective protein-protein interactions. This analysis revealed that the two proteins have non-overlapping protein partners: CMTR-1 showed a rich set of protein-protein interactions (Figure 3A-C); while CMTR-2 displayed a paucity of interactors (Figure 3D, E); none of them matching those we found for CMTR-1. The lack of overlap in protein interaction partners is consistent with the low levels of CMTR-2 in the nucleus (Figure 2E,F) in embryos, but this analysis fails to shed light on the functional significance of the cytoplasmic enrichment of CMTR-2.

C. elegans CMTR-1 showed broadly the same protein-protein interactions with a suite of pre-mRNA processing factors that were observed for mammalian cells

(Simabuco et al., 2019), with components of the spliceosome, heterogeneous

nuclear ribonucleo-proteins and ribosomal proteins being prominent interaction

partners (Figure 3A-C). We also saw an interaction with SNA-3, a component of the

spliced leader *trans*-splicing machinery (Fasimoye et al., 2022) that has previously

been shown to associate with nascent RNA polymerase II transcripts (Eijlers et al.,

2024). Many of the interactions were abolished by RNase treatment, consistent with

CMTR-1 interacting with them by virtue of binding to the same transcript. However,

there are notable interactions that were preserved in RNase treated samples,

included the previously well-defined CMTr1 interacting protein, DHX15 (DDX-15 in *C.*

elegans)(Inesta-Vaquera et al., 2018; Toczydlowska-Socha et al., 2018), and some

core components of the spliceosomal catalytic complex.

Figure 2. CMTR-1 is constitutively nuclear but CMTR-2 subcellular localisation

is developmentally regulated. Representative images showing the respective

expression of GFP tagged CMTR-1 (A, C, E) or CMTR-2 (B, D, F) in L2 larval stage

epidermis (A, B), L4 germlines (C, D) and early embryos (E, F). Wild type embryos

(G) show the level of embryonic background autofluorescence. Images in A – D and

E - G were taken at the same camera exposure settings. Scale bars indicate 20 μm.

(H) Western blot of GFP-tagged CMTR-1 and CMTR-2 in embryonic extracts probed

with anti-GFP polyclonal antibody. Specific signals are absent from wild type

embryos. Arrow indicates non-specific primary anti-GFP antibody immunoreactivity.

Amido black stained immunoblot indicates loading of protein in each lane.

Figure 3. CMTR-1 and CMTR-2 show distinct protein-protein interaction partners

Proteins enriched by co-immunoprecipitation with GFP::CMTR-1 (A, B, C) and GFP::CMTR-2 (D, E) from PE1001 and PE991 embryonic extracts treated without (A, D) and with RNase (B, E). Immunoprecipitations were performed in triplicate or quadruplicate using anti-GFP nanobodies coupled to agarose beads, or control agarose beads. Immunoprecipitated proteins were identified by label-free quantification as described (Eijlers et al 2024). Graphs show enrichment in anti-GFP nanobody immunoprecipitations (anti-GFP) compared to bead only controls (control) and the false discovery rate (y-axis). Proteins that were significantly enriched are shown in gold. The significance cutoff curve is drawn in black (false discovery rate <=0.05 and S0 of 0.1). TBH-1 (blue), a dopamine β-hydroxylase, is found in all immunoprecipitations but our previous work suggests that it non-specifically associates with GFP-tagged proteins (Eijlers et al., 2024; Fasimoye et al., 2022). (**C**) Heatmap summarising the enrichment of spliceosome components, RNA processing factors and RNA binding proteins in immunoprecipitation of GFP::CMTR-1. Proteins were grouped using the gene ontology annotation retrieved from UniprotKB

524   (Ashburner et al., 2000; Gene Ontology Consortium et al., 2023; The UniProt

525   Consortium et al., 2023). ne: not enriched. For complete lists of identified proteins

526   see Supplementary Table 2.

Loss of cOMe is the cause of the *cmtr-1* loss-of-function phenotype

The phenotypic impact of loss of *cmtr-1* compared to *cmtr-2* suggested that CMTR-1 function might be more important for cOMe modification of transcripts than CMTR-2. This was supported by our previous analysis, which showed that loss of *cmtr-2* did not detectably affect cOMe levels (Dix et al., 2022). To investigate the dependence of cOMe on CMTR-1, we assayed RNA derived from animals in which mNG^AID::CMTR-1 was knocked down on its own, and in combination with the *cmtr-2(tm4453)* loss-of-function mutation. Using a sensitive assay based on recapping mRNA with ^{32}P-alpha-GTP, followed by RNase I digestion to detect cOMe in small amounts of mRNA (Anreiter et al., 2020; Dix et al., 2022; Haussmann et al., 2022), we found that cOMe modified RNA (cap1) was almost completely abolished (Figure 4A). Note that using this assay, we were not able to detect cap2 modifications, consistent with a previous study that this cap2 is largely absent from *C. elegans* (Despic and Jaffrey, 2023). This assay also revealed that CMTR-2 makes a negligible impact on cap1 levels, as we previously demonstrated (Dix et al., 2022). Thus, CMTR-1 is the important determinant of cOMe in *C. elegans*, consistent with its strong loss-of-function phenotype.

We then looked specifically at cap1 in poly(A) mRNA derived from animals lacking functional CMTR-2 and depleted of mNG^AID::CMTR-1 by treatment with 5-Ph-IAA to find that cOMe is essentially absent (Figure 4B-G). Of note, the first transcribed nucleotide for the majority of *C. elegans* mRNAs will be the guanosine originating from the *trans*-spliced leaders. As shown by separation of the second dimension in solvent C, the spliced leader is also 2`-*O*-ribose methylated by CMTR-1 (Figure 4E-G), as expected for an RNA polymerase II transcript.

Previous studies in mammals have shown that around 50% of CMTR1 is in a

complex with DHX15, and that this interaction represses its methyltransferase

activity (Inesta-Vaquera et al., 2018; Toczydlowska-Socha et al., 2018). CMTR1 also

influences RNA polymerase II recruitment to transcription start sites and the

transcription of histone and ribosomal protein genes (Liang et al., 2022). We thus

wanted to confirm that the *cmtr-1* loss-of-function phenotype was due solely to loss

of cOMe; in other words that it was dependent on CMTR-1 methyltransferase activity.

To address this, we engineered a missense mutation, K244A, into the endogenous

cmtr-1 gene. The orthologous mutation in human CMTR1 (K239A) has previously

been show to abolish methyltransferase function (Bélanger et al., 2010; Feder et al.,

2003), and the lack of function of this mutation in *C. elegans* has been assessed in

transgene rescue experiments (Meisel et al., 2024). We carried out site-directed

mutagenesis of the *cmtr-1* allele that expresses CMTR-1 with a N-terminal GFP tag

(Figure 2), allowing us to monitor the impact of the K244A mutation on the

expression and localisation of CMTR-1 (Figure 4H,I). All worms homozygous for the

K244A (*fe175*) allele showed the same recessive sterile and growth defective

phenotypes displayed by the *syb3613* mutation, and these phenotypes were

complemented by the *rps-0::cmtr-1(+)* transgene (Figure 4J). Since these worms

specifically lack the methyltransferase activity of CMTR-1, this indicates that loss of

cOMe is the cause of the germline phenotypes.

572
Figure 4. CMTR-1 is required for cap-adjacent 2`-O-ribose methylation in *C.*

elegans. Analysis of cap-adjacent 2`-*O*-ribose RNA methylation with (control) and

without (5-Ph-IAA) mNG^AID::CMTR-1. (A) Recapping assay to detect mRNA using

^{32}P-αGTP. 5′ cap structures were separated on a 22% denaturing polyacrylamide

gels after digestion with RNase I (lanes 6-7, right). Markers (M1-4) are RNase I

digested ^{32}P-αGTP capped oligonucleotides. 2`-*O*-ribose methylation (N_m) was

added using vaccinia CMTr. Sequences of markers are shown on the left. (B - G)

Two dimensional thin-layer chromatograms (2D-TLCs) showing modifications of the

first cap-adjacent nucleotides prepared from total RNA isolated from worms

homozygous for the mNG^AID::*cmtr-1* and *cmtr-2(-)* alleles, grown on control (C, F)

or 5-Ph-IAA plates (D, G). (B, E) Schematic diagrams of a 2D-TLC, depicting the

positions of unmodified and 2`-*O*-ribose methylated nucleotides. One set of

chromatograms (C, D) was run in solvent B for the second dimension, another set (F,

G) was run in solvent C. 2`-O-ribose methylated nucleotides are indicated by arrows.

(H) Representative germline GFP fluorescence of an animal expressing GFP-tagged

CMTR-1(K244A), compared to GFP-tagged wild-type CMTR-1 protein. (J) Brood

counts of wild-type (data from Figure 1F) and animals homozygous for the K244A

mutation, in the absence and presence of the wild-type *cmtr-1* cDNA expressed

under the control of the *rps-0* promoter (*Ex-cmtr-1*). Counts were performed on five

individual animals per genotype.

Loss of CMTR-1 does not affect spliced leader *trans*-splicing

Approximately 85% of *C. elegans* genes produce transcripts that are *trans*-spliced to the SL1 or SL2 spliced leaders (Tourasse et al., 2017), and both spliced leaders also possess cOMe modifications (Figure 4E-G). In addition, we observed an interaction between CMTR-1 and SNA-3, an essential component of the SL1 *trans*-splicing machinery (Fasimoye et al, 2022, Eijlers et al, 2024) (Figure 3C). We therefore wanted to investigate whether cOMe had any functional impact on *trans*-splicing. We depleted mNG^AID::CMTR-1 in embryos by treating them with 5-Ph-IAA-AM, which is able to penetrate the egg-shell (Negishi et al., 2022) and prepared *in vitro* splicing extracts as described previously (Lasda et al., 2011). These were compared to extracts prepared from untreated embryos, and to PE1220 embryos depleted for SNA-3::AID^mNG, which, based on SNA-3's essential role, is predicted to impair spliced leader *trans*-splicing (Fasimoye et al., 2022). 5-Ph-IAA-AM treatment reduced mNG^AID::CMTR-1 and SNA-3::mNG^AID steady-state levels to undetectable levels and by 80%, respectively (Figure 5A).

We analysed the effect of protein depletion on spliced leader *trans*-splicing in embryonic extracts using two assays: 1) the steady state spliced leader *trans*-splicing levels of endogenous *rps-3* transcripts; and 2) the *trans*-splicing by embryonic extracts of an exogenous, synthetic *rps-3* transcript prepared by *in vitro* transcription. In the first assay, depletion of SNA-3 led to a 40.6-fold increase in non-*trans*-spliced, outron-containing endogenous *rps-3* RNAs, while the depletion of CMTR-1 had only a minor effect, leading to a 2.75-fold increase of non-*trans*-spliced *rps-3* RNA (Figure 5B).

SL1 *trans*-splicing of synthetic *rps-3* transcripts was analysed by PCR as previously

described (Lasda et al., 2011). The synthetic *rps-3* RNA used as substrate contains

plasmid sequence at the 3`-end allowing the use of primers that distinguish the

synthetic *rps-3* RNA from endogenous *rps-3* RNA (Figure 5C). Products P1, P2 and

P3 (Figure 5C, D) were identified by sequence analysis and are detected by the SL-

rps-3 primers that amplify SL *trans*-spliced synthetic RNA and the *rps-3* RNA control

primers that detect all forms of synthetic *rps-3* RNA. Importantly, SL-*rps-3* primer

products were only detected in reactions containing ATP necessary for spliced leader

trans-splicing to occur (Figure 5C). Analysis by qPCR shows that depletion of CMTR-

1 has a minor effect on *in vitro trans*-splicing of the exogenously added synthetic *rps-*

3 RNA, with near-identical amounts of product detected in the presence of ATP,

independent of whether CMTR-1 was depleted or not (Figure 5D). In contrast, the

depletion of SNA-3 led to a 280-fold reduction of detection of *trans*-spliced products

(Figure 5E). Thus, both assays failed to detect a clear, pronounced effect of CMTR-1

depletion on spliced leader *trans*-splicing, suggesting that neither cOMe nor CMTR-1

has a significant role in this process.

Figure 5. CMTR-1 is not required for spliced leader *trans*-splicing. (A) Western
blots showing levels of endogenous mNG^AID::CMTR-1 and SNA-3::mNG^AID
proteins in embryonic extracts prepared from PE1176 and PE1220 animals,
respectively. Embryos were control-treated with DMSO or with the auxin analogue
5-Ph-IAA-AM for 4h prior to extract preparation. Proteins were analysed by Western
blotting and detected by anti- mNeonGreen antibodies. Total protein visualised using
amidoblack staining was used to standardise the levels of mNeonGreen-tagged
proteins. The level of SNA-3::mNG^AID protein in 5-Ph-IAA-AM treated extract was
expressed as percentage of that detected in the DMSO-treated extract. The
mNG^AID::CMTR-1 was not detectable in 5-Ph-IAA-AM treated extracts (nd). Non-
specific bands are indicated with a star symbol. (B) Quantitative RT-PCR
measurement of outron retention of endogenous *rps-3* transcripts. The diagram
shows the location of primers that detect outron and internal regions of *rps-3* RNA.
RNA was isolated from embryonic extracts, reverse transcribed, and analysed by
quantitative PCR as described previously (Philippe et al., 2017). The graph shows
the ratio of endogenous *rps-3* outron to internal amplicon levels in PE1176 (gold) and
PE1220 (blue) extracts treated with 5-Ph-IAA-AM (+) or control treated with DMSO (-

) determined using the ΔC_T method (Schmittgen and Livak, 2008). Measurements

are technical replicates and their means (horizontal line, mean values are shown).

The analysis is representative of 2 independent measurements. (C) Substrate and

products of *in vitro* SL *trans*-splicing. Diagrammatic representation of the synthetic

rps-3 transcript (substrate S) with outron (ou) at the 5` end and the products P1, P2

and P3. The locations of primer pairs SL1-*rps-3* and *rps-3* RNA used for detection by

one-step qPCR and expected amplicon size are indicated. (D) Analysis of *in vitro* SL

trans-splicing by agarose gel electrophoresis. Synthetic *rps-3* RNA was incubated

with embryonic extracts from PE1176 embryos treated with 5-Ph-IAA-AM or control-

treated with DMSO (+/- 5-Ph-IAA-AM) as described. ATP and ATP regeneration

system were included where indicated (+/- ATP). Products were amplified by one-

step PCR, analysed by 1.8% agarose gel electrophoresis and visualised by staining

with ethidium bromide. (E) Analysis of *in vitro* SL *trans*-splicing by quantitative RT-

PCR. Synthetic *rps-3* RNA was incubated with embryonic extracts from PE1176

(gold) or from PE1220 embryos (blue) treated with 5-Ph-IAA-AM or control-treated

with DMSO (+/- 5-Ph-IAA-AM) as described. ATP and ATP regeneration system were

included where indicated (+/- ATP). Products were quantified by quantitative one-

step RT-PCR as described. The graph shows the ratio of *trans*-spliced SL1-*rps-3*

products standardised with respect to total synthetic *rps-3* RNA using the ΔC_T

method (Schmittgen and Livak, 2008). The *in vitro* processing reactions were done in

triplicates. Shown are the results for each reaction and the mean (horizontal line,

values are indicated).

Loss of cOMe leads to differential transcript levels for genes involved in the intracellular pathogen response and germline sex determination

To investigate the possible impact of loss of cOMe on gene expression, we compared transcript levels for 5-Ph-IAA treated to control-treated animals. We cultured synchronised populations of mNG^AID::CMTR-1 larvae, and began the 5-Ph-IAA treatment when the animals were at the third larval (L3) stage, corresponding to the developmental stage at which CMTR-1 is critical for the sperm-oocyte switch (Figure 1J). Both control and 5-Ph-IAA treated animals were harvested for RNA preparation at early L4 stage.

We performed differential gene expression analysis, comparing RNA-Seq derived from control versus 5-Ph-IAA treated animals. This revealed 1953 genes whose steady state transcript levels were significantly upregulated by mNG^AID::CMTR-1 depletion (FDR<=0.05, absolute log2 fold change >=1), and 3398 genes for which transcript levels were significantly reduced by the same treatment (FDR<=0.05, absolute log2 fold change >=1) (Supplementary Table 3).

Inspection of the genes that were significantly enriched more than four-fold in mNG^AID::CMTR-1 knock-down worms compared to controls revealed upregulation of around 80 genes, including *fbxa-158*, *pals-14*, and *pals-6*, which belong to a class of genes that constitute the intracellular pathogen response (IPR), a *C. elegans* transcriptional immune response to intracellular pathogens (Lažetić et al., 2023). IPR is induced by viral and microsporidia infections, but also mitochondrial dysfunction(Mao et al., 2022; Sowa et al., 2020).

To systematically investigate the possibility that IPR genes were generally upregulated in response to depletion of CMTR-1, we extracted the fold changes of

272 IPR genes that were upregulated more than four-fold in response to expression

of Orsay virus RNA-dependent RNA polymerase (Sowa et al., 2020)(Figure 6A).

These IPR genes showed statistically significantly higher transcript levels in

response to CMTR-1 depletion compared to non-IPR genes (P<<0.001), though the

effect size was modest (mean fold change: 0.697) (Figure 6A). The effect of loss of

CMTR-1 on innate immune gene expression was also supported by GeneOntology

(GO) term enrichment analysis (Figure 6B): the three most significantly

overrepresented GO terms among upregulated genes were "innate immune

response" (GO:0045087, qvalue = 2.02×10^{-8}), "response to external biotic stimulus"

(GO:0043207, qvalue = 3.92×10^{-7}), and "defense response" (GO:0006952, qvalue =

4.20×10^{-7}). Thus, loss of CMTR-1 and cOMe leads to transcriptional upregulation of

many intracellular pathogen response genes.

GO term enrichment analysis among downregulated genes in response to loss of

CMTR-1 highlighted a broad range of metabolic terms but, intriguingly, also terms

related to gametogenesis, including "gamete generation" (GO:0007276, qvalue =

3.59×10^{-13}) and "germ cell development" (GO:0007281, qvalue = 1.96×10^{-9})

(Supplementary Table 3).

To investigate whether transcripts from genes associated with germline sex

determination were downregulated in response to depletion of CMTR-1, we

extracted the fold changes of a set of 64 genes involved in germline sex

determination (Figure 6A). We found that this set of genes showed statistically

significantly (P<<0.001) lower fold changes than all other genes, though as for the

IPR gene response, the effect size was modest (mean fold change: -0.287).

Nonetheless, since germline development is known to be highly sensitive to shifts in

gene expression (Ellis, 2022), this overall trend of downregulation may explain the

germline defects that we observed (Figure 1).

We were unable to establish a clear correlation between transcript level changes and

the growth defects that we observe in animals depleted of CMTR-1. It is likely that

changes in the expression of a broad range of genes are responsible for this

phenotype.

Figure 6. Depletion of CMTR-1 results in upregulation of transcripts associated

with the innate immune response and downregulation of transcripts involved

in germline sex determination. (A) Upper panel, volcano plot showing differential

gene expression in animals subject to CMTR-1 depletion compared to controls (three

biological replicates). Intracellular pathogen response genes (IPR) are highlighted in

orange. Germline sex determination genes (Oocyte) are highlighted in blue. Lower

panel, box plot of the fold change data, showing the p-values (Welch Two Sample t-

test) for the differences between the means of the IPR and Oocyte gene sets and the

"other" (non-IPR/germline sex determination genes). The gene *eol-1*, which is part of

the IPR and was subsequently shown in this study to be epistatic to *cmtr-1* (Figure 7),

is also indicated. (B) Gene ontology enrichment analysis showing the significantly

enriched gene categories among upregulated genes in animals subject to CMTR-1

depletion compared to controls ranked by q-value.

Loss of the RNA decapping exonuclease EOL-1 bypasses the need for CMTR-1

To better understand the molecular and cellular significance of the global loss of cOMe, we performed an unbiased screen for suppressors of the *cmtr-1* loss-of-function phenotype, reasoning that such mutations might allow us to identify components involved in the cellular response to cOMe modified RNAs.

We mutagenised PE1176 worms with ethyl methanesulfonate (EMS) and grew the F2 generation on NGM plates containing 1 μM 5-Ph-IAA. We then screened for the presence of healthy gravid worms, which would indicate suppression of the *cmtr-1* loss-of-function phenotype. Suppression was confirmed by showing that the worms could be maintained over multiple generations on 5-Ph-IAA containing media. Only one strain was retained per F1 culture, to ensure independence of each suppressor mutation. From combined screens of approximately 20,000 haploid genomes, we recovered 27 independent mutants that were fertile when grown in the presence of 5-Ph-IAA.

We designed the screen with the expectation that we would recover loss of function mutations in the *TIR1(F79G)* transgene, which restore fertility simply by preventing 5-Ph-IAA-depdenent degradation of mNG^AID ::CMTR-1. Such mutations, which constitute a type of informational suppression (Mount and Anderson, 2000), could be distinguished from *bona fide cmtr-1* loss-of-function suppressors by the fact that they express mNG::AID^CMTR-1 in the presence of 5-Ph-IAA (Supplementary Figure 2). Using this secondary screen, we found that of the 27 mutants, 21 showed nuclear mNeonGreen fluorescence despite being grown on 5-Ph-IAA. We sequenced the TIR1 coding region from six of the putative TIR1 loss-of-function mutants and confirmed that all were missense mutations (Supplementary Figure 2).

The six mutants that lacked mNeonGreen fluorescence but were nonetheless fertile

were thus putative suppressors that bypass the requirement for CMTR-1. We

confirmed that they also suppressed the *cmtr-1(syb3613)* null allele: all six mutations

suppressed the sterile phenotype and growth defects conferred by the *syb3613*

allele confirming that they are epistatic to *cmtr-1* loss-of-function (Figure 7A).

To identify the gene(s) affected by these six mutations, we used a whole-genome

sequencing (WGS) mapping strategy based on the sibling-subtraction approach

described previously (Joseph et al., 2017). We mapped two of the suppressor

mutants, *fe152* and *fe154*, using this approach (Supplementary Figure 3). The

sibling-subtraction WGS mapping approach relies on the identification of variants

that are present in more than 90% of reads derived from the pooled broods of F2

suppressor animals, but absent, or at a low frequency in the reads derived from their

non-suppressor sibling broods. In practice, we found that focussing only on variants

found in 100% of suppressor pool reads was sufficient to narrow down the location of

the suppressor mutations to a region of chromosome V (Supplementary Table 4).

SnpEff and manual curation of these variants revealed predicted damaging variants

in a single gene, *eol-1*, allowing us to assign *fe152* and *fe154* as premature stop

codon mutations, W208Opal and W211Amber, respectively (Figure 7B).

Sanger sequencing of the *eol-1* locus in the other four suppressor mutants revealed

that each was a predicted damaging allele of *eol-1* (Figure 7B). A third nonsense

mutation is created by *fe174*, R168Opal, and two mutations, *fe155* and *fe172*, are

missense mutations, C162Y and E297K, respectively. The *fe153* allele was found to

be a deletion of 1456 bp that removes 940 bp upstream of the start codon as well as

half of the coding region. The deletion is accompanied by insertion of 349 bp of

sequence, part of which appears to be repetitive DNA found on multiple *C. elegans*

chromosomes.

Having identified the genetic location of the suppressor mutations, we crossed them

into a *cmtr-1* wild-type background, which revealed that they had wild-type brood

sizes (Figure 7A). Thus, loss of *eol-1* function is compatible with wild-type fertility. We

also did not detect any obvious phenotypes that would indicate defects in embryonic

or postembryonic development.

Although the recovery of six independent function altering mutations in the same

gene is strong evidence that loss of *eol-1* function suppresses the requirement for

CMTR-1 and thus cOMe, we wanted to formally confirm this. We aimed to delete the

eol-1 gene, using Cas9/CRISPR to generate double-strand breaks at either end of

the gene. However, we found that the guide RNA targeted to the 5`-end of the coding

region was ineffective, so we were not able to identify gene deletions. Nonetheless,

we identified two alleles, *fe176* and *fe177*, that created frameshifts just upstream of

the stop codon, both of which would be predicted to result in peptide extensions that

derive from translation into the 3`-UTR (Figure 7B). Proteins with such C-terminal

extensions are subject to "readthrough mitigation" mechanisms that lead to

downregulation of the affected protein (Arribere et al., 2016; Kesner et al., 2023;

Müller et al., 2023). Consistent with this, both alleles suppress the sterile phenotype

caused by depletion of CMTR-1, confirming that loss of *eol-1* function bypasses the

requirement for cOMe.

EOL-1 is a member of the DXO/Rai1 family of decapping enzymes (Chang et al.,

2012; Jiao et al., 2013; Mao et al., 2020; Shen et al., 2014), and some members are

also distributive 5` to 3` exonucleases, so that they can both decap and degrade

their target RNAs. The missense mutations, C162Y and E297K, alter residues that

are part of the predicted active site of EOL-1 (Figure 7C). Indeed, E297 is the

homologue of mammalian DXO E253, a critical catalytic residue, part of the so-called

Motif IV (Chang et al., 2012; Jiao et al., 2013). Thus EOL-1 loss of function

mutations which specifically abolish its decapping/exonuclease activity, completely

suppress the sterility caused by loss of CMTR-1, indicating that removing this activity

is likely the mechanism which mediates the suppression.

Previous studies have shown that EOL-1 is a prominent part of the intracellular

pathogen response. Given that loss of CMTR-1 led to enrichment of transcripts

associated with this response, we looked for evidence that *eol-1* transcripts are

enriched in CMTR-1 depleted animals compared to controls (Figure 6A) and found

that they are enriched 2.76-fold. EOL-1 is one of nine *C. elegans* DXO homologues

(Mao et al., 2020), and is part of a clade of paralogues (Figure 7D). Transcripts from

four of these are also upregulated by depletion of CMTR-1, though only two of them

are broadly upregulated as part of the intracellular pathogen response

(Supplementary Table 3).

Figure 7. Loss of function mutations in *eol-1* suppress the loss of *cmtr-1*

mutant phenotype. (A) Brood counts for the six *eol-1* alleles identified as

suppressors of *cmtr-1* loss-of-function in *cmtr-1(syb3613)* and *cmtr-1(+)* backgrounds.

Histograms show means and error bars show standard deviations for counts derived

from five adults for each genotype. (B) Schematic of EOL-1 showing the location of

the suppressor mutations and the two mutations generated by Cas9/CRISPR (*fe176*

and *fe177*). Shading indicates secondary structure assignments based on the

AlphaFold prediction (A5JYX9). Roman numerals indicate highly conserved motifs

necessary for decapping and exonuclease activity (Wang et al., 2015). (C) AlphaFold

prediction for EOL-1, showing location of residues affected by missense suppressor

mutations (green). Active site residues other than E197 are in yellow. (D) Phylogram

showing the relationship between EOL-1 and other DXO family members. Blue text

indicates proteins with transcripts that are significantly enriched in mNG^AID::CMTR-

1 depleted animals. Fold enrichment is given, along with adjusted *p* values in

brackets. Node statistical supports are given as aLRT (SH-like) values.

$cmtr\text{-}1(+); eol\text{-}1(+)$

$cmtr\text{-}1(\text{-}); eol\text{-}1(+)$

$cmtr\text{-}1(\text{-}); eol\text{-}1(\text{-})$

Figure 8. Cap-adjacent 2`-O-ribose methylation protects cellular transcripts
from EOL-1-mediated degradation. In this proposed model, EOL-1 is prevented
from decapping cOMe modified transcripts (M). In the absence of CMTR-1,
transcripts lack cOMe and thus become substrates for EOL-1. Transcripts for genes
that are sensitive to copy number changes, such as those involved in germline cell
fate, would be most susceptible to loss of cOMe. Loss of EOL-1 prevents the
degradation of these transcripts. Black shaded circle indicates the monomethyl
guanosine cap.

DISCUSSION

Insights into the role of cOMe in animal development requires that one study the effects of loss of the CMTR proteins in the context of whole organism biology. While work in mice and flies have advanced our understanding of cOMe biology, the fact that these RNA modifications are essential for mammalian embryogenesis, but dispensable in flies and some cultured mammalian cells is puzzling. By investigating cOMe function in *C. elegans*, we have obtained a broader evolutionary perspective of the role of these RNA modifications.

Unlike the situation in mammals (Furuichi et al., 1975; Werner et al., 2011), we find that CMTR-1 is the principal enzyme involved in adding cOMe to *C. elegans* RNAs, and consistent with this, has higher steady state nuclear levels compared to CMTR-2. Similarly consistent with these observations, CMTR-1 is essential while CMTR-2 is not. We did not find a function for CMTR-2 beyond augmenting that of CMTR-1, which suggests that CMTR-2 is also able to modify the first nucleotide, a substrate specificity also found for its *Drosophila* and human homologues in modifying canonical AGU starting transcripts (Anreiter et al., 2020; Dix et al., 2022; Haussmann et al., 2022).

The two nematode CMTR proteins have distinct sub-cellular localisations and interaction partners, suggesting that they are functionally differentiated. Much of this may be due to their differing subcellular localisations. As in mammals and *Drosophila* (Haussmann et al., 2022; Werner et al., 2011), we see stronger associations of CMTR-2 with the cytoplasm. This is most pronounced during embryogenesis where CMTR-2 is apparently excluded from the nucleus, something not reported in other animal cells. This observation warrants further investigation.

The protein interaction profile of *C. elegans* CMTR-1 is highly similar to its

mammalian homologue (Simabuco et al., 2019), with both proteins showing

prominent interactions with the spliceosome. The functional significance of these

observations is unclear but may simply reflect the fact that these proteins assemble

on nascent transcripts, though we cannot rule out a functional role for CMTR-1 in *cis*-

splicing of certain transcripts. However, our experiments do indicate that CMTR-1 is

not an essential component of nematode spliced leader *trans*-splicing. Despite the

interaction between CMTR-1 and SNA-3, neither of our functional assays indicates

that CMTR-1 is required for the activity of the *trans*-spliceosome.

A role for EOL-1 in degrading RNAs lacking cap-adjacent 2`-O-ribose

methylation

Since cOMe plays an essential physiological role in *C. elegans*, this has allowed us

to fully exploit the advantages of this genetically tractable model system to

understand the mechanistic basis of the cellular response to cOMe. Importantly, we

have identified EOL-1 as a key effector that recognises transcripts lacking cOMe. A

previous study showed that cOMe blocks the activity of mammalian DXO *in vitro*

(Picard-Jean et al., 2018). The genetic interaction between *cmtr-1* and *eol-1*

demonstrates that this interaction is physiologically significant. Since we find that

loss of EOL-1 effectively bypasses the requirement for CMTR-1, and hence cOMe,

our work provides clear support for the hypothesis that DXO proteins recognise and

destroy RNAs that lack cOMe (Figure 8). Thus, in animals lacking CMTR-1, all

cellular transcripts become potential targets, leading to changes in steady-state

transcript levels that we see correlated with defects in germline development.

What is the physiological significance of EOL-1 activity? In wild-type animals, most, if

not all, RNA polymerase II transcripts will possess this modification and thus would

be protected from EOL-1 degradation. Previous studies have proposed that DXO

proteins maintain RNA quality control, removing partially capped RNAs that lack

cOMe before they enter the cytoplasm (Jiao et al., 2013). They have also been

shown to have a high affinity for NAD-capped RNAs, which are targeted by DXO for

rapid degradation (Kiledjian, 2018). Such roles would be consistent with our data.

However, since EOL-1 is a prominent component of the intracellular pathogen

response, a role in innate immunity is also plausible; indeed, previous studies of this

protein show that it is intimately associated with responses to pathogens and cellular

dysfunction.

EOL-1 was first identified in screens for mutants with enhanced aversive behavioural

responses to pathogenic bacteria (Shen et al., 2014), and it is a notable member of

the intracellular pathogen response, induced by Orsay virus, a positive sense, single-

stranded RNA virus, and the microsporidian *Nematocida parisii* (Chen et al., 2017;

Tecle et al., 2021). Mutations associated with mitochondrial dysfunction also result in

elevated *eol-1* expression levels (Mao et al., 2020; Senchuk et al., 2018), and EOL-1

protein undergoes changes in its intracellular localisation, forming cytoplasmic foci in

response to mitochondrial defects (Mao et al., 2020). However, a mechanistic role for

EOL-1 in innate immunity has yet to be defined. Our work connecting EOL-1 to

cOMe suggests a possible function for this protein in the surveillance for RNAs

lacking cap1 structures. Mitochondrial dysfunction leads to the presence of RNAs in

the cytoplasm that lack cOMe (Mao et al., 2020), which would presumably be

substrates for EOL-1 degradation. Moreover, mitochondrial RNAs also possess high

levels of NAD-caps, making them preferential DXO family substrates (Bird et al.,

2018; Jiao et al., 2017). Likewise, Orsay virus RNAs, for instance, lack both a 5`

monomethyl guanosine cap and cOMe modifications (Ashe et al., 2013). Thus, the

identification and removal of pathogenic RNAs on the basis of cOMe absence is a

plausible function of this enzyme in *C. elegans* that warrants further exploration.

The relationship we have discovered between cOMe and EOL-1 may have broader

phylogenetic significance, with DXO proteins having a conserved role in RNA

surveillance based on cOMe status. In particular, this may explain the molecular

basis of the lethality observed in mammalian embryos lacking either of the two

CMTRs (Dohnalkova et al., 2023); in the light of our work, perhaps loss of cOMe in

mouse embryos renders many cellular transcripts targets of mammalian DXO. While

DXO transcripts are not upregulated by the loss of cOMe (Dohnalkova et al., 2023),

unlike those of *eol-1* in *C. elegans*, cellular levels of DXO may still be sufficient to

have an impact across the transcriptome. Thus, the embryonic lethality observed in

embryos lacking cOMe might be explained by the cumulative downregulation of

multiple transcripts. It will be important to determine whether loss of DXO can

suppress the embryonic lethality of loss of *Cmtr1* in mice.

Cap-adjacent 2`-O-ribose methylation and the innate immune response

Our findings highlight similarities between the functions of cOMe in nematodes and

mammals: as in mammals, loss of cOMe leads to the upregulation of genes

associated with an innate transcriptional immune response. This raises the

possibility that cOMe is a determinant that regulates an RNA-mediated immune

response in *C. elegans*. The upregulation of IPR genes that we observed in animals

depleted of CMTR-1 is intriguing given that loss of CMTR1 in mammals also results

in induction of the innate immune response (Daffis et al., 2010; Dohnalkova et al.,

2023; Schuberth-Wagner et al., 2015). In mammals, RNAs lacking cOMe are directly

sensed by RIG-I, which binds to double-stranded RNA substrates that lack cOMe

and leads to the activation of the type I interferon response (Devarkar et al., 2016;

Schuberth-Wagner et al., 2015); although the mechanism by which cellular RNAs are

converted into double-stranded RNAs remains unclear.

In *C. elegans*, there are three RIG-I homologues, DRH-1, DRH-2 and DRH-3 (Guo et

al., 2013), but DRH-1 appears to be functionally most similar to RIG-I in that it

recognises viral RNAs and is essential for the intracellular pathogen response to viral

infection (Sowa et al., 2020). Indeed, the double-stranded RNA sensing module (the

helicase and C-terminal domains) of RIG-I can functionally substitute for the

homologous region of DRH-1 in assays for viral load (Guo et al., 2013); and native

DRH-1 binds to double-stranded RNAs in a manner similar to RIG-I (Consalvo et al.,

2024). Thus, if the upregulation of the intracellular pathogen response is a direct

effect of appearance of RNAs lacking cOMe, then DRH-1is a good candidate

involved in their detection; it will be important to test whether loss of *drh-1* results in

abrogation of the intracellular pathogen response in animals lacking cOMe.

An alternative possibility is that the induction of the intracellular pathogen response

by loss of cOMe is indirect, caused by the overall impact that this loss has on global

gene expression patterns. The intracellular pathogen response is also induced by

proteotoxic stress (Bakowski et al., 2014; Panek et al., 2020; Reddy et al., 2017), so

the global, cumulative effects on gene expression caused by loss of cOMe, might

lead to an intracellular pathogen response due to disruption of proteostasis.

Distinguishing between direct and indirect effects of cOMe on the transcriptional

immune response in *C. elegans* will be important in determining the extent of

functional conservation of cOMe function between nematodes and mammals.

Multiple studies in diverse experimental systems have described roles for cOMe at

multiple steps in the regulation of gene expression, including transcription, splicing

and translation (Dönmez et al., 2004; Drazkowska et al., 2022; Haussmann et al.,

2022; Kuge et al., 1998; Liang et al., 2022; Yu et al., 1998). It is thus surprising that

in *C. elegans* that lack EOL-1, cOMe is largely dispensable, at least in laboratory

culture. However, our results are consistent with work showing that in certain

mammalian cell types, the impact of loss of cOMe is relatively minor (Dohnalkova et

al., 2023). Similarly, loss of cOMe in *Drosophila* does not result in profound cellular

impacts, with such animals being viable and fertile (Haussmann et al., 2022). The

ubiquity of cOMe and the conservation of the two CMTR proteins that catalyse it

shows that it must have an essential role across animal phylogeny. Perhaps its

original, ancestral role was in RNA-based immune surveillance, which has been

preserved in *C. elegans* and mammals (but apparently lost in *Drosophila*), with its

functions in gene expression having arisen due cellular adaptation to the presence of

cOMe on cellular RNAs. Alternatively, it may have been co-opted into roles in innate

immunity through convergent evolution in mammals and nematodes from an

ancestral function(s) in the regulation of gene expression. Better understanding of

the molecular impacts of cOMe loss in diverse physiologically relevant experimental

systems will be needed to distinguish these hypotheses.

DATA AVAILABILITY

The mass spectrometry proteomics data have been deposited to the

ProteomeXchange Consortium via the PRIDE (Perez-Riverol et al., 2022) partner

repository with the dataset identifier PXD058991 (See also Supplementary Table 2).

The RNA-Seq data used for the CMTR-1 depletion differential gene expression

analysis have been deposited in ArrayExpress under accession number E-MTAB-

14816. Sequence reads that were used in the sibling selection whole genome

sequence mapping of *eol-1(fe152)* and *eol-1(fe154)* are available in the NCBI

Sequence Read Archive under BioProject ID PRJNA1216994.

1004

SUPPLEMENTARY DATA STATEMENT

1005

1006 Supplementary Data are available at NAR Online.

1007

ACKNOWLEDGEMENTS

We thank Omar Alotaibi, Holly Armstrong, Sandeep Goli, Phurichaya Khunthong,

and Ursula Terech Rial for their assistance with the *cmtr-1* suppressor screen. We

also thank Max Bracken-Zmuda for help making the *cmtr-1(feK244A)* strain. Some

strains were provided by the Caenorhabditis Genetics Center, which is funded by

NIH Office of Research Infrastructure Programs (P40 OD010440). Proteomic

analysis was performed by Aberdeen Proteomics, and we thank Kate Burgoyne and

Craig Pattinson for technical support. We thank WormBase for providing the

community resource that facilitated the interrogation of *C. elegans* molecular

genetics used in this work. We acknowledge the support of the Maxwell HPC

computer cluster funded by the University of Aberdeen.

1019

AUTHOR CONTRIBUTIONS

1020

J.P., B.M. and M.S. conceived the research and provided oversight and leadership

responsibility for the research activity planning and execution. All authors were

involved in performing experiments and data collection. M.W., B.M. and J.P.

designed and implemented the computational analyses. E.C., P.E., M.K., M.W., I.H.,

M.S., B.M. and J.P. prepared figures and tables. E.C., P.E., B.M. and J.P. wrote the

manuscript, and all authors reviewed and edited the final manuscript.

1027

1028 **FUNDING**

1029 This work was supported by the UKRI Biotechnology and Biological Sciences

1030 Research Council (BBSRC) through the EastBio Doctoral Training Partnership

1031 (BB/T00875X/1) and a project grant awarded to M.S. (BB/X008193/1).

REFERENCES

1032

Adams JM, Cory S. 1975. Modified nucleosides and bizarre 5`-termini in mouse

myeloma mRNA. *Nature* **255**:28–33.

Andrews S. 2010. FastQC: a quality control tool for high throughput sequence data.

https://www.bioinformatics.babraham.ac.uk/projects/fastqc/

Anreiter I, Mir Q, Simpson JT, Janga SC, Soller M. 2020. New Twists in Detecting

mRNA Modification Dynamics. *Trends Biotechnol.*

doi:10.1016/j.tibtech.2020.06.002

Anreiter I, Tian YW, Soller M. 2023. The cap epitranscriptome: Early directions to a

complex life as mRNA. *Bioessays* **45**:e2200198.

Arribere JA, Bell RT, Fu BXH, Artiles KL, Hartman PS, Fire AZ. 2014. Efficient

marker-free recovery of custom genetic modifications with CRISPR/Cas9 in

Caenorhabditis elegans. *Genetics* **198**:837–846.

Arribere JA, Cenik ES, Jain N, Hess GT, Lee CH, Bassik MC, Fire AZ. 2016.

Translation readthrough mitigation. *Nature* **534**:719–723.

Ashburner M, Ball CA, Blake JA, Botstein D, Butler H, Cherry JM, Davis AP, Dolinski

K, Dwight SS, Eppig JT, Harris MA, Hill DP, Issel-Tarver L, Kasarskis A, Lewis

S, Matese JC, Richardson JE, Ringwald M, Rubin GM, Sherlock G. 2000.

Gene ontology: tool for the unification of biology. The Gene Ontology

Consortium. *Nat Genet* **25**:25–29.

Ashe A, Bélicard T, Le Pen J, Sarkies P, Frézal L, Lehrbach N, Félix M, Miska E.

2013. A deletion polymorphism in the *Caenorhabditis elegans* RIG-I homolog

disables viral RNA dicing and antiviral immunity. *Elife* **2**.

doi:10.7554/eLife.00994

1056 Bakowski MA, Desjardins CA, Smelkinson MG, Dunbar TL, Lopez-Moyado IF, Rifkin

1057   SA, Cuomo CA, Troemel ER. 2014. Ubiquitin-mediated response to

1058   microsporidia and virus infection in *C. elegans*. *PLoS Pathog* **10**:e1004200.

1059 Bélanger F, Stepinski J, Darzynkiewicz E, Pelletier J. 2010. Characterization of

1060   hMTr1, a human Cap1 2`-O-ribose methyltransferase. *J Biol Chem*

1061   **285**:33037–33044.

1062 Benjamini Y, Hochberg Y. 1995. Controlling the false discovery rate: A practical and

1063   powerful approach to multiple testing. *J R Stat Soc Series B Stat Methodol*

1064   **57**:289–300.

1065 Bird JG, Basu U, Kuster D, Ramachandran A, Grudzien-Nogalska E, Towheed A,

1066   Wallace DC, Kiledjian M, Temiakov D, Patel SS, Ebright RH, Nickels BE. 2018.

1067   Highly efficient 5` capping of mitochondrial RNA with NAD+ and NADH by

1068   yeast and human mitochondrial RNA polymerase. *Elife* **7**.

1069   doi:10.7554/eLife.42179

1070 Chang JH, Jiao X, Chiba K, Oh C, Martin CE, Kiledjian M, Tong L. 2012. Dxo1 is a

1071   new type of eukaryotic enzyme with both decapping and 5`-3`

1072   exoribonuclease activity. *Nat Struct Mol Biol* **19**:1011–1017.

1073 Chen K, Franz CJ, Jiang H, Jiang Y, Wang D. 2017. An evolutionarily conserved

1074   transcriptional response to viral infection in Caenorhabditis nematodes. *BMC*

1075   *Genomics* **18**:303.

1076 Cingolani P, Platts A, Wang LL, Coon M, Nguyen T, Wang L, Land SJ, Lu X, Ruden

1077   DM. 2012. A program for annotating and predicting the effects of single

1078   nucleotide polymorphisms, SnpEff. *Fly (Austin)* **6**:80–92.

Consalvo CD, Aderounmu AM, Donelick HM, Aruscavage PJ, Eckert DM, Shen PS,

Bass BL. 2024. *Caenorhabditis elegans* Dicer acts with the RIG-I-like helicase

DRH-1 and RDE-4 to cleave dsRNA. *Elife* **13**. doi:10.7554/eLife.93979

Daffis S, Szretter KJ, Schriewer J, Li J, Youn S, Errett J, Lin T-Y, Schneller S, Zust R,

Dong H, Thiel V, Sen GC, Fensterl V, Klimstra WB, Pierson TC, Buller RM,

Gale M Jr, Shi P-Y, Diamond MS. 2010. 2`-O methylation of the viral mRNA

cap evades host restriction by IFIT family members. *Nature* **468**:452–456.

Danecek P, Bonfield JK, Liddle J, Marshall J, Ohan V, Pollard MO, Whitwham A,

Keane T, McCarthy SA, Davies RM, Li H. 2021. Twelve years of SAMtools

and BCFtools. *Gigascience* **10**:giab008.

DeMott E, Dickinson D, Doonan R. 2021. Highly improved cloning efficiency for

plasmid-based CRISPR knock-in in *C. elegans*. *MicroPubl Biol* **2021**.

doi:10.17912/micropub.biology.000499

Despic V, Jaffrey SR. 2023. mRNA ageing shapes the Cap2 methylome in

mammalian mRNA. *Nature* **614**:358–366.

Devarkar SC, Wang C, Miller MT, Ramanathan A, Jiang F, Khan AG, Patel SS,

Marcotrigiano J. 2016. Structural basis for m7G recognition and 2`-O-methyl

discrimination in capped RNAs by the innate immune receptor RIG-I. *Proc*

Natl Acad Sci U S A **113**:596–601.

Dickinson DJ, Pani AM, Heppert JK, Higgins CD, Goldstein B. 2015. Streamlined

Genome Engineering with a Self-Excising Drug Selection Cassette. *Genetics*

200:1035–1049.

Dix TC, Haussmann IU, Brivio S, Nallasivan MP, HadzHiev Y, Müller F, Müller B,

Pettitt J, Soller M. 2022. CMTr mediated 2`-O-ribose methylation status of

cap-adjacent nucleotides across animals. *RNA* **28**:1377–1390.

Dohnalkova M, Krasnykov K, Mendel M, Li L, Panasenko O, Fleury-Olela F, Vågbø
    CB, Homolka D, Pillai RS. 2023. Essential roles of RNA cap-proximal ribose
    methylation in mammalian embryonic development and fertility. *Cell Rep*
    42:112786.

Doitsidou M, Poole RJ, Sarin S, Bigelow H, Hobert O. 2010. *C. elegans* mutant
    identification with a one-step whole-genome-sequencing and SNP mapping
    strategy. *PLoS One* **5**:e15435.

Dönmez G, Hartmuth K, Lührmann R. 2004. Modified nucleotides at the 5` end of
    human U2 snRNA are required for spliceosomal E-complex formation. *RNA*
    10:1925–1933.

Drazkowska K, Tomecki R, Warminski M, Baran N, Cysewski D, Depaix A, Kasprzyk
    R, Kowalska J, Jemielity J, Sikorski PJ. 2022. 2'-O-Methylation of the second
    transcribed nucleotide within the mRNA 5'`cap impacts the protein production
    level in a cell-specific manner and contributes to RNA immune evasion.
    Nucleic Acids Res **50**:9051–9071.

Eijlers P, Al-Khafaji M, Soto-Martin E, Fasimoye R, Stead D, Wenzel M, Müller B,
    Pettitt J. 2024. A nematode-specific ribonucleoprotein complex mediates
    interactions between the major nematode spliced leader snRNP and its target
    pre-mRNAs. *Nucleic Acids Res* **52**:7245–7260.

Ellis RE. 2022. Sex Determination in Nematode Germ Cells. *Sex Dev* **16**:305–322.

Ewels P, Magnusson M, Lundin S, Käller M. 2016. MultiQC: summarize analysis
    results for multiple tools and samples in a single report. *Bioinformatics*
    32:3047–3048.

Fasimoye RY, Spencer REB, Soto-Martin E, Eijlers P, Elmassoudi H, Brivio S,
    Mangana C, Sabele V, Rechtorikova R, Wenzel M, Connolly B, Pettitt J,

1129  Müller B. 2022. A novel, essential *trans*-splicing protein connects the

1130  nematode SL1 snRNP to the CBC-ARS2 complex. *Nucleic Acids Res*

1131  **50**:7591–7607.

1132 Feder M, Pas J, Wyrwicz LS, Bujnicki JM. 2003. Molecular phylogenetics of the

1133  RrmJ/fibrillarin superfamily of ribose 2`-O-methyltransferases. *Gene* **302**:129–

1134  138.

1135 Furuichi Y, Morgan M, Shatkin AJ, Jelinek W, Salditt-Georgieff M, Darnell JE. 1975.

1136  Methylated, blocked 5 termini in HeLa cell mRNA. *Proc Natl Acad Sci U S A*

1137  **72**:1904–1908.

1138 Galloway A, Cowling VH. 2019. mRNA cap regulation in mammalian cell function and

1139  fate. *Biochimica et Biophysica Acta (BBA) - Gene Regulatory Mechanisms*

1140  **1862**:270–279.

1141 Garrison E, Marth G. 2012. Haplotype-based variant detection from short-read

1142  sequencing. *arXiv:12073907 [q-bioGN]*.

1143 Gene Ontology Consortium, Aleksander SA, Balhoff J, Carbon S, Cherry JM,

1144  Drabkin HJ, Ebert D, Feuermann M, Gaudet P, Harris NL, Hill DP, Lee R, Mi H,

1145  Moxon S, Mungall CJ, Muruganugan A, Mushayahama T, Sternberg PW,

1146  Thomas PD, Van Auken K, Ramsey J, Siegele DA, Chisholm RL, Fey P,

1147  Aspromonte MC, Nugnes MV, Quaglia F, Tosatto S, Giglio M, Nadendla S,

1148  Antonazzo G, Attrill H, Dos Santos G, Marygold S, Strelets V, Tabone CJ,

1149  Thurmond J, Zhou P, Ahmed SH, Asanitthong P, Luna Buitrago D, Erdol MN,

1150  Gage MC, Ali Kadhum M, Li KYC, Long M, Michalak A, Pesala A, Pritazahra A,

1151  Saverimuttu SCC, Su R, Thurlow KE, Lovering RC, Logie C, Oliferenko S,

1152  Blake J, Christie K, Corbani L, Dolan ME, Drabkin HJ, Hill DP, Ni L, Sitnikov D,

1153  Smith C, Cuzick A, Seager J, Cooper L, Elser J, Jaiswal P, Gupta P, Jaiswal P,

Naithani S, Lera-Ramirez M, Rutherford K, Wood V, De Pons JL, Dwinell MR,

Hayman GT, Kaldunski ML, Kwitek AE, Laulederkind SJF, Tutaj MA, Vedi M,

Wang S-J, D'Eustachio P, Aimo L, Axelsen K, Bridge A, Hyka-Nouspikel N,

Morgat A, Aleksander SA, Cherry JM, Engel SR, Karra K, Miyasato SR, Nash

RS, Skrzypek MS, Weng S, Wong ED, Bakker E, Berardini TZ, Reiser L,

Auchincloss A, Axelsen K, Argoud-Puy G, Blatter M-C, Boutet E, Breuza L,

Bridge A, Casals-Casas C, Coudert E, Estreicher A, Livia Famiglietti M,

Feuermann M, Gos A, Gruaz-Gumowski N, Hulo C, Hyka-Nouspikel N, Jungo

F, Le Mercier P, Lieberherr D, Masson P, Morgat A, Pedruzzi I, Pourcel L,

Poux S, Rivoire C, Sundaram S, Bateman A, Bowler-Barnett E, Bye-A-Jee H,

Denny P, Ignatchenko A, Ishtiaq R, Lock A, Lussi Y, Magrane M, Martin MJ,

Orchard S, Raposo P, Speretta E, Tyagi N, Warner K, Zaru R, Diehl AD, Lee R,

Chan J, Diamantakis S, Raciti D, Zarowiecki M, Fisher M, James-Zorn C,

Ponferrada V, Zorn A, Ramachandran S, Ruzicka L, Westerfield M. 2023. The

Gene Ontology knowledgebase in 2023. *Genetics* **224**.

doi:10.1093/genetics/iyad031

Guo X, Zhang R, Wang J, Ding S-W, Lu R. 2013. Homologous RIG-I-like helicase

proteins direct RNAi-mediated antiviral immunity in *C. elegans* by distinct

mechanisms. *Proc Natl Acad Sci U S A* **110**:16085–16090.

Haline-Vaz T, Silva TCL, Zanchin NIT. 2008. The human interferon-regulated ISG95

protein interacts with RNA polymerase II and shows methyltransferase activity.

Biochem Biophys Res Commun **372**:719–724.

Haussmann IU, Wu Y, Nallasivan MP, Archer N, Bodi Z, Hebenstreit D, Waddell S,

Fray R, Soller M. 2022. CMTr cap-adjacent 2`-O-ribose mRNA

1178 methyltransferases are required for reward learning and mRNA localization to

1179 synapses. *Nat Commun* **13**:1209.

1180 Hills-Muckey K, Martinez MAQ, Stec N, Hebbar S, Saldanha J, Medwig-Kinney TN,

1181 Moore FEQ, Ivanova M, Morao A, Ward JD, Moss EG, Ercan S, Zinovyeva AY,

1182 Matus DQ, Hammell CM. 2022. An engineered, orthogonal auxin

1183 analog/AtTIR1(F79G) pairing improves both specificity and efficacy of the

1184 auxin degradation system in *Caenorhabditis elegans. Genetics* **220**.

1185 doi:10.1093/genetics/iyab174

1186 Hyde JL, Diamond MS. 2015. Innate immune restriction and antagonism of viral RNA

1187 lacking 2`-O methylation. *Virology* **479–480**:66–74.

1188 Inesta-Vaquera F, Chaugule VK, Galloway A, Chandler L, Rojas-Fernandez A,

1189 Weidlich S, Peggie M, Cowling VH. 2018. DHX15 regulates CMTR1-

1190 dependent gene expression and cell proliferation. *Life Sci Alliance*

1191 **1**:e201800092.

1192 Jiao X, Chang JH, Kilic T, Tong L, Kiledjian M. 2013. A mammalian pre-mRNA 5`end

1193 capping quality control mechanism and an unexpected link of capping to pre-

1194 mRNA processing. *Mol Cell* **50**:104–115.

1195 Jiao X, Doamekpor SK, Bird JG, Nickels BE, Tong L, Hart RP, Kiledjian M. 2017. 5'

1196 End Nicotinamide Adenine Dinucleotide Cap in Human Cells Promotes RNA

1197 Decay through DXO-Mediated deNADding. *Cell* **168**:1015-1027.e10.

1198 Joseph BB, Blouin NA, Fay DS. 2017. Use of a Sibling Subtraction Method (SSM) for

1199 Identifying Causal Mutations in *C. elegans* by Whole-Genome Sequencing.

1200 *G3* . doi:10.1534/g3.117.300135

1201 Keith G. 1995. Mobilities of modified ribonucleotides on two-dimensional cellulose

1202 thin-layer chromatography. *Biochimie* **77**:142–144.

Kelly WG, Xu S, Montgomery MK, Fire A. 1997. Distinct requirements for somatic

and germline expression of a generally expressed *Caernorhabditis elegans*

gene. *Genetics* **146**:227–238.

Kesner JS, Chen Z, Shi P, Aparicio AO, Murphy MR, Guo Y, Trehan A, Lipponen JE,

Recinos Y, Myeku N, Wu X. 2023. Noncoding translation mitigation. *Nature*

617:395–402.

Kiledjian M. 2018. Eukaryotic RNA 5`-End NAD+ Capping and DeNADding. *Trends*

Cell Biol **28**:454–464.

Kim D, Langmead B, Salzberg SL. 2015. HISAT: a fast spliced aligner with low

memory requirements. *Nat Methods* **12**:357–360.

Kimble J, Crittenden SL. 2007. Controls of germline stem cells, entry into meiosis,

and the sperm/oocyte decision in *Caenorhabditis elegans*. *Annu Rev Cell Dev*

Biol **23**:405–433.

Krueger F. 2015. Trim Galore!: A wrapper around Cutadapt and FastQC to

consistently apply adapter and quality trimming to FastQ files, with extra

functionality for RRBS …. *Babraham Institute*.

Kruse S, Zhong S, Bodi Z, Button J, Alcocer MJC, Hayes CJ, Fray R. 2011. A novel

synthesis and detection method for cap-associated adenosine modifications in

mouse mRNA. *Sci Rep* **1**:126.

Kuge H, Brownlee GG, Gershon PD, Richter JD. 1998. Cap ribose methylation of c-

mos mRNA stimulates translation and oocyte maturation in Xenopus laevis.

Nucleic Acids Res **26**:3208–3214.

Lasda EL, Kuersten S, Blumenthal T. 2011. SL *trans*-splicing in a *Caenorhabditis*

elegans in vitro extract. *Cold Spring Harb Protoc* **2011**:db.prot5574.

Lažetić V, Batachari LE, Russell AB, Troemel ER. 2023. Similarities in the induction

of the intracellular pathogen response in *Caenorhabditis elegans* and the type

I interferon response in mammals. *Bioessays* **45**:e2300097.

Li H, Durbin R. 2010. Fast and accurate short read alignment with Burrows–Wheeler

transform. *Bioinformatics* **25**:1754–1760.

Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G,

Durbin R, 1000 Genome Project Data Processing Subgroup. 2009. The

Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**:2078–2079.

Li Y, Lu J, Han Y, Fan X, Ding S-W. 2013. RNA interference functions as an antiviral

immunity mechanism in mammals. *Science* **342**:231–234.

Liang S, Silva JC, Suska O, Lukoszek R, Almohammed R, Cowling VH. 2022.

CMTR1 is recruited to transcription start sites and promotes ribosomal protein

and histone gene expression in embryonic stem cells. *Nucleic Acids Res*.

doi:10.1093/nar/gkac122

Liao Y, Smyth GK, Shi W. 2014. featureCounts: an efficient general purpose program

for assigning sequence reads to genomic features. *Bioinformatics* **30**:923–930.

Love MI, Huber W, Anders S. 2014. Moderated estimation of fold change and

dispersion for RNA-seq data with DESeq2. *Genome Biol* **15**:550.

Mao K, Breen P, Ruvkun G. 2022. The *Caenorhabditis elegans* ARIP-4 DNA helicase

couples mitochondrial surveillance to immune, detoxification, and antiviral

pathways. *Proc Natl Acad Sci U S A* **119**:e2215966119.

Mao K, Breen P, Ruvkun G. 2020. Mitochondrial dysfunction induces RNA

interference in *C. elegans* through a pathway homologous to the mammalian

RIG-I antiviral response. *PLoS Biol* **18**:e3000996.

Meisel JD, Wiesenthal PP, Mootha VK, Ruvkun G. 2024. CMTR-1 RNA

methyltransferase mutations activate widespread expression of a

dopaminergic neuron-specific mitochondrial complex I gene. *Curr Biol*.

doi:10.1016/j.cub.2024.04.079

Mello CC, Kramer JM, Stinchcomb D, Ambros V. 1991. Efficient gene transfer in *C.*

elegans: extrachromosomal maintenance and integration of transforming

sequences. *EMBO J* **10**:3959–3970.

Mount SM, Anderson P. 2000. Expanding the definition of informational suppression.

Trends Genet **16**:157.

Müller MBD, Kasturi P, Jayaraj GG, Hartl FU. 2023. Mechanisms of readthrough

mitigation reveal principles of GCN1-mediated translational quality control.

Cell **186**:3227-3244.e20.

Nance J, Frøkjær-Jensen C. 2019. The *Caenorhabditis elegans* Transgenic Toolbox.

Genetics **212**:959–990.

Negishi T, Kitagawa S, Horii N, Tanaka Y, Haruta N, Sugimoto A, Sawa H, Hayashi

K-I, Harata M, Kanemaki MT. 2022. The auxin-inducible degron 2 (AID2)

system enables controlled protein knockdown during embryogenesis and

development in *Caenorhabditis elegans*. *Genetics* **220**.

doi:10.1093/genetics/iyab218

Panek J, Gang SS, Reddy KC, Luallen RJ, Fulzele A, Bennett EJ, Troemel ER. 2020.

A cullin-RING ubiquitin ligase promotes thermotolerance as part of the

intracellular pathogen response in *Caenorhabditis elegans*. *Proc Natl Acad*

Sci U S A **117**:7950–7960.

Perez-Riverol Y, Bai J, Bandla C, García-Seisdedos D, Hewapathirana S,

Kamatchinathan S, Kundu DJ, Prakash A, Frericks-Zipper A, Eisenacher M,

1276    Walzer M, Wang S, Brazma A, Vizcaíno JA. 2022. The PRIDE database

1277    resources in 2022: a hub for mass spectrometry-based proteomics evidences.

1278    *Nucleic Acids Res* **50**:D543–D552.

1279 Philippe L, Pandarakalam GC, Fasimoye R, Harrison N, Connolly B, Pettitt J, Müller

1280    B. 2017. An *in vivo* genetic screen for genes involved in spliced leader *trans*-

1281    splicing indicates a crucial role for continuous *de novo* spliced leader RNP

1282    assembly. *Nucleic Acids Res* **45**:8474–8483.

1283 Picard-Jean F, Brand C, Tremblay-Létourneau M, Allaire A, Beaudoin MC,

1284    Boudreault S, Duval C, Rainville-Sirois J, Robert F, Pelletier J, Geiss BJ,

1285    Bisaillon M. 2018. 2`-O-methylation of the mRNA cap protects RNAs from

1286    decapping and degradation by DXO. *PLoS One* **13**:e0193804.

1287 Reddy KC, Dror T, Sowa JN, Panek J, Chen K, Lim ES, Wang D, Troemel ER. 2017.

1288    An intracellular pathogen response pathway promotes proteostasis in *C.*

1289    *elegans*. *Curr Biol* **27**:3544-3553.e5.

1290 Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, Preibisch

1291    S, Rueden C, Saalfeld S, Schmid B, Tinevez J-Y, White DJ, Hartenstein V,

1292    Eliceiri K, Tomancak P, Cardona A. 2012. Fiji: an open-source platform for

1293    biological-image analysis. *Nat Methods* **9**:676–682.

1294 Schmittgen TD, Livak KJ. 2008. Analyzing real-time PCR data by the comparative

1295    C(T) method. *Nat Protoc* **3**:1101–1108.

1296 Schuberth-Wagner C, Ludwig J, Bruder AK, Herzner A-M, Zillinger T, Goldeck M,

1297    Schmidt T, Schmid-Burgk JL, Kerber R, Wolter S, Stümpel J-P, Roth A, Bartok

1298    E, Drosten C, Coch C, Hornung V, Barchet W, Kümmerer BM, Hartmann G,

1299    Schlee M. 2015. A conserved histidine in the RNA sensor RIG-I controls

1300    immune tolerance to N1-2`-O-methylated self RNA. *Immunity* **43**:41–51.

Senchuk MM, Dues DJ, Schaar CE, Johnson BK, Madaj ZB, Bowman MJ, Winn ME,

Van Raamsdonk JM. 2018. Activation of DAF-16/FOXO by reactive oxygen

species contributes to longevity in long-lived mitochondrial mutants in

Caenorhabditis elegans. *PLoS Genet* **14**:e1007268.

Shen Y, Zhang J, Calarco JA, Zhang Y. 2014. EOL-1, the homolog of the mammalian

Dom3Z, regulates olfactory learning in *C. elegans*. *J Neurosci* **34**:13364–

13370.

Simabuco FM, Pavan ICB, Pestana NF, Carvalho PC, Basei FL, Campos Granato D,

Paes Leme AF, Zanchin NIT. 2019. Interactome analysis of the human Cap-

specific mRNA (nucleoside-2`-O-)-methyltransferase 1 (hMTr1) protein. *J Cell*

Biochem **120**:5597–5611.

Sowa JN, Jiang H, Somasundaram L, Tecle E, Xu G, Wang D, Troemel ER. 2020.

The *Caenorhabditis elegans* RIG-I homolog DRH-1 mediates the intracellular

pathogen response upon viral infection. *J Virol* **94**. doi:10.1128/jvi.01173-19

Stiernagle T. 2006. Maintenance of *C. elegans*. *WormBook* 1–11.

Tan X, Calderon-Villalobos LIA, Sharon M, Zheng C, Robinson CV, Estelle M, Zheng

N. 2007. Mechanism of auxin perception by the TIR1 ubiquitin ligase. *Nature*

446:640–645.

Tecle E, Chhan CB, Franklin L, Underwood RS, Hanna-Rose W, Troemel ER. 2021.

The purine nucleoside phosphorylase *pnp-1* regulates epithelial cell

resistance to infection in *C. elegans*. *PLoS Pathog* **17**:e1009350.

The UniProt Consortium, Bateman A, Martin M-J, Orchard S, Magrane M, Ahmad S,

Alpi E, Bowler-Barnett EH, Britto R, Bye-A-Jee H, Cukura A, Denny P, Dogan

T, Ebenezer T, Fan J, Garmiri P, da Costa Gonzales LJ, Hatton-Ellis E,

Hussein A, Ignatchenko A, Insana G, Ishtiaq R, Joshi V, Jyothi D,

Kandasaamy S, Lock A, Luciani A, Lugaric M, Luo J, Lussi Y, MacDougall A,
Madeira F, Mahmoudy M, Mishra A, Moulang K, Nightingale A, Pundir S, Qi G,
Raj S, Raposo P, Rice DL, Saidi R, Santos R, Speretta E, Stephenson J,
Totoo P, Turner E, Tyagi N, Vasudev P, Warner K, Watkins X, Zaru R, Zellner
H, Bridge AJ, Aimo L, Argoud-Puy G, Auchincloss AH, Axelsen KB, Bansal P,
Baratin D, Batista Neto TM, Blatter M-C, Bolleman JT, Boutet E, Breuza L, Gil
BC, Casals-Casas C, Echioukh KC, Coudert E, Cuche B, de Castro E,
Estreicher A, Famiglietti ML, Feuermann M, Gasteiger E, Gaudet P, Gehant S,
Gerritsen V, Gos A, Gruaz N, Hulo C, Hyka-Nouspikel N, Jungo F, Kerhornou
A, Le Mercier P, Lieberherr D, Masson P, Morgat A, Muthukrishnan V,
Paesano S, Pedruzzi I, Pilbout S, Pourcel L, Poux S, Pozzato M, Pruess M,
Redaschi N, Rivoire C, Sigrist CJA, Sonesson K, Sundaram S, Wu CH, Arighi
CN, Arminski L, Chen C, Chen Y, Huang H, Laiho K, McGarvey P, Natale DA,
Ross K, Vinayaka CR, Wang Q, Wang Y, Zhang J. 2023. UniProt: the
Universal Protein Knowledgebase in 2023. *Nucleic Acids Res* **51**:D523–D531.

Toczydlowska-Socha D, Zielinska MM, Kurkowska M, Astha N, Almeida CF,
Stefaniak F, Purta E, Bujnicki JM. 2018. Human RNA cap1 methyltransferase
CMTr1 cooperates with RNA helicase DHX15 to modify RNAs with highly
structured 5` termini. *Philos Trans R Soc Lond B Biol Sci* **373**:20180161.

Tourasse NJ, Millet JRM, Dupuy D. 2017. Quantitative RNA-seq meta-analysis of
alternative exon usage in *C. elegans*. *Genome Res* **27**:2120–2128.

Wang VY-F, Jiao X, Kiledjian M, Tong L. 2015. Structural and biochemical studies of
the distinct activity profiles of Rai1 enzymes. *Nucleic Acids Res* **43**:6596–6606.

Werner M, Purta E, Kaminska KH, Cymerman IA, Campbell DA, Mittra B, Zamudio
JR, Sturm NR, Jaworski J, Bujnicki JM. 2011. 2`-O-ribose methylation of cap2

in human: function and evolution in a horizontally mobile family. *Nucleic Acids*

Res **39**:4756–4768.

Wickham H. 2016. Ggplot2: Elegant graphics for data analysis, 2nd ed, Use R!

Cham, Switzerland: Springer International Publishing.

Yu G, Li F, Qin Y, Bo X, Wu Y, Wang S. 2010. GOSemSim: an R package for

measuring semantic similarity among GO terms and gene products.

Bioinformatics **26**:976–978.

Yu G, Wang L-G, Han Y, He Q-Y. 2012. clusterProfiler: an R package for comparing

biological themes among gene clusters. *OMICS* **16**:284–287.

Yu YT, Shu MD, Steitz JA. 1998. Modifications of U2 snRNA are required for snRNP

assembly and pre-mRNA splicing. *EMBO J* **17**:5783–5795.

Zhu A, Ibrahim JG, Love MI. 2019. Heavy-tailed prior distributions for sequence

count data: removing the noise and preserving large differences.

Bioinformatics **35**:2084–2092.

Züst R, Cervantes-Barragan L, Habjan M, Maier R, Neuman BW, Ziebuhr J, Szretter

KJ, Baker SC, Barchet W, Diamond MS, Siddell SG, Ludewig B, Thiel V. 2011.

Ribose 2`-O-methylation provides a molecular signature for the distinction of

self and non-self mRNA dependent on the RNA sensor Mda5. *Nat Immunol*

12:137–143.

1370

1371

1372 **SUPPLEMENTARY FIGURES**

1373

Supplementary Figure 1. 5-Ph-IAA depletion of mNG^AID::CMTR-1. (A) PE1176

animals were treated with 5-Ph-IAA or control-treated and subjected to Western blotting at

the indicated time points. Proteins were detected using anti-NeonGreen and anti-GAPDH

antibodies. Protein levels of mNG^AID::CMTR-1 were standardised relative to GADPH, with

the controls set to 1. Data represents 2 biological replicates. (B) Representative Western blot

for the data shown in A. (C) Western blot of mNG^AID::CMTR-1 depletion of replicates (R1-

4) used to generate RNA-Seq data (see Figure 6). In B and C, "+" indicates 5-Ph-IAA treated

samples and "-" control samples, respectively.

1382

1383

Supplementary Figure 2. Secondary screen to distinguish suppressors epistatic to

cmtr-1(-) **from** *TIR1* **loss-of-function mutants.** (A) The starting strain for the suppressor

mutagenesis screen (PE1176) is sterile, while strains carrying suppressor mutations (m,

unshaded, dashed box) that are epistatic to *cmtr-1(-)* restore fertility (blue ovals indicate

eggs inside gravid hermaphrodites). (B) We predicted that TIR1 loss-of-function mutants

(unshaded, dashed box) would also restore fertility to *cmtr-1(-)* animals, since these would

prevent depletion of mNG^AID::CMTR-1. To distinguish between these two suppressor

classes, we conducted a secondary screen based on mNeonGreen fluorescence. This was

absent in PE1176 and epistatic suppressor strains (panels show distal gonad arms of adult

hermaphrodites; nuclear mCherry::HIS-11 fluorescence was unaffected and serves to locate

the germline nuclei), but was present in *TIR1* loss-of-function mutants. (C) Location of

selected *TIR1* loss-of-function mutations identified from secondary screening (magenta).

View of the TIR1-LRR domain showing the auxin and substrate-binding pocket

(https://doi.org/10.2210/pdb2P1Q/pdb; (Tan et al., 2007)). Auxin is shown in green, the three

extended loops key for the formation of the pocket are shown in blue.

1399

Figure 3. Strategy used to isolate strains for sibling-subtraction/whole genome
sequencing mapping. Suppressor strains were crossed to the transgenic rescue
line, PE1053, which is homozygous for *cmtr-1(syb3613)* but rescued by the *feEx342*
cmtr-1(+) transgene. The Sup and Non-Sup lines were established from single F2
animals on the basis that F2 suppressor homozygotes were not dependent on the
feEx342 cmtr-1(+) transgene, while the broods of non-suppressor homozygotes
resembled those of the PE1053 grandparent strain. Note, *feEx342* shows relatively
poor rescue of the *cmtr-1* loss-of-function phenotype, so some transgenic animals
are sterile. Transgenic animals were recognised on the basis of the *myo-*
2p::tdTomato expression in the pharynx, which is also carried on the *feEx342*

1410 extrachromosomal transgenic array. Blue ovals indicate eggs inside gravid

1411 hermaphrodites.

1412 **SUPPLEMENTARY TABLES**

1413 **Supplementary Table 1.**

1414 Strains and oligonucleotides.

1415

1416 **Supplementary Table 2.**

1417 GFP::CMTR-1 and GFP::CMTR-2 immunoprecipitation data.

1418

1419 **Supplementary Table 3.**

1420 PE1176 5-Ph-IAA treated versus control differential gene expression and GO

1421 enrichment analysis.

1422

1423 **Supplementary Table 4.**

1424 Variant mapping for *fe152* and *fe154*. Only variants found in 100% of Sup reads, and

1425 absent in 100% of Non-Sup reads are shown. Highlights indicate variants found in

1426 both *fe152* and *fe154* Sup pools.

1427