C. elegans **food choice exhibits effort discounting-like behavior**

Jonathan R. M. Millet, Serge Faumont, Aaron B. Schatz, Amanda M. White, Kathy D. Chicas-Cruz,

Shawn R. Lockery*

Institute of Neuroscience, 1254 University of Oregon, Eugene OR 97403, USA

*Correspondence: shawn@uoregon.edu

ABSTRACT

Cost-benefit decisions are ubiquitous in both human and animal behavior. Economists have

developed formal models of cost-benefit decision-making by focusing on discounting behavior,

the devaluation of a reward based on the costs associated with it. The phylogenetic limits of

discounting behavior remain unknown. Here, we provide evidence that the nematode *C.*

elegans exhibits behavior closely resembling effort discounting. Given a choice between food

options that are easy or difficult to consume, worms devalue the latter in a manner predicted by

economic models. We identified a plausible mechanism for this behavior based on differential

rates of leaving food patches and demonstrated that this mechanism is disrupted by deficits in

dopamine signaling, as in rodents. Together, these results establish *C. elegans* as a potential

invertebrate model for discounting behavior and set new phylogenetic bounds on this type of

cost-benefit decision-making.

INTRODUCTION

Cost-benefit decisions are ubiquitous in animal behavior, driven by universal evolutionary

pressures to maximize survival and reproductive success. Foraging, mate selection, parental

investment, and territoriality each demand a balance between costs, such as energy

expenditure and risk, and benefits such as resources, mates, and offspring. A better

understanding of cost-benefit decision-making, including its biological basis, would advance a

variety of fields, including economics, psychology, neuroscience, and behavioral ecology.

Economic theorists have developed a variety of formal models of cost-benefit decision-making

by focusing on *discounting* behavior (Samuelson, 1937; Mazur and Coe, 1987). In this context,

discounting refers to the tendency of individuals to *devalue* a reward based on the costs

associated with it. Costs commonly investigated include delay, risk, and effort. In general, as cost

increases, perceived or subjective value decreases. For example, in the case of delay

discounting, receiving a guaranteed payment of $35 after 30 days is worth less than receiving

$25 today if a person accepts the smaller immediate reward. In risk discounting, participants

might choose to receive $50 for certain rather than a 50% chance of receiving $100. In effort

discounting, participants might choose to receive $1 with no effort rather than $5 after

repeatedly squeezing a stiff handgrip (Stanek and Richter, 2021).

Studies of discounting are not limited to humans. The literature on discounting in rodents is

particularly rich, with multiple paradigms for all three forms of discounting (Bailey, Simpson and

Balsam, 2016a; Salamone *et al.*, 2018; Gray *et al.*, 2019; Castrellon *et al.*, 2021). In models of

delay discounting, for example, mice choose between low-reward and high-reward nose pokes,

where the higher reward comes with a delay (Isles *et al.*, 2004). Risk discounting can be

modeled by giving subjects a choice between a low-probability lever yielding a large reward (4

food pellets) and a high-probability lever yielding a small reward (1 food pellet) 100% of the

time (St. Onge, Chiu and Floresco, 2010). Effort discounting can be modeled by giving rodents a

choice between low-reward (2 food pellets) and high-reward (4 food pellets) arms of a T-maze

where a physical barrier must be climbed to obtain the higher reward (Salamone, Cousins and

Bucher, 1994). However, behaviorists have also observed discounting in several species of

nonhuman apes, monkeys, lemurs, jays, chickens, pigeons, honeybees, and other animals (Isles

et al., 2004; Green, Myerson and Calvert, 2010; Hayden, 2016).

The phylogenetic limits of discounting behavior are unknown. We therefore asked whether

foraging decisions in the nematode worm *C. elegans* exhibit discounting-like behavior. *C.*

elegans is an omnivorous bacterivore. Worms swallow bacteria through rhythmic contractions

of the pharynx, a tube-shaped muscular organ that delivers bacterial particles to the gut; each

pharyngeal contraction is called a pump. *C. elegans* primarily inhabits rotting plant material

such as decaying fruits and stems (Frézal and Félix, 2015). Its natural habitat contains thousands

of different bacterial species (Samuel *et al.*, 2016). Each species has a characteristic nutritional

quality, defined in terms of the growth rate of individual worms cultured on that species (Avery

and Shtonda, 2003; Samuel *et al.*, 2016). *C. elegans* food preferences are sensitive not only to

nutritional quality but also to food density in a manner that satisfies quantitative economic

criteria for choices based on subjective value (Katzen *et al.*, 2023). *C. elegans* foraging behavior

has been used to investigate the neuronal and genetic basis of exploration-exploitation trade-

offs (Bendesky *et al.*, 2011; Milward *et al.*, 2011), a classical example of cost-benefit decision-

making. In the laboratory, worms feed on patches of bacteria grown on agarose plates. They

leave and re-enter a patch many times. Leaving rates are low on high-quality lawns but increase

if the bacterial food is depleted or is difficult to ingest (Shtonda and Avery, 2006; Bendesky *et*

al., 2011; Milward *et al.*, 2011; Olofsson, 2014; Scheer and Bargmann, 2023). This response

pattern suggests that leaving promotes exploration to find a better food source.

The fact that *C. elegans* exhibits subjective-value based choice and cost-benefit feeding

decisions suggested to us that it might also exhibit discounting behavior. Of the various forms of

discounting, we focused on effort discounting as it uniquely does not require learned

instrumental responses, like lever pressing for rewards, which are not feasible in *C. elegans*. We

modeled our experimental procedure on rodent T-maze studies, where one arm requires more

effort to reach the food reward (Salamone *et al.*, 2018). We baited the arms of a miniature T-

maze with selected densities of a single species of bacteria. One arm contained normal bacteria,

while the other arm contained a bacterial preparation in which the cells had been elongated,

which we predicted would make them more effortful to ingest.

We found that worms exhibit choice behavior that closely resembles effort discounting. Given a

choice between equal densities of normal and elongated bacteria, worms prefer normal

bacteria. This preference can be reversed by presenting elongated bacteria at a higher density

than normal bacteria, indicating that high food density compensates for greater effort, as in

rodent models. Furthermore, worms can be made indifferent to the two food options by

adjusting relative food densities. It is therefore possible to measure the extent of devaluation

and accurately predict novel indifference points. Preference for normal bacteria can be

attributed to an increased patch-leaving rate on elongated food. Additionally, typical levels of

discounting require intact dopamine signaling. Our results expand the repertoire of cost-benefit

behaviors in *C. elegans* and demonstrate that lower invertebrates are capable of effort

discounting-like behavior.

RESULTS

Elongated bacteria require more effort to eat

We conceptualized feeding effort in *C. elegans* as the amount of pharyngeal muscle activity required to ingest a given quantity of food. The starting point of our experiments was to make bacteria more effortful to consume. Computer simulations of pharyngeal mechanics predict that large bacterial cells are swallowed less efficiently than small cells (Avery and Shtonda, 2003). Consistent with this result, worms develop more slowly when cultured on large bacteria, possibly because they are acquiring nutrients less efficiently. These findings led to the hypothesis that larger bacteria require more effort to eat, necessitating more pharyngeal pumps per unit of food consumed. To test this hypothesis, we artificially increased the size of an easily ingested bacterium, *Comamonas* spp., using the antibiotic cephalexin. In Gram-negative bacteria such as *Comamonas*, cephalexin interferes with bacterial cell septation during mitosis, resulting in elongated, spaghetti-like bacterial filaments that nevertheless remain edible (Millet *et al.*, 2022).

Pharyngeal pumping rates in normal and elongated bacteria were measured in the reference strain N2 by placing individual worms in a microfluidic channel filled with a bacterial suspension and fitted with electrodes that record the electrical signals associated with each pharyngeal contraction (Lockery *et al.*, 2012). Bacteria were washed with cephalexin-free buffer before being fed to worms. Pumping frequencies in normal and elongated bacteria were equivalent (Fig. 1A; Table 1, row 1) indicating that worms expend approximately the same amount of effort per unit time feeding on normal and elongated food. To test the hypothesis that elongation interferes with food absorption, we measured nutrient uptake in worms cultured on normal or elongated bacteria, using fat stores as a proxy for carbohydrate and lipid consumption. This was done by staining fat stores with the specific label Oil-Red-O (Soukas *et al.*, 2009) and quantifying the density of staining after a two-day culture period extending from the L1 to the L4 stage. We found that fat levels were significantly lower in worms cultured on elongated food (Fig. 1B2; Table 1, row 2). Given that pumping rate was unchanged in the presence of normal versus elongated bacteria (Fig. 1A), the lower fat stores observed in worms fed elongated bacteria

suggest that the amount of nutrition incorporated per pump is lower in worms feeding on

elongated versus normal bacteria. This supports our hypothesis that elongation impairs food

transport, requiring longer feeding bouts and thus increased muscular activity to consume an

equivalent amount of food.

Our effort-discounting paradigm assumes that the normal and elongated bacteria are equivalent

except for the amount of effort required to consume them. For example, elongation could affect

nutrient content or olfactory attractiveness of bacteria. *C. elegans* food preferences are

sensitive to the nutrient content of bacteria (Feng *et al.*, 2025). We therefore asked whether

cephalexin treatment affects nutrient content. We focused on key macronutrients:

carbohydrates, lipids, and proteins. For protein and carbohydrates, we used colorimetric assays

(Materials and Methods; Fig. 1C,D). For lipids, we used phase separation to harvest and

measure lipid content (Materials and Methods; Fig. 1E). In all three tests, we detected no

differences between control and cephalexin-treated bacteria (Table 1, rows 3-5). We conclude

that cephalexin treatment does not alter bacterial nutritional composition.

Worms are attracted to particular species of bacteria by the characteristic mixture of volatile

organic compounds each species emits (Worthy *et al.*, 2018). Thus, another concern is that

cephalexin treatment might alter the odors produced by bacteria thereby changing their

attractiveness. To control for this possibility, we performed a T-maze assay (Materials and

Methods) in which patches of normal and elongated bacteria were placed at the ends of each

respective arm. The patches contained equal volumes of bacteria at an optical density (OD) of

1.0. Patches were spiked with sodium azide, a paralytic agent, to trap worms once they reached

a patch. Worms were placed at the base of the T-maze and allowed to explore it for 60 minutes

after which the number of worms in each patch was counted. A food preference index ($-1 \leq$

$I \leq 1$) was calculated, where positive and negative values indicate preference for normal and

elongated bacteria respectively, and $I \cong 0$ denotes indifference between the two options

(Materials and Methods). We found that worms under these conditions were indifferent (Fig.

1F2; Table 1, row 6). This result indicates that cephalexin treatment does not alter the relative

attractiveness of normal or elongated bacteria.

Figure 1

Figure 1. Establishment of an effort discounting paradigm in C. *elegans*.
A. Pharyngeal pumping frequency in suspensions of normal (CON) and elongated (CEPH) bacteria.
Symbols (box and whisker plot): *filled circles*, single-worm data points; *diamonds*, means;
numbers, sample size.
B1. Quantification of Oil-Red-O staining of lipid in worms cultured on normal (CON) or elongated
(CEPH) bacteria. Median worms from each condition are shown.

B2. Mean pixel intensity of Oil-Red-O staining of lipid in worms cultured on normal (CON) or
elongated (CEPH) bacteria. *Diamonds*, means; *numbers*, sample size.
C-D. Protein and carbohydrate in worms cultured on normal (CON) and elongated bacteria
(CEPH). Columns of data points of the same color are biological replicates run in parallel.
Symbols: *filled circles*, technical replicates (n = 6); *diamonds*, means of technical replicates.
E. Lipid concentration in worms cultured on normal (CON) and elongated bacteria (CEPH).
Symbols: *filled circles*, biological replicates (n = 3); *diamonds*, means of biological replicates.
F1. T-maze assay for relative attractiveness of normal and elongated bacteria. Food patches
contained sodium azide as a paralytic agent.
F2. Mean preference index in the T-maze assay. Error bars, ± 95% CI.

Worms devalue food that requires more effort to eat

Having demonstrated that elongated bacteria require more effort to eat are but equivalent to normal bacteria in terms of nutrition and attractiveness, we next asked whether *C. elegans* exhibits a preference for the normal, easier-to-eat bacteria – evidence of effort discounting-like behavior. We tested this by measuring food preference in the T-maze assay using the same methodology as in Fig. 1F but without the paralytic agent, allowing individual worms to sample both patches. One arm of the T-maze was baited with normal bacteria at an OD of 1.0, while the other arm contained elongated bacteria at ODs of 1.0, 1.5, or 2.0. Worms explored the maze for 60 minutes and the food preference index was computed based on worm counts at the end of this period. There was a significant overall effect of optical density of elongated bacteria on preference (Fig. 2B; Table 1, row 7). When normal and elongated bacteria were at the same concentration (OD 1), worms preferred normal bacteria ($I = 0.10$; Fig. 2B; Table 1, row 8). This demonstrates that worms do, in fact, devalue elongated food. When normal and elongated bacteria were at OD 1.0 and 2.0, respectively, worms preferred elongated bacteria ($I = -0.17$; Fig. 2B; Table 1, row 9). This suggests that, in the worm's valuation system, higher food density compensates for greater effort. This outcome was not a foregone conclusion. For instance, it was possible that no amount of elongated food would be sufficient to make worms prefer it. Our results are consistent with findings in rodents, where high-effort options are chosen when the food reward density is sufficiently high (Floresco, Tse and Ghods-Sharifi, 2008). When the optical densities of normal and elongated food were 1.0 and 1.5, respectively, worms were indifferent between the two options ($I = 0.03$; Fig. 2B; Table 1, row 10). Indifference points are also observed in rodent studies of effort discounting (Salamone, Cousins and Bucher, 1994; van den Bos *et al.*, 2006; Ostrander *et al.*, 2011). Overall, we conclude that in feeding decisions, worms take into account both relative effort and relative food density.

An economic model predicts novel indifference points

Indifference points are significant because, by definition, the presented options have equal perceived value, allowing for the quantification of devaluation. This quantification is only possible at an indifference point. For example, consider a hypothetical human scenario in which

a participant is asked to choose between receiving $20 immediately or $20 with the added cost

of climbing 20 flights of stairs. Most participants would choose the immediate $20. Given a

choice between $20 immediately and $100 at the cost of climbing 20 flights of stairs, some

participants might choose the $100 option. However, suppose that when given a choice

between $20 immediately and $80 at the cost of climbing 20 flights of stairs, a particular

participant is indifferent between the two options. For this person, under these circumstances,

we can conclude that the $20 option is perceived as equal in value to the $80 option. In other

words, the $80 option has been discounted (devalued) to $20.

Formally, at an indifference point,

$$V(E) = d(E)V_0$$

221   where E represents effort, V_0 represents the original value of the reward before devaluation

222   and $V(E)$ is the discounted value of V_0. The quantity $d(E)$ is the discount factor, which, in

223   economic models of discounting, is a function of E. From the above equation,

$$d(E) = \frac{V(E)}{V_0}$$

with $0 \le d(E) \le 1$; in the above example $d(E)$ = 20/80 or 1/4. In economic models, $d(E)$ is

inversely related to E but the precise form of $d(E)$ varies between models (Figure 2—figure

supplement 1).

In our experiments, E is determined by the degree of bacterial elongation. Although this

quantity is unknown, it remained unchanged because we used the same cephalexin exposure

protocol across experiments (see Materials and Methods). It is analogous to the fixed effort of

climbing 20 flights of stairs in the hypothetical example. Assuming a linear relationship between

optical density and perceived value, we can determine $d(E)$ from the indifference point shown

in Fig. 2B, where V_0 = OD 1.5 and $V(E)$ = OD 1.0, yielding $d(E)$ = 2/3. In the most basic models

of discounting, $d(E)$ is independent of V_0. Assuming independence, other values of V_0 can be

inserted into equation 1 to predict novel indifference points. For instance, setting V_0 = OD 3

predicts indifference at $V(E)$ = OD 2, and setting V_0 = OD 0.75 predicts indifference at $V(E)$ =

OD 0.5. That is, any pair of optical densities (normal versus elongated) in a 2:3 ratio should be

an indifference point. We tested these predictions by measuring preference in T-mazes baited

with normal and elongated food at OD ratios (normal:elongated) of 2:3 and 0.5:0.75, finding

indifference as predicted (Fig. 2C, Table 1, rows 11-14). The fact that standard discounting

models correctly predict indifference points reinforces the conclusion that *C. elegans* exhibits

discounting-like behavior.

Figure 2

Figure 2. Food preference in T-maze assays.

A. T-maze assay for preference for normal (Control) versus elongated bacteria (cephalexin).

B. Effect on preference of raising the optical density of elongated bacteria (CEPH). Mean preference index is shown for assays in which three different optical densities (OD) of elongated bacteria (CEPH) were paired with normal bacteria (CON) at OD 1.0. Worms are indifferent when elongated bacteria are at OD 1.5. *Numbers*, sample size. Error bars ± 95% CI

252   C. Tests of predicted indifference points. Mean preference index is shown for three different
253   pairs of optical densities of normal (CON) and elongated (CEPH) bacteria. The grey bar is the
254   indifference point found in panel B. The black bars are novel indifference points found by scaling
255   the optical density of elongated bacteria by the discount factor of 2/3 obtained from the grey
256   bar. *Numbers*, sample size. Error bars ± 95% CI.
257

    Figure 2—figure supplement 1

Model	Equation	Discount factor
Hyperbolic	$V(E) = \dfrac{V_0}{1 + \alpha E}$	$\dfrac{1}{1 + \alpha E}$
Generalized hyperbolic	$V(E) = \dfrac{V_0}{(1 + \alpha E)^s}$	$\dfrac{1}{(1 + \alpha E)^s}$
Exponential	$V(E) = \dfrac{V_0}{e^{\alpha E}}$	$\dfrac{1}{e^{\alpha E}}$

Figure 2—figure supplement 1. Standard models of effort discounting.
There are three main models of effort discounting, each with a unique equation relating effort
and reward. V_0 is the value of the reward if it were given effort-free. E is the level of effort
required to obtain the effortful reward. $V(E)$ is the discounted value of V_0 under conditions of
indifference, given level of effort E. The quantity α is a positive constant representing the
chooser's sensitivity to effort. The quantity s in the generalized hyperbolic model is the shape
parameter. Discount factors are constant at fixed E. In the graph, V_0 = 100, α = 0.05, and s = 1.5.
266

Effort discounting-like behavior is based on assessment of the local food environment

We next investigated the behavioral mechanism through which effort influences food preference, examining how worms interact with patches of normal versus elongated food. We studied these interactions at the microscopic level of individual worms to explain behavior at the macroscopic level of T-maze assays. This was done by performing food-patch leaving assays (Scheer and Bargmann, 2023) using either normal or elongated bacteria. Individual worms were transferred to a 1 cm diameter, nematode growth medium-filled arena containing a small patch (3–4 mm diameter) of normal or elongated bacteria (OD 1.0) at its center. The patch size was comparable to those in the T-maze assay. Each worm was video recorded for 15 minutes while it entered and left the patch multiple times (Fig. 3A).

For analysis, we took a kinetic approach to investigate the rates at which worms stochastically changed their position relative to the food patch. This approach is both simple, as it does not require assigning specific locomotory states at particular times to individual worms, and powerful, as it forms a bridge between the microscopic and macroscopic behavior of a stochastic system, thereby elucidating underlying dynamics. Accordingly, we defined three kinetic states based on occupancy of specific zones in the arena (Fig. 3B):

1. State F, *On Food*: both head and tail of the worm were in contact with the food patch.
2. State B, *At Border*: either the head or tail was in contact with the food patch, but not both.
3. State O, *Off Food*: neither head nor tail was in contact with the food patch.

This system contains four state transitions: F→B, B→O, O→B, and B→F. Behavior was analyzed in terms of state probabilities (P_F, P_B, P_O) and rate constants ($k_{FB}, k_{BO}, k_{OB}, k_{BF}$). State probability was defined as the fraction of total observation time the worm was in each state. Rate constants were defined as the number of transitions divided by the total time spent in the state of origin. Rate constants determine the probability per unit time of transitioning from one state to another and thus represent the underlying physical causes of state transitions.

We observed a variety of changes in kinetic parameters on elongated versus normal food. P_{B}

and P_{O} increased at the expense of P_{F}, consistent with decreased preference for elongated food

in T-maze assays (Fig. 3C; Table 1, rows 15,16). The rate constants exhibited two key features.

First, in control animals, between any pair of states, the mean rate constants leading to patch

entry (O→B→F transitions) were greater than the mean rate constants leading to patch leaving

(F→B→O transitions). In particular, $k_{\mathrm{BF}} > k_{\mathrm{FB}}$ and $k_{\mathrm{OB}} > k_{\mathrm{BO}}$ (Fig. D3 vs. D1 and Fig. D4 vs.

D2; Table 1, rows 17-23). This indicates a greater propensity to enter food patches than to leave

them, which is consistent with efficient foraging. Second, changes in rate constants brought

about by elongated food exhibited a distinctive pattern: the rates for patch leaving ($k_{\mathrm{FB}}, k_{\mathrm{BO}}$)

were increased whereas the rates for patch entry ($k_{\mathrm{OB}}, k_{\mathrm{BF}}$) were unchanged (Fig. 3D; Table 1,

rows 24-28). Thus, worms exhibited a heightened propensity to leave elongated food, consistent

with previous observations (Scheer and Bargmann, 2023). The fact that rates for patch entry

were unaffected by elongated bacteria rules out a mechanism in which worms, exiting patches

of elongated food (which is relatively undesirable), become reluctant to re-enter them. Finally,

we found that speed on elongated food increased, whereas speed off food remained unchanged

(Fig. 3E; Table 1, rows 29-33). Applying the kinetic data of individual worms to the population

level in T-maze assays suggests that effort discounting-like behavior could result from feeding

decisions based on the local food environment rather than a direct comparison of patch

contents across the arms of the T-maze.

Figure 3

Figure 3. Kinetic analysis of food-patch leaving assays.
A. Food-patch leaving assays for normal (control) and elongated (cephalexin-treated) bacteria.
Open circles, arenas; *colored circles*, food. A representative track of a single worm is
superimposed on each diagram.
B. Three-state kinetic model of behavior in the assay. Each state represents the indicated zone in
the arena.

C. Probability of states F, B, and O for normal (CON) and elongated (CEPH) bacteria. Significance
was assessed using compositional statistics as described in Fig. 3—figure supplement 1.
D. Rate constants for normal (CON) and elongated (CEPH) bacteria.
E. Locomotion speed for normal (CON) and elongated (CEPH) bacteria on or off food.

Figure 3—figure supplement 1. Statistical test of the effect of cephalexin on state probabilities in Figure 3C.

State probabilities are an instance of *compositional data*, meaning data that sum to a constant (1 in this case). Such data require special treatment because a decrease in one variable means an increase in at least one other variable. Accordingly, data were first subjected to an isometric log ratio transform (see Materials and Methods). The transformation quantifies the change in *balance* between subsets of the data. Balances are indicated by the numbers in the figure. Here, the control balance between P_F and the reciprocal probabilities P_B and P_O (1.47) is decreased by cephalexin treatment (0.85). A *t*-test on individual worm balances revealed that P_F significantly decreased while the joint P_B and P_O data significantly increased. In contrast, a similar approach showed that the balance between P_B and P_O was not significantly affected. This indicates that P_B and P_O increased by similar amounts.

Intact dopamine signaling is required for effort discounting-like behavior

A salient aspect of effort discounting in rodents is its sensitivity to changes in midbrain

dopamine signaling (Salamone *et al.*, 2018). Increased signaling promotes preference for high-

value, high-effort rewards, whereas decreased signaling promotes preference for low-value,

low-effort rewards (Bailey, Simpson and Balsam, 2016b; Salamone *et al.*, 2018; Hart and

Izquierdo, 2019); in other words, increased dopamine signaling results in greater effort

tolerance. Dopamine signaling is a candidate mechanism for the regulation of effort

discounting-like behavior in *C. elegans* because it affects how worms respond to the presence

and absence of food as they enter or leave food patches. Worms slow down when they

encounter a food patch, a response that requires intact dopamine signaling (Sawin,

Ranganathan and Horvitz, 2000). When worms exhaust their food supply or are removed from a

food patch, they engage in a bout of frequent, high-angled turns called area-restricted search

(Hills, Brockie and Maricq, 2004). This response also requires dopamine signaling.

To test whether dopamine signaling is involved in effort discounting-like behavior in *C. elegans*,

we examined food preference in a variety of mutants with altered dopamine signaling. The gene

cat-2 encodes a tyrosine hydroxylase essential to the main synthesis pathway for dopamine; *cat-*

2 mutants are commonly used to reduce dopamine signaling in *C. elegans*. The gene *dat-1*

encodes a dopamine transporter that mediates the reuptake of dopamine into presynaptic

neurons; *dat-1* mutants are used to increase dopamine signaling. The effects of reduced

dopamine signaling can also be investigated in dopamine receptor mutants. *C. elegans* has four

genes encoding dopamine receptors with homology to dopamine receptors in mammals. Of

these, *dop-1*, *dop-2*, and *dop-3* are the most thoroughly characterized to date. *dop-1* is a DRD1

homolog, whereas *dop-2* and *dop-3* are DRD2 homologs. The genes *dop-1* and *dop-3* are

required for normal responses to the presence of food (Chase, Pepper and Koelle, 2004); *dop-2*

is required for increased levels of activity in the absence of food (Bastien *et al.*, 2024). We found

a significant overall effect of strain on food preference measured in the T-maze assay (Fig. 4;

Table 1, row 34). Post hoc testing showed that *cat-2*, *dat-1*, *dop-1*, and *dop-2* were

indistinguishable from N2, whereas *dop-3* mutants were significantly different from N2 (Table 1,

row 35). In particular, they were indifferent between the two food options, indicative of

decreased effort discounting-like behavior (Table 1, row 36). We conclude that fully intact

dopamine signaling is required for normal effort discounting-like behavior in *C. elegans*, as in

rodents. However, the fact that impaired dopamine signaling rendered worms indifferent

between the food options suggests that decreased dopamine signaling increases tolerance for

high-effort food, in contrast to its effect in rodents. The absence of an effect of the *cat-2* and

dat-1 mutations is puzzling in view of the requirement for *dop-3*. One possibility is that residual

dopamine in the *cat-2* mutant (Sanyal *et al.*, 2004) is sufficient for normal effort discounting-like

behavior and that normal levels of dopamine signaling are saturating, such that *dat-1* mutations

have no effect.

Figure 4

380 **Figure 4. Effect of dopamine signaling mutations on preference for normal versus elongated**
381 **bacteria**. Data are mean preferences measured in T-maze assays. Dopamine signaling was
382 reduced by a nonsense mutation in *cat-2* or by deletions of three dopamine receptor genes
383 (*dop*). Dopamine signaling was increased by a deletion in *dat-1*. *Numbers*, sample size. Error
384 bars ± 95% CI.

Effort discounting-like behavior is not an adaptation to the laboratory environment

Finally, we considered the question whether effort discounting-like behavior may reflect

adaptation to the laboratory environment. The reference strain used in this study, N2, has been

cultured for over 50 years in the laboratory setting, where it is typically fed small, easily

consumed bacteria, such as *E. coli* strain OP50. Consequently, it may have lost the ability to

consume larger bacteria efficiently. To address this question, we tested whether any natural

isolate strains would prefer normal over elongated bacteria. Accordingly, we performed T-maze

assays on five such strains with normal and elongated food at OD 1.0. Natural isolate strains

proved to be challenging to work with as they frequently escaped the maze or burrowed in the

substrate. These traits reduced the number of countable worms per assay, leading to

considerable variance in the data. Nevertheless, there was a significant overall effect of strain

on mean preference index (Fig. 5; Table 1, row 37). Post hoc testing revealed that only the strain

JU775 had a preference index that was significantly different from N2 (Table 1, row 38) and that

JU775's preference was not distinguishable from zero (Table 1, row 39). Thus, JU775 was

indifferent between normal and elongated food. On the other hand, mean preference of DL238

was both different from zero (Table 1, row 40) and indistinguishable from N2 (Table 1, row 38).

That is, DL238 behaved like N2. We conclude that N2's preference is not an adaptation to the

laboratory environment. Between DL238 and JU775, there was a range of different responses,

suggesting the possibility of genetic variation in preference for normal versus elongated food.

This could reflect differences in effort tolerance or ease of consumption of elongated bacteria.

Figure 5. Preference for normal versus elongated bacteria in N2 and five wild-isolate strains.
Data are mean preferences measured in T-maze assays. N2 and DL238 are significantly different
from zero. *Numbers*, sample size. Error bars ± 95% CI.
409
410
411
412

DISCUSSION

Effort discounting is a form of cost-benefit decision-making with implications for economics, psychology, neuroscience, behavioral ecology, and other fields. To investigate effort discounting in *C. elegans*, worms were given a choice between two patches of the same species of bacteria. One patch contained normal bacteria, whereas the other patch contained elongated bacteria, which we showed to be more effortful to consume yet similar to normal bacteria in nutrition and attractiveness. When the two types of bacteria were presented at equal density, worms preferred normal bacteria. Three lines of evidence suggest that this preference is analogous to the devaluation of a reward based on increased effort to obtain it: (1) Preference can be reversed by increasing the density of the effortful option. (2) The relative density of normal and effortful food can be titrated to an indifference point. (3) Widely used economic models of discounting correctly predict novel indifference points. Points (1) (Floresco, Tse and Ghods-Sharifi, 2008) and (2) (Salamone, Cousins and Bucher, 1994; van den Bos *et al.*, 2006; Ostrander *et al.*, 2011), together with the requirement for intact dopamine (Salamone *et al.*, 2018) signaling, establish similarities to effort discounting in other animal models. The demonstration of effort discounting-like behavior in a lower invertebrate sets a new phylogenetic boundary on discounting behavior.

Relationship to previous work

The decision of whether to stay or leave a patch of food has been well studied in *C. elegans* (Shtonda and Avery, 2006; Bendesky *et al.*, 2011; Milward *et al.*, 2011; Olofsson, 2014; Scheer and Bargmann, 2023). Leaving is rare on rich patches of normal food but increases on depleted patches or when the mechanics of feeding have been genetically impaired. Both conditions would be expected to decrease feeding efficiency and thereby increase the amount of muscular effort required to obtain food. It is likely that patch-leaving rate in these studies and ours is linked to an internal signal of feeding efficiency. The nature of this signal remains to be determined, but it likely integrates pharyngeal pumping rate with the rate of food accumulation in the gut.

An increase in patch-leaving frequency in response to elongated bacteria has been reported

previously (Scheer and Bargmann, 2023). There are some significant differences between that

study and ours: (1) *E. coli* was used instead of *Comamonas*; (2) bacteria were elongated with the

antibiotic aztreonam which renders food inedible to *C. elegans* (Gruninger, Gualberto and

Garcia, 2008; Ben Arous, Laffont and Chatenay, 2009), whereas the cephalexin treatment we

used resulted in elongated but edible food (Millet *et al.*, 2022); (3) worms were video-recorded

for 40 minutes instead of 15 minutes; (4) food-patch leaving was defined using different

behavioral criteria; (5) patch-entry rates were not considered. Nevertheless, our results confirm

and extend the previous findings by showing that the effect of elongated food on patch-leaving

rate is robust across bacterial species, varied degrees of edibility of elongated food, and

differences in details of behavioral analysis.

Utility of economic modeling

Although *C. elegans* patch-leaving behavior has been described in detail before, placing it in an

explicit economic context is new. This context provided a quantitative framework that enriched

our study in two respects. First, it gave us a metric – the discount factor – for quantifying the

extent to which effortful food was devalued by the worm. We observed that worms were

indifferent when normal and elongated bacteria were at ODs 1.0 and 1.5, respectively, yielding a

discount factor of 2/3. Second, this framework made testable predictions. At the level of effort

determined by our cephalexin exposure protocol, any combination of optical densities of

normal and elongated food in a 2:3 ratio should be an indifference point. This prediction was

verified by measuring preference under conditions in which the optical density of elongated

food was both higher and lower than the original value of 1.5. This result shows that standard

models of effort discounting may apply to *C. elegans*. It will now be interesting to test whether

the discount factor is sensitive to effort, as assumed in many discounting models (Figure 1—

figure supplement 1). This could be done by identifying indifference points when effort is

manipulated by changing cephalexin exposure conditions to increase or decrease the extent of

elongation.

A model of effort-discounting like behavior

We propose the following simple model of effort discounting-like behavior in T-maze assays

based on our kinetic analysis of patch entry and exit. In the T-maze assay, worms preferentially

accumulate in the patch containing normal food. In theory, this could be because worms leave

patches of elongated food (F →B→O transitions) at a higher rate, enter patches of elongated

food (O →B→F transitions) at a lower rate, or both. Our kinetic analysis favors the former. We

found that the rates governing patch entry, k_{OB} and k_{BF}, were unchanged by the presence of

elongated food, whereas the rates governing patch leaving, k_{FB} and k_{BO}, were increased. The

increase in these two rate constants is likely the result of increased roaming, a state of arousal

characterized by increased speed and a greater tendency to leave food patches (Fujiwara,

Sengupta and McIntire, 2002; Ben Arous, Laffont and Chatenay, 2009; Flavell *et al.*, 2013; Scheer

and Bargmann, 2023). Our observation that speed increased on elongated food (Fig. 3E) is

consistent with increased roaming on these patches in the T-maze.

Role of dopamine signaling in effort discounting-like behavior

Fully intact dopamine signaling is required for normal effort discounting-like behavior in *C.*

elegans. This finding is broadly consistent with the known requirements of dopamine signaling

for foraging behavior in *C. elegans*, as our T-maze assay is essentially a foraging assay. *C. elegans*

dopamine receptors DOP-1, DOP-2, and DOP-3 are required for normal responses to the

presence or absence of food (Chase and Koelle, 2007; Suo, Culotti and Van Tol, 2009; Bastien *et*

al., 2024). Dopamine itself is required for area-restricted search in which worms, suddenly

finding themselves outside a food patch, engage in a period of frequent locomotory

reorientations in an attempt to relocate the food (Hills, Brockie and Maricq, 2004). However, the

relationship between dopamine signaling and effort discounting-like behavior in *C. elegans*

remains an area for future research. The fact that decreased dopamine signaling in the *dop-3*

mutant rendered worms indifferent between the food options suggests that decreased

dopamine signaling *increases* tolerance for high-effort food, in contrast to its effects in rodents.

This discrepancy may reflect a true difference in the function of dopamine in effort discounting

in *C. elegans* and rodents, or methodological differences. Manipulations of dopamine signaling

in rodent studies mainly involve acute experiments using pharmacological agents, whereas we

studied worms with chronic, lifelong alterations in dopamine signaling.

In summary, our study provides strong evidence that *C. elegans* exhibits effort discounting-like

behavior, reinforcing the idea that decision-making strategies based on effort-cost trade-offs are

evolutionarily conserved. By demonstrating that worms devalue effortful food like higher

organisms, we extend the phylogenetic boundary of this behavior. The application of an

economic framework allowed us to quantify this effect and make testable predictions,

confirming its robustness. Furthermore, our findings highlight the role of dopamine signaling in

modulating effort sensitivity, suggesting potential parallels with vertebrate systems. Future

research can now exploit the utility of *C. elegans* as a genetically tractable system to explore

how environmental factors and genetic variations influence effort-based decision-making in *C.*

elegans, providing deeper insights into the fundamental principles and molecular basis of cost-

benefit decisions.

530 **TABLES**

Row	Figure	Test	Effect or comparison tested	Units of replication or sampling	Number of replicates	Statistic	Statistic value	DF 1 or combined DF	p	Effect size metric	Effect size
1	1A	Mann-Whitney	Cephalexin	Worms	99, 98	U	4178	–	9.23E-02	Rank-biserial correlation (r)	0.13 (small)
2	1B2	Mann-Whitney	Cephalexin	Worms	44, 49	U	618	–	**3.20E-04**	Rank-biserial correlation (r)	0.43 (moderate)
3	1C	t-Test	Cephalexin	Biol. Replicate	3, 3	t	0.10	2	9.28E-01	Cohen's d	0.08 (small)
4	1D	t-Test	Cephalexin	Biol. Replicate	3, 3	t	0.10	2	9.31E-01	Cohen's d	0.08 (small)
5	1E	t-Test	Cephalexin	Tech. Replicate	3, 3	t	0.12	2	9.91E-01	Cohen's d	0.01 (small)
6	1F2	t-Test	Mean not = 0	Assay plates	108	t	1.65	104	3.56E-01	Cohen's d	0.16 (small)
7	2B	1-way ANOVA	OD ratios	Assay plates	137, 106, 73	F	19.07	2	**1.52E-08**	Eta squared	0.11 (moderate)
8	2B	t-Test	Mean not = 0	Assay plates	137	t	3.20	136	**1.69E-03**	Cohen's d	0.27 (small)
9	2B	t-Test	Mean not = 0	Assay plates	73	t	1.99	72	**2.06E-07**	Cohen's d	0.67 (large)
10	2B	t-Test	Mean not = 0	Assay plates	106	t	1.08	105	2.82E-01	Cohen's d	0.10 (small)
11	2C	1-way ANOVA	OD ratios	Assay plates	120, 106, 132	F	0.83	2	4.37E-01	Eta squared	0.00
12	2C	t-Test	Mean not = 0	Assay plates	120	t	0.71	119	4.80E-01	Cohen's d	0.06 (small)
13	2C	t-Test	Mean not = 0	Assay plates	106	t	1.08	105	2.82E-01	Cohen's d	0.10 (small)
14	2C	t-Test	Mean not = 0	Assay plates	132	t	0.64	131	5.22E-01	Cohen's d	0.06 (small)
15	3C	t-Test	P_F vs (P_O, P_B)	Single worm	66, 60	t	4.22	123.57	**4.62E-05**	Cohen's d	0.75 (medium)
16	3C	t-Test	P_O vs P_B	Single worm	66, 60	t	0.57	116.99	5.73E-01	Cohen's d	0.10 (small)
17	3D	1-way ANOVA	Transition type	Single worm	66, 60	F	24.71	3	**4.22E-14**	Eta squared	0.22 (large)
18	3D	Tukey	k_{FB}, k_{BF}	Single worm	66, 60	q	3.68	122	**<1.00E-07**	Cohen's d	2.199391 (large)
19	3D	Tukey	k_{BO}, k_{OB}	Single worm	66, 60	q	3.68	122	**4.30E-06**	Cohen's d	0.68 (medium)
20	3D	Tukey	k_{BO}, k_{BF}	Single worm	66, 60	q	3.68	122	**6.00E-07**	Cohen's d	1.60 (large)
21	3D	Tukey	k_{OB}, k_{BF}	Single worm	66, 60	q	3.68	122	9.77E-01	Cohen's d	0.051 (negligible)
22	3D	Tukey	k_{FB}, k_{BO}	Single worm	66, 60	q	3.68	122	4.86E-01	Cohen's d	0.91 (large)
23	3D	Tukey	k_{OB}, k_{FB}	Single worm	66, 60	q	3.68	122	**0**	Cohen's d	0.88 (large)
24	3D	MANOVA	Cephalexin	Single worm	66, 60	F	16.44	4121	**8.77E-11**	Eta squared (partial)	0.35
25	3D1	t-Test	Cephalexin	Single worm	66, 60	t	7.38	79.39	**1.36E-10**	Cohen's d	1.36 (large)
26	3D2	t-Test	Cephalexin	Single worm	66, 60	t	3.32	118.69	**1.18E-03**	Cohen's d	0.60 (medium)
27	3D3	t-Test	Cephalexin	Single worm	66, 60	t	0.28	123.25	7.83E-01	Cohen's d	0.05 (negligible)
28	3D4	t-Test	Cephalexin	Single worm	66, 60	t	1.38	95.7	1.70E-01	Cohen's d	0.24 (small)
29	3E	2-way ANOVA	Cephalexin	Single worm	66, 60	F	5.24	1	**2.29E-02**	Eta squared	0.02 (small)
30	3E	2-way ANOVA	Food	Single worm	66, 60	F	0.70	1	4.00E-01	Eta squared	0.00 (negligible)
31	3E	2-way ANOVA	Cephalexin × Food	Single worm	66, 60	F	6.66	1	**1.10E-02**	Eta squared	0.03 (small)
32	3E	t-Test	Cephalexin	Single worm	66, 60	t	3.68	115.91	**4.59E-04**	Cohen's d	0.66 (medium)
33	3E	t-Test	Cephalexin	Single worm	66, 60	t	0.19	123.94	8.47E-01	Cohen's d	0.034 (negligible)
34	4	1-way ANOVA	Strain	Assay plates	137, 179, 165, 160, 175, 180	F	7.69	3	**4.71E-05**	Eta squared	0.03 (small)
35	4	Dunnett test	N2 vs other strains	Assay plates	See above	q	2.78	649	**1.92E-02**	Cohen's d	0.50 (medium)
36	4	t-Test	Mean not = 0	Assay plates	180	t	0.65	159	5.14E-01	Cohen's d	0.05 (small)
37	5	1-way ANOVA	Strain	Assay plates	137, 137, 129, 184, 178, 146	F	3.30	5	**5.83E-03**	Eta squared	0.018 (small)
38	5	Dunnett test	N2 vs other strains	Assay plates	See above	q	3.03	905	**1.10E-02**	Cohen's d	0.68 (medium)
39	5	t-Test	Mean not = 0	Assay plates	146	t	0.36	145	1.03E-01	Cohen's d	0.13 (small)
40	5	t-Test	Mean not = 0	Assay plates	137	t	2.02	136	**4.56E-02**	Cohen's d	0.17 (small)

532 **Table 1.** Statistical information. Bold p values denote significant effects.

534 **MATERIALS AND METHODS**

535 **Key resources table**

Reagent type (species) or resource	Designation	Source or reference	Identifiers	Additional information
Strain (*E. coli*)	OP50	CGC[1]	RRID:WB-STRAIN:WBStrain00041969	Worm maintenance
Strain, strain background (*Comamonas* spp.)	DA1877	CGC	RRID:WB- STRAIN:WBStrain00040995	Behavioral testing
Strain (C. elegans)	N2	CGC	RRID:WB- STRAIN:WBStrain00000001	All WT assays
Strain (C. elegans)	*cat-2(e1112)*	CB	CB1112	Behavioral testing
Strain (C. elegans)	*dat-1(ok157)*	CGC	RM2702	Behavioral testing

Strain (C. elegans)	dop-1(vs101)	CGC	LX636	Behavioral testing
Strain (C. elegans)	dop-2(vs105)	CGC	LX702	Behavioral testing
Strain (C. elegans)	dop-3(vs106)	CGC	LX703	Behavioral testing
Drug	cephalexin hydrate	Thermo Fisher	J6317206	Bacteria elongation
Dye	Oil-Red-O	Sigma-Aldrich	00625	Lipid storage analysis
Software algorithm	Igor Pro	Wavemetrics	Version 9.05	Behavioral testing, image analysis
Software	R	R Core Team (2024)	Version 4.4.2 (2024-10-31)	Statistical analysis
Software	WormLab	MBF Bioscience	Version 2024	Video analysis

[1]*Caenorhabditis* Genetics Center.

Bacteria culture

Two bacterial species were used: *E. coli* (OP50) used for worm culture, and *Comamonas* spp.

(DA1877) for assays. Bacterial cultures were grown in liquid lysogeny broth (LB) under sterile

conditions at 37°C with agitation. For all assays, DA1877 was grown for one day, then "Control"

cultures were solubilized with M9 buffer (3g KH_2PO_4, 6g Na_2HPO_4, 5g NaCl, 1mL 1 mol/L $MgSO_4$,

H_2O to 1L); "cephalexin" cultures were supplemented with a solution of M9 and cephalexin

hydrate 97% to a concentration of 450 µmol/L and incubated for 2 hours. Cultures then

underwent three rounds of concentration by centrifugation, followed by rinsing with 10 mL of

M9 buffer before being resuspended to their final concentration. Concentration was defined as

optical density at 600 nm (OD), as measured with a cell density meter (Laxco, DSM, Bothell, WA,

USA). All measurements were performed on samples diluted into the linear range of the

instrument (OD 0.1–1.0).

Worm maintenance & synchronization

All strains were maintained at 20°C on 5 cm plates containing nematode growth medium (NGM)

seeded with *Escherichia coli* OP50 bacteria as a food source (Brenner, 1974). For

synchronization, 20 young adult worms were transferred to fresh 5 cm NGM plates. Adults were

removed from the plates after laying eggs for 3 hours. The plates were maintained at 20°C until

worms reached the young adult stage.

Pharyngeal pumping assay

Pharyngeal pumping was measured electrophysiologically using the ScreenChip microfluidic

system (InVivo Biosystems, Eugene, OR, USA). Synchronized young adults were preincubated

with control or cephalexin-treated bacteria at OD 1 for 20 minutes before being transferred into

the reservoir of a microfluidic device fitted with electrodes. Individual worms were moved one

at a time from the reservoir into the recording channel and given one to two minutes to

acclimate. Voltage transients associated with pharyngeal pumping were recorded for five

minutes. All recordings were performed within 90 minutes of the worms being loaded into the

microfluidic reservoir. Mean pumping frequency was extracted using custom code written in

Igor Pro (WaveMetrics, Lake Oswego, OR, USA).

Food-preference assay

T-mazes (Levichev *et al.*, 2023) were constructed by cutting masks from sheets of 2 mm thick

ethylene-vinyl acetate foam (Darice Craft Foam, Strongsville, OH, USA) using a fabric-cutting

machine (Cricut, South Jordan, UT, USA). Masks were placed on blank 5 cm NGM plates. Maze

arms were baited with 4.5 μL of various bacterial suspensions as described in the text.

Synchronized young adults were washed by five rounds of centrifugation (30 sec, 300 g)

followed by aspiration of the supernatant, then deposited by liquid transfer at the starting point

of the maze. A transparent plastic disc was placed over the maze to eliminate air currents.

Twelve plates were placed on a flatbed scanner (EPSON, model J221B, Los Alamitos, CA, USA)

and imaged after an elapsed time of 60 minutes. Imaged worms were counted manually. The

food preference index was computed as $I = (n_N - n_L)/(n_N + n_L)$, where n_N and n_L are the

number of worms in the normal and elongated patches, respectively. In some experiments, a

paralytic agent (sodium azide, NaN_3, 3 μL at 20 mM) was added to each food patch to prevent

animals from leaving the patch of food after reaching it. Sodium azide diffuses through the agar

over time, and its action is not instantaneous. To account for these effects, all worms within 5

mm of the end of the maze arm, rather than only those on food, were counted when calculating

the preference index.

Patch-leaving assay

Custom 8-well plates (25 mm × 45 mm) were fabricated from acrylic plastic. Wells (diameter: 10

mm; depth: 6 mm) were filled with NGM agar. Bacterial suspension (2 µL, OD 1) was placed at

the center of each well, forming a patch 3-4 mm in diameter. Synchronized one-day-old adult

worms were cleaned by transferring them to an unseeded NGM plate for several minutes. One

worm was transferred to each well using a pick and placed at the center of the patch. The well

plate was immediately placed on a stage 8 cm above a diffuse white-light source and imaged (2

or 5 frames per second, 23.4 µm/pixel) for 15 minutes using a video camera (AmScope, 1080P,

Irvine, CA, USA) fitted with a telephoto lens (Nikon USA, AF Micro Nikkor 60 mm, Melville, NY,

USA). The distance between the lens and the stage was 50 cm. Worm tracks and locomotion

parameters were extracted using WormLab software (MBF Bioscience, Williston, VT, USA) and

analyzed using custom routines in Igor Pro (WaveMetrics, Lake Oswego, OR, USA). To define the

worm's position with respect to the patch, its margin was traced manually.

Bacterial composition analysis

Bacteria were grown under sterile conditions at 37°C with agitation in 500 mL of LB overnight,

pelleted (5,000 rpm, 3 minutes), washed with water, and re-pelleted. Pellets were stored at -

80°C prior to analysis. Bacterial stocks were weighed and then lysed with CelLytic B Plus kit

(Sigma-Aldrich, CB0050, Saint Louis, MO, USA) on the day of the assay.

<u>Protein</u>. Protein was assayed with the Pierce BCA Protein Assay kit (Thermo Fisher Scientific,

A55864, Waltham, MA, USA). Readings were taken with a spectrophotometer (NanoDrop,

Thermo Fisher Scientific, ND-2000, Waltham, MA, USA) using the bicinchoninic acid assay

setting.

Carbohydrate. Carbohydrate was measured with the Abbexa Total Carbohydrate Assay Kit

(Abbexa, abx298986, Cambridge, UK). Spectrophotometer readings were made at OD 540 nm.

Lipid. Lipid was quantified using the method of Aggeler et al. (Aggeler, Then and Ghosh, 1987).

We prepared 100 µL of cell lysate and added: 500 µL methanol, 250 µL dichloromethane, and

100 µL distilled H_2O. The mixture was vortexed before adding an additional 375 µL of distilled

water. The solution was vortexed again to form an emulsion and then centrifugated at 4800 g

for 10 minutes. The bottom layer of solution was transferred to a pre-weighed tube and then

weighed again after the lipid layer dried. The difference in weight was used to measure the

quantity of lipid per 100 µL of cell lysate.

Fat storage analysis

Fat storage was assessed using the procedures of Soukas et al. (Soukas et al., 2009) and Stuhr

et al. (Stuhr and Curran, 2020). A 0.5% isopropanol solution of Oil-Red-O (Sigma-Aldrich, Oil-

Red-O powder 00625, Saint Louis, MO, USA) was prepared several days before the assay.

Bacterial cultures (Control and cephalexin) were prepared as described above and adjusted to

OD 3 before being loaded onto fresh NGM plates. Seventeen young adult worms were placed

on these plates and allowed to lay eggs for 5 hours before being removed. The plates were

incubated at 20°C for two days. L4 larvae were harvested and washed in PBS + 0.01% Triton X-

100. Worms were fixed with a 60% isopropanol solution and incubated with the Oil-Red-O

solution for two hours on a rotating plate. Worms were washed twice with PBS + Triton 0.01%

solution, first for 30 minutes, then for one hour, on a rotating plate. Stained worms were

imaged on a microscope slide at 32×, and staining intensity was quantified using custom

routines in Igor Pro (WaveMetrics, Lake Oswega, OR, USA).

Statistical analysis

General. Statistical details for each test (figure panel, statistical test, effect or comparison

tested, definition of units of replication, number of replicates, statistic value, degrees of

freedom, p-value, and effect size) are compiled in Table 1. These details are cited in the text

using the notation "Table 1, row *n*," where *n* is the row number in the table. For all

comparisons, the centers of distributions were their arithmetic mean or median. For dispersion

measures, we used 95% confidence intervals. The following statistical tests were employed

according to the experimental design: We used *t*-tests to determine whether the mean differed

from zero. To assess differences in pairs of means, we used *t*-tests and, when the data were not

normally distributed (according to the Shapiro-Wilk test), the Mann-Whitney U test. For

comparisons of three or more means we used 1-way or 2-way ANOVA with *post hoc* tests.

State probabilities. State probabilities are an instance of *compositional data*, meaning data that

sum to 1 (or 100%), which requires special treatment. For analysis, we subjected these data to

the isometric log ratio transform (Aitchison, 1986; Egozcue *et al.*, 2003) using the

"compositions" package in R (Team, 2020; van den Boogaart, Tolosana-Delgado and Bren, 2024).

Following transformation, *t*-tests were used to infer statistical significance (Figure 3—figure

supplement 1).

Box plots. The box represents the first and third quartiles of the data, with a horizontal line

representing the median. The whiskers extend to the maximum and minimum values following

the removal of extreme outliers. These were defined as points with values below $Q1 - 3 \times IQR$

or above $Q1 + 3 \times IQR$, where $Q1$ is the first quartile, $Q3$ the third quartile and IQR is the

interquartile range $Q3 - Q1$. A diamond symbol represents the mean.

Number of replicates. In the case of *assay plates*, a replicate is unique cohort of worms on an

assay plate, such as a T-maze. In the case of *worms*, a replicate is a unique animal tested once.

Nutrient replicates (Fig. 1C–E) were biological or technical, as indicated in Table 1. The minimum

sample size for the T-maze assays was based on pilot experiments that demonstrated the ability

to detect moderate to small effects with 10–30 replicates per experimental condition.

Previously published EPG data (Katzen *et al.*, 2021) showed that mutants or treatments could be

distinguished with approximately 20 replicates; however, to ensure the detection of small effect

sizes across experimental conditions, larger sample sizes were used.

<u>Effect size</u>. For *t*-tests we used Cohen's *d*. For Mann-Whitney U tests, we used the rank-biserial

correlation with $r = 2\,U/(n_1 n_2) - 1$, where U is the Mann-Whitney statistic and n_1 and n_2 are

group sizes. For 1-way ANOVA, we used eta squared with $\eta^2 = SS_{\text{between}}/SS_{\text{total}}$ where

SS_{between} is the sum of squares between groups and SS_{total} is the total sum of squares. For 2-

way ANOVA we used $\eta^2 = SS_{\text{effect}}/SS_{\text{total}}$ where SS_{effect} is the sum of squares for a factor or

interaction. For MANOVA we used partial eta squared with $\eta_p^2 = SS_{\text{effect}}/(SS_{\text{effect}} + SS_{\text{error}})$

where SS_{error} is sum of squares for error.

ACKNOWLEDGMENTS

This research was supported by NIH grants DA053817 and GM152169. Some strains were

provided by the CGC, which is funded by NIH Office of Research Infrastructure Programs (P40

OD010440).

COMPETING INTERESTS

S.R.L. has a competing financial interest in InVivo Biosystems, Inc., which produces the

ScreenChip system utilized in the pharyngeal pumping assay. The other authors have no

competing interests to declare.

REFERENCES

Aggeler, R., Then, R. L. and Ghosh, R. (1987) 'Reduced expression of outer-membrane proteins
in β-lactam-resistant mutants of Enterobacter cloacae', *Journal of General Microbiology*,
133(12), pp. 3383–3392. doi: 10.1099/00221287-133-12-3383.

Aitchison, J. (1986) *The Statistical Analysis of Compositional Data*. London: Chapman and Hall.

Ben Arous, J., Laffont, S. and Chatenay, D. (2009) 'Molecular and sensory basis of a food related
two-state behavior in C. elegans', *PLoS ONE*, 4(10), pp. 1–8. doi: 10.1371/journal.pone.0007584.

Avery, L. and Shtonda, B. B. (2003) 'Food transport in the C. elegans pharynx', *Journal of*
Experimental Biology. doi: 10.1242/jeb.00433.

Bailey, M. R., Simpson, E. H. and Balsam, P. D. (2016a) 'Neural substrates underlying effort,
time, and risk-based decision making in motivated behavior', *Neurobiology of Learning and*

702 *Memory*. Elsevier Inc., 133, pp. 233–256. doi: 10.1016/j.nlm.2016.07.015.

703 Bailey, M. R., Simpson, E. H. and Balsam, P. D. (2016b) 'Neural substrates underlying effort,
704 time, and risk-based decision making in motivated behavior', *Neurobiology of Learning and*
705 *Memory*. Elsevier Inc., 133, pp. 233–256. doi: 10.1016/j.nlm.2016.07.015.

706 Bastien, B. L. *et al.* (2024) 'nlr-1/CNTNAP regulates dopamine circuit structure and foraging
707 behaviors in C. elegans', *Communications biology*. Springer US, 7(1), p. 1248. doi:
708 10.1038/s42003-024-06936-6.

709 Bendesky, A. *et al.* (2011) 'Catecholamine receptor polymorphisms affect decision-making in C .
710 elegans', *Nature*. Nature Publishing Group. doi: 10.1038/nature09821.

711 van den Boogaart, K., Tolosana-Delgado, R. and Bren, M. (2024) *Compositional Data Analysis*.

712 van den Bos, R. *et al.* (2006) 'Rats assess costs and benefits according to an internal standard',
713 *Behavioural Brain Research*, 171(2), pp. 350–354. doi: 10.1016/j.bbr.2006.03.035.

714 Brenner, S. (1974) 'The genetics of Caenorhabditis elegans.', *Genetics*. Oxford University Press,
715 77(1), pp. 71–94. doi: 10.1093/genetics/77.1.71.

716 Castrellon, J. J. *et al.* (2021) 'Dopaminergic modulation of reward discounting in healthy rats: a
717 systematic review and meta-analysis', *Psychopharmacology*, 238(3), pp. 711–723. doi:
718 10.1007/s00213-020-05723-5.

719 Chase, D. L. and Koelle, M. R. (2007) 'Biogenic amine neurotransmitters in C. elegans.',
720 *WormBook : the online review of C. elegans biology*, pp. 1–15. doi: 10.1895/wormbook.1.132.1.

721 Chase, D. L., Pepper, J. S. and Koelle, M. R. (2004) 'Mechanism of extrasynaptic dopamine
722 signaling in Caenorhabditis elegans', *Nature Neuroscience*, 7(10), pp. 1096–1103. doi:
723 10.1038/nn1316.

724 Egozcue, J. J. *et al.* (2003) 'Isometric Logratio Transformations for Compositional Data Analysis',
725 *Mathematical Geology*, 35(3), pp. 279–300. doi: 10.1023/A:1023818214614.

726 Feng, L. *et al.* (2025) 'C. elegans interprets dietary quality through context-dependent
727 serotonergic modulation', *bioRxiv*. doi: https://doi.org/10.1101/2025.01.05.631367.

728 Flavell, S. W. *et al.* (2013) 'Serotonin and the neuropeptide PDF initiate and extend opposing
729 behavioral states in C. Elegans', *Cell*. Elsevier Inc., 154(5), pp. 1023–1035. doi:
730 10.1016/j.cell.2013.08.001.

731 Floresco, S. B., Tse, M. T. L. and Ghods-Sharifi, S. (2008) 'Dopaminergic and glutamatergic
732 regulation of effort- and delay-based decision making', *Neuropsychopharmacology*, 33(8), pp.

1966–1979. doi: 10.1038/sj.npp.1301565.

Frézal, L. and Félix, M. A. (2015) 'The Natural History of Model Organisms: C. elegans outside
the Petri dish', *eLife*. eLife Sciences Publications Ltd, 2015(4). doi: 10.7554/ELIFE.05849.001.

Fujiwara, M., Sengupta, P. and McIntire, S. L. (2002) 'Regulation of body size and behavioral
state of C. elegans by sensory perception and the egl-4 cGMP-dependent protein kinase',
Neuron, 36(6), pp. 1091–1102. doi: 10.1016/S0896-6273(02)01093-0.

Gray, J. C. *et al.* (2019) 'Genomic basis of delayed reward discounting', *Behavioural Processes*.
Elsevier, 162(November 2018), pp. 157–161. doi: 10.1016/j.beproc.2019.03.006.

Green, L., Myerson, J. and Calvert, A. L. (2010) 'Pigeons' Discounting of Probabilistic and
Delayed Reinforcers', *Journal of the Experimental Analysis of Behavior*, 94(2), pp. 113–123. doi:
10.1901/jeab.2010.94-113.

Gruninger, T. R., Gualberto, D. G. and Garcia, L. R. (2008) 'Sensory perception of food and
insulin-like signals influence seizure susceptibility', *PLoS Genetics*, 4(7). doi:
10.1371/journal.pgen.1000117.

Hart, E. E. and Izquierdo, A. (2019) 'Quantity versus quality: Convergent findings in effort-based
choice tasks', *Behavioural Processes*. Elsevier B.V., pp. 178–185. doi:
10.1016/j.beproc.2019.05.009.

Hayden, B. Y. (2016) 'Time discounting and time preference in animals: A critical review',
Psychonomic Bulletin and Review. Springer New York LLC, 23(1), pp. 39–53. doi:
10.3758/s13423-015-0879-3.

Hills, T., Brockie, P. J. and Maricq, A. V. (2004) 'Dopamine and Glutamate Control Area-
Restricted Search Behavior in Caenorhabditis elegans', *Journal of Neuroscience*, 24(5), pp.
1217–1225. doi: 10.1523/JNEUROSCI.1569-03.2004.

Isles, A. R. *et al.* (2004) 'Common genetic effects on variation in impulsivity and activity in mice',
Journal of Neuroscience, 24(30), pp. 6733–6740. doi: 10.1523/JNEUROSCI.1650-04.2004.

Katzen, A. *et al.* (2021) 'The nematode worm C. elegans chooses between bacterial foods
exactly as if maximizing economic utility', *bioRxiv 2021.04.25.441352*.

Katzen, A. *et al.* (2023) 'The nematode worm C. Elegans chooses between bacterial foods as if
maximizing economic utility', *eLife*, 12, pp. 1–43. doi: 10.7554/eLife.69779.

Levichev, A. *et al.* (2023) 'The conserved endocannabinoid anandamide modulates olfactory
sensitivity to induce hedonic feeding in C. elegans', *Current Biology*. Elsevier Inc., 33(9), pp.
1625-1639.e4. doi: 10.1016/j.cub.2023.03.013.

Lockery, S. R. *et al.* (2012) 'A microfluidic device for whole-animal drug screening using
electrophysiological measures in the nematode C. elegans', *Lab on a Chip*. Royal Society of
Chemistry, 12(12), pp. 2211–2220. doi: 10.1039/c2lc00001f.

Mazur, J. E. and Coe, D. (1987) 'Test of transitivity in choices between fixed and variable
reinforcer delays', *Journal of the Experimental Analysis of Behavior*, 47, pp. 287–297.

Millet, J. R. M. *et al.* (2022) 'C. elegans PEZO-1 is a mechanosensitive ion channel involved in
food sensation', *Journal of General Physiology*, 154(1). doi: 10.1085/jgp.202112960.

Milward, K. *et al.* (2011) 'Neuronal and molecular substrates for optimal foraging in
Caenorhabditis elegans'. doi: 10.1073/pnas.1106134109/-
/DCSupplemental.www.pnas.org/cgi/doi/10.1073/pnas.1106134109.

Olofsson, B. (2014) 'The olfactory neuron AWC promotes avoidance of normally palatable food
following chronic dietary restriction', *Journal of Experimental Biology*, 217(10), pp. 1790–1798.
doi: 10.1242/jeb.099929.

St. Onge, J. R., Chiu, Y. C. and Floresco, S. B. (2010) 'Differential effects of dopaminergic
manipulations on risky choice', *Psychopharmacology*, 211(2), pp. 209–221. doi:
10.1007/s00213-010-1883-y.

Ostrander, S. *et al.* (2011) 'Orbitofrontal cortex and basolateral amygdala lesions result in
suboptimal and dissociable reward choices on cue-guided effort in rats', *Behavioral
Neuroscience*, 125(3), pp. 350–359. doi: 10.1037/a0023574.

Salamone, J. D. *et al.* (2018) 'Dopamine, effort-based choice, and behavioral economics: Basic
and translational research', *Frontiers in Behavioral Neuroscience*. Frontiers Media S.A. doi:
10.3389/fnbeh.2018.00052.

Salamone, J. D., Cousins, M. S. and Bucher, S. (1994) 'Anhedonia or anergia? Effects of
haloperidol and nucleus accumbens dopamine depletion on instrumental response selection in
a T-maze cost/benefit procedure', *Behavioural Brain Research*, 65(2), pp. 221–229. doi:
10.1016/0166-4328(94)90108-2.

Samuel, B. S. *et al.* (2016) 'Caenorhabditis elegans responses to bacteria from its natural
habitats', *Proceedings of the National Academy of Sciences of the United States of America*, pp.
E3941–E3949. doi: 10.1073/pnas.1607183113.

Samuelson, P. (1937) 'A note on measurement of utility', *The Review of Economic Studies*, 4, pp.
155–161.

Sanyal, S. *et al.* (2004) 'Dopamine modulates the plasticity of mechanosensory responses in
Caenorhabditis elegans', *EMBO Journal*, 23(2), pp. 473–482. doi: 10.1038/sj.emboj.7600057.

Sawin, E. R., Ranganathan, R. and Horvitz, H. R. (2000) 'C. elegans locomotory rate is modulated
by the environment through a dopaminergic pathway and by experience through a serotonergic
pathway', *Neuron*. doi: 10.1016/S0896-6273(00)81199-X.

Scheer, E. and Bargmann, C. I. (2023) 'Sensory neurons couple arousal and foraging decisions in
Caenorhabditis elegans', *eLife*, 12, pp. 1–26. doi: 10.7554/eLife.88657.

Shtonda, B. B. and Avery, L. (2006) 'Dietary choice behavior in Caenorhabditis elegans', *Journal*
of Experimental Biology. doi: 10.1242/jeb.01955.

Soukas, A. A. *et al.* (2009) 'Rictor/TORC2 regulates fat metabolism, feeding, growth, and life
span in Caenorhabditis elegans', *Genes and Development*, 23(4), pp. 496–511. doi:
10.1101/gad.1775409.

Stanek, J. C. and Richter, M. (2021) 'Energy investment and motivation: The additive impact of
task demand and reward value on exerted force in hand grip tasks', *Motivation and Emotion*.
Springer US, 45(2), pp. 131–145. doi: 10.1007/s11031-020-09862-2.

Stuhr, N. L. and Curran, S. P. (2020) 'Bacterial diets differentially alter lifespan and healthspan
trajectories in C. elegans', *Communications biology*. Springer US, 3(1), p. 653. doi:
10.1038/s42003-020-01379-1.

Suo, S., Culotti, J. G. and Van Tol, H. H. M. (2009) 'Dopamine counteracts octopamine signalling
in a neural circuit mediating food response in C. elegans', *EMBO Journal*, 28(16), pp. 2437–
2448. doi: 10.1038/emboj.2009.194.

Team, R. . (2020) *R A language and environment for statistical computing, R Foundation for*
Statistical.

Worthy, S. E. *et al.* (2018) 'Identification of attractive odorants released by preferred bacterial
food found in the natural habitats of C. elegans', *PLoS ONE*, 13(7), pp. 1–14. doi:
10.1371/journal.pone.0201158.

822