Chronic stress impairs autoinhibition in neurons of the locus coeruleus to increase asparagine endopeptidase activity

Hiroki Toyoda[1,2,†,*], Doyun Kim[3,†], Byeong Geon Koh[3], Tomomi Sano[4], Takashi Kanematsu[4], Seog Bae Oh[3,5,*], and Youngnam Kang[6,*]

[†]These authors contributed equally to this work.

[1]Department of Oral Physiology, Graduate School of Dentistry, Osaka University, Osaka 565-0871, Japan

[2]Department of Physiology, School of Dentistry, Aichi Gakuin University, Nagoya 464-8650, Japan

[3]Department of Neurobiology & Physiology, School of Dentistry and Dental Research Institute, Seoul National University, Seoul 03080, Republic of Korea

[4]Department of Cell Biology, Aging Science, and Pharmacology, Faculty of Dental Science, Kyushu University, Fukuoka 812-8582, Japan

[5]ADA Forsyth Institute, Cambridge MA, 02142, USA

[6]Department of Behavioral Sciences, Graduate School of Human Sciences, Osaka University, Osaka 565-0871, Japan

*Correspondence:

Youngnam Kang: kang.yn.923@osaka-u.ac.jp

Department of Behavioral Sciences, Graduate School of Human Sciences, Osaka University,

1-2 Yamadaoka, Suita, Osaka 565-0871, Japan

Seog Bae OH: odolbae@snu.ac.kr

Department of Neurobiology & Physiology, School of Dentistry and Dental Research Institute, Seoul National University, Seoul 03080, Republic of Korea

Hiroki Toyoda: toyoda@dpc.agu.ac.jp

Department of Physiology, School of Dentistry, Aichi Gakuin University, Nagoya 464-8650, Japan

Abstract

Impairments of locus coeruleus (LC) are implicated in anxiety/depression and Alzheimer's disease (AD). Increases in cytosolic noradrenaline (NA) concentration and MAO-A activity initiate the LC impairment through production of NA metabolite, 3,4-dihydroxyphenyl-glycolaldehyde (DOPEGAL), by MAO-A. However, how NA accumulates in soma/dendritic cytosol of LC neurons has never been addressed despite the fact that NA is virtually absent in cytosol while NA is produced exclusively in cytoplasmic vesicles from dopamine by dopamine-β-hydroxylase. Since re-uptake of autocrine-released NA following spike activity is the major source of NA accumulation, we investigated whether and how chronic stress can increase the spike activity accompanied by NA-autocrine. Overexcitation of LC neurons is normally prevented by the autoinhibition mediated by activation of α2A-adrenergic receptor (AR)-coupled inwardly rectifying potassium-current (GIRK-I) with autocrine-released NA. Patch-clamp study revealed that NA-induced GIRK-I in LC neurons was decreased in chronic restraint stress (RS) mice while a similar decrease was gradually caused by repeated excitation. Chronic RS caused internalization of α2A-ARs expressed in cell membrane in LC neurons and decreased protein/mRNA levels of α2A-ARs/GIRKs in membrane fraction. Subsequently, chronic RS increased the protein levels of MAO-A, DOPEGAL-induced asparagine endopeptidase (AEP) and tau N368. These results suggest that chronic RS-induced overexcitation due to the internalization of α2A-ARs/GIRK is accompanied by $[Ca^{2+}]_i$ increases, subsequently increasing Ca^{2+}-dependent MAO-A activity and NA-autocrine. Thus, it is likely that internalization of α2A-AR increased cytosolic NA, as reflected in AEP increases, by facilitating re-uptake of autocrine-released NA. The suppression of α2A-AR internalization may have a translational potential for AD treatment.

Keywords: locus coeruleus; α2A adrenergic receptor; GIRK; autoinhibition; MAO-A; AEP

Background

Locus coeruleus (LC) is implicated in anxiety and depression (*Itoi and Sugimoto, 2010*), and is often the first place where Alzheimer's disease (AD)-related pathology appears (*Mather and Harley, 2016*). Noradrenaline (NA) is known to inhibit amyloid-induced oxidative stress, and caspase activation in cortex/hippocampus (*Counts and Mufson, 2010*). Thus, the impairment of LC-noradrenergic system is potentially involved in the AD pathogenesis (*Ross et al., 2015; Chen et al., 2022*).

It was reported that cytosolic NA in the soma/dendrites of LC neurons is degraded by monoamine oxidase A (MAO-A) into 3,4-dihydroxyphenyl-glycolaldehyde (DOPEGAL) (*Burke et al., 1999*), which generates apoptotic free radicals (*Burke et al., 1998*) and activates asparagine endopeptidase (AEP) to produce Aβ and hypephosphorylated tau, consequently leading to AD (*Burke et al., 1999; Kang et al., 2020*). However, there appears normally no NA except dopamine in the cytosol of LC neurons because NA is produced exclusively in cytoplasmic vesicles by dopamine-β-hydroxylase (DBH) (*Potter and Axelrod, 1963*) from dopamine which is taken up into cytoplasmic vesicles by vesicular monoamine transporter type 2 (VMAT2).

LC neurons release NA following action potential not only from the axon terminals but also from their cell bodies as autocrine (*Huang et al., 2007*), and autocrine-released NA activates α2 adrenergic receptor (α2-AR) expressed on the soma/dendrites (*Aoki et al., 1994*), resulting in activation of α2-AR-coupled inwardly rectifying K^+ (GIRK) channel (*Arima et al., 1998*) to cause autoinhibition (spike-afterhyperpolarization) (*Williams et al., 1985*). Thus, overexcitation of LC neurons is normally prevented by the autoinhibition.

As autocrine-released NA is taken up by NA-transporter (NAT) (*Torres et al., 2003*), uptaken-NA may be the only source of free NA to be metabolized by MAO-A in the cytosol of LC neurons. However, the amount of free NA to be metabolized by MAO-A in the cytosol is thought to be very small due to a much lower affinity of NA to MAO-A compared to VMAT2 (*Costa, 2012*). Therefore, it is not clear how free NA to be metabolized by MAO-A is increased in the soma/dendites of LC neurons in AD pathogenesis. Because the passive NA-leakage occurs from cytoplasmic vesicles in a manner that achieves a dynamic equilibrium with active NA-storage into cytoplasmic vesicles by VMAT2 (*Eisenhofer et al., 2004*), increased NA-uptake by NAT in response to increased NA-autocrine following $[Ca^{2+}]_i$ increases or overexcitation may increase free NA by facilitating the passive NA-leakage from cytoplasmic vesicles. Furthermore, the affinity of MAO-A for NA may be modifiable as MAO-A activity

is Ca^{2+}-dependent (*Cao et al., 2007*). Then, a question arises whether or not impairment of

autoinhibition that causes overexcitation and $[Ca^{2+}]_i$ increases can consequently increase AEP

activity through the production of DOPEGAL.

Neuronal activity in LC neurons is transiently increased by stress (*Valentino and Foote,*

1988) through activation of corticotropin releasing factor (CRF) receptor (*Valentino et al.,*

1983; McCall et al., 2015) together with glutamate receptors (*Valentino et al., 2001*). However,

even if firing activity is increased by synaptic activation (*Barcomb et al., 2022*), autoinhibition

would occur to suppress overexcitation in LC neurons. Nevertheless, it is known that long-

term overexcitation due to chronic stress ultimately causes degeneration of LC neurons

(*Kitayama et al., 2004*). Then, this raises an important question whether and how stress impairs

autoinhibition to cause sustained overexcitation. The whole mechanisms for the impairment

of LC would be revealed by solving these questions. Here we examined whether the

autoinhibition mediated by α2A-AR-coupled GIRK-I is impaired by overexcitation in LC

neurons or chronic stress, to clarify the mechanisms for NA accumulation and/or MAO-A

activity increase.

Results

Spike frequency adaptation in LC neurons is mediated by α2-AR-coupled GIRK channel

LC neurons invariably display spike-frequency adaptation (**Figure 1A, B**). We have

hypothesized that the autoinhibition would occur following respective spike firing and

contribute to the generation of the spike-frequency adaptation and the afterhyperpolarization

seen at the offset of current pulses.

When the cumulative number of spikes was plotted against the elapsed time from the

onset of the current pulse every 50 ms, the cumulative spike number increased non-linearly in

a saturation manner (**Figure 1C**). This non-linear relationship was fitted with a saturation

curve defined by Monod equation (see Extended Methods and Materials) in order to quantify

the spike-frequency adaptation. Because there was no significant difference in *b* between the

two groups of spike trains, the profile of the normalized cumulative spike number vs the elapsed

time remains the same regardless of firing frequency, suggesting that spike-frequency

adaptation occurs in the same way.

The spike-frequency adaptation observed was almost completely abolished by a blocker

of α2-AR, atipamezole (**Figure 1D, E**), which subsequently largely decreased pulse-

afterhyperpolarization (pulse-AHP) (**Figure 1D, E**, blue and red stars). The cumulative spike

number increased non-linearly in a saturation manner in the control condition while it increased

almost in a linear manner after atipamezole application (**Figure 1F**). In the relationship

between *a* and *b* examined in LC neurons, atipamezole invariably and significantly shifted both

a and *b* in a larger direction, and there was a significant difference in the relationship between

a and *b* (**Figure 1G**).

Bath application of NA induced a GIRK-I in response to the ramp pulse (**Figure 1H**, red

trace) whereas addition of atipamezole to NA almost completely abolished the NA-induced

GIRK-I (**Figure 1H**, blue trace), showing that GIRK-I reversed at –97 mV. The amplitudes of

the inward component of NA-induced GIRK-I were invariably larger than those of the outward

component, consistent with the inwardly rectifying property of GIRK-I. Atipamezole

significantly decreased the amplitudes of the inward and outward components (**Figure 1I**).

Taken together, NA-autocrine is likely to be caused following respective spikes in the LC

neuron and NA is accumulated around the LC neuron to activate α2-AR-coupled GIRK

channels, to cause spike-frequency adaptation by enhancing AHP.

Ca^{2+}-dependent rundown of NA-induced GIRK-I in LC neurons

In PC12 cells, heterologously expressed α2A-ARs undergo internalization following agonist

application (*Taraviras et al., 2002*). Such internalization of α2A-ARs would result in an

abolishment of GIRK-I or desensitization given the functional coupling between α2A-AR and

GIRK as in LC noradrenergic neurons. However, in LC neurons, NA-induced GIRK-I showed

almost no rundown/desensitization after repeating the ramp pulse 20 times in the persistent

presence of NA (**Figure supplement 1A-D**). Then, we next examined whether GIRK-I shows

rundown under the condition of "increased excitability" or increased [Ca^{2+}]$_i$, because

acute/chronic stress increases the excitability of LC neurons (*Borodovitsyna et al., 2018;*

Campos-Lira et al., 2018) and various channels/receptors are known to be internalized in a

Ca^{2+}-dependent manner (*Beattie et al., 2000; Aziz et al., 2012; Zaccor et al., 2020*).

To examine the Ca^{2+}-dependency of the rundown of GIRK-I, the respective ramp

pulses to evoke GIRK-I were preceded by various pulse trains which can induce Ca^{2+} currents

differentially (compare **Figure supplement 1F** and **Figure 2B**). After examining various pulse

protocols (**Figures supplement 1 and 2**), the best protocol to induce a prominent rundown of

GIRK-I was as follows (shown in **Figure 2B**). First, bath application of NA induced GIRK-I

(**Figure 2A,** red trace; 2) that showed the inwardly rectifying profile as revealed by subtraction

of the control response from that obtained after NA application (**Figure 2D**, red trace; 2−1).

Subsequently, the combined command pulse (a set of 5 trains of 20 pulses followed by a ramp

pulse; **Figure 2B**, upper trace) was applied every minute. The positive pulse trains evoked

large inward Ca^{2+} currents in the presence of extracellular 30 mM TEA (**Figure 2B**, lower

traces) whereas similar trains of the same positive pulse hardly induced Ca^{2+} currents when

applied in the absence of TEA (**Figure supplement 1F**, lower trace). With repetition of the

combined command pulse 20 times, the NA-induced GIRK-I (**Figure 2C**, red trace; 2) was

decreased to almost the same current as the control response (**Figure 2C**, blue trace; 3). Indeed,

no GIRK-I remained after 20 times repetition of the combined command pulse as revealed by

the subtraction between trace 1 and trace 3 (**Figure 2D**, blue trace; 3−1). Both the amplitudes

of the inward and outward components of the NA-induced GIRK-I promptly decreased to zero

along with repetition of the combined command pulse (**Figure 2E**, blue and red circles; **Figure**

2F). Thus, it is likely that NA-induced GIRK-I displays a marked rundown in a Ca^{2+}-dependent

manner. Furthermore, we confirmed that these rundowns were completely dependent on the

number and application timing of the command pulse in the presence or absence of TEA

(compare **Figure 2** with **Figures supplement 1 and 2**). Depending on the absence or presence

TEA, the single application of a set of 10 trains of 20 pulses differentially (slightly or

moderately, respectively) induced the rundown of NA-induced GIRK-I (compare **Figure**

supplement 1E-J and **Figure supplement 2A-E**). By contrast, the repetitive application of a

single train of 20 positive pulses in the presence of TEA markedly facilitated the rundown of

NA-induced GIRK-I (**Figure supplement 2F-K**). The relationship between the number of

spikes (i.e., positive pulse) and $[Ca^{2+}]_i$ was obtained by measuring Ca^{2+} transients using fura-2

(**Figure supplement 3**).

To investigate the effects of the rundown of GIRK-I on the spike-frequency adaptation,

spike trains were examined before and after the voltage-clamp experiment (**Figure 2A-F**).

After the rundown of GIRK-I was observed in the voltage-clamp experiment, the spike-

frequency adaptation was absent in the current clamp experiment, and the pulse-ADP was

largely reduced (**Figure supplement 4A-D**). The cumulative spike number that increased non-

linearly in a saturation manner before rundown of GIRK-I increased almost in a linear manner

after rundown of GIRK-I (**Figure supplement 4E, F**). Thus, the rundown of GIRK-I

invariably resulted in an abolishment of spike-frequency adaptation, similar to the effects of

atipamezole on spike-frequency adaptation (**Figure 1**). Therefore, it is likely that Ca^{2+}

increases can cause the rundown of NA-induced GIRK-I and almost abolished the spike-

frequency adaptation, leading to an increase in the excitability of LC neurons. These

observations suggest that $[Ca^{2+}]_i$ increases cause rundown of NA-induced GIRK-I to suppress

autoinhibition, leading to excitability increases that cause further increases in $[Ca^{2+}]_i$ to further

cause a rundown of the GIRK-I.

It was also examined whether the rundown of NA-induced GIRK-I can be caused solely

by $[Ca^{2+}]_i$ increases without activation of α2A-AR. We found that NA-induced GIRK-I

subjected to the combined command pulse 20 times in the presence of atipamezole was

significantly restored after washout of atipamezole in the presence of NA (**Figure supplement**

5A-D). This observation indicates that GIRK-I were not downregulated solely by $[Ca^{2+}]_i$

increases without activation of α2-AR, clearly revealing that both the activation of α2-AR and

$[Ca^{2+}]_i$ increases are necessary to cause the downregulation of GIRK-I. In agreement with this

result, the spike-frequency adaptation reflecting the autoinhibition was also preserved in spite

of $[Ca^{2+}]_i$ increases as long as activation of α2-AR is prevented (**Figures supplement 5E-J**).

We further examined whether the rundown of NA-induced GIRK-I can be prevented by the

presence of a GIRK channel blocker, tertiapin-Q. GIRK channel blockade did not prevent the

rundown of NA-induced GIRK currents following $[Ca^{2+}]_i$ increases (**Figure supplement 6A-**

H). In agreement with this result, the spike-frequency adaptation was not preserved by with

the blockade of GIRK channels during $[Ca^{2+}]_i$ increases (**Figure supplement 6I-N**). These

findings showed that α2-AR blockade but not GIRK blockade prevented the rundown of NA-

induced GIRK-I following $[Ca^{2+}]_i$ increases. Taken together, it is suggested that the rundown

of NA-induced GIRK-I may be brought about by the internalization of α2-AR in a Ca^{2+}-

dependent manner, regardless of opening or closing of GIRK channels.

β-Arrestin is involved in the rundown of NA-induced GIRK-I

We further examined whether the internalization of α2-AR is involved in the rundown of NA-

induced GIRK-I or not. As β-arrestin/AP2 complex is known to be involved in the agonist-

dependent internalization of GPCR (*DeGraff et al., 1999; Cottingham et al., 2011*), we

examined the effects of barbadin, which prevents AP2 from binding with β-arrestin by forming

barbadin/AP2 complex, on the rundown of NA-induced GIRK-I. With the patch pipettes filled

with barbadin-including internal solution, it was examined whether NA-induced GIRK-I

displays rundown in response to the same combined command pulse (**Figure 2H**, upper trace)

as that in the control experiment (**Figure 2B**, upper trace). The NA-induced GIRK-I (**Figure**

2G-I, red traces) was hardly decreased (**Figure 2H, I**, blue traces) after repeating the combined

pulse 20 times, in contrast to the control rundown (**Figure 2C, D**). Barbadin largely suppressed

the rundown of the both components of GIRK-I (**Figure 2J, K**). These results suggest that the
rundown of NA-induced GIRK-I is caused by the internalization of α2-AR-coupled GIRK
channels.
231
Restraint stress alters α2A-AR expression in LC neurons
We next investigated whether the expression of α2A-AR is altered in LC neurons by applying
restraint stress (RS) to mice. After mice were subjected to daily 12-h RS for 3 days followed
by non-stress condition for 2 days, the immunoreactivity for α2A-ARs (see **Figure supplement**
7) seen along the membrane of TH-positive LC neurons was apparently lower compared to the
control (**Figure 3A, B**). Consistent with this observation, Western blot analysis revealed
significantly lower protein levels of α2A-ARs in LC neurons obtained after 3-day RS compared
to the control (**Figure 3C**). Especially when α2A-ARs expression was examined in the
membrane fraction identified using Na^+/K^+-ATPase, the immunoreactivity for α2A-ARs
expression was apparently lower in LC neurons of RS mice than in those of the control mice
(**Figure 3D, E**; *, compare the rightmost lower panels). Consistent with this observation,
Western blot analysis revealed a significantly lower protein level of α2A-ARs in the membrane
fraction of LC neurons of RS mice compared to the control whereas there was no significant
difference in the cytosol between the control and RS mice (**Figure 3F, G**). Thus, the
internalization of α2A-ARs is not necessarily accompanied by increases in cytosolic levels of
α2A-ARs because the internalized receptors are either recycled or degraded (*Koenig and Ikeda,*
1996; Takei et al., 1996).
249
RS reduces NA-induced GIRK-I
As 3-day RS caused the internalization or downregulation of α2A-ARs, we next examined if
the daily 12-h RS for up to 5 days can differentially cause reductions of NA-induced GIRK-I
in LC neurons. NA-induced GIRK-I decreased with an increase in the RS period (**Figure 4A-**
E). The amplitudes of inward and outward components of GIRK-I decreased with the increase
in the period of RS (**Figure 4F, G**), in a way that is described by a saturation function (red
interrupted lines). The amplitudes of inward and outward components of GIRK-I in 1-day
mice were significantly smaller than those of the control mice, and those in 3-day RS mice
were significantly smaller than those in 1-day RS mice while no significant difference in those
between 3-day and 5-day RS mice (**Figure 4F, G**). Thus, the downregulation of GIRK-I was
increased as RS period is increased, but in a saturation manner.

Quantitative PCR analysis revealed that LC neurons expressed much higher mRNA

level of α2A-AR than that of α2C-AR (**Figure 4H**) and also expressed higher mRNA level of

GIRK1 compared to GIRK2 while GIRK3 was hardly expressed (**Figure 4I**). Consistent with

the significant decreases in NA-induced GIRK-I in the mice subjected to 3-day RS, mRNA

levels of α2A-AR, GIRK1 and GIRK2 in mice subjected to 3-day RS were significantly smaller

than those of the control mice (**Figure 4J-L**). Thus, RS downregulated the

expression/production of α2A-AR-coupled GIRK1 prominently in LC neurons.

Histological and biochemical changes in LC neurons after 5-day RS

Next, immunohistochemical experiments demonstrated that 5-day RS appeared to slightly

increase the expressions of TH and MAO-A in LC neurons (**Figure supplement 8A**).

Furthermore, AEP expression also appeared to increase in cytosol after 5-day RS compared to

the control (**Figure supplement 8B**). Because TH catalyzes the rate limiting step in the

synthesis of catecholamines, if the RS enhances TH activity, free NA concentration in LC

neurons would be directly and greatly increased. Therefore, we first carefully examined

whether TH is really upregulated in RS mice or not by performing Western blot analysis. No

significant increase in the protein level of TH was observed in 5-day RS mice compared to the

control (**Figure 5A**, upper panel; **Figure 5B**). This result clearly indicates that TH activity was

not significantly upregulated in 5-day RS mice, in spite of the TH immunohistochemical

experiment. We further examined whether protein levels of MAO-A and AEP are increased

or not by 5-day RS. Western blot analysis revealed significant increases in the protein levels

of MAO-A (**Figure 5A**, lower panel; **Figure 5C**) and of pro-AEP and active AEP (**Figure 5D-**

F) in LC neurons after 5-day RS compared to the control. Western blot analysis also revealed

a significant increase in the expression of tau and tau N368 fragment after 5-day RS (**Figure**

5G-I). Thus, an increased activity of active AEP caused a production of tau N368 fragments

as observed at 37 and 35 kDa (**Figure 5G**), which is consistent with a previous report (*Zhang*

et al., 2014). Relative protein levels of the tau N368 fragments (**Figure 5I**) were the mean

values obtained from the two band intensities, which may reflect the two N-terminal fragments

(upper and lower bands) generated by AEP cleavage at Asn-368 of the 4 and 3 repeat-tau

isoforms, respectively (*Goedert et al., 1989*). Finally, Y-maze test revealed that 5-day RS

decreased spatial memory (**Figure supplement 9**).

Discussion

We have found that LC neurons display spike-frequency adaptation which is caused by the autoinhibition mediated by α2A-AR-coupled GIRK-I following respective spikes (**Figure 1**) and that α2A-AR-coupled GIRK-I showed Ca^{2+}-dependent rundown due to the internalization of α2A-AR (**Figure 2**, **Figure supplement 5A-D** and **Figure supplement 6E-H**), leading to the abolishment of spike-frequency adaptation and the excitability increase (**Figure supplement 4**). Immunohistochemical examination and Western blot analysis revealed that α2A-AR expression in cell membrane was decreased following RS (**Figures 3 and 4J**). Consistently, NA-induced GIRK-I was decreased in LC neurons with an increase in the period of RS (**Figure 4A-G**). Following chronic RS, protein levels of MAO-A, pro-/active-AEP, and tau and tau N368 fragment were significantly increased (**Figure supplement 8A, B and Figure 5A, C-I**).

Based on these results, we propose a novel mechanism for how chronic RS impairs autoinhibition and increases cytosolic free NA to be metabolized by enhanced Ca^{2+}-dependent MAO-A activity, to consequently increase AEP activity and produce tau N368 protein (**Figure 6**). In non-stress mice, autocrine released NA is slowly taken up by NAT after binding to α2-ARs to cause autoinhibition (**Figure 6A**), while in chronic RS mice, autocrine-released NA may be directly and rapidly taken up by NAT due to the absence (internalization) of α2-ARs (**Figure 6B**). Such increased rate of re-uptake of NA by NAT would increase free NA concentration by facilitating the leakage of NA from cytoplasmic vesicles in the cell body, which would be metabolized by upregulated MAO-A activity to DOPEGAL, as revealed by the increased activity of AEP in chronic RS mice (**Figure 6B**). It may be further necessary to perform enzymatic assay to better demonstrate the AEP and MAO-A activities.

Spike frequency adaptation

LC neurons display spike-frequency adaptation. The mechanism for the spike-frequency adaptation may be different among neurons while M currents are usually responsible for the adaptation in hippocampal pyramidal neurons (*Aiken et al., 1995*). In LC neurons, the spike-frequency adaptation was mediated by α2A-AR-coupled GIRK-I as the spike-frequency adaptation was abolished by atipamezole (**Figure 1D-G**) and by causing the rundown of GIRK-I (**Figure supplement 4**). These results suggest that autocrine NA release following respective spikes activated α2A-ARs expressed in the soma/dendrites of LC neurons, and subsequently activated α2A-AR-coupled GIRK-I to cause autoinhibition as spike-frequency adaptation.

Thus, spike-frequency adaptation clearly reflects NA-autocrine.

Rundown of α2A-AR-coupled GIRK-I

α2A-AR-coupled GIRK-I displayed Ca^{2+}-dependent rundown in the presence of agonist. The

degree and time course of rundown of α2A-AR-coupled GIRK-I were dependent on the number

and timing of positive pulses applied under voltage-clamp condition to cause Ca^{2+} currents

(**Figure 2A-F** and **Figures supplement 1E-J, 2, 3**). Given the Ca^{2+}-dependency of the

rundown of autoinhibition, the excitability of LC neurons would increase as LC neurons were

excited more such as under RS condition, resulting in a vicious cycle between excitability

increase accompanied by $[Ca^{2+}]_i$ increase and rundown of autoinhibition/Ca^{2+}-dependent

internalization of α2A-ARs.

Internalization of α2A-ARs

Activation of GPCR by agonist binding usually causes the formation of GPCR/β-arrestin

complexes. Subsequently, the GPCR/β-arrestin complex is assembled through the interaction

between β-arrestin and AP2 in clathrin-coated pits (CCPs), which is formed by the association

of AP2 with phosphatidylinositol 4,5-bisphosphate (PIP2) at the plasma membrane, leading to

endocytosis of GPCRs (*Pierce and Lefkowitz, 2001*). PIP2 is locally produced from the

membrane phosphatidylinositol (PI) by PI4/PIP5 kinases in a Ca^{2+}-dependent manner

(*Sarmento et al., 2014*). Barbadin prevented the stable formation of sufficient GPCR/β-

arrestin/AP2 complexes in CCPs as barbadin prevents AP2 from binding with β-arrestin by

forming barbadin/AP2 complex, thereby hampering GPCR internalization (*Beautrait et al.,*

2017). In the present study, intracellular barbadin effectively suppressed the rundown of

GIRK-I (**Figure 2G-K**), indicating that the rundown of GIRK-I was caused by the

internalization of α2A-ARs in LC neurons.

Chronic RS increases the activities of MAO-A and pro-/active-AEP in the cell body

Chronic cold stress reduced α2-AR mediated inhibition of LC neurons (*Jedema et al., 2008*),

and chronic social stress downregulated α2-ARs in the LC (*Flugge, 1996*). Our present results

were consistent with those previous findings: Following RS, α2A-ARs expression was

decreased due to the internalization of α2A-ARs (**Figure 3**), together with the decreases in NA-

induced GIRK-I (**Figure 4A-G**) which mediates the autoinhibition. As long as RS persists, the

impairment of autoinhibition leads to the persistent overexcitation in LC neurons. This would

further increase NA-autocrine, because respective spikes were accompanied by NA-autocrine

as reflected in spike-frequency adaptation or autoinhibition.

As extracellular NA is mostly taken up into cytosol by NAT (*Torres et al., 2003*), it is

likely that under chronic RS the increased autocrine-released NA is directly and more rapidly

taken up by NAT due to the absence of α2A-ARs (**Figure 6B**). Such an increase in the rate of

NA uptake by NAT is similar to the effects of uptake inhibitor of transmitter ligand on the

activity of receptors. After re-uptake of NA by NAT, a large portion of NA in the cytosol is

restored in cytoplasmic vesicles by VMAT2 while only a small portion of NA can escape from

VMAT2 to be metabolized by MAO-A (*Eisenhofer et al., 1992*), because VMAT2 has a much

higher affinity for NA than MAO-A (*Bloom, 2006; Costa, 2012*). However, there is an

additional component of NA to be metabolized by MAO-A, as NA can leak from cytoplasmic

vesicles into cytosol (*Eisenhofer et al., 2004*). When re-uptake of NA by NAT is facilitated

due to the internalization of α2-ARs in response to NA-autocrine, active NA storage into

cytoplasmic vesicles by VMAT2 would increase. Then, the rate of NA leakage from

cytoplasmic vesicles would also increase because passive NA leakage from cytoplasmic

vesicles and active NA storage into cytoplasmic vesicles by VMAT2 are in a dynamic

equilibrium in cytoplasmic vesicles (*Eisenhofer et al., 2004*). Subsequently, such increases in

the rate of NA leakage would result in increases in a MAO-A metabolite, DOPEGAL

(*Goldstein, 2021*) (**Figure 6B**). Indeed, it has previously been demonstrated that a partial

inhibition of NAT in human subjects reduced the original level of a derivative of DOPEGAL

(3,4-dihydroxyphenylglycol; DHPG) in CSF by more than 50% (*Kielbasa et al., 2015*). Thus,

free NA to be metabolized by MAO-A would be increased when autocrine-released NA is

taken up by NAT. On the other hand, it is also known that the decreased VMAT2 levels,

generated in VMAT2 heterozygote mice, increased the vulnerability of nigral dopamine

neurons to MPTP (*Takahashi et al., 1997*). Thus, it is conceivable that NA leakage for free

NA accumulation is influenced by both NAT or VMAT2.

Therefore, under the chronic RS condition, more NA leaked from cytoplasmic vesicles is

likely to be metabolized by MAO-A, which activity is enhanced Ca^{2+}-dependently and by

chronic RS (*Cao et al., 2007*) (**Figures 5A, C and 6B**), to DOPEGAL compared to the control

condition, as revealed by the increased protein levels of pro-AEP and active-AEP (**Figure**

supplement 8B and Figure 5D-F). Thus, re-uptaken NA by NAT may largely be degraded to

DOPEGAL by the increased activity of MAO-A. Subsequently, DOPEGAL converts pro-AEP

into active-AEP to produce tau N368 protein (**Figure 5G-I**).

It has never been addressed how DOPEGAL production was caused in the cell body
of LC neurons while the DOPEGAL production in the cell bodies of LC neurons is well
established in AD patients (*Burke et al., 1999; Kang et al., 2020*) and in transgenic AD model
mice (*Kang et al., 2020*). We demonstrated for the first time that 5-day RS can impair the
autoinhibition by causing the internalization of α2A-AR and consequently induces active-AEP
to produce tau N368 fragment. Indeed, 5-day RS induced spatial memory impairment-like
responses in mice when assessed using Y-maze test (**Figure supplement 9**) while anxiety-like
behavior was also induced in mice after 3-day RS when assessed using an elevated plus maze
test (figure not shown), both suggesting an impairment of LC neurons. It is likely that, in
addition to NAT inhibitor, the suppression of the internalization of α2A-AR has a strong
translational potential for developing drugs that prevent the degeneration of LC.

Limitations of this study

One may argue against the usage of 3-week-old juvenile mice to study the neurodegenerative
disorder which is normally found in aged ones. However, a stringent autoinhibition together
with potent NAT activity were previously demonstrated in adult rats using intracellular
recordings (*Williams et al., 1985*). Thus, regardless of the age of mice, LC neurons showed
the autoinhibition that is mediated by the activity of α2A-AR-coupled GIRK channels. We
also confirmed that spike-frequency adaptation reflecting autoinhibition can be induced in adult
mice as prominently as in juvenile mice regardless of sexes (**Figure supplement 10**), although
we did not investigate the sex differences in RS effects (*Leistner and Menke, 2020*). The
downregulation of autoinhibition by the internalization of α2A-AR is agonist- and Ca^{2+}-
dependent regardless of the age of mice. However, it is not known if there is any difference in
the susceptibility of α2A-AR internalization to RS between juvenile and aged mice. Thus, a
further study may be necessary to examine if autoinhibition in LC neurons is more easily
downregulated in aged mice compared to juvenile ones.

It has also been reported that re-uptake of NA by NAT in LC neurons may decrease
with age (*Sanders et al., 2005; Mitchell et al., 2017*). However, NAT expression in LC in 3-
week-old juvenile mice was only slightly larger (by 4%) than in adult mice although the
difference was significant (*Mitchell et al., 2017*). Potent NAT activity stringently affecting
autoinhibition was reported in adult rats (*Williams et al., 1985*). Furthermore, partial inhibition
of NAT in human subjects largely reduced the original level of a derivative of DOPEGAL in
CSF as described above. Thus, based on the present study, it is likely that increased re-uptake

of NA by NAT following excessive NA-autocrine in the absence of α2A-AR, induced in
response to chronic RS, resulted in the increase in active-AEP and subsequent production of
tau N368. However, a further experiment to directly assess NAT activity would be necessary
to firmly justify our hypothesis.

To further validate the mechanisms which we have proposed in the present study
(Figure 6), direct quantification of the relationship between α2A-AR internalization and
increased cytosolic NA levels may be desirable. However, at present, the quantification of
such connection may not be possible because there appears to be no fluorescence probe to label
cytosolic NA. None of NA fluorescence probes, such as NS521, BPS3 or GRAB-NE, is
applicable to detect cytosolic free NA because NS521 and BPS3 anchor on plasma membrane
and GRAB-NE is dependent on GPCR activation (*Hettie and Glass, 2014; Feng et al., 2019;*
Mao et al., 2023).

Conclusion

Our study unveiled the cellular mechanism for the degeneration of LC neurons under chronic
RS as follows: Chronic RS impaired the autoinhibition in LC neurons by inducing Ca^{2+}- and
agonist-dependent internalization of soma/dendritic α2A-AR coupled with GIRK channels.
This subsequently caused overexcitation/$[Ca^{2+}]_i$ increases and increased Ca^{2+}-dependent
MAO-A activity and active AEP activity, leading to a production of tau N368 fragment
together with an impairment of spatial memory. These results clearly indicate that chronic RS
increases cytosolic free NA to be metabolized by MAO-A, presumably by facilitating the re-
uptake of autocrine-released NA due to the absence/internalization of α2A-AR. Thus, in
addition to NAT inhibitor, the suppression of α2A-AR internalization may have a strong
translational potential for developing drugs that prevent the degeneration of LC. However, the
present mechanism must be proven to be valid in adult or old mice, to validate its involvement
in the pathogenesis of AD.

Methods

Restraint stress

Three-weeks-old male C57BL/6 mice (JAPAN HAMAMATSU SLC, Hamamatsu, Japan)
were used and housed at $24 \pm 2°C$. Food and water were available ad libitum. The mice in the
stressed groups were restrained in a modified 50 ml clear polystyrene conical centrifuge with

multiple air holes for ventilation. The mice were subjected to 12-h restraint stress (RS; 20:00

p.m. to 8:00 a.m. of the next day) for 1, 2, 3 and 5 days. In the previous study, the duration of

RS widely ranged from 0.5-h (*Gray et al., 2010*) to 24-h (*Chu et al., 2016*) while the RS with

the duration less than 12-h has been applied daily for a period (6-h for 2 weeks (*Sakaguchi and*

Nakamura, 1990); 8-h for up to 16 weeks (*Sugama et al., 2016*); 12-h for 2 days (*Zhang et al.,*

2008)). As the prolonged stress is known to be a potential risk factor for AD (*Bisht et al.,*

2018), the protocol of daily 12-h RS is relevant for the study of functional and cytological

disorders of LC neurons. Once the restraint ended, the mice were returned to their home cages

with access to food and water ad libitum. The control group remained in their home cages

without the restraint procedure.

Slice preparation and electrophysiological recordings

Slices were prepared from 3-weeks-old male C57BL/6 mice to investigate the effects of RS for

varying periods on GIRK-I, and from C57BL/6 mice of either sex at 16–23 postnatal days (PD)

to investigate the effects of $[Ca^{2+}]_i$ increases on GIRK-I, similar to our previous study (*Toyoda*

et al., 2022). Using an Axopatch 200B (MDS Analytical Technologies, Sunnyvale, CA),

whole-cell recordings were made from LC neurons that were viewed under Nomarski optics

(BX51WI; Olympus, Tokyo, Japan). LC neurons were identified immunohistochemically and

electrophysiologically as we previously reported (*Toyoda et al., 2022*). The internal solution

of the patch pipettes had the following ionic composition (in mM): 123 K-gluconate, 18 KCl,

10 NaCl, 2 $MgCl_2$, 2 ATP-Na_2, 0.3 GTP-Na_3, 10 HEPES, and 0.2 EGTA; pH 7.3. The

membrane potential values given in the text were corrected for the junction potential (10 mV)

between the internal solution for the whole-cell recording (negative) and the standard

extracellular solution. The pipette resistances were 4–6 MΩ. The series resistance was < 10

MΩ. All recordings were made at room temperature (RT). Records of currents and voltages

were low-pass filtered at 5 kHz (3-pole Bessel filter), digitized at a sampling rate of 10 kHz

(Digidata 1322A, MDS Analytical Technologies) and stored on a computer hard disk. Spike-

frequency adaptation was quantitatively estimated by fitting with a saturation curve defined by

Monod equation (y = (**a** * x) / (**b** + x)) with a saturation level (**a**) and a half saturation constant

(**b**). The amplitudes of inward and outward components of NA-induced GIRK-I were

measured at –130 mV and –60 mV.

Procedure for immunohistochemical examination

C57BL6/J male mice (4-5 weeks) were perfused transcardially with 5 ml of PBS, followed by 10 ml of a freshly prepared 4% paraformaldehyde in PBS. The whole brain including LC (–9 to –10 mm from bregma) was dissected out and immersed in 4% paraformaldehyde solution for 4 h at 4°C fallowed by cryoprotection with 30% (w/w) sucrose in PBS.

Brainstem including LC regions was cut into 8-20 μm-thickness coronal sections on a freezing microtome (Leica Microsystems, Wetzlar, Germany). To examine the distribution of α2A-AR in LC neurons, mouse anti-TH (1:500; sc-25269, Santa Cruz Biotechnology, Santa Cruz, CA, USA), rabbit anti-sodium potassium ATPase (1:1000; ab76020, Abcam, Cambridge, MA, USA) and goat anti-α2A-AR (1:200; ab45871, Abcam) or rabbit anti-α2A-AR (1:500; RA14110, Neuromics, Northfield, MN, USA) were used as primary antibodies, while cy5 donkey anti-mouse IgG (1:500; cy-2500, Vector Laboratories, Burlingame, CA, USA), FITC donkey anti-rabbit IgG (1:500; 711-095-152, Jackson ImmunoResearch, West Grove, PA, USA) and Cy3 donkey anti-goat IgG (1:500; 706-165-147, Jackson ImmunoResearch) were used as secondary antibodies. We used the affinity purified IgG antibody for α2A-AR immunohistochemistry, in which antibody specificity was ensured as examined previously (*Stone et al., 1998; Tan et al., 2009*). To examine the expression in LC neurons, mouse anti-TH (1:500) and rabbit anti-MAO-A (1:2000; ab126751, Abcam) were used as primary antibodies, while FITC donkey anti-mouse IgG (1:500; 715-545-150; Jackson ImmunoResearch) for TH and Cy3 donkey anti-rabbit IgG (1:2000; 711-165-152; Jackson ImmunoResearch) were used as secondary antibodies. To examine the distribution of legumain in LC neurons, mouse anti-legumain (1:200; sc-133234, Santa Cruz) and rabbit anti-DBH (1:500; 22806, ImmunoStar, Hudson, WI, USA) were used as primary antibodies, while FITC donkey anti-mouse IgG (1:500; 715-545-150, Jackson ImmunoResearch) and Cy3 donkey anti-rabbit IgG (1:500; 711-165-152, Jackson ImmunoResearch) were used as secondary antibodies. The slices were coverslipped with mounting media (H1000, Vector Laboratories), and sections were visualized with a confocal microscope (LSM980, Carl Zeiss, Oberkochen, Germany) at 10x and 63x magnification.

Western blot study

Brain slices from the control and 5-day RS-treated mice were sectioned with vibratome (VT1000P, (Leica Microsystems) to obtain 500 μm-thickness coronal sections containing LC

region (bregma –5.34 mm to –5.80 mm). We performed western blot study only in 5-day RS
mice without examining in 3-day RS mice, under the assumption that for the degeneration-
related biochemical changes to reach sufficiently detectable levels, a high $[Ca^{2+}]_i$ condition due
to the rundown of GIRK-I have to be sustained for some period of time after the rundown of
GIRK-I reached a plateau level in 3-day RS mice (see Figure 4F, G). LC dissection was done
via micro punch using 1.5 mm biopsy punch (BP-15F, Kai medical, Gifu, Japan) bilaterally
under light microscope. Dissected LC were quickly frozen in liquid nitrogen and stored at
−80°C until processed. Because LC region was too small to collect sufficient tissue from one
mouse for the analysis, the punched LC tissues were collected from two to three mice to obtain
a protein lysate sample for each western blotting analysis. LC tissues were homogenized by
bead homogenizer in T-PER™ Tissue Protein Extraction Reagent (78510, Thermo Fisher)
containing protease and phosphatase inhibitor cocktails (MilliporeSigma, St. Louis, MO, USA).
The membrane and cytosol proteins from LC lysate were isolated using Mem-PER Plus Kit
(78840, Thermo Fisher) in accordance with the manufacture's protocol.   The protein
concentration of the lysate was determined using a Lowry protein assay (Bio-Rad, Hercules,
CA, USA). Protein samples (20 µg) were heated to 95°C for 5 min, separated by 12% sodium
dodecyl sulfate-polyacrylamide gel electrophoresis, and transferred to a polyvinylidene
fluoride membrane using a Transblot SD apparatus (Bio-Rad). After the blots had been washed
with tris-buffered saline containing Tween-20 (TBST, 10 mM Tris-HCl, 150 mM NaCl, 0.05%
Tween-20), the membranes were blocked with 2% BSA and 4% skim milk/TBST at RT for 1
h and incubated at 4°C overnight with primary antibodies, followed by an anti-rabbit IgG
horseradish peroxidase (HRP)-conjugated secondary antibody (1:5,000, 7074; Cell Signaling
Technology, Danvers, MA, USA) at RT for 1 h. The western blot membrane was stripped the
primary and secondary antibodies and reprobed with the anti-β-actin antibody followed by the
anti-rabbit IgG HRP-conjugated secondary antibody. The following primary antibodies were
used for the blots: rabbit anti-α2A-AR (1:500; AAR020, alomone), rabbit anti-GAPDH
(1:1000; LF-PA0018, Abfrontier, Seoul, Korea), mouse anti-beta actin (1:2000, A5441,
MilliporeSigma) and rabbit anti-sodium potassium ATPase (1:1000; ab76020, Abchem)
(**Figure 3C, F and G**); rabbit anti-TH (1:1600; GTX113016, GeneTex, Irvine, CA, USA),
rabbit anti-MAO-A (1:6400; ab126751, Abcam), mouse anti-legumain (AEP) (1:400; sc-
133234, Santa Cruz), mouse anti-TAU-5 (1:1500, Santa Cruz Biotechnology, SC-58860) and
mouse anti-beta actin (1:12800, 66009-1-Ig, Proteintec, Rosemont, IL, USA) (**Figure 5A, D**
and G); rabbit anti-tau AEP-cleaved (N368) (1:5000, ABN1703, MilliporeSigma) and mouse

anti-beta actin (1:1000, A5441, MilliporeSigma) (**Figure 5G, I**). Western blot analysis was performed by measuring band intensity with ImageJ 1.50i software (National Institutes of Health, Bethesda, MD, USA).

Quantitative real-time PCR

cDNA was synthesized with Super Script II Reverse Transcriptase (18064, Invitrogen, Carlsbad, CA, USA) with 1 mg of LC neuron RNA. Using a 7300 Gene Amp PCR system (Applied Biosystems, Carlsbad), quantitative real-time PCR was run in Power SYBR Green PCR master mix (4367656; Life Technologies, Carlsbad) under the following thermal cycling conditions; an initial step of 94°C/5 min, followed by 40 cycles of a series of steps of 94°C/45 s, 61°C/45 s, 72°C/45 s and ended with a series of steps of 95°C/60 s, 60°C/1 min, 95°C/15 s, 60°C/15 s. The following primers were used: α2A (forward GGAATCATGGCTGTGGAGAT, reverse CAGAAATCCCTTCCCTGTCA); α2C (forward TCATCGTTTTCACCGTGGTA, reverse GCTCATTGGCCAGAGAAAAG); GIRK1 (forward ACCCTGGTGGATCTCAAGTG, reverse GGCCACACAGGGAGTGTAGT); GRIK2 (forward CAACCTCAACGGGTTTGTCT, reverse ATCCTACCATGAAGGCGTTG); GIRK3 (forward CGTCTCACCTCTCGTCATCA, reverse CATCCACCAGGTACGAGCTT); GAPDH (forward CCAGAACATCATCCCTGCAT, reverse GCATCGAAGGTGGAAGAGTG).

Behavioral test

The control mice as non-stress group and mice exposed to RS for 1 day, 3 days and 5 days as 1-day, 3-day and 5-day RS groups were tested on the Y maze apparatus, consisted of three identical arms in a "Y" shape made of white acryl (35 cm in length, 5 cm in width, and 20 cm in height) to assess working spatial memory. All tests were performed between 9 AM and noon. Investigators were blinded to the group when performing experiments. Each mouse was placed at the extremity of the "start arm" and left for a training session of 15 minutes, with one of the 2 other arms closed with a white acrylic board (the closed arm was randomly chosen for each mouse). Then, the mouse was brought back to its cage for 5 minutes, the maze was cleaned, the closed arm was opened, and the mouse placed at the extremity of the start arm for a 5-minute test session, under video tracking. The time spent and distance traveled in each arm (start, familiar, and novel arms) were measured and analyzed with Smart 3.0 video tracking

software (Panlab, Harvard Apparatus, Holliston, MA, USA). Given their natural curiosity, mice are expected to spend more time exploring the new arm.

Drug application

Atipamezole as a blocker of α2-AR and tertiapin-Q as a GIRK blocker were bath-applied at 5-10 μM and 200 nM, respectively. Barbadin as an arrestin/AP2 blocker was added to the internal solution at a final concentration of 100 μM (*Beautrait et al., 2017*). All chemicals except tertiapin-Q were purchased from MilliporeSigma unless specified otherwise. Tertiapin-Q was purchased from Tocris Bioscience (Bristol, UK).

Statistical Analysis

Statistical analysis was performed using STATISTICA10J (StatSoft Japan, Tokyo, Japan). Numerical data were expressed as the mean ± SD. The statistical significance was assessed using Student's t-test, one-way ANOVA followed by post hoc Fisher's PLSD and Wilk's lambda. The significance level was set at 5% or less ($p < 0.05$). Statistical results are given as a p value, unless $p < 0.001$. All the statistical data contained in this study are presented in Table S1.

Acknowledgements

This research was funded by Japan Society for the Promotion of Science (26290006 and 21K06441 to Y.K.: 20K06926 and 23K06346 to H.T.). Y.K. was supported by Brain Pool Program (NRF) funded by the Korean government (Ministry of Science and ICT). H.T. was supported by Naito Foundation. This research was partly supported by the Basic Science Research Program through the National Research Foundation of Korea grants (NRF-2018R1A5A2024418 and NRF-2021R1A2C3003334 to S.B.O) funded by the Korean government (Ministry of Science and ICT).

Author Contributions

YK, HT, and SBO conceived and designed the study. HT, DK, BGK, and TS performed experiments. HT, DK, BGK, TS, TK, and YK analyzed data. YK, HT and SBO wrote the manuscript.

Ethics

619 This study was conducted following the Helsinki Declaration, and carried out according to the

620 ARRIVE guidelines. All experimental procedures were approved and performed in accordance

621 with the relevant guidelines of the Ethical Guidelines Committee at Osaka University, Seoul

622 National University and Kyushu University.

Data Availability

625 All data and/or analyses generated during the current study are available from the

626 corresponding author upon request.

Competing Interests

629 The authors declare no competing financial or non-financial interests.

References

Aiken SP, Lampe BJ, Murphy PA, Brown BS. 1995. Reduction of spike frequency adaptation
and blockade of M-current in rat CA1 pyramidal neurones by linopirdine (DuP 996), a
neurotransmitter release enhancer. *Br J Pharmacol* **115**:1163-1168.

Aoki C, Go CG, Venkatesan C, Kurose H. 1994. Perikaryal and synaptic localization of α_{2A}-
adrenergic receptor-like immunoreactivity. *Brain Res* **650**:181-204.

Arima J, Kubo C, Ishibashi H, Akaike N. 1998. α_2-Adrenoceptor-mediated potassium currents
in acutely dissociated rat locus coeruleus neurones. *J Physiol* **508 (Pt 1)**:57-66.

Aziz Q, Thomas AM, Khambra T, Tinker A. 2012. Regulation of the ATP-sensitive potassium
channel subunit, Kir6.2, by a Ca^{2+}-dependent protein kinase C. *J Biol Chem* **287**:6196-6207.

Barcomb K, Olah SS, Kennedy MJ, Ford CP. 2022. Properties and modulation of excitatory
inputs to the locus coeruleus. *J Physiol* **600**:4897-4916.

Beattie EC, Carroll RC, Yu X, Morishita W, Yasuda H, von Zastrow M, Malenka RC. 2000.
Regulation of AMPA receptor endocytosis by a signaling mechanism shared with LTD. *Nat
Neurosci* **3**:1291-1300.

Beautrait A, Paradis JS, Zimmerman B, Giubilaro J, Nikolajev L, Armando S, Kobayashi H,
Yamani L, Namkung Y, Heydenreich FM, Khoury E, Audet M, Roux PP, Veprintsev DB,
Laporte SA, Bouvier M. 2017. A new inhibitor of the β-arrestin/AP2 endocytic complex
reveals interplay between GPCR internalization and signalling. *Nat Commun* **8**:15054.

Bisht K, Sharma K, Tremblay ME. 2018. Chronic stress as a risk factor for Alzheimer's disease:
Roles of microglia-mediated synaptic remodeling, inflammation, and oxidative stress.
Neurobiol Stress **9**:9-21.

Bloom FE. 2006. Goodman & Gilman's the pharmacological basis of therapeutics (Brunton, L.
L. et al., eds). New York: McGraw-Hill; 2006: 283-304.

Borodovitsyna O, Joshi N, Chandler D. 2018. Persistent Stress-Induced Neuroplastic Changes
in the Locus Coeruleus/Norepinephrine System. *Neural Plast* **2018**:1892570.

Burke WJ, Kristal BS, Yu BP, Li SW, Lin TS. 1998. Norepinephrine transmitter metabolite
generates free radicals and activates mitochondrial permeability transition: a mechanism for
DOPEGAL-induced apoptosis. *Brain Res* **787**:328-332.

Burke WJ, Li SW, Schmitt CA, Xia P, Chung HD, Gillespie KN. 1999. Accumulation of 3,4-
dihydroxyphenylglycolaldehyde, the neurotoxic monoamine oxidase A metabolite of
norepinephrine, in locus ceruleus cell bodies in Alzheimer's disease: mechanism of neuron
death. *Brain Res* **816**:633-637.

Campos-Lira E, Kelly L, Seifi M, Jackson T, Giesecke T, Mutig K, Koshimizu TA, Hernandez
VS, Zhang L, Swinny JD. 2018. Dynamic modulation of mouse Locus coeruleus neurons
by vasopressin 1a and 1b receptors. *Front Neurosci* **12**:919.

Cao X, Wei Z, Gabriel GG, Li X, Mousseau DD. 2007. Calcium-sensitive regulation of
monoamine oxidase-A contributes to the production of peroxyradicals in hippocampal
cultures: implications for Alzheimer disease-related pathology. *BMC Neurosci* **8**:73.

Chen Y, Chen T, Hou R. 2022. Locus coeruleus in the pathogenesis of Alzheimer's disease: A
systematic review. *Alzheimers Dement (N Y)* **8**:e12257.

Chu X, Zhou Y, Hu Z, Lou J, Song W, Li J, Liang X, Chen C, Wang S, Yang B, Chen L, Zhang
X, Song J, Dong Y, Chen S, He L, Xie Q, Chen X, Li W. 2016. 24-hour-restraint stress
induces long-term depressive-like phenotypes in mice. *Sci Rep* **6**:32935.

Costa VM, Carvalho, F., Bastos, M.L., Carvalho, R.A., Carvalho, M., Remião, F. 2012.
Adrenaline and noradrenaline: Partners and actors in the same play. (Contreras, C.M.,eds).
In: Neuroscience-Dealing with Frontiers. InTech, 2012: 1-44.

Cottingham C, Chen Y, Jiao K, Wang Q. 2011. The antidepressant desipramine is an arrestin-
biased ligand at the α_{2A}-adrenergic receptor driving receptor down-regulation in vitro and
in vivo. *J Biol Chem* **286**:36063-36075.

Counts SE, Mufson EJ. 2010. Noradrenaline activation of neurotrophic pathways protects
against neuronal amyloid toxicity. *J Neurochem* **113**:649-660.

DeGraff JL, Gagnon AW, Benovic JL, Orsini MJ. 1999. Role of arrestins in endocytosis and
signaling of α_2-adrenergic receptor subtypes. *J Biol Chem* **274**:11253-11259.

Eisenhofer G, Kopin IJ, Goldstein DS. 2004. Catecholamine metabolism: a contemporary view
with implications for physiology and medicine. *Pharmacol Rev* **56**:331-349.

Eisenhofer G, Esler MD, Meredith IT, Dart A, Cannon RO, 3rd, Quyyumi AA, Lambert G,
Chin J, Jennings GL, Goldstein DS. 1992. Sympathetic nervous function in human heart as
assessed by cardiac spillovers of dihydroxyphenylglycol and norepinephrine. *Circulation*
85:1775-1785.

Feng J, Zhang C, Lischinsky JE, Jing M, Zhou J, Wang H, Zhang Y, Dong A, Wu Z, Wu H,
Chen W, Zhang P, Zou J, Hires SA, Zhu JJ, Cui G, Lin D, Du J, Li Y. 2019. A Genetically
Encoded Fluorescent Sensor for Rapid and Specific In Vivo Detection of Norepinephrine.
Neuron **102**:745-761 e748.

Flugge G. 1996. Alterations in the central nervous α2-adrenoceptor system under chronic
psychosocial stress. *Neuroscience* **75**:187-196.

Goedert M, Spillantini MG, Potier MC, Ulrich J, Crowther RA. 1989. Cloning and sequencing
of the cDNA encoding an isoform of microtubule-associated protein tau containing four
tandem repeats: differential expression of tau protein mRNAs in human brain. *EMBO J*
8:393-399.

Goldstein DS. 2021. The catecholaldehyde hypothesis for the pathogenesis of
catecholaminergic neurodegeneration: What we know and what we do not know. *Int J Mol*
Sci **22**.

Gray M, Bingham B, Viau V. 2010. A comparison of two repeated restraint stress paradigms
on hypothalamic-pituitary-adrenal axis habituation, gonadal status and central neuropeptide
expression in adult male rats. *J Neuroendocrinol* **22**:92-101.

Hettie KS, Glass TE. 2014. Coumarin-3-aldehyde as a scaffold for the design of tunable PET-
modulated fluorescent sensors for neurotransmitters. *Chemistry* **20**:17488-17499.

Huang HP, Wang SR, Yao W, Zhang C, Zhou Y, Chen XW, Zhang B, Xiong W, Wang LY, Zheng
LH, Landry M, Hokfelt T, Xu ZQ, Zhou Z. 2007. Long latency of evoked quantal transmitter
release from somata of locus coeruleus neurons in rat pontine slices. *Proc Natl Acad Sci U*
S A **104**:1401-1406.

Itoi K, Sugimoto N. 2010. The brainstem noradrenergic systems in stress, anxiety and
depression. *J Neuroendocrinol* **22**:355-361.

Jedema HP, Gold SJ, Gonzalez-Burgos G, Sved AF, Tobe BJ, Wensel T, Grace AA. 2008.
Chronic cold exposure increases RGS7 expression and decreases α2-autoreceptor-mediated
inhibition of noradrenergic locus coeruleus neurons. *Eur J Neurosci* **27**:2433-2443.

Kang SS, Liu X, Ahn EH, Xiang J, Manfredsson FP, Yang X, Luo HR, Liles LC, Weinshenker
D, Ye K. 2020. Norepinephrine metabolite DOPEGAL activates AEP and pathological Tau
aggregation in locus coeruleus. *J Clin Invest* **130**:422-437.

Kielbasa W, Pan A, Pereira A. 2015. A pharmacokinetic/pharmacodynamic investigation:
assessment of edivoxetine and atomoxetine on systemic and central 3,4-
dihydroxyphenylglycol, a biochemical marker for norepinephrine transporter inhibition. *Eur*
Neuropsychopharmacol **25**:377-385.

Kitayama IT, Otani M, Murase S. 2004. Contribution of the stress-induced degeneration of the locus coeruleus noradrenergic neurons to the pathophysiology of depression: a study on an animal model. *Acta Neuropsychiatr* **16**:190-199.

Koenig JH, Ikeda K. 1996. Synaptic vesicles have two distinct recycling pathways. *J Cell Biol* **135**:797-808.

Leistner C, Menke A. 2020. Hypothalamic-pituitary-adrenal axis and stress. *Handb Clin Neurol* **175**:55-64.

Mao L, Han Y, Zhang QW, Tian Y. 2023. Two-photon fluorescence imaging and specifically biosensing of norepinephrine on a 100-ms timescale. *Nat Commun* **14**:1419.

Mather M, Harley CW. 2016. The locus coeruleus: essential for maintaining cognitive function and the aging brain. *Trends Cogn Sci* **20**:214-226.

McCall JG, Al-Hasani R, Siuda ER, Hong DY, Norris AJ, Ford CP, Bruchas MR. 2015. CRH Engagement of the Locus Coeruleus Noradrenergic System Mediates Stress-Induced Anxiety. *Neuron* **87**:605-620.

Mitchell NC, Bowman MA, Gould GG, Koek W, Daws LC. 2017. Ontogeny of norepinephrine rransporter expression and antidepressant-like response to desipramine in wild-type and serotonin transporter mutant mice. *J Pharmacol Exp Ther* **360**:84-94.

Pierce KL, Lefkowitz RJ. 2001. Classical and new roles of β-arrestins in the regulation of G-protein-coupled receptors. *Nat Rev Neurosci* **2**:727-733.

Potter LT, Axelrod J. 1963. Properties of norepinephrine storage particles of the rat heart. *J Pharmacol Exp Ther* **142**:299-305.

Ross JA, McGonigle P, Van Bockstaele EJ. 2015. Locus coeruleus, norepinephrine and Aβ peptides in Alzheimer's disease. *Neurobiol Stress* **2**:73-84.

Sakaguchi T, Nakamura S. 1990. Duration-dependent effects of repeated restraint stress on cortical projections of locus coeruleus neurons. *Neurosci Lett* **118**:193-196.

Sanders JD, Happe HK, Bylund DB, Murrin LC. 2005. Development of the norepinephrine transporter in the rat CNS. *Neuroscience* **130**:107-117.

Sarmento MJ, Coutinho A, Fedorov A, Prieto M, Fernandes F. 2014. Ca^{2+} induces $PI(4,5)P_2$ clusters on lipid bilayers at physiological $PI(4,5)P_2$ and Ca^{2+} concentrations. *Biochim Biophys Acta* **1838**:822-830.

Stone LS, Broberger C, Vulchanova L, Wilcox GL, Hokfelt T, Riedl MS, Elde R. 1998. Differential distribution of α2A and α2C adrenergic receptor immunoreactivity in the rat spinal cord. *J Neurosci* **18**:5928-5937.

Sugama S, Sekiyama K, Kodama T, Takamatsu Y, Takenouchi T, Hashimoto M, Bruno C, Kakinuma Y. 2016. Chronic restraint stress triggers dopaminergic and noradrenergic neurodegeneration: Possible role of chronic stress in the onset of Parkinson's disease. *Brain Behav Immun* **51**:39-46.

Takahashi N, Miner LL, Sora I, Ujike H, Revay RS, Kostic V, Jackson-Lewis V, Przedborski S, Uhl GR. 1997. VMAT2 knockout mice: heterozygotes display reduced amphetamine-conditioned reward, enhanced amphetamine locomotion, and enhanced MPTP toxicity. *Proc Natl Acad Sci U S A* **94**:9938-9943.

Takei K, Mundigl O, Daniell L, De Camilli P. 1996. The synaptic vesicle cycle: a single vesicle budding step involving clathrin and dynamin. *J Cell Biol* **133**:1237-1250.

Tan M, Walwyn WM, Evans CJ, Xie CW. 2009. p38 MAPK and β-arrestin 2 mediate functional interactions between endogenous micro-opioid and α2A-adrenergic receptors in neurons. *J Biol Chem* **284**:6270-6281.

Taraviras S, Olli-Lahdesmaki T, Lymperopoulos A, Charitonidou D, Mavroidis M, Kallio J, Scheinin M, Flordellis C. 2002. Subtype-specific neuronal differentiation of PC12 cells transfected with α2-adrenergic receptors. *Eur J Cell Biol* **81**:363-374.

Torres GE, Gainetdinov RR, Caron MG. 2003. Plasma membrane monoamine transporters: structure, regulation and function. *Nat Rev Neurosci* **4**:13-25.

Toyoda H, Won J, Kim W, Kim H, Davy O, Saito M, Kim D, Tanaka T, Kang Y, Oh SB. 2022. The Nature of Noradrenergic Volume Transmission From Locus Coeruleus to Brainstem Mesencephalic Trigeminal Sensory Neurons. *Front Cell Neurosci* **16**:841239.

Valentino RJ, Foote SL. 1988. Corticotropin-releasing hormone increases tonic but not sensory-evoked activity of noradrenergic locus coeruleus neurons in unanesthetized rats. *J Neurosci* **8**:1016-1025.

Valentino RJ, Foote SL, Aston-Jones G. 1983. Corticotropin-releasing factor activates noradrenergic neurons of the locus coeruleus. *Brain Res* **270**:363-367.

Valentino RJ, Rudoy C, Saunders A, Liu XB, Van Bockstaele EJ. 2001. Corticotropin-releasing factor is preferentially colocalized with excitatory rather than inhibitory amino acids in axon terminals in the peri-locus coeruleus region. *Neuroscience* **106**:375-384.

Williams JT, Henderson G, North RA. 1985. Characterization of α2-adrenoceptors which increase potassium conductance in rat locus coeruleus neurones. *Neuroscience* **14**:95-101.

Zaccor NW, Sumner CJ, Snyder SH. 2020. The nonselective cation channel TRPV4 inhibits angiotensin II receptors. *J Biol Chem* **295**:9986-9997.

Zhang Y, Woodruff M, Zhang Y, Miao J, Hanley G, Stuart C, Zeng X, Prabhakar S, Moorman J, Zhao B, Yin D. 2008. Toll-like receptor 4 mediates chronic restraint stress-induced immune suppression. *J Neuroimmunol* **194**:115-122.

Zhang Z, Song M, Liu X, Kang SS, Kwon IS, Duong DM, Seyfried NT, Hu WT, Liu Z, Wang JZ, Cheng L, Sun YE, Yu SP, Levey AI, Ye K. 2014. Cleavage of tau by asparagine endopeptidase mediates the neurofibrillary pathology in Alzheimer's disease. *Nat Med* **20**:1254-1262.

 Figure legends

Figure 1. Atipamezole abolishes spike-frequency adaptation and blocks NA-induced

GIRK-I in LC neurons.

(**A, B**) Sample traces of spike trains induced in an LC neuron by injection of current pulses at

75 and 100 pA. Note the spike-frequency adaptation and pulse-afterhyperpolarization (pulse-

AHP).

(**C**) Plotting of the cumulative spike numbers vs the elapsed time from the onset of the current

pulses every 50 ms at 75 pA (blue circles) or 100 pA (red circles) ($n = 7$). Two-way RM

ANOVA, *$p < 0.001$. The saturation level (***a***) and the half saturation constant (***b***) were

measured by fitting the saturation function defined as y = (***a*** * x) / (***b*** + x) to the data points.

The values of ***a*** at 75 and 100 pA current pulses were 44.6 ± 16.2 and 52.5 ± 12.3, respectively

(paired t-test, $p = 0.010$). The values of ***b*** at 75 and 100 pA current pulses were 2.1 ± 0.7 and

2.1 ± 0.4, respectively (paired t-test, $p = 0.823$).

(**D, E**) Sample traces of spike trains induced in an LC neuron evoked by current pulses at 100

pA under control condition and after application of 10 µM atipamezole. Note that atipamezole

suppresses spike-frequency adaptation and pulse-AHP (blue and red stars).

(**F**) Plotting of the cumulative spike numbers vs the elapsed time during the current pulses

every 50 ms before and after application of atipamezole (blue and red circles, respectively) (n

= 10). Two-way RM ANOVA, *$p < 0.001$.

(**G**) Plotting of ***a*** against ***b***, which were measured by fitting the saturation function to the data

points in **F**. Both the values of ***a*** and ***b*** obtained by curve fitting to the data points after

application of atipamezole (open and filled red circles) were significantly shifted to larger

values than those obtained under control condition (open and filled blue circles) (paired t-test,

a and ***b***, [‡]$p = 0.003$ and [‡]$p = 0.002$, respectively), and there was a significant difference in the

relationship between ***a*** and ***b*** (Wilk's lambda, [§]$p = 0.002$) ($n = 10$).

(**H**) Top, Voltage command pulse. Lower panel, Superimposed current traces obtained under

control condition (1), after application of 100 µM NA (2), and after application of 10 µM

atipamezole (Atip) in addition to NA (3). The amplitudes of inward and outward components

in NA-induced GIRK-I were -70 ± 22 and 36 ± 11 pA, respectively (paired t-test, [‡]$p = 0.005$,

$n = 6$).

(**I**) A graph showing the effects of atipamezole on amplitudes of the inward and outward

components of NA-induced GIRK-I at -130 mV and -60 mV, obtained by subtraction of

currents recorded under control condition from those recorded after application of NA (2–1)

and from those recorded after application of NA and atipamezole (3–1). Inward component,

paired *t*-test, [‡]*p* < 0.001; outward component, paired *t*-test, [‡]*p* < 0.001 (*n* = 6).

Figure 2. Ca^{2+}-dependent rundown of NA-induced GIRK-I and its suppression by

barbadin.

(**A**) Upper panel, Ramp command pulse. Lower panel, Superimposed current traces obtained

under control condition (1) and after application of 100 μM NA for 5 min (2).

(**B**) Upper panel, A combined command pulse applied every minute; five trains of 20 positive

pulses (5 ms duration to 0 mV at 20 Hz) at an interval of 2 s in the presence of extracellular 30

mM TEA and intracellular 0.2 mM EGTA, which were followed by the ramp pulse after an

interval of 25 s. Lower panel, representative Ca^{2+} currents in response to positive pulses in the

presence of NA and 30 mM TEA.

(**C**) Upper panel, Ramp command pulse. Lower panel, Superimposed current traces obtained

after application of NA for 5 min (2) and in response to application of the 20th combined pulse

in the presence of NA and TEA (3).

(**D**) The I-V relationship of NA-induced GIRK-I obtained by subtraction of currents recorded

under control condition from that recorded after application of NA for 5 min (red trace, 2–1)

and that obtained by subtraction of the control current from that recorded in response to

application of the 20th combined pulse in the presence of NA and TEA (blue trace, 3–1).

(**E**) Plotting of amplitudes of inward components at –130 mV (blue circles) and outward

components at –60 mV (red circles) against time.

(**F**) Plotting of normalized amplitudes of inward components at –130 mV and outward

components at –60 mV before and 20 min after application of positive pulse trains in the

presence of NA and TEA. The amplitudes of inward components at –130 mV and those of

outward components at –60 mV were normalized by those recorded in the presence of NA

before applying positive pulse trains. Inward component, paired *t*-test, [‡]*p* < 0.001; outward

component, paired *t*-test, [‡]*p* < 0.001 (*n* = 8).

(**G**) Upper panel, Ramp command pulse. Lower panel, Superimposed current traces obtained

under control condition (1) and after application of 100 μM NA for 5 min (2) in LC neurons

dialyzed with 100 μM barbadin (β-arrestin/AP2 blocker).

(**H**) Upper panel, A combined command pulse applied every minute; five trains of 20 positive

pulses (5 ms duration to 0 mV at 20 Hz) at an inter-train interval of 2 s in the presence of

extracellular 30 mM TEA and intracellular 0.2 mM EGTA, which were followed by the ramp

pulse after an interval of 25 s. Lower panel, Superimposed current traces obtained after

application of NA for 5 min (2) and in response to application of the 20th combined pulse in

the presence of NA and TEA (3) in LC neurons dialyzed with barbadin.

(**I**) The I-V relationship of NA-induced GIRK-I obtained by subtraction of currents recorded

under control condition from that recorded after application of NA for 5 min (red trace, 2–1)

and that obtained by subtraction of the control current from that recorded in response to

application of the 20th combined pulse in the presence of NA and TEA (blue trace, 3–1).

(**J**) Plotting of normalized amplitudes (left vertical axis) of inward components at –130 mV

(filled blue circles) and outward components at –60 mV (filled red circles) against time ($n = 8$).

Amplitudes of GIRK-I were normalized to the values obtained before applying pulse trains.

Right vertical axis refers to the original amplitudes. Open blue and red circles represent the

normalized amplitudes of GIRK-I shown in **E**.

(**K**) Plotting of normalized amplitudes of inward components at –130 mV and outward

components at –60 mV before and 20 min after application of positive pulse trains in the

presence of NA and TEA ($n = 8$). The amplitudes of inward components at –130 mV and those

of outward components at –60 mV were normalized to those recorded in the presence of NA

before applying positive pulse trains. Open light pink circles represent results obtained under

the control condition in **F**. Inward component, unpaired t-test, *$p < 0.001$; outward component,

unpaired t-test, *$p < 0.001$.

Figure 3. RS reduces expression of α2A-ARs in LC Neurons.

(**A, B**) Confocal images of LC neurons showing immunoreactivities for TH and α2A-ARs,

together with a merged one in non-stress and 3-day RS mice. Arrowheads indicate the

membrane regions of TH-positive neurons, along which α2A-ARs were differentially

expressed between the control and the RS mice. Asterisks indicate LC neurons.

(**C**) Western blotting analyses showing the expression of α2A-ARs in non-stress and 3-day RS

mice ($n = 6$ and 7, respectively). Unpaired t-test, [†]$p < 0.001$.

(**D, E**) Confocal images of LC neurons in non-stress and RS mice. Upper panels from left to

right showing the respective immunoreactivity for TH, α2A-ARs (with rabbit anti-α2A-AR)

and Na^+-K^+-pump and an enlarged image of the region enclosed with a rectangle (arrowheads)

in its immediate left panel. Lower panels from left to right showing the merged image of TH

and α2A-ARs, that of TH, α2A-ARs and Na^+-K^+-pump, that of α2A-ARs and Na^+-K^+-pump

and an enlarged image of the region enclosed with a rectangle in its immediate left panel.

(**F, G**) Western blotting analyses showing expressions of α2A-ARs and Na$^+$-K$^+$-pump in

membrane fraction and those of α2A-ARs and β-actin in cytosol fraction in non-stress and 5-

day RS mice ($n = 3$ and 3 samples, respectively). Each sample represents the analysis result in

the LC tissues obtained from 2-3 mice. Membrane; unpaired t-test, $^†p = 0.016$. Cytosol;

unpaired t-test, $^†p = 0.083$.

Figure 4. RS reduces NA-induced GIRK-I in LC neurons.

(**A-E**) Representative traces of NA-induced GIRK-I obtained from LC neurons in non-stress

and 1-day, 2-day, 3-day, and 5-day RS mice.

(**F, G**) Amplitudes of inward and outward components of NA-induced GIRK-I obtained from

LC neurons in non-stress and 1-day, 2-day, 3-day, and 5-day RS mice ($n = 8, 7, 5, 6$ and 8,

respectively) decreased with the increase in the period of RS, in a way that can be described by

a saturation function (red interrupted lines). The saturation level ($\boldsymbol{a} + \boldsymbol{b}$) and the half saturation

constant ($\boldsymbol{c}$) were determined by fitting the saturation function, defined as y = $\boldsymbol{a} + (\boldsymbol{b}* $ x$) / (\boldsymbol{c} +$

x), to the data points. The values of $\boldsymbol{a}, \boldsymbol{b}$ and $\boldsymbol{c}$ for the inward component of GIRK-I were −151.1,

108, and 0.9, respectively, and those for the outward component of GIRK-I were 81.1, –67,

and 1.6, respectively. Inward component: one-way ANOVA, $p < 0.001$, post hoc fisher's

PLSD, 1-day; $^#p < 0.001$ vs Non-stress and $^*p < 0.05$ vs 3-day and 5-day, 2-day; $^#p < 0.001$ vs

Non-stress, 3-day; $^#p < 0.001$ vs Non-stress, 5-day; $^#p < 0.001$ vs Non-stress. Outward

component: one-way ANOVA, $p < 0.001$, post hoc fisher's PLSD, 1-day; $p = 0.004$ vs Non-

stress and $^*p < 0.05$ vs 3-day and 5-day, 2-day; $p = 0.015$ vs Non-stress, 3-day; $p < 0.001$ vs

Non-stress, 5-day; $p < 0.001$ vs Non-stress.

(**H**) Relative expressions of α2A and α2C mRNAs, normalized to GAPDH in LC neurons ($n =$

5). Paired t-test, $^‡p = 0.014$.

(**I**) Relative expressions of GIRK1, GIRK2, and GIRK3 mRNAs, normalized to GAPDH in

LC neurons ($n = 5$). One-way RM ANOVA, $p < 0.001$, post hoc fisher's PLSD; $^*p = 0.008$

for GIRK1 vs GIRK2, $^*p < 0.001$ for GIRK1 vs GIRK3, $^*p = 0.008$ for GIRK2 vs GIRK3.

(**J, K**) Relative expressions of α2A and GIRK1 mRNAs, respectively, normalized to GAPDH

in LC neurons in non-stress mice ($n = 8$), and 3-day RS mice ($n = 6$). α2A-AR: unpaired t-test,

$^†p = 0.020$; GIRK1: unpaired t-test, $^†p = 0.013$.

(**L**) Normalized relative expressions of GIRK2 mRNA in LC neurons in non-stress mice ($n =$

8) and 3-day RS mice ($n = 6$), normalized to the ratio of the mean value of the relative

expressions of GIRK1 mRNA to that of GIRK2 mRNA in LC neurons in non-stress mice (**I**).

Unpaired t-test, $^{\dagger}p = 0.037$.

Figure 5. Expressions of TH, MAO-A, DBH, AEP and tau proteins in LC neurons.

(**A**) Western blotting of TH and MAO-A in LC neurons obtained from non-stress and 5-day

RS mice.

(**B, C**) Western blot analysis revealing no significant change in TH expression ($n = 10$ samples)

but a significant increase in MAO-A expression ($n = 6$ samples) in 5-day RS group compared

to non-stress group. Unpaired t-test, $^{\dagger}p = 0.029$.

(**D**) Western blotting of pro-AEP and active AEP in LC neurons obtained from non-stress and

5-day RS mice.

(**E, F**) Western blot analysis revealing significant increases in pro-AEP and active AEP ($n = 6$

and $n = 6$ samples, respectively) in 5-day RS group compared to non-stress group. Unpaired

t-test, $^{\dagger}p = 0.034$ (**E**) and $^{\dagger}p = 0.047$ (**F**).

(**G**) Western blotting of tau and tau N368 fragment in LC neurons obtained from non-stress

and 5-day RS mice. Unpaired t-test, $^{\dagger}p = 0.025$.

(**H, I**) Western blot analyses revealing a significant increase in tau and tau N368 fragment in

LC neurons ($n = 8$, and $n = 6$, respectively) in 5-day RS group compared to non-stress group.

Each sample represents the analysis result in the LC tissues obtained from 2-3 mice. Unpaired

t-test, $^{\dagger}p < 0.05$.

Figure 6. Presumed cellular mechanisms for the degeneration of LC by stress.

(**A, B**) Differential free concentrations of NA to be metabolized by MAO-A into DOPEGAL

between control and stress conditions: Under the control condition (**A**), NA in the cytosol is

mostly taken up into the cytoplasmic vesicles by VMAT2 (thick arrow 3), rather than being

directly metabolized by MAO-A into DOPEGAL (thin interrupted arrow). Vesicular NA is

released from cell bodies as autocrine following $[Ca^{2+}]_i$ increases caused by action potentials,

and the released NA activates α2A-AR-coupled GIRK channels, causing autoinhibition.

Subsequently, the autocrine released NA is slowly taken up into the cytosol of LC neurons by

NAT after dissociation from α2A-ARs. Under the stress condition (**B**), an activation of CRF

receptors in LC neurons by stress inhibits leak K^+ channels and increases firing activities in LC

neurons, subsequently causing a larger $[Ca^{2+}]_i$ increase together with Ca^{2+}-induced Ca^{2+} release

(CICR). Impairment of autoinhibiton due to Ca^{2+} dependent internalization of α2A-ARs-

coupled GIRK channels leads to the persistent excitation in LC neurons, which enhances
autocrine release of NA (thick arrow 1). Subsequently, the excessively autocrine released NA
is taken up directly and rapidly by NAT into the cytosol without binding to α2A-ARs (thick
arrow 2). Such a facilitation of re-uptake of NA by NAT would increase active NA storage into
cytoplasmic vesicles by VMAT2 (thick arrow 3), while the rate of NA leakage from
cytoplasmic vesicles would also increase (arrow 4) due to a dynamic equilibrium in
cytoplasmic vesicles between active NA storage into cytoplasmic vesicles and passive NA
leakage from cytoplasmic vesicles. Subsequently, such an increase in the rate of NA leakage
would result in an increase in a MAO-A metabolite, DOPEGAL and AEP, leading to a
production of cleaved tau N368 fragment and an impairment of learning/memory.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6