1

# The relationship between cortisol, grid-like representations and path

# integration

Osman Akan[1*], Varna Chandreswaran[2], Anne Bierbrauer[3], Henry Soldan[1], Nikolai

Axmacher[2], Oliver T. Wolf[1], Christian J. Merz[1]

[1] Department of Cognitive Psychology, Institute of Cognitive Neuroscience, Faculty of

Psychology, Ruhr University Bochum, Bochum, Germany

[2] Department of Neuropsychology, Institute of Cognitive Neuroscience, Faculty of

Psychology, Ruhr University Bochum, Bochum, Germany

[3] Institute of Systems Neuroscience, Medical Center Hamburg-Eppendorf, Hamburg,

Germany

[*]Corresponding author

E-mail: osman.akan@rub.de

**Abstract**

Acute stress triggers the release of cortisol, which broadly affects cognitive processes.

Path integration, a specific navigational process, relies heavily on grid cells in the

entorhinal cortex (EC). The EC contains glucocorticoid receptors and is therefore likely

to be influenced by cortisol, though little is known about this relationship. Given the role

of the EC in neurological diseases such as Alzheimer's Disease, investigating the effects

of cortisol on this brain region may offer insights into how stress affects these diseases.

In this study, we examined the effects of cortisol on human path integration in thirty-nine

healthy participants across two sessions. On each day, they received either 20mg

cortisol or a placebo and performed a virtual homing task during functional magnetic

resonance imaging (fMRI). Cortisol markedly impaired path integration performance,

independent of path distance or the presence of spatial cues. Additionally, cortisol altered

navigational strategies, leading participants to navigate further away from landmarks,

which was associated with worse performance. FMRI results showed that cortisol

increased the activation of right caudate nucleus in the presence of landmarks. Using a

representational similarity analysis, we observed grid-like representations in the right

posterior-medial EC specifically on day one under placebo, but these were diminished

by cortisol. Grid-like representations facilitated performance over short distances but

hindered it over longer ones, suggesting that grid cells support PI specifically in case of

short trajectories. Overall, the study indicates that cortisol-induced disruption in grid cell

function in the EC may underly stress effects on path integration.

**1   Introduction**

Finding ourselves in situations where we lost orientation can easily invoke feelings of

discomfort or stress, but we also often experience the inverse relationship of stress

causing spatial disorientation, mainly through altering the way we navigate. For example,

under time pressure, we tend to use more familiar and less accurate trajectories (1). In

general, however, human spatial navigation has not gained much attention in the realm

of stress-related consequences, albeit being a type of cognition that is strongly based on

hippocampus (HC) and adjacent cortical regions and thus a major target of stress

hormones. Investigating the effects of stress on navigation may ultimately deepen our

understanding of neurological diseases, particularly since navigational deficits could

serve as early biomarkers for Alzheimer's Disease (2), and stress has been linked to

both its pathogenesis and progression (3).

The stress response involves the activation of the rapid sympathetic-adrenal-medullary

axis (SAM) and the slow hypothalamic-pituitary-adrenal axis (HPA), generally

conceptualized as an adaptive reaction (4). SAM axis activity initiates the release of

catecholamines (noradrenaline and adrenaline), while HPA axis activity leads to the

release of glucocorticoids (cortisol in humans, corticosterone in most rodents). Cortisol

effects are mainly exerted by binding to mineralocorticoid receptors (MRs) and

glucocorticoid receptors (GRs), while noradrenaline binds to adrenergic receptors (5, 6).

Thereby, acute stress or pharmacological cortisol administration affect several cognitive

domains including executive functions, episodic memory, fear conditioning, and decision

making (7–10).

During spatial navigation, different strategies can be applied depending on several

factors including environmental features (11). One strategy is path integration (PI), a

process mainly used in absence of spatial cues and for short trajectories because error

accumulation makes it less suited for longer trajectories. PI involves integration of self-

referential information to estimate the current position and orientation in relation to an

arbitrary reference point (12). On the neuronal sheet, PI has been related to head-

direction cells (13) and grid cell firing in the entorhinal cortex (EC; 14–17). Specifically,

the characteristic arrangement of grid cell firing fields in a regular hexagonal pattern (18,

19) may provide a general spatial metric of distances (20). In presence of spatial cues,

however, additional neural systems tuned to the specific type of cue (e.g., boundary or

landmark) are recruited, including HC (21) and posterior cingulate/retrosplenial cortex

(PC/RSC; 22). These additional systems may either stabilize grid cell firing (23) or

support complementary navigational strategies. Because processing in the striatum,

especially the caudate nucleus (CN), is involved in landmark-based navigation (24) and appears to be enhanced under acute stress (25), the presence of spatial cues likely moderates stress hormone effects on either performance or strategy use during navigational processes.

Crucially, the EC is an important mediator of the stress response itself (26–29) and the abundance of GRs in the EC (30) as well as its interdependence with the HC makes it a likely target of cortisol effects. Accordingly, acute stress impairs PI over long distances and in environments with little spatial information (31), and chronic stress is associated with impaired PI under similar conditions (32). These effects might reflect cortisol-induced alterations in grid cell activity in the EC. Furthermore, stress appears to alter navigational patterns, leading participants to opt for routes closer to a landmark (31). This suggests an increased recruitment of landmark-based strategies under stress, possibly due to enhanced striatal processing.

In this study, we aimed to investigate whether cortisol administration mimics the effects of stress on PI and to uncover the underlying neural mechanisms using functional magnetic resonance imaging (fMRI). To assess PI performance, we used an adapted version of the "Apple Game" paradigm implemented by Bierbrauer et al. (22). In different subtasks of this virtual PI task, participants can either rely only on visual flow (Pure PI) or on visual flow and a central landmark (Landmark PI; Fig. 1). To measure proxies of grid cell activity, we analyzed so called "grid-like representations" (GLRs; 33, 22, 34).

Our results strongly support the hypothesis that stress-induced deficits of PI are mediated via cortisol, but they do not show that cortisol changes the navigational pattern towards the landmark. The fMRI data demonstrated the importance of PC and CN during Landmark PI and for the CN during Landmark PI under cortisol. Additionally, they showed that the EC is most relevant during Pure PI under placebo. Importantly, our findings suggest that cortisol impairs GLRs, which were particularly beneficial for PI over short distances.

**A Subtasks**

Pure
Path Integration (Pure PI)

Landmark-supported
Path Integration (Landmark PI)

**B Trial procedure**

| Start phase | Outgoing phase (tree without apple) | Outgoing phase (tree with apple) | Incoming phase (max. 60 s) | Feedback (2 s) |

**C Task phases and distances**

Retrieval location

Outgoing phase
Outgoing distance

Incoming phase
Incoming distance

Goal location
Start phase

**D Performance measures**

Response path & location

Retrieval location

Rotation error

Correct path

Drop error

Distance error

Goal location

Figure 1. Experimental path integration task. (A) While the Pure PI subtask consisted only of a grassy plain, the Landmark PI subtask additionally contained a central lighthouse serving as spatial cue. (B) Each trial began with the "start phase", where participants navigated to a basket (goal location), the location of which they should encode. In the following "outgoing phase", they navigated to a variable number of trees (1-5) until reaching a tree containing an apple (retrieval location). Then, during the "incoming phase", participants had to find the way back to the goal location before receiving feedback via zero to three stars according to performance based on the drop error (see D). Basket and trees disappeared as soon as they were reached. (C) Outgoing phase (dashed black line) and incoming phase (dotted black line) were quantified according to their spatial distances: outgoing distance corresponded to the cumulated distance from goal to retrieval location (dashed red line), and incoming distance to the Euclidean distance between retrieval and goal location (dotted red line). (D) General PI performance was assessed via the drop error, which corresponded to the distance between response location (marked with an X) and goal location (solid red line). The drop error can further be differentiated into distance error, referring to the difference between retrieval-to-goal distance and retrieval-to-response distance (blue line), and rotation error, depicting the angle between the retrieval-to-goal path and the retrieval-to-response path (purple arc). Figure adapted from Bierbrauer et al. (22).

## 2   Results

We analyzed data from 39 men aged 19 – 34 years (24.18 ± 4.55 years; mean ± SD)

participating in a two-day within-subject crossover design. Participants received either

20 mg of cortisol or visually identical placebo tablets on day one, and the respective other

treatment on day two (one week later), before completing the Apple Game during fMRI

recordings on both days. The order of administration was counterbalanced across

participants (Fig. 2; STAR Methods). To examine effects of cortisol on PI, we conducted

univariate and multivariate fMRI data analyses and built several linear mixed models

(see Tab. S1 for an overview). In all mixed models, "subject" was added as random

factor, and age, testing day, and sequence (to control for order effects) as covariates.

**Figure 2. Experimental procedure.** All participants were tested in both conditions (cortisol and
placebo) on two testing sessions with an inter-session interval of one week. At the beginning of
day one, participants underwent a cognitive test battery, which lasted about 40 minutes.
Participants then received either 20 mg of cortisol, or a placebo. Afterwards, questionnaires (Q1)
assessing demographic, psychological and medical data were filled out. Participants then were
prepared for the fMRI scan and the PI task started approximately 40 minutes after cortisol/placebo
administration, before answering final questionnaires including a treatment guess and
assessment of individual navigational strategies (Q2). One week later, participants were invited
for the second testing session. The procedure on day two was similar, except that it did not contain
a cognitive test battery, included less questionnaires and involved the respective other
pharmacological intervention as compared to day one (crossover design). Numbers on x-axes

reflect time-points relative to tablet administration, ✎ collection of saliva sample, PI: path
integration.

*2.1    Cortisol impairs PI performance irrespective of path distance and availability of*
146            *spatial cues*

The analysis regarding the success of cortisol administration revealed elevated cortisol

concentrations under cortisol compared to placebo treatment for the entire PI task (Tab.

S2, STAR Methods). In the PI task, we assessed different measures for path distance,

PI performance and navigational pattern during the experimental task (STAR Methods).

For path distance, we focused on *incoming distance*, which refers to the Euclidean

distance between retrieval and goal location (Fig. 1C). For overall PI performance, we

assessed the drop error, i.e., the Euclidean distance between response location and goal

location (Fig. 1D). Moreover, to investigate the specific role of the landmark under

cortisol, we measured the distance between the goal and the spatial cue (goal-to-

landmark distance), and the mean Euclidean distance of the moving participant from the

landmark across all time points of the incoming phase (movement-to-landmark distance).

To investigate whether cortisol influences PI performance, we built a linear mixed model,

in which we investigated effects of subtask (two levels: Pure PI vs. Landmark PI), path

distance (i.e., incoming distance), and treatment (two levels: cortisol vs. placebo) on

overall PI performance (i.e., drop error) on the level of single trials. This analysis revealed

similar findings as in our previous work (31, 32, 22): We observed main effects of subtask

($F_{(1,4945)}$ = 432.88, $p$ < .001, $\eta_p^2$ = .080), and of incoming distance ($F_{(1,4945)}$ = 364.99, $p$ <

.001, $\eta_p^2$ = .069), indicating higher errors in Pure PI than Landmark PI and for longer

incoming distances, respectively. Besides, we found an interaction effect between

subtask and incoming distance ($F_{(1,4949)}$ = 44.99, $p$ < .001, $\eta_p^2$ = .009; Fig. 3A, left),

indicating a stronger relationship between incoming distance and drop error in Pure PI

as compared to Landmark PI ($t_{(4949)}$ = 6.71, $p$ < .001, $d$ = 0.095). Moreover, we found a

main effect of age ($F_{(1,36)}$ = 12.23, $p$ = .001, $\eta_p^2$ = .254; Fig. 3A, right), reflecting higher

drop errors in older age. No effects of testing day ($F_{(1,4945)}$ = 0.39, $p$ = .532, $\eta_p^2$ < .001) or

sequence ($F_{(1,36)}$ = 2.18, $p$ = .148, $\eta_p^2$ = .057) emerged, indicating the absence of an

order effect. Importantly, we found a main effect of treatment on drop error ($F_{(1,4945)}$ =

10.46, $p$ = .001, $\eta_p^2$ = .002; Fig. 3B, left), indicating that cortisol impaired overall PI

performance. Cortisol did not interact with other variables, suggesting a general effect

independent of spatial cues or path distance.

In a second step, we built linear mixed models only including Landmark PI to examine

the navigational pattern in the presence of the landmark under heightened cortisol levels.

These models aimed at investigating the role of goal-to-landmark distance and its

interaction with cortisol on PI performance, and at investigating whether cortisol affected

the employment of navigational strategies (irrespective of performance), respectively. In

the first model, we found that a longer distance between goal location and landmark

predicted worse PI performance, and cortisol did not moderate this effect ($F_{(1,2453)}$ =

348.20, $p$ < .001, $\eta_p^2$ = .124). Like in the main models, older age again significantly

reduced PI performance ($F_{(1,36)}$ = 6.30, $p$ = .017, $\eta_p^2$ = .149). In the second model,

importantly, we found a main effect of treatment on movement distance to the landmark

($F_{(1,2455)}$ = 4.57, $p$ = .033, $\eta_p^2$ = .002; Fig. 3B, right), showing that cortisol treatment led

participants to navigate further away from the landmark, suggesting that they used

landmark information less than under placebo.

The behavioral results so far demonstrated that cortisol impairs PI performance and

alters navigational strategy toward navigating further away from landmarks (if available).

To analyze whether these alterations were related to each other, we conducted a further

analysis examining the effect of movement distance to the landmark on the drop error.

Indeed, navigating further away from the landmark was detrimental for performance

($F_{(1,2453)}$ = 33.46, $p$ < .001, $\eta_p^2$ = .013 ;Fig. 3C, left), and this effect was equally observed

under placebo and under cortisol as evidenced by a lack of an interaction effect ($F_{(1,2463)}$

= 0.72, $p$ = .395, $\eta_p^2$ < .001 ;Fig. 3C, right). The cortisol-induced change in navigational

strategy may thus have contributed to the overall detrimental effect of cortisol on PI

performance.

**A Non-cortisol-related effects: effects of age, subtask and incoming distance on PI**

**B Effects of cortisol on PI performance and navigational patterns**

**C Relationship between PI performance and navigational patterns**

**Figure 3.** Predictors of PI performance and strategy use. (A) The effect of longer incoming distances leading to higher drop errors was more pronounced when no spatial cues were available (left), and older age led to higher drop errors (right). (B) CORT led to overall higher drop errors (left), and to navigating further away from the landmark in Landmark PI (right). (C) In Landmark PI, the movement distance to the landmark predicted performance, irrespective of treatment. Significant slopes are indicated via asterisks. Error bars and confidence bands represent SEM. Pure PI: pure path integration, Landmark PI: landmark-supported path integration, CORT: cortisol, PLA: placebo, vm: virtual meters, n. s.: not significant, ***p < .001, **p < .01, *p < .05.

*2.2    Cortisol enhances activation in the right CN in the presence of landmarks*

We tested the contrasts "Landmark PI > Pure PI", "CORT > PLA" and their interaction

"(Landmark PI > Pure PI)$_{CORT}$ > (Landmark PI > Pure PI)$_{PLA}$" – as well as the respective

reverse contrasts – during the outgoing and incoming phases. We performed an

exploratory whole-brain analyses and ROI analyses in left and right HC, left and right

CN, bilateral PC, and left and right pmEC for each contrast. For the whole-brain analysis,

statistical parametric maps were initially thresholded at a family-wise error (FWE)-

corrected α level of $p < 0.05$ across the whole brain and clusters were considered

significant at $p < 0.05$, FWE-corrected (extent threshold of five voxels). For ROI analyses,

the significance threshold was set to $p < 0.05$ on voxel level, corrected for multiple testing

within each ROI using FWE-correction with the small volume correction option of SPM12.

The exploratory whole-brain analyses revealed significantly higher activation in left

superior parietal lobule, bilateral precuneus (adjacent to PC) as well as in visual areas

including superior and middle occipital gyri during Landmark PI compared to Pure PI,

presumably reflecting the higher amount of visual information (see Fig. S2 and Tab. S3).

No significant clusters were found on whole brain level for the other contrasts.

The ROI analyses (see Fig. S3 for overview of masks) showed higher activation of PC

($t_{(35)} = 6.33$, $p_{FWE} < .001$) and right CN ($t_{(35)} = 4.45$, $p_{FWE} = .010$) in Landmark PI compared

to Pure PI (*LPI > PPI*; Fig. 4A), while no significant differences were found for the other

ROIs. Furthermore, no ROI showed differential activation in the contrast between cortisol

and placebo (*CORT > PLA*). Importantly, however, we observed higher activation in the

right pmEC ($t_{(35)} = 2.78$, $p_{FWE} = .045$), right CN ($t_{(35)} = 4.62$, $p_{FWE} = .006$), and left CN ($t_{(35)}$

$= 3.90$, $p_{FWE} = .035$) in the interaction contrast of Landmark PI vs. Pure PI during cortisol

compared to placebo (*[Landmark PI > Pure PI]$_{CORT}$ > [Landmark PI > Pure PI]$_{PLA}$*).

Pairwise post-hoc tests showed that these interactions reflected relatively higher

activation of the right CN under cortisol compared to placebo during Landmark PI ($t_{(35)} =$

$3.43$, $p_{Bonferroni} = .006$, $d = 0.550$) and during Pure PI compared to Landmark PI under

placebo ($t_{(35)} = 2.82$, $p_{Bonferroni} = .032$, $d = 0.291$), as well as a trend for a higher activation

of the right pmEC during Pure PI than Landmark PI under placebo ($t_{(35)} = 2.45$, $p_{Bonferroni}$

$= .077$, $d = 0.366$). No significant pairwise comparisons emerged for left CN (all $t \le 2.29$,

all $p_{Bonferroni} \ge .112$, all $d \le 0.360$; Fig. 4B). These findings indicate a reduced recruitment

of right CN during Landmark PI compared to Pure PI under placebo, but enhanced

recruitment during Landmark PI under cortisol compared to placebo, suggesting a

relatively increased relevance of this region in the presence of landmarks under

pharmacologically induced stress. Conversely, the findings suggest that the presence of

landmarks can reduce recruitment of the right pmEC. No other ROI exhibited higher

activation for this contrast, neither for any of the reverse contrasts.

**Figure 4. Univariate fMRI results.** (A) Significantly higher activation in PC and rNC during LPI
as compared to PPI. (B) Significantly higher differentiation between subtasks under different
treatments or between treatments in different subtasks in right CN, left CN and right pmEC. Error
bars represent SEM. PC: posterior cingulate, CN: caudate nucleus, pmEC: posterior-medial
entorhinal cortex, PPI: pure path integration, LPI: landmark-supported path integration, CORT:
cortisol, PLA: placebo, n.s.: not significant, ** $p < .01$, * $p < .05$, † $p < .10$.

*2.3    Cortisol impairs GLRs, which benefit PI performance specifically over short*
*distances*

To assess GLRs, we used a multivariate approach and focused on right pmEC,

consistent with previous studies (33, 22). Accordingly, due to their sixfold rotational

symmetry, grid cells should show similar activity during movements along directions that

differ by 60° (aligned movements) as compared to those differing by 30° (misaligned

movements; STAR Methods).

We first aimed to confirm the presence of GLRs by comparing activity during aligned vs.

misaligned movements (Fig. 5A) and then tested whether they were affected by cortisol.

Indeed, we found higher similarities between aligned movements (mod(α,60°) = 0°) than

misaligned movements (mod(α,60°) = 30°) across all days and treatments,

demonstrating the presence of GLRs in our study ($t_{(71)}$ = 2.92, $p$ = .002, $d$ = 0.345, Fig.

5B, top-left). Subsequent analyses revealed no main effect of treatment on GLRs ($F_{(1,67)}$

= 0.15, $p$ = .696, $\eta_p^2$ = .002), but we observed a significant treatment x day interaction

($F_{(1,67)}$ = 6.79, $p$ = .011, $\eta_p^2$ = .092; Fig. 5B, top-right). Due to the crossover design, the

interaction cannot easily be interpreted, as it may also include order effects. We thus

decided to examine differences between groups on day one (35). When doing this, post-

hoc tests revealed a significant effect of treatment ($t_{(67)}$ = 2.01, $p$ = .040, $d$ = 0.700),

indicating significantly higher GLRs during PLA than during CORT. Indeed, separate

analyses confirmed the presence of GLRs under placebo ($t_{(18)}$ = 2.38, $p_{Šídák}$ = .029, $d$ =

0.545, Fig. 5B, bottom-right) but not under cortisol treatment ($t_{(16)}$ = 0.20, $p_{Šídák}$ = .665, $d$

= 0.049). Control analyses of GLRs from day one under placebo (Fig. 5B, bottom-right)

confirmed that pattern similarity differed between aligned and misaligned movements

only for 6-fold rotational symmetry, but not for 4-fold, 5-fold, 7-fold, or 8-fold rotational

symmetries (all t ≤ 1.91, all p ≥ .073), and removing movements during same heading

directions (±15° from 0°) preserved (by trend) GLRs ($t_{(18)}$ = 1.57, $p$ = .067, $d$ = 0.36).

Subsequently, we investigated whether GLRs were behaviorally relevant using a linear

mixed model incorporating effects of subtask, path distance, and GLRs. Because we

only found GLRs on day one under placebo, these analyses focused on this condition.

We found similar effects as in the purely behavioral model (main effects of subtask,

incoming distance, age, and the subtask x incoming distance interaction) with the

addition of an interaction between incoming distance and GLRs ($F_{(1,1193)}$ = 10.47, $p$ =

.001, $\eta_p^2$ = .009; Fig. 5C, left). Post-hoc analyses revealed that GLRs are (by trend)

beneficial for performance (lower drop errors) in case of short incoming distances ($t_{(44)}$ =

2.29, $p_{Šídák}$ = .079), while they were detrimental for long incoming distances ($t_{(133)}$ = 2.45,

$p_{Šídák}$ = .046). For the sake of completeness, we repeated this analysis considering data

across days and treatments, which yielded in a more complex four-way interaction

between GLRs, subtask, incoming distance and treatment ($F_{(1,4560)}$ = 5.29, $p$ = .021, $\eta_p^2$

= .001; Fig. 5C, right). This interaction suggested that the effects of GLRs are limited to

environments without spatial cues (pure PI). More specifically, during pure PI under

placebo, GLRs tended to be beneficial for short distances, but detrimental over long

distances, similar to the findings from day 1 under placebo. In contrast, during pure PI

under cortisol, GLRs were beneficial over long distances. However, this last analysis

incorporating all data should be interpreted with caution, as GLRs were only reliably
observed on day 1 under placebo.

**Figure 5. Cortisol effects on GLRs and their behavioral relevance.** (A) Inner numbers
represent angular differences in 360° space, whereas outer numbers show differences in 60°
space (left panel). Higher pattern similarity was expected for angular differences of mod(α,60°) =

0° (rose; α as angular difference between two movement directions) than for angular differences
of mod(α,60°) = 30° (gray). The same result was expected when discarding pattern similarities of
movements for the same heading direction (dark rose). Two exemplary fMRI volumes
representing different movement directions (blue arrows; right panel). In the first volume, the
participant navigates at an angle of 0° with respect to the (arbitrary) reference axis, while
navigating at an angle of 90° in the second volume. This results in an angular difference of 90° in
360° space, corresponding to 30° in 60° space and thus mod(α,60°) = 30° (compare blue arrows
on the left panel). (B) Average pattern similarity over the entire time series of all voxels in the right
pmEC for all directions of aligned (rose) and misaligned (grey) fast runs (top-left panel).
Differences in pattern similarity depending on treatment and day (top-right panel), and average
pattern similarity (bottom-left panel) as well as control analyses (bottom-right panel) only for PLA
treatment on day one. (C) Relevance of GLRs for PI considering day 1 under placebo (left) and
across days and treatments (right). Significant slopes are indicated via asterisks. Min Incoming,
Mean Incoming, and Max Incoming refer to the minimum, mean and maximum incoming
distances, respectively, and were chosen as basis for estimating adjusted means of linear trends.
Error bars and confidence bands represent SEM. GLR: grid-like representation, pmEC: posterior-
medial entorhinal cortex, Pure PI: pure path integration, Landmark PI: landmark-supported path
integration, CORT: cortisol, PLA: placebo, vm: virtual meters, n.s.: not significant, ** $p < .01$, * $p$
$< .05$, † $p < .10$.

*2.4    Exploratory parametric modulation analysis of PI performance*

We further conducted a parametric modulation analysis using a general linear model

(GLM), including trial-wise inverted drop error values as regressor for the combined

outgoing and incoming phases. We did not distinguish between subtasks or treatments

as we were interested in general performance-related neural activity across conditions,

encompassing both whole-brain analysis and ROI analyses. The whole-brain analysis

yielded a set of clusters showing activation modulated by trial-wise drop error, while the

ROI analyses did not reveal any significant modulation of the BOLD signal (all $p_{FDR} >$

.630; Fig. S4 and Tab. S4).

## 3    Discussion

We investigated whether cortisol administration affects PI performance, navigational

patterns in the presence of a landmark, and associated brain activations in healthy young

participants. We found cortisol to impair overall PI performance and to promote a change

in navigational pattern towards navigating further away from the landmark. Because

navigating further away from the landmark was linked to worse performance, this cortisol-

induced change in navigational pattern likely contributed to the observed PI deficits. The

univariate fMRI data analysis indicated a higher activation of PC and right CN when PI

was supported by landmarks, and of right CN in this condition under cortisol. In the

absence of landmarks, when participants could solely rely on optic flow, right pmEC

showed relatively higher activation under placebo compared to cortisol. The multivariate

fMRI data analysis showed the presence of GLRs in the right pmEC under placebo on

day one, but not under cortisol, and while they were (on trend) beneficial for PI

performance over short distances, they were detrimental over long distances.

Our behavioral results showed that cortisol impairs overall PI performance, an effect

mediated by slight impairments of distance and rotation estimations (see Fig. S5). Even

though we expected a detrimental cortisol effect, we assumed it to be more pronounced

in environments with no spatial cues (Pure PI) and in trials with longer incoming

distances, because we had previously found acute stress effects to be moderated by

these variables (31). Cortisol is a relevant mediator of stress effects and known to

influence cognitive processes (8), but because its increase is only one part of the several

biological processes involved in the stress response, effects of cortisol and acute stress

are often not identical and sometimes even opposing (36). An important principle of

glucocorticoid effects is dose-dependency (37), referring to differential effects on

neurons in a concentration-depending manner. Dose-dependency interacts with the local

distribution pattern of MRs and GRs, which exhibit differences in their affinity to cortisol

(5). While MRs show greater affinity and are highly expressed in limbic structures, GRs

possess lower affinity and are more uniformly distributed across the brain, with high

abundance in the EC (30). Thus, high cortisol concentrations like in the current study

(much higher than average acute stress responses) may affect GRs in EC more strongly

and thus impair PI irrespective of moderators like environmental scarcity and path

distance. In another previous study, we did not find a main effect of chronically elevated

cortisol levels on PI performance, but subtle impairments of distance estimations under

conditions of spatial scarcity and long path distance (32). Again, cortisol concentrations

were in a low-to-mid range, potentially being insufficient to occupy enough GRs to impair

PI performance generally. We therefore propose that cortisol affects PI in a

concentration-dependent manner, where lower concentrations lead to subtle effects,

detectable in the absence of spatial cues and over long path distances, and where higher

concentrations induce severe effects, detectable irrespective of environmental properties

or path distance.

The role of landmarks during PI is to stabilize grid cell firing (23, 38), thereby improving

performance in general, an effect that is in accordance with our results. When examining

the navigational pattern during Landmark PI irrespective of performance, however, we

surprisingly found that cortisol led to navigating further away from the landmark. Notably,

this effect may partly explain the cortisol-induced impairment in overall PI performance,

because increased distances to the landmark generally predicted higher drop errors.

Conversely, in a previous study, we had shown that acute stress (on trend) changes the

navigational pattern towards the landmark, and we argued that this change might reflect

stress-induced enhanced use of landmark information, stronger recruitment of the

PC/RSC, or a shift towards striatal processing after acute stress (31). While the fMRI

data of our current study do not support the idea of higher recruitment of PC/RSC, we

indeed found higher activation in right CN under cortisol during Landmark PI compared

to Pure PI, supporting the claim of increased striatal processing. Mechanistically, acute

stress enhances the use of landmark information in the striatum via cortisol binding on

MRs (39), especially in participants showing high cortisol responses. Thus, we would

here expect converging effects of acute stress and cortisol, even though dose-

dependency might again play a relevant role. With high cortisol concentrations, like in

our current study, not only MRs are activated in the striatum, but also GRs, changing the

relative activation of MRs and GRs and thereby altering neuronal processing, an idea

referred to as *MR/GR ratio hypothesis* (40). However, while the potential concomitant

activation of both receptor types in this study might have contributed to the inconsistency

in findings, our task is not specifically designed for investigating landmark-based

strategies and their neuronal correlates, making it less comparable to studies targeting

this phenomenon explicitly. Further research is needed to clarify the role of cortisol

(concentration) on landmark use, both in general and specifically during PI.

The exploratory whole-brain analyses demonstrated higher activation in the left superior

parietal lobule, bilateral precuneus and in visual areas in the presence of landmarks. The

precuneus provides spatial information to HC and EC (41) and its higher activation along

with that of visual areas during Landmark PI presumably reflects the higher availability

of visual information. Additionally, the parametric modulation analysis of the drop error

revealed that these brain regions were also related to PI performance (among others).

The ROI analyses did not reveal any region that is modulated by PI performance, but

showed less deactivation of the PC and higher activation of the right CN during Landmark

PI, supporting the hypothesis of their crucial involvement in landmark processing (42, 38,

43, 22). Furthermore, we did not observe any ROI that shows generally more activation

under cortisol than placebo, but when incorporating subtask, we found effects in right CN

and right pmEC. Right CN showed higher activation under cortisol during Landmark PI,

suggesting that striatal processing might not be generally enhanced under stress, but

specifically when usage of stimulus-response type of strategies (landmark processing in

this case) is possible. Under placebo, the right pmEC showed less deactivation during

Pure PI (on trend), highlighting the relevance of this region in the absence of spatial cues

and suggesting that adding a landmark reduces its involvement. This is in accordance

with the idea that additional brain regions contribute to PI when spatial cues become

available (21, 23).

In the right pmEC, we further demonstrated the existence of overall GLRs, but to control

for potential order effects in our crossover design, we focused on differences between

treatments on day one only. There, we found GLRs under placebo, but not under cortisol

treatment, while control analyses specifically supported the presence of GLRs on day

one only under placebo. This finding is in accordance with the hypothesis that cortisol

affects the EC and impairs GLRs, thereby potentially inducing the several behavioral PI

deficits that we found in a series of studies including the current one (31, 32). More

specifically, because cortisol administration impairs inhibitory transmission in layer II of

the EC (44), we propose that cortisol-induced inhibition of grid cells in the medial EC, an

effect that correlates with the spacing of grid cells, causes behavioral PI deficits. Grid

cells, and the medial EC, are generally related to PI performance, and lesions in the

medial EC impair PI in rats (45, 46), presumably related to impaired distance estimations

(47). Our analyses showed a relationship between GLRs and PI performance that

depended on path distance, suggesting a beneficial role of GLRs for short incoming

distances, but a detrimental role of GLRs over long incoming distances conditions. Based

on previous studies (48, 22, 17), we assumed that GLRs should facilitate performance

specifically in environments with no spatial cues, while our findings suggest that path

distance is the more relevant moderator of GLR effects. In line with our finding, several

lines of research indicate that grid cells might support PI specifically in case of short

distances, including the emergence of grid distortions in larger environments (49), and

the proposition that grid cells might rather provide a narrow and local rather than a

general and global spatial metric (50). Over long incoming distances, grid distortions

might lead to increased error accumulation, making errors larger and more likely to occur,

a common observation in PI tasks, especially when no recalibration or anchoring of grid

cells is possible. However, while this explanation would imply that grid cells are non-

relevant over long distances, it does not cover the observation that they are detrimental,

necessitating further research on this aspect.

A few limitations of this study need to be addressed. First, due to increased power, we

used a within-subject crossover design. This can lead to results, where treatment and

order effects cannot be disentangled, which we accounted for at the price of reducing

sample size for GLR analyses. Also, stress effects in within-subject tasks differ from

those in between-subject designs (51), even though we bypassed this issue by

considering day one only (making it a pseudo between-subject design) for the stress

effect on GLRs. Second, cognitive fatigue can water down stress effects on cognition

(36, 51), and thus, the cognitive battery implemented before the task and the long

duration of the task might have influenced our results, suggesting that cortisol effects on

GLRs and PI could be even stronger than what we report here. Last, our sample

consisted of young healthy men, which is not representative for the entire population,

because gonadal hormones play a major in moderating effects of stress and cortisol (52).

However, we have shown cortisol effects on PI in an older, female sample (32), and thus

we can infer that behavioral effects of stress might be similar between sexes, while the

investigation of neuronal effects is warranted in more representative samples in future

studies.

*3.1    Conclusion*

Our current findings provide essential information about the underlying mechanisms of

stress effects on PI, supporting the hypothesis that they are strongly mediated by effects

of cortisol on the EC. Cortisol impaired overall PI performance in healthy human

participants, while navigational patterns were altered towards navigating further away

from the landmark. The fMRI data further demonstrated that cortisol increases activation

in right CN in the presence of landmarks and impairs GLRs in the right pmEC. The GLRs

correlated with better PI performance particularly over short distances.

465                              **Acknowledgements**

466    We thank Lukas Kunz for developing the experimental task and providing guidance for

467    data acquisition. We also thank Liv Hog, Miriam Seyfried, Leja Kornefell and Leander

468    Fester for their help with participant recruitment and data acquisition. Moreover, we thank

469    Tobias Otto and PHILIPS Germany for technical support.

470                              **Supplemental Information**

471    Document S1. Figures S1-S5 and Tables S1-S5.

**Materials and Methods**

*EXPERIMENTAL MODEL AND STUDY PARTICIPANT DETAILS*

*Participants*

We used a two-day within-subject crossover design and recruited 42 healthy men, two

of which had to be excluded due to technical failures of the joystick (see 2.5) and another

one due to unsuccessful experimental manipulation (no cortisol increase, see 2.5),

leaving a final sample size of $n$ = 39, aged 19 – 34 years (24.18 ± 4.55 years; mean ±

SD) and with a body mass index ranging between 18 and 29 kg/m² (23.95 ± 2.54 kg/m²;

mean ± SD). We recruited participants through online advertisements in social media

networks, mailing lists and in university classes at Ruhr University Bochum. Prior to

testing, participants were instructed about all study procedures and gave written

informed consent. The study was conducted in accordance with the Declaration of

Helsinki as approved by the medical ethics committee of Ruhr University Bochum.

*METHOD DETAILS*

*Experimental design*

Sample size was based on previous studies investigating PI (17), the functional

relevance of GLRs (21, 22), and effects of cortisol administration (e.g., 53). Exclusion

criteria comprised an acute or history of disease (i.e., neurological, psychiatric,

cardiovascular, immunologic), current or history of medical or psychological treatment,

drug use, female sex, previous experience with the PI paradigm (see experimental task),

or any contraindication for participation in MRI studies. We only included men because

of the well-known influence of sex hormones on the secretion of stress hormones (54)

and the related changes in neural functioning (55). All participants had normal or

corrected-to-normal vision and received a compensation of 10€/hour (60-70€ in total) or

course credits.

Participants received two tablets of 10 mg cortisol (hydrocortisone, Hoechst) on one

testing day, and two visually identical placebos on the other testing day in a double-blind

two-day within-subject crossover design. Order of conditions was counterbalanced

across participants. The dosage of 20 mg was chosen in accordance with previous

studies demonstrating effects of cortisol on behavioral and neuronal responses with

similar dosages (56–60). We assessed salivary cortisol using Salivettes (Sarstedt,

Nümbrecht, Germany) collected at several time-points (see Fig. 2). Saliva samples were

stored at -20 °C until assayed. Salivary cortisol concentrations were extracted from the

samples using a time-resolved fluorescence immunoassay (IBL, Hamburg, Germany) at

the local biochemical laboratory of Ruhr University Bochum and are reported in

nanomole per liter (nmol/l). Intra- and inter-assay coefficients of variations were below

6.1%.

*Experimental task*

The Apple Game is a virtual PI task that was implemented by Bierbrauer et al. (22) via

Unreal Engine 4 (Epic Games, version 4.11; see also Fig. 1 & Fig. S1). In this task, PI is

considered as the ability to integrate across several paths and to calculate home-coming

vectors based on visual cues only. A circular arena formed by an endless grassy plain

with a blue sky rendered at infinity, with a diameter of 13,576 virtual meters (vm), built

the environment. Briefly, each trial was composed of three phases. In the "start phase",

participants first moved to a basket and memorized its location (goal location). Then, the

"outgoing phase" followed, in which they navigated to a variable number of trees (1-5),

which appeared consecutively in different locations, until a tree containing a red apple

(retrieval location) was reached. The variable number of trees allowed manipulating the

path distance of the outgoing phase and thus difficulty. Basket and trees disappeared

upon arrival at the respective locations. During the "incoming phase", participants lastly

were asked to take the shortest path from the retrieval location back to the goal location.

They pressed a button when arriving at the presumed location (response location) and

received visual feedback via zero to three stars corresponding to the Euclidean distance

between response location and goal location (drop error; three stars for < 1,600 vm, two

stars for < 3,200 vm, one star for < 6,400 vm). To investigate the influence of spatial cues

on PI, the original version of the task contained three subtasks, of which we used two in

this study. The "Pure PI" subtask was composed only of a grassy plain, forcing

participants to solely rely on visual flow, while the "landmark-supported PI" (Landmark

PI) subtask contained a central lighthouse as spatial cue. Participants first played up to

eight training trials to get familiar with the task, during which a structural MRI scan (~ 6

min) was obtained (see 2.6). Then, a total of 64 experimental trials (32 per subtask)

divided into four blocks of 16 trials were presented, during which functional MRI scans

were assessed (see 2.6). The outgoing phase of the 32 trials of each subtask was

composed of six trials with 1, 2, 4 and 5 trees, respectively, and eight trials with three

trees in randomized order. Before every new trial, a fixation cross was presented with a

variable duration 5 – 7.5 s (randomly distributed).

*Experimental Procedure*

Testing sessions were performed after 1 p.m. to avoid interference with the cortisol

awakening response (61). On day one, upon arrival, participants read study information

and gave written informed consent, before undergoing a cognitive test battery, after

which the pharmacological intervention and a series of questionnaires followed (see Tab.

S5). About 30 minutes after pharmacological intervention, participants were prepared for

the fMRI session, during which the experimental PI task was conducted. Lastly,

participants answered a final set of questionnaires. The testing session on day two

followed one week later, and the procedure was essentially the same, except for the

absence of a cognitive test battery and the respective other pharmacological

intervention. Finally, participants were debriefed and compensated (see Fig. 2).

*Behavioral data*

To extract Apple Game data from computer-generated log-files we used MATLAB

(2021b, The MathWorks Inc., Massachusetts, US), including the Parallel Computing

Toolbox (v6.12) and the CircStat Toolbox (62). Statistical analyses were conducted in R

(63) using the lme4 (64), lmerTest (65) and emmeans (66) packages.

*MRI data*

We acquired MRI data during the whole experimental task at the Bergmannsheil hospital

in Bochum using a 3T Philips Achieva Scanner (Best, The Netherlands) with a 32-

channel head coil. High-resolution whole-brain structural brain scans were acquired

using a T1-weighted sequence with a 1 mm isotropic resolution, an FOV of 240 mm x

240 mm, and 220 transversally oriented slices during a total acquisition time (TA) of six

minutes and three seconds. Blood oxygenation level–dependent (BOLD) contrast

images were registered with a T2*-weighted gradient echoplanar imaging sequence with

2.5 mm isotropic resolution, TR = 2500 ms, TE = 30 ms, FA = 85°, FOV = 96 mm x 96

mm, 45 transversal slices in ascending order without slice gap, and TA = 17.32 ± 2.01

mins (mean ± SD), corresponding to 415.67 ± 48.27 (mean ± SD) volumes. Variation in

TA emerged because the task was self-paced, leading to different durations of runs. In

addition to three dummy scans preceding data acquisition, the first five images of each

session were discarded to allow for signal steady-state transition. We further used a shim

box of 60 mm x 60 mm x 60 mm around the medial temporal lobe to increase temporal

signal-to-noise ratio. The virtual environment was presented to participants via MR-

compatible liquid crystal display goggles (VisuaStim Digital, Resonance Technology Inc.,

Northridge, CA, USA) with a resolution of 800 × 600 pixels, and they navigated within

the virtual environment using an MR-compatible joystick (Nata Technologies, Coquitlam,

Canada).

fMRI data were preprocessed and subsequently analyzed using SPM12 implemented in

MATLAB (2021a, The MathWorks Inc., Massachusetts, US) and nilearn for Python

(Nilearn contributors, 2025). Preprocessing included slice time correction and spatial

realignment. For whole-brain and ROI analysis, fMRI scans were further normalized to

MNI space using parameters from the normalization procedure of the segmented

structural T1 image. Further, we applied spatial smoothing with a 5 mm isotropic

Gaussian kernel to the normalized fMRI data.

We used masks of the left and right HC, left and right CN, and bilateral PC (maximum

probability masks; probability threshold: 0.25; Harvard-Oxford Cortical and Subcortical

Structural Atlases, Harvard Center for Morphometric Analysis;

https://cma.mgh.harvard.edu/). Because the medial EC is especially related to grid-cell

activity in rodents and the human EC consists of structurally and functionally distinct

subparts, where the anterior-lateral EC (alEC) and the posterior-medial EC (pmEC)

represent the homolog of rodent lateral and medial EC, respectively, we focused our

analysis on left and right pmEC using masks created by Maass et al. (67).

*QUANTIFICATION AND STATISTICAL ANALYSIS*

*Behavioral data analysis*

One participant did not show any cortisol increase after cortisol administration and thus

was excluded from statistical analyses, leading to a sample size of $n$ = 39 for behavioral

analyses. We assessed several measures for path distance, PI performance and

navigational pattern during the experimental task. For path distance, two parametrical

measures were assessed: O*utgoing distance* represents the accumulated distance from

the goal to the retrieval location, while *incoming distance* refers to the Euclidean distance

between retrieval and goal location (Fig. 1C). These measures represent different

subcomponents of PI: Outgoing distance is relevant for keeping track of the travelled

path in relation to the starting point (i.e., the later goal location), and incoming distance

for calculating a direct vector to this goal location. Because previous studies indicated

incoming distance to be more closely related to EC activation (68, 42), and the results of

our previous work supported this view (32, 31), we again concentrated on this measure

as a proxy of path distance. Moreover, overall PI performance is represented by the drop

error, i.e., the Euclidean distance between response location and goal location (Fig. 1D).

Lastly, to investigate the specific role of the landmark under heightened cortisol levels,

we assessed two variables only applicable in Landmark PI. The first of these reflects the

distance between the goal and the spatial cue (goal-to-landmark distance), while the

other represents the mean Euclidean distance of the moving participant from the

landmark across all time points of the incoming phase (movement-to-landmark distance).

610              *fMRI data analysis*

We excluded three participants, whose ECs were only partly covered, most likely due to

susceptibility artifacts and other distortions, which are very strong in the medial temporal

lobe (69), leaving a final sample size of $n = 36$ for the fMRI data analysis. In the first

GLM, we modeled start phase, outgoing phase, incoming phase, and feedback

separately for each of the two subtasks and each of the two treatments (16 regressors

in total). The duration of these regressors corresponded to the duration of the respective

phases in each trial. We tested three contrasts (plus their respective reverse contrasts):

"Landmark PI > Pure PI", "CORT > PLA" and their interaction "(Landmark PI > Pure

PI)$_{CORT}$ > (Landmark PI > Pure PI)$_{PLA}$", specifically during the outgoing and incoming

phases, which represent the actual PI task. All regressors were convolved with the

hemodynamic response function before entering the GLM. As nuisance regressors, we

included motion parameters as estimated in the realignment procedure and a high-pass

filter (time constant = 128 s) was implemented. We performed whole-brain analyses on

each contrast. Contrast images from the first-level analysis of each participant were

entered into a GLM. Statistical parametric maps were initially thresholded at a family-

wise error (FWE)–corrected α level of $p < 0.05$ across the whole brain. We considered

clusters significant at $p < 0.05$, FWE-corrected (extent threshold of five voxels). For all

significant clusters, we provide maximum probability tissue labels with MNI coordinates

derived from the Neuromorphometrics atlas as implemented in SPM12 (www.oasis-

brains.org/; http://neuromorphometrics.com/). For ROI analyses, the significance

threshold was set to $p < 0.05$ on voxel level, corrected for multiple testing within each

ROI (FWE-corrected; using the small volume correction option of SPM12).

In a second GLM, we conducted an exploratory parametric modulation analysis of the

drop error, which included one single unmodulated trial regressor as well as one

parametric modulation regressor for trial-wise drop error modeling the combined

outgoing and incoming phases of each trial. The onsets of these regressors

corresponded to the onset of the outgoing phase and the modeled durations

corresponded to the added total durations of the outgoing and incoming phases of each

trial. We did not include separate regressors for each subtask (Landmark PI/Pure PI)

since we were interested in the modulation of neural activity by PI performance across

subtasks in this analysis. Drop error values were symmetrically inverted so that higher

values of the parametric regressor would reflect better task performance. In addition,

drop error values were normalized to range between 0 and 1 and mean-centered. Both

regressors were convolved with the hemodynamic response function before entering the

model. We included motion parameters resulting from realignment of functional images

during preprocessing as nuisance regressors. Additionally, we used a cosine model with

a high-pass frequency of 0.01 Hz to account for slow temporal drifts in the signal. Beta

maps for the contrast of drop error parametric modulation against baseline were

computed for each participant and each session. We performed a whole-brain analysis

on the first-level drop error parametric modulation contrast images. Contrast images from

each participant and each session were entered into a GLM which modeled the average

of the first-level parametric modulation contrast across participants while accounting for

subject-specific effects. Here, statistical parametric maps were thresholded at a false

discovery rate (FDR)-corrected α level of $p < 0.01$ across the whole brain. Clusters were

considered significant at $p < 0.01$, FDR-corrected and extending a minimum cluster size

of ten voxels. For all significant clusters, maximum probability tissue labels are reported

based on the peak voxel's MNI coordinates and segmentations derived from the

Harvard-Oxford atlas as implemented in nilearn. For the ROI analysis of drop error

parametric modulation, average $p$ values were corrected for the number of repeated tests

using FDR-correction at α < 0.01.

To assess GLRs during fMRI, we performed a representational similarity analysis (RSA),

using an approach analogous to previous studies (33, 22). This method assumes that

grid cells in EC, due to their sixfold rotational symmetry, show similar activity patterns

during movements that differ by 60° in angular space. Activity patterns should thus show

higher similarity for movements with an offset of $n*60°$, where $n = \{0, 1, ..., 6\}$, and lower

similarity for movements with an offset of $n*60° + 30°$, where $n = \{0, 1, ..., 5\}$. We refer to

the first condition as "mod($\alpha$,60°) = 0°" and to the second condition as "mod($\alpha$,60°) = 30°"

(where $\alpha$ represents the angular difference between movement directions, and "mod"

denotes the modulo operator; see Fig. 5A). The RSA involved the following steps: (i)

extracting the BOLD signal within the right pmEC ROI from the preprocessed fMRI

volumes without normalization or smoothing; (ii) calculating the mean orientation and

speed for each fMRI volume based on the trajectory data; (iii) excluding volumes where

movement speed was slower than the first tertile; (iv) averaging volumes into bins of 5°

orientation based on the mean orientation; (v) calculating the angular difference of the

movement orientation between the volume bins; and (vi) calculating Fisher z-

transformed Pearson correlations between the volume bins. We then compared the

pattern similarity of pairs of volume bins offset by ±15° from the mod($\alpha$,60°) = 0° condition

(aligned movements) with those offset by ±15° from the mod($\alpha$,60°) = 30 condition

(misaligned movements). Pattern similarity was expected to be higher for aligned

movements than for misaligned movements, thus reflecting GLRs. Two control analyses

were performed. First, to ensure that the increase in pattern similarity was specific to

sixfold symmetry, the same analysis was performed for other types of rotational

symmetry (4-fold, 5-fold, 7-fold and 8-fold). Second, the analysis was also performed

after excluding correlations that were ±15° from 0°, to ensure that increases in pattern

similarity were not primarily driven by movements in the same direction, which could

reflect head direction signals.

*Statistical analysis*

Our first statistical analysis examined the success of pharmacological intervention by

comparing cortisol concentrations using a rANOVA with time and pharmacological

intervention as within-subject factors. Because cortisol concentrations typically exhibit a

right-skewed distribution, we conducted a natural log (ln) transformation to obtain

normally distributed data. In case of violation of the sphericity assumption, we used

Greenhouse-Geisser adjustment (and for the sake of presentation, we rounded the

corrected degrees of freedom to the nearest whole number). Post-hoc pairwise

comparisons were performed using Bonferroni-corrected $t$-tests (or Welch's $t$-tests in

case of unequal variances).

To test the hypothesis postulating differential effects of cortisol depending on subtask

and path distance on PI performance, we then built a linear mixed model with PI

performance (i.e., drop error) on the level of single trials as criterion, and subtask (two

levels: Pure PI vs. Landmark PI), path distance (i.e., incoming distance), and treatment

(two levels: cortisol vs. placebo) as within-subject predictors (Model 1; Tab. S1).

To test the hypothesis concerning the effects of cortisol on the role of the landmark, we

built two further linear mixed models (Models 2A-B; Tab. S1). The first of these aimed at

investigating the role of goal-to-landmark distance and its interaction with cortisol on PI

performance and thus included PI performance on the level of single trials as criterion,

goal-to-landmark distance (only in Landmark PI), and treatment as within-subject

predictors. The second of these models aimed at investigating whether cortisol affected

the employment of navigational strategies (irrespective of performance) and included

movement-to-landmark distance (only in Landmark PI) on the level of single trials as

criterion, and treatment as within-subject predictor.

To evaluate the overall existence of GLRs, we conducted a one-tailed one-sample t-test

comparing activation of EC during aligned vs. misaligned movements against zero

(Model 3A; Tab. S1). To test whether GLRs were related to treatment, we conducted a

linear model with GLRs as criterion, treatment and day as within-subject predictors, and

age as covariate (Model 3B; Tab. S1). We conducted separate t-tests for the control

analyses (other symmetries or correction for heading direction). In the next step, to

examine the behavioral relevance of GLRs, we built a linear model with PI performance

on the level of single trials as criterion, and subtask, path distance, and GLRs as within-

subject predictors (Model 4; Tab. S1).

In all linear mixed models, "subject" was added as random factor, and age, testing day,

and sequence (to control for order effects) as covariates. For all analyses, we centered

within-subject parametric predictors (incoming distance, goal-to-landmark distance) to

the participant's mean and age to the grand mean of all participants (70). For analysis of

fixed effects, we always used type III sum of squares. In case of interactions between

parametric predictors, we discretized all but one parametric predictor and calculated

estimated marginal means ("emmeans" or "adjusted means") or estimated marginal

means of linear trends ("emtrends" or "conditional regression equations") based on the

minimum, mean and maximum values of these discretized predictors. For post-hoc tests

against zero or pairwise comparisons using Fisher's tests, we used Šídák-adjustment

correcting for number of discretized predictors (3), number of subtasks (2), number of

treatments (2), or a combination of those. For an approximation of degrees of freedom,

we used the Kenward-Roger method (and for the sake of presentation, we rounded them

to the nearest whole number). To estimate effect sizes, we used Partial Eta-Squared

($\eta_p^2$) for *F*-tests and Cohen's *d* for *t*-tests. Multicollinearity between predictors was not

problematic (all variance inflation factors < 5). Assumptions of statistical tests were

verified before application and all tests were conducted two-tailed (except for t-tests on

GLRs) at a significance level of $\alpha = 0.05$.

**References**

1. Brunyé TT, Wood MD, Houck LA, Taylor HA. The path more travelled: Time pressure
increases reliance on familiar route-based strategies during navigation. Q J Exp Psychol
(Hove) 2017; 70(8):1439–52.

2. Coughlan G, Laczó J, Hort J, Minihane A-M, Hornberger M. Spatial navigation deficits
- overlooked cognitive marker for preclinical Alzheimer disease? Nat Rev Neurol 2018;
14(8):496–506.

3. Justice NJ. The relationship between stress and Alzheimer's disease. Neurobiol
Stress 2018; 8:127–33.

4. Joëls M, Baram TZ. The neuro-symphony of stress. Nat Rev Neurosci 2009;
10(6):459–66. Available from: URL: https://www.nature.com/articles/nrn2632.

5. Herbert J, Goodyer IM, Grossman AB, Hastings MH, de Kloet ER, Lightman SL et al.
Do corticosteroids damage the brain? J Neuroendocrinol 2006; 18(6):393–411.

6. Kvetňanský R, Sabban EL, Palkovits M. Catecholaminergic systems in stress:
structural and molecular genetic approaches. Physiological Reviews 2009; 89(2):535–
606.

7. Merz CJ, Wolf OT. How stress hormones shape memories of fear and anxiety in
humans. Neuroscience and biobehavioral reviews 2022; 142:104901. Available from:
URL: https://www.sciencedirect.com/science/article/pii/S0149763422003906.

8. Shields GS, Bonner JC, Moons WG. Does cortisol influence core executive functions?
A meta-analysis of acute cortisol administration effects on working memory, inhibition,
and set-shifting. Psychoneuroendocrinology 2015; 58:91–103. Available from: URL:
https://pubmed.ncbi.nlm.nih.gov/25973565/.

9. Shields GS, Sazma MA, McCullough AM, Yonelinas AP. The effects of acute stress
on episodic memory: A meta-analysis and integrative review. Psychol Bull 2017;
143(6):636–75.

10. Wolf OT, Atsak P, Quervain DJ de, Roozendaal B, Wingenfeld K. Stress and
Memory: A Selective Review on Recent Developments in the Understanding of Stress
Hormone Effects on Memory and Their Clinical Relevance. J Neuroendocrinol 2016;
28(8). Available from: URL: https://onlinelibrary.wiley.com/doi/10.1111/jne.12353.

11. Jain D, Jakhalekar IR, Deshmukh SS. Navigational Strategies and Their Neural
Correlates. J Indian Inst Sci 2017; 97(4):511–25.

12. Wang RF. Building a cognitive map by assembling multiple path integration systems.
Psychon Bull Rev 2016; 23(3):692–702.

13. Valerio S, Taube JS. Path integration: how the head direction signal maintains and
corrects spatial orientation. Nat Neurosci 2012; 15(10):1445–53. Available from: URL:
https://www.nature.com/articles/nn.3215.

14. Banino A, Barry C, Uria B, Blundell C, Lillicrap T, Mirowski P et al. Vector-based
navigation using grid-like representations in artificial agents. Nature 2018;
557(7705):429. Available from: URL: https://www.nature.com/articles/s41586-018-0102-
6.pdf.

15. Gil M, Ancau M, Schlesiger MI, Neitz A, Allen K, Marco RJ de et al. Impaired path
integration in mice with disrupted grid cell firing. Nat Neurosci 2018; 21(1):81–91.

16. McNaughton BL, Battaglia FP, Jensen O, Moser EI, Moser M-B. Path integration and
the neural basis of the 'cognitive map'. Nat Rev Neurosci 2006; 7(8):663–78.

17. Stangl M, Achtzehn J, Huber K, Dietrich C, Tempelmann C, Wolbers T.
Compromised Grid-Cell-like Representations in Old Age as a Key Mechanism to Explain
Age-Related Navigational Deficits. Curr Biol 2018; 28(7):1108-1115.e6.

18. Fyhn M, Molden S, Witter MP, Moser EI, Moser M-B. Spatial representation in the
entorhinal cortex. Science 2004; 305(5688):1258–64.

19. Hafting T, Fyhn M, Molden S, Moser M-B, Moser EI. Microstructure of a spatial map
in the entorhinal cortex. Nature 2005; 436(7052):801–6.

20. Moser EI, Moser M-B, McNaughton BL. Spatial representation in the hippocampal
formation: A history. Nat Neurosci 2017; 20(11):1448–64.

21. Kunz L, Navarro Schröder T, Lee H, Montag C, Lachmann B, Sariyska R et al.
Reduced grid-cell-like representations in adults at genetic risk for Alzheimer's disease.
Science 2015; 350(6259):430–3.

22. Bierbrauer A, Kunz L, Gomes CA, Luhmann M, Deuker L, Getzmann S et al.
Unmasking selective path integration deficits in Alzheimer's disease risk carriers. Sci.
Adv. 2020; 6(35):eaba1394.

23. Hardcastle K, Ganguli S, Giocomo LM. Environmental boundaries as an error
correction mechanism for grid cells. Neuron 2015; 86(3):827–39.

24. Doeller CF, King JA, Burgess N. Parallel striatal and hippocampal systems for
landmarks and boundaries in spatial memory. Proc Natl Acad Sci U S A 2008;
105(15):5915–20.

25. Schwabe L, Wolf OT. Stress and multiple memory systems: from 'thinking' to 'doing'.
Trends Cogn Sci 2013; 17(2):60–8.

26. Umegaki H, Zhu W, Nakamura A, Suzuki Y, Takada M, Endo H et al. Involvement of
the entorhinal cortex in the stress response to immobilization, but not to insulin-induced
hypoglycaemia. J Neuroendocrinol 2003; 15(3):237–41.

27. Burtscher J, Copin J-C, Rodrigues J, Kumar ST, Chiki A, Guillot de Suduiraut I et al.
Chronic corticosterone aggravates behavioral and neuronal symptomatology in a mouse
model of alpha-synuclein pathology. Neurobiology of Aging 2019; 83:11–20.

28. Vouimba R-M, Yaniv D, Diamond DM, Richter-Levin G. Effects of inescapable stress
on LTP in the amygdala versus the dentate gyrus of freely behaving rats. Eur J Neurosci
2004; 19(7):1887–94.

29. Yaniv D, Vouimba R-M, Diamond DM, Richter-Levin G. Simultaneous induction of
long-term potentiation in the hippocampus and the amygdala by entorhinal cortex
activation: mechanistic and temporal profiles. Neuroscience 2003; 120(4):1125–35.

30. Sarrieau A, Dussaillant M, Moguilewsky M, Coutable D, Philibert D, Rostène W.
Autoradiographic localization of glucocorticosteroid binding sites in rat brain after in vivo
injection of [3H]RU 28362. Neurosci Lett 1988; 92(1):14–20.

31. Akan O, Bierbrauer A, Axmacher N, Wolf OT. Acute stress impairs visual path
integration. Neurobiol Stress 2023; 26:100561. Available from: URL:
https://www.sciencedirect.com/science/article/pii/S2352289523000498.

32. Akan O, Bierbrauer A, Kunz L, Gajewski PD, Getzmann S, Hengstler JG et al.
Chronic stress is associated with specific path integration deficits. Behav Brain Res
2023; 442:114305. Available from: URL:
https://www.sciencedirect.com/science/article/pii/S0166432823000232.

33. Bellmund JLS, Deuker L, Navarro Schröder T, Doeller CF. Grid-cell representations
in mental simulation. Elife 2016; 5.

34. Doeller CF, Barry C, Burgess N. Evidence for grid cells in a human memory network.
Nature 2010; 463(7281):657–61.

35. Lim C-Y, In J. Considerations for crossover design in clinical study. Korean Journal
of Anesthesiology 2021; 74(4):293–9. Available from: URL:
https://www.ncbi.nlm.nih.gov/pmc/articles/PMC8342834/.

36. Shields GS, Sazma MA, Yonelinas AP. The effects of acute stress on core executive
functions: A meta-analysis and comparison with cortisol. Neurosci Biobehav Rev 2016;
68:651–68.

37. Joëls M, Karst H, Sarabdjitsingh RA. The stressed brain of humans and rodents.
Acta Physiologica 2018; 223(2):e13066. Available from: URL:
https://onlinelibrary.wiley.com/doi/10.1111/apha.13066.

38. Mitchell AS, Czajkowski R, Zhang N, Jeffery KJ, Nelson AJD. Retrosplenial cortex
and its role in spatial cognition. Brain Neurosci Adv 2018; 2:2398212818757098.

39. Vogel S, Klumpers F, Navarro Schröder T, Oplaat KT, Krugers HJ, Oitzl MS et al.
Stress Induces a Shift Towards Striatum-Dependent Stimulus-Response Learning via
the Mineralocorticoid Receptor. Neuropsychopharmacol 2017; 42(6):1262–71. Available
from: URL: https://www.nature.com/articles/npp2016262.

40. de Kloet ER, Oitzl MS, Joëls M. Stress and cognition: are corticosteroids good or bad
guys? Trends Neurosci 1999; 22(10):422–6. Available from: URL:
https://www.sciencedirect.com/science/article/pii/S0166223699014381.

41. Rolls ET, Wirth S. Spatial representations in the primate hippocampus, and their
functions in memory and navigation. Prog Neurobiol 2018; 171:90–113. Available from:
URL: https://www.sciencedirect.com/science/article/pii/S0301008218300625.

42. Epstein RA, Patai EZ, Julian JB, Spiers HJ. The cognitive map in humans: spatial
navigation and beyond. Nat Neurosci 2017; 20(11):1504–13.

43. Chersi F, Burgess N. The Cognitive Architecture of Spatial Navigation: Hippocampal
and Striatal Contributions. Neuron 2015; 88(1):64–77.

44. Hartner JP, Schrader LA. Interaction of Norepinephrine and Glucocorticoids
Modulate Inhibition of Principle Cells of Layer II Medial Entorhinal Cortex in Male Mice.
Front Synaptic Neurosci 2018; 10:3.

45. Parron C, Save E. Evidence for entorhinal and parietal cortices involvement in path
integration in the rat. Exp Brain Res 2004; 159(3):349–59. Available from: URL:
https://link.springer.com/article/10.1007/s00221-004-1960-8.

46. van Cauter T, Camon J, Alvernhe A, Elduayen C, Sargolini F, Save E. Distinct roles
of medial and lateral entorhinal cortex in spatial cognition. Cereb Cortex 2013;
23(2):451–9.                    Available                    from:                    URL:
https://academic.oup.com/cercor/article/23/2/451/286831?login=true.

47. Jacob P-Y, Gordillo-Salas M, Facchini J, Poucet B, Save E, Sargolini F. Medial
entorhinal cortex and medial septum contribute to self-motion-based linear distance
estimation. Brain Struct Funct 2017; 222(6):2727–42. Available from: URL:
https://link.springer.com/article/10.1007/s00429-017-1368-4.

48. Newton C, Pope M, Rua C, Henson RN, Ji Z, Burgess N et al. Entorhinal-based path
integration selectively predicts midlife risk of Alzheimer's disease. Alzheimer's &
Dementia     2024;     20(4):2779–93.     Available     from:     URL:     https://alz-
journals.onlinelibrary.wiley.com/doi/full/10.1002/alz.13733.

49. Krupic J, Bauza M, Burton S, Barry C, O'Keefe J. Grid cell symmetry is shaped by
environmental geometry. Nature 2015; 518(7538):232–5. Available from: URL:
https://www.nature.com/articles/nature14153.

50. Ginosar G, Aljadeff J, Las L, Derdikman D, Ulanovsky N. Are grid cells used for
navigation? On local metrics, subjective spaces, and black holes. Neuron 2023;
111(12):1858–75.                    Available                    from:                    URL:
https://www.sciencedirect.com/science/article/pii/S0896627323002234.

51. Shields GS. Stress and cognition: A user's guide to designing and interpreting
studies. Psychoneuroendocrinology 2020; 112:104475. Available from: URL:
https://www.sciencedirect.com/science/article/pii/S0306453019304251.

52. Jentsch VL, Pötzl L, Wolf OT, Merz CJ. Hormonal contraceptive usage influences
stress hormone effects on cognition and emotion. Frontiers in Neuroendocrinology 2022;
67:101012.                    Available                    from:                    URL:
https://www.sciencedirect.com/science/article/pii/S0091302222000358.

53. Kinner VL, Wolf OT, Merz CJ. Cortisol alters reward processing in the human brain.
Horm Behav 2016; 84:75–83.

54. Kirschbaum C, Kudielka BM, Gaab J, Schommer NC, Hellhammer DH. Impact of
gender, menstrual cycle phase, and oral contraceptives on the activity of the
hypothalamus-pituitary-adrenal axis. Psychosomatic Medicine 1999; 61(2):154–62.

55. Goldfarb EV, Seo D, Sinha R. Sex differences in neural stress responses and
correlation with subjective stress and stress regulation. Neurobiol Stress 2019;
11:100177.

56. Pan D, Jentsch VL, Langer K, Hagedorn B, Höffken O, Wolf OT et al. What a
difference timing makes: Cortisol effects on neural underpinnings of emotion regulation.
Neurobiol     Stress     2023;     25:100544.     Available     from:     URL:
https://www.sciencedirect.com/science/article/pii/S2352289523000322.

57. Langer K, Jentsch VL, Wolf OT. Cortisol promotes the cognitive regulation of high
intensive emotions independent of timing. European Journal of Neuroscience 2022;

55(9-10):2684–98.                Available                from:                URL:
https://onlinelibrary.wiley.com/doi/10.1111/ejn.15182.

58. Hagedorn B, Wolf OT, Merz CJ. Cortisol before extinction generalization alters its
neural correlates during retrieval. Psychoneuroendocrinology 2022; 136:105607.
Available                                from:                                URL:
https://www.sciencedirect.com/science/article/pii/S0306453021004819.

59. Henckens MJAG, van Wingen GA, Joëls M, Fernández G. Time-dependent effects
of corticosteroids on human amygdala processing. J. Neurosci. 2010; 30(38):12725–32.
Available from: URL: https://www.jneurosci.org/content/30/38/12725.

60. Henckens MJAG, van Wingen GA, Joëls M, Fernández G. Time-dependent
corticosteroid modulation of prefrontal working memory processing. Proceedings of the
National Academy of Sciences of the United States of America 2011; 108(14):5801–6.

61. Clow A, Thorn EL, Evans P, Hucklebridge F. The awakening cortisol response:
methodological issues and significance. Stress 2004; 7(1):29–37.

62. Berens P. CircStat: A MATLAB Toolbox for Circular Statistics. J. Stat. Soft. 2009;
31(10):1–21.

63. R: A language and environment for statistical computing.: R Foundation for Statistical
Computing, Vienna, Austria.; 2021. Available from: URL: https://www.R-project.org/.

64. Bates D, Mächler M, Bolker B, Walker S. Fitting Linear Mixed-Effects Models Using
lme4. J. Stat. Soft. 2015; 67(1):1–48.

65. Kuznetsova A, Brockhoff PB, Christensen RHB. lmerTest Package: Tests in Linear
Mixed Effects Models. J. Stat. Soft. 2017; 82(13):1–26.

66. emmeans. Estimated Marginal Means, aka least square means. Version 1.7.2; 2022.
Available from: URL: https://CRAN.R-project.org/package=emmeans.

67. Maass A, Berron D, Libby LA, Ranganath C, Düzel E. Functional subregions of the
human entorhinal cortex. Elife 2015; 4.

68. Howard LR, Javadi AH, Yu Y, Mill RD, Morrison LC, Knight R et al. The hippocampus
and entorhinal cortex encode the path and Euclidean distances to goals during
navigation. Curr Biol 2014; 24(12):1331–40.

69. Olman CA, Davachi L, Inati S. Distortion and signal loss in medial temporal lobe.
PLOS ONE 2009; 4(12):e8160.

70. Enders CK, Tofighi D. Centering predictor variables in cross-sectional multilevel
models: a new look at an old issue. Psychol Methods 2007; 12(2):121–38.