Paramyxoviruses in Old World fruit bats (Pteropodidae): an open database and synthesis of sampling effort, viral positivity, and coevolution

Short title: Paramyxovirus sampling and prevalence in pteropodid bats

Maya M. Juman[1]*, Olivier Restif[1], Daniel J. Becker[2]

[1]Department of Veterinary Medicine, University of Cambridge, UK

[2]School of Biological Sciences, University of Oklahoma, Norman, OK, USA

*Corresponding author
Email: mmj38@cam.ac.uk

Note: This study involves no new data collection or live participants and therefore does not require an Ethics Statement.

Abstract

Over the past 25 years, research interest in the links between Old World fruit bats (family

Pteropodidae) and paramyxoviruses has driven a rapid proliferation of surveillance studies in this

system. We performed a systematic review and data synthesis of all published paramyxovirus

studies of wild pteropodids. Here, we present an open, static, PRISMA-compliant database called

pteroparamyxo, which includes 1,482 records of prevalence or seroprevalence from 119 studies

published between 1971 and 2023. Using this dataset, we examined biases in sampling effort,

predictors of viral positivity, tissue tropism, and pteropodid–paramyxovirus coevolution. We

found significant spatial and taxonomic bias in sampling effort, largely driven by

overrepresentation of *Pteropus*–henipavirus systems; in particular, African bats were

undersampled for paramyxoviruses relative to Asian and Oceanian taxa. On the viral side,

henipaviruses (specifically Hendra, Nipah, and Cedar viruses) were overrepresented relative to

pararubulaviruses. We also identified substantial non-random variability in paramyxovirus

prevalence and seroprevalence. Spleens and pooled fecal samples from pteropodids yielded the

highest PCR positivity, while samples from Oceania tended to have slightly higher

seroprevalence than those from Asia and Africa. Collection year was not a significant predictor

of positivity, suggesting limited overall change in paramyxovirus PCR prevalence or

seroprevalence over the last several decades. Finally, we found weak evidence of pteropodid–

paramyxovirus coevolution, supporting relatively frequent host-switching within this system.

Our study highlights critical sampling gaps to address in future surveillance studies and provides

preliminary evidence for sample and tissue types to prioritize in field- and museum-based

sampling.

Author Summary

Paramyxoviruses are a family of viruses that include the human measles and mumps viruses as well as emerging zoonoses like Hendra and Nipah henipaviruses. These henipaviruses spill over from wild fruit bats to domestic animals and humans in Australia and Asia, causing high fatality rates with no known human treatments or vaccines. The occurrence and prevalence of paramyxoviruses across the entire fruit bat family (Pteropodidae) has never been formally investigated, as published detection attempts have not yet been consolidated into a database. We conducted a systematic review of the existing literature on pteropodid–paramyxovirus sampling efforts and compiled an open database for researchers to reference and use. We then leveraged this novel database for a suite of analyses and demonstrated that Australian and Asian fruit bats are disproportionately sampled for paramyxoviruses while African species are understudied, despite the fact that African samples have slightly higher rates of paramyxovirus detection. We identified sample types that may be particularly likely to yield viral detections, a finding which can help guide future survey efforts using field studies as well as museum collections. We also demonstrate that paramyxoviruses appear liable to switch host species within Pteropodidae, challenging earlier hypotheses that these viruses are host specific.

Introduction

Anthropogenic change is driving increased viral transmission across species boundaries, including spillover of zoonotic viruses from wildlife to humans [1]. One particular viral family of interest is the *Paramyxoviridae*, negative-strand RNA viruses identified across vertebrate groups. Paramyxoviruses cause particularly high disease burdens in mammals and birds, with varying degrees of host specificity [2]. For example, the subfamily *Orthoparamyxovirinae* contains the genera *Morbillivirus* (e.g., host-specific measles virus and broadly infecting canine

distemper virus) and *Henipavirus* (e.g., zoonotic bat-borne Hendra and Nipah viruses, which

cause fatal encephalitis in horses and humans) (3). The subfamily *Rubulavirinae* contains the

genera *Orthorubulavirus* (e.g., mumps and human parainfluenza viruses) and *Pararubulavirus*

(e.g., Tioman virus, Menangle virus, and other recently isolated bat-borne viruses) (3). To

understand the host range, geographic range, and potential spillover risks of paramyxoviruses, a

more comprehensive picture of host–paramyxovirus relationships is critical.

Bats are known to host a variety of paramyxoviruses and have been suggested as the

ancestral hosts of *Paramyxoviridae* (4). In particular, Old World fruit bats (family Pteropodidae)

have been implicated as reservoir hosts of two highly pathogenic henipaviruses and other

potentially zoonotic paramyxoviruses. This ecologically and morphologically diverse bat family

of roughly 200 species is widespread across Asia, Oceania, and Africa (5). Immunity-related

genes in pteropodids have evolved at higher rates relative to other chiropteran families, and some

pteropodid-specific mutations are associated with decreased inflammatory responses to

pathogens (6). A balance of viral tolerance and immune defense strategies may explain these fruit

bats' apparent ability to host viruses that are highly pathogenic in other mammals (7).

Henipaviruses, including Hendra virus (8), Nipah virus (9), Ghanaian bat henipavirus (10), and

Cedar virus (11), have been detected in pteropodids. Novel pararubulaviruses have also been

isolated from pteropodids (12–15). Some studies have noted apparent clustering of related

paramyxoviruses in certain pteropodid species, raising the possibility of host specificity and

host-virus coevolution (16), though this has not been quantitatively tested. This growing body of

evidence linking pteropodids to paramyxoviruses has also prompted further research on the

ecological links between these bats and their viruses (17–20), which have implications for

downstream spillover risk (21,22).

Pteropodids are largely frugivorous and nectarivorous, and loss of native habitat has

driven many of these species to seek food sources in urban areas (e.g., public and private

gardens), increasing the likelihood of human–bat interactions (5). Climate and land-use change

have been linked to the spillover of henipaviruses from pteropodids (22,23), and the geographic

(and possibly host) ranges of paramyxoviruses appear to be expanding in certain cases—for

example, the recent Nipah virus spillovers and associated fatalities in South India (24,25). The

resulting interest in henipaviruses may have created a bias toward studying *Pteropus* bats in

Australia and South and Southeast Asia (23). Sampling disparities such as these, coupled with a

lack of consolidated data, prevent a deeper understanding of global changes in viral occurrence,

distributions, and spillover rates over time (26). Large, taxonomically broad host–virus

association databases partially address this issue, but these presence-only databases lack negative

results by nature and also often lack fine-grained metadata. While taxonomically specific

databases exist for other viral families like coronaviruses (27), interactions between pteropodids

and the broader *Paramyxoviridae* have not been comprehensively compiled, despite the fact that

these bats have been heavily sampled for henipaviruses and other paramyxoviruses (28–30). The

absence of such a synthesis limits the identification of patterns in viral prevalence, which is

critical for spillover prediction and mitigation (31).

To address these gaps and enable formal synthesis of pteropodid–paramyxovirus

associations, we built a standardized, PRISMA-compliant database of all published detection

attempts of paramyxoviruses in wild pteropodid bats, including metadata on viral taxonomy, host

taxonomy, sampling and detection methodologies, seasonality, location, and date, as well as

including data on negative results. We then used the resulting dataset to perform a series of

phylogenetic comparative analyses and formal meta-analyses to better understand pteropodid–

paramyxovirus relationships and the current state of the associated literature. To this end, we

investigated: (1) geographic and taxonomic (both host and viral) sampling gaps in this system;

(2) key predictors of paramyxovirus prevalence and seroprevalence in pteropodids, including

potential tissue tropism; and (3) the degree of pteropodid–paramyxovirus coevolution or host-

switching.

Results

Overview of dataset

Our final database consists of 1,482 individual records from 119 studies of

paramyxoviruses in pteropodids published between 1971 and 2023 (Supplementary Data 1).

Each record constitutes a detection attempt associated with a unique combination of the

following columns: host identification (at the finest taxonomic resolution available, typically

species), virus identification (at the finest taxonomic resolution available, typically genus or

species), detection type (single or pooled), study type (cross-sectional, longitudinal, or both),

general detection method (PCR, serology, or viral isolation), specific detection method (i.e., type

of PCR or serological assay), gene or antigen target, sample type (e.g., tissue, swab, blood,

urine), type of tissue tested, country, state/region, site (with coordinates if reported), sampling

year, and sampling month/season. Each record also includes the number of samples tested, the

number of positive samples, and the reported prevalence.

Only three of the 119 studies were published before the year 2000, reflecting the rapid

increase in interest in pteropodid–henipavirus relationships after the discovery of Hendra and

Nipah virus in the 1990s (Fig. 1). Of the total records, 66% involve PCR detection attempts, 27%

involve serological detection attempts, and 7% involve viral isolation attempts. Over 67% of

records focus on the viral subfamily *Orthoparamyxovirinae*, and 66% of records involve

henipaviruses specifically (the remaining 1% reflected "henipa-like" orthoparamyxoviruses).

Another 11% involve the viral subfamily *Rubulavirinae*, with the remaining records attempting

to detect any paramyxovirus (typically with PCR primers targeting the L gene). Detections of

unclassified sequences comprise under 5% of all records. Fine-grained spatial studies of

paramyxovirus prevalence are limited by available spatial metadata; only 21% of records have

associated latitude and longitude listed in the original studies. However, 68% include a specific

site (or multiple sites), 89% include an administrative area within a country (e.g., state, province,

or district), and 99% name a specific country, allowing for broader studies of geographic biases

in sampling effort (and potential georeferencing for future studies). The most common sample

types in our dataset are blood (29%), urine (27%), swabs (22%), and tissue (16%). Detailed

breakdowns of sampling effort and viral positivity by tissue type, detection method, and viral

subfamily are included in Supplementary Tables 1 and 2.

Fig. 1. Sampling effort of paramyxoviruses in Pteropodidae across time. (A) Studies by year

of publication. (B) Samples by year of publication (as collection year is often reported as a range

of time). The vertical dashed red and blue lines indicate the first isolations of Hendra and Nipah

virus, respectively, from *Pteropus* bats (8,9).

Spatial sampling biases

Our dataset includes studies in all three regions within the Pteropodidae range (Africa,

Asia, and Oceania) across 33 countries/territories (Fig. 2; Supplementary Table 3). However,

sampling effort has been highly heterogeneous across countries, with the number of studies per

country ranging from one to 29 (Fig. 2a) and the number of samples per country ranging from 21

(Gabon) to 58,590 (Australia) (Fig. 2b). In particular, large geographic sampling gaps are found

in central and southern Africa (Fig. 2). A generalized linear model (GLM) of binary sampling

effort and subsequent post-hoc comparisons did not reveal any statistically significant variation

across the three regions ($\chi^2 = 4.66$, $p = 0.10$, $R^2 = 0.03$) (Table 1a). However, the intensity of

sampling did vary across regions, in terms of both the number of studies per country ($\chi^2 = 26.23$,

$p < 0.0001$, $R^2 = 0.11$) and the number of samples per country ($\chi^2 = 147,816$, $p < 0.0001$, $R^2 =$

0.38). Specifically, there have been more pteropodid–paramyxovirus studies and samples from

Oceania than from Asia or Africa (Table 1b,c), a difference likely driven by an ongoing research

focus on the *Pteropus*–Hendra virus system in Australia. There have also been more samples

collected from Asia than from Africa (Table 1c).

Fig. 2. Geographic distribution of paramyxovirus sampling effort in pteropodids. The

aggregated Pteropodidae IUCN distribution is outlined in blue. Note that in the UN Geoscheme,

Saudi Arabia and Indonesia are classified as Asia. (A) Number of studies per country (ranging

from one to 29). (B) Number of samples tested per country (ranging from 21 to 58,590,

presented on a log scale).

Table 1. Post-hoc pairwise comparisons from GLMs of sampling effort by region. (A) binary sampling; (B) number of studies; and (C) number of samples across countries as a function of general geographic region. The *p*-values are adjusted for multiple comparisons using a Bonferroni correction. Confidence intervals are back-transformed from the log scale.

Comparison	Odds ratio (95% confidence interval)	z ratio	*p* value
A. Binary sampling effort (binomial response)			
Africa / Asia	1.39 (0.49-3.94)	0.76	1.00
Africa / Oceania	3.71 (0.74-18.54)	1.95	0.15
Asia / Oceania	2.67 (0.51-14.02)	1.42	0.47
B. Number of studies (Poisson response)			
Africa / Asia	0.69 (0.44-1.07)	-2.02	0.13
Africa / Oceania	0.30 (0.18-0.51)	-5.51	**<0.0001**
Asia / Oceania	0.44 (0.26-0.74)	-3.77	**0.0005**
C. Number of samples (Poisson response)			
Africa / Asia	0.29 (0.28-0.29)	-151.60	**<0.0001**
Africa / Oceania	0.06 (0.06-0.06)	-355.39	**<0.0001**
Asia / Oceania	0.20 (0.20-0.21)	-261.79	**<0.0001**

Taxonomic sampling biases

Pteropodids sampled for paramyxoviruses include 58 of the 194 species and 23 of the 43 genera in the family. Using a phylogeny of all the Pteropodidae (32), we found intermediate phylogenetic signal in binary sampling effort ($D = 0.85$), departing from both phylogenetic randomness ($p = 0.03$) and Brownian motion models of evolution ($p = 0$). We then applied phylogenetic factorization, a graph-partitioning algorithm (33), which identified one clade with a disproportionately high fraction of sampled species, including *Rousettus amplexicaudatus*, *R. spinalatus*, *R. leschenaultii*, *R. obliviosus*, *R. madagascariensis*, *R. aegyptiacus*, *R. lanosus*, and *R. celebensis*, as well as the genera *Epomops*, *Epomophorus*, *Micropteropus*, *Nanonycteris*, *Hypsignathus*, *Megaloglossus*, and *Myonycteris* (Fig. 3a, Table 2a). Among sampled species, the number of studies per species range from one to 24 and was randomly distributed across the host

phylogeny (Pagel's $\lambda = 0.00$), departing from Brownian motion clustering ($p < 0.0001$) but not

from phylogenetic randomness ($p = 1$). *Rousettus leschenaultii*, *R. obliviosus*, *R.*

madagascariensis, and *R. aegyptiacus* are overrepresented relative to other taxa, as have been

many species of *Pteropus*; however, *P. chrysoproctus*, *P. tonganus*, *P. neohibernicus*, and *P.*

temminckii have been significantly understudied (Fig. 3b, Table 2b). The number of samples per

species range from one to 25,047 and showed moderate phylogenetic signal ($\lambda = 0.59$), departing

from both Brownian motion clustering ($p < 0.0001$) and phylogenetic randomness ($p = 0.01$).

The single identified oversampled clade constitutes 16 *Pteropus* species, including those heavily

sampled for Hendra and Nipah viruses (e.g., *P. giganteus*, *P. lylei*, *P. poliocephalus*, *P. alecto*)

(Fig. 3c, Table 2c).

On the viral side, 16 of the 78 recognized species in the family *Paramyxoviridae* (3) have

been detected in Pteropodidae, comprising three genera: *Henipavirus*, *Orthorubulavirus*, and

Pararubulavirus. We found strong, highly conserved phylogenetic signal in binary sampling

effort ($D = -0.90$), departing from phylogenetic randomness ($p = 0$) but not from Brownian

motion models of evolution ($p = 0.99$). Phylogenetic factorization did not identify any viral

clades differing significantly in the fraction of detected species (Fig. 3d, Table 2d). Among

detected viruses, the number of studies per virus range from one to 55 and displayed strong

phylogenetic signal ($\lambda = 0.93$), departing from phylogenetic randomness ($p = 0.04$) but not from

Brownian motion ($p = 0.60$). Hendra, Nipah, and Cedar henipaviruses have been overstudied,

while Tioman, Teviot, and Menangle pararubulaviruses have been understudied (Fig. 3e, Table

2e). The number of samples per virus range from nine to 52,006 samples and also show strong

phylogenetic signal ($\lambda = 1.00$), departing from phylogenetic randomness ($p = 0.04$) but not from

Brownian motion ($p = 1.00$). Again, Hendra (52,006 samples), Nipah (46,089 samples), and

Cedar (2,139 samples) henipaviruses are overrepresented relative to all other detected

paramyxoviruses (Fig. 3f, Table 2f).

Fig. 3. Host and viral taxonomic biases in pteropodid–paramyxovirus sampling effort. (A)
Whether or not a bat species has been sampled for paramyxoviruses, across all Pteropodidae
(black dots indicate species that have been sampled); (B) number of studies per sampled bat

species; (C) logged number of samples per sampled bat species; (D) whether or not a viral

species has been detected in Pteropodidae (black dots indicate viruses that have been detected);

(E) number of studies screening for each detected viral species; (F) logged number of samples

screened for each detected viral species. Red and blue shading indicates clades with greater or

lesser sampling effort, respectively, in comparison to all other taxa (as identified by phylogenetic

factorization), with clade numbers corresponding to Table 2. In plots (B), (C), (E), and (F),

segment length represents relative sampling effort.

Table 2. Bat and virus clades differing significantly from the rest of their respective trees in terms of different measures of sampling effort, as identified by phylogenetic factorization. (A) Whether or not a bat species has been sampled for paramyxoviruses, across all Pteropodidae (0: not sampled; 1: sampled); (B) number of studies per sampled bat species; (C) logged number of samples per sampled bat species; (D) whether or not a viral species has been detected in Pteropodidae (no significant clades identified); (E) number of studies screening for each detected viral species; (F) logged number of samples screened for each detected viral species (Fig. 3). We report all significant clades detected through phylogenetic factorization (identified with Holm's sequentially rejective test at a family-wise error rate of 5%), the taxa and number of species within those clades, and the mean sampling effort for the clade compared to the rest of the tree. Bat taxonomy follows Upham et al. 2019 and viral taxonomy follows Rima et al. 2019.

Clade	Taxa	Number of species	Clade mean	Other mean
A) Binary sampling of Pteropodidae				
A1	*Rousettus amplexicaudatus, Rousettus spinalatus, Rousettus leschenaultii, Rousettus obliviosus, Rousettus madagascariensis, Rousettus aegyptiacus, Epomops, Epomophorus, Micropteropus, Nanonycteris, Hypsignathus, Megaloglossus, Myonycteris, Rousettus lanosus, Rousettus celebensis*	29	0.62	0.24
B) Number of studies of sampled pteropodids				
B1	*Pteropus chrysoproctus, Pteropus scapulatus, Pteropus vampyrus, Pteropus giganteus, Pteropus lylei, Pteropus seychellensis, Pteropus rufus, Pteropus poliocephalus, Pteropus hypomelanus, Pteropus conspicillatus, Pteropus alecto, Pteropus tonganus, Pteropus neohibernicus, Pteropus melanotus, Pteropus griseus, Pteropus temminckii*	16	8.44	3.38
B2	*Pteropus hypomelanus, Pteropus conspicillatus, Pteropus alecto*	3	15.67	4.18
B3	*Pteropus chrysoproctus, Pteropus tonganus, Pteropus neohibernicus, Pteropus temminckii*	4	1.00	5.06
B4	*Rousettus leschenaultii, Rousettus obliviosus, Rousettus madagascariensis, Rousettus aegyptiacus*	4	7.25	4.59
C) Logged number of samples of sampled pteropodids				
C1	*Pteropus chrysoproctus, Pteropus scapulatus, Pteropus vampyrus, Pteropus giganteus, Pteropus lylei, Pteropus seychellensis, Pteropus rufus, Pteropus poliocephalus, Pteropus hypomelanus, Pteropus conspicillatus, Pteropus alecto, Pteropus tonganus, Pteropus neohibernicus, Pteropus melanotus, Pteropus griseus, Pteropus temminckii*	16	6.59	3.91
D) Binary detection of paramyxoviruses in Pteropodidae				
E) Number of studies of detected paramyxoviruses				
E1	*Hendra henipavirus, Nipah henipavirus, Cedar henipavirus*	3	31	2.31
E2	*Tioman pararubulavirus, Teviot pararubulavirus, Menangle pararubulavirus*	3	6	8.08
F) Logged number of samples of detected paramyxoviruses				
F1	*Hendra henipavirus, Nipah henipavirus, Cedar henipavirus*	3	9.76	5.82

 Heterogeneity and predictors of paramyxovirus positivity

We fitted two initial, intercept-only phylogenetic meta-analysis models (one for PCR

prevalence and one for seroprevalence) that included random effects of within- and between-

study variation, host genus, genus-level phylogeny, and country (32,34). We observed significant

heterogeneity among positivity estimates in both the PCR dataset ($I^2 = 87.8\%$, $Q = 3,793.41$,

$p < 0.0001$) and the serological dataset ($I^2 = 96.5\%$, $Q = 9,770.71$, $p < 0.0001$). This heterogeneity

was mostly explained by between-study (65.0% PCR; 40.0% serology) and within-study (14.1%

PCR; 26.6% serology) effects. Bat phylogeny explained 0% of the heterogeneity in the PCR

dataset but 28.1% in the serological dataset, and other genus effects accounted for 0% of the

heterogeneity in both datasets. Country effects accounted for 8.7% and 1.8% of the heterogeneity

in the PCR and serological datasets, respectively. None of the fixed effects included in our

models were significantly associated with paramyxovirus prevalence or seroprevalence in

Pteropodidae, with all confidence intervals crossing zero and no significant post-hoc pairwise

comparisons; this may be due to the limited sample size of these separate datasets (particularly

serology; $n = 313$). However, in the PCR model, serum, feces, tissue, and urine appeared to have

slightly higher prevalence than the intercept (blood), and African and Asian samples had slightly

higher prevalence than the intercept (Oceania). Cross-sectional studies and longitudinal studies

appear to yield similar PCR prevalence rates. In the serology model, African and Asian samples

had slightly lower prevalence than samples from Oceania. In both models, Eidolinae,

Pteropodinae, and Rousettinae subfamilies had slightly higher viral detection than the intercept

(Cynopterinae), as did Macroglossusinae in the PCR model (Fig. 4).

Fig. 4. Phylogenetic meta-analysis model coefficients and 95% confidence intervals,

estimated with REML for each dataset. Colours indicate the variables included in each model.

Transparent confidence intervals indicate non-significant effects that cross zero (the vertical

dashed line). The intercept contains the following reference levels: cross-sectional (study type);

blood (sample type); region (Oceania); and Cynopterinae (host subfamily). Sample sizes are 786

PCR prevalence estimates and 313 seroprevalence estimates.

Because tissue type had to be excluded from the phylogenetic meta-analysis models due

to collinearity with other predictors, we separately investigated paramyxovirus tissue tropism in

Pteropodidae by calculating PCR prevalence for each tissue type. We found that PCR prevalence

was highest in spleen samples (6.9%), followed by pooled tissue samples from multiple organs

(6.5%), pooled fecal samples (6.5%), pooled swabs (5.4%), and pooled urine samples (4.2%)

(Fig. 5). All other tissue or sample types yielded around or under 2% prevalence.

Fig. 5. PCR prevalence of paramyxoviruses in different pteropodid tissues. Plotted by

relative sampling effort (represented by the log base 10 of the number of samples screened per

tissue type).

Cophylogenetic analysis

Lastly, we tested the dependence of the viral phylogeny on the host phylogeny using the

Procrustean Approach to Cophylogeny (PACo) (35). This matrix included 23 known PCR or

viral isolation links between 12 pteropodid species and 13 ICTV-ratified paramyxovirus species

(Fig. 6). Only three viral species (*Hendra henipavirus*, *Nipah henipavirus*, and *Tioman*

pararubulavirus) have been identified in more than one bat species, while over half of the

included bat species (7/12) have links to more than one virus (Fig. 6). When compared to 5,000

randomly permuted host–virus matrices, we found no statistically significant congruence

between the two phylogenies in our dataset ($m^2_{XY} = 9{,}827.11$, $p = 0.69$). Failure to reject the null

hypothesis indicates a high degree of host-switching by paramyxoviruses within Pteropodidae.

However, the links between *Pteropus* species and henipaviruses and some pararubulaviruses

(Teviot, Tioman, Menangle) had lower residuals and therefore demonstrated more support for

host–virus coevolution, as did the links between *Rousettus leschenaultii* and Tuhoko

pararubulaviruses (Fig. 6, Supplementary Table 4).

Fig. 6. Tanglegram indicating known links (PCR detections or viral isolations) between pteropodid species and ICTV-recognized paramyxovirus species. Darker links signify smaller residuals from PACo (suggesting more support for coevolution), while lighter links signify higher residuals (more support for host-switching) (Supplementary Table 4). The phylogeny on the left indicates evolutionary relationships among pteropodid species in this dataset, with *Eidolon helvum* and the *Pteropus* genus forming a sister clade to *Rousettus* species. The phylogeny on the right indicates evolutionary relationships among viral species in this dataset, with henipaviruses and pararubulaviruses forming two distinct clades.

Discussion

   The isolation of Hendra and Nipah viruses from flying foxes in the early 2000s sparked widespread research interest in the relationship between pteropodid bats and paramyxoviruses

(8,9). We sought to aggregate all published records of pteropodid–paramyxovirus associations

and prevalence into a PRISMA-compliant, open, static database (Supplementary Fig. 1). Our

database, named pteroparamyxo, includes 1,482 records of viral detection attempts from 119

studies published between 1971 and 2023, which allows for formal synthesis of sampling and

positivity across space, time, and host/viral taxonomy. This consolidation is critical for

addressing questions about changes in viral dynamics (26). Our focus on gathering detailed

metadata for a viral and host family-specific database will enable robust downstream models to

be trained on this taxonomic subset, an approach with higher predictive capacity than broader

models (36).

Spatial and taxonomic analyses of this dataset confirm suspected trends and biases in

sampling effort. Pteropodids in Australia and, to a lesser degree, South Asia (particularly India

and Bangladesh) have been heavily sampled for paramyxoviruses, in terms of both number of

studies and number of samples collected (Fig. 2). These trends are likely driven by studies of

Hendra and Nipah virus spillover events and high numbers of under-roost urine samples

collected following outbreaks (37,38). Much of Africa and Asia remains understudied for

paramyxoviruses in fruit bats, despite evidence of henipavirus spillover into human populations

in Cameroon, for example (39), a country which remains undersampled in our dataset (Fig. 2).

This sampling gap is also notable in light of the fact that African samples had slightly higher

PCR prevalence than samples from Oceania, and some African bat subfamilies (Eidolinae and

Rousettinae) had slightly higher PCR prevalence and seroprevalence than other subfamilies (Fig.

4). Taken together, this suggests missed opportunities for additional detection of

paramyxoviruses in African pteropodids (Fig. 4).

Similarly, taxonomic sampling biases abound on both the host and viral side. In terms of

binary sampling effort, a clade containing some *Rousettus* species and closely related genera

(*Epomops*, *Epomophorus*, *Micropteropus*, *Nanonycteris*, *Hypsignathus*, *Megaloglossus*,

Myonycteris) has a disproportionately high fraction of species sampled for paramyxoviruses.

Among sampled taxa, some species in the highly studied genera *Pteropus*, *Rousettus*, and

Eidolon are significantly overrepresented relative to other species; however, certain *Pteropus*

species have been understudied (e.g., largely island-dwelling species like *P. chrysoproctus*, *P.*

tonganus, *P. neohibernicus*, and *P. temminckii*) (Fig. 3; Table 2). The occurrence and prevalence

of paramyxoviruses in these bat species are currently unknown due to very small sample sizes,

and future surveillance studies could target these taxa accordingly. Those that are particularly

elusive or difficult to sample in the wild may be good candidates for museum-based viral

screening, an approach which allows for efficient and taxonomically broad sampling across

space and time (40). Further, while the scope of this study was limited to Pteropodidae,

collecting and consolidating viral data from non-pteropodid bats could allow for broader

analyses of bat–paramyxovirus coevolution. Preliminary evidence suggests that some

insectivorous bat families host paramyxovirus clades distinct from those in pteropodid bats, a

finding that could be more robustly assessed with additional sampling and explicit

cophylogenetic tests (4,41).

On the viral side, the majority of studies have focused on henipaviruses; consequently,

Hendra, Nipah, and Cedar viruses are overrepresented, while pararubulaviruses are

comparatively neglected in the literature (Fig. 3; Table 2). Outside of the viruses detected in

pteropodids, there is a vast diversity of other mammalian and vertebrate paramyxoviruses that do

not appear in our dataset (Fig. 3d), including paramyxoviruses identified in insectivorous bats

(42–44). Conversely, 70 records in our database involve unclassified paramyxovirus sequences

that could appear anywhere in the viral tree. Viral taxonomy can be unstable and often lags

behind host taxonomy (3); future ICTV updates may recognize additional bat-borne

paramyxovirus species. Still, based on published links between 13 recognized paramyxovirus

species and 12 named pteropodid species (other isolated viruses were generally attributed to

Pteropus spp.), we performed a cophylogenetic analysis that revealed little evidence of

phylogenetic congruence between hosts and viruses, contrary to tentative hypotheses of

pteropodid–paramyxovirus coevolution posed in earlier surveillance studies (16). The strongest

coevolutionary signal was found in *Pteropus–Henipavirus* systems, though this result was not

statistically significant and may be an artifact caused by heavy sampling of *Pteropus* species for

henipaviruses. There was more support for host-switching among other links (e.g.,

Rousettus/*Eidolon*–pararubulaviruses). These findings raise the possibility of paramyxovirus

expansion into new pteropodid hosts and geographic areas, which may have implications for

spillover into human and domestic animal populations.

Our results also hint at methodologies and sample types that might yield higher

paramyxovirus prevalence. In general, urine, tissue, serum, and fecal samples appeared to have

higher PCR prevalence than blood, whereas saliva and swab samples had lower prevalence (Fig.

4). A more in-depth look at tissue tropism suggested that spleen and pooled feces are

undersampled relative to their typical prevalence, as these tissues have the highest PCR

prevalence among specified sources (i.e., excluding pooled tissue) (Fig. 5; Supplementary Table

1). These sample types could be good priorities for future surveillance studies of

paramyxoviruses in pteropodids and offer specific options for both invasive (e.g., spleen

extraction during necropsy) and non-invasive (e.g., under-roost fecal collection (37)) sampling

approaches. Both cross-sectional and longitudinal studies yielded similar PCR prevalence (Fig.

4), suggesting that both are suitable approaches for surveillance studies and that the study type

can be selected based on the research question at hand.

Here, we have: (1) highlighted the major spatial and taxonomic sampling gaps in the

pteropodid–paramyxovirus system; (2) identified sample types, regions, host subfamilies, and

study types associated with higher prevalence or seroprevalence; and (3) found evidence of host-

switching by paramyxoviruses within Pteropodidae. Importantly, we found no evidence that

paramyxovirus prevalence or seroprevalence in pteropodids has changed in recent decades, as

collection year was not a significant predictor in our models (Fig. 4). However, syntheses and

meta-analyses like this one are limited in scope and statistical robustness by data gaps as well as

a lack of stratified and standardized metadata reported in existing studies. Where possible,

researchers should report the specific tissue types, localities, sampling dates, taxonomic

identifications, and other metadata included in pooled samples, and provide detection estimates

at as fine-grained a level as possible to enable more detailed analysis of pteropodid–

paramyxovirus associations. This will help inform sampling protocols for surveillance of wild

bats and museum specimens. Our published dataset constitutes a first step toward gathering,

standardizing, and stratifying surveillance data to guide future studies of paramyxoviruses, which

continue to pose threats to human health. Addressing the sampling gaps outlined here will be

critical in developing a more complete picture of these bat–virus systems.

Methods

Systematic review

To generate our database of paramyxovirus detection attempts in wild pteropodids, we followed the PRISMA protocol (Supplementary Fig. 1) (45). On August 3rd, 2023, we systematically searched PubMed and Web of Science with the following string: (pteropodid* OR pteropid* OR "fruit bat" OR "fruit bats" OR "flying fox" OR "flying foxes" OR *Acerodon* OR *Aethalops* OR *Alionycteris* OR *Aproteles* OR *Balionycteris* OR *Boneia* OR *Casinycteris* OR *Chironax* OR *Cynopterus* OR *Desmalopex* OR *Dobsonia* OR *Dyacopterus* OR *Eidolon* OR *Eonycteris* OR *Epomophorus* OR *Epomops* OR *Haplonycteris* OR *Harpyionycteris* OR *Hypsignathus* OR *Latidens* OR *Macroglossus* OR *Megaerops* OR *Megaloglossus* OR *Melonycteris* OR *Mirimiri* OR *Myonycteris* OR *Nanonycteris* OR *Neopteryx* OR *Nesonycteris* OR *Notopteris* OR *Nyctimene* OR *Otopteropus* OR *Paranyctimene* OR *Penthetor* OR *Pilonycteris* OR *Plerotes* OR *Ptenochirus* OR *Pteralopex* OR *Pteropus* OR *Rousettus* OR *Scotonycteris* OR *Sphaerias* OR *Stenonycteris* OR *Styloctenium* OR *Syconycteris* OR *Thoopterus*) AND (paramyxovirus* OR *Paramyxoviridae* OR henipavirus* OR Hendra OR Nipah OR rubulavirus* OR morbillivirus*). We identified 1,260 records in total, and proceeded to screen 863 titles and abstracts from non-duplicated records. Studies were excluded if they did not include new (i.e., not republished) data on paramyxovirus detection attempts in wild bats—therefore, any studies of captive bats or experimental studies of bat-derived cell lines were excluded. We identified 137 articles for further full-text screening, of which 119 met the inclusion criteria for our study.

Data collection

Our final database included 119 studies and 1,482 records. Where prevalence was not

reported in the original studies, it was calculated from the number of positive and tested samples

before meta-analysis. For viral isolation attempts where positivity rate or number of positive

samples was not reported, we included a binary variable (0: unsuccessful detection; 1: successful

detection). Therefore, the same bat may be included in multiple records, if the study tested

multiple samples (e.g., different tissue types or detection methods) per individual bat. For papers

where over 100 sampling events are detailed, sampling events were summarized by location,

species, and tissue type. When original studies did not separate samples by species, locations,

tissue types, or years, all information was included in one record (separated by commas in the

relevant cells). Before downstream analyses, these pooled records were filtered out when

relevant, but they remain in the database for researchers who may be interested in specific

species, tissue types, or sites. When original studies did not report all the metadata for a study

(e.g., sampling season), these were not imputed.

Spatial and taxonomic sampling biases

Our dataset included bat samples from 33 countries in three geographical regions (Africa,

Asia, and Oceania; countries classified according to the United Nations Geoscheme). All

analyses were conducted in R (version 4.2.1) (46). We first visualized the number of studies and

(log-scale) samples per country by building maps with the aggregated Pteropodidae IUCN range

map overlaid for context. We then fitted three GLMs to these country-level variables with region

as the categorical predictor and one of three respective response variables: binary sampling status

(whether or not a country had been sampled for paramyxoviruses in pteropodids) as a binomial

response, with all countries in these three regions included; the number of unique studies per

country as a Poisson-distributed response, with only sampled countries included; and the number

of samples per country as another Poisson-distributed response, with only sampled countries

included. We assessed the fit of GLMs using McFadden's R^2 ('performance' package) and made

individual region-to-region pairwise comparisons using the 'emmeans' package (47,48).

To assess taxonomic sampling biases, we first aligned all sampled bats ($n = 58$) and

viruses ($n = 16$) identified at the species level to host (32) and virus (3) phylogenies. To do so,

we relabeled species names to fit the published taxonomy where necessary (e.g., *Pteropus*

medius was relabeled as *P. giganteus* for the purpose of taxonomic backbone alignment, though

we retain the currently recognized names in our published database). For all bat and virus species

in the phylogenies, we created a binary response variable representing whether or not each

species had been sampled or sampled for, respectively. For those species present in our dataset,

we calculated the number of unique studies and total samples associated with each bat and virus

species. To assess the degree of phylogenetic signal in sampling effort across bats and viruses,

we used the 'caper' package (49) to calculate D (<0: highly conserved; 0: phylogenetic clustering

under Brownian motion; 0-1: intermediate phylogenetic signal; 1: phylogenetically random

sampling distribution; >1: overdispersed) for binary sampling effort and Pagel's lambda (λ) (0:

phylogenetically random sampling distribution; 0-1: intermediate phylogenetic signal; 1:

phylogenetic clustering under Brownian motion) for the $\log_{10}$-transformed number of studies and

samples. Finally, we used the package 'phylofactor' (Washburne et al. 2017) to apply

phylogenetic factorization, a graph-partitioning algorithm, to both the bat and virus phylogenies.

This approach partitioned binary sampling effort (binomial response), number of studies

(Poisson distribution), and number of samples (Gaussian distribution due to an extreme skew in

number of samples and consequent transformation via natural log) across the trees using iterative

GLMs, producing a list of bat or virus clades (containing at least three species) that differ

significantly from the rest of the tree in sampling effort. We used Holm's sequentially rejective

test with a family-wise error rate of 5% to account for multiple comparisons and determine the

number of statistically significant clades (50).

Meta-analysis of viral and antibody positivity

We used the 'metafor' package (34) to run two hierarchical meta-analysis models for two

separate viral positivity datasets: PCR detection (n = 786 records) and serological detection (n =

313 records). The response variables in our models were the Freeman-Tukey double arcsine-

transformed proportions of paramyxovirus PCR-positive or seropositive samples and their

sampling variances. Models were fitted using restricted maximum likelihood (REML) and

included bat genus, genus-level phylogeny (32), and country as random effects along with an

observation-level random effect nested inside a study-level random effect; all models were also

weighted by inverse sampling variance. We used genus instead of species as a random effect in

these models to retain records from bats not identified to the species level (e.g., unspecified

Australian *Pteropus* spp.). We calculated the contribution of heterogeneity (true variability rather

than noise) to variance in viral positivity (I^2) for the entire dataset, as well as the proportional

contribution for each random effect. Our models included all available non-collinear predictors.

For the PCR model, we included fixed effects of region (Oceania, Asia, Africa), study type (cross

sectional, longitudinal, or both), sample type (feces, serum, saliva, swab, tissue, various),

sampling year, and host subfamily (Cynopterinae, Eidolinae, Epomophorinae, Macroglossusinae,

Pteropodinae, Rousettinae); for the serology model, we included fixed effects of region (Oceania,

Asia, Africa), sampling year, and host subfamily (Cynopterinae, Eidolinae, Epomophorinae,

Macroglossusinae, Pteropodinae, Rousettinae). For records reflecting detection attempts across a

range of sampling years, the midpoint of the start and end years was used to calculate the average

"sampling year".

To assess paramyxovirus prevalence in different tissue types (as tissue type was a

collinear predictor in all possible models and therefore could not be included), we also calculated

overall viral prevalence per tissue type for records using PCR.

Cophylogenetic analysis

We lastly used PACo, with the 'paco' package, to assess coevolution between

paramyxoviruses and pteropodids and thus the dependence of the viral phylogeny on the host

phylogeny (35). For this analysis, we defined a known association as either a PCR detection or

viral isolation of an ICTV-recognized paramyxovirus species in a described species of

pteropodid, thereby excluding serological detections (due to lower viral specificity) and any PCR

or viral culture detection in bats recognized only to the genus level (e.g., unspecified Australian

Pteropus spp.). We conducted a goodness-of-fit test to evaluate how well a codivergence model

fits the matrix of known host–virus links, and then established its statistical significance by

fitting the model to 5000 randomizations of the matrix and testing the null hypothesis that host

and virus species are randomly associated. We generated residuals for each host–virus link

(smaller residuals indicate more support for coevolution, and higher residuals indicate more

support for host-switching). We then visualized the host–virus links with a tanglegram of all

known links in our dataset using the package 'phytools' (51).

References

1. Carlson CJ, Albery GF, Merow C, Trisos CH, Zipfel CM, Eskew EA, et al. Climate change

increases cross-species viral transmission risk. Nature. 2022;607(7919):555–62.

2. Pomeroy LW, Bjørnstad ON, Holmes EC. The evolutionary and epidemiological dynamics of

the *Paramyxoviridae*. J Mol Evol. 2008;66(2):98–106.

3. Rima B, Balkema-Buschmann A, Dundon WG, Duprex P, Easton A, Fouchier R, et al. ICTV

Virus Taxonomy Profile: *Paramyxoviridae*. J Gen Virol. 2019 Dec;100(12):1593–4.

4. Drexler JF, Corman VM, Müller MA, Maganga GD, Vallo P, Binger T, et al. Bats host major

mammalian paramyxoviruses. Nat Commun. 2012;3(1):796.

5. Giannini NP. Pteropodidae. In: Wilson DE, Mittermeier R, editors. Handbook of the Mammals

of the World. Barcelona: Lynx Edicions; 2019. p. 16–61.

6. Tian S, Zeng J, Jiao H, Zhang D, Zhang L, Lei C, et al. Comparative analyses of bat genomes

identify distinct evolution of immunity in Old World fruit bats. Sci Adv. 2023;9(18):0141.

7. Mougari S, Gonzalez C, Reynard O, Horvat B. Fruit bats as natural reservoir of highly

pathogenic henipaviruses: balance between antiviral defense and viral tolerance. Curr Opin Virol.

2022 Jun 1;54:101228.

8. Halpin K, Young PL, Field HE, Mackenzie JS. Isolation of Hendra virus from pteropid bats: A

natural reservoir of Hendra virus. J Gen Virol. 2000;81(8):1927–32.

9. Chua KB, Lek Koh C, Hooi PS, Wee KF, Khong JH, Chua BH, et al. Isolation of Nipah virus

from Malaysian Island flying-foxes. Microbes Infect. 2002;4(2):145–51.

10. Drexler JF, Corman VM, Gloza-Rausch F, Seebens A, Annan A, Ipsen A, et al. Henipavirus

RNA in African bats. PLoS ONE. 2009;4(7):6367.

11. Marsh GA, de Jong C, Barr JA, Tachedjian M, Smith C, Middleton D, et al. Cedar virus: a

novel Henipavirus isolated from Australian bats. PLoS Pathog. 2012;8(8):e1002836.

12. Chua KB, Wang LF, Lam SK, Crameri G, Yu M, Wise T, et al. Tioman virus, a novel

paramyxovirus isolated from fruit bats in Malaysia. Virology. 2001 May 10;283(2):215–29.

13. Baker KS, Todd S, Marsh GA, Crameri G, Barr J, Kamins AO, et al. Novel, potentially

zoonotic paramyxoviruses from the African straw-colored fruit bat *Eidolon helvum*. J Virol. 2013

Feb;87(3):1348–58.

14. Barr J, Smith C, Smith I, de Jong C, Todd S, Melville D, et al. Isolation of multiple novel

paramyxoviruses from pteropid bat urine. J Gen Virol. 2015;96(1):24–9.

15. Baker KS, Tachedjian M, Barr J, Marsh GA, Todd S, Crameri G, et al. Achimota

Pararubulavirus 3: A New Bat-Derived Paramyxovirus of the Genus Pararubulavirus. Viruses.

2020 Nov;12(11):1236.

16. Vidgen ME, de Jong C, Rose K, Hall J, Field HE, Smith CS. Novel paramyxoviruses in

Australian flying-fox populations support host–virus co-evolution. J Gen Virol.

2015;96(7):1619–25.

17. Plowright RK, Field HE, Smith C, Divljan A, Palmer C, Tabor G, et al. Reproduction and

nutritional stress are risk factors for Hendra virus infection in little red flying foxes (Pteropus

scapulatus. Proc R Soc B Biol Sci. 2008;275(1636):861–9.

18. Mortlock M, Geldenhuys M, Dietrich M, Epstein JH, Weyer J, Pawęska JT, et al. Seasonal

shedding patterns of diverse henipavirus-related paramyxoviruses in Egyptian rousette bats. Sci

Rep. 2021 Dec 20;11(1):24262.

19. Becker DJ, Eby P, Madden W, Peel AJ, Plowright RK. Ecological conditions predict the

intensity of Hendra virus excretion over space and time from bat reservoir hosts. Ecol Lett.

2023;26(1):23–36.

20. Juman MM, Gibson L, Suu-Ire RD, Languon S, Quaye O, Fleischer G, et al. Ecological and

Reproductive Cycles Drive Henipavirus Seroprevalence in the African Straw-Coloured Fruit Bat

(*Eidolon helvum*). Ecol Evol. 2024;14(11):e70555.

21. McKee CD, Islam A, Luby SP, Salje H, Hudson PJ, Plowright RK, et al. The ecology of

nipah virus in bangladesh: A nexus of land-use change and opportunistic feeding behavior in

bats. Viruses. 2021;13(2):169.

22. Eby P, Peel AJ, Hoegh A, Madden W, Giles JR, Hudson PJ, et al. Pathogen spillover driven

by rapid changes in bat ecology. Nature. 2023;613(7943):340–4.

23. Kessler MK, Becker DJ, Peel AJ, Justice NV, Lunn T, Crowley DE, et al. Changing resource

landscapes and spillover of henipaviruses. Ann N Y Acad Sci. 2018;1429(1):78–99.

24. Yadav PD, Shete AM, Kumar GA, Sarkale P, Sahay RR, Radhakrishnan C, et al. Nipah virus

sequences from humans and bats during Nipah outbreak. Emerg Infect Dis. 2019;25(5):1003–6.

25. Arunkumar G, Chandni R, Mourya DT, Singh SK, Sadanandan R, Sudan P, et al. Outbreak

investigation of Nipah Virus disease in Kerala. J Infect Dis. 2019;219(12):1867–78.

26. Carlson CJ, Brookson CB, Becker DJ, Cummings CA, Gibb R, Halliday FW, et al. Pathogens

and planetary change. Nat Rev Biodivers. 2025 Jan;1(1):32–49.

27. Cohen LE, Fagre AC, Chen B, Carlson CJ, Becker DJ. Coronavirus sampling and

surveillance in bats from 1996–2019: a systematic review and meta-analysis. Nat Microbiol.

2023 Jun;8(6):1176–86.

28. Young CCW, Olival KJ. Optimizing Viral Discovery in Bats. PLOS ONE. 2016 Feb

11;11(2):e0149237.

29. Becker DJ, Crowley DE, Washburne AD, Plowright RK. Temporal and spatial limitations in

global surveillance for bat filoviruses and henipaviruses. Biol Lett. 2019 Dec

11;15(12):20190423.

30. Crowley D, Becker D, Washburne A, Plowright R. Identifying Suspect Bat Reservoirs of

Emerging Infections. Vaccines. 2020 May 17;8(2):228.

31. Plowright RK, Parrish CR, McCallum H, Hudson PJ, Ko AI, Graham AL, et al. Pathways to

zoonotic spillover. Nat Rev Microbiol. 2017;15(8):502–10.

32. Upham NS, Esselstyn JA, Jetz W. Inferring the mammal tree: Species-level sets of

phylogenies for questions in ecology, evolution, and conservation. PLOS Biol.

2019;17(12):3000494.

33. Washburne AD, Silverman JD, Leff JW, Bennett DJ, Darcy JL, Mukherjee S, et al.

Phylogenetic factorization of compositional data yields lineage-level associations in microbiome

datasets. PeerJ. 2017 Feb 9;5:e2969.

34. Viechtbauer W. Conducting Meta-Analyses in R with the metafor Package. J Stat Softw.

2010 Aug 5;36:1–48.

35. Balbuena JA, Míguez-Lozano R, Blasco-Costa I. PACo: A Novel Procrustes Application to

Cophylogenetic Analysis. PLOS ONE. 2013 Apr 8;8(4):e61048.

36. Dallas TA, Becker DJ. Taxonomic resolution affects host−parasite association model

performance. Parasitology. 2021;148(5):584–90.

37. Giles JR, Peel AJ, Wells K, Plowright RK, McCallum H, Restif O. Optimizing noninvasive

sampling of a zoonotic bat virus. Ecol Evol. 2021;11(18):12307–21.

38. McKee CD, Islam A, Rahman MZ, Khan SU, Rahman M, Satter SM, et al. Nipah Virus

Detection at Bat Roosts after Spillover Events, Bangladesh, 2012–2019. Emerg Infect Dis. 2022

Jul;28(7):1384–92.

39. Pernet O, Schneider BS, Beaty SM, LeBreton M, Yun TE, Park A, et al. Evidence for

henipavirus spillover into human populations in Africa. Nat Commun. 2014 Nov 18;5(1):5342.

40. Colella JP, Bates J, Burneo SF, Camacho MA, Bonilla CC, Constable I, et al. Leveraging

natural history biorepositories as a global, decentralized, pathogen surveillance network. PLOS

Pathog. 2021 Jun 3;17(6):e1009583.

41. Mortlock M, Kuzmin IV, Weyer J, Gilbert AT, Agwanda B, Rupprecht CE, et al. Novel

Paramyxoviruses in Bats from Sub-Saharan Africa, 2007–2012. Emerg Infect Dis. 2015

Oct;21(10):1840–3.

42. Kurth A, Kohl C, Brinkmann A, Ebinger A, Harper JA, Wang LF, et al. Novel

Paramyxoviruses in Free-Ranging European Bats. PLOS ONE. 2012 Jun 21;7(6):e38688.

43. Wilkinson DA, Temmam S, Lebarbenchon C, Lagadec E, Chotte J, Guillebaud J, et al.

Identification of novel paramyxoviruses in insectivorous bats of the Southwest Indian Ocean.

Virus Res. 2012 Dec 1;170(1):159–63.

44. Larsen BB, Gryseels S, Otto HW, Worobey M. Evolution and Diversity of Bat and Rodent

Paramyxoviruses from North America. J Virol. 2022 Feb 9;96(3):e0109821.

45. Moher D, Liberati A, Tetzlaff J, Altman DG, PRISMA Group. Preferred reporting items for

systematic reviews and meta-analyses: the PRISMA statement. PLoS Med. 2009 Jul

21;6(7):e1000097.

46. R Core Team. R: A language and environment for statistical computing [Internet]. Vienna,

Austria: R Foundation for Statistical Computing; 2022. Available from: https://www.R-

project.org/

47. Lüdecke D, Ben-Shachar MS, Patil I, Waggoner P, Makowski D. performance: An R Package

for Assessment, Comparison and Testing of Statistical Models. J Open Source Softw. 2021 Apr

21;6(60):3139.

48. Lenth RV. emmeans: Estimated Marginal Means, aka Least-Squares Means. R package

version 1.8.2 [Internet]. 2022. Available from: https://CRAN.R-project.org/package=emmeans

49. Orme D, Freckleton R, Thomas G, Petzoldt T, Fritz S, Isaac N, et al. caper: Comparative

Analyses of Phylogenetics and Evolution in R. R package version 1.0.3 [Internet]. 2023.

Available from: https://CRAN.R-project.org/package=caper

50. Holm S. A Simple Sequentially Rejective Multiple Test Procedure. Scand J Stat.

1979;6(2):65–70.

51. Revell LJ. phytools 2.0: an updated R ecosystem for phylogenetic comparative methods (and

other things). PeerJ. 2024 Jan 5;12:e16505.

Supporting Information Captions

Supplementary Fig. 1. PRISMA reporting diagram for study selection and screening.

Supplementary Table 1. Database breakdown by sample type and detention method.

Sample size (records and number of samples), percentage of records with at least one positive

sample, and average prevalence by detection method, sample type, and viral subfamily.

Supplementary Table 2. Database breakdown by type of paramyxovirus. Number of studies,

host species, and samples tested for different paramyxoviruses, with positivity rates reported for

serological and PCR detection attempts as well as number of successful viral isolations.

Supplementary Table 3. Database breakdown by country. Number of studies, number of

samples, and mean viral prevalence and seropositivity rates for all countries.

Supplementary Table 4. Results from PACo cophylogenetic analysis of paramyxoviruses in

Pteropodidae. Jackknifed squared residuals and upper 95% confidence intervals for each known

pteropodid–paramyxovirus link from PACo.