Complexity shapes uniqueness: Neuropil volumes and synaptic clusters shape behavioural

plasticity under challenging environments in the invasive Argentine ants

Srikrishna Narasimhan[1], María Eugenia Villar[2], Violette Chiara[1,3], Ignacio Arganda-Carreras[4,5,6,7],

Sara Arganda[2,8], Magdalena Witek[1*], Iago Sanmartín-Villar[1,9*§]

[1]*Laboratory of Social and Myrmecophilous Insects, Museum and Institute of Zoology, Polish*

Academy of Sciences, Twarda 51/55, 00-818, Warsaw, Poland

[2]*Departamento de Biología y Geología, Física y Química Inorgánica, Área de Biodiversidad y*

Conservación, Universidad Rey Juan Carlos, Tulipán s/n, 28933 Móstoles, Spain

[3]Aquatic Ecology, Lund University, Lund, Sweden

[4]*Department of Computer Science and Artificial Intelligence, University of the Basque Country*

(UPV/EHU), Manuel Lardizabal 1, Donostia, 20008, Basque Country, Spain

[5]*IKERBASQUE, Basque Foundation for Science, Plaza Euskadi 5, Bilbao, 48009, Basque Country,*

Spain

[6]*Donostia International Physics Center (DIPC), Manuel Lardizabal 4, Donostia, 20018, Basque*

Country, Spain

[7]*Biofisika Institute (CSIC-UPV/EHU), Barrio Sarriena, Leioa, 48940, Basque Country, Spain*

[8]*Instituto de Investigación en Cambio Global (IICG-URJC), Universidad Rey Juan Carlos, Tulipán*

s/n, 28933 Móstoles, Spain

[9]*ADIBECO, Biodiversidad y Gestión ambiental, Universidad de León, 24007, León, Spain*

*Joint last authors

§Corresponding author: Iago Sanmartín-Villar, *Departamento de Biodiversidad y Gestión*

ambiental, Universidad de León (León, Spain), isanv@unileon.es

Abstract:

Adaptation to new environments is key for organisms' survival, but also for their invasiveness in

their introduced areas. Behaviour is considered the fastest phenotype allowing adaptation, but

its plasticity can involve costs as neural development. Although individuals' investment in

cognition was pointed out as unnecessary for colony behaviour in eusocial insects, recent studies

are highlighting the behavioural dependence on neural traits in eusocial insects. The costs of

producing behavioural and neural plastic offspring could exceed the investments of eusocial

insects, in which one or certain reproductives must produce multiple offspring. Thus, we wanted

to analyse the link between the neuroanatomy and the behavioural variability of *Linepithema*

humile, an invasive species organised in supercolony units containing millions of individuals, to

understand its adaptive mechanisms. We repeatedly tested same aged callow workers of *L.*

humile in behavioural tests of increasing environmental complexity and analysed the volume of

their brain functional areas (neuropils) and the synaptic clusters abundance in the mushroom

body calices (information processing). Given the potential large cost of plasticity, we expected to

find homogeneous interindividual neuronal structures and behavioural responses. Although *L.*

humile is considered a monomorphic species, body size conditioned behavioural and neural traits

and determined individuals efficiency in exploring simple environments. Contrary to our

expectations, the increase in environmental complexity revealed the behavioural plasticity of

Linepithema humile workers as well as its correlation with neuropil volumes and synaptic

clusters. Our results highlight the relevance of the central complex and the mushroom bodies on

exploration efficiency rather than optic and olfactory lobes. Behavioural plasticity under complex

environments relied on the synaptic connections of the olfactory processing area (dense lip),

while individuals with higher number of synaptic connections on the visual processing area

(collar) explored complex environments less efficiently. Our results suggest that behavioural

differences that correlate with morphological traits might promote adaptive mechanisms in

simple environments, whereas neurologically based plastic behavior may be necessary to adapt

in complex environments.

Keywords: Argentine ant, social insects, personality, microglomureli, volumetry, Neuroethology.

Introduction

Species adaptation to the environment has been a revolutionary concept in the understanding

of Biology since the publication of *On the Origin of Species* (Darwin, 1859). Gradual adaptations,

often driven by the accumulation of mutations over many generations, allow physiological and

morphological traits to evolve at the species level. However, evolutionary change can also occur

rapidly, enabling organisms to adapt to sudden environmental shifts, such as those caused by

climate change, or to establish themselves in new regions. A notable example of the latter is the

introduction of non-native species, often a byproduct of globalization, which has become a

significant ecological and economic challenge (*e.g.* Bertelsmeier, 2021; López et al., 2023). To

become established in a new area, introduced species must cope with a range of diseases,

resources, competitors, and conditions. In the case of the alien invasive species, these

adaptations not only allow them to survive in the new place, but to expand their range and

colonize new areas (*e.g.* Prentis et al., 2008; Stapley et al., 2015).

One of the mechanisms that enable species to rapidly modify the way they interact with the

environment is through their behaviour (reviewed in Snell-Rood, 2013). At the population level,

these behavioural adaptations can arise from two different mechanisms. One of them consists of

the difference of interindividual behavioural patterns (personality; Dall et al., 2004) and their

match with the environmental conditions, which determines the fitness of the individuals.

Another mechanism consists in the intraindividual changes in behaviour through processes and

experiences (plasticity; Dingemanse et al., 2010). While these concepts may directly apply to

solitary species, the proximate causes of behavioural variability and the fitness outcome of its

dynamics differs in eusocial organisms. In such species, evolutionary pressures and selections do

not act at the individual but at the colony level (*e.g.*, mole rats, Jacobs & Jarvis, 1996; ants,

Delgado & Solé, 2000). On one hand, interindividual behavioural variability in eusocial organisms

often relies on task speciation (Jeanson & Weidenmüller, 2014), which is frequently related with

individuals' age (polyethism; reviewed in Beshers & Fewell, 2001), a form of intraindividual

behavioural plasticity through development (also see Jeanson 2019). On the other hand, eusocial

behaviour is strongly influenced by colony-level dynamics, as new behaviours often emerge as

collective phenomena (reviewed in Lihoreau et al., 2012; Navas-Zuloaga et al., 2022).

However, in the last years a growing number of studies are demonstrating the importance of

interindividual behavioural differences (*e.g.* Maák et al., 2020; Trigos-Peral et al., 2023) and

intraindividual behavioural plasticity (reviewed in Lucon-Xiccato et al., 2024) in the context of

coping with environmental challenges at the individual level in eusocial species. Whether due to

phenotype, task specialization or perceived experiences, proximal mechanisms such as

neuroanatomical development, are responsible for the variability of insect behavior. Although

insect morphology stops changing after metamorphosis, their brains show large plasticity in the

adult stage. For instance, the relative volumes of certain neuronal regions (*e.g.* Gronenberg et

al., 1996) or even the total volume of ants' brains can increase through maturation and

experience in the adult stage (*Camponotus* sp.. Gronenberg et al., 1996; Yilmaz et al., 2016b;

Oecophylla smaragdina and *Formica subsericea*, Kamhi et al., 2017). In polymorphic species,

larger workers show larger optical lobes - necessary when foraging - and smaller workers show

higher ratio of collars of the mushroom body to optic lobes, *i.e.* more processing than perception

of the visual cues (Arganda et al., 2020). The navigation under natural skylight involves the

increase of the synaptic buttons of visual projection neurons and volume of the central complex

in *Cataglyphus nodus* (Grob et al., 2017). When learning, *Acromyrmex ambiguus* shows a

temporal increase in the synaptic cluster (microglomureli) in the olfactory region rather than in

the optic region of the mushroom body (Falibene et al., 2015).

The production, development, and maintenance of neuronal plasticity (*e.g.* new or larger neural

routes) involves resource allocation and costs for the individual (Jeanson & Weidenmüller, 2014).

Behavioural processes such as memory and learning allow faster adaptation (Fawcett et al., 2013)

but involves the organisation of the brain functional areas (neuropils) in charge of processing

sensory information (mushroom bodies of insects; Aso et al., 2014; Li & Strausfeld, 1999). Some

of these costs could scale to the progenitor level, which might invest more resources to allow

individuals coping with neuroanatomic plasticity, leading to a trade-off related with the quantity

and quality of the offspring production (Clutton-Brock et al., 1985; Einum & Fleming, 2004.).

Therefore, we expected an inverse relationship between the behavioural plasticity and the

colony size on eusocial animals, *i.e.* queens investing in workers' quality in small colonies but in

workers' quantity in large colonies (Kramer & Schaible, 2013). Under this scenario, large colonies

might rely more on the polyethism and the behavioural traits arised by the collective than in the

individual behavioural properties.

The Argentine ant (*Linepithema humile*, Mayr 1868) could be one of the most representative

examples to illustrate trade-offs between the benefits and costs of behavioural variability due to

the millions of members that form its supercolonies (non-competitive colonies; reviewed in

Moffett, 2012) and their necessity to adapt to the introduced areas in which they behave as

invasive. We therefore used the Argentine ant to highlight the individual capacities to interact

with the environment and its dependence on neuroanatomy. At present, we do not know how

differences and neuroethological plasticity of workers cause or affect colony adaptation in

invasive species. We expect that in a polygynous superorganism (Hölldobler & Wilson, 1990) in

which workers are almost non-genetically related (Keller & Fournier, 2002) and with the

dimension of the Argentine ant introduced supercolonies (reviewed in Angulo et al., 2024), the

behavioural variability should rely more on genetic or developmental differences among and

within the supercolony members than in the individual capacity to modulate behavioural

responses. On the contrary, some characteristics of the *L. humile* workers such as the rapid

consolidation of long-term memory (Wagner et al., 2023) and traits found in other ant species

like learning (Czaczkes et al., 2013; Jeanson & Weidenmüller, 2014), personality and tool use

(Maák et al., 2020) or task syndromes (Trigos-Peral et al., 2023), suggest a high relevance of

individual cognition in problem-solving in eusocial organised species (also shown in bees,

Lemanski et al., 2021; bumblebees, Hill et al., 2023). Unraveling the underlying mechanisms

involved in resolving environmental challenges faced by the introduced *L. humile* supercolony

may help to prevent and control the continued expansion (reviewed in Angulo et al., 2024) and

long-term stabilisation (Baratelli et al., 2023) of this ant species.

We performed standardised laboratory experiments with different environmental complexities

to analyse the behavioural strategies used by the Argentine ants. We considered exploratory

behaviour as a basic mechanism to solve problems in new environments (open field test,

response to an object and solving maze). We focused on the relationship between the individual

neuroanatomy and the behavioural plasticity in the face of environmental challenges to

understand potential dependences of behaviour on the brain and thus, colony costs associated

with individual production. Argentine ants have managed to spread widely in their introduced

areas owing to their high activity, population density, and capacity to monopolise larger foraging

territories (Carpintero & Reyes-López, 2008; Human & Gordon, 1996), so we expected

experimental ants to show high level of exploratory behaviour. As a possible strategy of reducing

developmental costs in maintaining a large population (Sanmartín-Villar et al., 2021), we

expected ants showing low inter- and intraindividual behavioural variability when increasing the

environment complexity and sparse relationship between the neuroanatomy (microglomureli

organisation and brain region volumes). In other words*,* we expected that the colony interaction

with the environment will be a product of individuals' trial and errors and without cognitive

processes associated (Lihoreau et al., 2012).

Material and Methods

Sampling collection and laboratory maintenance

Linepithema humile colonies were collected in NW Spain between the 23[rd] and 28[th] of August

2022 in five locations separated by ca. 7 km (in latitudinal order; Catoira: 42°40'18"N, 8°43'41";

Rianxo: 42°38'43", 84°9'14"; Carril: 42°37'14", 8°46'21"; Chazo: 42°36'27", 8°51'25"; Vilanova:

42°34'00", 8°49'42"). Queens, workers, and larvae were transferred with the soil of their nests

to plastic containers with the inner part of the walls covered by polytetrafluoroethylene (Fluon®)

to prevent ants from escaping. The contents of each collected nest were spread in plastic pot

bases (97 x 40 x 7 cm) also covered by fluon®. We added plastic tubes filled with water and cotton

to provide humidity to the ants and food *ad libitum* following the recipe based on Bhatkar &

Whitcomb (1970) and improved according to Csata *et al.* (2020). This food was used throughout

the experiment. Over the course of five days, all ants successively entered into the tubes and

were transferred to new plastic containers and transported, without soil, to the Museum and

Institute of Zoology of Warsaw (Poland). Ant nests ("source nest" hereafter) were maintained in

temperature chambers at 26°C with a 12:12h light:dark period, the optimal temperature from

egg to adult development of *L. humile* (Abril et al., 2010). Food was replaced in the source nest

three times per week until the end of the experiment.

To control the age of the focal individuals and avoid behavioural and neural differences due to

polyethism (*e.g.*, Fahrbach et al., 1998; Kamhi et al., 2019; Tripet & Nonacs, 2004) and different

experiences (*e.g.*, Weidenmüller et al., 2009), we followed the development of last instar larvae

until their metamorphosis. To facilitate this process and to standardise the experiences lived by

focal individuals in their larval stage, we created homogeneous experimental groups by

transferring from the source colonies 20 workers and five last-instar larvae to Petri dishes (Ø=

5.7 cm) containing a 1.5 ml Eppendorf tube filled with water and cotton and the tube's lid filled

with food. Twenty experimental group replicates were initially isolated from each source nest. In

cases where two callow workers emerged in the same group, the newest callow worker was

transferred to fresh experimental groups. We reached a maximum of 50 and minimum of 20

experimental groups for the source nests. We checked each experimental group daily for new

emergences. One day after emerging, focal individuals were marked with pen markers (Edding

Creative 751©). Focal individuals were easily identified even one day after their emergence

because the colour of their cuticle was lighter than the one of their nestmates. Behavioural tests

started ten days after the emergence of focal individuals, a period that we considered optimal to

guarantee the development of the locomotive workers' function (see also Abril et al., 2010).

Behavioural test

To study inter- and intra individuals' behavioural variability, we exposed each focal individual (N=

172) to a series of three consecutive 10-minute tests. Five observations of the three different

tests, one per day for five consecutive days, were made (N= 860 tests) at 23.75±0.73 °C and

54.67±2.78 % of humidity, testing each individual over a similar time frame. Tests were always

conducted in the following order for each individual and observation: (i) open field test, focused

on the percentage of new area explored; (ii) object exposure test, focused on the interaction with

new objects; and (iii) maze test, focused on the decision making on whether to explore further in

a complex environment. A total of 13 individuals were not exposed to the objects and mazes, *i.e.*

they remained for 30 min in the open field (see below), to control the effect of the exposure to

new objects on the behaviour and brain structure. A total of 20 individuals were not exposed to

any of the behavioural tests as a control for possible inherent factors of our procedure such as

manipulation and isolation (Bernadou et al., 2018; Seid & Junge, 2016) that could cause some

changes to the brain.

Behavioural tests were performed in separate Petri dishes inside three 60 x 70 x 60 cm wooden

boxes with white walls to limit visual references and thus, without distal clues for the focal ants

(see Harris et al., 2010). Ants' behaviour was recorded from the top of the boxes with cameras

(two Sony FDR-AX53; one Canon Legria HF M56) and tracked with AnimalTA software (Chiara &

Kim, 2023). A maximum of 10 ants were recorded at the same time for a single camera. Before

each series of tests started, we transferred the experimental groups from the temperature

chambers and placed them in the wooden boxes for 10 minutes as pre-acclimatisation.

Open field test

Following the 10 min of acclimatisation, focal ants were carefully isolated into their respective

empty Petri dishes (Ø= 5.7 cm) using a brush. Petri dishes were covered with their lids after the

addition of each individual. These dishes possessed a red circle of 1 cm diameter drawn on the

centre and outer part of the bottom dish as a reference for placing the object in the following

test (see next paragraph). We assumed that ants did not perceive this circle due to its position

(no odour because it was out of the dish). Recordings started before the introduction of the first

ant and lasted 10 min after the introduction of the last ant in the experimental setup. By doing

so, we were able to register the behaviour of all focal individuals since their contact with the new

arena. This procedure was also applied to the other behavioural tests. Upon introducing the last

ant, we closed the wooden boxes with a lateral white wooden cover.

Object exposure test

Following the 10-minute open field test, we removed the cover of the wooden boxes and

introduced an object in each Petri dish. The objects were added in the centre of the arena,

matching its contour with the red circle designed. After adding the object to the last dish, the

cover of the wooden box was put back and we recorded the ants' behaviour for 10 mins. Objects

consisted of a three-level circular pyramid (Ø= 1 cm; Fig. 1A) designed on Tinkercad

(https://www.tinkercad.com/) and printed on resin (Anycubic Eco UV Resin White) using

Chitubox® and a 3-D printer (Elegoo Mars 2P). The objects were painted in blue (Pinty Plus Aqua;

Ref. 320). We consider that the shape, material, colour, and odour were uncommon in the focal

species environment and thus, relevant as a novelty stimulus. After the end of each test, the

objects were washed with hand soap and tap water and allowed to dry under sunlight.

Maze test

Following the 10 min object exposure test, ants were individually transferred to a new Petri dish

(Ø= 5.7 cm) respectively containing a white resin-printed maze (height 4 mm, Ø= 5.7 cm; Fig. 1B).

The maze consisted of a circular platform with two inner chambers (start and end) connected by

a concentric path of 3 mm wide. The path was composed of a sequence of five 1 cm segments.

Each segment ended in a Y-choice with the next two branches oriented 130˚ from the previous

path. Opposed at an angle of 100˚ to each of the segments, the maze possessed a dead-end 1

cm segment. In this way, the maze was composed of six Y-choices connected by the concentrical

path. Half of the mazes had dead-ends directed toward the left and the other half had dead-ends

directed toward the right. On each Y-choice, ants had the choice between selecting a dead-end

path or to continue advancing through the maze. The three vertices of the Y-choice were

expanded to the interior of the path as projections constraining it to 2 mm. This forced the focal

ants to face the two possible choices and avoid them selecting a choice based on the possible

thigmotaxis (wall-following) used to guide them through the previous path (see Czaczkes, 2018).

Focal ants were introduced to the ovoid start chamber of the maze (2.1 x 1 cm) with only one

exit towards the maze path. We covered the maze with a plexiglas layer to avoid ants escaping.

Ants that selected six times the same direction following the concentric path through the six

consecutive Y-choices encountered the ending reward chamber. This reward chamber was

circular ($\emptyset$= 1 cm) and positioned in the middle of the Petri dish (occupying the same place as the

red circle of the open field test and the object on the object exposure test). Before each test, a

drop of sucrose was added in this reward chamber (1M; see Czaczkes, 2018). The ending chamber

was the only section of the maze without resin floor, to allow a better removal of the sucrose

drop. The sucrose was added to stimulate the ants that found the end chamber for future trials,

although our aim was not to test learning (no training).

To analyse the variability of choice strategy under novel environments, focal ants were exposed

on the three first consecutive observations to mazes with a concentric path oriented in one

direction and to mazes oriented in the opposite direction on the last two observations. By this,

we could check if ants were habituated to the maze and how they reacted after a change, which

we assumed is a more demanding cognitive process (central integration; see Menzel & Giurfa,

2001) and thus, the opportunity to compare personality traits and cognition based on different

mechanisms (see Carere & Locurto, 2011). After 10 min of recording, ants were returned to their

original experimental group and then to the temperature chamber until the next day, in which

tests were repeated.

Ants belonging to the control group were maintained 30 minutes in empty arenas (open field) to

control the effect of experiencing new or/and complex environments (object, maze) while

considering the effects of experimentation (manipulation and isolation). Similar to the

experimental group, we made 5 observations over 5 consecutive days.

After 16 days of their emergence (immediately after the last test on the 5th day for focal ants),

individuals were isolated in 15 ml falcon tubes and placed in a thermally isolated box with ice

packs until brain extraction.

Brain extraction and fixation

Brain extraction was done following the protocol of Ott (2008). Focal and control ants (N=205)

were anaesthetised on ice. Immediately after decapitation, the eye span of each ant was

measured using a micrometre under a binocular lens (Olympus SZH10). Brain extractions were

performed in HEPES solution (150 mM NaCl, 5 mM KCl, 5 mM CaCl$_2$, 25 mM sucrose, 10 mM

HEPES) in silicone-coated Petri dishes. Extracted brains were individually fixed in 1% Zn-PFA

solution overnight and continuously shaken (60 rpm; Thermo Fisher) at room temperature. The

brains were then rinsed in HEPES solution six times for 10 minutes each. Subsequently, the brains

were fixed in Dent's Fixative (1:4 DMSO-Methanol solution) at room temperature for 1 hour and

30 minutes. After fixation, the brains were stored in 100% methanol at -20°C until the

immunostaining procedure. Once all brains were collected, they were shipped to the University

Rey Juan Carlos (Spain) under cold conditions to ensure preservation.

Immunohistochemical analysis and imaging

To image the neuroanatomy of the extracted brains, they were rehydrated in 0.1 M Tris buffer

and seated in a block and permeabilization solution of normal goat serum (NGS) for 1 hour

(PBSTN solution: 5% NGS + 0.005% sodium azide in 0.2% Triton-X phosphate buffered saline

[PBST]) at room temperature on a shaker. After this step, brains were incubated on the shaker

during four nights at room temperature in the primary antibody solution (SYNORF1 –

AB_2315426, Developmental Studies Hybridoma Bank – diluted 1:30 in PBSTN). Then, brains

were washed six times (10 minutes/wash) in 0.2% PBST and incubated for another three nights

protected from light on a shaker in AlexaFluor488 (ThermoFisher) Goat anti-Mouse IgG (H+L)

Cross-Adsorbed ReadyProbes™ Secondary Antibody (1:500 in PBSTN). Once immunolabeled,

brains were washed again for six times in 0.2% PBST (10 minutes/wash) and dehydrated in a

series of ethanol:PBS solutions (10 minutes each in 30%, 50%, 70%, 95%, 100%, 100% solutions).

Then, they were either stored overnight at -20˚C or directly mounted as follows. To mount the

brains, they were cleared in methyl salicylate (removing the ethanol solution) and seated in wells

of stainless-steel slides containing methyl salicylate. Brains were imaged on an Olympus FV-3000

laser scanning confocal microscope. Whole brain images were taken with either a 20X (NA = 0.8)

or a 40X (NA = 0.95) objective, in z-steps of 2 µm thickness. Close ups of the right mushroom body

calyces (both lateral and medial) were taken with an oil immersion 100x objective, in z-steps of

0.5 µm thickness. We selected regions where the mushroom body calyces and the peduncle was

visible.

Neuroanatomic analyses

<u>Volumetry</u>

We aimed to quantify relative volumes of the following functionally distinct brain neuropils (Ito

et al., 2014): the optic lobe (with lamina excluded - because it was not always present after

dissection), the antennal lobe, the central complex (including central body, protocerebral bridge

and nodes), and the mushroom body, where both the medial and lateral calyces, the peduncle,

and lobules were separately identified. To avoid possible human bias and to save labelling time,

a statistical template of *L. humile* brains was created by co-registering ten of the extracted brains.

For these ten brains, the mentioned functional neuropils and the rest of the undifferentiated

central brain were manually labelled in Amira (FEI – Amira 3D 2022.2) in the right hemisphere of

each brain (only the central complex was labelled in both hemispheres, but only half of its volume

was considered in the analyses). These manually obtained labels were used to create consensus

labels in the template, allowing for automatic labelling of the rest of the registered brains.

Automatic labels were smothered with Amira and manually corrected when needed. To calculate

relative volumes of each neuropil, we calculated the volume of each region and of the whole

brain in Amira (module Material Statistics) and computed the ratio between them. The ratio

between primary (sensory) and secondary (processing) neuropils was calculated by dividing the

volume of the optic lobes and antennal lobes by the volume of the mushroom bodies and the

central complex.

Template construction and automatic labelling of brains

Following Arganda-Carreras et al. protocol (2017, 2018), the *L. humile* brain template was created

using diffeomorphic space. Diffeomorphisms, which are smooth and invertible mappings of 3-D

images, are commonly utilised in computational neuroimaging due to their topological properties

that facilitate brain co-registration and the measurement of structural variation (Arganda-

Carreras et al., 2017; Klein et al., 2009). The brain template was constructed from 3-D average-

shape brains derived from the co-registration of 10 original brains obtained through confocal

scans.

Initially, variations in illumination among the original brain images were corrected using

histogram matching (Yoo, 2004). Co-registration involved an initial affine transformation to

maximise mutual information between brain volumes, identifying common neuropil shapes. This

was followed by iterative refinement using local non-rigid transformations to maximise the cross-

correlation of voxel intensities among the co-registered brains. The final average shapes were

generated by computing the voxel-wise median of the co-registered brains.

After creating the anatomical template, we applied the same diffeomorphic transformations to

each label image of the original 10 brains, as performed on their corresponding brain anatomy.

This was followed by per-voxel majority voting across all deformed label images of the same brain

to produce consensus labels.

To automatically label new brain images, we registered individual brain images against the

template using the same two-step method described above: an initial affine registration

maximising mutual information, followed by a non-rigid registration optimising cross-correlation.

The inverse transformations were then applied to the template's regional labels, automatically

assigning label values to the individual grey value brain images registered against the template.

All steps were implemented using the Advanced Normalisation Tools (ANTs) software (Avants et

al., 2011) after converting grey and label images to the open NRRD format in Fiji (Schindelin et

al., 2012).

<u>Microglomeruli</u>

Microglomeruli numbers were counted manually in close-ups of the right mushroom body

calyces using Amira. Close-up images of the right mushroom body calyces were analyzed to

manually measure the number of microglomeruli. Using Amira, three cylindrical regions (each

spanning 5 µm along the z-axis) were selected for analysis within each calyx: one in the collar and

two in the lip. For the lip, regions were differentiated into the non-dense (volume: 256.288 µm³)

and dense areas (volume: 97 µm³). The medial region of the calyx was located using the

pedunculus as a reference, and circles were drawn on ten consecutive slices in this region.

Microglomeruli contours were analyzed in slice 5 of the ten marked slices. A masking threshold

of 700 was applied in Amira to identify microglomeruli contours, and the generated labels were

smoothed using the Smooth Label tool (size: 3). This step helped standardize the contours of the

microglomeruli within the cylindrical regions.

Images of slice 5, with the contours generated in Amira, were then imported into ImageJ for

counting. Using the Multi-Point Tool in ImageJ (Schindelin et al., 2012), microglomeruli were

marked within the defined contours. For validation, 3-D images in Amira were reviewed to

confirm that all labeled contours corresponded to microglomeruli.

Statistical analyses

All analyses were performed using R version 4.4.2 (www.r-project.org). Assumptions of normality

were checked for every variable using the *descdist* function. Non-parametric analyses were used

instead of data transformation. When comparing means, we used ANOVA (*aov*), Kruskal-Wallis

(*kruskal.test*), or Wilcoxon tests (*wilcox.test*) considering the dependency of the variables. Mean

values are shown with standard errors (x̄±SE). When comparing regression lines, we used Linear

Mixed Models (LMMs) for models following Normal distribution (*lmer*) and Generalised Linear

Mixed Models (GLMMs) using Template Model Builder (*glmmTMB*) for those following Negative

Binomial (left-skewed and overdispersed continuous events) and Beta distribution (skewed

continuous proportions variables ranged between 0 and 1). Knowing the dependency on body

size of locomotion in ants (*e.g.*, Hurlbert et al., 2008), we used eye span as a proxy of body size

as a covariable for all the analyses. We tested the dependency of the measured behavioural

variables on body size (covariable) and across observations (fixed factor). We added the nest

source and the ant worker identification code as random factors to take into account the

potential variability among workers from different colonies (*e.g.* see colony personalities in

Pinter-Wollman, 2012; behavioural differences in the populations surrounding the selected

populations in Sanmartín-Villar et al., 2022) and to avoid pseudoreplication (individual repeated

measures across observations).

To analyse the underlying mechanisms driving the measured variables on the three behavioural

tests, we first analysed the effect of the cited factors on the locomotive traits: the average moving

speed (calculated by discounting stationary events) and the proportion of time moving on each

observation. All the tracks were smoothed on AnimalTA with Salvitzky-Golay filter with window

length 15 and polyorder 2 to reduce the noise of the tracks. We considered the ant moving if its

speed was above 2mm/s.

We used exploration (percentage of the total area visited in 10 min) as a variable to analyse the

ant efficiency (or willingness) (i) to cover an unknown area (open field test), (ii) to cover a known

area (after 10 min of acclimatisation) under the presence of a new object (object exposure test),

and (iii) to discover unvisited paths (maze test). Based on the measurements of 25 ants' sizes, we

considered that ants were exploring an area of 5 mm^2 around them for these exploration

analyses. The exploration and the use of the inner part of the arena (38.48 mm^2) were compared

between experimental and control individuals by comparing the variables obtained in the open

field (10 first min of observation) and the exposure to the object (10 next min) with the first 10

min and next 10 min of observation in a control open field.

We used the proportion of time spent interacting with the object as a proxy of

neophobia/neophilia. We used the relative number of choices allowing maze exploration (those

following the concentric path; after now, "ratio of choices to reward") as a proxy of ant

exploration in complex scenarios. We consider that the ant made a choice if it crossed both the

projections on each Y junction of the maze. To calculate the ratio of choices to reward, we first

divided the total number of choices allowing maze exploration by two to avoid considering the

data of individuals crossing backwards a choice point. In the case of obtaining decimal values, the

numbers were rounded to the next superior complete digit. For instance, a value of three units

meant that one ant crossed three times the same choice, two forward and one backwards. By

dividing by two (3/2= 1.5) and rounding to the next superior complete digit we obtain the number

of forward crosses (2). Later, we divided the outcome variable by the total number of choices

(total number of choices allowing maze exploration plus the total number of choices driving to a

dead-end segment). We analysed the direction (left or right) of the first path selected in the maze

to disentangle if this selection was conditioned by (i) lateralization preference (*e.g.* trend to select

left or right paths; see Wystrach et al., 2016); (ii) previous choices performed in the previous

observation (*e.g.* starting to the same direction as in the previous observation); (iii) the openness

of the path (assuming that focal ants were able to perceive the end of the path in the selection

moment, they could have a preference to select a path that allows continuing exploring or going

to a dead-end path).

To analyse if individuals showed different behavioural patterns across time (intraindividual

differences in behavioural plasticity) and if these patterns differed among individuals

(interindividual differences in behavioural plasticity) we compared the AIC values of models with

and without observations as random slopes for each individual intercept (see Sanmartín-Villar et

al., 2021). AIC values are shown as the difference (Δ) between $AIC_{without\ random\ slopes}$ - $AIC_{with\ random}$

$_{slopes}$. Models with random slopes were only selected under ΔAIC positive values higher than 2

units (the rule of thumb when selecting AIC models, *e.g.* Mazerolle, 2006). The marginal (R^2m;

variance explained by the fixed factors) and conditional (R^2c; variance explained by the full

model) coefficient of determination are given for the most explanatory model (lower AIC value).

To analyse the consistency of behaviours, we analyse unpredictability, repeatability, and

coefficients of variation. We calculated unpredictability following the residual individual standard

deviation (riSD), a standard deviation that "estimates variation around the expected values at

every time step instead of estimating the amount of variation around the average of all of the

observations" (Stamps et al., 2012). We selected this method among others (see Cleasby et al.,

2015) because its simplicity and the similarity of the produced results respect other methods that

we found in previous experiments (e.g., Sanmartín-Villar et al., 2021).

We calculated repeatability using the *rptGaussian* function (*rptR,* Stoffel et al., 2017; Suppl.

Material), which is based on the Intraclass Correlation Coefficient (see Nakagawa & Schielzeth,

2010). Due to criticisms of using this coefficient without taking into account the measurement

errors (see Dingemanse et al., 2022), we also calculated the coefficient of inter- and

intraindividual variation (formulae 4 and 5 in Dingemanse et al., 2022) for the behaviours that

we consider more explanatory of each test (exploration in opened field test, object interaction in

object exposure test, and the ratio of choices to reward in the maze test).

Behavioural syndromes (the correlation of personality traits; Bell, 2006) were analysed by adding

one behavioural measure as variable and other as covariable of GLMMs. We compared the

correlation between exploration in open fields and the interaction with the new objects and both

of them with the maze participation (ant willingness to enter into the maze after being added in

the starting area), the ratio of choices to reward, and the number of times that ants reached the

end of the maze.

To test the correlation between behaviours dependent of different information processing (see

Carere & Locurto, 2011), we compared the proportion of time spent on the object for the first

time (neophilia/neophobia) with the total number of choices and the ratio of choices to reward

in the fourth maze test (maze performance after maze flip).

The behavioural relationship with neuropils relative volume and number of microglomeruli was

analysed using MANOVA tests (*manova*), adding the behavioural means or variances as

dependent variables and the neuropils relative volume or the number of microglomeruli from

the measured brain areas as independent variables. We analysed by the same method the effect

of worker size on the neuropil's relative volumetry and number of microglomeruli. We also

performed Principal Components Analysis (PCA) to analyse which variables explain most of the

variance in behaviour, neuropils relative volume and number of microglomeruli. We then used

the resulting PCs to compare behaviour with the two brain traits. We could not compare the PCs

from the two brain measures in the same analyses because of the quality of the images (some

good for volume analyses but not for microglomeruli and *vice versa*).

Results

<u>Behavioural variability</u>

Test performance and individual differences

Body size affected locomotive traits. Larger ants were faster and spent more time moving in the

behavioural tests (Table 1). The locomotive traits changed across observations. The ants' average

speed decreased across observations, but the proportion of time spent moving showed different

patterns across time: they did not change in the open field, increased in the object exposure test,

and decreased in the maze (Table 1).

Workers' exploration depended on body size but showed similar and consistent interindividual

patterns across time. Larger ants explored a larger proportion of area in the open field and the

maze, but arena exploration was not dependent on ant size in the object exposure test (Table 1).

Individual's exploration pattern was constant through observations (no intraindividual

differences) in the open field (60.60±0.69% of the area) and maze (26.81±0.74%) but it showed

a marginal increase in object exposure tests (1st observation: 53.22±1.71%, 5th observation:

57.71±1.67%; Table 1). In the three behavioural tests, the exploration pattern through

observations showed no differences among individuals (no interindividual differences; Table 2;

Figure 1C).

The addition of the object represented a change in the exploration of the arena's areas and its

interaction highlighted different behavioural patterns. Smaller ants used more the inner zone of

the experimental dish when the object was absent (open field and control group in the object

exposure test) but body size did not affect the interaction with the object in the object exposure

test (Table 1). On the first observation, ants were more attracted to the new object (9.84±1.21%

of the time interacting) than to the inner area of the dish in the open field during the 10 previous

minutes (5.67±0.97%; V= 7816, p= 0.003). The proportion of time spent in the inner zone of the

dish by the control group (object absent) was even lower (2.73±1.58%), but no statistical

differences were found with the experimental group due to the sample size (W= 686, p= 0.258).

The time spent interacting with the object increased through observations (5th observation:

18.28±1.24%; F= 25.13, *p*< 0.001). This increase was more pronounced than the increase of the

use of the inner zone in the open field or for control ants in the object exposure test (object

absent; 5[th] observation < 6.17±10.82%). The increase of the proportion of time interacting with

the object across observations differed among ants (interindividual differences; Table 2; Figure

1D). Unexpectedly, ants exposed to the object (focals) modified the use of the inner zone of the

dish in the open field (object absent) through observations, while those never exposed (controls)

did not change it (Table 1). After 10 min of test, *i.e.,* in the object exposure test, control individuals

(object absent) also increased the use of the inner zone of the dish though observations (Table

1), although this increase was lower than the observed in the experimental group and in the open

field test.

Focal workers showed different strategies toward the rewarded path. The ratio of choices for

rewarded path differed among focal workers (interindividual differences) both before

(28.26±1.16%; Figure 1E) and after flipping the maze direction (27.51±1.56%; Figure 1F).

Individual choices followed such different trends that it was not possible to statistically detect an

overall effect across observations (F< 0.20, *p*> 0.689). Ant size did not affect the ratio of choices

for rewarded path (Z= -0.35, *p*= 0.725). More detailed information about maze performance and

strategies of path selection are explained in Supplementary Material.

Behavioural syndromes

Behaviours measured in different tests were correlated within individuals. Ants that explored

more in the open field spent more time interacting with the new object (Table 1). This correlation

was not affected by the observation number (Table 1). Ants that explored more in the open field

participated more in the maze exploration (t= -4.37, df= 149, p< 0.001) and reached more times

the reward (χ^2_2= 18.18, p< 0.001), but they did not reach faster the reward of the maze after

having reached it once (Z= 0.61, p= 1). Ants' exploration or object interaction were not correlated

with the ratio of choices to reward (Table 1).

We found no behavioural syndromes involving different information processing. The first

interaction with the object was not correlated with maze participation ($F_{1;149}$= 0.075, p= 0.785)

or with the ratio of choices to reward of the fourth observation (first encounter of the flipped

maze) (S= 627247, p= 0.915, r= 0.01).

Unpredictability

Focal workers behavioural unpredictability was independent in the three behavioural tests, and

it was not affected by their behavioural performance. No correlations were found between the

unpredictability of the exploration in open field, object interaction, and the ratio of choices to

reward (t< [0.51], df= 154, p> 0.635). No differences in unpredictability were found between

control and focal individuals for their exploration in the open field ($F_{1;165}$= 0.12, p= 0.731). The

introduction of a new object decreased the unpredictability of the ants' use of the inner part of

the arena when the object was present - or after 10 min of test for the control group - (control-

focal: $F_{4;165}$= 4.75, p= 0.031), while the unpredictability to use the inner part did not change in

the exploration on the open field test ($F_{4;165}$= 0.12, p= 0.731). No differences in unpredictability

for the object interaction were found for individuals increasing or decreasing the interaction over

the observations (t= -0.53, df= 154, p= 0.592). The unpredictability of the ratio of choices to

reward was similar for those individuals increasing or decreasing this ratio considering all

observations or only those before maze flip (t< 1.36, df= 137.04, p> 0.177). Ant's exploration and

the ratio of choices to reward were similarly unpredictable according to the ant size (Table 1),

but the interaction with the new object was less unpredictable in larger ants (Table 1).

Coefficients of variation

The behavioural variability observed among focal and control ant workers was observed in the

object exposure tests, but not in the open field. The coefficient of intraindividual variation for

exploration was similar among control (21.52±3.23%) and focal ants (23.80±1.09%; $F_{1;165}$= 0.30,

p= 0.586). No differences were found for worker size (t= -1.63, p= 0.193). The coefficient of

intraindividual variation for the interaction with the inner area differed between control (without

object; 0.91±0.09%) and focal ants (with object; 13.67±0.80%; $F_{1;164}$= 5.40, p= 0.021) in the object

exposure test. Smaller ants showed higher variation (t= 12.70, p< 0.007). The coefficient of

intraindividual variation for the ratio of choices to reward (23.06±0.92%) did not depend on

worker size (t= -1.41, df= 154, p= 0.162).

The coefficient of interindividual variation was similar among the control (33.35%) and the focal

group (20.03%; Z= 1.31; critical value= 1.96) for exploration but strongly differed for the

interaction with the object (control: 33.16%; focal: 75.57%; Z= -3.63; critical value= 1.96). Ants

showed an interindividual variation of 43.17% for the ratio of choices to reward.

<u>Neuroanatomy</u>

Volumetry

The relative volume of the cited functional neuropils was only analysed in 38 focal workers (daily

exposed to the three behavioural tests over five days) due to logistics. Unfortunately, the relative

neuropils volumes of the control group (not exposed to behavioural tests) could only be analysed

in two individuals and were therefore discarded.

The relative volumes of the different neuropils were uncorrelated in most cases. No significant

correlation was found between the body size and the relative volume of the brain neuropils

measured ($F_{1;30}$ < 2.21, p> 0.148). The relative volumes of the medial and lateral mushroom body

calyces were positively correlated (t= 5.09, df= 36, p> 0.001, r= 64.68%). The relative volume of

the central complex and the optic lobe was negatively correlated (t= -2.39, df= 36, p= 0.022). No

correlations were found between the relative volume of (i) the optic and antennal lobes, (ii) the

central complex and the mushroom body, (iii) and the mushroom body and the sensory (optic or

antennal) lobes (t< [1.37], p> 0.181, r< 22.19%).

Microglomeruli

The number of microglomeruli of the analysed regions (collar, dense lip, non-dense lip) were

positively correlated (t > 3.75, p < 0.001, r > 54.65). The number of microglomeruli present in the

collar showed a positive correlation with the body size ($F_{1;24}$= 5.42, p= 0.029). No significant effect

of worker size was found for the microglomeruli of the lip ($F_{1;24}$< 0.69, p= 0.414).

Behaviour-brain correlation

The relative volume of the neuropils conditioned the behavioural performance in complex

environments. Ant workers with higher relative optic lobe volumes moved faster in the presence

of the new object ($F_{1;29}$= 7.43, p= 0.011; Figure 2A). Ant workers with lower primary:secondary

neuropils ratio showed higher exploration when exposed to the object ($F_{1;29}$= 4.84, p= 0.036;

Figure 2B) and to the maze ($F_{1;29}$= 5.30, p= 0.029; Figure 2C). No other significant correlations

were found between the neuropils' relative volume and the mean and variance of speed or

exploration in the three behavioural tests performed ($F_{1;29} < 3.02$, $p = 0.093$). The regression slopes

of the ratio of choices to reward before the maze flip were negatively correlated with the relative

volume of the medial mushroom body calyx ($F_{1;29} = 7.65$, $p = 0.010$; Figure 2D; but not with the

lateral calyx: $F_{1;29} = 2.68$, $p = 0.113$) and positively correlated with the relative volume of the

central complex ($F_{1;29} = 10.98$, $p = 0.002$; Figure 2E). No significant differences were found when

comparing behavioural and brain volumetry PCs ($F_{1;35} < 1.73$, $p > 0.196$).

The number of microglomeruli was similar among control and focal ants and among those

showing different mean behaviours. No differences were found between the number of

microglomeruli of individuals exposed (N= 38) or not exposed (control; N= 5) to behavioural tests

in any of the three analysed brain regions ($F_{1;38} < 1.64$, $p > 0.209$), but a trend suggested that

control individuals showed higher variance than the focal ones in the number of microglomeruli

present in the collar ($F_{1;38} = 3.90$, $p = 0.056$). The number of microglomeruli of the three regions

analysed was not correlated with the average speed ($F_{1;24} < 1.20$, $p > 0.334$) or the average

exploration ($F_{1;24} < 1.17$, $p > 0.345$) performed on the three behavioural tests. Nor were

correlations found with the mean object interactions and the mean ratio of choices to reward

before and after maze flip ($F_{1;24} < 1.79$, $p > 0.189$).

However, the number of microglomeruli differed among individuals showing different

behavioural patterns under complex environments. The number of microglomeruli of the three

regions analysed was not correlated with the variance of speed ($F_{1;24} < 1.20$, $p > 0.334$) or

exploration ($F_{1;24} < 1.17$, $p > 0.345$) performed on the three behavioural tests. Individuals with

higher number of microglomeruli on the dense lip showed higher variance ($F_{1;24} = 6.50$, $p = 0.018$;

Figure 3A), more unpredictability (riSD: $F_{1;24}$= 7.87, p= 0.010; Figure 3B) on the object interaction,

and higher increase of the ratio of choices to reward before maze flip ($F_{1;24}$= 9.10, p= 0.006; Figure

3C). No correlations were found between the object interaction variance and the number of

microglomeruli from the collar or non-dense lip ($F_{1;24}$< 1.23, p> 0.278), nor between the slopes

of the ratio of choices to reward before maze flip and the non-dense lip ($F_{1;24}$= 1.01, p= 0.326).

Individuals with higher numbers of microglomeruli in the collar showed decreased slopes of the

ratio of choices to reward before maze flip ($F_{1;24}$= 5.81, p= 0.024; Figure 3D). The variance after

maze flip was not correlated with the number of microglomeruli in any region ($F_{1;24}$< 0.85, p>

0.367). No significant differences were found when comparing behavioural and microglomeruli

PCs ($F_{1;13}$< 1.18, p> 0.339).

Discussion

The behavioral variability observed at the individual level suggests two mechanisms by which

Linepithema humile colonies manage to explore new environments: through efficiency

associated with body size and neural processing areas. On one hand, larger ants showed higher

speed, movement, exploration and more microglomeruli on the collar (visual perception) while

they did not show larger relative neuropil volumes. Smaller ants explored the arena differently,

visiting its inner zone more frequently. Behavioural differences linked to body size could increase

the efficiency of colony exploration due to different trial-and-error strategies. On the other hand,

individuals who invested more in processing than in perception were more efficient at exploring

complex environments, especially those who processed olfactory rather than visual stimuli.

Complex scenarios resolution may be favoured by this small number of individuals in whom the

correlation between behaviour and neuroanatomy was observed (see dots outside the dashed

line in Figure 2D-E, Figure 3C-D). This possible evolutionary strategies supports the hypothesis of

the colony's investment reduction in the behavioural and neural plasticity of its members in *L.*

humile (e.g., Sanmartín-Villar et al., 2021), limiting costs to cope with complex environments

through a neural demanding mechanism in only a few individuals, while in a non-neural

demanding mechanism in most them. Our results suggest that exploration at the colony level

might depend on the variability among-individuals, but interestingly, in the case of our study this

variability was not dependent on the individual adult development - as division of labour in

eusocial animals is often explained - because we analysed workers of the same age after

emergence. Future studies should corroborate in the field the consequences of the mechanisms

proposed here for local adaptation, invasiveness, and coping with climate change.

The expected similar behaviour among workers and lack of correlation between behaviour and

neuroanatomical traits were only observed at low environmental complexity. Individuals showed

different and dependent responses of their neuroanatomy when exposed to complex

environmental challenges, such as novel experiences and choices. Sensory neuropils hardly

affected behavior, only detecting faster exploration after object exposure in individuals with

larger visual lobes which support the navigation improvement through volume increase of

primary lobes as seen in other species of ants (*Cataglyphus nodus*; Grob et al., 2017, *Cataglyphis*

fortis; Schmitt et al., 2016, *Camponotus rufipus*; Yilmaz et al., 2017). Processing areas (central

complex and mushroom bodies) determined exploration and maze progress. Progress in the

maze was positively dependent on the number of synaptic clusters in the dense lip (olfactory

processing), but negatively dependent on those in the collar (visual processing). Individuals who

showed more dense lip synaptic clusters showed higher behavioral variability (variance and

unpredictability). These results highlight the relevance of high processing areas in the resolution

of complex environments and the different functions of different brain regions. Specialized

behavioral strategies could be expected from a higher neuropil volume due to the development

of synaptic clusters, but also a lower number of these clusters due to pruning (the concentration

of microglomeruli in a more volumetric button due to specialization; Changeux and Danchin,

1976; Cowan et al., 1984). The correlations found between exploration efficiency and the number

of microglomeruli could highlight the diversity of behavioral strategies (variance,

unpredictability) rather than learning processes, although they coil highlight long term memory

formation due to the increase of microglomeruli density (Falibene et al., 2015).

In contrast to the expected, *L. humile* workers showed low percentage of exploratory

performance, exploring an average of 60% ca. of the open field, interacting with the object up to

18% of time, and selecting rewarded paths in 27% ca. of cases. Although *L. humile* is considered

a monomorphic species, body size conditioned behavioural and neural traits. The allometry for

speed and body size is a common phenomenon in ants due to the higher motion of individuals

with longer legs and higher metabolic rate (Hurlbert et al., 2008), but we did not expect to see

such a strong difference within a monomorphic species. Assuming size-dependent differences,

we expected that body size correlate with risk-taking decisions as it occurs in fish (e.g. Darby &

McGhee, 2019; but see Brown & Braithwaite, 2004), turtles (Ibáñez et al., 2013), or snakes (Mayer

et al., 2016). Therefore, we expected a body size dependency on the interaction with the novel

object (but see the mismatch between neophobia and boldness in Carter et al., 2012), the use of

the inner part of the arena (assumed as a bolder trait for some authors; Carlson & Langkilde,

2013; Detrain et al., 2019; Sneddon, 2003; Valle, 1970; Walsh & Cummins, 1976), and the

participation on the maze. However, none of these traits correlated with body size, suggesting

that *L. humile* workers may not evaluate the environment challenges according to their body

characteristics.

The interaction with the object was less variable and unpredictable for larger workers, suggesting

a more homogeneous behaviour. In addition, larger workers reached more times the reward

inside the maze but decreased their ratio of choices to reward through observations. Although

we could not determine the overall volume changes in the brains based on experience (see

Gronenberg et al., 1996) due to our methodology, our results support the relationship between

body size, brain development and foraging behaviour shown on *Atta cephalotes* (Arganda et al.,

2020), highlighting our findings that larger individuals were better in simple tasks like exploration

and navigation but not in complex tasks like in the maze and also being associated with larger

volume ratio of primary to secondary neuropil. All these results suggest that large workers

succeeded to better solve environmental complexity (higher exploration and maze-solving, more

constant object interaction) due to their higher locomotive traits, but not due to the

neuroanatomy that might underlie their cognitive skills (no correlation was found between speed

or exploration or body size and neuropil relative volumes or microglomeruli abundance). In fact,

our results suggest that large workers only had more microglomeruli in the collar than smaller

workers (therefore, no more microglomeruli on the lip than smaller workers). We consider

improbable that larger workers could be able to better solve the tests in which they were exposed

due to their higher secondary processing of visual stimuli in comparison with the chemical stimuli

(more microglomeruli on the collar, but no more on the lip; Falibene et al., 2016; Gordon et al.,

2018, 2019; Gordon & Traniello, 2018; Seid & Wehner, 2008) due to the lack of visual references

and similitude among choices. In any case, our results support the idea that body size can

influence environmental exploration in monomorphic species as it was shown in other species

such as *Lasius niger* (*e.g.*, Grześ et al., 2016; Okrutniak et al., 2020). Therefore, future behavioural

studies should control for the effect of body size in monomorphic ant species but also analyse

whether worker body size is related to the spatial complexity of the environment as an

evolutionary strategy.

Focal workers acclimatised to the experimental setup, reducing their average speed, increasing

the interaction with the object and decreasing their mobility in the maze over the observations.

This suggests that ants were able to modify their behaviour according to previous experiences,

as was previously shown in ants of the same species (Wagner et al., 2023). However, we cannot

discard that the obtained outcomes depend on other factors such as fatigue or developmental-

related changes. Previous works done with repeated observations on open fields showed

different patterns (reduction of locomotion in a ten repetition test in *L. niger*, Sanmartín-Villar et

al., 2021; increase of locomotion in a five repetition test in *L. humile*, Sanmartín-Villar et al.,

2021). If ants recognise the object from the previous tests, the object introduced could only be

considered as novel on the first observation. Despite the lack of novelty and apparently lack of

functionality of the object (except for climbing), the focal workers interacted more with it

through the observations, especially the workers that showed higher exploration, while the

opposite happened in the maze, in which workers could find a reward. Note that we did not

starve focal individuals, and a lack of food motivation could interfere with performance in the

maze. Still, we expected that ants would increase their maze performance while losing the fear

to navigate through it and especially after the reduction of their speed through observations,

which may could be due to the speed-accuracy trade-off and its dependency on neural structures

(reviewed in Standage et al., 2014).

The increase of the environmental complexity through the addition of an object in a relatively

known arena (experienced through 10 minutes) triggered behavioural differences among

workers and modified the apparently characteristic homogeneous locomotion pattern of the *L.*

humile workers in open fields (also seen in Sanmartín-Villar et al., 2021). The unpredictability of

the object interaction, *i.e.*, its deviation through the expected pattern, correlated with the neural

area in charge of processing chemical stimuli (dense lip) and the mean and variance of the object

interaction correlated with the relative volume of the central complex, which is in charge of

navigation (Honkanen et al., 2019; reviewed in Webb & Wystrach, 2016). The increase of the

object interaction and movement (marginally exploration) could fit with the expected result of

environmental enrichment (reviewed in Newberry, 1995), in which individuals improve their

cognitive skills after the addition of objects. Individuals exposed to the object used more the

inner zone of the area in posterior tests when the object was not present while control individuals

showed this pattern with a delay of 10 min, suggesting an improvement of the area exploration

after previous stimuli. However, we cannot discard that the behaviours observed were caused

by the stress produced when adding a new element in a known area. Considering the correlation

trend of the object interaction with the central complex (navigation centre), we propose a third

explanation in which workers used the object as a spatial reference to which they recurred to

orient in an arena lacking other spatial clues.

The difficulty of the maze test highlighted the relation between neuroanatomy and behaviour.

Within the mushroom bodies (in charge of information processing, memory and learning), the

maze performance before the flip was only correlated with the relative volume of the medial

calyx, and not with the lateral calyx. As medial and lateral calyces are considered to perform the

same function, our results might be a suggestion for further research trying to disentangle

distinctive information processing among them. The relative volume of the central complex

(navigation centre) also correlates with maze performance before the flip. The difficulty of the

maze due to its design and the lack of references accomplished the aim of exposing ants towards

an exercise focused on exploration rather than learning (we did not train the ants before the

tests). Workers with more microglomeruli in dense lip showed higher ratio of choices to reward

before maze flip, which suggests that the presence of more synaptic routes - but smaller - in this

region responsible for chemical processing allowed individuals to improve their efficiency on the

maze. We discarded that the experience on the maze was responsible for the synaptic cluster

development during the tests because of the similar microglomeruli found between control and

focal workers, although more research must be performed to conclude it. In contrast, the number

of microglomeruli of the collar negatively correlated with the ratio of choices to reward before

maze flip. In addition, the relative volume of the central complex (more relevant in the maze

performance) was negatively correlated with the relative volume of the optic lobes. We interpret

these results as a conflict between the role of primary and higher order processing regions in the

studied behaviours: one region in charge of perceiving and processing of visual stimuli (optical

lobes), and two in charge of the higher order processing of stimuli (the chemical information in

the non-dense lip in the mushroom body calyces, and the visual, chemical and mechanosensory

information required for orientation in the central complex). Individuals that relied more on

visual stimuli were probably not able to disentangle the orientation on the maze due to the exact

symmetry of all the choices. However, individuals relying more on the chemical stimulus could

better interpret their own pheromones or other odors present in the maze (although backwards

behaviour was uncommon). Surprisingly, no correlations were found with the antennal lobes,

which we expected to have a higher relevance in the ants' orientation.

The relevance of the microglomeruli of the dense lip in behaviour (unpredictability of the object

interaction, increase of rewarded path choice) opposes the results found in honey bees (Van Nest

et al., 2017), in which only the microglomeruli of the non-dense lip (also olfactory) were related

with foraging behaviour. Thermal fluctuations affect the microglomeruli density in the non-dense

lip in the mushroom body calyx of *Camponotus mus* but were less evident in dense lip and visual

collar (Falibene et al., 2016). If the correlations between these experiments are comparable, this

would highlight phylogenetic differences in the neuroethology of the species, with behaviour

being dependent on different neural structures across insect and ant species. However, all these

experiments highlight the expected relevance of the olfactory stimulus over the visual one to

drive behaviour.

Our results suggest that individual variability depends on neuronal development and highlight

behavioural plasticity and the strategies to cope with environments with different degree of

challenge in *Linepithema humile*, skills likely related with its adaptation in introduced areas.

Although we could not observe the associated neuronal changes with each exposure to

behavioural tests, likely to decipher in detail, the mechanisms involved in their behavioural

patterns, we provide the first neuroethological clue to understand the invasive strategies of this

species. We attempted to dissect and correlate the behavioural and neuronal plasticity of an

invasive species to cope with novel scenarios, providing a framework for understanding the

dynamics for colony fitness in the introduced areas. We are convinced that the present results

could be the basis for future control and management programs dealing with *L. humile* invasion.

For instance, neural disruptors could affect the behaviour of worker ants differently depending

on their size and on the complexity of their environment.

Acknowledgements

We are grateful to Xoan Sanmartín Pazos for his help in the field work and to the staff of the

Łomna Biological Station (MiIZ) for allowing us to produce uninterrupted work even in national

holidays and heavy snow conditions. This research is part of the project No.

2021/43/P/NZ8/03306 co-funded by the National Science Centre and the European Union

Framework Programme for Research and Innovation Horizon 2020 under the Marie Skłodowska-

Curie grant agreement no.945339. Part of this research was supported by the 2017 program for

attracting and retaining talent of Comunidad de Madrid (2021-5A/BMD-20960). For the purpose

of Open Access, the author has applied a CC-BY public copyright licence to any Author Accepted

Manuscript (AAM) version arising from this submission.

Contributions: M.W. and I.S.V. conceived the ideas and designed the methodology; V.C and I.S.V.

collected the samples; S.N, V.C., and I.S.V conducted the experiments; M.E.V. and S.A. performed

the protocols and conducted the brain imaging; S.N collected the behavioural data; M.E.V., I.A.C.,

and S.A. collected the brain-related data; I.S.V. analysed the data; S.N., M.E.V., S.A., I.A.C., and

I.S.V. wrote the manuscript; M.W., S.A., and I.S.V. lead the funding acquisition, project

administration, and supervision. All authors contributed critically to the drafts and gave final

approval for publication.

References

Abril, S., Oliveras, J., & Gómez, C. (2010). Effect of temperature on the development and survival of the

Argentine ant, *Linepithema humile*. *Journal of Insect Science*, *10*(97), 1–13.

https://doi.org/10.1673/031.010.9701

Angilletta, M. J., Roth, T. C., Wilson, R. S., Niehaus, A. C., & Ribeiro, P. L. (2008). The fast and the fractalous:

Speed and tortuosity trade off in running ants. *Functional Ecology*, *22*(1), 78–83.

Angulo, E., Guénard, B., Balzani, P., Bang, A., Frizzi, F., Masoni, A., Abril, S., Suarez, A. V., Hoffmann, B.,

Benelli, G., Aonuma, H., Lach, L., Mothapo, P. N., Wossler, T., & Santini, G. (2024). The Argentine ant,

Linepithema humile: Natural history, ecology and impact of a successful invader. *Entomologia Generalis*,

41–61. https://doi.org/10.1127/entomologia/2023/2187

Arganda, S., Hoadley, A. P., Razdan, E. S., Muratore, I. B., & Traniello, J. F. A. (2020). The neuroplasticity of

division of labor: Worker polymorphism, compound eye structure and brain organization in the leafcutter

ant *Atta cephalotes*. *Journal of Comparative Physiology A*, *206*(4), 651–662.

https://doi.org/10.1007/s00359-020-01423-9

Arganda-Carreras, I., Gordon, D. G., Arganda, S., Beaudoin, M., & Traniello, J. F. A. (2017). Group-wise 3D

registration based templates to study the evolution of ant worker neuroanatomy. *2017 IEEE 14th*

International Symposium on Biomedical Imaging (ISBI 2017), 429–432.

https://doi.org/10.1109/ISBI.2017.7950553

Arganda-Carreras, I., Manoliu, T., Mazuras, N., Schulze, F., Iglesias, J. E., Bühler, K., Jenett, A., Rouyer, F.,

& Andrey, P. (2018). A statistically representative atlas for mapping neuronal circuits in the *Drosophila*

adult brain. *Frontiers in Neuroinformatics*, *12*. https://doi.org/10.3389/fninf.2018.00013

Aso, Y., Sitaraman, D., Ichinose, T., Kaun, K. R., Vogt, K., Belliart-Guérin, G., Plaçais, P.-Y., Robie, A. A.,

Yamagata, N., Schnaitmann, C., Rowell, W. J., Johnston, R. M., Ngo, T.-T. B., Chen, N., Korff, W., Nitabach,

M. N., Heberlein, U., Preat, T., Branson, K. M., … Rubin, G. M. (2014). Mushroom body output neurons

encode valence and guide memory-based action selection in *Drosophila*. *eLife*, *3*, e04580.

https://doi.org/10.7554/eLife.04580

Avants, B. B., Tustison, N. J., Song, G., Cook, P. A., Klein, A., & Gee, J. C. (2011). A reproducible evaluation

of ANTs similarity metric performance in brain image registration. *NeuroImage*, *54*(3), 2033–2044.

https://doi.org/10.1016/j.neuroimage.2010.09.025

Baratelli, E., Tillberg, C., Suarez, A., Menke, S., Naughton, I., & Holway, D. (2023). Variation in Argentine

ant *(Linepithema humile)* trophic position as a function of time. *Biological Invasions*, *25*(1), 133–140.

https://doi.org/10.1007/s10530-022-02898-0

Bell, A. M. (2006). Future directions in behavioural syndromes research. *Proceedings of the Royal Society*

B: Biological Sciences, *274*(1611), 755–761. https://doi.org/10.1098/rspb.2006.0199

Bernadou, A., Schrader, L., Pable, J., Hoffacker, E., Meusemann, K., & Heinze, J. (2018). Stress and early

experience underlie dominance status and division of labour in a clonal insect. *Proceedings of the Royal*

Society B: Biological Sciences, *285*(1885), 20181468. https://doi.org/10.1098/rspb.2018.1468

Bertelsmeier, C. (2021). Globalization and the anthropogenic spread of invasive social insects. *Current*

Opinion in Insect Science, *46*, 16–23. https://doi.org/10.1016/j.cois.2021.01.006

Beshers, S. N., & Fewell, J. H. (2001). Models of division of labor in social insects. *Annual Review of*

Entomology, *46*(Volume 46, 2001), 413–440. https://doi.org/10.1146/annurev.ento.46.1.413

Bhatkar, A., & Whitcomb, W. H. (1970). Artificial diet for rearing various species of ants. *The Florida*

Entomologist, *53*(4), 229–232. https://doi.org/10.2307/3493193

Brown, C., & Braithwaite, V. A. (2004). Size matters: A test of boldness in eight populations of the poeciliid

Brachyraphis   episcopi.   Animal   Behaviour,   *68*(6),   1325–1329.

https://doi.org/10.1016/j.anbehav.2004.04.004

Carere, C., & Locurto, C. (2011). Interaction between animal personality and animal cognition. *Current*

Zoology, *57*(4), 491–498. https://doi.org/10.1093/czoolo/57.4.491

Carlson, B. E., & Langkilde, T. (2013). Personality traits are expressed in bullfrog tadpoles during open-

field trials. *Journal of Herpetology*, *47*(2), 378–383. https://doi.org/10.1670/12-061

Carpintero, S., & Reyes-López, J. (2008). The role of competitive dominance in the invasive ability of the

Argentine ant (*Linepithema humile*). *Biological Invasions*, *10*(1), 25–35. https://doi.org/10.1007/s10530-

007-9103-3

Carter, A. J., Marshall, H. H., Heinsohn, R., & Cowlishaw, G. (2012). How not to measure boldness: Novel

object and antipredator responses are not the same in wild baboons. *Animal Behaviour*, *84*(3), 603–609.

https://doi.org/10.1016/j.anbehav.2012.06.015

Changeux J-P, Danchin A. 1976. Selective stabilisation of developing synapses as a mechanism for the

specification of neuronal networks. *Nature*, 264:705–712. https://doi.org/10.1038/264705a0

Chiara, V., & Kim, S.-Y. (2023). AnimalTA: A highly flexible and easy-to-use program for tracking and

analysing animal movement in different environments. *Methods in Ecology and Evolution*, *14*(7), 1699–

1707. https://doi.org/10.1111/2041-210X.14115

Cleasby, I. R., Nakagawa, S., & Schielzeth, H. (2015). Quantifying the predictability of behaviour: Statistical

approaches for the study of between-individual variation in the within-individual variance. *Methods in*

Ecology and Evolution, *6*(1), 27–37. https://doi.org/10.1111/2041-210X.12281

Cowan W, Fawcett J, O'Leary D, Stanfield B. 1984. Regressive events in neurogenesis. *Science* 225:1258–

1265. https://doi.org10.1126/science.6474175

Csata, E., Gautrais, J., Bach, A., Blanchet, J., Ferrante, J., Fournier, F., Lévesque, T., Simpson, S. J., &

Dussutour, A. (2020). Ant foragers compensate for the nutritional deficiencies in the colony. *Current*

Biology, *30*(1), 135-142.e4. https://doi.org/10.1016/j.cub.2019.11.019

Czaczkes, T. J. (2018). Using T- and Y-mazes in myrmecology and elsewhere: A practical guide. *Insectes*

Sociaux, *65*(2), 213–224. https://doi.org/10.1007/s00040-018-0621-z

Czaczkes, T. J., Grüter, C., Ellis, L., Wood, E., & Ratnieks, F. L. W. (2013). Ant foraging on complex trails:

Route learning and the role of trail pheromones in Lasius niger. *The Journal of Experimental Biology*,

216(Pt 2), 188–197. https://doi.org/10.1242/jeb.076570

Dall, S. R. X., Houston, A. I., & McNamara, J. M. (2004). The behavioural ecology of personality: Consistent

individual   differences   from   an   adaptive   perspective.   *Ecology   Letters*,   *7*(8),   734–739.

https://doi.org/10.1111/j.1461-0248.2004.00618.x

Darby, N. A., & McGhee, K. E. (2019). Boldness is affected by recent experience with predation cues and

body   size   in   mosquitofish.   *Behavioural   Processes*,   *164*,   143–149.

https://doi.org/10.1016/j.beproc.2019.05.007

Darwin, C., 1809-1882. (1859). *On the origin of species by means of natural selection, or preservation of*

favoured races in the struggle for life. London : John Murray, 1859.

https://search.library.wisc.edu/catalog/9934839413602122

Delgado, J., & Solé, R. V. (2000). Self-synchronization and task fulfilment in ant colonies. *Journal of*

Theoretical Biology, *205*(3), 433–441. https://doi.org/10.1006/jtbi.2000.2077

Detrain, C., Pereira, H., & Fourcassié, V. (2019). Differential responses to chemical cues correlate with task

performance in ant foragers. *Behavioral Ecology and Sociobiology*, *73*(8), 107.

https://doi.org/10.1007/s00265-019-2717-5

Dewitt, T. J., Sih, A., & Wilson, D. S. (1998). Costs and limits of phenotypic plasticity. *Trends in Ecology &*

Evolution, *13*(2), 77–81. https://doi.org/10.1016/s0169-5347(97)01274-3

Dingemanse, N. J., Hertel, A. G., & Royauté, R. (2022). Moving away from repeatability: A comment on

Stuber et al. *Behavioral Ecology*, *33*(3), 488–489. https://doi.org/10.1093/beheco/arac006

Dingemanse, N. J., Kazem, A. J. N., Réale, D., & Wright, J. (2010). Behavioural reaction norms: Animal

personality meets individual plasticity. *Trends in Ecology & Evolution*, *25*(2), 81–89.

https://doi.org/10.1016/j.tree.2009.07.013

Dussutour, A., Deneubourg, J.-L., & Fourcassié, V. (2005). Amplification of individual preferences in a social

context: The case of wall-following in ants. *Proceedings of the Royal Society B: Biological Sciences*,

272(1564), 705–714. https://doi.org/10.1098/rspb.2004.2990

Fahrbach, S. E., Moore, D., Capaldi, E. A., Farris, S. M., & Robinson, G. E. (1998). Experience-expectant

plasticity in the mushroom bodies of the honeybee. *Learning & Memory*, *5*(1), 115–123.

https://doi.org/10.1101/lm.5.1.115

Falibene, A., Roces, F., & Rössler, W. (2015). Long-term avoidance memory formation is associated with a

transient increase in mushroom body synaptic complexes in leaf-cutting ants. *Frontiers in Behavioral*

Neuroscience, *9*. https://doi.org/10.3389/fnbeh.2015.00084

Falibene, A., Roces, F., Rössler, W., & Groh, C. (2016). Daily thermal fluctuations experienced by pupae via

rhythmic nursing behavior increase numbers of mushroom body microglomeruli in the adult ant brain.

Frontiers in Behavioral Neuroscience, *10*. https://doi.org/10.3389/fnbeh.2016.00073

Gordon, D. G., & Traniello, J. F. A. (2018). Synaptic organization and division of labor in the exceptionally

polymorphic   ant   *Pheidole   rhea*.   *Neuroscience   Letters*,   *676*,   46–50.

https://doi.org/10.1016/j.neulet.2018.04.001

Gordon, D. G., Zelaya, A., Arganda-Carreras, I., Arganda, S., & Traniello, J. F. A. (2019). Division of labor

and brain evolution in insect societies: Neurobiology of extreme specialization in the turtle ant *Cephalotes*

varians. *PLOS ONE*, *14*(3), e0213618. https://doi.org/10.1371/journal.pone.0213618

Gordon, D. G., Zelaya, A., Ronk, K., & Traniello, J. F. A. (2018). Interspecific comparison of mushroom body

synaptic complexes of dimorphic workers in the ant genus *Pheidole*. *Neuroscience Letters*, *662*, 110–114.

https://doi.org/10.1016/j.neulet.2017.10.009

Grob, R., Fleischmann, P. N., Grübel, K., Wehner, R., & Rössler, W. (2017). The role of celestial compass

information in *Cataglyphis* ants during learning walks and for neuroplasticity in the central complex and

mushroom bodies. *Frontiers in Behavioral Neuroscience*, *11*. https://doi.org/10.3389/fnbeh.2017.00226

Gronenberg, W., Heeren, S., & Hölldobler, B. (1996). Age-dependent and task-related morphological

changes in the brain and the mushroom bodies of the ant *Camponotus floridanus*. *Journal of Experimental*

Biology, *199*(9), 2011–2019. https://doi.org/10.1242/jeb.199.9.2011

Grześ, I. M., Okrutniak, M., & Grzegorzek, J. (2016). The size-dependent division of labour in monomorphic

ant *Lasius niger*. *European Journal of Soil Biology*, *77*, 1–3. https://doi.org/10.1016/j.ejsobi.2016.08.006

Harris, S., Ramnarine, I. W., Smith, H. G., & Pettersson, L. B. (2010). Picking personalities apart: Estimating

the influence of predation, sex and body size on boldness in the guppy *Poecilia reticulata*. *Oikos*, *119*(11),

1711–1718. https://doi.org/10.1111/j.1600-0706.2010.18028.x

Hill, L., Gérard, M., Hildebrandt, F., & Baird, E. (2023). Bumblebee cognitive abilities are robust to changes

in colony size. *Behavioral Ecology and Sociobiology*, *77*(2), 25. https://doi.org/10.1007/s00265-023-

03299-6

Hölldobler, B., & Wilson, E. O. (1990). *The Ants*. Harvard University Press.

Honkanen, A., Adden, A., Freitas, J. da S., & Heinze, S. (2019). The insect central complex and the neural

basis of navigational strategies. *The Journal of Experimental Biology*, *222*(Pt Suppl 1), jeb188854.

https://doi.org/10.1242/jeb.188854

Human, K. G., & Gordon, D. M. (1996). Exploitation and interference competition between the invasive

Argentine ant, *Linepithema humile*, and native ant species. *Oecologia*, *105*(3), 405–412.

https://doi.org/10.1007/BF00328744

Hurlbert, A. H., Ballantyne Iv, F., & Powell, S. (2008). Shaking a leg and hot to trot: The effects of body size

and temperature on running speed in ants. *Ecological Entomology*, *33*(1), 144–154.

https://doi.org/10.1111/j.1365-2311.2007.00962.x

Ibáñez, A., Marzal, A., López, P., & Martín, J. (2013). Boldness and body size of male Spanish terrapins

affect their responses to chemical cues of familiar and unfamiliar males. *Behavioral Ecology and*

Sociobiology, *67*(4), 541–548. https://doi.org/10.1007/s00265-012-1473-6

Ito, K., Shinomiya, K., Ito, M., Armstrong, J. D., Boyan, G., Hartenstein, V., Harzsch, S., Heisenberg, M.,

Homberg, U., Jenett, A., Keshishian, H., Restifo, L. L., Rössler, W., Simpson, J. H., Strausfeld, N. J., Strauss,

R., & Vosshall, L. B. (2014). A systematic nomenclature for the insect brain. *Neuron*, *81*(4), 755–765.

https://doi.org/10.1016/j.neuron.2013.12.017

Jacobs, D. S., & Jarvis, J. U. M. (1996). No evidence for the work-conflict hypothesis in the eusocial naked

mole-rat (*Heterocephalus glaber*). *Behavioral Ecology and Sociobiology*, *39*(6), 401–409.

https://doi.org/10.1007/s002650050307

Jeanson, R. (2019). Within-individual behavioural variability and division of labour in social insects. *Journal

of Experimental Biology*, *222*(10), jeb190868. https://doi.org/10.1242/jeb.190868

Jeanson, R., & Weidenmüller, A. (2014). Interindividual variability in social insects—Proximate causes and

ultimate consequences. *Biological Reviews of the Cambridge Philosophical Society*, *89*(3), 671–687.

https://doi.org/10.1111/brv.12074

Kamhi, J. F., Ilieş, I., & Traniello, J. F. A. (2019). Social complexity and brain evolution: Comparative analysis

of modularity and integration in ant brain organization. *Brain Behavior and Evolution*, *93*(1), 4–18.

https://doi.org/10.1159/000497267

Kamhi, J. F., Sandridge-Gresko, A., Walker, C., Robson, S. K. A., & Traniello, J. F. A. (2017). Worker brain

development and colony organization in ants: Does division of labor influence neuroplasticity?

Developmental Neurobiology, *77*(9), 1072–1085. https://doi.org/10.1002/dneu.22496

Keller, L., & Fournier, D. (2002). Lack of inbreeding avoidance in the Argentine ant *Linepithema humile*.

Behavioral Ecology, *13*(1), 28–31. https://doi.org/10.1093/beheco/13.1.28

Klein, A., Andersson, J., Ardekani, B. A., Ashburner, J., Avants, B., Chiang, M.-C., Christensen, G. E., Collins,

D. L., Gee, J., Hellier, P., Song, J. H., Jenkinson, M., Lepage, C., Rueckert, D., Thompson, P., Vercauteren,

T., Woods, R. P., Mann, J. J., & Parsey, R. V. (2009). Evaluation of 14 nonlinear deformation algorithms

applied to human brain MRI registration. *NeuroImage*, *46*(3), 786–802.

https://doi.org/10.1016/j.neuroimage.2008.12.037

Kramer, B. H., & Schaible, R. (2013). Life span evolution in eusocial workers—A theoretical approach to

understanding the effects of extrinsic mortality in a hierarchical System. *PLoS ONE*, *8*(4), e61813.

https://doi.org/10.1371/journal.pone.0061813

Lemanski, N. J., Cook, C. N., Ozturk, C., Smith, B. H., & Pinter-Wollman, N. (2021). The effect of individual

learning on collective foraging in honey bees in differently structured landscapes. *Animal Behaviour*, *179*,

113–123. https://doi.org/10.1016/j.anbehav.2021.06.033

Li, Y., & Strausfeld, N. J. (1999). Multimodal efferent and recurrent neurons in the medial lobes of

cockroach mushroom bodies. *Journal of Comparative Neurology*, *409*(4), 647–663. Scopus.

https://doi.org/10.1002/(SICI)1096-9861(19990712)409:4<647::AID-CNE9>3.0.CO;2-3

Lihoreau, M., Latty, T., & Chittka, L. (2012). An exploration of the social brain hypothesis in insects.

Frontiers in Physiology, *3*. https://doi.org/10.3389/fphys.2012.00442

López, D. N., Fuentes-Contreras, E., Ruiz, C., Ide, S., & Estay, S. A. (2023). A bug's tale: Revealing the history,

biogeography and ecological patterns of 500 years of insect invasions. *NeoBiota*, *81*, 183–197.

https://doi.org/10.3897/neobiota.81.87362

Lucon-Xiccato, T., Carere, C., & Baracchi, D. (2024). Intraspecific variation in invertebrate cognition: a

review. *Behavioral Ecology and Sociobiology*, 78(1), 1. https://doi.org/10.1007/s00265-023-03413-8

Maák, I., Roelandt, G., & d'Ettorre, P. (2020). A small number of workers with specific personality traits

perform tool use in ants. *eLife*, *9*, e61298. https://doi.org/10.7554/eLife.61298

Mayer, M., Shine, R., & Brown, G. P. (2016). Bigger babies are bolder: Effects of body size on personality

of hatchling snakes. *Behaviour*, *153*(3), 313–323. https://doi.org/10.1163/1568539X-00003343

Mazerolle, M. (2006). Improving data analysis in herpetology: Using Akaike's Information Criterion (AIC)

to assess the strength of biological hypotheses. *Amphibia-Reptilia*, *27*(2), 169–180.

https://doi.org/10.1163/156853806777239922

Menzel, R., & Giurfa, M. (2001). Cognitive architecture of a mini-brain: The honeybee. *Trends in Cognitive*

Sciences, *5*(2), 62–71. https://doi.org/10.1016/S1364-6613(00)01601-6

Moffett, M. W. (2012). Supercolonies of billions in an invasive ant: What is a society? *Behavioral Ecology*,

23(5), 925–933. https://doi.org/10.1093/beheco/ars043

Nakagawa, S., & Schielzeth, H. (2010). Repeatability for Gaussian and non-Gaussian data: A practical guide

for biologists. *Biological Reviews*, *85*(4), 935–956. https://doi.org/10.1111/j.1469-185X.2010.00141.x

Navas-Zuloaga, M. G., Pavlic, T. P., & Smith, B. H. (2022). Alternative model systems for cognitive variation:

Eusocial-insect colonies. *Trends in Cognitive Sciences*, *26*(10), 836–848.

https://doi.org/10.1016/j.tics.2022.06.011

Newberry, R. C. (1995). Environmental enrichment: Increasing the biological relevance of captive

environments. *Applied Animal Behaviour Science*, *44*(2), 229–243. https://doi.org/10.1016/0168-

1591(95)00616-Z

O'Donnell, S., Donlan, N. A., & Jones, T. A. (2004). Mushroom body structural change is associated with

division of labor in eusocial wasp workers (*Polybia aequatorialis*, Hymenoptera: *Vespidae*). *Neuroscience*

Letters, *356*(3), 159–162. https://doi.org/10.1016/j.neulet.2003.11.053

Okrutniak, M., Rom, B., Turza, F., & Grześ, I. M. (2020). Body size differences between foraging and

intranidal workers of the monomorphic ant *Lasius niger*. *Insects*, *11*(7), Article 7.

https://doi.org/10.3390/insects11070433

Ott, S. R. (2008). Confocal microscopy in large insect brains: Zinc–formaldehyde fixation improves synapsin

immunostaining and preservation of morphology in whole-mounts. *Journal of Neuroscience Methods*,

172(2), 220–230. https://doi.org/10.1016/j.jneumeth.2008.04.031

Pinter-Wollman, N. (2012). Personality in social insects: How does worker personality determine colony

personality? *Current Zoology*, *58*(4), 580–588. https://doi.org/10.1093/czoolo/58.4.580

Pratt, S. C., Brooks, S. E., & Franks, N. R. (2001). The use of edges in visual navigation by the ant *Leptothorax*

albipennis. *Ethology*, *107*(12), 1125–1136. https://doi.org/10.1046/j.1439-0310.2001.00749.x

Prentis, P. J., Wilson, J. R. U., Dormontt, E. E., Richardson, D. M., & Lowe, A. J. (2008). Adaptive evolution

in invasive species. *Trends in Plant Science*, *13*(6), 288–294. https://doi.org/10.1016/j.tplants.2008.03.004

Rössler, W. (2019). *Neuroplasticity in desert ants (Hymenoptera: Formicidae) – importance for the*

ontogeny of navigation. https://doi.org/10.25849/MYRMECOL.NEWS_029:001

Sandoz, J. C. (2011). Behavioral and neurophysiological study of olfactory perception and learning in

honeybees. *Frontiers in Systems Neuroscience*, *5*. https://doi.org/10.3389/fnsys.2011.00098

Sanmartín-Villar, I., Cruz Da Silva, E., Chiara, V., Cordero-Rivera, A., & Lorenzo-Carballa, M. O. (2022).

Genetic divergence and aggressiveness within a supercolony of the invasive ant *Linepithema humile*.

NeoBiota, *77*, 125–153. https://doi.org/10.3897/neobiota.77.90852

Sanmartín-Villar, I., Csata, E., & Jeanson, R. (2021). Variability in activity differs between castes in the ant.

Ecological Entomology, *46*(6), 1373–1378. https://doi.org/10.1111/een.13075

Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., Preibisch, S., Rueden, C.,

Saalfeld, S., Schmid, B., Tinevez, J.-Y., White, D. J., Hartenstein, V., Eliceiri, K., Tomancak, P., & Cardona, A.

(2012). Fiji: An open-source platform for biological-image analysis. *Nature Methods*, *9*(7), 676–682.

https://doi.org/10.1038/nmeth.2019

Seid, M. A., & Junge, E. (2016). Social isolation and brain development in the ant *Camponotus floridanus*.

The Science of Nature, *103*(5), 42. https://doi.org/10.1007/s00114-016-1364-1

Seid, M. A., & Wehner, R. (2008). Ultrastructure and synaptic differences of the boutons of the projection

neurons between the lip and collar regions of the mushroom bodies in the ant, *Cataglyphis albicans*. *The*

Journal of Comparative Neurology, *507*(1), 1102–1108. https://doi.org/10.1002/cne.21600

Sneddon, L. U. (2003). The bold and the shy: Individual differences in rainbow trout. *Journal of Fish*

Biology, *62*(4), 971–975. https://doi.org/10.1046/j.1095-8649.2003.00084.x

Snell-Rood, E. C. (2013). An overview of the evolutionary causes and consequences of behavioural

plasticity. *Animal Behaviour*, *85*(5), 1004–1011. https://doi.org/10.1016/j.anbehav.2012.12.031

Stamps, J. A., Briffa, M., & Biro, P. A. (2012). Unpredictable animals: Individual differences in

intraindividual variability (IIV). *Animal Behaviour*, *83*(6), 1325–1334.

https://doi.org/10.1016/j.anbehav.2012.02.017

Standage, D., Blohm, G., & Dorris, M. C. (2014). On the neural implementation of the speed-accuracy

trade-off. *Frontiers in Neuroscience*, *8*. https://doi.org/10.3389/fnins.2014.00236

Stapley, J., Santure, A. W., & Dennis, S. R. (2015). Transposable elements as agents of rapid adaptation

may explain the genetic paradox of invasive species. *Molecular Ecology*, *24*(9), 2241–2252.

https://doi.org/10.1111/mec.13089

Stoffel, M. A., Nakagawa, S., & Schielzeth, H. (2017). rptR: Repeatability estimation and variance

decomposition by generalized linear mixed-effects models. *Methods in Ecology and Evolution*, *8*(11),

1639–1644. https://doi.org/10.1111/2041-210X.12797

Trigos-Peral, G., Maák, I. E., Ślipiński, P., & Witek, M. (2023). Behavioral and morphological traits

influencing variation in task performance of *Camponotus vagus* ants. *Insectes Sociaux*, *70*(4), 451–461.

https://doi.org/10.1007/s00040-023-00937-0

Tripet, F., & Nonacs, P. (2004). Foraging for work and age-based polyethism: The roles of age and previous

experience on task choice in ants. *Ethology*, *110*(11), 863–877. https://doi.org/10.1111/j.1439-

0310.2004.01023.x

Valle, F. P. (1970). Effects of Strain, Sex, and Illumination on Open-Field Behavior of Rats. *The American*

Journal of Psychology, *83*(1), 103–111. https://doi.org/10.2307/1420860

Van Nest, B. N., Wagner, A. E., Marrs, G. S., & Fahrbach, S. E. (2017). Volume and density of microglomeruli

in the honey bee mushroom bodies do not predict performance on a foraging task. *Developmental*

Neurobiology, *77*(9), 1057–1071. https://doi.org/10.1002/dneu.22492

Wagner, T., Galante, H., Josens, R., & Czaczkes, T. J. (2023). Systematic examination of learning in the

invasive ant *Linepithema humile* reveals fast learning and long-lasting memory. *Animal Behaviour*, *203*,

41–52. https://doi.org/10.1016/j.anbehav.2023.06.012

Walsh, R. N., & Cummins, R. A. (1976). The open-field test: A critical review. *Psychological Bulletin*, *83*(3),

482–504. https://doi.org/10.1037/0033-2909.83.3.482

Webb, B., & Wystrach, A. (2016). Neural mechanisms of insect navigation. *Current Opinion in Insect*

Science, *15*, 27–39. https://doi.org/10.1016/j.cois.2016.02.011

Weidenmüller, A., Mayr, C., Kleineidam, C. J., & Roces, F. (2009). Preimaginal and adult experience

modulates the thermal response behavior of ants. *Current Biology*, *19*(22), 1897–1902.

https://doi.org/10.1016/j.cub.2009.08.059

Wystrach, A., Lagogiannis, K., & Webb, B. (2016, October 18). *Continuous lateral oscillations as a core*

mechanism for taxis in Drosophila larvae. eLife; eLife Sciences Publications Limited.

https://doi.org/10.7554/eLife.15504

Clutton-Brock, T. H., Albon, S. D., & Guinness, F. E. (1985). Parental investment and sex differences in

juvenile mortality in birds and mammals. *Nature*, *313*(5998), 131–133.

https://doi.org/10.1038/313131a0

Einum, S., & Fleming, I. A. (n.d.). *Environmental unpredictability and offspring size: Conservative versus*

diversified bet-hedging.

Fawcett, T. W., Hamblin, S., & Giraldeau, L.-A. (2013). Exposing the behavioral gambit: The evolution of

learning and decision rules. *Behavioral Ecology*, *24*(1), 2–11. https://doi.org/10.1093/beheco/ars085

Van Nest, B. N., Wagner, A. E., Marrs, G. S., & Fahrbach, S. E. (2017). Volume and density of

microglomeruli in the honey bee mushroom bodies do not predict performance on a foraging task.

Developmental Neurobiology, *77*(9), 1057–1071. https://doi.org/10.1002/dneu.22492

Yilmaz, A., Lindenberg, A., Albert, S., Grübel, K., Spaethe, J., Rössler, W., & Groh, C. (2016a). Age-related

and light-induced plasticity in opsin gene expression and in primary and secondary visual centers of the

nectar-feeding ant *Camponotus rufipes*. *Developmental Neurobiology*, *76*(9), 1041–1057.

https://doi.org/10.1002/dneu.22374

Yilmaz, A., Lindenberg, A., Albert, S., Grübel, K., Spaethe, J., Rössler, W., & Groh, C. (2016b). Age-related

and light-induced plasticity in opsin gene expression and in primary and secondary visual centers of the

nectar-feeding   ant   *Camponotus   rufipes*.   *Developmental   Neurobiology*,   *76*(9),   1041–1057.

https://doi.org/10.1002/dneu.22374

Yoo, T. S. (2004). *Insight into Images: Principles and Practice for Segmentation, Registration, and Image*

Analysis. AK Peters Ltd.

1083

1084 **Table 1**. Statistical values (*Value*: t, F or Z) for the behavioural variables measured. *Ratio choices*

1085 *reward*: relative number of times ants selected the path connecting to the reward instead of the

1086 dead-end paths of the maze.

Variables	Test	Value	DF (F)	P
Speed × Size	Open field	26.78	1;154.36	**<0.001**
	Object exposure	31.55	1;148.50	**<0.001**
	Maze	6.69	1;151.23	**0.011**
Speed × Observation	Open field	11.83	1;621.48	**<0.001**
	Object exposure	9.04	1;600.23	**0.003**
	Maze	17.10	1;605.13	**<0.001**
Time Moving (%) × Size	Open field	11.17	1;153.78	**0.001**
	Object exposure	5.86	1;154.07	**0.017**
	Maze	2.83		**0.005**
Time Moving (%) × Observation	Open field	3.29	1;621.71	**0.070**
	Object exposure	10.67	1;622.01	**0.001**
	Maze	-2.92		**0.004**
Exploration × Size	Open field	3.46		**<0.001**
	Object exposure	1.76		0.078
	Maze	2.72		**0.007**
Exploration × Observation	Open field	0.45		0.656
	Object exposure	1.90		0.057
	Maze	1.45		0.141
Object interaction × Size	Object exposure	-0.93		0.354
Object interaction × Observation	Object exposure	8.25		**<0.001**
Inner arena use × Size	Open field	3.33		**<0.001**
	Object exposure (control)	-2.82		**0.005**
Inner arena use × Observation	Open field	3.03		**0.002**
	Open field (control)	0.57		0.572
	Object exposure (control)	2.27		**0.026**
Maze participation × Size	Maze	1.91	1;149	0.169
Maze participation × Observation	Maze	2.62	1;603	0.106
Ratio choices reward × Size	Maze	0.51		0.608
Ratio choices reward × Observation	Maze	-0.35		0.725
Exploration × Object interaction	Open field & Object exposure	2.88		**0.004**
Exploration × Object interaction × Observation	Open field & Object exposure	0.29		0.773
Exploration × Ratio choices reward	Open field & Maze	1.25		0.212
Object interaction × Ratio choices reward	Object exposure & Maze	1.34		0.182

Exploration unpredictability × Size	Open field	1.91	1;152	0.169
Object interaction unpredictability × Size	Object exposure	5.44	1;152	**0.021**
Rewarded path choice unpredictability × Size	Maze	0.00	1;152	0.926
Exploration unpredictability × Object interaction unpredictability	Open field & Object exposure	0.48	1;154	0.635
Exploration unpredictability × Ratio choices reward unpredictability	Open field & Maze	-0.09	1;154	0.930
Object interaction unpredictability × Ratio choices reward unpredictability	Object exposure & Maze	0.51	1;154	0.611

1087

1088 **Table 2**. Intraindividual differences. *Ratio choices reward*: relative number of times ants selected

1089 the path connecting to the reward instead of the dead-end paths of the maze.

Variable	Test	ΔAIC	R^2m	R^2c
Exploration	Open field		0.06	0.49
	Object exposure		0.01	0.11
	Maze		-0.22	-2.82
Object interaction	Object exposure	9.02	0.16	1.60
Ratio choices reward before flip	Maze	73.55		
Ratio choices reward after flip	Maze	52.72		

1090

Figure 1. Behavioural tests and patterns of *L. humile* workers (regression models) across

observations. **A-B**: Schematic representation of the novel object (A) and maze (B) used in the

behavioural tests. Both objects can be downloaded for printing from

https://www.tinkercad.com/things/2lNxDsoOoAD-neophobia-object-miiz and

https://www.tinkercad.com/things/7IELVyTIwXr-maze-miiz. **C**: Exploration in the open field test.

D: Interaction with the object. **E**: Relative number of times ants selected the path connecting to

the reward instead of the dead-end paths of the maze (*Ratio choices reward*) before maze

direction flip. **F**: Relative number of times ants selected the path connecting to the reward instead

of the dead-end paths of the maze (*Ratio choices reward*) after maze direction flip.

Figure 2. Correlation between brain volumetry and behaviour of *L. humile* workers: correlation

between the relative volume of the optic lobe and the mean speed performed (**A**); correlation

between the ratio primary:secondary neuropils and the mean exploration performed in presence

of the object (**B**), on the maze (**C**), and the slopes for the relative number of times ants selected

the path connecting to the reward instead of the dead-end paths of the maze (*Ratio choices*

reward) before flip (**D**); correlation between the relative volume of the central complex and the

slopes for the relative number of times ants selected the path connecting to the reward instead

of the dead-end paths of the maze before flip (**E**). Continuous red: linear regression between the

two variables. Dashed orange: zero value to facilitate the visual comparison of negative and

positive slopes. F: different sections of a *L. humile* brain showing the neuropils analysed (synapsin

in green).

Figure 3. Correlation between the number of microglomeruli and behaviour of L. humile workers:

correlation between the number of microglomeruli of the dense lip and the variance (**A**) and

unpredictability (**B**) of the object interaction, and the slopes of choices before maze flip (**C**);

correlation between the number of microglomeruli of the collar and the slopes of choices before

maze flip (**D**).

Figure 1

A

1118

1119

1120   2.5 mm

B

13.75 mm

☐ Catoira ☐ Carril ☐ Chazo ☐ Rianxo ☐ Vilanova

1121

1122

1123   **Figure 2**

1124

1125

1126

1127 **Figure 3**

1128

1129

1130