1 **Brain neuromarkers predict self- and other-related mentalizing across**
2 **adult, clinical, and developmental samples**
3

4 Dorukhan Açıl[1,2,3], Jessica R. Andrews-Hanna[4,5], Marina Lopez-Sola[6,7], Mariët van
5 Buuren[8], Lydia Krabbendam[8], Liwen Zhang[9], Lisette van der Meer[10,11], Paola Fuentes-
6 Claramonte[12,13], Edith Pomarol-Clotet[12,13], Raymond Salvador[12,13], Martin Debbané[14,15],
7 Pascal Vrticka[16], Patrik Vuilleumier[17], David A. Sbarra[4], Andrea M. Coppola[4], Anita
8 Tusche[18, 19], Lars O. White[3], Tor D. Wager[20], & Leonie Koban[21]*
9

10
[1] Max Planck Institute for Human Cognitive and Brain Sciences, Leipzig, Germany
[2] Department of Child and Adolescent Psychiatry, Psychotherapy, and Psychosomatics, Leipzig
University, Leipzig, Germany
[3] Department of Clinical Child and Adolescent Psychology and Psychotherapy, University of
Bremen, Bremen, Germany
[4] Department of Psychology, University of Arizona, Tucson, Arizona, USA
[5] Cognitive Science, University of Arizona, Tucson, Arizona, USA
[6] Department of Medicine, School of Medicine and Health Sciences, Institute of Neurosciences,
University of Barcelona, Spain
[7] Institut d'Investigacions Biomèdiques August Pi i Sunyer (IDIBAPS), Barcelona, Spain
[8] Department of Clinical, Neuro and Developmental Psychology, Faculty of Behavioral and
Movement Sciences, Institute for Brain and Behavior Amsterdam, Vrije Universiteit Amsterdam,
The Netherlands
[9] Institute for Medical Imaging Technology, Ruijin Hospital, Shanghai Jiao Tong University School
of Medicine, Shanghai, China
[10] Department of Clinical and Developmental Neuropsychology, University of Groningen,
Groningen, The Netherlands
[11] Department of Psychiatric Rehabilitation, Lentis Zuidlaren, The Netherlands
[12] FIDMAG Germanes Hospitalàries Research Foundation, Barcelona, Spain
[13] Centro de Investigación Biomédica en Red de Salud Mental (CIBERSAM) ISCIII, Barcelona,
Spain
[14] Developmental Clinical Psychology Research Unit, Faculty of Psychology and Educational
Sciences, University of Geneva, Switzerland
[15] Research Department of Clinical, Educational and Health Psychology, University College
London, United Kingdom
[16] Department of Psychology, University of Essex, Colchester, United Kingdom
[17] Laboratory of Behavioural Neurology and Imaging of Cognition, Department of Neuroscience,
University Medical Center, University of Geneva, Geneva, Switzerland
[18] Department of Psychology, Queen's University, Kingston, ON, Canada
[19] Center for Neuroscience Studies, Queen's University, Kingston, ON, Canada
[20] Department of Psychological and Brain Sciences, Dartmouth College, Hanover, NH, USA
[21] Lyon Neuroscience Research Center (CRNL), CNRS, Inserm, Université Claude Bernard Lyon
1, Bron France
44
45
46 Running Head: BRAIN SIGNATURES OF MENTALIZING
47
48 *Please address correspondence to:
49 Dr. Leonie Koban, CRNL, Institut des Épilepsies IDEE, 59 Boulevard Pinel, 69500 Bron,
50 France; email: leonie.koban@cnrs.fr

Abstract

Human social interactions rely on the ability to reflect on one's own and others' internal states and traits—a process known as mentalizing. Impaired or altered mentalizing is a hallmark of multiple psychiatric and neurodevelopmental conditions. Yet, replicable and easily testable brain markers of mentalizing have so far been lacking. Here, we apply an interpretable machine learning approach to multiple datasets (total N=390) to train and validate fMRI brain signatures that predict i) mentalizing about the self, ii) mentalizing about another person, and iii) both types of mentalizing. Self-mentalizing and other-mentalizing classifiers had positive weights in anterior/medial and posterior/lateral brain areas, respectively, with accuracy rates of 82% and 77% for out-of-sample prediction. The classifier trained across both types of mentalizing showed 98% predictive accuracy and separated (mental) attributional from factual inferences. Classifier patterns revealed better self/other separation in healthy adults compared to individuals with schizophrenia and with increasing age in adolescence. Together, our findings reveal consistent and separable neural patterns subserving trait-based mentalizing about self and others—present at least from the age of adolescence and functionally altered in severe neuropsychiatric disorders. These mentalizing signatures hold promise as candidate neuromarkers of social-cognitive processes in different contexts and clinical conditions.

Author Note

This work was funded by a Starting Grant from the European Research Council (ERC, 101041087) to LKo, a German Academic Exchange Service (DAAD) doctoral grant and a Network of European Neuroscience Schools (NENS) exchange fellowship to DA, an R01 grant from the U.S. National Institutes of Mental Health (R01MH125414-01) to JAH and DAS, a Junior Leader Fellowship from "la Caixa" Foundation (LCF/BQ/PR22/11920017) to PFC, a Consolidator Grant from the European Research Council (ERC, 648082) to LKr, R37, R01 support from the U.S. National Institutes of Mental Health (R37MH076136 to TDW, MH116026 to TDW and L. Chang [PI], R01EB026549 to TDW and M. Lindquist [MPIs]), an NIMH grant (P50MH094258-01A1) to R. Adolphs, and a CIC Brain and Mental Health Chair from the Neurodis foundation to AT. LvdM acknowledges a European Science Foundation EURYI grant (044035001) that funded her doctoral studies (PI: A. Aleman). Views and opinions expressed are however those of the authors only and do not necessarily reflect those of the European Union or the European Research Council. Neither the European Union nor the granting authority can be held responsible for them. The funders had no role in study design, data analysis, manuscript preparation, or publication decisions.

86   **Introduction**

Mentalizing—representing and inferring the psychological states of oneself and

others—is a fundamental process for adaptive navigation through the social world[1,2].

Delineating the brain systems involved in mentalizing about self and others is important

for understanding brain health and dysfunction, as atypical mentalizing is characteristic

for many neurodevelopmental and psychiatric conditions[3–9]. Many studies have examined

the neural correlates of mentalizing, suggesting an interplay of different brain regions,

including medial prefrontal cortex (mPFC), temporoparietal junction (TPJ), and

precuneus[10–14]. However, predictive brain measures of mentalizing that can be easily

applied to new data to decode the degree of self-related and other-related processing are

still lacking. Most cognitive and affective processes cannot be captured by activity in

individual brain regions, as they are reflected in patterns of brain activity distributed across

multiple regions and systems, which can be harnessed by pattern-based decoding[15,16].

For instance, recent work demonstrates that distributed brain activity and connectivity

patterns, as indexed by fMRI, enable us to decode the intensity of pain[17], drug and food

craving[18], sustained attention[19], depressive rumination[20], and clinically relevant behaviors

and outcomes[21–23]. As opposed to the standard brain mapping approach that maps task

features onto brain activity, these predictive brain activity patterns or '*brain signatures'* are

multivariate models that utilize data across the whole brain to make formally testable

population-level predictions across subjects and datasets[15,24]. The predictions address the

involvement and/or the intensity of a mental process. Here, we apply this brain signature

approach to predict mentalizing about oneself or other people.

Mentalizing, as a hidden state, would arguably benefit from such an approach.

Thinking about self and others is inherently multilayered and multidimensional[12,25,26].

Recent conceptualizations of self- and social-cognition point to multiple dimensions, such

as about oneself versus others, affective versus cognitive[6,12,27] and multiple layers, such

as observing behaviors, and processing at the state and trait levels[25]. Moreover, social

cognition terms, including but not limited to mentalizing (e.g., empathy, perspective

taking), suffer from inconsistent usage in the literature and a lack of consensual

definitions[28]. Developing brain signatures of mentalizing could potentially help account for

this heterogeneity, offering the possibility of testing whether the proposed subdimensions

of mentalizing are subserved by dissociable neurobiological patterns. In turn, these

signatures carry the potential for validating the multiple facets of mentalizing[15]. However,

it remains unclear—especially considering the complexity of the mentalizing construct—

whether distributed neural patterns can reliably predict mentalizing using brain images.

A central question is whether mentalizing about oneself and mentalizing about

others (self- and other-mentalizing hereafter) are reliably distinguishable based on brain

activity. Recent electrophysiological evidence suggests that self- and other-mentalizing

activate overlapping cortical areas following a similar temporal sequence: TPJ, medial

temporal gyrus (MTG)/temporal poles (TP), precuneus/posterior cingulate cortex (PCC),

mPFC, in a roughly posterior to anterior temporal order[29,30]. Conversely, differences in

brain activity between self- and other-mentalizing have also been observed. In the mPFC,

more dorsal areas coincide with other-related processing and ventral areas with self-

related processing, with research initially supporting a linear[31] but more recently a

curvilinear[32] ventral-to-dorsal gradient for self versus other within the mPFC. Further, self-

referential thought typically elicits activation in cortical midline structures, such as anterior

cingulate cortex (ACC), subcortical areas, including thalamus, striatum and caudate

nucleus, and, to a lesser extent, in insula, temporal poles, and ventrolateral prefrontal

cortex[31–36] (vlPFC). In contrast, other-referential thought typically elicits activation in TPJ,

middle and superior temporal gyri extending to temporal poles[10,11,32,37–39], and to a lesser

degree in supplementary motor area (SMA), left inferior and medial frontal gyri and medial

orbitofrontal cortex[34,40]. Collectively, the literature remains inconsistent regarding the

separation between self- and other-mentalizing, and it is unclear whether generalizable

brain models of self- versus other-mentalizing can be identified.

To address these gaps, we leverage fMRI and machine learning to develop four

distinct brain signatures, 1) the Mentalizing Signature (MS) for mentalizing overall (i.e.,

thinking about either the self or another person versus non-mentalizing control conditions),

2) the Self-Referential Signature (Self-RS) to detect specifically self-related thought (here

referred to as "self-mentalizing"), 3) the Other-Referential Signature (Other-RS) to detect

other-related thought (here referred to as "other-mentalizing"), and 4) the Self-vs-Other

Referential Signature (SvO-RS) to specifically separate self- and other-related

mentalizing. While the overall Mentalizing Signature aimed at developing a broad marker

of social reflection that establishes the benchmark ceiling performance and may

generalize across a wide range of social processes, the other three signatures (Self-RS,

Other-RS, and SvO-RS) allowed us to test whether dissociable and generalizable neural

patterns underlie distinct targets of mentalizing.

152   To this end, we used data from 11 cohorts and seven independent fMRI studies.

153 The training and testing datasets used variants of a standard trait-evaluation task with

154 three conditions, which involved reflecting on personality traits/statements describing

155 oneself or another person. Conceivably, personality traits represent the enduring mental

156 states that can be inferred from information collected by social perceptual systems[41]

157 across multiple observations[12,25]. Trait representations are used to predict others' future

158 behavior and become the basis of the "mental-self"[26]. Moreover, trait evaluations are one

159 of the few mentalizing tasks used in the literature to include a self-condition, as other

160 commonly used tasks (e.g., mental attribution, false-belief, perspective-taking tasks) only

161 permit mentalizing about others. Thus, trait evaluation is an ideally suited task to study a

162 key mentalizing process, namely representing stable psychological states of self and

163 others.

164   We first used standard machine learning algorithms—support vector machines—

165 to develop and cross-validate multivariate classifiers (signatures) of mentalizing about

166 traits in a sample of healthy adults. In a second step, we further validated these signatures

167 in seven independent cohorts from different laboratories, countries, and scanners, and

168 with different sample characteristics (healthy, adolescent, and clinical populations),

169 allowing us to test their generalizability and predictive validity in adolescent and clinical

170 populations. Third, we tested the signatures in two additional independent datasets that

171 used different social cognition tasks to see whether mentalizing signatures would

172 generalize to other social contexts in which mentalizing is likely to occur. Finally, we

173 assessed whether local patterns of brain activity in several regions of interest (ROIs)

174 contain sufficient information to predict self- and other-referential mentalizing. Together,

175 the results of these analyses inform us about the functional neural organization of self-

176 versus other-related mentalizing and provide us with distributed brain signatures that can

177 serve as brain targets for measuring and intervening on mentalizing-related brain

178 processes in future studies.

180        **Results**

181 **Data overview**

182   The study included a total of 1122 contrast images from 390 participants and 11

183 independent cohorts, including six samples of healthy adults (*n*=227), two samples of

184 healthy adolescents (*n*=105; M_{age}=12.9 and 16), and three samples of adults with clinical

185 diagnosis of either schizophrenia (*n*=40) or bipolar disorder (*n*=18; see Fig. 1A and

Supplementary Table 1). Thus, this study combined seven independent studies that
constituted the training dataset, testing datasets, and the extension datasets (see Fig. 1A).
The training and testing datasets included participants completing variants of a self- and
other-referential judgement task. The two extension datasets included participants
performing a social feedback task and an inference task. Studies 1-5 and 7 included three
conditions, namely a Self-condition, an Other-condition, and a nonsocial Control condition,
whereas Study 6 included two conditions in one study (attributional versus factual
inferences; Study 6a) and a 2x2 design in the second (Study 6b; attributional versus
factual inferences x social versus nonsocial scenes). Contrast images were computed for
each condition (versus implicit baseline) and rescaled using L2-norm to standardize the
scale of beta (regression coefficient) weights across participants, studies, and scanners.
197

Cross-validated training results

The training dataset (Study 1a) consisted of n=21 adult participants who completed
a trait-evaluation task using a block design (similar to ref[42]; see Fig. 1C). In each block,
participants were presented with several trait adjectives: In the Self-reflection condition,
they were asked to rate the degree to which each trait adjective described themselves. In
the Other-reflection condition, they had to rate how much each adjective described
another person—a confederate with whom the participant had previously interacted during
a decision-making task[43]. In the non-mentalizing Control condition, participants indicated
the number of syllables in the trait adjectives. To reduce the influence of non-mentalizing-
related brain regions (e.g., visual cortex) and opportunistic classification based on features
not related to mentalizing, we used an inclusive mask of brain regions related to social
processing (see Fig. 1B).

We used support vector machines (SVM) with default parameters (to avoid
overfitting) and 10-fold cross-validation[44] to train three distinct mentalizing signatures
using the masked contrast images from the training dataset. The folds were defined at the
subject level, so that all contrast images from a given subject were assigned to the same
fold. The Self-Referential Signature (Self-RS) was trained to detect Self-reflection (versus
the two remaining conditions), the Other-Referential Signature (Other-RS) to detect Other-
reflection (versus the two remaining conditions), the Mentalizing Signature (MS) to detect
both types of mentalizing versus the Control condition, and the Self-vs-Other Referential
Signature (SvO-RS) to differentiate Self- versus Other-related processing (see Fig. 1C).

All four signatures showed excellent cross-validated (out-of-sample) prediction
accuracy (100% accuracy in two-alternative forced-choice tests, $p<.001$, *Cohen's d* for the
Self-RS = 3.18, the Other-RS = 2.45, the MS = 4.92; the SvO-RS = 2.45; AUC = 1.00 for
all signatures). To identify which voxels contributed most reliably to the mentalizing
signatures, we used bootstrapping (5000 samples) to obtain one *p*-value per voxel and
displayed the thresholded weight maps using false discovery rate (FDR) correction at
$q < .05$ and cluster extent $k > 10$ voxels. Brain regions with significant positive voxel
weights for the Self-RS (see Fig. 2) included the ventromedial (vmPFC) and dorsomedial
(dmPFC) prefrontal cortex, frontal eye fields, ventral ACC, frontal operculum, anterior
insula, thalamus, and caudate nucleus (see Supplementary Table 2). For the Other-RS,
significant positive weights were found in left vlPFC, left superior temporal sulcus (STS),
bilateral TPJ, and precuneus/PCC (see Fig. 2 and Supplementary Table 3). For the SvO-
RS (see Fig. 3 and Supplementary Table 5), brain regions with significant positive voxel
weights (thus, associated more with the Self than the Other condition) included
vmPFC/ACC, left anterior insula, left inferior frontal sulcus, left caudate nucleus, left
middle temporal cortex, and bilateral thalamus. Significant negative clusters in the SvO-
RS (thus, associated more with the Other than the Self condition) were found in
precuneus, bilateral TPJ/inferior parietal cortex, right PCC, left STS, and left inferior frontal
gyrus. The MS had significant positive clusters in mPFC (both dorsal and medial), bilateral
vlPFC, dorsal ACC (dACC), frontal operculum, bilateral SMA, bilateral MTG, bilateral TP,
left STS, middle cingulate gyrus (MCC), bilateral PCC, precuneus, bilateral angular
gyrus/TPJ, and subcortical areas, including right caudate nucleus, left anterior insula, and
bilateral thalamus (see Fig. 2 and Supplementary Table 4).
242

Validation in independent samples
Next, we validated the four brain signatures on several completely independent
studies: two samples of healthy adults (Studies 4a & 5a), two adolescent samples (Studies
2 & 3), one cohort of participants with bipolar disorder (Study 5c), and two cohorts of
participants with schizophrenia (Studies 4b & 5b). All datasets included comparable trait
evaluation tasks and fMRI block designs with three conditions, with some small variations
in task designs between studies (see Fig. 1A). To obtain pattern expression values, we
computed the matrix dot product between the mentalizing signatures and each subject-
level contrast image from these datasets, yielding one scalar value per individual contrast
image and per signature (see Fig. 1C). These pattern expression values were then used

to test the predictions of the mentalizing signatures. The weighted average prediction

accuracy in two-alternative forced-choice (2AFC) tests, across all independent testing

datasets, was 81.52% for the Self-RS (+/- 6.17% average STE; significant in 10 out of 14

validation tests [7 samples*2 comparisons]), 77.25% for the Other-RS (+/- 6.02% average

STE; significant in 12 out of 14 tests), 97.87% for the MS (+/- 1.79% average STE;

significant in all 14 tests), and 77.73% for the SvO-RS (+/- 7.14% average STE; significant

in 6 out of 7 tests), suggesting overall high prediction accuracy of the four signatures, even

in new samples, although with some variations between datasets (see Fig. 2, Fig. 3 &

Supplementary Table 6). There were no sex effects on pattern expression or signature

fit—defined as the difference between the pattern expression for the true condition and

the false condition in each signature (see Methods).

Lower self/other separation in participants with schizophrenia

The results above show that the signatures significantly predicted mentalizing in

most samples, including clinical samples. Yet, schizophrenia, in particular, is often

associated with impaired social cognition and altered self-perception[45,46]. Thus, we next

tested how well the signatures separated self- from other-related mentalizing in two of the

testing datasets (Study 4 & Study 5) that included both participants with schizophrenia

(total n = 40) and matched healthy participants (total n = 48). As expected, the three

signatures, the Self-RS (M_{HC} = .32, M_{SCZ} = .12; β = .21, *STE* = .07, *CI* = [.07, .35], $t(86)$ =

2.99, p = .004), the Other-RS (M_{HC} =.43, M_{SCZ} = .24; β = .19, *STE* = .07, *CI* = [.04, .33],

$t(86)$ = 2.57, p = .012), and the SvO-RS (M_{HC} =.39, M_{SCZ} = .19; β = .20, *STE* = .07,

CI = [.07, .34], $t(86)$ = 2.97, p = .004) showed better discrimination between self- and

other-referential mentalizing in healthy adults, compared to adults with schizophrenia (i.e.,

a greater positive difference between the Self-RS responses in the Self compared to the

Other condition, and a greater difference between the Other-RS responses for the Other

compared to the Self condition). No group differences were found in the correct

discrimination of Mentalizing versus Control by the MS (p = .29, see Fig. 4). These results

indicate that, compared to healthy adults, participants with schizophrenia have less

differentiated brain patterns between self- and other-related thought, indicating the

potential clinical utility of the signatures.

For completeness, we also compared the discrimination of self- versus other-

related mentalizing activity in participants with bipolar disorder (n = 18) versus healthy

adults (n = 15) using data from Study 5. However, participants with bipolar disorder did

not differ significantly from controls for any of the signatures.

Better self-other separation with increasing age in adolescents

Next, we explored potential developmental differences in the classifier

performances by testing whether the ability of the classifiers to separate self- from other-

related mentalizing depended on the age of the participants in the two adolescent samples

(N = 105). We combined data from Study 2 (age 12-18 years; M_{age} = 16) and Study 3 (age

11-14 years; M_{age} = 12.9). We found that (controlling for sex and the source study) older

adolescents had better self/other separation for the three signatures trained to distinguish

these two conditions: the Self-RS (β = .08, STE = .03, CI = [.018, .14], t(102) = 2.55, p =

.012), the Other-RS (β = .06, STE = .03, CI = [.001, .12], t(102) = 2.03, p = .045), and the

SvO-RS (β = .07, STE = .03, CI = [.01, .13], t(102) = 2.31, p = .023; see Fig. 5). In contrast,

there were no significant associations between age and performance of the MS. These

results suggest that, with increasing age, adolescents' brain activity becomes more

differentiated for self- versus other-related mentalizing.

Testing the signatures during attributional versus factual inference

So far, we have shown that the signatures performed well in several independent

datasets from different labs, but all using a similar explicit mentalizing task. In order to test

whether signatures are implicated specifically in mentalizing and not in broader social or

even nonsocial reflections, we tested them in an additional dataset (Study 6a & 6b) of

n=109 healthy adults making attributional versus factual inferences about social and

nonsocial stimuli (see Fig. 6A for the task). Attributional inferences (e.g., "Is Ada proud of

herself?"), especially in a social context, require mentalizing, or in other words, they

require representing hidden mental states. Factual inferences (e.g., "Is Ada tall?"), on the

other hand, refer to directly observable features and should not require mentalizing.

Therefore, we tested whether the Other-RS and the MS could predict attributional against

factual inference conditions successfully.

Participants in Study 6a (n = 59) were asked to make these two types of inferences

on social images showing faces or hands with a question prompt (e.g., is the person proud

of themselves?). Attributional (versus factual) inference conditions were predicted with

98% accuracy (2AFC) by both the Other-RS and the MS (see Fig. 6C; p < .001, AUC =

1.00, sensitivity = 98%; specificity = 98%, $Cohen's\ d$ = 2.62 for the Other-RS and 3.94 for

the MS). Study 6b had a 2 (attributional versus factual) x 2 (social versus nonsocial)

design. The Other-RS predicted the attributional inferences against factual inferences with

82% accuracy in the social condition ($p < .001$, *Cohen's d* = .95, AUC = .85, sensitivity =

82%, specificity = 82%) and 66% accuracy in the nonsocial condition ($p = .03$, *Cohen's d*

= .61, AUC = 0.73, sensitivity = 66%, specificity = 55%; see Fig. 6D). The MS predicted

the attributional versus factual inference with 100% accuracy in the social condition ($p <$

.001, *Cohen's d* = 3.56, AUC = 1.00, sensitivity = 100%, specificity = 100%) and 88%

accuracy in the nonsocial condition ($p < .001$, *Cohen's d* = 1.21, AUC = 0.92, sensitivity =

88%, specificity = 88%; see Fig. 6D). Further, we tested how well the signatures separate

social from nonsocial inferences (averaged across attributional and factual conditions).

While the Other-RS did not significantly separate social from non-social inferences (56%

accuracy, $p = .48$, Cohen's $d = 0.27$, AUC = 0.62, sensitivity = 56%, specificity = 56%),

the MS separated social from nonsocial inferences with almost perfect accuracy (98%

correct, $p < .001$, Cohen's $d = 2.96$, AUC = 1.00, sensitivity = 98%, specificity = 98%; see

Fig. 6B). Together these results suggest that the mentalizing signatures are sensitive to

processes that require a reflection about internal or hidden states, and–at least for the

MS–especially for social agents.

Testing the signatures in a social feedback task

To examine how the signatures would respond to social interaction tasks in which

mentalizing is not directly instructed but likely implicated, we next tested their performance

in an fMRI dataset (Study 7) of *n*=49 romantic partners performing a social feedback task.

Participants were led to believe that the task was designed to understand how people rate

other people and interpret others' ratings. Therefore, participants first indicated how likable

they felt each of 20 individuals (confederates) was, who supposedly rated the participants

and their partners in return. To set up a context facilitating spontaneous mentalizing,

participants were told that their partners saw the ratings on the screen simultaneously, in

a separate room. The task included conditions in which participants received feedback

about their own likeability (self-feedback), the likeability of their partner (partner-feedback),

as well as a no-feedback control condition. The control condition consisted of trials where

no feedback was given, with a prior explanation that such trials signaled that the rater (the

individual on the screen) was not shown the participants' picture or did not supply their

rating in the time required by the rating task.

Here, we tested whether the signatures' responses paralleled the target of the

feedback conditions. For instance, to provide evidence of successful extension, the Self-

RS should show higher pattern expression values in the self-feedback condition as

opposed to the other two conditions. As expected, the Self-RS responded more strongly

to the Self-condition, $F(2,96) = 14.214$, $p < .001$, $\eta_p^2 = .23$ (Fig. 7). Bonferroni-corrected

pairwise comparisons showed that the Self-feedback condition ($M = .16$) had significantly

higher Self-RS expression scores than both the Partner-feedback ($M = -.09$, $p=.01$) and

Control ($M = -.36$, $p < .001$) conditions. Additionally, the Self-RS also produced higher

pattern expression values for the Partner-feedback ($M = -.09$) compared to Control

condition ($M = -.36$, $p = .04$). Similarly, the MS successfully discriminated both Feedback

conditions against Control condition, $F(2,96) = 14.711$, $p < .001$, $\eta_p^2 = .24$ (Fig. 7).

Bonferroni-corrected pairwise comparisons indicated that both Self-feedback ($M = .17$,

$p < .001$) and Partner-feedback ($M = .12$, $p = .002$) conditions had significantly higher

pattern expression scores than the Control condition ($M = -.17$). However, the Other-RS

did not produce any significant differences between task conditions, $F(2,96) = 2.003$, $p =$

$.14$ (Fig. 7). Together this suggests a modulation of the MS and the Self-RS (but not the

Other-RS) even in a task where mentalizing was not directly instructed, in line with the

idea that feedback to oneself or one's partner should lead to more mentalizing related

brain activity.

Local patterns of self- and other-related mentalizing

We trained local classifiers in regions of interest (ROI) associated with mentalizing

to gain further insight into how mentalizing-related information is processed locally and

which regions can predict self- versus other-related mentalizing. The whole brain patterns

indicate which regions have the most reliable positive and negative contributions to

different forms of mentalizing, but they do not necessarily show which regions are *not*

involved and do not inform us whether local patterns alone can predict the target of

mentalizing (self or other). Training and testing classifiers for brain regions implicated in

the literature can inform us in this regard.

To this end, we included ten ROIs (see Fig. 8) previously associated with

mentalizing as shown in an automated term-based meta-analysis (NeuroSynth[47]): mPFC,

two clusters in anterior middle temporal gyrus (aMTG) bilaterally, two clusters in TPJ

bilaterally, a cluster covering precuneus and posterior cingulate cortex (precuneus/PCC),

left SMA, and three clusters in the cerebellum. We trained four types of classifiers in each

of these ten ROIs. In other words, we trained each ROI to perform the four following

classification tasks separately: 1) Self-mentalizing (versus the two remaining conditions);

2) Other-mentalizing (versus the two remaining conditions), 3) both mentalizing conditions

(self and other) against the Control condition, and 4) Self-mentalizing against Other-

mentalizing. The ROI classifiers were trained and cross-validated in our training dataset

(Study 1a; n = 21) and tested in the combined sample (n = 211) of participants from all

testing datasets (including all healthy, developmental, and clinical cohorts; for detailed

results, see Supplementary Table 7).

As expected, all ten regions showed excellent classification accuracy as the

general mentalizing classifiers in the training dataset, and all predicted mentalizing

successfully in the testing dataset (see blue bars in Fig. 8). However, not all regions could

significantly differentiate Self-reflection and Other-reflection conditions. While mPFC,

aMTG, TPJ, and precuneus/PCC were capable of differentiating self and other conditions,

the cerebellum clusters and left SMA failed at this task (see orange bars in Fig. 8). When

trained for self-mentalizing (see red bars in Fig. 8), the right aMTG cluster could not

capture any consistent configurations for Self-mentalizing. While TPJ subregions were

successful here against control conditions, they could not predict Self- against Other-

mentalizing, implying a lack of exclusive neural patterns for self-mentalizing within TPJ

subregions. Finally, when trained for Other-mentalizing, the Cerebellum subregions and

left SMA were incapable of picking up consistent patterns (see violet bars in Fig. 8).

Together, our analyses suggest that mPFC (including both vmPFC and dmPFC),

TPJ, aMTG, and precuneus/PCC clusters encode both self- and other-mentalizing, and

that the left SMA and Cerebellum clusters most likely are involved in domain-general

operations during mentalizing. Besides, our findings suggest that TPJ and aMTG are more

strongly associated with other-mentalizing, while mPFC and precunes/PCC clusters are

unique in the sense that they are consistently involved in both self- and other-related

mentalizing.

Leave-One-Site-Out classifiers trained on all datasets

Finally, to ensure that the high accuracies were not an artifact of features unique

to the training study (e.g., scanner, sample), we trained new classifiers that combined data

across all five training and testing datasets using a leave-one-site-out (LOSiO) cross-

validation approach. We combined images from n = 232 participants from Studies 1 to 5

that used a trait-evaluation task, pooling data from healthy participants, adolescents, and

clinical samples. We used the same parameters to train SVM models corresponding to
the four mentalizing signatures.

Cross-validated prediction accuracies of all four classifiers were very good to
excellent (see Fig. 9A-D): 83% averaged accuracy (2AFA) for the Self-RS$_{LOSiO}$, 91% for
the Other-RS$_{LOSiO}$, 96% for MS$_{LOSiO}$, and 90% for the SvO-RS$_{LOSiO}$ in two-alternative
forced-choice tests, $p < .001$ (averaged *Cohen's d* for the Self-RS$_{LOSiO}$ = 1.10, for the
Other-RS$_{LOSiO}$ = 1.36, for the MS$_{LOSiO}$ = 0.64, for the SvO-RS$_{LOSiO}$ = 1.19; averaged AUC
for the Self-RS$_{LOSiO}$ = 0.90, for the Other-RS$_{LOSiO}$ = 0.95, for the MS$_{LOSiO}$ = 0.93, for the
SvO-RS$_{LOSiO}$ = 0.93). Thus, these results show that the high prediction accuracies
achieved by the mentalizing signatures were not inflated by task-specific idiosyncrasies
but rather reflect consistent, robust multivariate patterns that are achieved using different
combinations of the training dataset.

Finally, we applied the predictions of the LOSiO classifiers to the inference task
(Study 6) to test how well they classify attributional versus factual and social versus
nonsocial conditions (see Fig. 9E). Both Other-RS$_{LOSiO}$ and MS$_{LOSiO}$ predicted social
attributional (versus factual) statements in Study 6A with 97% accuracy ($p < .0001$;
Cohen's *d* for Other-RS$_{LOSiO}$ = 2.75 and for MS$_{LOSiO}$ = 2.42). In Study 6b, the Other-RS$_{LOSiO}$
separated attributional versus factual statements in the social condition with 84% accuracy
($p < .001$, Cohen's $d = 1.32$), whereas this classification was non-significant in the
nonsocial condition (accuracy = 56%, $p = .48$). The MS$_{LOSiO}$ predicted attributional versus
factual statements in the social condition with 96% accuracy $p < .001$, Cohen's $d = 2.16$)
and in the nonsocial condition with 66% accuracy ($p = .033$, Cohen's $d = 0.73$). Finally,
MS$_{LOSiO}$ and Other-RS$_{LOSiO}$ separated social from non-social inferences with 96% and 94%
accuracy respectively ($p < .001$, Cohen's *d* for MS$_{LOSiO}$ = 2.12, and for Other-
RS$_{LOSiO}$ = 2.5). Together, this illustrates that, similar to the mentalizing signatures, the
LOSiO classifiers are most sensitive to attributional processing in the social context, while
also capturing nonsocial attributional processing to some degree.

449   **Discussion**

Mentalizing—reflecting about others' and one's own internal states—is a
fundamental capacity required to function in a social world. Transdiagnostically,
mentalizing deficits typify an array of psychopathological conditions[3–9]. Here, we
developed novel brain signatures of self- and other-related mentalizing (*Self-RS*, *Other-
RS, and SvO-RS*), as well as mentalizing in general (the '*Mentalizing Signature'* or *MS*),

combining data from eleven independent cohorts (*N*=390). The four signatures showed
good to excellent classification accuracies in independent data and provided several
insights into the brain circuits of trait-based mentalizing. First, mentalizing about traits
appears to coincide with a specific pattern of brain activity, with reliably dissociable
activation patterns for self- and other-related mentalizing, based on both whole-brain
activity as well as within key nodes of the social brain, especially mPFC, precuneus/PCC,
TPJ and aMTG. Second, the signatures significantly predicted mentalizing not only in
healthy adults, but also in adolescents and in individuals with schizophrenia and with
bipolar disorder, demonstrating its utility across different (including clinical) samples.
Third, our results point to the relevance of these mentalizing signatures for a range of
different research questions in social, developmental, and clinical neuroscience, by
showing (i) engagement of the signatures in independent social cognition tasks, (ii) that
the differentiation of self- and other-related mentalizing increases in transition from
adolescence to adulthood, and (iii) that it is less pronounced in individuals with
schizophrenia compared to healthy controls. Thus, our data provide us with clearly defined
brain models[15], applicable across contexts to assess the engagement of mentalizing-
related brain regions in different experimental conditions and their alteration in clinical and
neurodevelopmental conditions.

The Mentalizing Signature (MS) successfully predicted mentalizing across
different contexts and extended to tasks without explicit mentalizing instructions (e.g.,
spontaneous mentalizing). It performed very well in separating higher-level inferences
about others and hidden states from factual inferences, with higher classification in social
compared to non-social contexts. The MS (but not the Other-RS) also significantly
separated social from nonsocial inferences. Together, these findings demonstrate its
sensitivity to mental state compared to external state attributions. The regions that
contributed to the MS most positively included mPFC, precuneus, TPJ, STS, and many
other regions previously associated with mentalizing[10,13,29,38,48], providing convincing face
validity of this signature. Its high predictive performance in testing datasets implies that
mentalizing recruits consistent and reliable neural configurations in the brain, not only in
healthy controls, but also in clinical cohorts and developing adolescents. The MS also
extended to other tasks (social-feedback and attributional versus factual inferences),
predicting conditions that potentially evoke implicit, spontaneous mentalizing. This
suggests that implicit and explicit forms of mentalizing largely share a common neural
basis, in keeping with previous research[49]. Thus, subnetworks of mentalizing (implicit-

explicit) may be building on a common anatomical architecture that is here covered by the
MS.

Recent work proposes that self- and other-mentalizing activate common neural
constellations or representations and recruit largely overlapping regions[29,30,48]. In contrast,
our findings suggest that, while they share a large common core (as evident in the MS),
they can also be distinguished reliably based on both whole-brain and local activation
patterns. The Self-RS and the SvO-RS had positive weights in anterior mentalizing regions
around cortical midline structures, such as mPFC, ACC, thalamus, caudate nucleus,
frontal eye fields, and insula, extending to the frontal operculum. This pattern is in line with
previous meta-analyses[31,34,36]. The mPFC has been consistently related to mentalizing,
and especially its ventral part, with self-related processing[33,39,42,50,51]. The ventral-to-dorsal
linear[31] and curvilinear gradient[32] hypotheses propose that mPFC is involved in both self-
and other-person related processing. Accordingly, our results show that mPFC is involved
in both self- and other-related mentalizing with different neural configurations, and that the
information encoded in mPFC seems to predict self-mentalizing more consistently.
Besides, self-processing recruited reward-processing circuits (e.g., striatum), in line with
the idea that introspection about oneself is intrinsically rewarding[52–54]. The involvement of
subcortical areas (e.g., thalamus) and cortical areas associated with interoception (e.g.,
insula) is in line with previous work suggesting that one has access to
affective/physiological information during self-mentalizing, which are less pertinent during
mentalizing about others[55], and that interoceptive information might be an important
contribution to one's sense of self[26,56,57].

The Other-RS and (negative weights of) the SvO-RS subsumed more posterior
and lateral regions, such as TPJ, precuneus/PCC, STS, and vlPFC, resonating with
previous findings[10,11,13,32,37–39]. Interestingly, our results concerning the precuneus/PCC do
not align with the known subdivisions of the default-mode[58] or mentalizing networks[30]. This
region is thought to be part of the midline core of the default-mode network[58] and of the
medial subsystem of the mentalizing network[30], with stronger associations with self-
processing. However, here in our results, while the precuneus/PCC cluster also contains
information about self-mentalizing, it appears more strongly associated with other-related
mentalizing, along with the areas that are part of the lateral subdivisions of both networks
(e.g., TPJ, temporal lobes). A potential explanation here could be the involvement of
precuneus in mental orientation/imagery[59,60].

The mentalizing signatures generalized to data from different types of cohorts,

including healthy adults, schizophrenia and bipolar samples, and adolescents. This shows

that the overall neurobiological configurations of self- and other-related mentalizing are

comparable across these populations and that the signatures have utility in different types

of study populations. However, the Self-RS, the Other-RS, and the SvO-RS signatures

also showed sensitivity to clinical status, since self/other differentiation in the pattern

responses was significantly less pronounced for both patterns in participants with

schizophrenia compared to matched healthy controls. This finding is noteworthy and in

line with clinical observations of alterations in self-perception, mentalizing, and the ability

to discriminate between self- and other-generated thought[36,45,46,61]. The responses in the

bipolar sample did not differ significantly from those of the healthy adults but given the

limited sample size of the bipolar group in the present analysis, future work is needed to

assess more fine-grained (and potentially context-dependent) differences in mentalizing

responses in bipolar disorder, as well as in other psychiatric and neurodevelopmental

disorders with mentalizing deficits.

In the two adolescent samples (aged 11 to 18 years), greater age was associated

with better self/other differentiation of the Self-RS, the Other-RS, and the SvO-RS. In other

words, responses of these brain signatures to self- and other-related mentalizing were

more different from each other in older adolescents. Considering that the signatures were

developed using an adult dataset, this reflects an ongoing development of the mentalizing

neurobiology[62,63] to become more adult-like and more differentiated for different targets of

mentalization, during the age span covered in our study (11-18 years). Future studies

could test the role of pubertal development instead of chronological age, and test the

signatures in younger children, and those with developmental disorders.

Of note, the mentalizing signatures were also implicated in making inferences

about nonsocial targets—albeit to a lesser degree compared to social targets (see Fig. 6D

and Fig 9E). An earlier study[64] that used a similar inference task as Study 6 found that

social and nonsocial attributions displayed overlapping activations in dmPFC, STS, TPJ,

and lateral orbitofrontal cortex. A common denominator in both conditions may reflect

domain-general attributional processing[65], which at least partially builds on semantic

cognition[66]. Likewise, processes supporting mentalizing may also more broadly subserve

extraction of conceptual and semantic meaning for nonsocial stimuli[67,68], in line with the

default mode network's putative meaning-making function[51,69,70]. Thus, the mentalizing

signatures may involve components of both domain-general representational processing,

domain-specific social processing, and target-specific (self and other) processing
components.

We note that the signatures' classification performance in the cross-validated
training is consistently higher than in the independent testing datasets. This could
potentially be due to leakage of information across the different participants in the training
dataset, who all performed the same task in the same scanner and experimental settings.
However, the additionally developed leave-one-site-out (LOSiO) classifiers that combined
data across five studies performed similarly to the brain signatures that are developed
using data from only the training dataset (Study 1). This provided evidence that the results
of this study are not an artifact of our analytic choices and that the methodological
approach fits the aims of the study well, as the results are robust with different
permutations. The LOSiO classifiers showed consistently high classifications for the
original training dataset (Study 1), which further provided support for the use of this study
for training. Yet, because the LOSiO classifiers used information from n=232 participants
during model development, it has potential to generalize very well to independent
contexts. A key feature of the brain signature approach[15] employed here is that the brain
models are not necessarily final products, but tunable, improvable working models.
Therefore, we endorse the use of LOSiO classifiers in future research along with the main
mentalizing signatures for further evaluation of their utility in diverse settings.

The high separability of self- and other-referential processing in Study 1 might also
reflect the fact that the Other-condition in the mentalizing task was about a relatively
unfamiliar person (a confederate). This may lead to a greater difference in self- versus
other-related processing compared to studies in which mentalizing was about a familiar or
close person—which is often more closely related to the self and likely also engages self-
referential processing. This could potentially also explain why the Other-classifier was less
successful in separating other-related from self-related feedback in the sample of romantic
couples (Study 7), since participants may consider their partners as an extension of
themselves. Relatedly, the lower performance of signatures in Study 5 (compared to other
testing datasets) could be due to the use of physical attributes (in addition to personality
traits), which rely less on mentalizing.

Of note, the mentalizing signatures developed here primarily capture trait-related
mentalizing. Here, we provide initial evidence that they extend beyond explicit reflections
on trait-level representations. However, they also show above-chance classification in
nonsocial inferences (see Fig 6D and Fig. 9E), which might raise concerns about their

specificity to social agents. Future studies are needed to further test how well the

mentalizing signatures separate mentalizing about other agents from other higher-order,

conceptual, and abstract cognitive processing. Future studies should also test their

specificity, as well as their validity in other forms (e.g., affective, spontaneous, or non-

verbal) of mentalizing. A natural next step could be to test whether other dimensions of

mentalizing[6,12] (e.g., cognitive versus affective) or different types of mental state content[71]

(i.e., stable versus transient or beliefs versus preferences) are also associated with robust,

replicable multivariate patterns. Recent work[72,73] investigated valence and self-relevance

as two key components of internal thought. It is an open question how the brain integrates

valence with different targets during mentalizing. Finally, future research can also build on

this work by testing the signatures in other mentalizing tasks (e.g., false-belief, emotion

imagery), on a greater variety of targets[74], and in other related mental processes (e.g.,

autobiographical memory retrieval).

In conclusion, we trained and validated four brain signatures that predict self-

related, other-related, and both types of mentalizing in multiple independent datasets that

used different variants of a standard trait-mentalizing task in diverse populations. Our

findings imply that self- and other-mentalizing use largely dissociable neural mechanisms

that build on a foundational overall mentalizing capacity. Indeed, the four mentalizing

signatures possess potential for use as predictive neuromarkers of mentalizing about self

and others, for example by testing how brain-circuits of mentalizing are engaged or

modulated by different types of experimental conditions, and how they might be altered in

different psychiatric and neurodevelopmental populations.

613   **Methods**

Participants

The present study pooled data from a total of N=390 participants (119 females and

162 males, M_{age}= 25.5, SD_{age}= 12.9) from eleven cohorts participating in seven

independent studies (see Fig. 1A and Supplementary Table 1). These eleven cohorts

comprised six samples of healthy (neurotypical) adults, two samples of healthy

adolescents, and three samples of adults with a diagnosed psychiatric condition. While

most of the datasets have been previously published separately, the analyses reported

here were not published previously, and the seven studies have not been previously

combined.

The training sample (Study 1[43]) consisted of n=21 healthy adults (10 women and

11 men, M_{age} = 23.5) who were recruited at the University of Geneva, Switzerland. One

additional participant with structural abnormalities in the brain was excluded from the

original study and the present analysis.

Study 2[75] included n=44 healthy adolescents (23 females and 21 males, M_{age} = 16,

SD_{age} = 1.86, age range = 12.01 to 18.84) who were recruited from secondary schools in

Geneva, Switzerland. In the original study, one additional participant was excluded due to

structural abnormalities in the brain, three due to non-completion of the paradigm, one

due to signs of substance use, and five due to excessive movement.

Study 3[76] included n=61 healthy adolescents (27 females and 34 males, M_{age} =

12.9, SD_{age} = 0.43, age range = 11.61 to 14.22) who were recruited for a longitudinal

project from secondary schools in the Netherlands. An additional 18 participants were

excluded from the original study due to excessive movement, incorrect task completion,

or measurement errors. Additionally, 5 participants were not included in this study (original

study n = 66) because they did not provide explicit consent for data sharing.

Study 4[77,78] included n=33 healthy adults (14 women and 19 men, M_{age} = 41.7) and

n=23 adults with schizophrenia (7 women and 16 men, M_{age} = 37). The two cohorts were

matched on age, sex, and a measure of general intelligence. The patients were recruited

from a local psychiatric hospital in Barcelona, Spain, based on diagnostic interviews. The

schizophrenia diagnosis was confirmed using the Structured Clinical Interview for DSM

Disorders[79] (SCID).

Study 5[80] included three types of cohorts: Healthy adults (n=15, 6 women and 9

men, M_{age} = 33.3), adults with schizophrenia (SZ, n=17, 6 women and 11 men, M_{age} =

35.5), and adults with bipolar disorder (BD, n=18, 9 women and 9 men, M_{age} = 40.3). The

clinical cohorts were recruited from a local hospital in the north of the Netherlands. The

diagnoses of the patients were confirmed using the Mini International Neuropsychiatric

Interview-Plus[81] 5.0.0 (MINI-Plus). All BD patients were chosen among those who had a

history of at least one psychotic episode. All three cohorts were matched with one another

on age, sex, level of education, and a measure of general intelligence. The SZ and BP

patients were additionally matched on the level of cognitive and clinical insight as

measured by the Schedule of Assessment of Insight[82]-Expanded version (SAI-E, clinical

insight) and the Beck Cognitive Insight Scale[83] (BCIS, cognitive insight).

Study 6a[84] included 59 healthy adults (26 female and 33 male; M_{age}= 28.3), and

Study 6b[84] included 50 (19 female and 31 male; M_{age}= 33.6) healthy adults from the Los

Angeles metropolitan area; these final sample sizes reflect exclusions described below.

One participant from each study was excluded due to poor task performance (>70%

missed trials), and four additional participants from Study 6b were excluded due to outlier

behavioral scores (>3 SDs from the mean, n=1) and excessive scanner motion (n=3). All

participants were right-handed, had normal or corrected-to-normal vision, spoke English

fluently, and had IQs in the normal range (assessed via the WASI-II).

Study 7[85] included 56 healthy adults in romantic relationships recruited from the

Tucson, Arizona community and surrounding areas. Seven participants were excluded

from the study due to missing or inadequate imaging data, yielding a final sample size of

n=49 (26 women and 23 men, M_{age} = 22.6). Community members were eligible to

participate if they had been in a romantic relationship for at least six months, had no

contraindications for MRI scanning, and did not meet criteria for active psychosis or mania

at the time of screening. Both members of the couple completed all components of the

study, including the social feedback fMRI task.

All participants gave written informed consent and were compensated for their

participation via monetary means or gifts. All studies were approved by the respective

institutional ethics committees. For additional information, please refer to the original

studies and Supplementary Table 1.

Tasks

Training dataset

We used a trait evaluation task, adapted from ref[42] as the training mentalizing task.

Participants were asked to rate the extent to which certain personality adjectives (e.g.,

'talkative', 'daring') described themselves (Self-condition) or a same-sex confederate

(Other-condition), with whom they thought they were interacting in a preceding decision-

making task[43]. In the Control condition, they were asked to count the syllables for each

trait adjective.

Participants met the confederate in person at the beginning of the experimental

session, where they were briefed that one of them would be tested in the scanner, the

other one in a separate room, and that they would interact via computer interface. They

first participated in a social decision-making task resembling a serial dictator paradigm,

whereby the participant was given the choice to share or keep the resources with the

confederate in a series of trials[43]. After this task, participants were introduced to the trait-

evaluation task and were asked to rate their co-player.

The task consisted of 150 trials in total, presented in 30 blocks (ten per condition).
Each block contained five trials and lasted for 20 seconds, with an inter-block interval
(fixation cross) of 8 seconds. Half of the blocks only contained negative and the other half
only positive adjectives. Each word appeared once for each condition (50 adjectives were
used in total).

Each trial started with a 3-second cue above the fixation cross, reminding the
participant of the condition (self, other, or syllables), which was followed by the
presentation of the adjective and a Likert scale (1-4) on which the participant was expected
to select using a button press device. A word was presented every 4 seconds. If the
participant made a choice sooner, the word would disappear for the remaining time of the
4 seconds before the next trial. The order of the blocks and words displayed in each block
was randomized.

Testing and extension datasets

All testing datasets included similar mentalizing tasks with the same three
conditions (Self, Other, Control) as the training task and utilized a block design (for an
overview, see Fig. 1A and Supplementary Table 1). The main difference between
validation tasks was the stimuli that were presented: Studies 2 and 3 used trait adjectives,
Study 4 (with two cohorts) used trait statements, and Study 5 (with three cohorts) used a
mix of trait and physical statements. The studies also used a variety of *others* in the Other-
condition: Best/close friend (Study 2), similar and dissimilar classmate (Study 3), relative
or close friend (Study 5), and acquaintance (Study 4). Studies 2 and 5 collected responses
using Likert scales, and Studies 3 and 4 asked for binary choices (yes/no). Finally, the
Control conditions also varied between studies. Studies 4 and 5 presented general
knowledge statements; Study 3 asked participants to search for the letter in words; Study
2 asked them to count the syllables in words.

Extension datasets (Studies 6 and 7) used different tasks to test how well the
mentalizing signatures extend to tasks that involve other forms of social (or nonsocial)
processing. Studies 6a and 6b used an inference task based on visual stimuli[84] (see Fig.
6A). In Study 6a, participants made rapid yes/no judgments in response to social scenes
(intentional hand actions and emotional faces) that required either attributional inferences
about hidden/internal states (mentalizing) or factual inferences about observable features,
resulting in a 1 (stimulus type: social) × 2 (inference type: attributional, factual) factorial
design. Study 6b included an additional nonsocial condition using images of natural

scenes (e.g., rain pouring out of a gutter), each paired with attributional or factual inference

prompts, resulting in a 2 (stimulus type: social, nonsocial) × 2 (inference type: attributional,

factual) design. Study 7 (see Fig. 7A) investigated spontaneous empathy when negative

or positive feedback thought to be from other participants (confederates) was directed to

participants' romantic partners (or one's self). Participants received positive and negative

feedback either about themselves (self-condition), their romantic partners (other-

condition), or no feedback (control-condition). The original study (see ref[85] for more

details) included more conditions, such as directing feedback to both self and partner,

which are not analyzed in the present manuscript.

fMRI data acquisition, preprocessing, and first-level analysis

Training dataset

The training MRI images were acquired on a 3T Magnetom TIM Trio whole-body

scanner (Siemens, Germany) with the product 12-channel head coil. A T1-weighted

MPRAGE sequence (TR = 1900ms, TI = 900 ms, TE = 2.27 ms, voxel size 1 x 1 x 1 mm)

was used to acquire structural anatomical images. Functional images were obtained using

a standard T2-weighted echo-planar imaging sequence (2D-EP, TR = 2100 ms, TE = 30

ms, flip angle 80°, voxel size 3.2 x 3.2 x 3.2 mm) that scanned the whole brain in 36

sequential slices. An automated shimming procedure was included to minimize magnetic

field inhomogeneities.

As it is the standard practice in the development of brain signatures[15,17,18,21], we

used statistical maps from first-level analysis (e.g., single-subject contrast images) as

input for training and testing the classifiers. SPM8 (Wellcome Department of Imaging

Neuroscience, UCL, London, UK) and Matlab® (The MathWorks Inc.) were used for image

preprocessing and first-level analysis. A standard preprocessing pipeline was performed,

which included spatial realignment and reslicing of the functional images, coregistration,

unified segmentation and normalization to the Montreal Neurological Institute (MNI) echo

planar imaging template (voxel size: 2 mm^3), and finally spatial smoothing using an 8 mm

Full Width at Half Maximum (FWHM) Gaussian kernel.

During first-level analysis, we included six task (block) regressors that were

composed of 3 task conditions by positive and negative valence. The task regressors were

convolved with a canonical hemodynamic response function. We also included six

additional regressors of no interest to control for motion. A high-pass frequency filter

(128s) and autocorrelation corrections (using restricted maximum likelihood and an

autoregressive model) were used in model estimation.

Testing datasets

We conducted a non-systematic literature review to identify recent fMRI studies of

comparable mentalizing tasks with at least three conditions (Self, Other, and non-

mentalizing Control condition) and in which participants rated trait adjectives or

statements. We emailed the authors of seven studies that we identified and received

positive responses from four of them. Data from these four studies were included in the

current study as testing datasets. In addition, to test the signatures in a different type of

task, we included an unpublished dataset[85] (Study 7) and data from recently published

work[84] (Study 6) as extension datasets. Please refer to the original studies (see

Supplementary Table 1) for details of image acquisition, preprocessing, and first-level

analysis.

The authors provided the subject-level contrast images of the different

experimental conditions (versus the implicit baseline). Voxel weights were normalized

within each image using L2-norm, and contrast images were resampled onto the same

image space as the training dataset using linear resampling. For Study 3, which included

two different Other-conditions (similar and dissimilar classmates), we averaged the

contrast images across these two conditions, resulting in a single Other-condition, as in

the training and the other testing datasets (analyzing them separately did not alter the

results). Study 6 included two types of social conditions: hands and faces. As we were not

interested in the social processing of hand- or face-related information, we averaged the

contrast images of these conditions, creating a single Social Condition image per

participant.

Data Analysis

Cross-validated training

Using 10-fold cross-validation, we trained three support-vector-machine (SVM)

classifiers in Study 1 that discriminate each condition from the other two conditions: The

Self-RS was trained to separate the Self-condition from Other and Control conditions, the

Other-RS to separate the Other-condition from Self and Control conditions, and the MS

was trained to separate both mentalizing conditions (Self and Other) from the Control

condition. In addition, we trained one signature to specifically separate the Self from the

Other condition (the SvO Signature). The input for SVMs was subject-level contrast

images for each of the three task conditions (versus implicit baseline). The cross-validation

folds were organized to ensure that the three contrast images from each participant were

kept together within a single fold (nine folds with two participants and one fold with three

participants). To reduce the possibility that classifiers opportunistically used non-

mentalizing related processes (e.g., visual information), we applied a mask in the training

dataset that includes key social-cognition regions (see Fig. 1B). This mask was computed

as the union of six term-based meta-analytic maps (association and uniformity maps for

"mentalizing", "self-referential", and "social", downloaded from NeuroSynth[47] on

06/06/2024) to cover a wide range of brain areas typically activated by social and self-

related processing. While the use of a mask was intended to minimize involvement of the

confounding voxels that are not directly related to social cognition, this might raise the

question of how more extensive, whole-brain classifiers might perform. As shown in

Supplementary Fig. 1, applying a whole-brain gray matter mask yielded highly similar

results.

SVMs were chosen based on their high performance for binary linear classification

problems in high-dimensional and low-sample data[86–88]. Linear SVM is a commonly used

method in cognitive neuroscience in decoding problems[48,89,90], which is computationally

efficient and provides interpretable, robust weight maps. The algorithm fits a hyperplane

that classifies true and false classes by assigning weights for each feature (i.e., each

voxel). The fitting is performed for each of 10 folds, in which the data of 90% of participants

are used for fitting the classifier, and the resulting classifier is tested on the remaining 10%

hold-out participants' data, allowing for assessing its classification performance in

independent hold-out data. Because using a one versus the rest approach in SVM (e.g.,

Self versus Other and Control, see Fig. 1C) may add bias into the model by favoring the

majority class, we fitted weighted SVM models with a ridge parameter of .5 to train the

Self-RS, the Other-RS, and the MS (but not the SvO-RS). To avoid overfitting, the SVMs

were otherwise trained using default parameters (regularization parameter C = 1, linear

kernel function, number of folds = 10). The cross-validated distance from the hyperplane

of hold-out images was used to calculate the receiver operating characteristic (ROC)

curves and the accuracy for each classification. Each SVM results in a weight map with

one value per voxel. These voxel weights are effectively the predictive weights of the true

class, yielding brain signatures of mentalizing that can be applied to other brain images to

obtain a single pattern expression score per image. The classification accuracies in the

training dataset were tested using two-alternative forced-choice predictions and binomial

tests.

We used bootstrapping to train SVM classifiers, producing stable model estimates

and voxel-wise p-values. In the first step, cross-validated training generated prediction

accuracies and a final model output (i.e., weight map) that is trained on all samples. In the

next step, the bootstrapping procedure resampled the training data 5000 times with

replacement, yielding statistical maps with two-tailed, uncorrected p-values. These maps

were then thresholded using FDR-correction at $q < .05$ with a minimum cluster size of 10

voxels. Note that corrected maps were only used for visualization and inference. The

unthresholded weight maps constitute the mentalizing signatures and, therefore, are used

for analysis.

Validation in independent datasets

The four mentalizing signatures were applied to the testing datasets by computing

the pattern similarity values as the matrix dot product between mentalizing signatures and

the subject-level (1st level) contrast images of each study, yielding one scalar value per

condition and participant and signature. The predictions followed a binary forced-choice

principle using paired observations. The signatures' classification accuracies were

assessed using Receiver Operating Characteristic (ROC) analysis and binomial tests

using a two-sided significance threshold of $p < .05$. The accuracy rate, area under the

curve (AUC), and specificity/sensitivity obtained from these analyses provide the

benchmark measures of binary classifications in multivariate modeling[44]. We also report

Cohen's d in the text as a more standardized measure to quantify prediction effect sizes

across studies and samples. Cohen's d is calculated as the mean of differences between

condition values (either pattern expression or distance from the hyperplane) divided by

the pooled standard deviation of differences (where difference =

input_values[binary_class] - input_values[~binary_class]). Because the three signatures

(Self-RS, Other-RS, and MS) were each tested in two comparisons (e.g., Self versus

Other and Self versus Control), the main text reports the average statistical values across

these comparisons, with the complete results provided in the Supplementary

Tables. Similarly, when accuracy statistics from a classifier were combined across

conditions or studies, we report the weighted average values in the main text and present

the full set of statistics in the Supplementary Material.

To examine the effects of individual differences (e.g., sex and age) on pattern

expression values, we used linear mixed-effects models. We pooled data from five testing

datasets that used the same task (n = 211; Studies 2–5). The signature fit ("true minus

false class score") was defined as the difference between the pattern expression for the

true condition and that for the false condition. Specifically, we subtracted the false-

condition value from the true-condition value for each signature: Self-RS (Self – Other),

Other-RS (Other – Self), SvO-RS (Self – Other), and MS ([Self + Other]/2 – Control). Age

and sex were entered as fixed effects, and study was included as a random intercept in

the mode: *fit ~ age + sex + (1 | study)*.

Group comparisons of pattern expression values. To quantify how well

signatures discriminate true versus false conditions and to easily compare self/other

separation across groups, we used the signature fits for each signature, the calculation of

which is explained in the section above. For statistical comparisons between participants

with schizophrenia and healthy controls, we ran linear mixed effects models for each of

the signatures, with the signature fits ('true minus false class score') as the dependent

variable. Group constituted the fixed effect variable (healthy controls [n=48] versus

schizophrenia sample [n=40]). Because the data came from two different studies (Study

4 and Study 5), study was included as a random effect variable. Therefore, the model

equation was "*signature_fit ~ group + (1|study)*".

To test whether the bipolar sample (n=18) differed from healthy adults (n=15) in

Study 5 regarding their pattern expression values, we performed independent samples t-

tests for each of the signatures consecutively. The dependent variable was the "signature

fit" as above.

Associations with age. We tested whether the adolescents' age was associated

with self-other discrimination using linear mixed effects models. We included age as the

fixed effect, Study (Study 2 and Study 3) as the random effect, and sex as the control

variable (fixed effect). The dependent variable was the difference of the true minus false

classes for each of the signatures, as detailed above. Therefore, the model formula was

"*signature_fit ~ age + sex + (1|study)*".

Testing the signatures in extension tasks

In order to test how well the signatures generalized to the two extension tasks, we

applied them to the subject-level contrast images for each experimental condition by

computing their dot products. When a signature is applied to another context (e.g., a

different task), one could apply either i) a binary prediction test or ii) a linear-effects model

to test for condition differences, depending on the study aim. Binary prediction is arguably

most appropriate when there is a ground truth or clear directional expectation, whereas

linear models are better suited to more distant applications (e.g., novel tasks or

populations). Accordingly, we tested the signatures in Study 6 using i) binary prediction

tests, expecting the signatures to selectively predict the attributional statements involving

mentalizing, consistent with the original study[84]. We followed the same principles for

predictions as outlined above (validation in independent tests), where pattern expression

values and predictions are subjected to ROC analyses and binomial tests. In Study 7, we

instead used ii) linear-effects models, submitting pattern expression values to

repeated-measures ANOVAs across the three social-feedback conditions, followed by

Bonferroni-corrected t-tests for pairwise comparisons.

Region-of-Interest (ROI) Analyses

To test the ability of local patterns to predict mentalizing and to separate self-

related versus other-related mentalizing, we trained and cross-validated ROI classifiers

using a parallel approach to the whole-brain classifiers. We downloaded a term-based

meta-analytic map for 'Mentalizing' from NeuroSynth[47] on 14/09/2022 that included 151

studies. We selected clusters that contained more than 200 voxels, resulting in the

following ten ROIs: mPFC, bilateral TPJ, bilateral anterior MTG, precuneus/PCC, right

SMA, and three clusters in the cerebellum. In each ROI, we trained and 10-fold cross-

validated four classifiers in the training dataset and validated them in the remaining

datasets, to (i) predict Self-mentalizing (versus the two remaining conditions), (ii) predict

Other-mentalizing (versus the two remaining conditions), (iii) predict mentalizing (Self- and

Other-mentalizing versus Control condition), and (iv) differentiate Self- versus Other-

processing. We tested these ROI classifiers in the testing datasets using the same

parameters and analytic approach as outlined above in the main whole-brain analyses.

Leave-One-Site-Out classifiers trained on all datasets

To fully benchmark how well our approach generalizes across independent

datasets, we trained additional classifiers based on the pooled cohorts from all datasets

that used a variation of the trait evaluation task (Studies 1 to 5; *n*=232). We defined five

folds, each representing one study, so that the classifiers were trained on four studies and

tested on the remaining study in each fold. The cross-validated training of bootstrapped

classifiers followed the same analytic flow as the main classifiers (see cross-validated

training above). Besides the number of folds, the only other change was the number of

bootstrapped samples, which we limited to 2500 samples for computational efficiency. We

additionally tested these classifiers in the inference task from Study 6 using the binary

prediction tests as described above for the main signatures.

General statistical approach

All data analysis and data visualizations were performed using Matlab® R2022b

software and the Canlab toolbox (https://github.com/canlab). Violin plots in Fig.4 and Fig.

7 were generated using the MATLAB function daviolinplot[91]. Scatter plots in Fig. 5 were

generated in R Statistical Software[92] (v12.1) using ggplot2[93]. Statistical inference used a

significance threshold of $p < 0.05$, unless otherwise noted.

Data availability statement

Source data for all figures and single-subject contrast images of the training

dataset (Study 1) are available at https://figshare.com/s/86b77a31fec7cd0e0925.

Contrast images from Study 4a can be accessed via OpenNeuro

(https://openneuro.org/datasets/ds001618/versions/1.0.1) and data from Study 6 are

available through the NIHM Archive (https://nda.nih.gov/edit_collection.html?id=2643).

Deidentified data from Studies 2, 3, 4b, 5, and 7 can be made available in line with

applicable data protection and privacy regulations upon reasonable request to the

corresponding authors.

Code availability statement

Mentalizing signatures for use in future studies, as well as custom MATLAB scripts

for analyses are available at: https://github.com/ldmk/2025_MentalizingSignatures

Author Contributions

LK, TDW, JAH, and MLS conceived the project. DA and LK conceptualized

the research goals, built the collaborations, analyzed the data, and prepared the first draft

of the manuscript. DA created the figures. LK, JAH, DAS, AMC, MvB, LK, LZ, LvdM, PFC,

EPC, RS, MD, PVr, AT, and PVu designed the experiments and provided data. LK, DA,

and LOW acquired funding for the project. LK supervised the project. All authors

contributed to the writing and editing of the paper.

References

1.   Moore, C. & Frye, D. The Acquisition and Utility of Theories of Mind. in *Children's*

Theories of Mind: Mental States and Social Understanding (eds Frye, D. & Moore,

C.) 1–14 (Lawrence Erlbaum, Hillsdale, NJ, 1991).

2.   Wellman, H. M. *Making Minds: How Theory of Mind Develops*. (Oxford University

Press, 2014).

3.   Brüne, M. & Brüne-Cohrs, U. Theory of mind—evolution, ontogeny, brain

mechanisms and psychopathology. *Neurosci. Biobehav. Rev.* **30**, 437–455 (2006).

4.   Debbané, M. *et al.* Attachment, Neurobiology, and Mentalizing along the Psychosis

Continuum. *Front. Hum. Neurosci.* **10**, (2016).

5.   Gray, K., Jenkins, A. C., Heberlein, A. S. & Wegner, D. M. Distortions of mind

perception in psychopathology. *Proc. Natl. Acad. Sci.* **108**, 477–479 (2011).

6.   Luyten, P., Campbell, C., Allison, E. & Fonagy, P. The Mentalizing Approach to

Psychopathology: State of the Art and Future Directions. *Annu. Rev. Clin. Psychol.*

16, 297–325 (2020).

7.   Sharp, C. Mentalizing Problems in Childhood Disorders. in *Handbook of*

Mentalization-Based Treatment 101–121 (John Wiley & Sons, Ltd, 2006).

doi:10.1002/9780470712986.ch4.

8.   Sloover, M., van Est, L. A. C., Janssen, P. G. J., Hilbink, M. & van Ee, E. A meta-

analysis of mentalizing in anxiety disorders, obsessive-compulsive and related

disorders, and trauma and stressor related disorders. *J. Anxiety Disord.* **92**, 102641

(2022).

9.   Johnson, B. N., Kivity, Y., Rosenstein, L. K., LeBreton, J. M. & Levy, K. N. The

association between mentalizing and psychopathology: A meta-analysis of the

reading the mind in the eyes task across psychiatric disorders. *Clin. Psychol. Sci.*

Pract. **29**, 423–439 (2022).

10. Frith, C. D. & Frith, U. The Neural Basis of Mentalizing. *Neuron* **50**, 531–534

(2006).

11. Saxe, R. Uniquely human social cognition. *Curr. Opin. Neurobiol.* **16**, 235–239

(2006).

12. Schurz, M. *et al.* Toward a hierarchical model of social cognition: A neuroimaging

meta-analysis and integrative review of empathy and theory of mind. *Psychol. Bull.*

147, 293–327 (2021).

13. Van Overwalle, F. & Baetens, K. Understanding others' actions and goals by mirror

and mentalizing systems: A meta-analysis. *NeuroImage* **48**, 564–584 (2009).

14. Yang, D. Y.-J., Rosenblau, G., Keifer, C. & Pelphrey, K. A. An integrative neural

model of social perception, action observation, and theory of mind. *Neurosci.*

Biobehav. Rev. **51**, 263–275 (2015).

15. Kragel, P. A., Koban, L., Barrett, L. F. & Wager, T. D. Representation, Pattern

Information, and Brain Signatures: From Neurons to Neuroimaging. *Neuron* **99**, 257–

273 (2018).

16. Rosenberg, M. D., Casey, B. J. & Holmes, A. J. Prediction complements explanation

in understanding the developing brain. *Nat. Commun.* **9**, 589 (2018).

17. Wager, T. D. *et al.* An fMRI-Based Neurologic Signature of Physical Pain. *N. Engl.*

J. Med. **368**, 1388–1397 (2013).

18. Koban, L., Wager, T. D. & Kober, H. A neuromarker for drug and food craving

distinguishes drug users from non-users. *Nat. Neurosci.* **26**, 316–325 (2023).

19. Rosenberg, M. D. *et al.* A neuromarker of sustained attention from whole-brain

functional connectivity. *Nat. Neurosci.* **19**, 165–171 (2016).

20. Kim, J. *et al.* A dorsomedial prefrontal cortex-based dynamic functional connectivity

model of rumination. *Nat. Commun.* **14**, 3540 (2023).

21. López-Solà, M. *et al.* Towards a neurophysiological signature for fibromyalgia. *Pain*

158, 34–47 (2017).

22. Gabrieli, J. D. E., Ghosh, S. S. & Whitfield-Gabrieli, S. Prediction as a Humanitarian

and Pragmatic Contribution from Human Cognitive Neuroscience. *Neuron* **85**, 11–26

(2015).

23. Koban, L. *et al.* An fMRI-Based Brain Marker of Individual Differences in Delay

Discounting. *J. Neurosci.* **43**, 1600–1613 (2023).

24. Woo, C. W., Chang, L. J., Lindquist, M. A. & Wager, T. D. Building better

biomarkers: brain models in translational neuroimaging. *Nat. Neurosci.* **20**, 365

(2017).

25. Tamir, D. I. & Thornton, M. A. Modeling the Predictive Social Mind. *Trends Cogn.*

Sci. **22**, 201–212 (2018).

26. Qin, P., Wang, M. & Northoff, G. Linking bodily, environmental and mental states in

the self—A three-level model based on a meta-analysis. *Neurosci. Biobehav. Rev.*

115, 77–95 (2020).

27. Corradi-Dell'Acqua, C., Hofstetter, C. & Vuilleumier, P. Cognitive and affective

theory of mind share the same local patterns of activity in posterior temporal but not

medial prefrontal cortex. *Soc. Cogn. Affect. Neurosci.* **9**, 1175–1184 (2014).

28. Quesque, F. *et al.* Defining key concepts for mental state attribution. *Commun.*

Psychol. **2**, 1–5 (2024).

29. Tan, K. M. *et al.* Electrocorticographic evidence of a common neurocognitive

sequence for mentalizing about the self and others. *Nat. Commun.* **13**, 1919 (2022).

30. Wang, Y. *et al.* A large-scale structural and functional connectome of social

mentalizing. *NeuroImage* **236**, 118115 (2021).

31. Denny, B. T., Kober, H., Wager, T. D. & Ochsner, K. N. A meta-analysis of

functional neuroimaging studies of self- and other judgments reveals a spatial

gradient for mentalizing in medial prefrontal cortex. *J. Cogn. Neurosci.* **24**, 1742–

1752 (2012).

32. Parelman, J. M. *et al.* Overlapping Functional Representations of Self- and Other-

Related Thought are Separable Through Multivoxel Pattern Classification. *Cereb.*

Cortex https://doi.org/10.1093/CERCOR/BHAB272 (2021)

doi:10.1093/CERCOR/BHAB272.

33. Fossati, P. *et al.* In Search of the Emotional Self: An fMRI Study Using Positive and

Negative Emotional Words. *Am. J. Psychiatry* **160**, 1938–1945 (2003).

34. Murray, R. J., Debbané, M., Fox, P. T., Bzdok, D. & Eickhoff, S. B. Functional

connectivity mapping of regions associated with self- and other-processing. *Hum.*

Brain Mapp. **36**, 1304–1324 (2014).

35. Northoff, G. *et al.* Self-referential processing in our brain—A meta-analysis of

imaging studies on the self. *NeuroImage* **31**, 440–457 (2006).

36. Van Der Meer, L., Costafreda, S., Aleman, A. & David, A. S. Self-reflection and the

brain: A theoretical review and meta-analysis of neuroimaging studies with

implications for schizophrenia. *Neurosci. Biobehav. Rev.* **34**, 935–946 (2010).

37. Arioli, M., Cattaneo, Z., Parimbelli, S. & Canessa, N. Relational vs representational

social cognitive processing: a coordinate-based meta-analysis of neuroimaging data.

Soc. Cogn. Affect. Neurosci. **18**, nsad003 (2023).

38. Tamir, D. I., Bricker, A. B., Dodell-Feder, D. & Mitchell, J. P. Reading fiction and

reading minds: the role of simulation in the default network. *Soc. Cogn. Affect.*

Neurosci. **11**, 215–224 (2016).

39. Wagner, D. D., Chavez, R. S. & Broom, T. W. Decoding the neural representation of

self and person knowledge with multivariate pattern analysis and data-driven

approaches. *Wiley Interdiscip. Rev. Cogn. Sci.* **10**, (2019).

40. Arioli, M., Cattaneo, Z., Ricciardi, E. & Canessa, N. Overlapping and specific neural

correlates for empathizing, affective mentalizing, and cognitive mentalizing: A

coordinate-based meta-analytic study. *Hum. Brain Mapp.* **42**, 4777–4804 (2021).

41. Molapour, T. *et al.* Seven computations of the social brain. *Soc. Cogn. Affect.*

Neurosci. **16**, 745–760 (2021).

42. Kelley, W. M. *et al.* Finding the Self? An Event-Related fMRI Study. *J. Cogn.*

Neurosci. **14**, 785–794 (2002).

43. Koban, L., Pichon, S. & Vuilleumier, P. Responses of medial and ventrolateral

prefrontal cortex to interpersonal conflict for resources. *Soc. Cogn. Affect. Neurosci.*

9, 561–569 (2014).

44. Scheinost, D. *et al.* Ten simple rules for predictive modeling of individual differences

in neuroimaging. *NeuroImage* **193**, 35–45 (2019).

45. Bora, E., Yucel, M. & Pantelis, C. Theory of mind impairment in schizophrenia:

Meta-analysis. *Schizophr. Res.* **109**, 1–9 (2009).

46. Sprong, M., Schothorst, P., Vos, E., Hox, J. & Engeland, H. V. Theory of mind in

schizophrenia: Meta-analysis. *Br. J. Psychiatry* **191**, 5–13 (2007).

47. Yarkoni, T., Poldrack, R. A., Nichols, T. E., Van Essen, D. C. & Wager, T. D. Large-

scale automated synthesis of human functional neuroimaging data. *Nat. Methods* **8**,

665–670 (2011).

48. Oosterwijk, S., Snoek, L., Rotteveel, M., Barrett, L. F. & Scholte, H. S. Shared states:

using MVPA to test neural overlap between self-focused emotion imagery and other-

focused emotion understanding. *Soc. Cogn. Affect. Neurosci.* **12**, 1025–1035 (2017).

49. Van Overwalle, F. & Vandekerckhove, M. Implicit and explicit social mentalizing:

dual processes driven by a shared neural network. *Front. Hum. Neurosci.* **7**, (2013).

50. Heatherton, T. F. *et al.* Medial prefrontal activity differentiates self from close others.

Soc. Cogn. Affect. Neurosci. **1**, 18–25 (2006).

51. Koban, L., Gianaros, P. J., Kober, H. & Wager, T. D. The self in context: brain

systems linking mental and physical health. *Nat. Rev. Neurosci.*

https://doi.org/10.1038/s41583-021-00446-8 (2021) doi:10.1038/s41583-021-00446-

8.

52. Chavez, R. S., Heatherton, T. F. & Wagner, D. D. Neural Population Decoding

Reveals the Intrinsic Positivity of the Self. *Cereb. Cortex* cercor;bhw302v1 (2017)

doi:10.1093/cercor/bhw302.

53. Northoff, G. & Hayes, D. J. Is Our Self Nothing but Reward? *Biol. Psychiatry* **69**,

1019–1025 (2011).

54. Tamir, D. I. & Mitchell, J. P. Disclosing information about the self is intrinsically

rewarding. *Proc. Natl. Acad. Sci.* **109**, 8038–8043 (2012).

55. Maresh, E. L. & Andrews-Hanna, J. R. Putting the "Me" in "Mentalizing": Multiple

Constructs Describing Self Versus Other During Mentalizing and Implications for

Social Anxiety Disorder. in *The Neural Basis of Mentalizing* (eds Gilead, M. &

Ochsner, K. N.) 629–658 (Springer International Publishing, Cham, 2021).

doi:10.1007/978-3-030-51890-5_33.

56. Babo-Rebelo, M. & Tallon-Baudry, C. Interoceptive signals, brain dynamics, and

subjectivity. in *The Interoceptive Mind: From Homeostasis to Awareness* (eds

Tsakiris, M. & De Preester, H.) 46–62 (Oxford University Press, Oxford, UK, 2018).

57. Garfinkel, S. N., Nagai, Y., Seth, A. K. & Critchley, H. D. Neuroimaging Studies of

Interoception and Self-Awareness. in *Neuroimaging of Consciousness* (eds Cavanna,

A. E., Nani, A., Blumenfeld, H. & Laureys, S.) 207–224 (Springer, Berlin,

Heidelberg, 2013). doi:10.1007/978-3-642-37580-4_11.

58. Andrews-Hanna, J. R., Reidler, J. S., Sepulcre, J., Poulin, R. & Buckner, R. L.

Functional-Anatomic Fractionation of the Brain's Default Network. *Neuron* **65**, 550–

562 (2010).

59. Peer, M., Salomon, R., Goldberg, I., Blanke, O. & Arzy, S. Brain system for mental

orientation in space, time, and person. *Proc. Natl. Acad. Sci.* **112**, 11072–11077

(2015).

60. Schurz, M., Radua, J., Aichhorn, M., Richlan, F. & Perner, J. Fractionating theory of

mind: A meta-analysis of functional brain imaging studies. *Neurosci. Biobehav. Rev.*

42, 9–34 (2014).

61. Potvin, S., Gamache, L. & Lungu, O. A Functional Neuroimaging Meta-Analysis of

Self-Related Processing in Schizophrenia. *Front. Neurol.* **10**, (2019).

62. Crone, E. A. & Fuligni, A. J. Self and Others in Adolescence. *Annu. Rev. Psychol.*

71, 447–469 (2020).

63. Fehlbaum, L. V., Borbás, R., Paul, K., Eickhoff, S. B. & Raschle, N. M. Early and

late neural correlates of mentalizing: ALE meta-analyses in adults, children and

adolescents. *Soc. Cogn. Affect. Neurosci.* **17**, 351–366 (2022).

64. Spunt, R. P. & Adolphs, R. Folk Explanations of Behavior: A Specialized Use of a

Domain-General Mechanism. *Psychol. Sci.* **26**, 724–736 (2015).

65. Spunt, R. P. & Adolphs, R. A new look at domain specificity: insights from social

neuroscience. *Nat. Rev. Neurosci.* **18**, 559–567 (2017).

66. Binney, R. J. & Ramsey, R. Social Semantics: The role of conceptual knowledge and

cognitive control in a neurobiological model of the social brain. *Neurosci. Biobehav.*

Rev. **112**, 28–38 (2020).

67. Andrews-Hanna, J. R. & Grilli, M. D. Mapping the Imaginative Mind: Charting New

Paths Forward. *Curr. Dir. Psychol. Sci.* **30**, 82–89 (2021).

68. Baetens, K., Ma, N., Steen, J. & Van Overwalle, F. Involvement of the mentalizing

network in social and non-social high construal. *Soc. Cogn. Affect. Neurosci.* **9**, 817–

824 (2014).

69. Andrews-Hanna, J. R., Smallwood, J. & Spreng, R. N. The default network and self-

generated thought: component processes, dynamic control, and clinical relevance.

Ann. N. Y. Acad. Sci. **1316**, 29–52 (2014).

70. Yeshurun, Y., Nguyen, M. & Hasson, U. The default mode network: where the

idiosyncratic self meets the shared social world. *Nat. Rev. Neurosci.* **22**, 181–192

(2021).

71. Defendini, A. & Jenkins, A. C. Dissociating neural sensitivity to target identity and

mental state content type during inferences about other minds. *Soc. Neurosci.* **18**,

103–121 (2023).

72. Kim, H. J., Lux, B. K., Lee, E., Finn, E. S. & Woo, C.-W. Brain decoding of

spontaneous thought: Predictive modeling of self-relevance and valence using

personal narratives. *Proc. Natl. Acad. Sci.* **121**, e2401959121 (2024).

73. Kim Lux, B., Andrews-Hanna, J. R., Han, J., Lee, E. & Woo, C.-W. When self comes

to a wandering mind: Brain representations and dynamics of self-generated concepts

in spontaneous thought. *Sci. Adv.* **8**, eabn8616 (2022).

74. Courtney, A. L. & Meyer, M. L. Self-Other Representation in the Social Brain

Reflects Social Connection. *J. Neurosci.* **40**, 5616–5627 (2020).

75. Debbané, M. *et al.* Brain activity underlying negative self- and other-perception in

adolescents: The role of attachment-derived self-representations. *Cogn. Affect. Behav.*

Neurosci. **17**, 554–576 (2017).

76. van Buuren, M. *et al.* Neural correlates of self- and other-referential processing in

young adolescents and the effects of testosterone and peer similarity. *NeuroImage*

219, 117060 (2020).

77. Fuentes-Claramonte, P. *et al.* Shared and differential default-mode related patterns of

activity in an autobiographical, a self-referential and an attentional task. *PLOS ONE*

14, e0209376 (2019).

78. Fuentes-Claramonte, P. *et al.* Brain imaging correlates of self- and other-reflection in

schizophrenia. *NeuroImage Clin.* **25**, 102134 (2020).

79. First, M. B. Structured Clinical Interview for the DSM (SCID). in *The Encyclopedia*

of Clinical Psychology 1–6 (John Wiley & Sons, Ltd, 2015).

doi:10.1002/9781118625392.wbecp351.

80. Zhang, L., Opmeer, E. M., Ruhé, H. G., Aleman, A. & van der Meer, L. Brain

activation during self- and other-reflection in bipolar disorder with a history of

psychosis: Comparison to schizophrenia. *NeuroImage Clin.* **8**, 202–209 (2015).

81. Sheehan, D. V. *et al.* The Mini-International Neuropsychiatric Interview (M.I.N.I.):

the development and validation of a structured diagnostic psychiatric interview for

DSM-IV and ICD-10. *J. Clin. Psychiatry* **59 Suppl 20**, 22-33;quiz 34-57 (1998).

82. Kemp, R. & David, A. Insight and compliance. in *Treatment compliance and the*

therapeutic alliance (ed. Blackwell, B.) 61–84 (Harwood Academic Publishers,

1997).

83. Beck, A. T., Baruch, E., Balter, J. M., Steer, R. A. & Warman, D. M. A new

instrument for measuring insight: the Beck Cognitive Insight Scale. *Schizophr. Res.*

68, 319–329 (2004).

84. Tusche, A., Spunt, R. P., Paul, L. K., Tyszka, J. M. & Adolphs, R. Neural signatures

of social inferences predict the number of real-life social contacts and autism

severity. *Nat. Commun.* **14**, 4399 (2023).

85. Ma, S. *et al.* When Empathy Gets Tough: Neural Responses to Overcoming the Self

in a Novel Paradigm Predict Everyday Prosocial Behavior. (2024)

doi:10.31234/osf.io/qcj45.

86. Cox, D. D. & Savoy, R. L. Functional magnetic resonance imaging (fMRI) "brain

reading": detecting and classifying distributed patterns of fMRI activity in human

visual cortex. *NeuroImage* **19**, 261–270 (2003).

87. Misaki, M., Kim, Y., Bandettini, P. A. & Kriegeskorte, N. Comparison of

multivariate classifiers and response normalizations for pattern-information fMRI.

NeuroImage **53**, 103–118 (2010).

88. Mourão-Miranda, J., Bokde, A. L. W., Born, C., Hampel, H. & Stetter, M.

Classifying brain states and determining the discriminating activation patterns:

Support Vector Machine on functional MRI data. *NeuroImage* **28**, 980–995 (2005).

89. Woo, C.-W. *et al.* Separate neural representations for physical pain and social

rejection. *Nat. Commun.* **5**, 5380 (2014).

90. Steardo, L. *et al.* Application of Support Vector Machine on fMRI Data as

Biomarkers in Schizophrenia Diagnosis: A Systematic Review. *Front. Psychiatry* **11**,

588 (2020).

91. Karvelis, P. violin and raincloud plots (https://github.com/povilaskarvelis/DataViz),

GitHub.

92. R Core Team. R: A language and environment for statistical computing. R

Foundation for Statistical Computing, Vienna, Austria. (2024).

93. Wickham, H. ggplot2: Elegant Graphics for Data Analysis. Springer-Verlag. (2016).

1211

1212

1213   **Figures**

1214
Figure 1. Study design and analytic approach. A) The present study included data from seven
independent studies (eleven cohorts, total *N*=390). Training and testing datasets included a trait-
evaluation task with three conditions: A Self-condition, an Other-condition, and a nonsocial Control
condition. Extension datasets constituted the novel instances to test signatures in distinct tasks. B)
Display of the inclusive mask of brain areas related to social cognition. The single-subject contrast
images included in this study are masked using this mask. For results with a whole-brain grey
matter mask, please refer to Supplementary Fig. 1. C) The training dataset (Study 1) included
contrast images of *n* = 21 participants performing a trait-evaluation task with three conditions (Self,
Other, Control). We used a 10-fold cross-validated linear SVM to train four mentalizing signatures.

The Self-Referential Signature (Self-RS) was trained to predict the Self condition versus the two

remaining conditions. The Other-Referential Signature (Other-RS) was trained to predict the Other

condition versus the two remaining conditions. The mentalizing signature (MS) was trained to

separate the Self and Other conditions from the Control condition. The Self-vs-Other Referential

Signature (SvO-RS) was trained to separate the Self and Other conditions. For the validation in

independent datasets, we applied the signatures to the single-subject contrast images from Studies

2-7, by computing the dot product between the weight maps and the contrast images.

1231

1232

1233

Figure 2. Model weights, cross-validated training, and validation results of the mentalizing signatures. On the top row of each box, weight maps illustrate the positive and negative weights of A) the Mentalizing Signature (MS), B) the Self-Referential Signature (Self-RS), and C) the Other-Referential Signature (Other-RS). Voxels significant at an FDR-corrected threshold (*q* < .05, minimum cluster size of *k* = 10) are highlighted by black outlines and brighter colors.

1239 The left and middle plots in each box depict the receiver operating characteristic (ROC) plots from

1240 the i) cross-validated training and ii) testing datasets. The last column shows the accuracies of the

1241 mentalizing signatures in each testing dataset separately, with color representing the study, and

1242 shape illustrating the cohort type. The one-sided error bars below shapes illustrate the standard

1243 error. The star shows the weighted average accuracy across all seven datasets. To ensure that our

1244 cross-validated training and validation results were not influenced by the social-processing mask

1245 that we used, we repeated our analyses with the whole-brain (grey-matter masked) images, (see

1246 Supplementary Fig. 1). To test if a different feature selection method would yield similar results, we

1247 trained classifiers using recursive feature elimination (RFE), which we report in Supplementary

1248 Fig. 2.

1249

Figure 3. Model weights, cross-validated training, and validation results of the Self-vs-Other Classifier (SvO-RS). The weight map (first row) illustrates the positive and negative weights, as well as the voxels significant at an FDR-corrected threshold ($q < .05$, minimum cluster size of $k = 10$). The second row depicts the receiver operating characteristic (ROC) plots from the i) cross-validated training and ii) validation stages, and iii) the accuracies of the SvO-RS in each testing dataset. The cohort types are shape-coded, and the studies are color-coded in the validation accuracy plot. The one-sided error bars below shapes illustrate the standard error. The star shows the weighted average accuracy across all seven datasets.

1262
1263

Figure 4. Self/other discrimination based on classifier responses in healthy adults compared to individuals with schizophrenia. Signature fits are measured as the pattern expression of true minus false class, separately for each signature in healthy adults (*n*=48, in dark grey) and adults with schizophrenia (*n*=40, in light grey). Self/other discrimination was found to be better in Self-RS, Other-RS, and SvO-RS. Linear mixed effects are used to test the group differences, controlling for different cohorts. Self-RS: *β* = .21, *STE* = .07, *CI* = [.07, .35], *p* = .004. Other-RS: β = .19, *STE* = .07, *CI* = [.04, .33], *p* = .01, SvO-RS: β = .20, *STE* = .07, *CI* = [.07, .34], *p* = .004. Dots indicate values per person, and boxplots mark the median, lower, and upper quartiles. Outliers (> 3 SD from the mean) are marked with an "×". Whiskers extend from the minimum to the maximum data points, excluding outliers. Asterisks indicate statistical significance: *p < .05.

1274
1275

Figure 5. Association between self/other discrimination and age of adolescents (n = 105).
Older age correlated significantly with better differentiation of self- and other-related mentalizing,
controlling for cohort (Self-RS [β = .08, STE = .03, CI = [.012, .018], p = .01]; Other-RS [β = .06,
STE = .03, CI = [.001, .12], p = .047]; SvO-RS [β = .07, *STE* = .03, *CI* = .009 to .12, *p* = .02]. Black
lines display the linear regression fits, shaded areas the 95% confidence intervals. Dots show
values per participant.

1285

Figure 6. Extension of the Other-RS and the MS to an independent inference task (total n=109). A) Task design. Study 6a had a 1x2 design (attributional versus factual inferences regarding social scenes). Study 2 had a 2x2 design (attributional versus factual inferences x social versus nonsocial scenes). B) Binary classification performances of the MS and the Other-RS in social versus nonsocial attributions (Study 6b). C) Binary classification results from Study 6a for separating attributional versus factual inferences. D) Binary classification results from Study 6b, suggesting that the Other-RS and MS signatures are more sensitive to attributional (versus factual) inferences that require reflection about internal states—especially those concerning social agents.

1294

1295

Figure 7. Extension of three signatures to an independent social feedback task (n=49).
A) Participants received feedback for themselves, for their partners, or no feedback (Control
condition). B) Signature responses for each of the three conditions, demonstrating higher Self-RS
responses to feedback concerning oneself than the partner and compared no feedback, and higher
mentalizing-related brain responses (MS) for self- and partner-directed compared to no feedback.
Dots indicate values per person, and boxplots mark the median, lower, and upper quartiles. Outliers
(> 3 SD from the mean) are marked with an "×". Whiskers extend from the minimum to the maximum
data points, excluding outliers. Asterisks indicate significant Bonferroni-corrected pairwise
comparisons following an initial repeated measures ANOVA test within each signature: *p < .05;
***p < .001.

1306

Figure 8. Results of the ROI analysis. Cross-validated training and validation results for ten ROIs

derived   from   a   term-based   meta-analysis   ('mentalizing',   NeuroSynth[47]):   mPFC,

precuneus/posterior cingulate cortex (precuneus/PCC), temporoparietal junction (TPJ) right, TPJ

left, anterior middle temporal gyrus (aMTG) right, aMTG left, supplementary motor area (SMA) left,

cerebellum right, cerebellum left, and a bilateral cerebellum cluster. The ROIs that were

consistently predictive of both self- and other-related mentalizing were: mPFC, precuneus/PCC,

TPJ, aMTG. Four different classifiers were trained for each ROI: (i) to predict Self exclusively, (ii)

to predict Other exclusively, (iii) to predict overall mentalizing, and (iv) to discriminate the Self

versus Other. If an ROI classifier had significant accuracy in the cross-validated training dataset, it

was further tested across the testing datasets. The cross-validated training accuracies are

illustrated by the wider bars in the background. The validation accuracies are illustrated by the

thinner bars at the front. The error bars represent the standard error of the prediction. Classification

performances significantly above the chance level (50%) are marked with asterisks. The surface

images illustrate the weights of the classifiers that were trained to discriminate Self versus Other.

Orange-yellow colors show areas with positive weights towards Self, blueish colors show areas

associated with positive weights towards Other. If a specific ROI did not yield significant training

and independent classification accuracy for this task, then the area covering this ROI is displayed

in green color.

1326

1327
Figure 9. Leave-one-site-out (LOSiO) classifiers. We trained additional classifiers that combined
data across all datasets that used a trait-evaluation task (n=232) to rule out that the results were
driven by idiosyncratic features of the original training dataset. The classifiers were trained using
the same analytic strategy as the main mentalizing signatures, but using five folds, with each study
being held out in one-fold. A) Self-classifier, B) Other-classifier, C) Mentalizing-classifier, D) Self-
vs-Other classifier. Red stars show the out-of-sample accuracy across the whole sample (n=232),
while the colored circles represent the cross-validated accuracy for each study when classifiers
were trained on all other studies' data. The one-sided error bars in accuracy plots illustrate the

standard error. E) ROC plots for classifications in the inference task (Study 6) that involve

attributional versus factual statements in social and nonsocial settings.

1338