Sparse innervation and local heterogeneity in the vibrissal corticostriatal projection

Abbreviated title : Functional somatosensory corticostriatal connectivity

Authors and affiliations:

Kenza Amroune[1], Lorenzo Fontolan[2], Agnès Baude[1], David Robbe[2] and Ingrid Bureau[1]*

[1] Aix-Marseille Université, INSERM, INMED, Marseille, 13009, France

[2] Aix-Marseille Université, INSERM, INMED, Turing Centre for Living Systems, Marseille, 13009, France

corresponding author email address: ingrid.bureau@inserm.fr

Abstract

The density and overlap of cortical axons in the dorsolateral striatum (DLS) have suggested that striatal

neurons integrate widespread information from cortical regions that are functionally related. However,

in vivo, DLS neuronal responses to sensory stimuli have shown unexpectedly high selectivity, raising

questions about the actual degree of input convergence of functional corticostriatal projection on

individual striatal cells. Here, we investigated this question by focusing on the projections from different

whisker cortical columns, as they overlap in the striatum and are co-active during behavior. Using *ex vivo*

patch-clamp recordings in the DLS and glutamate uncaging for focal stimulations in the barrel cortex, we

were able to map the location of presynaptic neurons to individual striatal projection neurons (SPNs).

We found that each SPN was innervated by cells located in a small number of whisker cortical columns

scattered across the barrel field in the slice. Connectivity of single SPNs with cortical neurons was thus

highly discontinuous horizontally, despite the presence of more potential connections. Moreover,

connectivity patterns were specific to each cell, with neighboring SPNs sharing few common clusters of

presynaptic cells in the cortex. Despite this sparse and distinct innervation of individual SPNs, the

projection was topographically organized at the population level. Finally, we found similar innervation

patterns for D1 and D2-type SPNs, but observed distinct differences in synaptic strength at connections

with specific cortical layers, notably with the associative layer 2/3. Our results suggest that the high

convergence of somatosensory inputs to the striatum, enabled by diffuse and overlapping cortical

innervation, is accomplished through sparse yet complementary connectivity to individual SPNs.

Introduction

The dorsolateral striatum receives dense glutamatergic innervation from all sensory cortical areas. Its presumed function is to monitor the animal's ongoing sensory information, contributing to the modulation of sensory-guided behaviors and the learning of voluntary tasks (Pennartz et al., 2009; Znamenskiy and Zador, 2013; Sippy et al., 20158). The observations that cortical neurons are overabundant compared to striatal projection neurons (SPN) (Oorschot 1996), that axonal projections originating from functionally linked cortical regions overlap in the striatum (Flaherty and Graybiel, 1991; Alloway et al., 1999; Wright et al., 1999; Hooks et al., 2018) and that synaptic glutamatergic inputs must overcome strong feedforward inhibition (Mallet et al., 2005; Pidoux et al., 2011), have led to the idea that striatal neurons integrate dense and broad cortical inputs. In fact, studies examining the density of axonal boutons relative to the single cell dendritic spines have predicted that several thousand cortical neurons could potentially connect every SPN if synapses were formed promiscuously (Kincaid et al., 1998). However if synapses were formed in a selective manner, this would reduce the degree of convergence of cortical input onto individual SPNs and potentially increase the heterogeneity of inputs between SPNs (Kincaid et al., 1998), a scenario with important implications for how cortical information is integrated in the striatum. These alternative connectivity models schematized in Fig. 1 remained to be tested functionally.

Here, we investigated this question at the level of the projections to SPNs originating from the barrel field in the mouse primary somatosensory cortex (S1). Vibrissal information is transmitted to the cortical columns of S1 through parallel sensory channels, each corresponding to an individual whisker (reviewed in Adibi 2019). In S1, layer 4 contains optically dense, barrel-shaped structures whose organization mirrors the spatial arrangement of whiskers on the snout. This feature provides a framework for studying neuronal connectivity in relation to the whisker cortical columns *ex vivo*. However, this model has not yet been used to examine connectivity within the functional vibrissal corticostriatal projections, leaving the degree of input convergence in this pathway unclear. On the one hand, axons of neurons located in different whisker cortical columns overlap heavily in the dorsal striatum, suggesting that SPNs could receive dense and broad vibrissal cortical input (Wright et al., 1999; Alloway et al., 2000). On the other hand, cell activity evoked by whisker deflection showed notable heterogeneity in the DLS: neurons were either tuned to multiple whiskers, selective to a single whisker, or yet non-responsive (Carelli and West, 1991; Pidoux et al., 2011). These observations prompted us to investigate whether the vibrissal

innervation of SPNs violated Peters' rule stating that connectivity is proportional to the overlap of axons

and dendrites (Peters, 1979; Braitenberg and Schüz, 1998).

SPNs comprise two distinct populations, defined by their projection targets and expression of dopamine

receptors: D1 receptors in the direct pathway and D2 receptors in the indirect pathway (Smith et al.,

1998; Kravitz et al., 2010) (neurons are hereafter referred to as D1 and D2 SPNs). Studies have found

that inputs received by D1 SPNs were stronger than those received by D2 SPNs (Kress et al., 2013; Reig

and Silberberg, 2014; Parker et al., 2016; Filipović et al., 2019). These observations may indicate that

inputs converge to different degrees to D1 and to D2 SPNs.

To investigate the connectivity and input convergence on SPNs, we used a functional mapping technique

that combines patch-clamp recordings and laser scanning photostimulation (LSPS) with glutamate

uncaging on a novel slice preparation, which preserved the organization of the vibrissal corticostriatal

projections, from the superficial layers of the barrel cortex to the DLS. The sites in the barrel cortex

where LSPS evoked synaptic currents indicated the location of neuronal somas that were presynaptic to

the recorded SPN. We found overall discrete innervations, as presynaptic neurons were thinly-scattered

across the barrel field, or present in a single cortical column. SPNs located nearby in the same slice had

distinct patterns of innervation, suggesting that connections on each SPN were fewer than the local

potential connections. No difference was found between the innervation patterns of D1 and D2 SPNs,

but the strength of their connections with certain layers differed.

Results

A novel preparation for investigating functionally the organization of projections from the barrel

cortex to individual SPNs in the DLS

To investigate the spatial organization of sensory projections to SPNs, we have developed a parasagittal

somatosensory corticostriatal slice, which visibly contained ~ 8 whisker barrels (Methods; Fig. 2A and

Supplemental Fig. 1A). Biocytin-labeled projections extended from the top of the barrel cortex to the

DLS (Fig. 2A,B). To further validate the angles used in the preparation, we virtually re-sectioned a brain

from the Allen Mouse Connectivity Atlas in which pyramidal cells of the barrel cortex had been labeled

in vivo (i.e., prior to slicing; Supplemental Fig. 1B). Cortical axons extended into the striatum in a similar

manner in both the reconstituted and acute slices (compare Fig. 2A and Supplemental Fig. 1B),

supporting the idea that projections from the present cortical columns targeting the recorded striatal

region were largely preserved in the acute slice. Animals were Drd1a-tdTomato hemizygous mice in

which D1 receptor expressing neurons were labeled. D1$^+$ and D1$^-$ (D2$^+$) SPNs were recorded in the

whole-cell voltage–clamp configuration (n = 101 cells, N = 54 mice). Simultaneously, an ultraviolet laser

beam was directed at every site of a 29 x 16 pixel grid (2.1 × 1.1 mm, 75 µm spacing) to uncage

glutamate over the barrel cortex, excite cortical neurons, and thereby reveal cells presynaptic to the

recorded SPN (Fig. 2C,D). When glutamate was uncaged on an excitatory cortical neuron innervating the

recorded SPN, it elicited short-lived excitatory postsynaptic currents (EPSCs; < 50 ms duration) (Fig. 2D).

We and others have shown that only subthreshold synaptic events are elicited with this method,

implying that the evoked EPSCs are mono-synaptic and that feedforward inhibition is prevented

(Shepherd et al., 2003; Bureau et al., 2008). In fact, glutamate uncaging excited cortical neurons above

their firing threshold at a maximal distance of 75 µm from their cell body (Fig. 1E; Supplemental Fig. 2A),

on 2.8 ± 0.3 sites of the LSPS grid (mean ± sem; L5 pyramidal cells, n = 40; 75 µm spacing grid; see

Methods), consistent with a direct (i.e., non synaptic) and nearly somatic excitation (data per cell type in

the Methods and Supplemental Fig. 2). Thus, the sites where LSPS evoked EPSCs in SPNs contained or

were adjacent to the cell bodies of cortical neurons that were directly connected to the recorded cell.

These sites are hereafter referred to as "connected sites". EPSCs had small amplitudes, 40 ± 1 pA (n =

550), consistent with the presence of only one or a few presynaptic cells (≤ 5) at each connected site of

the map.

For each SPN, an input map was assembled from the excitatory response evoked at the uncaging sites of

the LSPS grid (Fig. 2D, top). From this input map, a binary map was derived which reported the presence

or lack of evoked EPSC, referred to as "connectivity map". We first analyzed the ensemble of

connectivity maps to characterize the global organization of the projections from the barrel field in the

slice and checked that their properties were consistent with those described by anatomical studies (Fig.

2F). When all SPN connectivity maps were superimposed and overlaid on the barrel cortex (see

Methods; Fig. 2F), the connected sites principally highlighted the layer (L) 5a. Indeed, ~ 65 % of all SPNs

received input from this layer (Fig. 2G). Responses were elicited less frequently when stimulations were

in L4, L5b, L2/3 and in L6. We found that 50 % of L5a cells fired action potentials when stimulations were

at the bottom of L4, indicating that a fraction of the relatively high connectivity rate seen at the bottom

of L4 was in fact with nearby L5a pyramidal cells (Fig. 2E; Supplemental Fig. 2A). In contrast, photo-

stimulations in L2/3 never elicited the firing of L5a cells (n = 11; Fig. 2E), indicating that all EPSCs evoked

by stimulations in these superficial layers arose from direct synaptic connections between L2/3

pyramidal cells and SPNs. EPSCs evoked with stimulations in L2/3 to L5b had similar amplitudes (Fig. 2H),

suggesting that L5a dominated these other layers thanks to a greater connectivity with SPNs principally.

However, L6-EPSCs were smaller compared to L5a and L5b-EPSCs (Kruskal-Wallis, $H(4) = 22.8$, $p =$

0.0001; Dunn post-hoc test and Šidák correction, $p = 0.0001$ and 0.034; Fig. 2H). These observations are

aligned with the anatomical studies which have identified L5a as the primary, although not exclusive,

source of cortical innervation in the DLS (Wise and Jones, 1977; Cowan and Wilson, 1994; Wall et al.,

2013).

In the superimposed connectivity maps, the ensemble of connected sites observed across recordings

formed a band in the barrel cortex that was larger than the region in the DLS containing the recorded

SPNs (2.5 mm vs. 1 mm, Fig. 2F). An opposite asymmetry was observed when analyzing the striatal

region connected to individual cortical columns: sites within a single cortical column (~ 150 μm in this

preparation) were connected to SPNs distributed across a larger striatal region, spanning about 600 μm

(Fig. 2I). These two asymmetric relationships (converging vs. diverging), summarized in Fig. 2J, implied

that functional projections from adjacent cortical columns overlapped in the DLS, which is consistent

with their known anatomy (Wright et al., 1999; Alloway et al., 2000; Hooks et al., 2018). Thus,

projections from the cortical columns in the slice overlapped in the striatum, which allowed us to

investigate the degree of convergence of whisker cortical inputs onto individual SPNs.

Low convergence of projections from the barrel cortex to individual SPNs in the DLS

To investigate the number and position of whisker cortical columns from which individual SPNs received

synaptic inputs, each connectivity map was collapsed along its vertical axis and the horizontal

distribution of connected sites was characterized using the following metrics (Fig. 3A): 1- the input field,

or the overall region within the barrel field in the slice from which an SPN receives inputs. Its width is

the distance between the SPN most distant connected sites.; 2- the cluster, a region in the collapsed

connectivity map constituted of adjacent connected sites; 3- the cortical column, or its proxy: 1 to 2

adjacent connected sites in the collapsed connectivity map (i.e. 75-150 μm width). The connected sites

were organized in 1 to 6 clusters, 1.9 ± 0.1 on average in the input fields (Fig. 3B). Typically, the width of

clusters was about one cortical column (171 ± 7 μm) and it rarely exceeded two adjacent columns (Fig.

3C). Finally, SPNs were innervated by cells distributed in close to 3 columns (2.7 ± 0.2), though some

SPNs with several clusters (54 %) were innervated by as many as eight columns (Fig. 3D). The distance

between clusters was 75 to 900 μm, 265 ± 28 μm on average (n = 55; Fig. 3C). Thus, SPNs could receive

inputs from clusters of cortical cells separated by substantial connectivity gaps, often spanning several

columns. Heterogeneous spacing and number of clusters contributed to the variability of the input fields

width, which ranged between 0.075 and 1.6 mm (average, 535 ± 42 µm; Fig. 3E). Strikingly, 45 % of the

input fields with several clusters produced no synaptic response upon stimulation (map collapsed along

the vertical axis; Fig. 3F, bottom). Moreover, the number of clusters was not increased proportionally in

very large input fields, > 1.4 mm, in which connectivity gaps amounted to more than 50 % of space in

the horizontal axis (Fig. 3F, bottom). Altogether, these findings suggest a connectivity pattern

characterized by a few small patches of presynaptic cortical cells scattered across the axis of barrels.

It could be that small input fields that contained a single cortical column (75-150 µm) were obtained

when recordings were performed on the edges of the striatal region with cortical innervation in the

slice. Contrary to this hypothesis, we found that SPNs with small input fields and those with broader

ones were intermingled along the horizontal axis in the striatum (open and solid symbols in Fig. 3G,

respectively). In fact, the width of the input fields remained relatively constant across SPN positions in

the slice (Fig. 3G, top; Spearman correlation coefficient, R = 0.05, p = 0.61). This suggests that the

position of our recordings did not impact the SPN input fields. Furthermore, SPNs receiving input from a

single column were often located near others receiving input from multiple ones (Fig. 3D), reinforcing

that the low functional connectivity with barrel columns in the slice was genuine in these cases.

Next, we investigated these connectivity patterns along their vertical axis. Individual SPNs had

connected sites distributed in 2.4 ± 0.1 cortical layers on average (2.9 ± 0.1 for SPNs with several

clusters). But these sites were not necessarily vertically aligned within a cortical column (examples in Fig.

2D and 3A). In fact, in 55 % of the cases, a column of the barrel field innervated an SPN through

projections from a single cortical layer. This suggests that neurons projecting to an SPN are dispersed in

the barrel cortex in such a way that the transmitted inputs may originate from distinct layers as well as

from different whisker cortical columns. Our findings are summarized in Fig. 3H, with schematics of the

average and principal connectivity patterns.

Individual SPNs have unique innervation

The above analysis has revealed great heterogeneity between SPN innervations. We then investigated

whether innervations were more similar when SPNs were close to each other. To address this question,

we compared the connectivity patterns of SPNs recorded in the same slice (Fig. 4A). First, we observed

that the majority of SPN pairs had their input fields overlapping in the barrel cortex (n = 70; Fig. 4B). The

ratio was the highest, 77 %, for SPNs that were neighbors in the striatum (< 100 µm, n = 33), but

remained above 50 % for cell distances of 400-500 µm. We next examined to what extent the connected

sites were shared or vertically aligned between the connectivity maps. In contrast to input fields,

alignment or overlap of connected sites was only observed when SPNs were less than 300 μm apart (Fig.

4B,C). Moreover, this concerned only a small fraction of the pair's connected sites: when SPNs were less

than 100 μm apart, only 18 ± 2 % of the connected sites in the two connectivity maps were vertically

aligned and 7.7 ± 1.6 % of them overlapped (Fig. 4B,C). Overall, these results indicate that DLS SPNs

could receive inputs from the same domain in the barrel cortex and yet have patterns of cortical

innervation without or little redundancy. They support a connectivity model in which synaptic

connections on each SPNs are sparser than the potential connections, and with significant heterogeneity

between SPNs in terms of their whisker-related cortical inputs.

Topographic organization of SPN functional vibrissal innervation

The sparse innervation patterns with little similarities of neighboring SPNs might be inconsistent with a

topographic organization. To examine the global organization of functional projections, connectivity

maps of SPNs were superimposed, and the connected sites were labeled based on the SPN positions in

the striatum (Fig. 5A). SPNs located laterally/medially in the dorsal striatum had an input field whose

connectivity center of mass (CM, see Methods) was in the lateral/medial part of the barrel cortex (Fig.

5A,B). Thus, the functional projections from the barrel cortex were topographically organized in the DLS,

consistent with anatomy (Flaherty and Graybiel, 1991; Alloway et al., 1999; Wright et al., 1999; Hooks et

al., 2018). Given the orientation of the slice (Supplemental Fig. 1A), a shift of the connectivity CM on the

horizontal axis of the barrel field corresponded principally to a shift between whisker arcs (e.g. E1 to E8

barrel). The connectivity CMs of two adjacent SPNs could be 1 mm apart in the barrel cortex (Fig. 5B),

indicating that the topographic organization was not as precise as for intracortical projections (Shepherd

and Svoboda, 2005; Erlandson et al., 2015). This finding, together with previous analyses, supports a

connectivity model that falls between the two illustrated in the middle and right panels of Fig. 1

(selective, broad and loosely topographic)

EPSC amplitudes were heterogeneous within each input map (Fig. 5C; ex Fig. 2D). To test the hypothesis

of a topographic organization influencing synaptic strength, we examined whether strong connections

were associated with presynaptic cells located close to the connectivity CM. If such an organization of

synaptic strengths existed, we would expect it to improve the correlation between the CM of synaptic

input maps and SPN positions. However, we found nearly identical correlations whether taking the

connectivity CM or the synaptic input CM, which is based on the connectivity pattern weighted by EPSC

amplitudes (see Methods; R = 0.60 vs. 0.62; F(1) = 115.08, *p* = 0.98, ANCOVA; Fig. 5D). While residuals

from the correlation curve tended to be smaller using synaptic input CMs, the difference was not

statistically significant (- 9 ± 7 μm, p = 0.084, Wilcoxon, SPNs with >1 cluster). This was because synaptic

input CMs were only slightly shifted from the connectivity CMs, by 45 ± 5 μm for SPNs with two or more

clusters (Fig. 5D). These findings indicate that EPSC amplitudes showed no consistent pattern relative to

the connectivity CM of the input field. This confirms the connectivity patterns as principal determinants

of the topographic organization of somatosensory corticostriatal projections.

Finally, given that previous studies have revealed that responses of some striatal neurons exhibited

strong selectivity for specific whiskers *in vivo* (Carelli and West, 1991; Pidoux et al., 2011), we

investigated whether, in cases where SPNs were innervated by multiple clusters, one cluster might

dominate the others. Comparing the sum of EPSCs corresponding to clusters, we found substantial

variability, with the strongest cluster providing input that was 1.12 to 55 times greater than other

clusters within the input field (× 6.8 ± 1.4 on average; Fig. 5E). Thus, connectivity patterns with multiple

clusters may still permit some SPNs to have a preference for a particular whisker cortical column, due to

the heterogeneity between clusters.

D1 and D2 SPNs receive similar input from the barrel cortex

Finally, we investigated whether innervation patterns differed between the two populations of SPNs,

expressing D1 or D2 receptors. Based on previous studies, unlabeled neurons in Drd1a-tdTomato

hemizygous mice were principally SPNs expressing the D2 receptor (Bertran-Gonzalez et al., 2008; Ade

et al., 2011; Enoksson et al., 2012; Thibault et al., 2013; Cao et al., 2018). As previously reported (Wall et

al., 2013), projections to D1 and D2 SPNs were spatially intermingled along the axis of barrels (Fig. 6A,B ;

n = 47 and 54, cells respectively). We found that D1 and D2 SPNs had similar input fields: first, their

widths were not different (Fig. 6C). In addition, they received input from a similar number of cortical

columns, which were organized in clusters of similar widths (Fig. 6D,E). Paired analysis of D1 and D2

SPNs recorded in the same slice also showed that their input fields did not significantly differ in width,

number of cortical columns or cluster width (p = 0.44 – 0.50 – 0.26, Wilcoxon; Fig. 6C-E). None of these

parameters exhibited a clear relationship between subtypes (R = 0.13 – 0.30 – -0.003; p = 0.43 – 0.058 –

0.99, Pearson; Fig. 6C-E). Finally, both types of SPNs received input from each cortical layer in similar

proportions (Fig. 6F). Thus, the heterogeneity in patterns of innervation is not due to a difference

between SPN subtypes. Overall, there was no difference in the total inputs received by D1 and D2 SPNs

(EPSC sum or $EPSC^T$; Fig. 6G). In addition, the principal cluster was similarly dominant over the other

clusters (Fig. 6H). However, differences were observed in a layer-specific manner (Fig. 6I). Whereas L5-

EPSC^T was similar in D1 and D2 cells, differences were found in L2/3 and L6: L2/3-EPSC^T was larger in D2

SPNs (132 ± 36 vs. 53 ± 22 pA, p = 0.005, Mann-Whitney) but L6-EPSC^T was larger in D1 SPNs (37 ± 10 vs.

16 ± 2 pA, p = 0.038). Further analysis suggested a difference in the strength of synapses as principal

mechanism. Indeed, L2/3-EPSCs evoked at single sites were larger in D2 SPNs (48 ± 11 pA vs. 27 ± 5 pA;

p = 0.043) whereas L6-EPSCs were larger in D1 SPNs (24 ± 2 pA vs. 16 ± 2 pA ; p = 0.038).

Discussion

We investigated the organization of functional projections originating in the barrel cortex and targeting

projection neurons in the dorsal striatum in a novel preparation. We found overall little input

convergence as cortical neurons innervating an SPN were either located in a single cortical column or

thinly scattered across the barrel field. Each SPN had a distinct pattern of innervation from the barrel

cortex, albeit with occasional overlap for SPNs closely positioned in the DLS. D1 and D2 presented

similar patterns of innervations.

Sparse sensory innervation of single SPNs

As previously revealed by tracings of corticostriatal projections, the primary source of innervation of

SPNs in this study is the upper part of L5 in the barrel cortex, L5a, although, SPNs are occasionally

innervated by cells located in deeper and superficial layers (Wise and Jones, 1977; Reiner et al., 2003;

Wall et al., 2013; Guo et al., 2017; Yamashita et al., 2018; Bertero et al., 2022). Given the abundance of

cortical axons in the striatum, it has been assumed that a large number of cortical cells innervate each

SPN, potentially up to few thousands if synaptic connections were made promiscuously (Kincaid et al.,

1998). On the other hand, it has also been reported that neurons in the DLS fire in response to the

stimulation of a single body-part in monkeys and rodents *in vivo*, even of a single whisker in some cases

(Carelli and West, 1991; Jaeger et al., 1995; Cho and West, 1997). This finding suggested that each

striatal neuron was in fact innervated by one small subregion of S1, although feedforward inhibition

could have masked a broader selectivity (Pidoux et al., 2011). Here, the LSPS connectivity maps indicate

that convergence from the barrel cortex to individual SPNs is indeed low as 60 % of SPNs responded to

the stimulation of only one or two whisker cortical columns present in the preparation. The LSPS

combined with glutamate uncaging maps projections from neuronal cell bodies within the slice.

Therefore, our study characterized connectivity patterns rather than the full extent of connectivity with

the barrel cortex, as projections from columns outside the slice were not activated. A number of critical

findings argue that innervation on individual SPNs is genuinely sparse: i) The cortical columns present in

the slice corresponded to different whisker arcs, an axis associated with the greatest overlap of cortical

axons in the striatum (Alloway et al., 1999), and therefore where input convergence could have been

high. This organization was visible in the global SPN LSPS connectivity map (Fig. 2J). ii) The peak of

connectivity was with L5a as previously documented, not with a layer closer to striatum, as this would

be the case if the native organization of the projection was altered in the preparation. iii) The connected

sites were scattered along broad horizontal and vertical axes, implying that projections in between were

preserved. Critically, the fact that adjacent SPNs had different connectivity patterns in the same slice

further supported the presence of a pool of potential connections larger than the number of actual

connections made on each SPN. iv) Connectivity clusters were sparser in larger input fields, suggesting

that the number of presynaptic cells to an SPN was regulated and reached a plateau. v) EPSCs were

small, consistent with one or a few presynaptic cells at each connected site. Altogether, our results

support the model in which, within a loosely topographical organization of projections, each SPN

integrates limited and heterogeneous rather than exhaustive inputs transmitted by the barrel cortex

(Fig. 7). The speckled connectivity pattern of individual SPNs, arising from the abundant and diffuse

cortical innervation in the DLS, suggests that sensory corticostriatal synapses are established through a

selective and/or competitive process.

Specific cortical input to individual SPNs

A peculiar feature of the corticostriatal innervation is the imbalance between the number of cortical

cells projecting to the striatum and the number of striatal cells, thought to be at a ratio of 10 to 1

(Oorschot, 1996). Based on this ratio and the ultrastructure of the corticostriatal connection, the density

of axonal boutons and spines, C. Wilson's group concluded that the probability of an SPN being

contacted by one given axon entering the domain of its dendritic arborization was low, 0.04 - 1.4 %,

depending on the model. Hence, the chance that two SPNs were innervated by the same cortical axon

was even lower according to this model. Consistent with this, here, we found that overlaps between the

connectivity maps of SPNs were rare. This indicates that SPNs primarily integrate distinct inputs

originating from the barrel cortex. When overlaps were observed, they occurred only for SPNs located

within 250 µm of each other. However, such overlaps involved only a small fraction of their inputs, as

the connectivity patterns of nearby SPNs were largely interdigitated. In addition, the input fields

continued to intersect significantly beyond 250 µm and up to 500 µm of separation between SPNs. This

means that relatively distant SPNs can share a general region of connectivity in the barrel cortex without

any of their presynaptic partners residing within the same cortical column. Interestingly, the observation

that two SPNs must be less than 250 μm apart to share some of their inputs was not predicted by the

known ultrastructure of corticostriatal connections. In fact, it predicted the opposite: that two SPNs

would more likely to be innervated by the same cortical cell if their dendrites extended into different

striatal subregions, as this would increase the likelihood of each SPN contacting boutons of this cortical

cell, as these are sparsely distributed in the striatum (Kincaid et al., 1998). One possible explanation for

this discrepancy could be that, in our experiments, pairs of SPNs with shared connectivity were

innervated by different neurons that were co-localized in the barrel cortex (< 75 μm apart). Such events

could also be at the origin of the striatal clusters described *in vivo,* in which neurons had similar, yet non-

identical, sensory selectivity (Carelli and West, 1991; Jaeger et al., 1995).

Similarities and specificities in the patterns of projection to SPNs of the direct and indirect pathways

Electrophysiological studies have shown that D1 SPNs exhibit larger responses than D2 SPNs to whisker

deflections (Reig and Silberberg, 2014; Filipović et al., 2019). Also, the optostimulation of cortical

efferents in brain slices showed a bias towards D1 SPNs (Kress et al., 2013; Parker et al., 2016). In our

study, we had contrasting results. There was no difference in the innervation of D1 and D2 SPNs, neither

in the rate at which cells were connected by layers, nor in the organization of the projection. The

principal layer innervating SPNs, L5a, delivered inputs of similar strengths to D1 and D2 SPNs. However,

layers with lower incidence in the pattern of innervation showed bias in the amplitude of responses:

stimulations in L2/3 induced stronger responses in D2 cells whereas stimulations in L6 activated D1 cells

more strongly. We have shown that stimulation in L2/3 activates the L2/3 projection only. Similarly, the

L6-evoked EPSCs are consistent with the axon collaterals that L6 corticofugal projection neurons have in

the striatum (Guo et al., 2017; Bertero et al., 2022). The larger L2/3 inputs received by the D2 SPNs is

intriguing because of the role of this layer in higher-order integration processes, such as those activated

during operant sensory discrimination tasks (Kwon et al., 2016; Han and Helmchen, 2024; Oryshchuk et

al., 2024). Moreover, the activity of L2/3 cells during mouse displacement is distinctive, exhibiting a

sustained response to ongoing whisker-wall contacts, as opposed to the transient response observed in

L5 (Ayaz et al., 2019). In L6, the population of corticofugal projection neurons is diverse, with some

responding to whisking and sensory stimuli, while others are active during transitions into quiet periods

of behavior (Dash et al., 2022; Vélez-Fort et al., 2014). Further investigation is needed to specifically

examine projections from the upper and lower cortical layers and how they influence the dynamics of

sensory integration in the direct and indirect pathways of the DLS. However, these data illustrate the

diversity of paths of sensory integration involving D1 and D2 SPNs.

In conclusion, we found a low and sparse connectivity within the cortical vibrissal projections to
individual SPNs. While projections respected a loose topography, the innervation patterns of individual
SPNs displayed substantial connectivity gaps and heterogeneity. Since the inputs of each SPN represent
only a limited subset of whisker columns, sensory integration must occur at the population level, with
each SPN representation complementing the representations of its neighbors (Fig. 7). These
observations raise the hypothesis of a selective or competitive process underlying the formation of
corticostriatal synapses.

Materials and Methods

Materials and Methods

Animals and ethics
Hemizygous male and female B6.Cg-Tg(Drd1a-tdTomato)6Calak/J mice (JAX stock #016204; Ade et al.,
2011) were used on postnatal day 22-43, in accordance with institutional guidelines and the French
Ministry of Research (APAFIS#27242).

Brain slices preparation and electrophysiology.
Mice were deeply anesthetized with isoflurane (4 %) prior to cervical dislocation and decapitation. We
prepared corticostriatal slices (350 μm thick) from the brain left hemisphere, based on stereotaxic
coordinates placing the striatal cells 1-2 mm anteriorly to the projection neurons in the barrel cortex
(Aronoff et al., 2010; de la Torre-Martinez et al., 2023). Parasagittal slices were cut with a 60° angle from
the midline and a 10° angle in the dorso-ventral axis (Supplemental Fig. 1A) in a chilled cutting solution
containing (in mM): 110 choline chloride, 25 NaHCO3, 25 D-glucose, 11.6 sodium ascorbate, 7 MgCl2,
3.1 sodium pyruvate, 2.5 KCl, 1.25 NaH2PO4, and 0.5 CaCl2 (Sigma Aldrich). Slices were then transferred
to artificial cerebrospinal fluid (ACSF) containing (in mM): 127 NaCl, 25 NaHCO3, 25 D-glucose, 2.5 KCl, 1
MgCl2, 2 CaCl2, and 1.25 NaH2PO4, aerated with 95% O2/5% CO2. Slices were first incubated at 34 °C
for 30 min and then maintained at room temperature for 20 min prior to use. Slices used for LSPS (1-2
per animal) contained barrels in the L4 of cortex, the globus pallidus, its external segment (GPe), the
internal capsule, the ventral posteromedial nucleus of thalamus and the anterior hippocampus. ~ 8
barrels (5-13) were visible in the slice. 2 rows of barrels may have been superimposed. At the end of
each experiment, a picture of the slice was saved in order to superimpose it digitally to other slices,
according to visual landmarks (Adobe Photoshop; Adobe Inc.). SPNs were visualized under infrared and
fluorescent lights in a BX61WI microscope (Olympus) and patched with borosilicate electrodes (3–6 MΩ)
and recorded in the voltage-clamp whole-cell configuration using a Multiclamp 700A amplifier (Axon
Instrument, Molecular Devices). The holding membrane potential was − 80 mV. The intracellular

solution contained (in mM) 128 Cs-methylsulfate, 4 MgCl2, 10 HEPES, 1 EGTA, 4 Na2ATP, 0.4 Na2GTP, 10

Na-phosphocreatine, 3 ascorbic acid; pH 7.25; 290-300 mOsm. Cells in L5 and L2/3 of the barrel cortex

were recorded in the current-clamp mode, with an intracellular solution in which Cs-methylsulfate was

replaced by K-methylsulfate. All experiments were performed at room temperature (21°C).

LSPS with glutamate uncaging

LSPS was performed as described previously (Bureau et al., 2006). Recirculating (2 mL/min) ACSF

solution contained (in mM): 0.2 MNI-caged glutamate (Tocris), 0.005 CPP [()-3-(2-carboxipiperazin-4-yl)

propyl-1-phosphonic acid], 4 $CaCl_2$, and 4 $MgCl_2$. Focal photolysis of caged glutamate was accomplished

with a 2 ms 20 mW pulse of a pulsed UV (355 nm) laser (DPSS Lasers Inc.) through a 0.16 NA 4 ×

objective (Olympus). 25 mW laser pulses were used for stimulating cortical neurons in mice older than

P30 to maintain their excitation at a similar level than in younger mice (Supplemental Fig. 2B). The

stimulus pattern for mapping the corticostriatal projections was 464 positions spaced by 75 μm on a 29

× 16 grid (2.1 × 1.1 mm) over barrel cortex. The corticostriatal slice and the LSPS grid were oriented in

such a way that layer 5a was laid out horizontally. UV stimuli were applied every 700 ms and their

successive positions on the LSPS grid were such as to maximize the time between stimulations of

neighboring sites. Electrophysiological traces consisted of 100 ms baseline, a 450 ms window followed

by a -5 mV 100 ms test pulse. A minimum of 2 and up to 4 stimulations were performed at each site at

several minutes intervals. Custom software for instrument control and acquisition (Suter et al. 2010)

was written in Matlab (Mathworks). Excitation profiles of pyramidal neurons were generated under

similar conditions except that cells were recorded in current-clamp mode and glutamate was uncaged

on a smaller 8 × 8 grid covering their soma and dendrites (50 μm spacing; 350 × 350 μm). In a subset of

L5a cell recordings, a 8 × 24 grid was used to stimulate up in L1 (50 μm spacing).

Analysis of LSPS data

Synaptic input maps of neurons were constructed by taking the peak amplitude of EPSCs detected in a

50 ms time window starting at stimulation onset for each position in the LSPS grid. Measures were

averaged across repetitions of stimulations (2-4). The threshold for EPSC detection was 3 standard

deviations from baseline, or 9.2 ± 0.4 pA. To disambiguate evoked responses from spontaneous activity,

synaptic responses occurring less than 2 times across repetitions of maps were set to zero. Averaged

maps were superimposed taking L5a as reference in the vertical axis and the junction of the GPe, dorsal

striatum and internal capsule as reference in the horizontal axis (Ref_{hor}). In order to detect connectivity

clusters and center of mass in the LSPS map, we used the binary version of the map ('connectivity map')

reporting the location of connected and non-connected sites (i.e., yielding EPSCs or none in the

recorded SPN). To detect clusters, the binary map was collapsed along its vertical axis. A cluster

comprised 1 or more consecutive connected sites that was framed by 1 or more non-connected sites.

Thus, a cluster here may include connected sites that were not adjacent on the vertical axis in the

original map, and may combine synaptic inputs from different layers. The number of cortical columns

was estimated based on the number of consecutive connected sites (ConsSites) in the collapsed map

and the mean width of a barrel in the slice, 150 µm, which is equal to 2 connected sites in the map.

Hence, the number of cortical columns = Σ (ceiling(ConsSites/2)). The connectivity center of mass was

computed based on the uncollapsed map as follows : Σ (Σ_{vert} connected sites × lateral distance from

Ref_{hor}) /Σ (Σ_{vert} connected sites). The synaptic input center of mass was computed based on the original

map, after thresholding, as follows: Σ ($mean_{vert}$ EPSC × lateral distance from Ref_{hor}) /Σ ($mean_{vert}$ EPSC).

Traces from current clamp recordings were analyzed to count the number of action potentials (APs)

elicited upon glutamate uncaging. In the 8 × 8 grid, 50 µm spacing, total number of spikes was 2.2 ± 0.2

for L2/3 pyramidal cells (n = 26), 5.4 ± 0.6 for regular spiking (RS) pyramidal cells in L5a (n = 30) and 9.9

± 1.6 for non-RS pyramidal cells in L5b (n = 10). The number of spikes elicited by uncaging at single sites

was 1 or 2 and on average: 1.04 ± 0.04 for L2/3 cells, 1.03 ± 0.02 for L5 RS cells and 1.04 ± 0.02 for L5

non-RS cells. As spacing in the stimulation grid was 50 µm in these particular experiments, we used the

following equation to compute the number of sites with AP (sAP) in a stimulation grid with a spacing of

75 µm (i.e. as in connectivity maps): $sAP_{75} = sAP_{50} \times (75/50)^2$.

Labeling of cortical cells

In a somatosensory corticostriatal slice prepared as described above, biocytin (2 %) was injected by

iontophoresis in the L5a through a patch electrode (200 ms, 2.5 Hz) for 2 x 10 minutes (Chang, LoTurco

and Nisenbaum, 2000). The slice was incubated for 4 hours at room temperature in ACSF and

transferred to 4 % paraformaldehyde at 4°C overnight. It was rinsed in PBS containing 0.3 % Triton X-100

(PBS-T), incubated overnight at room temperature in streptavidin-AlexaFluor488 (1:1000 in PBS-T) and

mounted using Vectashield mount medium. Image acquisition was performed using a LSM 800 confocal

Zeiss microscope.

Statistical analysis

All data are expressed as mean ± SEM in the text. Median and 25-75[th] percentiles are shown in Figures.

N and n are the numbers of animals and neurons, respectively. Unless stated otherwise, paired or

unpaired non parametric tests, and Kruskal-Wallis test for repeated measures followed by Dunn post-

hoc test and Šidák correction for multiple comparisons were used. *p* < 0.05 is considered as statistically
significant.

Acknowledgments:

This work was supported by funding from the Institut National de la Santé et de la Recherche Médicale
and a grant from the Agence Nationale de la Recherche (Corticostriatal, ANR-20-CE16-0002). K.
Amroune is supported by fellowships from the Ministère de l'Enseignement Supérieur et de la
Recherche, from the governmental "France 2030" program via A*Midex (Initiative d'Excellence d'Aix-
Marseille Université, AMX-19-IET-004) and by ANR funding (ANR-17-EURE-0029). L. Fontolan is
supported by funding from Excellence Initiative of Aix Marseille Université - A*MIDEX (Turing Centre for
Living Systems). We thank the staff of the animal and genotyping facilities, the histology and imaging
platforms of INMED, the members of the CBGB group for their help and support and Elodie Fino, Rosa
Cossart, Ede Rancz, Roustem Khazipov, Thomas Morvan, Jean-Luc Gaiarsa, Corinne Beurrier and Lydia
Kerkerian Le Goff for their critical reading of the manuscript.

References

Ade KK, Wan Y, Chen M, Gloss B, Calakos N (2011) An Improved BAC Transgenic Fluorescent Reporter
Line for Sensitive and Specific Identification of Striatonigral Medium Spiny Neurons. Front Syst
Neurosci 5:32.

Adibi M (2019) Whisker-Mediated Touch System in Rodents: From Neuron to Behavior. Front. Syst.
Neurosci 13:40.

Alloway KD, Crist J, Mutic JJ, Roy SA (1999) Corticostriatal Projections from Rat Barrel Cortex Have an
Anisotropic Organization that Correlates with Vibrissal Whisking Behavior. J Neurosci 19:10908–
10922.

Alloway KD, Mutic JJ, Hoffer ZS, Hoover JE (2000) Overlapping corticostriatal projections from the rodent
vibrissal representations in primary and secondary somatosensory cortex. J Comp Neurol
428:51–67.

Aronoff R, Matyas F, Mateo C, Ciron C, Schneider B, Petersen CCH (2010) Long-range connectivity of
mouse primary somatosensory barrel cortex. Eur J Neurosci 31:2221–2233.

Ayaz A, Stäuble A, Hamada M, Wulf M-A, Saleem AB, Helmchen F (2019) Layer-specific integration of
locomotion and sensory information in mouse barrel cortex. Nat Commun 10:2585.

Benison AM, Ard TD, Crosby AM, Barth DS (2006) Temporal patterns of field potentials in vibrissa/barrel
cortex reveal stimulus orientation and shape. J Neurophysiol 95:2242–2251.

Bertero A, Verrillo L, Apicella AJ (2022) A Novel Layer 4 Corticofugal Cell Type/Projection Involved in
Thalamo-Cortico-Striatal Sensory Processing. J Neurosci 42:1383–1405.

Bertran-Gonzalez J, Bosch C, Maroteaux M, Matamales M, Hervé D, Valjent E, Girault J-A (2008)
Opposing patterns of signaling activation in dopamine D1 and D2 receptor-expressing striatal
neurons in response to cocaine and haloperidol. J Neurosci 28:5671–5685.

Braitenberg V, Schüz A (1998) Cortex: Statistics and Geometry of Neuronal Connectivity. Springer

Bureau I, Shepherd GMG, Svoboda K (2008) Circuit and plasticity defects in the developing
somatosensory cortex of FMR1 knock-out mice. J Neurosci 28:5178–5188.

Bureau I, von Saint Paul F, Svoboda K (2006) Interdigitated paralemniscal and lemniscal pathways in the
mouse barrel cortex. Plos Biol 4:e382

Cao J, Dorris DM, Meitzen J (2018) Electrophysiological properties of medium spiny neurons in the
nucleus accumbens core of prepubertal male and female Drd1a-tdTomato line 6 BAC transgenic
mice. J Neurophysiol 120:1712–1727.

Carelli RM, West MO (1991) Representation of the body by single neurons in the dorsolateral striatum of
the awake, unrestrained rat. J Comp Neurol 309:231–249.

Chang SL, LoTurco JJ and Nisenbaum LK (2000) In vitro biocytin injection into perinatal mouse brain: a
method for tract tracing in developing tissue. J Neurosci Methods 97:1-6

Cho J, West MO (1997) Distributions of single neurons related to body parts in the lateral striatum of the
rat. Brain Res 756:241–246.

Cowan RL, Wilson CJ (1994) Spontaneous firing patterns and axonal projections of single corticostriatal
neurons in the rat medial agranular cortex. J Neurophysiol 71:17–32.

Dash S, Autio DM, Crandall SR (2022) State-Dependent Modulation of Activity in Distinct Layer 6
Corticothalamic Neurons in Barrel Cortex of Awake Mice. J Neurosci 42:6551-6565

de la Torre-Martinez R, Ketzef M, Silberberg G (2023) Ongoing movement controls sensory integration in
the dorsolateral striatum. Nat Commun 14:1004.

Enoksson T, Bertran-Gonzalez J, Christie MJ (2012) Nucleus accumbens D2- and D1-receptor expressing
medium spiny neurons are selectively activated by morphine withdrawal and acute morphine,
respectively. Neuropharmacology 62:2463–2471.

Erlandson MA, Manzoni OJ, Bureau I (2015) The Functional Organization of Neocortical Networks
Investigated in Slices with Local Field Recordings and Laser Scanning Photostimulation. Plos One
10:e0132008.

Filipović M, Ketzef M, Reig R, Aertsen A, Silberberg G, Kumar A (2019) Direct pathway neurons in mouse
dorsolateral striatum in vivo receive stronger synaptic input than indirect pathway neurons. J
Neurophysiol 122:2294–2303.

Flaherty AW, Graybiel AM (1991) Corticostriatal transformations in the primate somatosensory system.
Projections from physiologically mapped body-part representations. J Neurophysiol 66:1249–
1263.

Flaherty AW, Graybiel AM (1994) Input-output organization of the sensorimotor striatum in the squirrel
monkey. J Neurosci 14:599–610.

Guo C, Peng J, Zhang Y, Li A, Li Y, Yuan J, Xu X, Ren M, Gong H, Chen S (2017) Single-axon level
morphological analysis of corticofugal projection neurons in mouse barrel field. Sci Rep 7:2846.

Hooks BM, Papale AE, Paletzki RF, Feroze MW, Eastwood BS, Couey JJ, Winnubst J, Chandrashekar J,
Gerfen CR (2018) Topographic precision in sensory and motor corticostriatal projections varies
across cell type and cortical area. Nat Commun 9:3549.

Jaeger D, Gilman S, Aldridge JW (1995) Neuronal activity in the striatum and pallidum of primates
related to the execution of externally cued reaching movements. Brain Res 694:111–127.

Kincaid AE, Zheng T, Wilson CJ (1998) Connectivity and Convergence of Single Corticostriatal Axons. J
Neurosci 18:4722.

Kravitz AV, Freeze BS, Parker PRL, Kay K, Thwin MT, Deisseroth K, Kreitzer AC (2010) Regulation of
parkinsonian motor behaviours by optogenetic control of basal ganglia circuitry. Nature
466:622–626.

Kress GJ, Yamawaki N, Wokosin DL, Wickersham IR, Shepherd GMG, Surmeier DJ (2013) Convergent
cortical innervation of striatal projection neurons. Nat Neurosci 16:665–667.

Kwon SE, Yang H, Minamisawa G, O'Connor DH (2016) Sensory and decision-related activity propagate in
a cortical feedback loop during touch perception. Nat Neurosci 19:1243–1249.

Mallet N, Le Moine C, Charpier S, Gonon F (2005) Feedforward inhibition of projection neurons by fast-
spiking GABA interneurons in the rat striatum in vivo. J Neurosci 25:3857–3869.

Oorschot DE (1996) Total number of neurons in the neostriatal, pallidal, subthalamic, and substantia
nigral nuclei of the rat basal ganglia: a stereological study using the cavalieri and optical disector
methods. J Comp Neurol 366:580–599.

Oryshchuk A, Sourmpis C, Weverbergh J, Asri R, Esmaeili V, Modirshanechi A, Gerstner W, Petersen CCH,
Crochet S (2024) Distributed and specific encoding of sensory, motor, and decision information
in the mouse neocortex during goal-directed behavior. Cell Rep 43:113618.

Parker PRL, Lalive AL, Kreitzer AC (2016) Pathway-Specific Remodeling of Thalamostriatal Synapses in
Parkinsonian Mice. Neuron 89:734–740.

Pennartz CMA, Berke JD, Graybiel AM, Ito R, Lansink CS, van der Meer M, Redish AD, Smith KS, Voorn P
(2009) Corticostriatal Interactions during Learning, Memory Processing, and Decision Making. J
Neurosci 29:12831–12838.

Peters A (1979) Thalamic input to the cerebral cortex. Trends Neurosci 2:183–185.

Pidoux M, Mahon S, Deniau J-M, Charpier S (2011) Integration and propagation of somatosensory
responses in the corticostriatal pathway: an intracellular study *in vivo*: Sensory responses in the
corticostriatal pathway. J Physiol 589:263–281.

Ramaswamy S, Markram H (2015) Anatomy and physiology of the thick-tufted layer 5 pyramidal neuron.
Front Cell Neurosci 9:233.

Reig R, Silberberg G (2014) Multisensory integration in the mouse striatum. Neuron 83:1200–1212.

Reiner A, Jiao Y, Del Mar N, Laverghetta AV, Lei WL (2003) Differential morphology of pyramidal tract-
type and intratelencephalically projecting-type corticostriatal neurons and their intrastriatal
terminals in rats. J Comp Neurol 457:420–440.

Shepherd GMG, Pologruto TA, Svoboda K (2003) Circuit analysis of experience-dependent plasticity in
the developing rat barrel cortex. Neuron 38:277–289.

Shepherd GMG, Svoboda K (2005) Laminar and columnar organization of ascending excitatory
projections to layer 2/3 pyramidal neurons in rat barrel cortex. J Neurosci 25:5670–5679.

Shepherd GMG, Stepanyants A, Bureau I, Chklovskii D, Svoboda K (2005) Geometric and functional
organization of cortical circuits. Nat Neurosci 8:782–790

Sippy T, Lapray D, Crochet S, Petersen CCH (2015) Cell-Type-Specific Sensorimotor Processing in Striatal
Projection Neurons during Goal-Directed Behavior. Neuron 88:298–305.

Smith Y, Bevan MD, Shink E, Bolam JP (1998) Microcircuitry of the direct and indirect pathways of the
basal ganglia. Neuroscience 86:353–387.

Thibault D, Loustalot F, Fortin GM, Bourque M-J, Trudeau L-É (2013) Evaluation of D1 and D2 Dopamine
Receptor Segregation in the Developing Striatum Using BAC Transgenic Mice. Plos One
8:e67219.

Vélez-Fort M, Rousseau CV, Niedworok CJ, Wickersham IR, Rancz EA, Brown APY, Strom M, Margrie TW
(2014) The stimulus selectivity and connectivity of layer six principal cells reveals cortical
microcircuits underlying visual processing. Neuron 83:1431-43

Wall NR, De La Parra M, Callaway EM, Kreitzer AC (2013) Differential innervation of direct- and indirect-
pathway striatal projection neurons. Neuron 79:347–360.

Wise SP, Jones EG (1977) Cells of origin and terminal distribution of descending projections of the rat
somatic sensory cortex. J Comp Neurol 175:129–157.

Wright AK, Norrie L, Ingham CA, Hutton EAM, Arbuthnott GW (1999) Double anterograde tracing of
outputs from adjacent "barrel columns" of rat somatosensory cortex. Neostriatal projection
patterns and terminal ultrastructure. Neuroscience 88:119–133.

Yamashita T, Vavladeli A, Pala A, Galan K, Crochet S, Petersen SSA, Petersen CCH (2018) Diverse Long-
Range Axonal Projections of Excitatory Layer 2/3 Neurons in Mouse Barrel Cortex. Front
Neuroanat 12:33.

Znamenskiy P, Zador AM (2013) Corticostriatal neurons in auditory cortex drive decisions during
auditory discrimination. Nature 497:482–485.

Figure 1. Potential connectivity patterns within the corticostriatal projection.
A rich and overlapping cortical innervation of the striatum (lines) permits different connectivity patterns: Left, synapses are formed promiscuously. In this case convergence is high as each SPN receives broad cortical input, from many cortical origins (colored bars on top). Middle and right, only a fraction of the potential connections are actually formed and, in these cases, convergence is lower. The selected presynaptic neurons are either scattered (middle) or topographically (right) positioned in the cortex. Increasing the selectivity of the corticostriatal connections while keeping their broad origin (middle) could generate maximal heterogeneity in the patterns of input for SPNs sharing the same striatal volume, due to a large number of combinations of cortical origins. In the case of a topographic innervation (right), neighboring SPNs form synapses with presynaptic neurons located in the same cortical region, reducing the local heterogeneity.

Figure 2. A novel slice preparation for investigating the spatial organization of the somatosensory projections to the DLS neurons.

A. A somatosensory corticostriatal slice in which axons from L5 neurons were labeled with biocytin iontophoresis. Black star, the electrode position in L5a. Right, focus on a region in the striatum.

B. Right, Example dendritic arbors of L5 cortical cells labeled with biocytin in the slice.

C. Montage of a corticostriatal slice (left) and layout of the experiment (right). A SPN was recorded in

the dorsolateral striatum while cortical neurons were photostimulated with LSPS. The grid of LSPS (blue)

was positioned on the barrel cortex. GPe, globus pallidus, external segment; IC, internal capsule; hipp.,

hippocampus.

D. Top, examples of synaptic input maps for individual SPNs showing one or four clusters of input (SPN 1

and 2, respectively). In the color maps, each pixel of color indicates the peak amplitude of EPSCs

detected within a 50 ms window after the stimulus onset. The different cortical layers are represented

by solid white vertical lines on the left side of the map. The green boxes at the top of the maps, the

clusters of sites in the connectivity maps collapsed in the vertical axis whose stimulation evokes EPSCs.

Bottom, EPSC traces evoked at the sites indicated by letters in the maps above. Two repetitions are

superimposed (black and orange). Vertical dashed lines, the stimulus onsets (2 ms stimulus).

E. LSPS evoked excitation of two L5a pyramidal neurons recorded in current-clamp mode. LSPS was a 8 x

8 (left) or 24 x 8 (right) 50 µm spacing grid. Cortical neurons are positioned at the center (white

triangles). Traces with an action potential are in red. Bottom, Average number of action potentials

evoked at every site of the LSPS grid.

F. Overlay of the sites in the barrel cortex (green polygons) where stimulations evoked EPSCs in SPNs

(green symbols; n =101 cells, N = 54 mice). The red dashed vertical line is the reference (Ref$_{hor}$ in the

Methods) used for aligning slices across experiments horizontally.

G. Contribution of each cortical layer to the SPN innervations. The 16 rows of the grid correspond to

different cortical layers (Layer 2/3: 1-6; L4: 7-9; L5a: 10; L5b: 11-13; L6: 14-16). The horizontal white

band is L5a.

H. Amplitude of SPN EPSCs as a function of their laminar origin. Median (red) and 25-75th percentiles

(boxes). * Kruskal-Wallis *p* = 0.000125, Dunn-Šidák posthoc tests, *p* = 0.0001 and 0.03415.

I. Bottom, Positions of SPNs in the striatum for each position of connected sites on the horizontal axis of

the LSPS grid. Top, maximal distance between SPNs (d$_{SPN}$) for every connected site position in the barrel

cortex on the horizontal axis, binned every 150 µm, in other words the width of the projection zone of

one cortical column within the striatum

J. Schematic of the slice with the size of the cortical and striatal regions that are connected by

projections to SPNs (pink) and the size of the striatal domain with functional projections from a single

cortical column (shades of green).

Figure 3. Sparse functional projections from the barrel cortex to individual SPNs in the DLS.

A. Example slice with 3 recordings. The hatched horizontal band is L5a, the polygons are the stimulation sites evoking EPSCs, the solid boxes on top are the connectivity clusters and the open rectangle the input field in the collapsed connectivity map. The circles at the bottom mark the positions of SPNs on the horizontal axis. Recordings are color coded.

B. Fraction of SPNs receiving inputs from 1 to 6 clusters of projections in the barrel cortex. In inset, median (red) and 25-75th percentiles (box). Clusters are defined as the ensemble of contiguous sites in the connectivity map collapsed in the vertical axis, whose stimulation evokes EPSCs (see examples in A, boxes at the top). n = 101 cells, N = 54 mice.

C. Median (red) and 25-75th percentiles (box) of cluster width and spacing.

D. Left, Fraction of cells receiving inputs from 1 to 8 cortical columns in the barrel area. In inset, median (red) and 25-75th percentiles (box). Right, Number of cortical columns innervating individual SPNs. Symbols are cells, lines are slices. Light gray symbols, slices with ≥ 2 recordings. Dark symbols, one recording per slice.

E. Fraction of SPNs with input fields from 0.075 to 1.6 mm.

F. Top to bottom, For every input field width, the number of clusters, cluster width (expressed in cortical
column and μm) and proportion of connectivity gap, in input fields collapsed along the vertical axis. Gray
symbols, cells with ≥ 2 connectivity clusters. Other symbols, cells with 1 cluster.
G. Bottom, width of input fields in the barrel cortex for different positions of SPNs in the striatum. SPNs
with an input field the size of a single or of several cortical columns are shown (open and solid symbols,
respectively). Top, The average width of input fields as a function of SPN positions binned every 250 μm.
H. Schematized average and principal connectivity patterns. In pie charts, the SPN proportions.
638

Figure 4. Individual SPNs have unique cortical innervation
A. Four example slices with 2 to 4 recordings. Same as in Fig. 2A.
B. Left, fraction of pairs with overlap in their input field. Right, width of overlap as a function of the
horizontal distance separating cells. n = 70 pairs.
C. Same as in B for the fraction of vertical alignment between SPN connectivity maps.
D. Same as in B for the fraction of overlap between SPN connectivity maps.

Figure 5. Topographic organization of the functional vibrissal innervation of SPNs in the DLS

A. Overlay of the sites in the barrel cortex where stimulations evoked EPSCs in SPNs. The colors, yellow to blue, indicate the position of the SPNs (gray circles) along the horizontal axis in the striatum (axis at the bottom, 0 is Ref_{hor} in the Methods; n = 101, N = 54). Blue shades are for lateral SPNs.

B. Position of the connectivity center of mass (CM) as a function of the SPN position in dorsal striatum. Zero on the x and y axis is the position of the vertical dashed line shown in A.

C. Ratio of the largest EPSC over the smallest EPSC for each recording. Median (red) and 25-75th percentiles (box).

D. Position of the synaptic input CM as a function of the SPN position in dorsal striatum. In light gray, the connectivity CM.

 E. EPSC sum obtained in the strongest connectivity cluster relative to others sum. Cells with ≥ 2 clusters, n = 55.

Figure 6. D1 and D2 SPNs have similar patterns of innervation from the barrel cortex

A. Overlay of the sites in the barrel cortex where stimulations evoked EPSCs in D1 (blue) or D2 SPNs
(red). The shades, light to dark, indicate the position of the SPNs (bottom circles) along the medio-lateral
axis in the dorsal striatum (axis at the bottom). D1 cells, n = 47, N = 36; D2 cells, n = 54, N = 37.

B. Position of the connectivity center of mass (CM) as a function of the SPN position in the dorsal

striatum (D1, red; D2, blue). R = 0.70 (D1, n = 47, N = 36); R = 0.57 (D2, n = 54, N = 37).

C. Left, D2 SPN input field width as a function of the D1 SPN input field width, in the same slice. The

black symbol and lines are the median value and 25-75th percentiles. Large symbols indicate n = 2 to 3

pairs. n = 42 pairs. Right, medians (thick lines) and 25-75th percentiles (boxes) of D1 and D2 input field

widths, across all cells. n = 101.

D. Same as in C for the number of cortical columns.

E. Same as in C for the width of connectivity clusters.

F. Contribution of each cortical layer to the SPN innervations. The horizontal hatched band is L5a.

G. EPSC sum for D1 and D2 SPNs. Outliers were not shown for clarity (1 D1, 2 D2, 1.2-1.5 nA)

H. EPSC sum obtained from the strongest cluster relative to others. Cells with ≥ 2 clusters. D1, n = 24;

D2, n = 30.

I. Top, D1 and D2 SPN EPSC sum as a function of laminar origin. For each layer, only cells with inputs are

included. Outliers are not shown for clarity (3 D1, 2 D2, 350-700 pA). * indicates a significant difference

(p = 0.005 and p = 0.008; Mann-Whitney). Bottom, fraction of cells with input.

Figure 7. Connectivity pattern within the somatosensory corticostriatal projection to SPNs.

Left, Our results support the model in which each SPN integrates limited and heterogeneous rather than

exhaustive inputs transmitted by the barrel cortex, intermediate between the middle and right panels in

Fig. 1. Right, Each SPN representation of the whisker cortical columns complements the representations

of its neighbors.

Supplemental Figure 1.

A. Left, the corticostriatal slice generated from the Allen mouse brain reference atlas (CutNII custom-angle slice visualization tool; G. Csucs). bf, barrel field. str, striatum. VPL, ventral posterolateral thalamic nucleus. VPM, ventral posteromedial thalamic nucleus. Center, the section angles illustrated on horizontal and longitudinal sections of the brain. Right, the section angle with respect to the orientations of the arcs and rows in the barrel field of the left hemisphere (Allen mouse brain atlas).
B. A simulated somatosensory corticostriatal slice generated using data from the Allen mouse connectivity atlas (https://connectivity.brain-map.org/projection/experiment/cortical_mp/293728197). eGFP was expressed in L2/3 and L5 pyramidal cells principally, in 3-4 arcs of the D and E rows of whisker columns (whole brain visualization in the green box at the bottom left; the dashed lines indicate the slice orientation). Acquisitions of serial two-photon tomography were stacked in Fiji and the reconstituted brain was sectioned using the angles to produce the acute somatosensory corticostriatal slices (BigDataViewer plugging). The red box identifies the "experimental slice" with intact projections from the barrel cortex to the striatum. Fluorescence in the striatum (blue) and in the corpus callosum (cc, pink) was quantified on consecutive max projections (300 µm thick) obtained in the anterior-posterior

703 axis (A-P). Note that slices, being from the opposite hemisphere to our experiments, have the reverse
704 orientation in the medio-lateral axis compared to Fig. 2A,C.
705
706

Supplemental Figure 2.
A. Vertical (left) and horizontal (right) excitation profile of L5 pyramidal cells recorded in current clamp,
in the juveniles and adolescents combined (20 and 25 mW stimulation, respectively). Evoked action
potentials (APs) were summed along each column (left) or line (right) of the LSPS grid (50 µm spacing).
Solid line, median. In gray, 25-75th percentiles. n = 30, N = 14.
B. Total number of APs evoked in the stimulation grid for L5 (left) and L2/3 (right) pyramidal cells in
juvenile (juv, P22-30) and adolescent (ado, P31-41) mice. A higher stimulation intensity was used for
adolescent mice (25 mW instead of the standard -std - intensity, 20 mW), so that the total number of
APs matched between the two age groups. L5, juv, n = 22, N = 9 ; ado std, n = 22, N = 13; ado high: n = 8,
N = 5; L2/3, juv: n = 16, N=3; ado std and high n = 10, N = 2; * p < 0.001, Mann-Whitney.
718