Modulation of SAL retrograde signaling promotes yield and water productivity responses
in dynamic field environments

Andrew F. Bowerman[1*], Marten Moore[1*], Arun Yadav[1*†], Jing Zhang[1], Matthew D.
Mortimer[1], Zuzana Plšková[1], Estee E. Tee[1,2], Eng Kee Au[1], Derek P. Collinge[1], Gonzalo
M. Estavillo[1,3], Crispin A. Howitt[3], Kai X. Chan[1], Greg J. Rebetzke[3], Barry J. Pogson[1†]

[1] Research School of Biology, The Australian National University, Canberra, 2601, ACT,
Australia

[2] The John Innes Centre, Norwich Research Park, Norwich, NR4 7UH, United Kingdom

[3] CSIRO Agriculture and Food, Canberra, 2601, ACT, Australia

[*] AFB, MM and AY contributed equally to this work

[†] Corresponding Authors: Barry J. Pogson barry.pogson@anu.edu.au, Arun Yadav
arunkumar102@gmail.com. BJP is the nominated corresponding author for the article.

ORCID details: AFB: 0000-0003-1729-7843. MM: 0000-0002-3423-2403. AY: 0000-
0003-1350-9191. JZ: 0000-0001-8780-069X. MDM: 0000-0002-8135-9319. ZP: 0000-
0001-8954-559X. EET: 0000-0003-2613-5977. EKA: 0009-0002-2841-7310. DPC:
0000-0002-3940-3777. GME: 0000-0003-4634-5103. CAH: 0000-0002-1533-6602.
KXC: 0000-0003-3554-7228. GJR: 0000-0001-7404-0046. BJP: 0000-0003-1869-2423.

Short title: "SAL retrograde signaling modulation affects wheat yield in field
environments"

Twitter: @BiologyANU, @barry_pogson, @ngryDocB. Bluesky:
@biologyanu.bsky.social, @ngrydocb.bsky.social

Keywords: Drought, retrograde signaling, wheat, SAL, PAP, chloroplast,

photosynthesis,

Introduction: 998

Materials and Methods: 1464

Results: 2857

Discussion: 1675

Total: 6994

Figures: 6. Tables: 3

Summary

• Chloroplast-to-nucleus retrograde signaling enables rapid stress responses in
plants, but whether these signals accumulate to affect crop performance across
entire growing seasons under field conditions remains unknown.

• We generated wheat mutants with targeted deletions in specific SAL gene copies
from two distinct homeologous groups (TaSAL1 and TaSAL2), creating lines with
enhanced stress signal responsiveness. We tested these lines across 15 field
trials spanning diverse Australian environments with varying temperatures,
rainfall, and irrigation regimes, measuring physiological responses, yield,
biomass, and water productivity.

• Lines with TaSAL2 gene deletions showed 4-8% yield improvements with
enhanced water productivity, while TaSAL1 deletions reduced yields. The
TaSAL2 mutants maintained superior photosynthetic function under drought
stress, showed improved relative water content, and demonstrated enhanced
yield stability across environments. Canopy temperature measurements revealed
dynamic stomatal regulation, with increased closure during midday stress periods
but normal aperture under benign conditions. Significantly, specific SAL
modifications enhanced photosynthetic efficiency and stress resilience without
traditional yield penalties.

• Targeted modification of specific SAL homeologous groups can simultaneously
improve both yield and stress tolerance in wheat. This demonstrates that
retrograde signaling integrates environmental information across the plant
lifecycle, and highlights the importance of locus-specific targeting and multi-
environment field validation for crop modifications.

Introduction

Climate variability poses an escalating threat to global food security, demanding urgent advances in crop stress tolerance (Bowerman *et al.*, 2023). Modern crops face an unprecedented combination of abiotic stresses - from extreme temperatures and water availability to excess light and nutrient imbalances - often occurring simultaneously or in rapid succession. Plants coordinate their responses to these challenges through intricate cellular signaling networks, where even subtle changes in key pathways can trigger extensive transcriptional and physiological adaptations. Among these networks, chloroplast-to-nucleus retrograde signaling plays a particularly crucial role in integrating environmental cues with cellular responses (for review, see Pogson *et al.* (2008; Chan *et al.*, 2016b; Crawford *et al.*, 2018; Calderon & Strand, 2021)). These signals are essential for coordinating cellular responses to transient stress, and for optimal plastid operation given the number of plastid-located protein complexes that are nuclear encoded and subject to damage by oxidative stress. (Jan *et al.*, 2022).

The SAL enzyme, an adenosine phosphatase (EC:3.1.3.7), regulates the levels of 3'-phosphoadenosine 5'-phosphate (PAP), a chloroplast-to-nucleus signal molecule (for review, see Jan *et al.* (2022)). SAL converts PAP into adenosine monophosphate (AMP), and is a negative regulator of excess light and drought tolerance in *Arabidopsis* (*Arabidopsis thaliana* (L.) Heynh.). Under drought or light stress, AtSAL1 becomes inactivated by oxidation, allowing PAP to accumulate and move between organelles, increasing up to 30-fold in *Arabidopsis* (Estavillo *et al.*, 2011; Chan *et al.*, 2016a). Elevated PAP affects the expression of a raft of genes, partly through inhibition of exoribonucleases (XRNs) and resulting promotion of RNA polymerase II read-through at transcription termination sites (Estavillo *et al.*, 2011; Gigolashvili *et al.*, 2012; Crisp *et al.*, 2018; Ashykhmina *et al.*, 2019).

Studies of retrograde signaling in general (and SAL in particular) have largely focused on gene knockout lines in model species. Without a functional SAL protein, such knockouts constitutively accumulate PAP to high levels without stress induction, approximately 20-fold in *Arabidopsis alx8* mutants (Estavillo *et al.*, 2011). These lines exhibit increased drought stress tolerance (40-50% longer survival compared to wild

type) and improved water use efficiency with a decreased growth rate, dwarf phenotype,

and leaf morphology changes (Rossel *et al.*, 2006; Wilson *et al.*, 2009). PAP itself has

been implicated directly in the control of stomatal aperture closure across diverse taxa

with functioning stomata, from mosses and ferns to monocots and dicots (Zhao *et al.*,

2019). PAP bypasses and augments canonical ABA signaling; this includes exogenous

and endogenous addition of PAP to ABA-insensitive *Arabidopsis* lines restoring

stomatal closure (Pornsiriwong *et al.*, 2017). Broader effects of knocking out AtSAL1

include changes in hormone levels, induction of phosphate starvation responsive gene

expression, and modified metabolome including altered levels of osmoprotective

compounds (Rossel *et al.*, 2006; Wilson *et al.*, 2009; Hirsch *et al.*, 2011). Thus,

determining the transient stress-responsive functions of PAP *bona fide* is confounded

by the persistently elevated concentrations of PAP in mutants, which result in

developmental anomalies (Phua *et al.*, 2018a).

Recent studies in wheat (*Triticum aestivum* L.), which possesses a more complex SAL

gene family than *Arabidopsis*, show varied outcomes from attempts to knock down all

SAL copies simultaneously, including purported yield penalties alongside improved

resilience. VIGS-mediated silencing improved drought tolerance markers (Manmathan

et al., 2013), while CRISPR-based knockouts reported enhanced stress resistance

based on osmotic shock experiments (Abdallah *et al.*, 2022) and yield penalties in pots

subject to competitive glasshouse conditions (Mohr *et al.*, 2022); noting that pot-based

yields are perilous measures (Passioura, 2006). None of these studies considered the

roles of different SAL homeoforms, assayed any traits in the field, or measured the

impacts of the genetic modification on the enzyme or its substrate. Nor did they

consider whether retrograde signaling is a transient process or is integrated across the

life cycle.

Indeed, more broadly retrograde signaling research has focused on standardized

conditions with single stresses, rather than the complex, unpredictable stress

combinations crops face in the field, and ignored if it has a cumulative effect on the

entire life cycle of plants in response to multifactorial stresses (for review of

multifactorial stress, see Zandalinas & Mittler (2022)). The SAL-PAP pathway's role in

coordinating chloroplast stress response, regulating hormonal signaling, and metabolic
responses makes it ideal for studying these complex conditions (Phua *et al.*, 2018b).
However, this necessitates a different strategy to simple loss-of-function genetic
modifications and single stresses in controlled environments.

The challenges we meet in this report are to first test the null hypothesis that short-term
changes mediated by retrograde signaling pathways have no ongoing impact on crops
grown in dynamic and varied environments. Second, to consider how do we best
translate findings in a model plant under controlled environments into crops that are
selected for high and stable yields across multiple environments? Our strategy was to
create knockouts of select TaSAL homeologues in wheat. SAL is typically in excess
(Chan *et al.*, 2016a), which means that PAP levels remain low unless the stress is
significant, or AtSAL1 transcript levels are reduced by more than 80% (Phua *et al.*,
2018a). By reducing one to two of the six homeologues present in cv. Chara, we
engineered plants with slightly diminished overall SAL activity, thereby lowering the
threshold for PAP signaling. We postulate that such primed plants would respond more
dynamically to environmental perturbations. We utilized heavy ion bombardment (HIB)
to knockout *TaSAL* genes, enabling us to assess these materials in natural
environments during a period when gene-edited field trials were not authorized (2015-
2019) and producing a crop that is globally far more deployable to industry. Our
evaluation of the primed retrograde signaling plants in natural environments was
conducted across 15 distinct settings, covering four different cereal-growing districts
over five years, characterized by varying rainfall and irrigation regimes. Subsequently,
we incorporated the observations into a multi-environment trial (MET) model to examine
whether modulation of retrograde signaling influences altered lifetime responses to
diverse and dynamic natural environments. An additional aspect of the study involves
an evaluation of the trade-offs in yield resilience for genes that can modulate both stress
responses and developmental processes.

Materials and Methods

Phylogeny and expression analysis

Sequence homologues of AtSAL1 (NP) were identified from Liliopsida using Uniprot BLAST (e-value<0.1). A sequence similarity network was constructed and filtered to retain sequences with ≥ 40% identity that clustered with *Arabidopsis* (*Arabidopsis thaliana* (L.) Heynh.) SAL1/2/3/4 references, and sequences between 250-600 amino acids in length. Wheat (*Triticum aestivum* L.) sequences were updated using homologues identified from Gramene, and AHL enzyme sequences were added as an out-group. Multiple sequence alignment was performed using T-COFFEE, refined with MAFFT, and manually curated. Sequences with non-consensus insertions/deletions were removed. Phylogenetic reconstruction used ModelFinder for model selection, with multiple replicates and ultrafast bootstraps. The final phylogeny was selected based on outgroup contiguity and visualized in R. Detailed protocols and software versions are provided in Supporting Methods S1.

Expression data were retrieved from the Wheat RNA-seq Database https://plantrnadb.com/wheatrna/ (Zhang *et al.*, 2020) and from the database of Loudya *et al.* (2021). Gene co-expression analysis was performed using ATTED-II, using all members of each TaSAL group as seed sequences. The top 100 co-expressed genes for TaSAL1 and TaSAL2 were then used in ShinyGo v0.80 to summarise GO terms related to these co-expressed genes (Ge *et al.*, 2020).

Identification of *TaSAL* mutants and generation of double null lines

A wheat population mutagenised by heavy ion bombardment (HIB) was used to search for mutations in each *TaSAL* homeologues (Regina *et al.*, 2015). Rather than single nucleotide changes which are seen in chemical mutation-based populations, HIB mutations result in large deletion mutations, usually encompassing entire genes. A total of 3000 individual M3 HIB lines were screened using homeoform specific primers for deletions in each *TaSAL* gene by PCR, resulting in the identification of five primary null lines, three deletions on chromosome 4A and two on chromosome 5D. Primer sequences are listed in Supporting Information Table S1. Multiple crosses were

generated using combinations of each of these null lines; all line details are found in

Supporting Information Table S2. Deletions were characterised using two targeted

genotype-by-sequencing (tGBS) approaches, completed by Agriculture Victoria

(AgriBio, Bundoora, Australia). Briefly, one tGBS assay was constructed from a subset

of the Infinium 90K SNP chip (Chidzanga *et al.*, 2021), and the second tGBS method

was constructed using exome sequencing results from approx. 1000 wheat lines

(Sharma *et al.*, 2022). Combined, this maximised the marker density and maximised our

chances of identifying deletions. See Supporting Information files details of deletion

characterisation.

Plant growth and drought stress in glasshouse

All lines were grown in a controlled greenhouse with day-length extension to 16h at

24/16°C day/night temperature and 60% relative humidity. Pots of 25 cm (diameter) ×

30 cm (h) were filled with compost-based potting mix and arranged in a randomised

complete block design. All pots were watered daily to pot capacity until drought stress

was imposed at the 3-4 leaf stage (21 days). Two drought regimes were used. **Cyclic**

Drought: At day 0 of drought, water was withheld in droughted plants until day 21 of the

first drought and then re-watered to their 100% soil water capacity as per Bowne *et al.*

(2012). **Terminal drought**: All pots were saturated with water at day 0 and designated

drought-stressed plants were not watered up to 42 days of drought. All control plants

were maintained at 100% water capacity. Where required, soil weight measurements

were collected to compare water use during drought periods.

Field trial management

Performance of genotypes was assessed under a series of field trials undertaken

across four seasons at four managed environment facilities in New South Wales and

Victoria, Australia. Details of facilities can be found in Rebetzke *et al.* (2013), and are

indicated, along with temperature and rainfall variation between and within sites in

Figure 4. Multiple trials were performed in each site/year with varying access to

irrigation to ensure variation in water availability between trials. All designs were

produced using a complete randomised block design approach. In all experiments, all

the plots were treated with 30 mm of pre-sowing irrigation and entries were sown at an

optimal 3 to 5 cm sowing depth into 6-m long, 0.17-m spaced, 8-row plots at seeding

rates consistent with local practice (i.e. 120 seeds per m^2 at Narrabri, and 180 seeds

per m^2 at Yanco and Condobolin). Where irrigation was supplied, it was calculated at

the site to target 25% above mean rainfall. Nutrients were supplied at sowing as Starter

15® (14% N: 12.7% P: 11% S) applied at 103 kg.ha^{-1}. Additional nitrogen was applied as

needed to meet crop demand. Plots were end-trimmed to 5.4 m length and border rows

removed prior to machine harvesting for plot yield estimates.

PAP quantification, SAL activity assay and protein quantification

PAP was quantified using a modified version from Ashykhmina et al. (2019).

Metabolites were extracted from liquid nitrogen-frozen leaf tissue (30-50 mg) using

chloroform:methanol and citrate-phosphate buffer extraction. PAP was quantified by

HPLC following chloracetaldehyde derivatization using commercial PAP as a standard

(Sigma-Aldrich, A5763). Separation was performed on a Kinetex XB-C18 column

(Phenomenex) using tetrabutylammonium hydrogensulfate and acetonitrile buffers. SAL

activity was measured using protein extracts from leaf tissue incubated with PAP

substrate at 37°C, with enzymatic activity determined by measuring PAP degradation

via HPLC. Detailed protocols are provided in Supporting Method S2.

ABA quantification

Leaf ABA was quantified using a modified extraction protocol. Approximately 100 mg of

flash-frozen leaf tissue was homogenized in a Tissue Lyser II (Qiagen, Germany) with a

steel ball bearing at 19.5 Hz for 2 minutes. The tissue was extracted twice with 2 mL

cold 80% methanol by overnight shaking (25 rpm, 4°C) followed by centrifugation

(17,000 rcf, 4°C). After each extraction, supernatants were vacuum concentrated to 500

μL using a miVac Duo Concentrator (GeneVac, UK). A final methanol wash step was

performed and concentrated to 500 μL. Combined concentrated extracts (1.5 mL total)

were further reduced to 500 μL under vacuum. Extracts were acidified to pH 2.5 with 0.5

mM HCl and partitioned against equal volumes of ethyl acetate. The organic phase was

dried under vacuum and resuspended in 1 mL Tris-Buffered Saline (TBS). ABA was

quantified using the Phytodetek ABA Kit (Agdia Inc, USA) according to manufacturer's

instructions, with absorbance measured at 405 nm (Tecan Group, Switzerland). Final

ABA concentrations were expressed as ng/g dry weight.

Photosynthesis, stomatal conductance, relative water content and carbon isotope

discrimination

An open gas-exchange system with a 2 × 3 cm clamp-on leaf cuvette was used to

simultaneously measure stomatal conductance and photosynthesis (Li-6400; Li-Cor,

Inc., Lincoln, USA). The topmost expanded leaf was chosen for taking measurements.

Measurements were taken after leaves were adapted in the chamber for approximately

10 minutes at 380 μmol CO_2 and light intensity of 1000 μmol/m/s, block temperature of

24°C, and ambient humidity approximately at 55%.

Relative water content was measured as in Wilson *et al.* (2009), with leaf sections cut

and weighed to measure fresh weight (F_W). Leaves were incubated in water at 4°C in

dark conditions, blotted, and re-weighed to measure turgid weight (T_W). Finally, leaves

were dried at 80°C overnight, weighed to determine dry weight (D_W), and relative water

content was calculated as

$$RWC = \frac{F_W - D_W}{T_W - D_W}$$

Samples of dried and ground plant material were weighed to 1.5 mg (within 0.2 mg) into

folded tin foil cups. These were then combusted in a Carlo-Erba EA-1110 Elemental

Analyser (Carlo Erba, Milan, Italy) providing a controlled oxidation and reduction. Gases

were separated in the helium carrier using a packed gas chromatograph (HayeSep D

80/100, Grace Davison, Australia) at room temperature and passed to a stable isotope

ratio mass spectrometer (Isoprime, Micromass U.K., Manchester, U.K.). A pulse of

reference CO_2 succeeded each CO_2 sample peak, to correct for pulse to pulse variation.

Standards used were a C3 beet sucrose (Beet89) with a $\delta^{13}C$ of -24.62‰ Vienna

Peedee Belemnite (VPDB) and ANU C4 cane sucrose with an accepted value of -

0.45‰ VPDB. Each run commenced with two of each sucrose standard and proceeded

with pairs of standards every 10 samples. Final corrections were made using the slope
and offset derived from the two standards. Values were then converted to Δ form using

$$\Delta(\text{‰}) = \frac{\delta_a - \delta_p}{1 + \delta_p} \times 1000$$

where δ_a and δ_s are the $\delta^{13}C$ values for atmosphere (-8‰) and the sample, respectively
(Farquhar *et al.*, 1989).

Hyperspectral imaging and leaf temperature measurements

Hyperspectral measurements were taken using ASD FieldSpec 3 (Boulder, CO, USA).
Photosynthetic parameters were then estimated using the Wheat Physiology Predictor
at https://wheatpredictor.appf.org.au/ (Furbank *et al.*, 2021).

Canopy temperatures were measured at Narrabri, NSW, on two consecutive days in
2017 and once in 2019 using infrared cameras deployed on unmanned aerial vehicles
(UAVs). Six measurements were taken over seven hours of daylight to capture diurnal
temperature patterns. Data were analysed using multivariate repeated measures linear
mixed effects models, with plot as a random effect and time, genotype, and water
treatment as fixed effects. Best linear unbiased estimates (BLUEs) were calculated for
each time point.

Statistics

All statistics were performed using R (R Core Team, 2021). Field analysis was
performed using ASREML-R (Butler, 2022) or the 'lme4' package (Bates *et al.*, 2015).
Multi-environment trials were analysed using the two-stage approach of Smith *et al.*
(2001) facilitated by the use of the 'ASRtriala' package (Gezan *et al.*, 2022). Models
best fitting a trait in each trial were first constructed before combining using the Factor
Analytic framework, identifying the optimal number of factors for each trait.

Results

Identification of null alleles in *TaSAL* subtypes

Previously 5-7 SAL genes were identified in different wheat varieties (Mohr *et al.*, 2022). Phylogenetic analysis of monocot SAL homologues showed independent duplications in both Pooideae and Panicoideae subfamilies of the Poaceae, with retention suggesting functional importance of both groups (Figure 1a). We designated these as TaSAL1 (5A/B/D) and TaSAL2 (4A/7A/7D), with the 4A copy resulting from a known 4AL/7BS translocation (Devos *et al.*, 1995). Chara cv., used herein, contained six copies across two paralogous groups: 5A/B/D and 4A/7A/7D.

Analysis of public RNA-seq based expression data revealed that TaSAL1 homeologues showed 10-fold higher expression than TaSAL2 and different developmental profiles (Figure 1b, Supporting Information Fig. S1) (Loudya *et al.*, 2021). Co-expression analysis of the TaSALs revealed enrichment in distinct functional associations: TaSAL1 with sulfur metabolism (11-fold), DNA repair (15-fold), and aminoacyl-tRNA pathways (62-fold), while TaSAL2 associated with carbon (14-fold) and amino acid metabolism (37-fold), suggesting roles in photosynthesis and energy metabolism (Figure 1c).

Given this evidence of differential expression and coexpression, we aimed to investigate the relative impact of targeted reductions in TaSAL1 and TaSAL2. We screened a Chara HIB population (M3 generation) for single gene knockouts (Supporting Information Fig. S2). Five lines were identified, two in TaSAL1-5D (designated SAL1.1 and SAL1.2) and three in TaSAL2-4A (SAL2.1, SAL2.2, and SAL2.3), and systematic crosses between these lines were generated (Supporting Information Table S2).

Genomic characterisation of background deletions in the 12 lines was performed using two targeted Genotype-by-Sequencing (tGBS) methods to genotype these lines and identify deletions. The GBSs confirmed the PCR demonstrating that all lines had complete deletions in the respective TaSAL homeologue (Supporting Information Dataset 1). The size of the deletions varied from minimal additional deletions of other loci, through to line SAL2.1, which was shown to have an additional large deletion in Chromosome 1B. Other than this line and its associated double mutants, no other

incidental mutations were shown to alter traits analysed in this study, nor were any other

gene deletions identified in all mutants that would be consistent with the observed

phenotypes.

Selective knockout of SAL genes lowers SAL activity under stress

To ascertain the effects of SAL mutations on PAP concentrations and enzymatic

activity, lines were assessed under controlled glasshouse conditions with adequate

water and drought schedules (detailed in Methods). The quantification of PAP levels in

wheat has not been previously reported and required optimisation of HPLC conditions,

including column selection (Kinetex XB-C18 vs standard C18), gradient simplification,

and increased acetonitrile concentration to improve separation of co-eluting compounds

that interfered with PAP detection.

Under sufficient water, there was no significant difference in PAP levels (Supporting

Information Fig. S3) or SAL activity between mutant and control lines. However, under

moderate Cyclic Drought, most mutant lines showed significant decreases in SAL

activity of up to 60% compared to Chara, and several double mutant lines showed

increased PAP accumulation at anthesis (Figure 2, Supporting Information Fig. S3).

These genotype-dependent differences were confirmed statistically (genotype:treatment

interaction: PAP content $F_{(12,52)}=2.42$, $p<0.05$; SAL activity $F_{(12,52)}=4.01$, $p<0.001$).

Relative water content (RWC) was measured at 7, 14 and 21 days under the Cyclic

Drought regime (Figure 2, Supporting Information Fig. S4B), and modeled using

ASREML-R as a repeated measures experiment by using a uniform multivariate model.

By day 14 four of five single nulls demonstrated increased RWC compared to Chara, as

did two double null lines. At 21 days, two of three TaSAL2 lines maintained this water

advantage, as did one of two TaSAL1 null lines.

To determine whether improved drought performance resulted from reduced water use

or genuine stress tolerance, we analysed pot weight depletion during drought stress

(days 0-25; Supporting Information Fig. S5). Mixed-effects modelling revealed no

significant genotype×time interaction ($F_{(12,481)}=0.059$, $p=1.000$), with water depletion

rates ranging from 1.26 ± 0.23%/day to 1.46 ± 0.23%/day across all genotypes. No

significant differences were detected between any TaSAL mutant and Chara.

These findings demonstrate that all genotypes experienced equivalent rates of soil

water depletion, ruling out water use avoidance as an explanation for improved drought

performance. The data support a tolerance mechanism whereby TaSAL mutants

maintain superior physiological function whilst experiencing identical soil water. This

priming of the SAL-PAP pathway creates lines that respond more dynamically to

drought while maintaining normal function under unstressed conditions.

Physiological analysis of unstressed and stressed TaSAL germplasm

No obvious changes to crop phenological traits were identified in TaSAL germplasm.

Single mutants maintained comparable biomass, plant height, and tillering. From the 3rd

leaf stage, plants were grown in two different watering regimes for 35 days. The well-

watered regime was to sufficiency (normalized to pot weight), while drought was

enforced by the Cyclic Drought regime. Single mutant lines had comparable plant height

to Chara within regimes, whereas most double mutants were reduced in stature

(Supporting Information Fig. S4A). The Treatment and Genotype interaction was

significant ($F_{(12,43.4)}$ = 2.6, $p < 0.05$), indicating that drought stress had differential

impacts on genotypes, with TaSAL mutants maintaining higher relative water content

than Chara under water limitation (Supporting Information Fig. S4A).

To assess drought effects on photosynthesis (A) and stomatal conductance (g_s), we

subjected four lines at the 5/6 leaf stage plants to two regimes, well-watered (control)

and withholding all water for up to 42 days (Terminal Drought regime). While

photosynthesis remained stable in the control regime (Figure 3, Supporting Information

Fig. S6), it declined significantly in Chara after 28 days of drought. TaSAL deletion lines

maintained higher photosynthetic rates compared to Chara over the drought period

($F_{(3,11)}$ = 6.67, $p = 0.008$). Stomatal conductance was also examined in the same

experiment. While drought reduced g_s across all lines ($F_{(1,24)}$ = 89.25, $p < 0.0001$, η_G^2 =

0.53), there were no genotype-specific differences. This maintenance of photosynthesis

despite similar g_s reductions suggests improved photosynthetic capacity under water

limitation in TaSAL mutants as seen in Figure 3.

The ratio of photosynthesis to water use is known as water use efficiency and was

investigated given the changes in *A* and RWC. A life-time integrated measure of water

use efficiency is carbon isotope discrimination. This was determined for grain collected

from plants grown under Terminal Drought and control conditions and presented as

delta (Δ). Δ is calculated using the carbon isotope ratio of both the plant samples and of

air (Farquhar *et al.*, 1989), and is positive in nature; typically C3 plants have a Δ of

approximately 20×10^{-3} or 20‰. Two-way ANOVA found no significant interaction

between the genotypes and the water regime, but both water regime and genotype

effects were significant and large ($F_{(1,29)}$ = 147.26, $p < 0.001$, η^2 = 0.84 (partial) and

$F_{(5,29)}$ = 10.05, $p < 0.001$, η^2 = 0.63 (partial), respectively). Drought increased Δ by

approximately 2.45 compared to watered control. Under both conditions, line SAL2.1

and a double mutant composed of line SAL1.1×SAL2.1 (DM) both have significantly

lower Δ than the Chara control. Additionally, under drought line SAL1.2 also has a

significantly reduced Δ (Table 1). In summary, given the lack of difference in g_s

combined with a reduction in Δ levels, we conclude improved maintenance of

photosynthetic capacity in TaSAL mutants under water-limited conditions.

We then measured ABA content of leaf material under both control and Terminal

Drought conditions on all single mutants to evaluate if there is any interaction between

ABA and PAP (Supporting Information Fig. S7). A three-way repeated measures

ANOVA was performed to evaluate the interaction of growth conditions and ABA

content, using paired sampling at days 0 and 28 of the experiment. There was a

statistically significant three-way interaction between genotype, treatment and time, ($F_{(5,}$

$_{40)}$ = 3.163, $p = 0.017$, η_g^2 = 0.16). At 0 days, interaction of genotype and treatment was

not significant, nor treatment, but some significance was applied to genotype alone ($F_{(1,}$

$_{40)}$ = 3.71, $p < 0.05$); conversely, genotype, treatment and their interaction were all

significant. Simple-simple main effects of genotype were found to be significant under

drought, and pairwise comparison to Chara control, using a Holm correction, found all

lines had significantly lower ABA content after 28 days drought. We identified that the

dilution of TaSAL had the effect of inhibiting the ABA increase due to drought seen in

the Chara control line. Critically, since all genotypes depleted soil water at equivalent

rates, this reduced ABA accumulation reflects enhanced stress tolerance rather than

stress avoidance, indicating that TaSAL mutants require lower stress hormone

concentrations to maintain physiological function under identical water limitation

conditions. Of interest is the wide variance in values for Chara, implying a higher stress

susceptibility compared to the single mutant lines. This observation is consistent with an

hypothesis of PAP-mediated resilience limiting stress sensitivity and consequential ABA

induction in the mutant lines.

Examination of retrograde signaling in field environments

Field trials were conducted at four distinct agroecological sites, Birchip, Yanco,

Condobolin, and Narrabri, between 2015 and 2019 (Figure 4). The sites encompass

essentially the breadth of the cereal growing regions of south-east Australia, separated

by ~1,000 km. The sites have varied soil types, average rainfall (decreasing with

latitude from 550 to 320 mm per year) and temperature profiles (from average monthly

highs of 28.9 to 24.3°C) (Summary Tables S3 to S7). The sites were used over five

years, resulting in large seasonal variation. See Methods for details on field

management and trial types. Collectively, this enabled assessment in 15 different

growth environments spanning years of both high and low water availability,

temperature and light. For example, actual total water availability ranged from 190 mm

to 725 mm across the growing seasons; actual October average temperatures during

grain fill ranged from 21 to 30°C for highs, and establishment lows for April ranged from

6 to 14°C (Figure 4, Supporting Information Tables S3-S7).

Field-based phenological assessments revealed no substantive changes in growth and

morphology relative to the parent line, Chara. The one exception was line SAL2.1 (the

line with substantive off-target deletions) and its associated double mutants, which

showed a shorter and spindly stature compared to Chara and the other two TaSAL2

deletion alleles, consistent with it having more off-target lesions (Supporting Information

Dataset 1).

To assess stomatal function in field conditions, we measured canopy temperature as a
proxy for stomatal aperture. Canopy temperature reflects stomatal aperture, which PAP
can directly and rapidly regulate (Wilson *et al.*, 2009; Estavillo *et al.*, 2011; Pornsiriwong
et al., 2017). Exogenous application of PAP closes wheat and barley stomata within
minutes, as rapidly as ABA (Pornsiriwong *et al.*, 2017; Zhao *et al.*, 2019), though PAP
acts through Respiratory Burst Oxidase Homolog D (RBOHD) rather than RBOHF for
apoplastic ROS production (Tee *et al.*, 2025). We measured canopy temperatures six
times over seven hours of daylight. Measures were collected at Narrabri, NSW, on two
consecutive days in 2017 and once in 2019 using infrared cameras deployed on drones
and helicopters.

TaSAL mutant canopy temperatures showed distinct diurnal patterns compared to
Chara (Figure 5, Supporting Information Fig. S8 and S9). In the morning, both 2017 and
2019 trials showed no significant differences between any genotypes and Chara,
indicating no constitutive alterations in stomatal aperture under cooler conditions.
During the middle of the day, however, TaSAL lines showed significantly increased
canopy temperatures compared to Chara, indicating greater stomatal closure. In 2017
trials, the pattern of significance differed between water treatments: under rainfed
conditions, lines SAL2.1 and the SAL1.1×SAL2.1 double mutant showed significant
increases, while under water-supplemented conditions, lines SAL1.2, SAL2.1, and
SAL1.1 were highly significant, with SAL2.2 and the SAL1.1×SAL2.1 double mutant
also showing significance. Temperature increases reached up to 1.2°C above Chara. In
2019 trials, conducted under higher ambient temperatures (maximum 27-31°C vs 19-
22°C in 2017), only lines SAL2.1 and the SAL2.3×SAL1.1 double mutant showed
significant increases at 1:15 pm under irrigated conditions, though the absolute
temperature differences were greater under rainfed conditions where significance was
not achieved. Significantly, the increased temperature/greater closure during the mid-
day period is consistent with an enhanced environmental sensitivity - more protective of
water losses - than exhibited by the controls. By the afternoon/evening, temperature
differences between genotypes returned to non-significant levels in both trial years.

This pattern is consistent with the lines having enhanced environmental sensitivity
rather than baseline stomatal defects, differing from prior publications in pot
experiments for complete knockout/down lines. The enhanced midday stomatal closure
suggests increased sensitivity to peak evapotranspiration conditions when demand is
highest.

Photosynthesis-related traits were predicted for field grown materials using
hyperspectral imaging, enabling the measurement of photosynthetic parameters at
scale. Hyperspectral data was collected for the two trials at Narrabri 2019 (Supporting
Information Fig. S10). Predictions of photosynthesis-related traits were produced using
the Wheat Physiology Predictor (Furbank *et al.*, 2021). This predicts leaf mass per area
(LMA), nitrogen content per unit area (N_{area}) and mass (N_{mass}), maximum rate of
carboxylation of photosynthesis at ambient temperature (V_{cmax}), 25°C (V_{cmax25}) and
corrected for nitrogen (V_{cmax25}/N_{area}, electron transport rate (J), CO_2 assimilation rate
(Photo), chlorophyll content (SPAD) and stomatal conductance (g_s/Cond).

Linear mixed effects models found no changes in V_{cmax} measures in either trial, with the
exception of one double null (SAL1.2×SAL2.2) in rainfed conditions, and another
double SAL2.3×SAL1.1 when expressed as V_{cmax25}/N_{area}. Maximum rates of
carboxylation were more stable under water supplementation for those lines than
rainfed, but no large differences could be identified for these traits. Leaf mass and
nitrogen per unit area were increased in line SAL1.2, or double nulls derived from line
SAL1.2 under rainfed conditions. These lines also exhibited an increased
SPAD/chlorophyll content. One line, SAL1.2×SAL2.3, demonstrated a consistent
increase in stomatal conductance, but no other lines exhibited this change. Overall,
photosynthesis was not constitutively altered in ambient conditions, consistent with no
constitutive changes in PAP levels.

Analysis of life-time integrated measures of retrograde signaling

Yield, biomass and water productivity were assessed as life-time integrated traits of
plant performance, reasoning that if the effect of repeated transient differences in

retrograde signaling is cumulative, altered PAP responsiveness will be reflected in the

alteration of these traits in different environments, compared to Chara (Figure 4).

These data were assessed by a two-stage multi-environment trial (MET) analysis of 15

trials that varied substantially for temperature, rainfall, irrigation or lack thereof, and

latitude. We used a Factor Analytic approach (Smith *et al.*, 2001; Gezan *et al.*, 2022).

As expected, variation between environments impacted yield, biomass and water

productivity (individual trial BLUEs presented in Figure 6; Supporting Information Table

S8); more interestingly, yield differences between SAL genotypes and Chara exhibited

considerable variability. For example, line SAL2.2 had a range of yield differences, from

an increase of over 30% in Condobolin 2017, down to a low of -7% in very hard

conditions in Birchip 2018.

Predictions of yield averaged across trials found one line, SAL2.2, had a significant

increase in yield compared to control (approximately 4%), while its sister, SAL2.3,

demonstrated an increased yield of 8%. In the integrated MET analysis this was not

significant, due to the small number of trials it was included in, however in those trials it

was grown, there was a statistically relevant increase when analysed as single sites

(Table 2). In contrast to the TaSAL2 yield gain, the TaSAL1 subtype had a consistent

yield penalty compared to Chara. (Table 2).

A multivariate 3rd order Factor Analytic model was used to combine these results with

appropriate weighting, allowing direct prediction of best linear unbiased predictions

(BLUPs) for all lines and to predict stability of yield across sites. FA1 accounted for 90%

of the variance. Together the three factors in the analytic model accounted for

approximately 95% of all variance across the environments. Model biplot indicating

cultivar (points) suitability to environments (arrows) and genetic covariance, indicating

the magnitude of genetic variation in each environment and the degree of genetic

correlation between environments, is included in Supporting Information Fig. S11. We

tested the model using environmental factors, such as latitude, rainfall, irrigation, and

temperature, via PCA or Random Forests and they did not explain FA1; rather, it was

best explained by the genotypes, namely deletions in SAL homologues.

Water productivity measures yield per unit of actual available water for a given site's
growing season (kg/Ha/mm). Line SAL2.2 showed higher water productivity compared
to the Chara control, demonstrating its improved yield given the same availability of
water (Table 2). Line SAL2.3 also showed improved water productivity compared to
Chara, though this improvement was not statistically significant in the MET analysis, but
was for the individual sites. Again, the TaSAL1 subtype mutants showed significantly
poorer water productivity compared to the Chara control and the TaSAL2 types.

Biomass was also measured in select field environments. Unlike yield and water
productivity, few differences in biomass were observed. Line SAL1.2 had a significant
increase compared to Chara, while the SAL1.1×SAL2.1 double mutant was smaller.
Overall, no patterns could be seen in biomass changes between lines and the Chara
control.

Stability of yield is a measure of the amount of variance of yield across different
environments for a given genotype. It was calculated using the Superiority measure (P_i)
of Lin & Binns (1988), using the mean square difference between a cultivar's yield and
the maximum yield in each environment. This measure is less affected by unbalanced
trials than other stability measures. The results indicated that the stability of the TaSAL2
null lines has improved compared to the Chara control, whereas the TaSAL1 lines have
had a decrease in stability (Table 3). This could reflect improved environmental sensing
for TaSAL2 and impaired sensing for TaSAL1 lines compared to Chara.

Discussion

Enhanced Crop Performance Through Precision Stress Signaling

There has been a view in crop improvement that there is often an inherent trade-off
between yield potential and stress resilience. Development of high-yielding varieties can
introduce environmental yield penalties or instability through inadvertent selection under
optimal conditions (Dietz *et al.*, 2021). For example, lettuce breeding led to shallower
root systems that perform well under irrigation but poorly in stress (Jackson, 1995). In
wheat, such trade-offs manifest in multiple traits including nutrition, flowering time, and
architecture (Dwivedi *et al.*, 2021). Even traits selected specifically for stress tolerance

can penalize yield - as seen with stem carbohydrate storage, which improves drought

survival but reduces grain-filling efficiency in good seasons (Blum, 1998; Rebetzke *et*

al., 2008).

Our findings indicate that this perceived trade-off can be overcome through precisely

modulating stress signaling pathways to enhance both productivity and resilience.

Rather than constitutive changes that improve stress tolerance at the cost of growth, we

created wheat lines with primed stress responses - plants that detect and respond to

environmental stress earlier and more sensitively, without developmental costs. This

enhanced environmental sensing translated to improved yield and water productivity,

water use efficiency and sustained photosynthesis under stress conditions. This

tolerance rather than avoidance mechanism provides the mechanistic basis for the

enhanced field performance observed across diverse water-limited environments.

To assess if retrograde signaling variation was functioning at field scale, we used

canopy temperature as a proxy for stomatal responses. TaSAL mutants showed

increased canopy temperatures specifically during midday periods, with no differences

morning or evening, indicating enhanced rather than constitutive regulation (Figure 5).

Temperature differences were more pronounced under rainfed versus irrigated

conditions (approaching 1°C), consistent with primed lines dynamically regulating

stomatal aperture to optimise photosynthesis and water conservation.

Across 15 diverse field environments, wheat lines with targeted reductions in specific

TaSAL2 gene copies consistently outperformed the parent cultivar. Such lines showed

yield improvements of 4-8% - a substantial gain considering current yield improvement

rates. Critically, these same lines also demonstrated enhanced water productivity and

stress tolerance.

However, the precision of genetic targeting proved crucial. While lines with TaSAL2

deletions consistently improved, those with TaSAL1 deletions showed yield penalties,

demonstrating that enhanced performance requires precise pathway modulation rather

than broad disruption of stress signaling. The contrasting effects of TaSAL1 versus

TaSAL2 deletions highlight that targeted modification of specific loci, rather than
complete pathway disruption, is essential for achieving these benefits.

Priming Environmental Responsiveness

Knockouts of TaSAL2 or TaSAL1 did not lower SAL activity or increase PAP in
unstressed conditions, but in response to water deficit did lower relative SAL activity in
most lines by up to 60% and could elevate PAP levels. The mild attenuation of SAL
activity is expected to allow PAP levels to have a greater capacity to rise early in stress.
In turn, this will lead to a modified pattern of stress signals then communicated to the
nucleus. What is significant is that titrating SAL activity raises PAP only under stress,
thereby generating a set of lines that prime retrograde signal responsiveness.

Compared to *Arabidpsis sal1* knockout mutants and previous wheat studies that
knocked out all or most TaSAL genes (Rossel et al., 2006; Wilson et al., 2009;
Pornsiriwong et al., 2017; Mohr et al., 2022; Abdallah et al., 2022), our primed, selective
knockdown lines exhibited minimal growth phenotypes during glasshouse evaluations
(Supporting Information Fig. S4). Under both watered and water-deficit conditions,
growth, tiller numbers, and plant height remained largely consistent with the Chara
control, suggesting that partial TaSAL reduction avoids the developmental penalties
associated with complete pathway disruption whilst enabling enhanced stress
responses. The primed lines demonstrated improved water use efficiency, maintained
relative water content, and sustained photosynthetic rates under drought conditions.
Most significantly, deletion of either TaSAL subtype enabled continued photosynthetic
assimilation during water limitation - demonstrating the benefits of selective over
complete pathway disruption (Figure 3, Supporting Information Fig. S6). This
maintenance of photosynthesis reveals the sophisticated interplay between chloroplast-
mediated sensing and ABA-independent PAP regulation of stomatal function
(Pornsiriwong et al., 2017), allowing dynamic rather than constitutive stress responses.

The molecular basis for this enhanced environmental sensing lies in PAP's dual role as
both a stress signal and stomatal regulator (Pornsiriwong *et al.*, 2017; Tee *et al.*, 2025).
PAP accumulates rapidly under oxidative stress when SAL enzyme activity is reduced

(Estavillo et al. 2011, Chan et al. 2016a). In our partially knocked-down lines, this
established mechanism creates a more sensitive environmental detection system.
Unlike constitutive stress responses, this priming mechanism is expected to allow PAP
levels to rise earlier and reach higher peaks during stress episodes, enabling plants to
detect and respond to environmental perturbations before damage occurs.

The transient nature of PAP accumulation in primed lines reflects the sophisticated
regulation of this pathway. Previous work in *Arabidopsis* demonstrated that
approximately 80% reduction in AtSAL1 activity is required before significant PAP
accumulation occurs (Phua et al., 2018a); our knockdown lines operate far below this
level. This creates a dynamic system where early PAP increases likely trigger rapid
stress responses (Estavillo et al., 2011) that could subsequently restore SAL activity
through feedback mechanisms, potentially suppressing further PAP accumulation. Such
regulatory feedback loops explain why PAP increases appear modest in single-
timepoint measurements despite the enhanced stress responsiveness observed in our
physiological assessments.

The equivalent water depletion rates across all genotypes during controlled drought
experiments definitively establish that enhanced performance results from improved
stress tolerance rather than reduced water consumption. This tolerance mechanism
explains the paradoxical observation of reduced ABA accumulation in TaSAL mutants
under drought: rather than indicating reduced stress exposure (avoidance), the lower
ABA levels reflect enhanced cellular resilience that reduces the need for ABA-mediated
stress responses. PAP acts directly on stomatal guard cells, bypassing traditional ABA
signaling pathways (Pornsiriwong et al., 2017), enabling more efficient stress responses
that maintain cellular function without requiring elevated stress hormone concentrations.

This tolerance rather than avoidance mechanism identified under controlled conditions
provides the mechanistic basis for the enhanced field performance observed across
diverse water-limited environments. PAP-mediated stomatal control enables dynamic
water management throughout the day. Field measurements of canopy temperature
revealed that primed TaSAL lines increased stomatal closure specifically during midday
stress periods while maintaining normal aperture during benign morning and evening

conditions. This temporal regulation may optimise the trade-off between CO_2 uptake for

photosynthesis and water conservation, explaining the observed improvements in both

water use efficiency and sustained photosynthetic rates under water limitation.

Gene-Specific Engineering Enables Targeted Pathway Modulation

The two homeologous groups were found to be the result of a conserved duplication

that we conclude has occurred twice in the Poaceae. The conservation of the

duplication in the Pooideae, distinct tissue expression and co-expression analysis

patterns, along with contrasting effects in field assessments of knockouts of TaSAL1

and TaSAL2 subtypes, are evidence towards specialization within subgroups; however,

the nature of this subgrouping is far from clear, be it simply distinct expression levels or

reflects subfunctionalisation, which may be suggested by the different co-expression

profiles for TaSAL1 and TaSAL2.

Co-expression analysis provided molecular evidence for functional specialisation

between TaSAL homeologue groups. TaSAL1 homeologues showed highest

expression in vegetative tissues and were enriched for fundamental cellular processes

including aminoacyl-tRNA biosynthesis and porphyrin metabolism, consistent with their

high expression during active vegetative growth phases (Soprano *et al.*, 2018; Adjei *et*

al., 2023). In contrast, TaSAL2-associated networks showed stronger enrichment for

metabolic flexibility pathways including amino acid metabolism, energy metabolism, and

secondary metabolism (phenylpropanoid biosynthesis).

Potentially, this molecular evidence could support the divergence of function between

gene families, with TaSAL1 specialized for supporting vegetative growth and TaSAL2

for environmental stress adaptation. These pathway enrichments align with established

roles of branched-chain amino acid degradation (Pires *et al.*, 2016), amino acid

metabolism coordination (Joshi *et al.*, 2010; Bowne *et al.*, 2012; Yadav *et al.*, 2019),

and phenylpropanoid biosynthesis (Dong & Lin, 2021) in plant adaptation to variable

water availability.

From Transient Signals to Lifetime Performance

Traditionally, operational retrograde signaling has been studied under controlled conditions, revealing regulation of oxidative stress genes over seconds to hours (Chan *et al.*, 2010; Crawford *et al.*, 2018). However, chloroplasts continuously sense environmental conditions, and constitutive alterations to retrograde signals lead to sustained developmental changes, as in the case of β-cyclocitral (Jiang & Dehesh, 2021; Sierra *et al.*, 2022). This raises a fundamental question: can transient retrograde signals that operate on minutes-to-hours timescales accumulate to influence plant performance across an entire growing season? If so, primed retrograde responsiveness should manifest as altered performance differences across diverse field environments, indicating integration of multiple environmental fluctuations, rather than a forgotten response to single acute events.

Critically, the fact that both homeologous groups show altered stability relative to Chara - TaSAL2 beneficial and TaSAL1 detrimental - demonstrates that priming the responsiveness of retrograde signaling has successfully altered environmental responses across the entire growing season (Table 3). This interpretation is further supported by our Factor Analytic model, where FA1 accounted for 90% of the variance across environments, strongly suggesting that the specific TaSAL gene modifications are fundamental to explaining the differential environmental responses observed. This differential environmental responsiveness across multiple field seasons represents the strongest evidence that these lines are truly primed for enhanced retrograde signaling, with the direction of the stability change determined by the specific functional role of each homeologous group. The contrasting stability profiles demonstrate that targeted modification of specific TaSAL genes has altered how plants integrate and respond to environmental information across their lifecycle, validating our hypothesis that retrograde signaling effects accumulate beyond transient stress responses.

Conclusions

Modulating retrograde signaling in field environments results in locus-specific differences in biomass, yield, water productivity, and yield stability. The TaSAL1 locus improves resilience at the expense of yield, whereas the TaSAL2 locus results in better resilience, 4-8% gains in yield and thus improved yield stability. Therefore, improving the dynamic nature of retrograde signaling, such as the priming of the SAL-PAP pathway herein, can enhance responsiveness to dynamic environments by maintaining photosynthesis whilst acclimating to transient stress and water deficits. Critically, we show that retrograde signals integrate environmental information throughout the plant lifecycle, not just during acute stress episodes. This precision approach to pathway modulation demonstrates that selective gene targeting can develop climate-adapted crops without yield penalties.

Acknowledgements

The authors acknowledge and thank Dr Zhongyi Li, formerly of CSIRO Agriculture & Food, for access to the HIB population and for his advice. We also thank Dr Hilary Stuart-Williams of the Farquhar Laboratory and ANU's Stable Isotope Laboratory for assistance and guidance of δ^{13}C isotope discrimination analysis. We thank Professor Owen Atkin for his helpful discussion and encouragement. This work was supported by the Australian Grains Research and Development Corporation (GRDC) through grant ANU00020. AFB, MM and BJP were supported by an Australian Research Council Australian Laureate Fellowship (FL190100056). BJP, KXC and ZP are supported by the Australian Research Council Training Centre for Accelerated Future Crops Development (IC210100047).

Financial disclosure

None reported.

Competing Interests

None Declared

Author Contributions

AY, GJR, GME and BJP conceived the experiment(s), MM, AY, JZ, MDM, ZP, EET,

EKA, CAH and DPC conducted the experiment(s), AFB, MM and BJP analysed the

results. AFB, MM, KXC and BJP wrote and reviewed the manuscript. BJP and AY share

corresponding authorship. AFB, MM and AY contributed equally to this work.

Data Availability

The data that support the findings of this study are available in the Supporting

Information of this article. Genotyping data are provided in Supporting Information

Dataset 1, field trial environmental characteristics are detailed in Supporting Information

Tables S3-S7, and comprehensive yield, biomass and water productivity data across all

trials are presented in Supporting Information Table S8. Additional raw phenotypic

measurements are available from the corresponding authors upon request for research

purposes.

Abdallah NA, Elsharawy H, Abulela HA, Thilmony R, Abdelhadi AA, Elarabi NI. **2022**. Multiplex CRISPR/Cas9-mediated genome editing to address drought tolerance in wheat. *GM Crops & Food* **16**: 1–17.

Adjei MO, Luo J, Li X, Du J, Luan A, Li S, Ma J. **2023**. Function of ALA content in porphyrin metabolism regulation of ananas comosus var. bracteatus. *International Journal of Molecular Sciences* **24**.

Ashykhmina N, Lorenz M, Frerigmann H, Koprivova A, Hofsetz E, Stührwohldt N, Flügge U-I, Haferkamp I, Kopriva S, Gigolashvili T. **2019**. PAPST2 Plays Critical Roles in Removing the Stress Signaling Molecule 3′-Phosphoadenosine 5′-Phosphate from the Cytosol and Its Subsequent Degradation in Plastids and Mitochondria. *The Plant Cell* **31**: 231–249.

Bates D, Mächler M, Bolker B, Walker S. **2015**. Fitting Linear Mixed-Effects Models Using lme4. *Journal of Statistical Software* **67**: 1–48.

Blum A. **1998**. Improving wheat grain filling under stress by stem reserve mobilisation. *Euphytica* **100**: 77–83.

Bowerman AF, Byrt CS, Roy SJ, Whitney SM, Mortimer JC, Ankeny RA, Gilliham M, Zhang D, Millar AA, Rebetzke GJ, *et al.* 2023. Potential abiotic stress targets for modern genetic manipulation. *The Plant Cell* **35**: 139–161.

Bowne JB, Erwin TA, Juttner J, Schnurbusch T, Langridge P, Bacic A, Roessner U. **2012**. Drought Responses of Leaf Tissues from Wheat Cultivars of Differing Drought Tolerance at the Metabolite Level. *Molecular Plant* **5**: 418–429.

Butler D. **2022**. asreml: Fits the Linear Mixed Model.

Calderon RH, Strand Å. **2021**. How retrograde signaling is intertwined with the evolution of photosynthetic eukaryotes. *Current Opinion in Plant Biology* **63**: 102093.

Chan KX, Crisp PA, Estavillo GM, Pogson BJ. **2010**. Chloroplast-to-nucleus communication. *Plant Signaling & Behavior* **5**: 1575–1582.

Chan KX, Mabbitt PD, Phua SY, Mueller JW, Nisar N, Gigolashvili T, Stroeher E,
Grassl J, Arlt W, Estavillo GM, *et al.* **2016a**. Sensing and signaling of oxidative stress
in chloroplasts by inactivation of the SAL1 phosphoadenosine phosphatase.
Proceedings of the National Academy of Sciences **113**: E4567–E4576.

Chan KX, Phua SY, Crisp P, McQuinn R, Pogson BJ. **2016b**. Learning the
Languages of the Chloroplast: Retrograde Signaling and Beyond. *Annual Review of*
Plant Biology **67**: 25–53.

Chidzanga C, Fleury D, Baumann U, Mullan D, Watanabe S, Kalambettu P, Pontre
R, Edwards J, Forrest K, Wong D, *et al.* **2021**. Development of an Australian Bread
Wheat Nested Association Mapping Population, a New Genetic Diversity Resource for
Breeding under Dry and Hot Climates. *International Journal of Molecular Sciences* **22**:
4348.

Crawford T, Lehotai N, Strand Å. **2018**. The role of retrograde signals during plant
stress responses. *Journal of Experimental Botany* **69**: 2783–2795.

Crisp PA, Smith AB, Ganguly DR, Murray KD, Eichten SR, Millar AA, Pogson BJ.
2018. RNA Polymerase II Read-Through Promotes Expression of Neighboring Genes in
SAL1-PAP-XRN Retrograde Signaling. *Plant Physiology* **178**: 1614–1630.

Devos KM, Dubcovsky J, Dvořák J, Chinoy CN, Gale MD. **1995**. Structural evolution
of wheat chromosomes 4A, 5A, and 7B and its impact on recombination. *Theoretical*
and Applied Genetics **91**: 282–288.

Dietz K-J, Zörb C, Geilfus C-M. **2021**. Drought and crop yield. *Plant Biology* **23**: 881–
893.

Dong NQ, Lin HX. **2021**. Contribution of phenylpropanoid metabolism to plant
development and plant–environment interactions. *Journal of Integrative Plant Biology*
63: 180–209.

Dwivedi SL, Reynolds MP, Ortiz R. **2021**. Mitigating tradeoffs in plant breeding.
iScience **24**: 102965.

Estavillo GM, Crisp PA, Pornsiriwong W, Wirtz M, Collinge D, Carrie C, Giraud E,
Whelan J, David P, Javot H, *et al.* 2011. Evidence for a SAL1-PAP Chloroplast
Retrograde Pathway That Functions in Drought and High Light Signaling in Arabidopsis.
Plant Cell **23**: 3992–4012.

Farquhar GD, Ehleringer JR, Hubick KT. **1989**. Carbon Isotope Discrimination and
Photosynthesis. *Annual Review of Plant Physiology and Plant Molecular Biology* **40**:
503–537.

Furbank RT, Silva-Perez V, Evans JR, Condon AG, Estavillo GM, He W, Newman
S, Poiré R, Hall A, He Z. **2021**. Wheat physiology predictor: predicting physiological
traits in wheat from hyperspectral reflectance measurements using deep learning. *Plant*
Methods **17**: 108.

Ge SX, Jung D, Yao R. **2020**. ShinyGO: a graphical gene-set enrichment tool for
animals and plants (A Valencia, Ed.). *Bioinformatics* **36**: 2628–2629.

Gezan SA, Galli G, Murray D. **2022**. ASRtriala: An R package with complementary
functions for single and multi-environment trial analyses.

Gigolashvili T, Geier M, Ashykhmina N, Frerigmann H, Wulfert S, Krueger S,
Mugford SG, Kopriva S, Haferkamp I, Flügge U-I. **2012**. The *Arabidopsis* Thylakoid
ADP/ATP Carrier TAAC Has an Additional Role in Supplying Plastidic
Phosphoadenosine 5′-Phosphosulfate to the Cytosol. *The Plant Cell* **24**: 4187–4204.

Hirsch J, Misson J, Crisp PA, David P, Bayle V, Estavillo GM, Javot H, Chiarenza
S, Mallory AC, Maizel A, *et al.* 2011. A Novel fry1 Allele Reveals the Existence of a
Mutant Phenotype Unrelated to 5′->3′ Exoribonuclease (XRN) Activities in Arabidopsis
thaliana Roots (E Newbigin, Ed.). *PLoS ONE* **6**: e16724.

Jackson LE. **1995**. Root architecture in cultivated and wild lettuce (*Lactuca* spp.). *Plant,*
Cell & Environment **18**: 885–894.

Jan MS, Liu ZX, Rochaix JD, Sun XW. **2022**. Retrograde and anterograde signaling in
the crosstalk between chloroplast and nucleus. *Frontiers in Plant Science* **13**: 24.

Jiang J, Dehesh K. **2021**. Plastidial retrograde modulation of light and hormonal
signaling: an odyssey. *New Phytologist* **230**: 931–937.

Joshi V, Joung JG, Fei Z, Jander G. **2010**. Interdependence of threonine, methionine
and isoleucine metabolism in plants: Accumulation and transcriptional regulation under
abiotic stress. *Amino Acids 2010 39:4* **39**: 933–947.

Lin CS, Binns MR. **1988**. A Superiority Measure of Cultivar Performance for Cultivar ×
Location Data. *Canadian Journal of Plant Science* **68**: 193–198.

**Loudya N, Mishra P, Takahagi K, Uehara-Yamaguchi Y, Inoue K, Bogre L, Mochida
K, López-Juez E**. **2021**. Cellular and transcriptomic analyses reveal two-staged
chloroplast biogenesis underpinning photosynthesis build-up in the wheat leaf. *Genome
Biology* **22**: 1–30.

Manmathan H, Shaner D, Snelling J, Tisserat N, Lapitan N. **2013**. Virus-induced
gene silencing of Arabidopsis thaliana gene homologues in wheat identifies genes
conferring improved drought tolerance. *Journal of Experimental Botany* **64**: 1381–1392.

Mohr T, Horstman J, Gu YQ, Elarabi NI, Abdallah NA, Thilmony R. **2022**. CRISPR-
Cas9 Gene Editing of the Sal1 Gene Family in Wheat. *Plants* **11**: 2259.

Passioura JB. **2006**. Viewpoint: The perils of pot experiments. *Functional Plant Biology*
33: 1075.

Phua SY, Pornsiriwong W, Chan KX, Estavillo GM, Pogson BJ. **2018a**.
Development of strategies for genetic manipulation and fine-tuning of a chloroplast
retrograde signal 3′-phosphoadenosine 5′-phosphate. *Plant Direct* **2**: 15.

Phua SY, Yan DW, Chan KX, Estavillo GM, Nambara E, Pogson BJ. **2018b**. The
Arabidopsis SAL1-PAP Pathway: A Case Study for Integrating Chloroplast Retrograde,
Light and Hormonal Signaling in Modulating Plant Growth and Development? *Frontiers
in Plant Science* **9**: 8.

**Pires MV, Júnior AAP, Medeiros DB, Daloso DM, Pham PA, Barros KA, Engqvist
MKM, Florian A, Krahnert I, Maurino VG, *et al.*** **2016**. The influence of alternative

pathways of respiration that utilize branched-chain amino acids following water shortage
in arabidopsis. *Plant, Cell & Environment* **39**: 1304–1319.

Pogson BJ, Woo NS, Förster B, Small ID. **2008**. Plastid signalling to the nucleus and
beyond. *Trends in Plant Science* **13**: 602–609.

Pornsiriwong W, Estavillo GM, Chan KX, Tee EE, Ganguly D, Crisp PA, Phua SY,
Zhao C, Qiu J, Park J, *et al.* 2017. A chloroplast retrograde signal, 3'-
phosphoadenosine 5'-phosphate, acts as a secondary messenger in abscisic acid
signaling in stomatal closure and germination. *eLife* **6**: 34.

R Core Team. **2021**. R: A Language and Environment for Statistical Computing.

Rebetzke GJ, Chenu K, Biddulph B, Moeller C, Deery DM, Rattey AR, Bennett D,
Barrett-Lennard EG, Mayer JE. **2013**. A multisite managed environment facility for
targeted trait and germplasm phenotyping. *Functional Plant Biology* **40**: 1.

Rebetzke GJ, Van Herwaarden AF, Jenkins C, Weiss M, Lewis D, Ruuska S, Tabe
L, Fettell NA, Richards RA. **2008**. Quantitative trait loci for water-soluble
carbohydrates and associations with agronomic traits in wheat. *Australian Journal of*
Agricultural Research **59**: 891–905.

Regina A, Berbezy P, Kosar-Hashemi B, Li S, Cmiel M, Larroque O, Bird AR,
Swain SM, Cavanagh C, Jobling SA, *et al.* 2015. A genetic strategy generating wheat
with very high amylose content. *Plant Biotechnology Journal* **13**: 1276–1286.

Rossel JB, Walter PB, Hendrickson L, Chow WS, Poole A, Mullineaux PM, Pogson
BJ. **2006**. A mutation affecting *ASCORBATE PEROXIDASE 2* gene expression reveals
a link between responses to high light and drought tolerance. *Plant, Cell & Environment*
29: 269–281.

Sharma DL, Bhoite R, Reeves K, Forrest K, Smith R, Dowla MANNU. **2022**.
Genome-wide superior alleles, haplotypes and candidate genes associated with
tolerance on sodic-dispersive soils in wheat (*Triticum aestivum* L.). *Theoretical and*
Applied Genetics **135**: 1113–1128.

Sierra J, McQuinn RP, Leon P. **2022**. The role of carotenoids as a source of
retrograde signals: impact on plant development and stress responses. *Journal of*
Experimental Botany **73**: 7139–7154.

Smith A, Cullis B, Gilmour A. **2001**. Applications: The Analysis of Crop Variety
Evaluation Data in Australia. *Australian & New Zealand Journal of Statistics* **43**: 129–
145.

Soprano AS, Smetana JHC, Benedetti CE. **2018**. Regulation of tRNA biogenesis in
plants and its link to plant growth and response to pathogens. *Biochimica et Biophysica*
Acta (BBA) - Gene Regulatory Mechanisms **1861**: 344–353.

Tee EE, Fairweather SJ, Vo HM, Zhao C, Breakspear A, Kimura S, Carmody M,
Wrzaczek M, Bröer S, Faulkner C, *et al.* 2025. SAL1-PAP retrograde signaling
orchestrates photosynthetic and extracellular reactive oxygen species for stress
responses. *The Plant Journal* **122**: e70271.

Wilson PB, Estavillo GM, Field KJ, Pornsiriwong W, Carroll AJ, Howell KA, Woo
NS, Lake JA, Smith SM, Harvey Millar A, *et al.* 2009. The nucleotidase/phosphatase
SAL1 is a negative regulator of drought tolerance in Arabidopsis. *The Plant Journal* **58**:
299–317.

Yadav AK, Carroll AJ, Estavillo GM, Rebetzke GJ, Pogson BJ. **2019**. Wheat drought
tolerance in the field is predicted by amino acid responses to glasshouse-imposed
drought. *Journal of Experimental Botany* **70**: 4931–4948.

Zandalinas SI, Mittler R. **2022**. Plant responses to multifactorial stress combination.
New Phytologist **234**: 1161–1167.

Zhang H, Zhang F, Yu Y, Feng L, Jia J, Liu B, Li B, Guo H, Zhai J. **2020**. A
Comprehensive Online Database for Exploring ~20,000 Public *Arabidopsis* RNA-Seq
Libraries. *Molecular Plant* **13**: 1231–1233.

Zhao C, Wang Y, Chan KX, Marchant DB, Franks PJ, Randall D, Tee EE, Chen G,
Ramesh S, Phua SY, *et al.* 2019. Evolution of chloroplast retrograde signaling

facilitates green plant adaptation to land. *Proceedings of the National Academy of*

Sciences **116**: 5015–5020.

Supporting information

Additional supporting information may be found in the online version of the article at the publisher's website.

Dataset 1 Genotype-by-sequencing results for all mutants in study.

Fig. S1 Comparison of variation in wheat SAL gene expression across developing seedling leaves.

Fig. S2 Representative agarose gel of screening for TaSAL mutants via PCR.

Fig. S3 PAP content under irrigated and drought conditions in glasshouse experiments.

Fig. S4 Plant height and relative water content under drought.

Fig. S5 Soil weight changes during drought

Fig. S6. Photosynthesis and stomatal conductance measures.

Fig. S7 ABA content of glasshouse-grown wheat material under Control and Drought conditions.

Fig. S8 Leaf/canopy temperature results from Narrabri 2019.

Fig. S9 Leaf/canopy temperature results from Narrabri 2017.

Fig. S10 Hyperspectral-derived estimates of photosynthesis related traits.

Fig. S11 Factor analytic model visualization of multi-environment trial analysis - Biplot and Genetic Covariance Matrix heatmap.

Table S1 Table of primers used in discovery of *TaSal* null lines from HIB population.

Table S2 Line identifiers and *TaSal* gene knockout.

Table S3 Narrabri environmental characteristics for all years of trials.

Table S4 Condobolin environmental characteristics for all years of trials.

Table S5 Yanco environmental characteristics for all years of trials.

Table S6 Birchip environmental characteristics for all years of trials.

Table S7 Characterising average and actual water conditions per site and year,

including irrigation where used.

Table S8 Yield, Biomass and Water Productivity Best Linear Unbiased Predictions for

all trials.

Method S1 Phylogenetic analysis.

Method S2 PAP extraction, preparation, and quantification by HPLC.

Figure Captions

Figure 1. Wheat SAL gene diversity. (a) Phylogeny of Liliopsida SAL sequences.
Black diamonds indicate bootstrap support >95% based on 1000 ultrafast bootstrap
replicates. Purple: TaSAL1 (chr 5), green: TaSAL2 (chr 4A/7A/7D). (b) Expression of
wheat SAL genes. (c) GO enrichment for TaSAL groups. Significance = $-\log_{10}(p$-value).

Figure 2. TaSAL activity (a) and relative water content (b) under drought stress.
(a) SAL activity after 21d drought. Mean ± SD. Two-way ANOVA with FDR correction.
(b) Relative water content after 21d drought. Mean ± Half Least Significant Difference
(HLSD) from ASREML-R repeated measures model. *$p<0.05$, **$p<0.01$, ***$p<0.001$.

Figure 3. Photosynthesis vs stomatal conductance under water stress.
Relationship between photosynthetic assimilation and stomatal conductance under
watered control (Day 28) and Terminal Drought conditions (Days 28, 35). TaSAL
mutants maintain higher assimilation rates despite similar conductance reductions
compared to Chara. Mixed effects analysis. Ellipses show SE.

Figure 4. Trial environmental characteristics. (a) Trial locations across southeastern
Australia. (b) Average rainfall patterns: (i) monthly averages and (ii) annual totals by
site. (c) Actual rainfall variation: (i) monthly rainfall by trial and (ii) total water availability
per trial. (d-f) Temperature patterns: (i) average monthly temperatures and (ii) actual
temperature ranges across trials, showing (d) overall temperature patterns, (e)
maximum temperatures, and (f) minimum temperatures.

Figure 5. Canopy temperature, Narrabri 2017. Diurnal canopy temperature patterns
measured using UAV-based infrared cameras under rainfed and water-supplemented
conditions. TaSAL mutants showed increased midday temperatures vs Chara,
indicating enhanced stomatal closure during peak evapotranspiration demand.
Multivariate repeated measures linear mixed effects model. Values are BLUEs ± SE. *p
< 0.05, **$p < 0.01$, ***$p < 0.001$.

Figure 6. Field trial performance of TaSAL mutants. BLUEs (Best Linear Unbiased
Estimators) for yield, biomass, and water productivity across trials. TaSAL2 lines
outperformed Chara; TaSAL1 lines showed reduced performance.

Tables

Table 1: Carbon isotope discrimination (ΔC) in glasshouse conditions.

Line	ΔC Control	ΔC Drought
Chara	20.59 ± 0.12	18.43 ± 0.18
SAL2.1	18.77 ± 0.17***	16.39 ± 0.16***
SAL2.2	20.57 ± 0.84	17.78 ± 0.16
SAL1.1	20.02 ± 0.39	18.28 ± 0.51
SAL1.2	19.83 ± 0.31*	17.08 ± 0.25*
SAL1.1×SAL2.1	19.29 ± 0.28***	16.43 ± 0.22***

966 Mean ± SE, n=3 per Line/Condition. Two-way ANOVA with post-hoc comparisons to
967 Chara control. *p<0.05, **p<0.01, ***p<0.001 vs Chara.

969 **Table 2: Yield, Biomass and Water Productivity Best Linear Unbiased Predictions**
970 **(BLUPs)**

Genotype	Yield (t/Ha)	Biomass (t/Ha)	Water Productivity (kg/Ha/mm Water)
Chara	4.70 ± 0.03	9.41 ± 0.11	14.20 ± 0.10
SAL2.2	4.89 ± 0.03***	9.38 ± 0.11	14.34 ± 0.10***
SAL2.3	5.09 ± 0.20†	9.83 ± 0.46	14.29 ± 0.51†
SAL1.1	4.46 ± 0.03***	9.69 ± 0.11	13.28 ± 0.10***
SAL1.2	4.57 ± 0.03***	9.78 ± 0.11*	13.83 ± 0.10***
SAL1.1×SAL2.1	3.85 ± 0.03***	8.02 ± 0.14***	11.68 ± 0.10***

971 **Notes:** Mean values of random effects ± SE. *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$ vs
972 Chara. †Not significant in MET model, but significant in trials present.

974 **Table 3: Cultivar superiority (P_i) estimates for yield.**

Entry	Stability (P_i)
SAL2.3	0.272
SAL2.2	0.488
Chara	0.706
SAL1.2	0.927
SAL1.1	1.087
SAL1.1×SAL2.1	2.362

975 Main lines investigated only included. Smaller values indicate greater stability of yield
976 across environments.

Canopy Temperature