1 Utility of a quantitative approach to microbial dysbiosis using machine learning in an
2 African American cohort with self-reported hair loss
3

4 Judy Malas[1], Sebastian Reczek[1], Imani Porter[2,3], Yolanda M. Lenzy[4], Victoria Barbosa[5],
5 Jack A. Gilbert[6], Crystal Porter[3], Jarrad Hampton-Marcell[1]
6
[1]Department of Biological Sciences, University of Illinois Chicago, Chicago, IL
[2]Department of Genetics, Stanford University, Stanford, CA,
[3]Mane Insights, Naperville, IL
[4]Lenzy Dermatology, Chicopee, MA
[5]Department of Medicine, University of Chicago, Chicago, IL
[6]Department of Pediatrics and Scripps Institution of Oceanography, University of
California San Diego, La Jolla, CA
14
Corresponding author:
Jarrad Hampton-Marcell, PhD
Department of Biological Sciences
University of Illinois at Chicago
Telephone: 708-368-6533
Email: marcell2@uic.edu
21
22

23 **Conflicts of Interest**: None declared.
24

25 **IRB approval status:** Reviewed and approved by The University of Chicago Biological
26 Sciences Division institutional review board; #IRB16-1541.
27
28 **Keywords:** dysbiosis; hair loss; alopecia; machine learning; microbiome; sequencing;
29 cohort

30
31
32

Abstract

Microbial dysbiosis has been identified as a therapeutic target for several dermatologic conditions. However, the concept of dysbiosis is poorly defined, limiting reproducibility. We developed a reproducible metric for dysbiosis that can be applied to research within dermatology. Thirty-six individuals from an African-American cohort with self-reported hair loss without formal clinical diagnoses provided scalp swabs from both afflicted (hair loss) and normal (no hair loss) sites. The scalp microbiome was characterized via 16S rRNA gene sequencing. A dysbiosis score that considers the proportion of all taxa within the samples was calculated. Further, we identified the taxa most associated with dysbiosis using both a machine learning random forest classifier and a negative binomial mixed effects model to control for participant age. Sites of hair loss exhibited an increase in microbial diversity, in particular for individuals older than 40 years. We identified a core set of OTUs assigned to 7 genera that were significant contributors to increased scalp dysbiosis. This work demonstrates the utility of applying a quantitative approach to dysbiosis and provides a framework that can be applied to other microbiome-associated conditions.

Introduction. Recent research has revealed that changes in the microbiome are associated with several dermatologic conditions including different types of alopecia [1], dandruff, seborrheic dermatitis [2], atopic dermatitis [3], scalp psoriasis, and folliculitis decalvans [4]. The term "dysbiosis" is often invoked when describing microbial associations with disease. However, in most studies the definition of "dysbiosis" lacks clarity; its use ranges from a broad shift in the microbial community composition to a change in a specific taxon [5]. Dysbiosis is readily applied to the gut microbiome and therefore is frequently equated with decreased microbial diversity [5]. This conflation with decreased diversity may lead to confusion, as increased diversity is associated with aging and disease in dermatologic conditions [6, 7]. With treatments for skin disorders targeting microbial "dysbiosis" on the rise [8-10], there is a need for a more quantitative approach for the concept of dysbiosis.

We built a reproducible microbial dysbiosis score and identified the taxa differentiating dysbiotic from non-dysbiotic scalp sites, without *a priori* selection of the organisms associated with dysbiosis. This work involved a cohort of thirty-six African-American individuals with self-reported hair loss. We posited that skin sites exhibiting hair loss would show significant shifts in microbiome composition, indicative of dysbiosis development. Individuals provided both afflicted (hair loss) and normal (no hair loss) scalp swabs and the bacterial 16S rRNA gene was sequenced.

Methods

Recruiting and Sample Collection. Adults with self-reported hair loss were recruited for the study without restrictions in ethnic background. Exclusion criteria included participants with complete hair loss, the presence of open sores, drainage and/or injury to the scalp, and cancer or chemotherapy treatment. Participants were instructed to use a sterile swab (Puritan catalog # BD 25-806 10WC) to collect two samples: one from an afflicted (hair-loss) site and one from a normal site, swabbing the areas for 10 seconds each to collect microbial biomass. All qualifying participants completed a hair density scale; a questionnaire where participants rated their degree of hair loss using an anchored line scale from 1 to 10. An index score of 1 represented a participant whose scalp was readily viewable, and 10 represented a scalp completely covered by hair fibers [11, 12]. Participants also completed a hair stress scale, which contained anchored line scales where they indicated their level of stress on a scale from 1 – 10 [13]. Study protocols were approved and overseen by The University of Chicago Biological Sciences Division institutional review board, Chicago, Illinois (protocol number IRB16-1541). A signed consent was obtained from all participants prior to sampling and questionnaire completion.

Library preparation and sequencing. Swabbed samples were placed inside of a sterile Whirl Pak® bag and mailed to the testing facility. Upon arrival, samples were stored in a -80°C freezer until DNA extraction. Genomic DNA was extracted using the recommended protocols of the Earth Microbiome Project (EMP) and American Gut Project [14, 15]. The V4 region of the 16S rRNA gene (515F-806R) was amplified. Barcoded primer sets were adapted for the Illumina MiSeq System as described by [16].

Statistical analysis. All statistical analyses were performed using R v4.4.2 and
visualized with the package 'ggplot2' v 3.5.1 [17]. Sequences were quality-filtered using
QIIME v1.9.1 [18]. Operational taxonomic units (OTUs) were defined and assigned
taxonomy by clustering sequences against 97% OTU reference data set from the
Greengenes database [19, 20]. Alpha diversity measures were generated using
'microbiome' package v1.28.0 [21]. Pearson correlations were calculated to determine
the linear relationship between age and alpha diversity. Age groups were further split
into 20-year intervals (< 40, 40-59, 60+) and a Wilcoxon rank-sum test was conducted
to evaluate differences in alpha diversity between age groups.

Dysbiosis score. Dysbiosis scores were calculated using the R package 'dysbiosisR'
v1.0.4 (https://github.com/microsud/dysbiosisR). Our dysbiosis score is a modification of
the score reported in [22]. The microbial dysbiosis score is defined as the difference
between (the Aitchison distance between the test sample and the healthy class
centroid) and (the Atichison distance between the test sample and the diseased class
centroid). In other words, the dysbiosis score for each sample measures its closeness
to the group (hair-loss afflicted or normal) mean. The dysbiosis score **D** of the test
sample **z**, is defined as

$$D(z;\ \mu C_N, \mu C_A) = d_A(z,\ \mu C_N) - d_A(z,\ \mu C_A)$$

where μC_N and μC_A are the centroids (mean composition) of the normal and afflicted
samples, respectively and d_A is the Aitchison distance either between the test sample
and the μC_N or μC_A. A dysbiosis score of 0 indicates that the test sample is at equal
distance from the center of both class (afflicted and normal) centroid, and higher scores
indicate more deviation from normal. Calculated dysbiosis scores were classified into
two groups of either dysbiosis (score > 0) or normobiosis (score < 0). Code used to
calculate dysbiosis scores along with all microbial analyses conducted is publicly
available (https://github.com/jud-m/Defining-microbial-dysbiosis).

Determination of taxa contributing to dysbiosis. To determine the bacterial taxa that
differentiated dysbiotic hair sites from the normal hair sites, a random forest (RF)
machine learning classifier was used to determine which data features (OTUs) were
most important for classifying dysbiotic versus normobiosis samples using the R
package 'randomForest' v4.7-1.2. An out-of-bag random forest model was used to
assess the importance of each microbial OTU by removing each OTU in turn and
observing its effect on the classification accuracy. The100 most important OTUs (out of
6,009 total OTUs) for classification were considered further. In addition, because age
can be a confounding variable in determining the microbial taxa associated with
dysbiosis, a negative binomial mixed-effect model (NBMM) was used to model the
microbiome count data using the R package 'NBZIMM' v1.0 [23]. The NBMM identified
differentially abundant taxa in the afflicted hair sites relative to normal sites while
controlling for the effect of age. As a final check, a Wilcoxon rank-sum test was
conducted on the log10 abundance of the OTUs that were flagged in both the NBMM
and the random forest classifier. OTUs that were not differentially abundant according to
the Wilcoxon rank-sum test were excluded.

Results
150
Cohort characteristics. The majority of recruited participants were African American
(36 of 38) and the remaining two participants were Caucasian. Caucasian participants
were excluded from further analyses as microbiome differences are associated with
ethnicity [24], and there were not enough Caucasian participants to control for this
confounding variable. The average age of the participants was 51.6 years (sd +/- 13.5).
The cohort included twenty-nine female and seven male participants. The state of the
hair among participants included: natural (n = 20), relaxed (n = 8), color (n = 5), keratin
treatment (n = 1), and other (n = 2). None of the participants wore hair prostheses. Both
scalp density (Pearson, p = 0.0019, R^2 = -0.51) and scalp stress (Pearson, p = 0.073,
R^2 = -0.31) decreased with age but only the correlation with scalp density was
statistically significant. The frequency of scalp cleansing reported by participants
included monthly (n = 7), biweekly (n = 16), weekly (n = 12), and daily (n = 1).
163
Microbial diversity. Following quality filtering,1,720,130 microbial sequences
comprising 6,009 OTUs were observed for 67 samples (35 afflicted, 32 normal).
Firmicutes (50.6%), Actinobacteria (33%) and Proteobacteria (11.6%) were the most
abundant phyla. *Staphylococcus, Corynebacterium*, and *Propionibacterium* were the
most abundant genera (**Figure S1**). Shannon alpha diversity, a measure of richness
and evenness of species within a sample, was significantly higher in the afflicted hair
sites (p = 0.004). Shannon alpha diversity was positively correlated with age in both
afflicted and normal hair sites (R^2 = 0.62 and 0.4, respectively; **Fig. 1A**). When
categorized by 20-year intervals, participants above 40 years of age have significantly
higher Shannon diversity values in the afflicted sites (**Fig. 1B**). Neither hair state
(including treatments) nor cleansing frequency were significantly associated with alpha
diversity of the afflicted hair sites.
176
Dysbiosis. Individual samples were assigned a dysbiosis score **(Fig. 2)**. Calculated
dysbiosis scores were classified into groups of either dysbiosis (score > 0) or
normobiosis (score < 0). It was determined that the majority, but not all, sites of hair loss
are dysbiotic (**Fig. 2**). Based on the threshold set, 24 of 35 afflicted samples were
classified as dysbiotic (68.5%) and 30 of 32 samples normal samples as normobiosis
(93.75%).
183
Core taxa contributing to dysbiosis. A RF machine learning classifier was used to
determine the most important OTUs for distinguishing the dysbiotic from normobiosis
samples. The overall error rate for the RF was 20.9%. The model was better at
classifying normobiosis sites; dysbiotic sites were classified to an accuracy of 57.6%,
while the normobiosis sites were correctly classified in 92.6% of samples. Twenty-seven
unique genera were assigned to the 100 most important OTUs for classification (**Fig.
S2**). The mean decrease accuracy (MDA), a measure that indicates the percentage
decrease in the random forest model accuracy when a specific OTU is excluded, was
>10% for OTUs assigned to the genera *Streptococcus* (66%), *Corynebacterium*
(34.2%), *Anaerococcus* (15.8%), *Haemophilus* (14.3%), *Peptoniphilus* (12.5%),
Staphylococcus (11.9%), *Actinomyces* (11.4%), and *Rothia* (10.5%) (**Fig. S2**).

To control for the potential confounding effect of age, a NBMM was generated
each of the 6009 OTUs were independently tested for differential abundance. A total of
100 OTUs belonging to 29 unique genera were found to be differentially abundant in the
afflicted sites compared to the normal sites, the majority of which (83%) were more
abundant in the afflicted sites (**Fig S3)**. OTUs associated with dysbiosis from the RF
were compared to the differentially abundant genera from the NBMM (**Fig. S4**). Taxa
that were not identified as differentially abundant by the NBMM were excluded. After
conducting a Wilcoxon-rank-sum test to confirm differential abundance, 20 OTUs
assigned to 7 genera were determined to be significantly associated with dysbiosis
while controlling for age (**Fig 3**).
205
206 **Discussion**
207
We have developed a microbial dysbiosis score as a proxy for scalp health. Sparse
areas on the scalp are associated with dysbiosis in individuals experiencing self-
reported hair loss. In addition, we determined a core set OTUs belonging to 7 genera
that are significantly more abundant in dysbiotic sites.
Dysbiosis has been broadly defined as a condition differing from the healthy or
normal state [25]. The taxonomic composition of the skin microbiome is a result of a
multitude of factors including host physiology, environment, lifestyle, and underlying
pathobiology [26, 27]. In addition, the microbiome is affected by factors such as skin
thickness, folds, and the density/type of hair follicles and glands [24, 27, 28]. Therefore,
"normal" is subject to interindividual variation, rendering an informal definition of
dysbiosis ineffective. Here, we have defined dysbiosis quantitatively. This approach
allows for a gradient approach to characterizing dysbiosis so researchers can quantify
the degree to which a particular sample differs from normal and set different thresholds
depending on study goals.
Bacterial genera commonly identified on healthy human scalp are
Propionibacterium (P. acnes), Staphylococcus (S. epidermidis), and *Corynebacterium.*
Relative changes in the abundance of these bacteria have been implicated in dysbiosis,
though there are inconsistencies between studies (Table 1). Discrepancies may reflect
an environmental, disease state, or ethnicity-specific response; most reported studies,
including this one, focus on a particular population of subjects (Table 1). In addition,
studies often aggregate data at the genus level, though different species within a genus
can be either beneficial or antagonistic to human health. Therefore, considering the
abundance of the entire genus may be too coarse of a metric. Here, we evaluated
abundance differences at the OTU level, which allowed for the differentiation of OTUs
within a genus that expressed different patterns of association. In this study, taxa of the
genera *Corynebacterium* and *Staphylococcus* were less abundant in hair loss sites
when evaluated in aggregate (**Fig S1**)*,* however, some OTUs within these genera were
more abundant in hair loss sites (**Fig S3)**. *Corynebacterium* increases and decreases
have been associated with scalp-related conditions (Table 1); here, only a few
Corynebacterium OTUs contributed to dysbiosis (**Fig 3**), potentially accounting for the
differences between studies. Therefore, we recommend that future work evaluate
taxonomic abundances without aggregating sequence data at the genus level.

Our results indicate that an increase in less abundant residents of the scalp, in
addition to an increase of certain OTUs within the genus *Corynebacterium,* is
associated with the symptom of hair loss. Taxa within the genera *Acinetobacter,*
Anaerococcus, Corynebacterium, Finegoldia, Gamella, Haemophilus, Rothia, and
Streptococcus were identified as the most significantly associated with dysbiosis (**Fig.**
3). Several of these genera can be highly pathogenic though the pathogenicity is
species-dependent. The genera *Acinetobacter* and *Anaerococcus* have previously been
identified as more abundant in the scalp dermis of patients with Alopecia Areata [30].
Streptococcus, which is implicated in many infectious diseases including cellulitis and
toxic shock syndrome [34], contributed most to dysbiosis (**Fig. S2; Fig 3**).

While this work does not include formal clinical diagnoses for our subjects, our
findings show that the symptom of hair loss can serve as an indicator of scalp health.
Hair shafts promote the growth of gram-negative biofilms while inhibiting the growth of
gram-positive *Staphylococcus* species [28]. We observed a decrease in *S. epidermidis*
OTUs in the afflicted sites, which could have allowed colonization of other, potentially
harmful species. In addition, antimicrobial proteins (lysozyme and histones) have been
extracted from human hair [35]. Therefore, the loss of hair in the afflicted sites may have
opened the door to enhanced colonization and thus increased diversity due to the loss
of protective biofilms and/or antimicrobial properties of the hair. Microbial diversity
increased significantly at the afflicted sites for those older than 40 years (**Fig. 1**), which
may indicate some degree of protection from colonization of harmful bacteria in younger
individuals. Alternatively, the older individuals in the cohort may have had differing
underlying scalp disorders. For example, androgenetic alopecia is more common in
older individuals, and this condition may elicit a different microbial response compared
to traction alopecia, which is the predominant alopecia type in younger individuals of
African descent [36]. Future work should compare hair loss sites from more individuals
with clinical diagnoses.

CONCLUSIONS

Here, we have corroborated previous work that scalp-related disorders are
associated with increased microbial diversity and have shown that this diversity is
spatially significant on the scalp within individuals and driven in several taxa including
Streptococcus. We have shown that within a genus, taxa can be more or less abundant
depending on the species, and we recommend against aggregating 16S sequence data
at the genus level for differential abundance analyses. Importantly, we have developed
a reproducible dysbiosis score that can be used to inform dermatological diagnoses and
compare research findings meaningfully. The limited research into treatment modalities
that target the scalp microbiome has shown promise [7, 37] and future work should
explore targeting the reduction of dysbiosis-associated taxa.

ACKNOWLEDGEMENTS

We like to acknowledge Argonne's ACT-SO High School Research Program for
providing a platform for Ms. Imani Porter to conduct research at Argonne National
Laboratory. We would also like to acknowledge Sarah Owen and Stephanie Moormann
of the Environmental Sample Preparation and Sequencing Facility at Argonne National
Laboratory for their expertise.

References

1. Suzuki, K., et al. *Scalp Microbiome and Sebum Composition in Japanese Male Individuals with and without Androgenetic Alopecia.* Microorganisms, 2021. **9**, DOI: 10.3390/microorganisms9102132.

2. Saxena, R., et al., *Comparison of Healthy and Dandruff Scalp Microbiome Reveals the Role of Commensals in Scalp Health.* Frontiers in Cellular and Infection Microbiology, 2018. **8**.

3. Seite, S., et al., *Microbiome of affected and unaffected skin of patients with atopic dermatitis before and after emollient treatment.* J Drugs Dermatol, 2014. **13**(11): p. 1365-72.

4. Shah, R.R., et al., *Scalp microbiome: a guide to better understanding scalp diseases and treatments.* Arch Dermatol Res, 2024. **316**(8): p. 495.

5. Hooks, K.B. and M.A. O'Malley, *Dysbiosis and Its Discontents.* mBio, 2017. **8**(5).

6. Luna, P.C., *Skin Microbiome as Years Go By.* American Journal of Clinical Dermatology, 2020. **21**(1): p. 12-17.

7. Saxena, R., et al., *Longitudinal study of the scalp microbiome suggests coconut oil to enrich healthy scalp commensals.* Scientific Reports, 2021. **11**(1): p. 7220.

8. *Autologous skin microbiota transplantation: Development of a technique to transfer representative and viable cutaneous microbial communities.* Journal of the American Academy of Dermatology, 2018. **79**(3): p. AB55.

9. Comito, N., et al., *50550 2-Week Application of Topical Colloidal Oatmeal Gel Cream Supports the Growth of Skin Commensal Staphylococcus Epidermidis in Subjects with Mild-to-Moderate Atopic Dermatitis.* Journal of the American Academy of Dermatology, 2024. **91**(3): p. AB99.

10. *Using a specific emollient to manage skin microbiome dysbiosis.* Journal of the American Academy of Dermatology, 2016. **74**(5): p. AB89.

11. Courtois, M., et al., *Ageing and hair cycles.* British Journal of Dermatology, 1995. **132**(1): p. 86-93.

12. Barman, J.M., I. Astore, and V. Pecoraro, *The Normal Trichogram of the Adult.* Journal of Investigative Dermatology, 1965. **44**(4): p. 233-236.

13. Meilgaard, M., G. Civille, and B. Carr, *Sensory evaluation techniques 4th ed CRC Press LLC.* New York, 2007.

14. McDonald, D., et al., *American Gut: an Open Platform for Citizen Science Microbiome Research.* mSystems, 2018. **3**(3): p. 10.1128/msystems.00031-18.

15. Gilbert, J.A., J.K. Jansson, and R. Knight, *The Earth Microbiome project: successes and aspirations.* BMC Biology, 2014. **12**(1): p. 69.

16. Caporaso, J.G., et al., *Ultra-high-throughput microbial community analysis on the Illumina HiSeq and MiSeq platforms.* The ISME Journal, 2012. **6**(8): p. 1621-1624.

17. Wickham, H., *ggplot2 : Elegant Graphics for Data Analysis*, in *Use R!,*. 2016, Springer International Publishing : Imprint: Springer,: Cham. p. 1 online resource (XVI, 260 pages 232 illustrations, 140 illustrations in color.

18. Caporaso, J.G., et al., *QIIME allows analysis of high-throughput community sequencing data.* Nat Methods, 2010. **7**(5): p. 335-6.

19. McDonald, D., et al., *An improved Greengenes taxonomy with explicit ranks for ecological and evolutionary analyses of bacteria and archaea.* Isme j, 2012. **6**(3): p. 610-8.

20. Edgar, R.C., *Search and clustering orders of magnitude faster than BLAST.* Bioinformatics, 2010. **26**(19): p. 2460-1.

21. Lahti, L. and S. Shetty, *Microbiome package.* 2017, Bioconductor

22. AlShawaqfeh, M.K., et al., *A dysbiosis index to assess microbial changes in fecal samples of dogs with chronic inflammatory enteropathy.* FEMS Microbiology Ecology, 2017. **93**(11): p. fix136.

23. Zhang, X. and N. Yi, *NBZIMM: negative binomial and zero-inflated mixed models, with application to microbiome/metagenomics data analysis.* BMC Bioinformatics, 2020. **21**(1): p. 488.

24. Dimitriu Pedro, A., et al., *New Insights into the Intrinsic and Extrinsic Factors That Shape the Human Skin Microbiome.* mBio, 2019. **10**(4): p. 10.1128/mbio.00839-19.

25. Wei, S., et al., *Determining Gut Microbial Dysbiosis: a Review of Applied Indexes for Assessment of Intestinal Microbiota Imbalances.* Appl Environ Microbiol, 2021. **87**(11).

26. Harris-Tryon, T.A. and E.A. Grice, *Microbiota and maintenance of skin barrier function.* Science, 2022. **376**(6596): p. 940-945.

27. Grice, E.A. and J.A. Segre, *The skin microbiome.* Nature Reviews Microbiology, 2011. **9**(4): p. 244-253.

28. Kerk, S.K., et al., *Bacteria Display Differential Growth and Adhesion Characteristics on Human Hair Shafts.* Frontiers in Microbiology, 2018. **9**.

29. Gallo, R.L. and T. Nakatsuji, *Microbial symbiosis with the innate immune defense system of the skin.* J Invest Dermatol, 2011. **131**(10): p. 1974-80.

30. Pinto, D., et al., *Scalp bacterial shift in Alopecia areata.* PLOS ONE, 2019. **14**(4): p. e0215206.

31. Won, E.J., et al., *A Potential Predictive Role of the Scalp Microbiome Profiling in Patients with Alopecia Areata: Staphylococcus caprae, Corynebacterium, and Cutibacterium Species.* Microorganisms, 2022. **10**(5).

32. Jung, D.-R., et al., *Comparative analysis of scalp and gut microbiome in androgenetic alopecia: A Korean cross-sectional study.* Frontiers in Microbiology, 2022. **13**.

33. Pyle, H.J., et al., *A Pilot Study Characterization of the Scalp Microbiome in Central Centrifugal Cicatricial Alopecia Shows Shift in Corynebacterium.* Skin Appendage Disorders, 2024: p. 1-8.

34. Brouwer, S., et al., *Pathogenesis, epidemiology and control of Group A Streptococcus infection.* Nature Reviews Microbiology, 2023. **21**(7): p. 431-447.

35. Adav, S.S., et al., *Studies on the Proteome of Human Hair - Identification of Histones and Deamidated Keratins.* Scientific Reports, 2018. **8**(1): p. 1599.

36. Khumalo, N.P., et al., *Determinants of marginal traction alopecia in African girls and women.* Journal of the American Academy of Dermatology, 2008. **59**(3): p. 432-438.

386   37.   Rinaldi, F., A. Trink, and D. Pinto, *Efficacy of Postbiotics in a PRP-Like Cosmetic*
387   *Product for the Treatment of Alopecia Area Celsi: A Randomized Double-Blinded*
388   *Parallel-Group Study.* Dermatol Ther (Heidelb), 2020. **10**(3): p. 483-493.

Table 1. Summary of findings from studies on the association between microbial
dysbiosis and scalp-related dermatological conditions.

Ref	Population	Condition	Diversity	*Staphylococcus*	*Corynebacterium*	*Propionibacterium / Cutibacterium*
[7]	Indian women	Dandruff	Increase	Increase in *S. epidermidis*	No change	Decrease in *P. acnes*
[30]	Italian men	Alopecia Areata	Increase	Decrease in Genus Decrease in *S. epidermidis*	No change	Increase in Genus Increase in *P. acnes*
[1]	Japanese men	Androgenetic Alopecia	No change	Increase	Decrease	Increase
[31]	Patients recruited from hospital in Korea	Alopecia Areata	Increase	Decrease in Genus Decrease in *S. carprae*	Increase	Increase in severe cases
[32]	Korean women and men	Androgenetic Alopecia	Increase	Decrease in men only	Increase in men only	No change
[33]	Black women	Central Centrifugal Cicatricial Alopecia	No Change	No change	Increase	No change
This study	African American men and women	Hair loss	Increase	Increase in several OTUs Decrease in *S. epidermidis*	Increase in several OTUs	Decrease in *P. acnes*

Figure 1. A) Linear regression of age of participants with Shannon Alpha diversity of the
scalp microbiome. Red points are the afflicted sites, black points are the normal sites.
B) Shannon alpha diversity of the afflicted and normal hair site. Each panel represents
an age group 1) Less than 40, 2) 40-49 and 3) 60+.

Figure 2. Dysbiosis scores plotted with threshold (white dotted line). A dysbiosis score
of 0 indicates that the test sample is at equal distance from the center of both class
(afflicted and normal) centroid, and higher scores indicate more deviation from normal.

Figure 3. Relative abundances of core taxa differentiating dysbiotic from normobiosis
hair sites while controlling for age differences. Within each genus, only OTUs that are
significantly associated with dysbiosis are displayed.

B