1 **Anterior-posterior polarity signals differentially regulate regeneration-competence of the**
2 **tapeworm *Hymenolepis diminuta*.**
3

4 Elise McCollough Nanista[1], Landon Elizabeth Poythress[1], Isabell Reese Skipper[1], Trevor
5 Haskins[1], Marieher Felix Cora[2] and Tania Rozario[1]*
6
7 [1]University of Georgia, Athens, GA, USA
8 [2]University of Puerto Rico-Cayey, PR, USA
9
10 Running title: Tapeworm regeneration-competence
11
12 * corresponding author
13 tania.rozario@uga.edu
14 500 D.W. Brooks Dr., Athens, GA 30602, USA
15
ORCIDS:
EMN: 0009-0002-3844-876X
LEP: 0009-0007-5836-5714
IRS: 0009-0009-1869-7292
TH: 0009-0007-1528-3807
MFC: 0009-0004-9971-0826
TR: 0000-0002-9971-5211
23

Keywords: regeneration, tapeworm, planaria, flatworm, Wnt signaling

Abstract

Competence to regenerate lost tissues varies widely across species. The rat tapeworm,

Hymenolepis diminuta, undergoes continual cycles of shedding and regenerating thousands of

reproductive segments to propagate the species. Despite its prowess, *H. diminuta* can only

regenerate posteriorly from a singular tissue: the neck or germinative region (GR). What cells and

signaling pathways restrict regeneration competence to the GR? In this study, we show that the

head regulates regeneration-competence by promoting maintenance of the GR and inhibiting

proglottid formation in a distance-dependent manner. Anterior-posterior (A-P) patterning within

the GR provide local signals that mediate these head-dependent responses. *βcat1* is necessary for

stem cell maintenance, proliferation and proglottidization. On the other hand, *sfrp* is necessary for

maintaining the GR at its proper length. Our study demonstrates that the head organizes a balance

of pro- and anti-regeneration signals that must be integrated together and therefore control

competence to regenerate.

Summary statement

We uncover how the tapeworm head organizes pro- and anti-regeneration signals that control

competence to regenerate at prolific rates.

Introduction

Regeneration occurs over a wide range of biological scales from regeneration of axons at the cellular scale to whole body regeneration in planarians and hydra (Bely and Nyberg, 2010). Tapeworms represent a fascinating case study in the field of regeneration. They descend from a monophyletic clade (Neodermata) exclusively populated by parasites within the Playhelminthes phylum (Laumer et al., 2015). As a sister group to planarians, it is tempting to assume that shared regenerative abilities have enabled tapeworms to achieve their monstrous growth potential and reproductive prowess. In the normal life cycle of many adult tapeworms, large pieces of their body are shed to disperse the developing embryos for consumption by an intermediate host. Simultaneously, the adult tapeworm continues to replenish the lost tissue in the intestine. Studies with the rat tapeworm, *Hymenolepis diminuta*, demonstrated that serially transplanting anterior pieces of tapeworms into new host intestines could greatly extend the lifespan of this parasite (Goodchild, 1958; Read, 1967). *H. diminuta* can regenerate all lost reproductive segments (proglottids) and may not inherently need to age and die. However, regeneration only occurs from one anatomical structure: the neck or germinative region (GR) (Rozario et al., 2019). *H. diminuta* regeneration is dependent on a large and heterogeneous population of adult stem cells that are distributed throughout the body and do not preferentially reside in the GR (Rozario et al., 2019). By transplanting cells from regeneration-competent and regeneration-incompetent regions into lethally irradiated tapeworm GRs, we previously demonstrated that proglottid regeneration is independent of stem cell source (Rozario et al., 2019), suggesting that signals within the GR play important roles in regulating tapeworm regeneration.

A plethora of signals likely operate in the GR and control stem cell proliferation, survival, differentiation, as well as proglottidization and competence to regenerate. What these signals are and how they operate is largely unknown. Here we show that regeneration competence is tightly linked to signals from the head. The head plays seemingly contradictory roles as it is necessary to maintain the GR but negatively regulates proliferation and proglottid regeneration. We find two genes typically associated with Wnt signaling, *sfrp and βcatenin1*, as critical mediators of each head-dependent phenotype. Our study demonstrates that for successful proglottid regeneration to occur, a balance of pro- and anti-regeneration signals must be overcome to enable GR regeneration

and then initiation of proglottids. *H. diminuta* provides a fascinating example of how extrinsic

signals can promote or restrict the ability to regenerate.

Results

***H. diminuta* is competent to regenerate the GR.**

The basic adult body plan of *H. diminuta* includes a head (scolex), an unsegmented GR and a

strobilated body made of metameric proglottids (Fig. 1A). *H. diminuta* cannot regenerate its

anterior or head but can regenerate proglottids from the GR when amputated outside of the GR

(Rozario et al., 2019). The GR is the only regeneration-competent tissue, but it is unclear if the

GR itself can regenerate. Nuclei staining with DAPI enables a quick and easy way to observe and

measure the GR, which is defined as the region between the base of the head and the first

proglottid. When tapeworms are continuously grown in vitro, the GR length is typically 1 mm

(Rozario et al., 2019). After acclimating worms to in vitro culture conditions for 3 days, we

amputated within the GR (0.5 mm) vs. outside the GR (2 mm) and grew the anterior fragments for

15 days. In both cases, proglottids were regenerated (Fig. 1A). Regenerates that were amputated

within the GR, successfully formed 1 mm long GRs 15 days post amputation (dpa) comparable to

regenerates that were amputated outside of the GR (Fig. 1B). The 0.5 mm fragments first

regenerated GRs within 2-4 days before new proglottids were added (Fig. 1C). Once proglottids

were formed, the rate of proglottid regeneration was comparable between 0.5 mm and 2 mm

fragments, at 14-15 proglottids per day (Fig. 1C). Growth curves for both regenerates were also

comparable and non-linear since individual proglottids increase in length as they mature (Rozario

et al., 2019). Thus, when amputated within the GR, *H. diminuta* is capable of regenerating the

posterior GR and subsequently proglottids.

The GR is patterned by differentially expressed genes along the anterior-posterior (A-P) axis

(Rozario et al., 2019). If *H. diminuta* can regenerate its GR, then A-P polarized patterns should be

restored following amputation. We performed whole mount in situ hybridization (WISH) for three

posterior-enriched transcripts: *βcatenin1* (*βcat1*), *frizzled5/8* (*fzd5/8*), and *tumor necrosis factor*

receptor-16 (*tnfr16*) and one anterior-enriched transcript: *secreted frizzled related protein* (*sfrp*)

(Fig. 1D; uncut). At 0 dpa, all markers were reduced or lost but normal A-P polarized patterns in

successful regenerates were restored by at least 10 dpa demonstrating normal GR patterning (Fig.

1D). These data indicate that *H. diminuta* can successfully regenerate its GR and reestablish AP

patterning within 10 days.

GR regeneration fails after serial amputation.

Regeneration of proglottids is a stem cell-driven process and can occur after multiple rounds of serial amputation (Goodchild, 1958; Read, 1967; Rozario et al., 2019). Is GR regeneration similarly robust? We amputated worms within the GR and allowed them to regenerate for 18 days before re-amputating within the GR (Fig. 2A). Subsets of each group were fixed and analyzed to capture the variation and reproducibility of the amputations performed. Amputations were targeted at the half-way point or ~0.5 mm in worms that were briefly immobilized in cold media. We find that tissues extend slightly after heat-killing and fixation resulting in slightly longer 0 dpa fragments: 0.63±0.07 mm and 0.61±0.16 mm after each amputation (Fig. 2B). The percentage of GR length retained calculated over the mean GR length of uncut worms was close to the target range of 50% and was not statistically significant between the two amputations (48±9% after cut 1 and 56±11% after cut 2) (Fig. 2C). Despite making comparable cuts at each timepoint, marked differences in regeneration were observed after the second amputation within the GR (Fig. 2D-F). Successful regenerates were almost always produced after cut 1 but ranged wildly between 10-100% regeneration after cut 2 (Fig. 2E). When worms did regenerate after cut 2, they were significantly shorter (Fig. 2D) and produced fewer proglottids (Fig. 2F). The additional challenge of serial amputation revealed that GR regeneration does fail under certain conditions.

Signals from the head regulate proglottid regeneration and maintenance of the GR.

When amputated within the GR, the wound site was closer to the head. Could the head be exerting a negative effect on regeneration? Following head amputation, regenerates exhibited increased growth and proglottid regeneration (Fig. 3A) indicating that signals from the head do inhibit growth. Using thymidine analog F-*ara*-EdU incorporation to quantify the density of proliferating cells in the GR, we found that head amputation resulted in increased proliferation, which can explain the increased growth (Fig. 3B). In uncut worms, proliferation density in the GR increases with distance from the head (Fig. 3C) further supporting our conclusion that the head provides anti-proliferation signals. Upon head amputation and in vitro culture for 3 days, the increasing proliferation density along the GR A-P axis was no longer significant (Fig. 3D). However, the curves did not fully flatten and the pattern of decreased proliferation toward the most anterior was still detectable. This suggests that while signals from the head do regulate proliferation in the GR, the effects are mediated by factors resident within the GR as well.

Many regenerating animals respond by increasing proliferation at the wound site. Is this true in *H. diminuta*? Following amputation in the GR, we quantified proliferation at both anterior- and posterior-facing wounds and found increased proliferation (Fig. S1). However, the local increase in proliferation was equivalent regardless of head presence (Fig. S1) indicating that amputation itself does not explain the increased proliferation in -head regenerates. This implies that the head does indeed negatively regulate proliferation.

In addition to proliferation within the GR, maintenance of GR tissue must also be preserved as regeneration cannot occur without it. Previous results have shown that GR maintenance depends on the head; after head amputation, the GR is eventually lost, the regenerates become fully proglottidized and cannot continue adding proglottids (Rozario et al., 2019). To further characterize how the head affects proglottid regeneration and GR maintenance, four amputation schemes were compared: + head, ½ head, -head, and -0.5 mm anterior (Fig. 4A). Of these four schemes, making transverse cuts through the head was the most challenging. A sampling of ½ head fragments were fixed immediately and stained with anti-SYNAPSIN antibodies to visualize the nervous system. Though the amount of head tissue removed did vary, the brain was always retained (Fig. S2A: yellow arrowheads). After making the four different amputations at the anterior region, we tested how the four fragment types would regenerate following serial amputation. All worms were amputated posteriorly to obtain 2 mm fragments that were grown in vitro for 14 days, then re-amputated for 3 more cycles (Fig. 4A). The posteriors of regenerates were fixed and analyzed to determine the number of proglottids regenerated after every cycle. As previously reported (Rozario et al., 2019), tapeworms regenerated after serial amputation when the head was present (Fig. 4B). The ½ head regenerates consistently displayed increased proglottid regeneration compared to +head regenerates (Fig. 4B). Both +head and ½ head regenerates maintained their GRs throughout 2 months of in vitro culture. When the head was amputated completely (-head), proglottid regeneration eventually ceased though some regenerates still had GRs after the 4[th] amputation (Fig. 4B). However, the GR lengths in -head regenerates that were not fully proglottidized were significantly shorter (Fig. S2B) indicating that complete proglottidization was imminent. Previously, we observed rapid loss of the GR after head amputation, but anterior GR tissue was also removed in those experiments (Rozario et al., 2019). Cognizant of regional

differences along the A-P axis of the GR, we were careful to make minimal head amputations that

retained as much anterior GR as possible in this study. We find that the GR persists for much

longer than previously reported but does eventually recede. On the other hand, fragments in which

both the head and anterior GR were removed (-0.5 mm) became fully proglottidized by the second

amputation or earlier (Fig. 4B). In sum, GRs were maintained if the brain region was retained but

GRs were lost when more and more anterior tissue was removed (Fig. 4C).

The number of proglottids regenerated was also dependent on amputations at the head though our

analysis was complicated by GR loss in -head and -0.5mm fragments. After the first amputation,

a clear dose-dependent response was detected; when more head tissue was retained, the number of

proglottids decreased (Fig. S2C). This trend was maintained after serial rounds of amputation when

comparing +head and ½ head regenerates where GRs were not lost (Fig. S2D). After the 4^{th} round

of amputation, the remaining head area plotted against number of proglottids for +head and ½ head

regenerates revealed a negative correlation between head tissue abundance and proglottidization

(Fig. 4D). We also measured the effect of head tissue abundance on proliferation within the GR

using F-*ara*-EdU. A general trend of increasing proliferation with decreasing head and anterior

tissue emerged though differences between +head and ½ head were not statistically significant

(Fig. 4E-F). Taken together, the head effects on maintaining the GR and regulating proliferation

can be separated. Signals from the brain region are necessary to maintain the GR whereas

proliferation and proglottidization are regulated by head-dependent signals in a dose-dependent

manner.

βcatenin1 is necessary for proglottid regeneration and stem cell maintenance.

What is the molecular basis for the head-dependent effects on tapeworm regeneration? The Wnt

signaling pathway has been implicated in A-P polarity across many systems (Petersen and

Reddien, 2009) and plays roles in regulating cell proliferation, survival, and differentiation (Nusse

and Clevers, 2017). Multiple members of the Wnt signaling pathway were found to be either

anterior-enriched or posterior-enriched after RNA sequencing along the A-P axis of the GR

(Rozario et al., 2019). As βcatenin protein is a key downstream effector of Wnt signaling, we

decided to examine its function further. *H. diminuta* has three *βcatenin* paralogs as do other

parasitic flatworms (Montagne et al., 2019). BLASTp comparisons of putative *H. diminuta*

βcatenin paralogs with published sequences for other flatworms (*Schmidtea meddterranea,*
Schistosoma mansoni and Ecchinococcus multilocularis) demonstrated that *Hdim-βcat1*
(WMSIL1_LOCUS14475) is the rat tapeworm ortholog of *βcat1* (Table S1). *Hdim-βcat1* retains
expected conserved domains including multiple armadillo repeats and the DSGxxSxxx[S/T]xxxS
motif for CKI/GSK-phosphorylation necessary for targeting by the destruction complex
(Montagne et al., 2019; Valenta et al., 2012) (Fig. S3A). Capacity to bind α-catenin cannot be
ruled in or out at present (Fig. S3B; discussed further below). Thus *Hdim-βcat1* likely mediates
Wnt signaling and may also function in cell adhesion.
208
Where is *Hdim-βcat1* (hereafter *βcat1*) expressed in *H. diminuta*? In 6-day-old adults, *βcat1*
expression is largely absent in the head and anterior GR but progressively increases toward the
posterior GR (Fig. 5A). *βcat1* is also expressed throughout the body, in transverse stripes at
proglottid boundaries and within a subset of accessory reproductive structures (Fig. S4). The broad
expression pattern throughout the worm indicates that *βcat1* is expressed in many tissues and is
not restricted to a particular cell type. The posterior-biased pattern of *βcat1* expression was
restored after successful GR regeneration (Fig. 1D) leading us to wonder if *βcat1* is necessary for
proliferation and proglottidization from the GR. We performed RNAi by injecting double-stranded
RNA (dsRNA) targeting *βcat1* throughout the GR. 2 mm anterior fragments were amputated and
grown in vitro to assay regenerative ability. A dramatic growth defect was captured just 7 dpa/10
days post injection (dpi) (Fig. 5B). Decreased growth was evident by 3 dpa and this RNAi
treatment was ultimately lethal. The proglottids formed were extremely small and impossible to
quantify but there was a marked decrease in worm lengths following *βcat1* RNAi (Fig. 5C). RNAi
of *βcat1* reduced transcript levels to 8.7±2.0% (Fig. 5D). As *βcat1*-dependent Wnt signaling is
mediated through degradation of βCAT1 protein by the destruction complex, we investigated
expression of a key destruction complex component: *axin1*. Previous work has demonstrated that
tapeworm *E. multilocularis,* AXIN proteins directly interact with Emul-βCAT1 and
overexpression of Emul-AXIN1 results in degradation of Emul-βCAT1 (Montagne et al., 2019).
We find that after *βcat1* RNAi, *axin1* expression is markedly reduced to 18.8±11.1% of normal
levels (Fig. 5D). This suggests that canonical Wnt signaling is inhibited following *βcat1* RNAi
though other roles for *βcat1* cannot be ruled out.

The strong decrease in growth after *βcat1* RNAi suggests that the cycling stem cell population in

the GR is likely dependent on *βcat1*. We measured proliferation in the GR and detected a dramatic

decrease in dividing cells compared to controls (Fig. 5E). This suggests that the stem cell

population likely cannot be maintained in the absence of *βcat1*. To visualize the stem cell

population, we performed WISH for two known stem cell markers in *H. diminuta*:

minichromosome maintenance complex component-2 (*mcm2*) and *laminB receptor* (*lbr*) (Rozario

et al., 2019). Both stem cell markers showed decreased expression indicating that *βcat1* is required

to maintain stem cells in the GR (Fig. 5F).

GR maintenance requires *sfrp*.

Our analyses so far suggest that signals from the head set up gene expression patterns within the

GR with posteriorly enriched *βcat1* playing indispensable roles in maintaining stem cells.

However, it is unclear if *βcat1* regulates stem cell maintenance through Wnt signaling, cell

adhesion or both. If Wnt signaling is necessary, the head may segregate Wnt inhibitors toward the

anterior. Previously, RNA sequencing of A-P polarized transcripts in the GR found that *sfrp* was

anterior-enriched (Rozario et al., 2019). Like other tapeworms, *H. diminuta* has one true *sfrp*

family member and a second highly divergent member, *sfrp-like*, which was not investigated

further in this study. Interpro (Blum et al., 2020) predicts that Hdim-SFRP retains both frizzled

and netrin domains albeit with some variation (Fig. S5; discussed further below). Since SFRP can

inhibit Wnt signaling (Leyns et al., 1997; Rattner et al., 1997; Wang et al., 1997), we decided to

pursue its potential role in regeneration.

By WISH, *sfrp* expression is strikingly anterior-enriched. We detected *sfrp* expression sparsely

within the head and strongly throughout the GR with expression tapering off posteriorly (Fig. 6A).

In adult tapeworms, *sfrp* expression is largely absent from the strobilated body except at the lateral

ends of proglottids (Fig. S4). Perhaps knocking down *sfrp* would result in the opposite phenotypes

observed from *βcat* RNAi and would phenocopy head amputation. We injected dsRNA into the

head, amputated 2 mm anterior fragments, reinjected dsRNA into the head and collected worms at

10-12 dpa (Fig. 6B). Contrary to our expectations, a modest decrease in worm lengths and

proglottid regeneration were observed (Fig. 6C-D). Strikingly, we discovered that the GRs were

significantly shortened following *sfrp* RNAi (Fig. 6E-F). This is the first time GR length has been

affected by RNAi of any target gene. Injecting dsRNA into the head does not preclude diffusion

into the GR but the knockdown is likely to be anteriorly biased. If so, we may inadvertently miss

other potential roles for *sfrp* toward the posterior GR. We injected dsRNA throughout the GR and

observed the same phenotypes (Fig. S6A-D). Both injection schemes resulted in similar *sfrp*

knockdown efficacy (Fig. S6E). While the levels of knockdown did vary substantially, GR length

reduction was consistently observed.

Our results indicate that *sfrp* is necessary for proper GR maintenance but is unlikely to be involved

in head-dependent inhibition of proliferation or proglottidization. Instead, *sfrp* likely mediates

head-dependent GR maintenance. Accordingly, we observed decreased *sfrp* expression just 3 days

after head amputation (Fig. 6G). This correlation supports the conclusion that the failure to

maintain the GR after head amputation is mediated through anterior-enriched *sfrp* expression.

Decreased proglottid regeneration after *sfrp* RNAi does not support the hypothesis that *sfrp*

functions as a Wnt inhibitor. Proliferation within the GR was not significantly changed after *sfrp*

RNAi (Fig. S5D). However, the shortening of the GR would bring the posterior boundary of the

GR closer to the head. Thus, *sfrp* RNAi may indirectly exacerbate the anti-proglottidization and

anti-proliferation effects from the head.

Amputation distance from the head determines whether pro- or anti-regeneration signals

dominate.

The mechanisms that govern the two head-dependent effects on regeneration have begun to

emerge. The presence of the head is necessary for *sfrp* expression and GR maintenance. However,

the presence of the head also inhibits proliferation and proglottidization, which are dependent on

βcat1. Perhaps when the posterior wound is too close to the head, the growth inhibitory effect of

the head dominates, and GR regeneration fails. This suggests that regeneration would fail more

when the posterior cut is made at decreasing distance from the head. To test this, worms were

allowed to regenerate after 0.5 mm amputation within the GR and then cut a second time at 4

different distances from the head. All samples were cut at ~1mm and then subsets were

progressively shaved closer and closer toward the head (Fig. 7A; cuts A-D). Subsets of fragments

were fixed at 0 dpa and we verified that cut A-D progressively retained 87±5%, 64±7%, 47±9%,

and 22±4% of the GR (Fig. 7B). As expected, GR and proglottid regeneration plummeted at cut B

or C over three independent experiments (Fig. 7C). Surprisingly, the smallest fragments (cut D)

regenerated equivalently to the longest fragments (cut A) (Fig. 7C-D and Fig. S7). Fragments from

cut A and D achieved different absolute lengths at 18 dpa but the fold change in growth was

comparable (Fig. 7E). Also, no significant difference is GR length was detected in regenerates

from cut A or D (Fig. 7F). Thus, the inhibitory effect of the head was overcome when enough

posterior GR tissue was removed. These results suggest that there are yet more signals to be

uncovered that explain how regeneration is promoted and restricted in tapeworms.

While there is still much to discover, we propose the following working model (Fig. 8A). The head

likely produces direct or indirect inhibitor(s) of a posterior signal that we dub factor X. Factor X

may be posterior Wnts, other signaling pathways that crosstalk with Wnt signaling or Wnt-

independent signals. The head promotes *sfrp* expression, which is necessary to maintain the GR.

The posterior GR produces a counteracting signal that results in stable maintenance of the GR

length. Factor X acts via *βcat1* to promote proglottid formation. Proglottids only form at sufficient

distance from the head where high *βcat1* activity is present. Upon head amputation (Fig. 8B), *sfrp*

expression is inhibited causing the posterior boundary of the GR to creep up towards the anterior.

Without the head, inhibition of factor X is relieved and overstimulation of *βcat1* results in

increased proglottid regeneration until the GR is fully lost.

Our experiments also demonstrated that cutting within the GR produced highly mixed results,

which can now be explained by our model. Tapeworms grown in vitro are adjusting to a new

environment and all tissue dimensions change with time. The various amputation schemes within

the GR used throughout this paper revealed that depending on the state of signaling gradients from

the head and GR, we pushed regeneration competence toward a pro- or anti-regeneration state.

When cuts were made "mid" GR at an unfavorable distance from the head (Fig. 8C), inhibition of

factor X dominated leading to failure to regenerate proglottids. However, when a minimal

fragment was produced (Fig. 8D), perhaps the absence of factor X negates the inhibitory functions

of the head. The head could promote establishment of *sfrp* and proper growth of the GR. At a

sufficient distance from the head, factor X is now stimulated and can promote proglottid

regeneration. The exact players, direct and indirect relationships, as well as additional molecular

mechanisms predicted by our working model still need to be tested but the general take-home

holds: a balance of pro- and anti-regeneration factors operate in the GR of *H. diminuta*.

Discussion

Regenerative abilities vary widely even amongst species that are competent to regenerate. Thus, it is important to understand the factors that both promote and inhibit regeneration. The flatworm phylum offers opportunities to study variations in regenerative abilities. While the free-living planarian *S. mediterranea* can regenerate from almost any amputation fragment, regeneration in *H. diminuta* is far more limited. This study explores how a balance of activating and inhibiting signals operate in the regeneration-competent region: the GR.

We show that in addition to regenerating proglottids, *H. diminuta* can also regenerate the posterior GR following amputation. However, the challenge of serial amputation within the GR revealed that there are anti-regeneration factors that were previously unknown. By testing amputation schemes at the anterior, we revealed that the head plays at least two seemingly contradictory roles: 1) the head promotes GR maintenance and 2) the head inhibits proliferation within the GR and proglottid regeneration. Interestingly, these two roles can be separated both anatomically and through molecular regulators.

GR maintenance depends on the brain-region without which the GR will eventually be lost. Loss of the GR is gradual unless the anterior GR is also removed. The head likely acts as an organizer that relays signals within the GR. In this model, total loss of the GR will occur after anterior-enriched patterns within the GR turnover and cannot be reestablished following head amputation. One critical mediator of head-dependent GR maintenance is *sfrp*. Like head amputation, RNAi of *sfrp* results in shortened GRs. However, *sfrp* RNAi does not phenocopy the growth spurts observed after head amputation. The mechanism by which *sfrp* maintains the GR is more mysterious. While *sfrp* is primarily known as a Wnt antagonist, it has also been shown to bind and stimulate frizzled receptors, interact with other signaling pathways and modulate cell-matrix interactions by binding integrins (Bovolenta et al., 2008; Mii and Taira, 2011). There is evidence that *sfrps* can stimulate Wnt and BMP signaling in the zebrafish retina (Holly et al., 2014). In Drosophila, *sfrps* can modulate Wnt signaling in a biphasic manner: inhibiting Wnt signaling when expressed at high levels while promoting Wnt signaling when expressed at low levels (Üren et al., 2000). In *H. diminuta*, the *sfrp* RNAi phenotypes were not the opposite of the *βcat1* RNAi phenotypes as one

might predict if *sfrp* and *βcat1* were acting solely as Wnt antagonist and Wnt effector respectively.

Nonetheless, it is still possible that *sfrp* is inhibiting Wnt signaling. In many flatworms, Wnt

ligands are secreted from both anterior and posterior domains (Armstrong et al., 2025; Koziol et

al., 2016; Reddien, 2018; Soria et al., 2020). Expression of *sfrp* throughout the GR may be required

as a permissive signal to tune Wnt signaling to a moderately low level so that it can be acted upon

differentially by other modulators. Alternatively, *sfrp* may be functioning independent of Wnt

signaling. Future studies investigating the RNAi phenotypes of other Wnt antagonists, Wnt ligands

and frizzled receptors will be necessary to determine how Wnt signaling regulates individual cell-

and tissue-level behaviors at the GR that are necessary for regeneration.

A further complication is that the biochemical properties and interactors of SFRP proteins in

flatworms in not well understood. Both frizzled and netrin domains are readily identifiable in

flatworm SFRPs. The netrin domain should contain 6 conserved cysteine residues (Bhat et al.,

2007; Chong et al., 2002) and conflicting conclusions have been asserted as to whether parasitic

flatworm SFRPs satisfy this requirement (Armstrong et al., 2025; Koziol et al., 2016; Riddiford

and Olson, 2011). Our alignments predict that all 6 cysteine residues are present, however, the 6[th]

residue is displaced by three animo acids in flatworms but not humans (Fig. S5). The three

additional amino acids are also present in free living *S. mediterranea* and *Smed-sfrp1* is generally

accepted as a Wnt antagonist. While planarian *sfrp* paralogs show complex expression patterns

(Gurley et al., 2010), *Smed-sfrp1* is an anterior marker that robustly responds to Wnt inhibition

and loss of posterior identity (Doddihal et al., 2024; Hill and Petersen, 2018; Reuter et al., 2015;

Scimone et al., 2016; Sureda-Gómez et al., 2015). RNAi of *Smed-sfrp1* shifts the gradient of Smed-

βCAT1 protein expression anteriorly supporting *Smed-sfrp1* function as a Wnt antagonist

(Stückemann et al., 2017). However, other Wnt antagonists such as *Smed-notum* demonstrate

much clearer evidence of Wnt inhibition. *Smed-notum* RNAi results in tail regeneration in place

of heads (Petersen and Reddien, 2011) as opposed to *Smed-βcat1* RNAi, which results in head

regeneration in place of tails (Gurley et al., 2008; Iglesias et al., 2008; Petersen and Reddien,

2008). Thus, it remains possible that a muted or more ambiguous role in Wnt signaling by *sfrp1* is

a shared feature in free-living and parasitic flatworms.

In addition to GR maintenance, the head negatively regulates both proliferation in the GR and

proglottid regeneration in a dose-dependent and distance-dependent manner with increased

proliferation farther from the head. This observation suggests that the head may be the source of

growth inhibitors that diffuse or are differentially received along the A-P axis of the GR. We find

that at least one posterior-enriched factor, *βcat1*, is necessary for stem cell maintenance in the GR.

Knockdown of *βcat1* results in a dramatic inhibition of growth, proglottid regeneration and

ultimately death. Recently, a study in the liver fluke, *Fasciola hepatica,* demonstrated that RNAi

of *Fhep-βcat1* resulted in stunted juvenile growth, decreased proliferation and worm death

(Armstrong et al., 2025), phenocopying our observations. Thus, roles for *βcat1* in regulating stem

cell proliferation and/or survival may be a common theme in parasitic flatworms as it is in many

metazoans as well as in neoplastic transformations (Mohammed et al., 2016; Steinhart and Angers,

2018) .

In addition to canonical Wnt signaling, β-catenin proteins have additional roles in cell adhesion

through linking cadherins with the actin cytoskeleton via α-catenin. These two functions have been

reported in many but not all species. In *S. mediterranea,* there is complete separation of function

by two catenin paralogs: Smed-βCAT1 mediates Wnt signaling whereas Smed-βCAT2 mediates

cell adhesion (Chai et al., 2010; Su et al., 2017). Accordingly, Smed-βCAT1 does not have α-

catenin binding sites (Chai et al., 2010; Su et al., 2017). The presence/absence of α-catenin binding

sites in βCAT1 from tapeworms is more ambiguous. Previously, a likely α-catenin binding region

demarcated by 10 putative critical residues within a 26 amino acid span was identified for βCAT1

and βCAT2 in flatworms (Montagne et al., 2019). The 10 residues were identified as necessary for

α-catenin binding to mammalian plakoglobin and/or βcatenin using alanine scanning mutagenesis

(Aberle et al., 1996; Pokutta and Weis, 2000). When compared to human βcatenin (CTNB1), we

and others find strong conservation of this α-catenin binding region in βCAT2 from multiple

flatworms (Fig. S3B) (Montagne et al., 2019). While Smed-βCAT1 is clearly very divergent at

this site, two tapeworms (*H. diminuta* and *E. multilocularis*) show more similarity to the human

reference (Fig. S3B) (Montagne et al., 2019). Thus, we cannot rule out the possibility that βCAT1

in tapeworms functions in both Wnt signaling and cell adhesion, unlike their free-living

counterparts. Future studies investigating the biochemical interactions and functional roles of all

βcatenin paralogs will help resolve the mechanisms regulating stem cell maintenance in parasitic

flatworms.

The role of *βcat1* in posterior identity is well conserved across many taxa (Petersen and Reddien,

2009). In planarian species that are deficient in head regeneration, knockdown of *βcat1* alone can

rescue head regeneration indicating that failure to regenerate the head in wild type worms was due

to the inability to overcome the posteriorizing effects of *βcat1* (Liu et al., 2013; Sikes and

Newmark, 2013; Umesono et al., 2013). *H. diminuta* cannot regenerate its head and we initially

assumed that tapeworm *βcat1* would be acting in a homologous fashion to planarians. Because

βcat1 is required for stem cell maintenance in *H. diminuta,* potential roles in posteriorization could

not be uncovered. It remains possible that Wnt signaling is necessary to maintain GR posterior

identity in *H. diminuta*. Extensive RNAi manipulations of Wnt signaling components that are

beyond the scope of this study will help reveal a more holistic understanding of Wnt signaling in

this system. However, tapeworms and planarians do differ in that *Smed-βcat1* is not necessary for

stem cell maintenance.

The differences in utilization of *βcat1* between different species of platyhelminths is potentially

very interesting. Perhaps segregating functions in posteriorization vs. stem cell maintenance in *S.*

mediterranea has enabled more nimble remodeling of axes that makes this species so remarkable

at regenerating from almost any amputated fragment. In intestinal tapeworms, the head contains

suckers that are necessary for attachment without which the tapeworm will simply be extruded

with the feces. Thus, maintenance of the head and proglottid regeneration should not be easily

decoupled in tapeworms. Considering the wider role for βcatenin in regulating stem cell

proliferation and survival across metazoa, it would be easy to suppose that planarian phenotype is

divergent. However, studies in the acoel *Hofstenia miamia*, a representative of a sister taxa to all

bilaterians, show that RNAi of *Hmia-βcat1* is necessary for maintaining posterior identity but

functions in regulating stem cells, proliferation, or growth were not reported (Srivastava et al.,

2014; Tewari et al., 2019). Considering that *Hmia-βcat1* RNAi experiments were performed in the

context of regeneration, inhibition of growth is unlikely to have been missed with the caveat the

RNAi cannot be fully penetrant. Thus, there is evidence that the planarian phenotype is ancestral.

A deeper understanding of how catenin paralogs have evolved across platyhelminths and how they

have functionally specialized will be illuminating. Such investigations would enrich our

understanding of how strategies to regulate axial identity and stem cell proliferation/survival have

evolved in regeneration-competent species.

Our work has revealed that a balance of opposing signals from anterior and posterior ends of the

GR are necessary for proper regeneration. It makes intuitive sense that initiation of proglottids

must be inhibited close to the head if the GR is to be maintained as an unsegmented tissue. A

recent study from the mouse bile duct tapeworm, *Hymenolepis microstoma,* has described discrete

expression of Wnt and Hedgehog signaling components in a signaling quartet (SQ) within the

transition zone from GR to fully elaborated proglottids (Jarero et al., 2024). It is possible the SQ

initiates proglottid formation and represents the unknown X factor predicted by our study. How

the SQ responds to head-dependent signals and vice versa is currently unknown and warrants

future investigation.

Our study also shows that regulation of proliferation dynamics is dependent on signals from the

head. Why should the head provide anti-proliferation signals at all? One intriguing possibility is

that distinct subpopulations of stem cells within the GR respond differentially to signals from the

head. Such signals may affect the balance of self-renewal vs. differentiation. Alternatively, a slow-

cycling pluri- or multi-potent subpopulation of stem cells may require these head-dependent

signals. There is already evidence for different cycling rates in stem cells of parasitic flatworms

(Herz et al., 2024; Wang et al., 2018) . In this study, we observed that amputations made

progressively closer to the head inhibited regeneration until enough posterior GR tissue was

removed, which restored competence to regenerate. Perhaps the anterior portion of the GR contains

the minimal cohort of stem cells required to regenerate the GR and seed new proglottids. These

anterior stem cells may represent a slow-cycling population with greater potency. These ideas are

highly speculative, but investigating how head-dependent signals impact stem cell subpopulations

in the GR may shed light on the regulation of stem cell potency in tapeworms and potentially other

species.

Materials and Methods

Animal care and use

The life cycle of *H. diminuta* was maintained in house using mealworm beetles (*Tenebrio molitor*) and Sprague-Dawley rats for the larval and adult stages respectively. Rats were fed 200-400 infective cysticercoids by oral gavage in ~0.5 mL 0.85% NaCl. Typically, adult tapeworms were collected 6 days post infection by euthanizing infected rats in a CO_2 chamber and flushing the small intestine with 1X Hanks Balanced Salt Solution (HBSS: 21-023-CV; Corning). Rodent care was performed in accordance with protocols approved by the Institutional Animal Care and Use Committee (IACUC) of the University of Georgia (A2023 10-019-Y1-A0).

In vitro parasite culture

Adult tapeworms or amputated fragments were grown in vitro in biphasic cultures as described before (Rozario et al., 2019). 2.5X MEM essential amino acids (M5550; Sigma-Aldrich) was added to the liquid culture: Working Hanks 4 (WH4: HBSS/4 g/L glucose/antibiotic-antimycotic). This ameliorated inconsistencies between batches of blood used for culturing but was not strictly necessary.

In situ hybridization and other staining

Heat-kills were performed by swirling tapeworms in 75°C water for a few seconds before fixing (4% formaldehyde in Phosphate Buffered Saline with 0.3% TritonX-100 (PBSTx)) for 30 min-2 hr at room temperature or overnight at 4°C. Worms were either dehydrated into methanol and stored (-30°C) or used directly for staining. For simple phenotyping of growth and regeneration, worms were stained in 1 μg/mL DAPI (D9542; Sigma-Aldrich)/PBSTx overnight at 4°C and cleared in mounting solution (80% glycerol/10 mM Tris pH7.5/ 1 mM EDTA) overnight before mounting and imaging. Previously published methods were employed for in situ hybridization (Rozario et al., 2019) and immunostaining (Rozario and Newmark, 2015). Anti-Syn antibodies (3C11; Developmental Studies Hybridoma Bank) were used at 1:200 with Tyramide Signal Amplification (TSA).

F-*ara*-EdU uptake and staining

F-*ara*-EdU pulses were performed in 0.1 µM F-*ara*-EdU (T511293; Sigma-Aldrich)/1%

DMSO/WH4 at 37°C for 1 hr. Staining was performed as previously described (Ishan et al., in

press) with the following specifics: 1) 15 min proteinase-K digestion and 2) 10 min TSA reaction.

<u>Imaging and image processing</u>

Confocal imaging was performed on a Zeiss LSM 900 with the following objectives: 20X/Plan-

Apochromat/0.8 M27/FWD=0.55mm and 63X/Plan-Apochromat/1.40 Oil DIC M27. WISH and

DAPI-stained worms were imaged on a Zeiss AxioZoom V16 Microscope. Fiji (Schindelin et al.,

2012) was used for brightness/contrast adjustments, maximum-intensity projections and

measurements of lengths, areas, and numbers of proglottids. F-*ara*-EdU+ cells were quantified

using Imaris (Oxford Instruments) spot finder as previously described (Ishan et al., in press) .

<u>RNAi</u>

To perform knockdowns, tapeworms were microinjected with dsRNA (1-1.5 µg/µl in HBSS), 2

mm anterior fragments were amputated 3 days later and cultured in vitro. Injections were

performed using a Femtojet 4i (Eppendorf) at 500 hPa for 0.3-1s. dsRNA was synthesized as

previously described (Rouhana et al., 2013) from ~1 kb PCR products from cDNA cloned into

pJC53.2 vector (Collins et al., 2010). Experimenters were blinded to the dsRNA used throughout

microinjections, staining and image analysis.

<u>qRT-PCR</u>

To exclude genomic DNA, primers were designed to flank introns or span an intron-exon junction.

Amplicon range was 133-277 bp. All primers were tested on cDNA +/- reverse transcriptase and

gel electrophoresis to confirm the presence of a single product. Primer efficiencies were calculated

from 2-fold serial dilutions of cDNA.

For RNA extractions, 3 mm anteriors from 4-5 worms were cut, submerged in 200 µL TRIzol

Reagent (15596026; Invitrogen) and processed according to manufacturer's instructions with these

exceptions: 1) samples were homogenized with a motorized pestle twice in a semi-solid state then

centrifuged at >12K RCF for 5 min to obtain 180 µL clean supernatant, 2) RNA pellets were

washed twice with 75% ethanol-DEPC and 3); pellets were resuspended in 40 µL nuclease-free

water. DNAse (M6101; Promega) treatment was performed for 1 hr at 37°C. RNA was cleaned

using RNA Clean & Concentrator-5 kit (R1016; Zymo Research) and the concentration measured

on a Nanodrop spectrophotometer. cDNA synthesis was performed, according to manufacturer's

protocol, using Oligo(dT)$_{20}$ primers and iScript Select cDNA Synthesis Kit (1708897; Bio-Rad)

with undiluted RNA. cDNA was further diluted with 5-25 µL nuclease-free water depending on

the number of reactions needed.

qRT-PCR was performed using GoTaq MasterMix (A6001; Promega) according to manufacturer's

instructions and scaled to 25 µL reactions with 2 µL cDNA template for each of 3 technical

replicates. Each reaction had 0.4 µM primer pairs, 0.3 µM CXCR in 1X master mix. 7500 Real

Time PCR System (Applied Biosystems) was used for 40 amplification cycles: 15 s at 95°C and 1

min at 60°C. Relative gene expression change was calculated using the PFAFFL equation

(Hellemans et al., 2007; Vandesompele et al., 2002) to account for different primer efficiencies.

Internal normalization was done on the geometric mean of two previously published standards:

60S ribosomal protein L13 (*60Srpl13*) and *myosin heavy chain* (*mhc*) (Rozario et al., 2019) .

<u>Statistical Analysis</u>

Graph Pad Prism 10 was used for all statistical analyses. All experiments were repeated at least

twice. Error bars, statistical tests, number of replicates (N) and sample sizes (n) are indicated in

corresponding figure legends. P-values: ns = not significant, * = $p < 0.05$, **=$p < 0.01$, ***=$p < 0.001$,

****=$p < 0.0001$.

Acknowledgements

We are sincerely thankful to Phil Newmark and Melanie Issigonis (Morgridge Institute for

Research, WI). Phil's unwavering support in the initialization of these projects and continued

advice was instrumental in our success. We are also grateful for Melanie's critical evaluation of

our manuscript and valuable feedback. We thank the Biomedical Microscopy Core at the

University of Georgia, Athens, GA for necessary microscopy services.

Competing interests

No conflicts to report.

Funding

This work was funded by a National Institute of Allergy and Infectious Disease (NIAID) grant

(DP2 AI 154416-01) to Tania Rozario.

Data and resource availability

All relevant data and resources can be found within the article and its supplementary information.

Primers and transcripts used are listed in Table S2. All source data is provided in Table S3.

References

Aberle, H., Schwartz, H., Hoschuetzky, H. and Kemler, R. (1996). Single Amino Acid Substitutions in Proteins of the armadillo Gene Family Abolish Their Binding to α-Catenin (∗). *J. Biol. Chem.* **271**, 1520–1526.

Armstrong, R., Marks, N. J., Geary, T. G., Harrington, J., Selzer, P. M. and Maule, A. G. (2025). Wnt/β-catenin signalling underpins juvenile *Fasciola hepatica* growth and development. *PLOS Pathog.* **21**, e1012562.

Bely, A. E. and Nyberg, K. G. (2010). Evolution of animal regeneration: re-emergence of a field. *Trends in Ecology & Evolution* **25**, 161–170.

Bhat, R. A., Stauffer, B., Komm, B. S. and Bodine, P. V. N. (2007). Structure–Function analysis of secreted frizzled-related protein-1 for its Wnt antagonist function. *J. Cell. Biochem.* **102**, 1519–1528.

Blum, M., Chang, H.-Y., Chuguransky, S., Grego, T., Kandasaamy, S., Mitchell, A., Nuka, G., Paysan-Lafosse, T., Qureshi, M., Raj, S., et al. (2020). The InterPro protein families and domains database: 20 years on. *Nucleic Acids Res.* **49**, D344–D354.

Bovolenta, P., Esteve, P., Ruiz, J. M., Cisneros, E. and Lopez-Rios, J. (2008). Beyond Wnt inhibition: new functions of secreted Frizzled-related proteins in development and disease. *J. Cell Sci.* **121**, 737–746.

Chai, G., Ma, C., Bao, K., Zheng, L., Wang, X., Sun, Z., Salò, E., Adell, T. and Wu, W. (2010). Complete functional segregation of planarian β-catenin-1 and -2 in mediating Wnt signaling and cell adhesion. *The Journal of biological chemistry* **285**, 24120–24130.

Chong, J. M., Üren, A., Rubin, J. S. and Speicher, D. W. (2002). Disulfide Bond Assignments of Secreted Frizzled-related Protein-1 Provide Insights about Frizzled Homology and Netrin Modules. *J. Biol. Chem.* **277**, 5134–5144.

Collins, J. J., Hou, X., Romanova, E. V., Lambrus, B. G., Miller, C. M., Saberi, A.,
Sweedler, J. V. and Newmark, P. A. (2010). Genome-wide analyses reveal a role for
peptide hormones in planarian germline development. *PLoS biology* **8**, e1000509.

Doddihal, V., Mann, F. G., Ross, E. J., McKinney, M. C., Guerrero-Hernández, C.,
Brewster, C. E., McKinney, S. A. and Alvarado, A. S. (2024). A PAK family kinase and
the Hippo/Yorkie pathway modulate WNT signaling to functionally integrate body axes
during regeneration. *Proc. Natl. Acad. Sci.* **121**, e2321919121.

Goodchild, C. G. (1958). Transfaunation and Repair of Damage in the Rat Tapeworm,
Hymenolepis diminuta. *The Journal of parasitology* **44**, 345–351.

Gurley, K. A., Rink, J. C. and Alvarado, A. S. (2008). β-catenin defines head versus tail
identity during planarian regeneration and homeostasis. *Science* **319**, 323–327.

Gurley, K. A., Elliott, S. A., Simakov, O., Schmidt, H. A., Holstein, T. W. and Alvarado, A.
S. (2010). Expression of secreted Wnt pathway components reveals unexpected complexity of
the planarian amputation response. *Developmental biology* **347**, 24–39.

Hellemans, J., Mortier, G., Paepe, A. D., Speleman, F. and Vandesompele, J. (2007). qBase
relative quantification framework and software for management and automated analysis of
real-time quantitative PCR data. *Genome Biol.* **8**, R19.

Herz, M., Zarowiecki, M., Wessels, L., Pätzel, K., Herrmann, R., Braun, C., Holroyd, N.,
Huckvale, T., Bergmann, M., Spiliotis, M., et al. (2024). Genome-wide transcriptome
analysis of *Echinococcus multilocularis* larvae and germinative cell cultures reveals genes
involved in parasite stem cell function. *Front. Cell. Infect. Microbiol.* **14**, 1335946.

Hill, E. M. and Petersen, C. P. (2018). Positional information specifies the site of organ
regeneration and not tissue maintenance in planarians. *eLife* **7**, e02238.

Holly, V. L., Widen, S. A., Famulski, J. K. and Waskiewicz, A. J. (2014). Sfrp1a and Sfrp5
function as positive regulators of Wnt and BMP signaling during early retinal development.
Dev. Biol. **388**, 192–204.

Iglesias, M., Gomez-Skarmeta, J. L., Saló, E. and Adell, T. (2008). Silencing of *Smed-*
βcatenin1 generates radial-like hypercephalized planarians. *Development* **135**, 1215–1221.

Ishan, M., Skipper, I. R. and Rozario, T. (2025). Combined fluorescent in situ hybridization
and F-ara-EdU staining on whole mount *Hymenolepis diminuta. Biol. Methods Protoc.*
bpaf011. (in press)

Jarero, F., Baillie, A., Riddiford, N., Montagne, J., Koziol, U. and Olson, P. D. (2024).
Muscular remodeling and anteroposterior patterning during tapeworm segmentation. *Dev.*
Dyn. **253**: 998–1023.

Koziol, U., Jarero, F., Olson, P. D. and Brehm, K. (2016). Comparative analysis of Wnt
expression identifies a highly conserved developmental transition in flatworms. *BMC biology*
14, 1.

Laumer, C. E., Hejnol, A. and Giribet, G. (2015). Nuclear genomic signals of the
"microturbellarian" roots of platyhelminth evolutionary innovation. *eLife* **4**, e05503.

Leyns, L., Bouwmeester, T., Kim, S.-H., Piccolo, S. and Robertis, E. M. D. (1997). Frzb-1 Is
a Secreted Antagonist of Wnt Signaling Expressed in the Spemann Organizer. *Cell* **88**, 747–
756.

Liu, S. Y., Selck, C., Friedrich, B., Lutz, R., Vila-Farré, M., Dahl, A., Brandl, H.,
Lakshmanaperumal, N., Henry, I. and Rink, J. C. (2013). Reactivating head regrowth in a
regeneration-deficient planarian species. *Nature* **500**, 81–84.

Mii, Y. and Taira, M. (2011). Secreted Wnt "inhibitors" are not just inhibitors: Regulation of
extracellular Wnt by secreted Frizzled-related proteins. *Dev., Growth Differ.* **53**, 911–923.

Mohammed, M. K., Shao, C., Wang, J., Wei, Q., Wang, X., Collier, Z., Tang, S., Liu, H.,
Zhang, F., Huang, J., et al. (2016). Wnt/β-catenin signaling plays an ever-expanding role in
stem cell self-renewal, tumorigenesis and cancer chemoresistance. *Genes Dis.* **3**, 11–40.

Montagne, J., Preza, M., Castillo, E., Brehm, K. and Koziol, U. (2019). Divergent Axin and
GSK-3 paralogs in the beta-catenin destruction complexes of tapeworms. *Development genes*
and evolution **11**, 1–14.

Nusse, R. and Clevers, H. (2017). Wnt/β-Catenin Signaling, Disease, and Emerging
Therapeutic Modalities. *Cell* **169**, 985–999.

Petersen, C. P. and Reddien, P. W. (2008). *Smed-βcatenin-1* is required for anteroposterior
blastema polarity in planarian regeneration. *Science* **319**, 327–330.

Petersen, C. P. and Reddien, P. W. (2009). Wnt signaling and the polarity of the primary body
axis. *Cell* **139**, 1056–1068.

Petersen, C. P. and Reddien, P. W. (2011). Polarized notum activation at wounds inhibits Wnt
function to promote planarian head regeneration. *Science* **332**, 852–855.

Pokutta, S. and Weis, W. I. (2000). Structure of the Dimerization and β-Catenin- Binding
Region of α-Catenin. *Mol. Cell* **5**, 533–543.

Rattner, A., Hsieh, J.-C., Smallwood, P. M., Gilbert, D. J., Copeland, N. G., Jenkins, N. A.
and Nathans, J. (1997). A family of secreted proteins contains homology to the cysteine-rich
ligand-binding domain of frizzled receptors. *Proc. Natl. Acad. Sci.* **94**, 2859–2863.

Read, C. P. (1967). Longevity of the tapeworm, *Hymenolepis diminuta*. *J Parasitol* **53**, 1055–6.

Reddien, P. W. (2018). The Cellular and Molecular Basis for Planarian Regeneration. *Cell* **175**,
327–345.

Reuter, H., März, M., Vogg, M. C., Eccles, D., Grífol-Boldú, L., Wehner, D., Owlarn, S.,
Adell, T., Weidinger, G. and Bartscherer, K. (2015). β-Catenin-Dependent Control of
Positional Information along the AP Body Axis in Planarians Involves a Teashirt Family
Member. *Cell reports* **10**, 253–265.

Riddiford, N. and Olson, P. D. (2011). Wnt gene loss in flatworms. *Development genes and*
evolution **221**, 187–197.

Rouhana, L., Weiss, J. A., Forsthoefel, D. J., Lee, H., King, R. S., Inoue, T., Shibata, N.,
Agata, K. and Newmark, P. A. (2013). RNA interference by feeding in vitro-synthesized
double-stranded RNA to planarians: methodology and dynamics. *Dev. Dyn.* **242**, 718–730.

Rozario, T. and Newmark, P. A. (2015). A confocal microscopy-based atlas of tissue
architecture in the tapeworm *Hymenolepis diminuta*. *Experimental parasitology* **158**, 31–41.

Rozario, T., Quinn, E. B., Wang, J., Davis, R. E. and Newmark, P. A. (2019). Region-
specific regulation of stem cell-driven regeneration in tapeworms. *eLife* **8**, e48958.

Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T.,
Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B., et al. (2012). Fiji: an open-source
platform for biological-image analysis. *Nat. Methods* **9**, 676–682.

Scimone, M. L., Cote, L. E., Rogers, T. and Reddien, P. W. (2016). Two FGFRL-Wnt circuits
organize the planarian anteroposterior axis. *eLife* **5**, 905.

Sikes, J. M. and Newmark, P. A. (2013). Restoration of anterior regeneration in a planarian
with limited regenerative ability. *Nature* **500**, 77–80.

Soria, C. L. D., Lee, J., Chong, T., Coghlan, A., Tracey, A., Young, M. D., Andrews, T.,
Hall, C., Ng, B. L., Rawlinson, K., et al. (2020). Single-cell atlas of the first intra-
mammalian developmental stage of the human parasite *Schistosoma mansoni*. *Nat. Commun.*
11, 6411.

Srivastava, M., Mazza-Curll, K. L., Wolfswinkel, J. C. van and Reddien, P. W. (2014).
Whole-Body Acoel Regeneration Is Controlled by Wnt and Bmp-Admp Signaling. *Current*
biology : CB **24**, 1107–1113.

Steinhart, Z. and Angers, S. (2018). Wnt signaling in development and tissue homeostasis.
Development **145**, dev146589.

Stückemann, T., Cleland, J. P., Werner, S., Vu, H. T.-K., Bayersdorf, R., Liu, S.-Y.,
Friedrich, B., Jülicher, F. and Rink, J. C. (2017). Antagonistic Self-Organizing Patterning
Systems Control Maintenance and Regeneration of the Anteroposterior Axis in Planarians.
Developmental cell **40**, 248-263.e4.

Su, H., Sureda-Gomez, M., Rabaneda-Lombarte, N., Gelabert, M., Xie, J., Wu, W. and
Adell, T. (2017). A C-terminally truncated form of β-catenin acts as a novel regulator of
Wnt/β-catenin signaling in planarians. *PLoS Genet.* **13**, e1007030.

Sureda-Gómez, M., Pascual-Carreras, E. and Adell, T. (2015). Posterior Wnts Have Distinct
Roles in Specification and Patterning of the Planarian Posterior Region. *International journal*
of molecular sciences **16**, 26543–26554.

Tewari, A. G., Owen, J. H., Petersen, C. P., Wagner, D. E. and Reddien, P. W. (2019). A
small set of conserved genes, including *sp5* and *Hox*, are activated by Wnt signaling in the
posterior of planarians and acoels. *PLoS Genet.* **15**, e1008401.

Umesono, Y., Tasaki, J., Nishimura, Y., Hrouda, M., Kawaguchi, E., Yazawa, S.,
Nishimura, O., Hosoda, K., Inoue, T. and Agata, K. (2013). The molecular logic for
planarian regeneration along the anterior-posterior axis. *Nature* **500**, 73–76.

Üren, A., Reichsman, F., Anest, V., Taylor, W. G., Muraiso, K., Bottaro, D. P.,
Cumberledge, S. and Rubin, J. S. (2000). Secreted Frizzled-related Protein-1 Binds Directly
to Wingless and Is a Biphasic Modulator of Wnt Signaling*. *J. Biol. Chem.* **275**, 4374–4382.

Valenta, T., Hausmann, G. and Basler, K. (2012). The many faces and functions of β-catenin.
EMBO J. **31**, 2714–2736.

Vandesompele, J., Preter, K. D., Pattyn, F., Poppe, B., Roy, N. V., Paepe, A. D. and
Speleman, F. (2002). Accurate normalization of real-time quantitative RT-PCR data by
geometric averaging of multiple internal control genes. *Genome Biol.* **3**, research0034.1.

Wang, S., Krinks, M., Lin, K., Luyten, F. P. and Moos, M. (1997). Frzb, a Secreted Protein
Expressed in the Spemann Organizer, Binds and Inhibits Wnt-8. *Cell* **88**, 757–766.

Wang, B., Lee, J., Li, P., Saberi, A., Yang, H., Liu, C., Zhao, M. and Newmark, P. A.

(2018). Stem cell heterogeneity drives the parasitic life cycle of *Schistosoma mansoni. eLife*

7, e35449.

 Figure legends

 Fig. 1. The germinative region (GR) regenerates after a single amputation. (A) DAPI-stained
 images of anterior fragments cut within the GR (0.5 mm) or outside of it (2 mm) at 15 dpa.
 Quantification of growth in length and number of proglottids regenerated at 15 dpa. N=3, n= 6-10
 per condition; t-test. (B) DAPI-stained image of 0.5 mm anterior fragment at 15 dpa showing
 regenerated GR. Yellow arrowhead marks the first proglottid. Quantification of GR lengths from
 A; t-test. (C) Quantification of proglottids regenerated and length every 2 days from 0.5 mm vs. 2
 mm anterior fragments. Change in proglottids was fitted with a simple linear regression and
 showed 14-15 proglottids regenerated per day. Change in length was fitted with a nonlinear curve
 as each proglottid increases in length. n=5-7 fragments per group per timepoint. (D) WISH for
 transcripts with A-P polarized expression patterns 3 days after acclimation to in vitro culture
 (uncut, n = 7-10 per transcript) then amputated at 0.5 mm and stained at 0 dpa (n= 7-8 per
 transcript) and 10 dpa (n= 5-13 per transcript). Asterisks mark pigmented debris that adhered to
 cut ends. Error bars= SD.

 Fig. 2. Serial amputation within the GR negatively affects regenerative ability. (A) Scheme
 for serial amputation within the GR. (B-C) GR lengths and percentage of GR retained compared
 to uncut whole worms at 0 dpa. n=7-10 per timepoint per experiment. (D) DAPI-stained
 regenerates 18 dpa for each cut. (E-F) Quantification of regeneration success rate and number of
 proglottids regenerated (n= 8-10 per timepoint per experiment). Error bars= SD; t-test.

 Fig. 3. The head inhibits proglottid regeneration and cell proliferation. (A) DAPI-stained
 regenerates from 0.5 mm amputations +/- head at 15 dpa. Quantification of proglottids regenerated
 from a representative experiment. n= 7; t-test. (B) Confocal micrographs from maximum intensity
 projections of +/- head worms at 3 dpa immunostained for neuronal marker SYN after a 1 hr pulse
 of F-*ara*-EdU to label proliferating cells. Quantification of the density of proliferating cells within
 the GR from a representative experiment. Each sample was normalized to the mean proliferation
 density in the corresponding +head group. n= 5-6 per group per timepoint; t-test. (C-D) Maximum
 intensity projections of worm anteriors after 1 hr pulse of F-*ara*-EdU either with the head (C) or -
 head 3 dpa (D). Yellow dotted lines represent 200 μm-wide regions used for quantification of

proliferation density. H= head. Quantification across 3 cohorts, n=11 and 13; one-way ANOVA

with Dunnett's multiple testing correction. Error bars= SD.

Fig. 4. Effect of head or head-adjacent amputations on proliferation and regeneration. (A)

Scheme for results shown in B-C and Fig. S2. Step 1: Four amputation schemes were tested: intact

head (+head), transverse cut through half the head while retaining the brain (½ head), full head

amputation (-head) and cut 0.5 mm from the anterior into the GR (-0.5 mm). Step 2: all worms

were amputated posteriorly to obtain 2 mm fragments. Step 3: fragments were grown in vitro for

14 days. Step 4-5: regenerates were collected and recut to obtain 2 mm anterior fragments that

were cultured in vitro while the posteriors were fixed and stained with DAPI to measure the

number of proglottids regenerated. (B) Quantification of proglottids regenerated. All samples were

normalized to the mean of the +head worms set to 100%. Bars indicate the means from 2-3

experiments for all samples after cut 1 and only for samples with visible GRs for cut 2-4. Fully

proglottidized worms without a GR are represented with a red dot. (C) Percentage of regenerates

that still maintained a visible GR from B. (D) After cut 4, proglottids regenerated from +head and

½ head samples from one experiment were plotted against the area of the head remaining measured

from DAPI stained worms. Linear regression fitted. (E-F) Quantification of F-*ara*-EdU+ cells

normalized to area within the GR 3 dpa. All samples were normalized to the mean of the +head

worms set to 100%. Means from 3-4 experiments analyzed with two-way ANOVA (mixed model)

and Tukey's multiple corrections test (E). All pooled data from analyzed with one-way ANOVA

compared to +head with Dunnett's multiple comparison test (n= 21-28). Error bars= SD.

Fig. 5. *βcat1* is a posterior-biased factor that is necessary to maintain the cycling stem cell

population in the GR. (A) Schematic and WISH of *βcat1* expression at the GR from a wild type

6-day-old adult. (B) DAPI-stained 2 mm anterior regenerates after single injection of dsRNA at 7

dpa. (C) Quantification of worm lengths after RNAi at 7 dpa. Bars show means and grey circles

represent individual worms. N= 3, n= 23-24 per group; one-way ANOVA with Dunnett's multiple

comparison test. (D) qRT-PCR from 3 RNAi experiments. Target expression normalized to two

endogenous controls and compared to *GFP* RNAi set at 1 (dotted line). (E) Representative

maximum intensity confocal projections at the GR stained after 1 hr F-*ara*-EdU uptake. Dotted

yellow lines are the worm outlines. Quantification of F-*ara*-EdU+ cells normalized to area within

an 800 µm-wide cropped region directly posterior to the head at 6-7 dpi. All samples were

normalized to the mean of *GFP* RNAi worms set to 100%. Bars show means and grey circles

represent individual worms. N= 3, n= 16-20 per group; one-way ANOVA with Dunnett's multiple

comparison test. (F) WISH for cycling stem cell markers *mcm2* and *lbr* at 6 dpi (N=3). Error bars=

SD.

Fig. 6. *sfrp* is an anterior-biased factor that regulates GR length. (A) Schematic and WISH of

sfrp expression at the GR from a wild type 6-day-old adult. (B) RNAi scheme for C-F. Double

injections of dsRNA were performed into the head and 2 mm anterior fragments were allowed to

regenerate for 10-12 days. (C) Representative DAPI-stained regenerates after RNAi. (D)

Quantification of proglottids regenerated after RNAi normalized to mean from *GFP* RNAi set to

100%. Bars show means and grey circles represent individual worms. N= 3, n= 25-26 per group;

one-way ANOVA with Dunnett's multiple comparison test. (E) Representative DAPI-stained

images showing GR lengths (between yellow arrowheads) after RNAi. (F) Quantification of GR

lengths from D. Bars show means and grey circles represent individual worms; one-way ANOVA

with Dunnett's multiple comparison test. (G) qRT-PCR for *sfrp* expression normalized to two

endogenous controls compared +head set at 1. N=3; t-test. Error bars= SD.

Fig. 7. How amputations at increasing distance from the head influence competence to

regenerate. (A) Regenerates from 0.5 mm anterior fragments were grown for 18 days then

amputated a second time at 1 mm (cut A). A subset was fixed to quantify GR length and a subset

was grown in vitro for 18 days. The remaining fragments were cut again to shave off ~20-25%

more GR (cut B) and the process repeated successively for cut C and cut D. Darkfield images

showing the live fragments before transfer to in vitro culture show that the wound site is

progressively closer to the head with successive cuts. (B) GR lengths of each fragment measured

from DAPI-stained 0 dpa fragments and represented as a percentage of the GR relative to uncut

worms from the same cohort. N=3, n= 3-9; one-way ANOVA. (C) Percent of fragments that

regenerated proglottids after 18 days from 3 experiments. Each experiment represented by the

same color in B and C. (D-F) Comparisons of worm length (D), growth fold change (E) and GR

length (F). N=3, n= 20 and 17; t-test. Error bars = SD.

Fig. 8. Models for regeneration in *H. diminuta*. (A) Working model for a minimal complement
of factors within the GR. X represents unknown factor(s) that directly/indirectly promote *βcat1*
activity. (B-D) Graphical representation using the working model to explain three scenarios
described in this study. For all: solid black activating or inhibiting arrows indicate activity in situ
whereas dotted grey arrows indicate the associated function without spatial reference.

Fig. S1. Proliferation response at wound site. Worms were amputated at 0.5 mm within the GR.
Fragments were analyzed at both the posterior-facing wound (Pfw) and anterior-facing wound
(Afw) at 1 dpa following a 1 hr pulse of F-*ara*-EdU. Proliferation density was quantified from a
box width of 100 μm from the cut site to capture the local proliferation response to amputation.
Quantification from one representative experiment is shown, n= 5-7; error bars= SD, one-way
ANOVA with Tukey's multiple testing correction.

Fig. S2. How head tissues influence regeneration. (A) Widefield fluorescent images of ½ head
fragments at 0 dpa stained with anti-SYN antibodies. Yellow arrowheads point to the
brain/cephalic ganglia. (B) GR lengths at the end of a representative serial amputation experiment.
n= 5-8; one-way ANOVA with Dunnett's multiple comparison test. (C) Quantification of
proglottids regenerated after cut 1, normalized to +head means. N=3; n=7-17; statistical
significance for each of 3 experiments calculated using one-way ANOVA and was the same for
all 3. (D) Quantification of proglottids regenerated normalized to +head means. N=2-3, n=5-16;
means (black circles) and statistical significance for each experiment using t-test are shown. Error
bars= SD.

Fig. S3. Domain analysis of *H. diminuta* βCAT1. (A) Sequence alignment of βCAT1 from four
flatworm species (*Hd: H. diminuta, Em: E. multilocularis, Sm: S. mansoni, Smed: S.*
mediterranea). The conserved DSGxxSxxx[S/T]xxxS motif for CKI/GSK-phosphorylation (red)
is present as are armadillo repeats (lilac). The boundaries of both domains are shown according to
S. mediterranea (Su et al., 2017). The shading indicates identical residues between 4 species (grey)
or 3 species (blue). The N- and C-termini are more divergent between all species. (B) Sequence
alignment of 26 amino acids covering a putative α-catenin binding domain previously reported
(Montagne et al., 2019). 10 critical residues for α-catenin binding (Aberle et al., 1996; Pokutta and

Weis, 2000) are marked by the symbol α (green). In reference to human CTNB1, dark grey shading

highlights identical residues. Chemically similar residues marked in light grey for aromatic (F, Y,

W, H), aliphatic (V, I, L), positively charged (R, K, H), negatively charged (D, E), polar but not

charged (N, Q, T, S, Y, C) and hydrophobic (V, I, L, M, A, F, P, W, G) residues, also in reference

to human CTNB1.

Fig. S4. Wild type WISH expression patterns for *βcat1* and *sfrp*. Representative micrographs

spanning different stages of development in 6-day-old worms with anterior facing left.

Fig. S5. Domain analysis of *H. diminuta* SFRP. BLASTp alignment of SFRPs from flatworms

and human SFRP-1,-2 and -5 as references (*Hd: H. diminuta, Em: E. multilocularis, Smed: S.*

mediterranea). The frizzled domain (orange) and netrin domain (blue) are marked at the most

inclusive positions reported for human SFRP1 using Interpro (Blum et al., 2020). Conserved

cysteines are highlighted in yellow.

Fig. S6. Phenotypes observed after *sfrp* RNAi. (A) Scheme for RNAi with dsRNA injections

throughout the GR. (B-C) Quantification of proglottid regeneration (B) and GR length (C) from

two independent experiments; one-way ANOVA with Dunnett's multiple comparison test. (D)

Quantification of F-*ara*-EdU+ cells normalized to area within an 800 μm-wide cropped region

directly posterior to the head or to the first proglottid. Combined from single and double injections

of dsRNA throughout the GR all from experiments with significant *sfrp* knockdown. All samples

were normalized to the mean of *GFP* RNAi worms set to 100%. Bars show means and grey circles

represent individual worms. N= 3, n= 21-23 per group; one-way ANOVA with Dunnett's multiple

comparison test. (E) qRT-PCR from 3 RNAi experiments where dsRNA was injected at the scolex

(Sc-biased) or throughout the GR. Expression of *sfrp* was normalized to two endogenous controls

compared to *GFP* RNAi set at 1 (dotted line). Error bars= SD.

Fig. S7. Comparison of regenerates from longest (cut A) vs. shortest (cut D) fragments after

re-amputation. (A) DAPI-stained regenerates at 18 dpa. (B) Proglottid regeneration at 18 dpa.

N=3, n= 20 and 17; t-test, error bars = SD.

Table S1. *βcatenin* paralogs. BLASTp analysis indicates that *H. diminuta* has three *βcatenin*

paralogs with orthologs in other flatworms.

Table S2. Primers and transcripts used in this study.

Table S3. All source data used in this study.

Figure 2

A

18 days

18 days

?

B GR length at 0 dpa

ns

length (mm)

1.0
0.8
0.6
0.4
0.2
0.0

cut 1   cut 2

C GR retained at 0 dpa

ns

Percentage (%)

cut 1   cut 2

D

cut 1

5 mm

cut 2

5 mm

E Regeneration success

Percentage (%)

cut 1   cut 2

F Proglottid regeneration

**

proglottids

cut 1   cut 2

Figure 3

Figure 4

Figure 5

A βcat1 · βcat1 · 0.5 mm

B *GFP* RNAi · *βcat1* RNAi · 5 mm · 5 mm

C Worm length
ns ***
length (mm)
UN · *GFP* RNAi · *βcat1* RNAi

D *βcat1* RNAi
Relative expression
βcat1 · axin1
target

E *GFP* RNAi · F-*ara*-EdU · *βcat1* RNAi · F-*ara*-EdU

Proliferation 6-7 dpi
Normalized percentage F-*ara*-EdU+ cells/mm²
ns **
UN · *GFP* RNAi · *βcat1* RNAi

F RNAi:
mcm2
GFP · 14/14
βcat1 · 16/16 · 200 µm

lbr
GFP · 15/15
βcat1 · 13/13 · 200 µm

Figure 6

A

sfrp

B

sfrp dsRNA → 2 mm → *sfrp* dsRNA → 10-12 dpa

C

GFP RNAi · *sfrp* RNAi

5 mm · 5 mm

D

Proglottid regeneration

% proglottids normalized to *GFP* RNAi

ns · *

UN · *GFP* RNAi · *sfrp* RNAi

E

GFP RNAi · *sfrp* RNAi

0.5 mm · 0.5 mm

F

GR length

GR length (mm)

ns · ****

UN · *GFP* RNAi · *sfrp* RNAi

G

sfrp at 3 dpa

Relative expression

*

+ head · - head

Figure 7

A 2nd amputation

cut A

cut B

cut C

cut D

1st amputation

18 days

0.5 mm

B % GR included

C Regeneration success

D Mean Body Length

E Growth Fold Change

F Mean GR length

Figure 8

A **Working Model**

sfrp ┈┈▷ GR

X?

high *βcat1*

proglottids

B **Scenario 1: head amputation**

GR inhibited

HIGH X

proglottids promoted

C **Scenario 2: cut "mid" GR**

x → FAIL

proglottids inhibited

D **Scenario 3: closest cut to head (cut D)**

NO X → establish *sfrp* → X

high *βcat1*

Amputation induced proliferation

Figure S2

A Rostellum amputated but suckers partially intact

SYN

Amputated through suckers

SYN

B GR after cut 4

****

ns

GR length (mm)

1.5

1.0

0.5

0.0

+head   1/2 head   -head

C After first amputation

****

*

Normalized proglottids to + head (%)

+ head   1/2 head   -head

D After subsequent amputations

Normalized proglottids to + head (%)

cut 2

**
***
*

+ head   1/2 head

cut 3

ns
ns

+ head   1/2 head

cut 4

*
*

+ head   1/2 head

A

This is a protein sequence alignment figure (Figure S3A) showing multiple sequence alignment of beta-catenin (Bcat1) homologs from four species:
- Hd_Bcat1_WMSIL1_LOCUS14475
- Em_Bcat1_EmuJ_001007700.1
- Sm_Bcat1_Smp_023550.1
- Smed_Bcat1_ABW79875.1

The alignment highlights the **CKI/GSK-3 phosphorylation sites** (labeled in red, with a red box around the DSGIQS...THTPS region around positions 75-95) and the **armadillo repeats** (labeled in purple below the alignment).

B

Figure S4

Figure S5

```
                                      20                    40                    60
Hd_sfrp_HD_tophat_13573  M- - - - - - - - -  - - - - LYLLLL  IFIAITSRGN  TVGRWDWNR-  - - -VNKTSVT  TAPFGGGAFG  43
Em_sfrp_EmuJ_000838700.1 M- - - - - - - - -  - - - - LPLVIL  FAIS-AAQAN  TVGRWDWHRG  GGAKNKTAVT  SSPFGGTSLS  46
Smed_sfrp1_ABY85212.1    MEMTKI - - - -  - - - - FPLSLL  LFISNV- - - -  - - - - - - - - - -  - - - - - - - - - -  - - - - - - - - - -  18
Human_SFRP1_sp|Q8N474.1| MGIGRSEGGR  RGAALGVLLA  LGAALLAVGS  - - - - - - - - - -  - - - - - - - - - -  - - - - - - - - - -  30
Human_SFRP5_sp|Q5T4F7.3| MRAAAAGGGV  RTAALALLL-  - GALHWAPAR  - - - - - - - - - -  - - - - - - - - - -  - - - - - - - - - -  28
Human_SFRP2_sp|Q96HF1.2| MLQGPG- - - -  - - - - -SLLLL  FLASHCCLGS  - - - - - - - - - -  - - - - - - - - - -  - - - - - - - - - -  21

                                      80                   100                   120
Hd_sfrp_HD_tophat_13573  RLTTADADMI  SNEPPQSPYF  SDWNRLVSG-  - YGSQ- RCYK  IPKELKLCHK  IGYDLMVLPN  100
Em_sfrp_EmuJ_000838700.1 RLTTADVHMI  SNEPPQSPYF  SDWNRLVSG-  - YGSQ- RCYK  IPRGMKLCHK  IGYDFMVLPN  103
Smed_sfrp1_ABY85212.1    - - - - - - - - - -  - - - - - -NSFI  GDWQAFQSG-  - YTID- QCYQ  IPDNFTLCSN  VGYRLMVLPN  59
Human_SFRP1_sp|Q8N474.1| - - - - - - - - - -  ASEYDYVSFQ  SDIGPYQSGR  FYTKPPQCVD  IPADLRLCHN  VGYKKMVLPN  80
Human_SFRP5_sp|Q5T4F7.3| CEEYDYYGWQ  AE- - PLH- GR  SYSKPPQCLD  IPADLPLCHT  VGYKRMRLPN  75
Human_SFRP2_sp|Q96HF1.2| - - - - - - - - - -  ARGLFLFG- Q  PDF- - - - - - -  SY- KRSNCKP  IPANLQLCHG  IEYQNMRLPN  62
```

Frizzled domain

```
                                     140                   160                   180
Hd_sfrp_HD_tophat_13573  SLEHEGMDEV  ITQSEVWLTL  VNLGCHDELE  RFLCSLYAPV  CIRGYHEKLI  QPCRELCESV  160
Em_sfrp_EmuJ_000838700.1 SLEHETLDEA  ITQSEVWLTL  VNLGCHDELK  RFLCSLYAPV  CINGYHEKLI  QPCRELCESV  163
Smed_sfrp1_ABY85212.1    YIQHESIEEA  AQHSKVWMGL  VNTQCHVDIK  KFICSLYAPV  CISNQRNQKV  PPCRELCVTV  119
Human_SFRP1_sp|Q8N474.1| LLEHETMAEV  KQQASSWVPL  LNKNCHAGTQ  VFLCSLFAPV  CLD- - - - RPI  YPCRWLCEAV  136
Human_SFRP5_sp|Q5T4F7.3| LLEHESLAEV  KQQASSWLPL  LAKRCHSDTQ  VFLCSLFAPV  CLD- - - - RPI  YPCRSLCEAV  131
Human_SFRP2_sp|Q96HF1.2| LLGHETMKEV  LEQAGAWIPL  VMKQCHPDTK  KFLCSLFAPV  CLDDL- DETI  QPCHSLCVQV  121

                                     200                   220                   240
Hd_sfrp_HD_tophat_13573  RAACLPTMTT  FGLGWPDIVK  CSKFPQAPQE  LCIPPNKQRN  KTVVLKPETR  - CSGCVDKPT  219
Em_sfrp_EmuJ_000838700.1 RAACLPTMTT  FGLGWPDIVK  CSKFPQAPQE  LCIPLNKHKN  KTIILNPETR  - CSGCIDRPT  222
Smed_sfrp1_ABY85212.1    KESCLPSMKM  YGFDWPTIMK  CSKFPQFKNS  LCIP- - - - -  K .VV- - PGKK  - CDLCMKIAS  170
Human_SFRP1_sp|Q8N474.1| RDSCEPVMQF  FGFYWPEMLK  CDKFP- - EGD  VCIAMTPPNA  TEASKPQGTT  VCPPCDNELK  194
Human_SFRP5_sp|Q5T4F7.3| RAGCAPLMEA  YGFPWPEMLH  CHKFPL- DND  LCIAVQFGHL  P- ATAPPVTK  ICAQCEMEHS  189
Human_SFRP2_sp|Q96HF1.2| KDRCAPVMSA  FGFPWPDMLE  CDRFPQ- DND  LCIPLASSDH  LLPATEEAPK  VCEACKNKND  180
```

Netrin domain

```
                                     260                   280                   300
Hd_sfrp_HD_tophat_13573  YES- AIGSFC  TADVAVRSKV  LDLIPVPSTK  GQNTTYRIRT  DGRGGIFKM-  - - - PKNVNKE  274
Em_sfrp_EmuJ_000838700.1 YES- AIGSFC  TADVVIRAKV  LDLVPTLPSS  GHNTTHRIRT  TGRVGAFKL-  - - - PKTLNSI  277
Smed_sfrp1_ABY85212.1    YEV- IANRFC  LSPIVIRAKI  KRIIP- - - TS  GN- - AIQIIL  QKKSKFLKF-  - - - QANFSQN  220
Human_SFRP1_sp|Q8N474.1| SEA- IIEHLC  ASEFALRMKI  KEV- - - - - KK  ENGDKKIV- -  PKKKKPLKLG  PIKKKDLKKL  246
Human_SFRP5_sp|Q5T4F7.3| ADG- LMEQMC  SSDFVVKMRI  KEI- - - - - KI  ENGDRKLIGA  QKKKKLLKPG  PLKRKDTKRL  243
Human_SFRP2_sp|Q96HF1.2| DDNDIMETLC  KNDFALKIKV  KEI- - - - - TY  INRDTKIILE  TKSKTIYKLN  GVSERDLKKS  235

                                     320                   340                   360
Hd_sfrp_HD_tophat_13573  A- NLDFRMTC  DCPIIKAAMT  RSKGPGRWLL  MGKLNEDGRS  VTVLHLSQPS  RQNKGIKRAI  333
Em_sfrp_EmuJ_000838700.1 V- NLEFEMEC  DCPIIRAALT  RQKGPGRWLM  MGKLNDDRRT  VTVQHISQPS  RQNAGIKRAI  336
Smed_sfrp1_ABY85212.1    A- - LELNLKC  NCTNLKFKQR  ALKG- - RWII  MAQIDSDNKA  V- VNFISKWK  RKKAEFKHSM  275
Human_SFRP1_sp|Q8N474.1| VLYLKNGADC  PCHQLDNLS-  - - - - - HHFLI  MGR- - KVKSQ  YLLTAIHKWD  KKNKEFKNFM  298
Human_SFRP5_sp|Q5T4F7.3| VLHMKNGAGC  PCPQLDSLA-  - - - - -GSFLV  MGR- - KVDGQ  LLLMAVYRWD  KKNKEMKFAV  295
Human_SFRP2_sp|Q96HF1.2| VLWLKDSLQC  TCEEMNDIN-  - - - - - APYLV  MGQ- - KQGGE  LVITSVKRWQ  KGQREFKRIS  287

                                     380                   400                   420
Hd_sfrp_HD_tophat_13573  KEMRHQPDTL  CKVDLTRSSP  VVHPRTVAGR  PAANGAQISQ  EQKQQHQQLQ  PRRNGSNLSK  393
Em_sfrp_EmuJ_000838700.1 RDIRRRPDSL  CKVDLMHPNP  VAHSRAISSL  SSPSSMQSLR  PRSHQSRRYR  P- - - - -HLSP  391
Smed_sfrp1_ABY85212.1    KIIQNFGHIV  CKSKLPDGIS  L- RERYAAYL  KMNNQLKYSN  ERKLQTRKFR  - - - - - - - - -K  325
Human_SFRP1_sp|Q8N474.1| KKMKNHE- - -  CPTFQSVF- -  - - - -K- - - - -  - - - - - - - - - -  - - - - - - - - - -  - - - - - - - - - -  314
Human_SFRP5_sp|Q5T4F7.3| KFMFSYP- - -  CSLYYPFFYG  AAEPH- - - - -  - - - - - - - - - -  - - - - - - - - - -  - - - - - - - - - -  317
Human_SFRP2_sp|Q96HF1.2| RSIRKLQ- - -  C- - - - - - - - -  - - - - - - - - - -  - - - - - - - - - -  - - - - - - - - - -  - - - - - - - - - -  295

                                     440                   460                   480
Hd_sfrp_HD_tophat_13573  EERRRRRQLR  RRRNE- - - AG  AQQQQHPSNV  TLSSTNSQES  QQPVNDMPIG  VNTQHPYNQQ  450
Em_sfrp_EmuJ_000838700.1 EERRRR- QLR  RRRNHPRGSS  PEAQQRAANV  TPSRKPPQLQ  QQPINDIPIL  TPTPRPFSHH  450
Smed_sfrp1_ABY85212.1    QKSRK- - - - -  - - - - - - - - -S  VKPNNRSAGI  S- - - - - - - - -  - - - - - - - - - -  - - - - - - - - - -  342
Human_SFRP1_sp|Q8N474.1| - - - - - - - - - -  - - - - - - - - - -  - - - - - - - - - -  - - - - - - - - - -  - - - - - - - - - -  - - - - - - - - - -  314
Human_SFRP5_sp|Q5T4F7.3| - - - - - - - - - -  - - - - - - - - - -  - - - - - - - - - -  - - - - - - - - - -  - - - - - - - - - -  - - - - - - - - - -  317
Human_SFRP2_sp|Q96HF1.2| - - - - - - - - - -  - - - - - - - - - -  - - - - - - - - - -  - - - - - - - - - -  - - - - - - - - - -  - - - - - - - - - -  295
```

A *sfrp* dsRNA → 2 mm → *sfrp* dsRNA → 10-12 dpa

injection zone

injection zone

B Proglottid Regeneration

Exp 1   Exp 2

Normalized proglottids to *GFP* RNAi (%)

UN   *GFP* RNAi   *sfrp* RNAi

C GR length

Exp 1   Exp 2

GR length (mm)

UN   *GFP* RNAi   *sfrp* RNAi

D Proliferation in GR

Normalized percentage F-*ara*-EdU+ cells/mm²

UN   *GFP* RNAi   *sfrp* RNAi

E *sfrp* knockdown efficacy

Relative expression

Sc-biased   GR

A

cut A → 18 dpa

cut D → 18 dpa

5 mm

5 mm

B

Proglottid regeneration

proglottids

**

A   D