Shared potential metabolism trends in degraded soils and type 2 diabetes gut microbiomes

Craig Liddicoat[1,*], Bart A. Eijkelkamp[1], Timothy R. Cavagnaro[1], Jake M. Robinson[1], Kiri Joy Wallace[2], Andrew D. Barnes[2], Garth Harmsworth[3], Damien J. Keating[4], Robert A. Edwards[1] and Martin F. Breed[1]

[1]College of Science and Engineering, Flinders University, Bedford Park, South Australia, Australia

[2]Te Aka Mātuatua – School of Science, University of Waikato, Hamilton, Aotearoa New Zealand

[3]Manaaki Whenua – Landcare Research, Palmerston North, Aotearoa New Zealand

[4]College of Medicine and Public Health, Flinders University, Bedford Park, South Australia, Australia

*Corresponding author: Craig Liddicoat; craig.liddicoat@flinders.edu.au

ORCIDs: CL: 0000-0002-4812-7524, BAE: 0000-0003-0179-8977, TRC: 0000-0002-9922-5677, JMR: 0000-0001-8108-3271; KJW: 0000-0003-1236-6846, ADB: 0000-0002-6499-381X, GH: 0000-0002-7041-1847, DJK: 0000-0002-0154-8305, RAE: 0000-0001-8383-8949, MFB: 0000-0001-7810-9696

Competing Interest Statement: The authors declare no competing interest.

Keywords: Healthy microbiome, type 2 diabetes, metagenomics, compound processing potential, soil biodiversity, human health, metabolism, branched-chain fatty acids

Abstract

Microbiome-mediated impacts of global change on human metabolic health remain understudied. Type 2 diabetes (T2D) is shaped by nutrition, host and environmental factors, with rapidly increasing global prevalence. Soil microbiomes shift with ecosystem degradation and may influence human metabolism through altering food quality and gut microbiomes, including metabolite exposures without requiring colonization. Here, we investigated functional overlaps between soil microbiomes across ecosystem degradation gradients (USA, Australia) and gut microbiomes in T2D versus normal health (Sweden, China). We developed a method to translate metagenomic functional pathways to potential metabolism of biochemical compounds. In silico trend analyses revealed consistent shifts relevant to energy harvesting and management. Both T2D gut microbiomes and degraded soil microbiomes exhibited increased potential metabolism for sugars and decreased potential metabolism for lignin and monomethyl branched-chain fatty acids. Our findings suggest ecosystem degradation may contribute to T2D pathogenesis through nutrient-depleted food and/or adverse shaping of gut microbiome functional capacities.

Introduction

Microbiomes connect ecosystems and humans under One Health[1, 2]. Yet, microbiome-mediated impacts of accelerating global change[3] on human metabolic health have received little attention. Type 2 diabetes (T2D) is a major chronic multifaceted metabolic disease, characterized by insulin resistance and high blood sugar, and linked to host, nutritional and environmental factors[4, 5, 6]. As the health and economic burden of T2D rise rapidly worldwide[7, 8], understanding its complex pathogenesis is critical. The human gut microbiome is integral to our metabolism[9, 10] with a vast enzymatic repertoire that complements human digestive capabilities[11] and produces metabolites that modulate appetite, gut motility, hormonal signaling and gene expression[10]. Gut microbiomes shift in T2D and with metformin treatment[12], while fecal microbiome transplantations can improve clinical outcomes[13, 14]. Given that long-term diet and environmental exposures have major influences on the gut microbiome[15, 16], which is intricately linked to metabolism and T2D risk, there is a pressing need to identify upstream nutritional and environmental factors that influence this crucial interface to human metabolic health.

Beyond diet and lifestyle factors, environmental exposures are increasingly recognized as contributors to T2D risk[5], including shifts from rural to urban areas[17, 18, 19, 20]. Microbiomes are a hypothesized link between environments and gut-associated health[21, 22]. Soils are the dominant reservoir of microbial life[23, 24], and exposure to diverse soil microbiomes (for example, via direct contact, food and airborne dust), especially from a young age, can shape gut microbiomes[16, 21, 25, 26, 27]. In wild primates, gut communities vary with environmental heterogeneity, especially soil differences[28]. Soil microbiomes also shape food quality by driving nutrient cycling and gathering[29], incorporating microbial metabolites into plants[30], and influencing gut microbiota of ruminants[31]. Beyond the microbes per se, microbiomes also comprise important metabolites (for example, biochemical compounds, constituent molecules) and potentially transferable genetic material[32] with diverse structural, functional and nutritional roles[9, 30, 33]. In soils and the gut, a vast range of metabolites with

varying bioactivities may be accumulated, depleted and/or transferred via microbial activity and

turnover[9, 34]. This means microbiomes can extend their influence via metabolites and necromass

without requiring direct colonization. While the survival of soil microbes in the gut may be low or

uncertain, the shaping of gut microbiomes by soil microbiome exposures is well-recognized[1, 35].

Moreover, soil microbiome composition and functional capacities may shift in generalizable ways with

ecosystem degradation, for example, via urbanization and land-use change, with potential to impact

human health[36, 37, 38, 39]. Together, these phenomena create pathways whereby degraded environments

may adversely shape soil microbiomes, food quality and gut microbiomes, and potentially contribute to

metabolic dysfunction in T2D.

In ongoing work to define 'healthy microbiomes', approaches based on community-scale

functional capacity (rather than taxonomy) should provide greater insight[40, 41]. Notably, many processes

involve assemblages using shared resources in extracellular space[9]. Striving for deeper functional

insights and interpretability, metabolite- or compound-oriented approaches[38] aim to move beyond

metabolic pathways towards measures related to specific chemical compounds (such as carbohydrates,

lipids, proteins, often linked to microbial feedstocks or by-products; Supplementary Fig. 1).

Compound-scale insights should offer more foundational information (as compounds can underpin

multiple functional pathways), benefit from known links to biochemistry and host health, and may

point to new mechanistic hypotheses and therapeutic targets. However, existing metabolite prediction

frameworks[42, 43, 44] rely on organism-specific and environment-specific approaches unsuited to highly

biodiverse, poorly characterized and varying environments, such as soils[38]. Therefore, there is a need

for new, compound-oriented methods to examine normal versus aberrant functioning in comparative

analyses across soil and gut microbiomes.

In this integrative in silico study, we explored potential links between ecosystem degradation

and metabolic alterations in the T2D gut by examining functional overlaps between soil microbiomes

across ecosystem degradation gradients and gut microbiomes in T2D versus normal health (datasets are

described in Supplementary Table 1). We translated metagenomic functional gene-associated

biochemical pathways to 'compound processing potential' (CPP, %), reflecting potential metabolism of

individual compounds. While not measuring compounds directly, our method quantified the capacity of

metagenomes to metabolize (or process) them. Reactants and products were not differentiated, as

compounds can be inputs and outputs of various processes. Examining (1) 30 selected compounds

relevant to soil and gut health (Supplementary Table 2) and (2) via exhaustive compound-wide trend

analyses, we identified compounds relevant to energy harvesting and management that shared potential

metabolism (i.e., CPP) trends. In a staged approach, we first examined soils from a forest restoration

chronosequence[45] (from the United States, US; spanning degraded post-mining, reforested, and

unmined reference sites), then progressively considered overlapping CPP trends in T2D versus control

cohorts from Sweden[46] and then China[47]. Lastly, we sought corroboration of overlapping trends in a

'disturbed versus natural' soil dataset from the Australian Microbiome Initiative (AMI[48], described in

ref.[38]; see Supplementary Table 1).

Results

Trends with ecosystem degradation

For the 30 selected compounds we assessed trends with ecosystem degradation in the chronosequence

soils (US) (Supplementary Table 2; Supplementary Figs. 2-31) and found increased potential

metabolism for ($P < 0.001$) D-fructose, sucrose, L-arabinose, D-glucose, xylose and chitin; also ($P <$

0.01) galactose, mannose and hydrogen sulfide; and ($P \leq 0.05$) glucosamine and arabinoxylan

(hemicellulose). We found decreased potential metabolism with ecosystem degradation for ($P < 0.001$)

lignin, acetate and carbon dioxide; also ($P < 0.01$) glycogen, butyrate, trimethylamine N-oxide and

trimethylamine; and ($P \leq 0.05$) propionate and methane.

Compound-wide trend analyses identified 2,122 compounds with significant CPP trends ($n =$

757 decreased, $n = 1,365$ increased with ecosystem degradation; compound and trend data are in

Supplementary Material files). Carbon-containing compounds were mapped using chemical formula

ratios oxygen:carbon (O:C), hydrogen:carbon (H:C) and nitrogen:carbon (N:C) (Fig. 1) to visualize

energetically and chemically similar compounds that trend together (see example plot that visualizes all

carbon-containing compounds within a single sample in Supplementary Fig. 32).

Trends in type 2 diabetes

For the 30 selected compounds, we found consistent trends in the Swedish (SWE) and Chinese (CHN)

cohort datasets (Supplementary Table 2; Supplementary Figs. 2-31, 33) for increased potential

metabolism in T2D for D-fructose (SWE: $P \leq 0.05$, CHN: $P < 0.01$), L-arabinose (SWE: $P \leq 0.05$,

CHN: $P < 0.001$), and amylopectin (SWE and CHN: $P \leq 0.05$); and decreased potential metabolism in

T2D for lignin (SWE and CHN: $P \leq 0.05$).

Numerous trends were only found in one T2D cohort or the other. Increased potential

metabolism was found in T2D in the Swedish cohort for sucrose and starch ($P \leq 0.05$), and in the

Chinese cohort for mannose ($P < 0.001$); glucose, galactose and xylose (all $P < 0.01$). Decreased

potential metabolism was found in T2D in the Chinese cohort for acetate, propionate, butyrate, carbon

dioxide, hydrogen sulfide, indole (all $P < 0.001$); menaquinone (Vitamin K2), glucosamine ($P < 0.01$);

methane, p-cresol and serotonin (all $P \leq 0.05$). We found conflicting potential metabolism trends for

ammonia in T2D, which increased in Swedish ($P < 0.01$) but decreased in Chinese ($P < 0.001$) cohorts.

Fig. 1 | Compound-wide trend analysis results from restoration chronosequence soil (US) metagenomes. Points represent carbon-containing compounds with decreased[†] (aqua, n = 720) or increased[†] (red, n = 1,255) potential metabolism with ecosystem degradation ([†]trends with ecosystem degradation are interpreted as the opposite to displayed trends with revegetation age). Each compound is mapped via its chemical formula O:C, H:C and N:C elemental ratios, where clusters of points indicate energetically and chemically similar compounds. The O:C by H:C panels display a limited extent to emphasize data-rich regions. N:C ratios are displayed as three panels to represent the third dimension. Rectangular zones denoting compound classes offer a visual guide only, adapted from[49, 50]. Chrosequence samples[45] spanned planting ages 6, 12, 22, 31 years and unmined (UM), n = groups of 3.

The compound-wide trend analyses in T2D Swedish and Chinese cohorts found substantially more compounds with decreased potential metabolism in the Chinese cohort (compounds and trend data are in Supplementary Material files; carbon-containing compounds are visualized in Supplementary Figs. 34-35). In the T2D (SWE) data, 1269 compounds had trending potential

metabolism (n = 911 decreased, *n* = 358 increased). For the T2D (CHN) data a total of 4292

compounds showed trending potential metabolism (*n* = 3984 decreased, *n* = 308 increased). We note,

to explore the most interesting (potentially significant) results, the T2D compound-wide associations

were not subject to *P*-adjustment. From visual inspection, the chemical ratio mapping (Supplementary

Figs. 34-35) showed limited commonality in CPP trends in the vicinity of carbohydrates and

widespread reduced CPP in the T2D (CHN) dataset.

Consistent trends with ecosystem degradation and T2D

In the 30 selected compounds, we found consistent CPP trends with ecosystem degradation and T2D in

the chronosequence soils (US), T2D (SWE), and T2D (CHN) datasets, for increased potential

metabolism in D-fructose and L-arabinose, and decreased potential metabolism in lignin

(Supplementary Table 2).

From overlapping compound-wide trend results from the chronosequence soils (US) and T2D

(SWE) datasets, based on matching ModelSEED compound identifiers, 304 compounds displayed CPP

trends in both datasets (detailed in Supplementary Material files), from which 286 carbon-containing

compounds were visualized in Fig. 2. There, we observed two prominent clusters along straight lines in

chemical mapping space: 'Sugars' (L-arabinose, melibiose, manninotriose, D-lyxulose, sucrose,

D-fructose) that showed increased CPP with T2D (SWE) and ecosystem degradation (US)

(Supplementary Table 3); and 'BCFA-ACPs' comprising 35 compounds dominated by branched-chain

fatty acid (BCFA) – acyl-carrier proteins (ACPs) (Supplementary Table 4), that showed reduced CPP

with T2D (SWE) and ecosystem degradation (US). Importantly, the BCFA-ACPs were all involved in

fatty acid biosynthesis and dominated by monomethyl unsaturated (-enoyl) BCFAs that were not found

elsewhere in the datasets beyond this trend group. The Sugars and BCFA-ACPs are described in detail

in Supplementary Material Supporting Information, Supplementary Figs. 36-40, and trends shown in

Fig. 3.

Fig. 2 | Overlapping compound-wide trend results from restoration chronosequence soils (US)

and Swedish (SWE) T2D cohort. Chemical ratio mapping of carbon-containing compounds with

overlapping potential metabolism trends from Fig. 1 and Supplementary Fig. 34). Clusters of each

color represent energetically and chemically similar compounds occurring in each of the four trend

groups shown. Total (and carbon-containing) compounds in the four trend groups are: decreased

potential metabolism with T2D (SWE) and ecosystem degradation, n = 70 (n = 66 red points);

increased potential metabolism with T2D (SWE) and ecosystem degradation, n = 89 (n = 81 green

points); decreased potential metabolism with T2D (SWE) and ecosystem restoration, n = 95 (n = 90

orange points); increased potential metabolism with T2D (SWE) and ecosystem restoration, n = 50 (n =

49 blue points). Expanded views of 'Sugars' and 'BCFA-ACPs' (i.e., branched-chain fatty acid - acyl-

carrier proteins) are in Supplementary Figs. 36, 38. The plot format is further described in Fig. 1.

Fig. 3 | Key potential metabolism trends. Sugars (a-c), lignin (d-f), and branched-chain fatty acid –

acyl carrier proteins (BCFA-ACPs, listed in Supplementary Tables 4-5) (g-i), within the restoration

chronosequence (US; spanning planting ages 6, 12, 22, 31 years and unmined, UM, n = groups of 3) (a,

d, g), T2D (Swedish cohort, SWE; T2D met- = type 2 diabetes not treated with metformin, n = 33;

Normal = normal healthy subjects, n = 43) (b, e, h), and T2D (Chinese cohort, CHN; T2D met- n = 30;

Normal n = 52) (c, f, i). Sugars and BCFA-ACPs were groupings identified via matching trends in T2D

(SWE) and with ecosystem degradation (Fig. 2). Sugars were graphed as the sum of compound

processing potential (CPP, %) for six compounds (L-arabinose, melibiose, D-fructose, sucrose,

manninotriose and D-lyxulose), however subsequent testing showed only three of these sugars (L-

arabinose, melibiose, D-fructose) displayed individual trends in T2D (Chinese cohort, CHN)

(Supplementary Table 3). Lignin CPP values were $\log_{10}$-transformed to improve display.

Further matching of parallel compound-wide association trends across chronosequence soils

(US), T2D (SWE) and T2D (CHN) datasets identified 76 compounds, with $n = 60$ decreased, $n = 16$

increased CPP with ecosystem degradation T2D (Fig. 4; see Supplementary Material files). Even after

filtering due to intersecting the T2D (CHN) dataset, the increased CPP for sugars: L-arabinose,

melibiose, D-fructose still remained as trending, together with 6-phosphosucrose, an intermediate in

sucrose biosynthesis. The signals of decreased potential metabolism for lignin and all 35 BCFA-ACPs

also remained.

Lastly, corroboration of the majority of observed overlapping trends was found in the disturbed

versus natural AMI soils dataset. In disturbed AMI soils, we observed increased potential metabolism

of L-arabinose, D-fructose, melibiose and 6-phosphosucrose, and decreased potential metabolism of

lignin and all 35 of the BCFA-ACPs (all $P = {<}0.001$; Supplementary Fig. 41). From the 76 compounds

with matching potential metabolism trends with ecosystem degradation (US), T2D (SWE) and T2D

(CHN), a total of 65 compounds (85.5 %) displayed shared potential metabolism trends in the AMI

disturbed soils (Supplementary Table 5).

Fig. 4 | Relative importance of compounds with consistent trends across restoration chronosequence soils (US) and T2D (Swedish and Chinese) datasets. Points indicate increased or decreased potential metabolism (or compound processing potential, CPP) with ecosystem degradation (soil) and in T2D (gut) microbiomes. Compounds including branched-chain fatty acid-acyl carrier proteins (BCFA-ACPs), sugars that feature widely across many functional themes and pathways, and lignin are highlighted in the plot. Trends with ecosystem degradation are interpreted as the inverse of restoration (based on Kendall's tau ordinal correlation with revegetation age classes). Measures of significance (y-axis) from the Wilcoxon difference testing of CPP in T2D (versus normal), and the number of level 3 (L3) subsystems (functional themes) linked to each compound, are calculated as

mean values from the Swedish and Chinese cohorts. Details for all $n = 76$ compounds are in

Supplementary Table 5.

Discussion

We identified biologically relevant overlapping trends in the potential metabolism of compounds

within soil and human gut microbiomes, supporting the hypothesis that nutrient-depleted food (i.e., low

BCFA and lignin content, high sugar content) and/or excessive microbiome exposures associated with

degraded ecosystem soil may contribute to the development of type 2 diabetes. These linkages warrant

further investigation due to the rising burden of T2D and increasing global change impacts on soil

microbiomes – which underpin food quality and shape human gut microbiomes. An alternative or

complementary explanation may be that similar ecological stresses and dysfunction in soil and gut

ecosystems may be driving convergence in microbiome functional traits. Our compound-oriented

approach facilitates linkages to a wealth of published literature that links chemical compounds with

metabolism, health and wider biological processes in humans and the environment. Also, mapping of

compounds into the bioenergetically informed O:C, H:C and N:C coordinate space helped to reveal

meaningful patterns of simultaneously trending compounds (i.e., sugars, BCFA-ACPs) with soil-

ecosystem quality and T2D versus normal health. Our results align with the One Health framework,

revealing plausible functional linkages between ecosystem condition and human metabolic health, via

the microbiome. This work contributes necessary research towards finding synergies and co-benefits in

simultaneously advancing UN Sustainable Development Goals, including improving theoretical models

of nature–human health relationships[51, 52].

Do branched-chain fatty acids connect healthy ecosystems and healthy people?

Our results suggest that microbiomes in degraded ecosystems and T2D have reduced capacity for

monomethyl BCFA (mmBCFA) biosynthesis. This may reflect a shift in the biochemical composition

of bacterial assemblages as mmBCFAs are key components that enhance bacterial membrane fluidity,

and mmBCFA synthesis capacity enables membrane fluidity-related adaptive responses for survival

and proliferation through environmental stresses[53, 54]. BCFA levels vary among bacterial taxa, being

commonly higher in Gram-positive bacteria, present in some Gram-negative species, and especially

high in *Bacillus* and *Bifidobacterium*[53, 55]. Within ecosystems, BCFAs are present in lipid/waxy plant

compounds, animal (including human) tissues and fluids, ruminant products (milk, cheese, meat), fish,

organs, blood serum, skin, hair, adipose tissue, colostrum, breast milk, *vernix caseosa* of newborns, and

at very high levels in saprotrophic fungi such as *Conidiobolus* spp. found in soil and decaying organic

matter[54, 56, 57]. Therefore, variation in BCFA synthesis capacity could result from changes in microbial

taxonomic composition as ecosystems degrade.

Monomethyl BCFAs possess multiple beneficial bioactivities for human health and link

bacterial and human physiology[56]. Beneficial bioactivities of mmBCFA include lipid-lowering,

reducing metabolic disorder risk, maintaining normal insulin-producing β cell function and insulin

sensitivity, anti-inflammation, cytotoxicity to cancer cells and regulation of development[54, 55, 56].

Bacterially-derived mmBCFAs facilitate direct biochemical uptake and cross-domain metabolism

between the gut microbiome and animal hosts[58, 59, 60]. In the nematode *Caenorhabditis elegans* and

mammalian tissue culture cells, mmBCFA have been shown to be critical for triggering the host

mechanistic target of rapamycin complex 1 (mTORC1)[59, 60], a central regulator of cell growth and

metabolism[61]. The host mTORC1 responds to environmental signals (for example, amino acid

availability, growth factors, energy levels, stress) to coordinate cellular status. However, mTORC1

dysregulation is closely linked to diabetes, cancer, and neurodegenerative disorders[61]. Ref. [62] reported

high capacities for human fetal intestinal epithelial cells (enterocytes) to incorporate long-chain BCFA

into membrane phospholipids. Following uptake by human intestinal epithelial cells, further

interactions between BCFA and host endogenous BCFA synthesis and metabolism may occur[54, 56],

including post-translational modifications to cell proteins, which alter intestinal cell function[63]. While

precise mechanisms are uncertain, it is possible that mmBCFA in the gut microbiome may modulate

host cellular metabolism and physiological functioning via these interactions.

Microbial sources play a dominant role in the supply of health-associated mmBCFA[55, 56]. Key

inputs to BCFA synthesis include branched-chain amino acids which can only be made in bacteria,

plants, and fungi, but not in animals[64]; or shorter BCFA, which can be elongated. Microbes (for

example, rumen bacteria) also contribute to BCFA synthesis in animals[56]. Therefore, both diet and gut

microbiome synthesis provide major sources of BCFA to humans[55, 56]. Within soil and gut

microbiomes, the interchange of BCFA between substrates and microbes can be expected because fatty

acids are a high-value energy source, and bacteria assimilate environmental BCFA into their bodies[65,

66]. In the gut, the constant release of mmBCFAs will occur from cell membranes with bacterial

turnover from competitive interactions[67] and secretion of fatty acids and their derivatives[58]. A portion

of these gut bacteria may be supplied from external ecosystem exposures[21, 22], while BCFA-

synthesizing bacteria may be further shaped by diet and other factors (for example, high BCFA-content

Bifidobacterium spp. proliferate in infants receiving breast milk[68]).

As ecosystems degrade, there may be several reasons why BCFA-synthesizing organisms

decline. Ecosystem degradation is widely associated with reductions in soil organic matter[3, 69, 70, 71].

Indeed, the more degraded chronosequence soils we analyzed had lower soil organic content than

unmined sites[72]. Soil organic matter comprises a continuum of progressively decomposed organic

compounds, commonly including amino acids and fatty acids[73]. Amino acids are released from proteins

in the soil at rates correlated with soil organic matter pools and soil protein concentrations[74], and

increasing with ecosystem succession[75]. Natural decomposition and recycling networks of saprotrophic

fungi (including high BCFA-content spp.) may also be degraded, likely reducing BCFA content and

turnover in soil microbiomes. Diverse plant litter contributes to increased soil carbon, higher microbial

abundance, diversity, decomposition and turnover, which strengthens over time[76]. Taken together, these

influences may reduce access to and recycling of both branched-chain amino acids and fatty acids for

soil microbiomes in degraded ecosystems, thereby potentially reducing ambient human exposure to

health-promoting microbial mmBCFA synthesis capacity (Fig. 5). Degraded agricultural soils might

also produce crops and ruminant products (for example, meat, dairy) with reduced mmBCFA content,

that impact gut microbiomes via diet. However, these ideas require further investigation.

Fig. 5 | Hypothesis linking ecosystem degradation and type 2 diabetes. Intact natural ecosystems

typically have greater biodiversity, woody plant debris (including lignin), fungal recycling and soil

organic matter (SOM) pools. SOM releases branched-chain amino acids (BCAA) which are used to

make precious monomethyl branched-chain fatty acids (mmBCFA) used in bacterial cell membranes,

stored by fungi, and needed to regulate animal cell metabolism via the mechanistic target of rapamycin

complex 1 (mTORC1) among other pathways. With ecosystem degradation, soil organic matter

decomposes into simple sugars. Degraded soil microbiomes have increased capacity to metabolize

simple sugars, decreased capacity to metabolize lignin (potentially impacting production of short-chain

fatty acids, SCFAs), and decreased mmBCFA synthesis capacity (potentially linked to mmBCFA

content in microbial bodies, necromass and the wider ecosystem). If people consume nutrient-depleted

food (low mmBCFA and lignin content, high sugar content) and/or are increasingly exposed to soil

microbiomes associated with degraded ecosystems, it may contribute to metabolic shifts in the gut

microbiome consistent with type 2 diabetes.

Identified compounds linked to energy harvesting

We found that potential metabolism of sugars L-arabinose, D-fructose, melibiose and the intermediate

compound 6-phosphosucrose each increased with ecosystem degradation and T2D. Additionally, we

found the potential metabolism of sucrose, glucose, galactose, xylose and mannose increased in more

degraded ecosystems. Sugars are the most abundant organic compounds in the biosphere and are key

carbon and energy sources for soil microbes[77]. Sugars in soil can either be derived from primary

sources (for example, decomposition of plant litter, rhizodeposits, root exudates) or secondary sources

(for example, microorganisms and their residues)[77]. Previous work suggests that galactose, glucose,

mannose, arabinose and xylose are chief components of sugar residues found within soil organic

matter[78]. Our detection of increased potential metabolism of simple sugars (largely monosaccharides

and disaccharides) likely reflects enhanced microbial decomposition of soil organic matter within more

degraded ecosystems[69].

In the gut microbiome, carbohydrate metabolism contributes up to 10% of the host's overall

energy extraction and is recognized as an important contributor to insulin resistance and the

development of obesity and prediabetes[79]. Ref. [79] found increased gut microbiome metabolism of

monosaccharides, including fructose, galactose, mannose and xylose, significantly correlated with

insulin resistance. Interestingly, in soils and humans, enhanced sugar metabolism is also associated

with more opportunistic bacteria[80]. Increased relative abundances of opportunists (including fast-

growing, potential pathogens) have been associated with disturbance in urban greenspace soils[37] and in

human-altered land uses[81]. If microbiome components promoting enhanced sugar metabolism are

transferred from degraded ecosystems to the human gut, they may foster opportunistic or pathogen-like

traits – potentially contributing to increased energy harvest and metabolic imbalances associated with

T2D.

Identified compounds linked to energy management

We found altered potential metabolism for compounds with plausible links to energy storage via fatty

acids, gut hormone secretion and appetite regulation that decreased in both degraded ecosystems and

T2D. This raises the question whether nutrient-depleted foods and/or excessive exposures to degraded

ecosystem soils that are deficient in potential metabolism of compounds involved in energy

management may shape gut microbiomes that are also deficient in this way. We discuss above how gut

microbiome mmBCFA content and biosynthesis capacity might modulate host cellular metabolism via

central mTORC1 regulation, alteration of enterocyte phospholipid composition and function, and/or

supply of BCFA for further host endogenous fatty acid synthesis and BCFA-mediated metabolism.

Dietary fiber, such as lignin, is known to decrease plasma ghrelin, known as the hunger

hormone, and increase other key gut regulatory hormones: cholecystokinin, glucagon-like-peptide-1

(GLP-1), and peptide YY[82]. Short-chain fatty acids (SCFAs) are produced when gut microbes ferment

non-digestible fibers, and these SCFAs bind to receptors in the colon, leading to the production of

hormones that regulate food intake, body weight, and energy metabolism[10, 82]. Similar trends to lignin

appeared for sinapyl alcohol and coumaryl alcohol (both precursors to lignin) in the compound-wide

results spanning all datasets. Potential metabolism for the dietary fiber pectin also decreased in T2D

(significantly in the Swedish cohort, and marginally in the Chinese cohort), but not in degraded

ecosystems. We searched for oligosaccharides (for example, fructan, inulin, galactan) but did not detect

these compounds in our datasets. Additional compounds that we identified remain to be explored

(Supplementary Table S5).

Disentangling dietary and environmental influences on T2D gut microbiomes

Altered potential metabolism for compounds detected within gut microbiomes may arise from dietary

and environmental exposures, gut environment and functions, neurological influences, among other

factors[83]. Where we found matching trends across all case studies (for example, L-arabinose, D-

fructose, melibiose, BCFA-ACPs, lignin), this may be consistent with an alignment between

environmental exposures and dietary inputs. In the scenario where compounds consistently showed in

both T2D cohorts, but not in the restoration chronosequence soils, this suggests dietary effects alone.

For example, increased potential metabolism results for amylopectin suggests that excessive

consumption of starch-rich foods may contribute to T2D. For many compounds we found alignment in

potential metabolism between the ecosystem condition and T2D gut (Chinese cohort), but not the T2D

gut (Swedish cohort) – this may point to greater environmental exposures and detrimental health

influence in some population groups. Yet another contrast occurred in the case of ammonia, where

potential metabolism trends were opposing in the T2D cohorts, in the absence of any ecosystem

condition effect, which suggests influences from dietary exposures alone associating with T2D. We

raise further noteworthy implications for community connection to soils and ecosystems, and study

limitations in the Supplementary Material, Supporting Information.

Conclusions

We found overlapping trends in the potential metabolism of compounds with beneficial bioactivities, and involved in energy harvesting and management, within separate model case study datasets of soil-ecosystem degradation from the United States and Australia, and T2D gut microbiomes from Swedish and Chinese cohorts. Our results support existing studies implicating, at least in part, roles for microbiomes, sugars, lignin (dietary fiber) and mmBCFA in metabolism and T2D. Additionally, our findings could inform new T2D interventions; and suggest that degraded soil microbiomes may contribute to metabolic imbalances found in T2D via nutrient-depleted food (for example, low mmBCFA and lignin content, high sugar content) and/or excessive environmental exposures associated with degraded ecosystems. Our method used to examine microbiome potential metabolism at the resolution of individual compounds has raised new biologically interpretable and testable hypotheses regarding potential links between soil, ecosystem and T2D gut health that warrant further transdisciplinary investigation.

Methods

Case study data

We examined previously published soil and human gut metagenome case study datasets (detailed in Supplementary Table 1). Restoration chronosequence[45] soil samples spanned planting ages 6, 12, 22, 31 years and unmined (UM) with n = groups of 3. T2D datasets were from Swedish[46] (T2D n = 33, Normal n = 43) and Chinese[47] (T2D n = 30, Normal n = 52) cohorts. We used metadata from ref. 12 to exclude subjects treated with metformin or diagnosed with prediabetes. From the Chinese cohort, we only included subjects >50 years old to compare with the mature age (69-72 years) Swedish subjects. Australian Microbiome Initiative (AMI) disturbed (n = 29) versus natural (n = 55) soils were surface depth (0–10 cm), from the temperate climate zone, with 7.5–45 % clay content (i.e., avoiding very

sandy and very clayey soils) as described in ref. 38. Disturbed soils were from agricultural land uses

(cereals, cotton, horticulture, irrigated pasture), while natural soils were from conservation and national

park land uses.

Metagenome functional profiling via compound processing potential (CPP)

Our CPP analysis approach used a purpose-built bioinformatics workflow (Fig. 6) that redistributes

values of functional potential relative abundances (%), from the scale of functions (i.e., biochemical

pathways that often comprise multiple linked chemical reactions), down to the level of individual

chemical compounds. These compounds are conceptually participating in a weighted array of

intersecting chemical reactions that represent the collective functional capacity of a metagenome.

Fig. 6 | Overview of the compound processing potential (CPP) analysis workflow. Metagenomic

functional potential pathway relative abundances were translated to quantify hypothetical potential

metabolism at the scale of individual compounds. CPP % values for carbon-containing compounds can

be mapped using chemical formula ratios to visualize clusters of bioenergetically similar compounds

that trend together.

Measures of CPP were derived via steps outlined in Fig. 6. Firstly, following ref. 38, using high performance computing resources[84, 85] raw metagenomic sequences were accessed, trimmed for quality control, and human case study sequences that mapped using Bowtie2[86] to the human genome (GRCh38.p14/hg38) were excluded (low levels were observed; Supplementary Material, Supporting Information). Then good quality read 1 sequences were classified to SEED subsystem[87] functional annotations and relative abundances using SUPER-FOCUS software[88]. Every SUPER-FOCUS function (output row) was translated to one or more corresponding chemical reaction(s) using a purpose-built R-script algorithm based on ModelSEED database lookup tables (from https://github.com/ModelSEED/ModelSEEDDatabase; accessed 10 Aug 2022). Linking of functions (and sub-functions where present) to chemical reactions was based on either: full matching of functional hierarchies (using subsystem-class, -subclass, -name and -role); detection of Enzyme Commission (EC) number; or matching of SUPER-FOCUS function name within ModelSEED lookup tables for reactions (reaction name or alias), subsystems (role), or reaction-pathways (external reaction name). Reactions were then linked to corresponding compounds using ModelSEED database tables ('reactions.tsv', 'compounds.tsv').

Advancing from the previous method used in ref. 38, here we derived CPP measures at the resolution of individual compounds (all software parameters and supporting R code for CPP calculations are available at: https://github.com/liddic/cpp3d_t2d):

(1) Function level relative abundances from SUPER-FOCUS (which sum to 100% in each sample) were allocated to compounds by dividing values equally across all sub-functions and reactions, and dividing again across all corresponding compounds, with weightings to account for stoichiometric coefficients for each compound within each reaction, with equal consideration to reactants and products.

(2) Relative abundances (%) for respective unique compounds (which may arise across multiple

reactions and functions) were then collated and summed to provide a summary CPP relative

abundance (%) for each compound. (Here, conversion to CPP data format successfully

recovered typically 65 to 70% of the initial 100% function potential per sample; Supplementary

Table 1).

(3) For mapping visualization of CPP patterns for carbon-containing compounds, elemental ratios

of O:C, H:C, and N:C were calculated. Such chemical mapping frameworks have been used to

display and interpret groupings of energetically and chemically similar compounds[38, 49].

Statistical Analysis

We used a staged approach to explore overlapping potential metabolism (i.e., CPP) trends in the soil

and T2D gut samples. We progressively examined the restoration chronosequence soils (US), then

Swedish and Chinese T2D cohorts, and lastly the AMI soils to corroborate findings. Further detail is in

Supplementary Material and at https://github.com/liddic/cpp3d_t2d:

(1) *Thirty selected compounds* were initially explored for CPP trends, chosen due to their

relevance to soil and/or gut health (Supplementary Table 2). In chrosequence soils (US), CPP

trends were evaluated using Kendall's tau rank correlation tests based on ordinal revegetation

age classes. We interpreted CPP trends with ecosystem degradation as the inverse of calculated

trends with increasing revegetation age (as informed by previous work showing highly

disturbed post-mining soil microbiomes progressively shift in their structure with ecosystem

restoration towards intact natural condition[36]). In T2D versus normal health, CPP trends (i.e.,

differences) were evaluated using non-parametric Wilcoxon rank-sum (Mann-Whitney) tests,

due to often differing variance of groups (T2D vs. normal) and to avoid normality assumptions.

All data were plotted to assist interpretation.

(2) ***Compound-wide association tests*** were also performed to exhaustively scan CPP trends for all compounds. This used the same approach as above, with further steps to automate analyses. In correlation testing of chrosequence soils, compounds that did not have relative abundance data for at least a quarter of the total sample number were excluded due to low data. In this chrosequence soil dataset, strong CPP trends were discovered, and only significant results based on Benjamini and Hochberg adjusted P-values were considered further. In the subsequent compound-wide association analysis of T2D gut samples, compounds were only included if at least one or other of the T2D or normal groups contained at least 50% non-zero CPP values. Testing was based on $\log_{10}$-transformed CPP % values, after zero replacement with small positive values equal to half the minimum non-zero CPP % of all samples. One-sided (directional) difference tests were based on whether the median value in T2D was greater or less than the median value for normal healthy subjects. To retain and explore the most interesting (potentially significant) results from the compound-wide T2D analyses, these data were not subject to P-adjustment for multiple testing.

(3) ***Overlap analysis.*** Compounds with CPP % trends in both the restoration chronosequence (US) and T2D (SWE) case studies were identified, based on matching ModelSEED compound identifiers. Our focus was to highlight compounds that displayed consistent increased or decreased CPP values with ecosystem degradation and T2D.

(4) ***Visualising compound-wide association test results.*** Compounds with CPP % trends in the restoration chronosequence (US) and T2D (SWE) gut microbiomes, and overlaps between the two, were mapped using their chemical formula elemental ratios of O:C, H:C, and N:C. From this mapping space, prominent localized clusters of consistently trending compounds (i.e., with ecosystem degradation and T2D) were identified and further analyzed to determine trends in their collective (summed) CPP % and to interpret their possible metabolic importance.

(5) *Trend consolidation and validation.* The above steps were repeated with the second T2D

(CHN) cohort, which refined the list of compounds with parallel CPP trends with ecosystem

degradation (US) and in both T2D datasets. Trend and contextual information for these

compounds was reported and visualized to represent the significance ($-\log_{10}[P\text{-value}]$; as used

in genome-wide association study Manhattan plots) of association with ecosystem degradation

and T2D, and the number of functional themes (level 3 subsystems) they are linked to. Here,

data for T2D were visualized using mean results from the Swedish and Chinese cohort datasets.

To corroborate parallel CPP trends from the restoration chronosequence (US) and both T2D

datasets, a focused analysis of CPP trends only for previously identified compounds was

performed using the disturbed versus natural AMI soils.

(6) *Beta diversity ordinations.* Supplementary analyses using the restoration chronosequence (US)

and T2D Swedish cohort dataset were performed to examine community-scale compositional

patterns and differences between sample groups and analysis approaches, comparing

conventional functional potential (%) data (termed 'functions') versus CPP (%) data. This

analysis used principal coordinate analysis (PCoA) ordinations based on Bray-Curtis distances,

and testing for differences between groups via permutational analysis of variance

(PERMANOVA) and beta dispersion testing for homogeneity of groupings using the vegan R

package[89]. These data are presented in the Supplementary Material, Supporting Information and

Supplementary Fig. 42.

Data availability

The datasets used are available from NCBI Sequence Read Archive (accessions PRJEB1786 and

PRJNA422434), MG-RAST (project mgp16379), and soil metagenomes from the Australian

Microbiome Initiative data portal (https://data.bioplatforms.com/organization/australian-microbiome).

Code availability

Code used to support this study is available from: https://github.com/liddic/cpp3d_t2d .

Acknowledgments

We acknowledge and thank the authors of previously published metagenomics datasets used here, which enabled comparison of samples spanning human and environmental health. This includes the Australian Microbiome initiative which was supported by funding from Bioplatforms Australia and the Integrated Marine Observing System (IMOS) through the Australian Government's National Collaborative Research Infrastructure Strategy (NCRIS), Parks Australia through the Bush Blitz program funded by the Australian Government and BHP, and the CSIRO. We acknowledge Indigenous peoples and the traditional custodians of the lands on which the case study datasets originate, and our authors live and work. This work was supported by the People, Cities and Nature research program (Aotearoa New Zealand Ministry of Business, Innovation and Employment, grant UOWX2101).

References

1. Muhummed, A.M., Lanker, K.C., Yersin, S., Zinsstag, J. & Vonaesch, P. One Health, One Microbiome. *Microbiome* **13**, 216 (2025).

2. Banerjee, S. & van der Heijden, M.G.A. Soil microbiomes and one health. *Nature Reviews Microbiology* **21**, 6-20 (2022).

3. IPBES. The IPBES assessment report on land degradation and restoration. Bonn, Germany; 2018.

4. Pearson, E.R. Type 2 diabetes: a multifaceted disease. *Diabetologia* **62**, 1107-1112 (2019).

5. Beulens, J.W.J. *et al.* Environmental risk factors of type 2 diabetes—an exposome approach. *Diabetologia* **65**, 263-274 (2022).

6. Lu, X. *et al.* Type 2 diabetes mellitus in adults: pathogenesis, prevention and therapy. *Signal Transduction and Targeted Therapy* **9**, 262 (2024).

7. Ong, K.L. *et al.* Global, regional, and national burden of diabetes from 1990 to 2021, with projections of prevalence to 2050: a systematic analysis for the Global Burden of Disease Study 2021. *The Lancet* **402**, 203-234 (2023).

8. Sun, H. *et al.* IDF Diabetes Atlas: Global, regional and country-level diabetes prevalence estimates for 2021 and projections for 2045. *Diabetes Res Clin Pract* **183**, 109119 (2022).

9. Daisley, B.A. *et al.* Emerging connections between gut microbiome bioenergetics and chronic metabolic diseases. *Cell Reports* **37**, 110087 (2021).

10. Chao, J., Coleman, R.A., Keating, D.J. & Martin, A.M. Gut Microbiome Regulation of Gut Hormone Secretion. *Endocrinology* **166** (2025).

11. Rowland, I. *et al.* Gut microbiota functions: metabolism of nutrients and other food components. *European Journal of Nutrition* **57**, 1-24 (2018).

12. Forslund, K. *et al.* Disentangling type 2 diabetes and metformin treatment signatures in the human gut microbiota. *Nature* **528**, 262-266 (2015).

13. Chen, L. *et al.* Fecal microbiota transplantation ameliorates type 2 diabetes via metabolic remodeling of the gut microbiota in db/db mice. *BMJ Open Diabetes Research & Care* **11**, e003282 (2023).

14. Wu, Z. *et al.* Fecal microbiota transplantation reverses insulin resistance in type 2 diabetes: A randomized, controlled, prospective study. *Frontiers in Cellular and Infection Microbiology* **Volume 12 - 2022** (2023).

15. Gilbert, J.A. *et al.* Current understanding of the human microbiome. *Nature Medicine* **24**, 392 (2018).

583   16.   Wopereis, H., Oozeer, R., Knipping, K., Belzer, C. & Knol, J. The first thousand days – intestinal
584   microbiology of early life: establishing a symbiosis. *Pediatric Allergy and Immunology* **25**, 428-438
585   (2014).
586

587   17.   Attard, S.M. *et al.* Multilevel examination of diabetes in modernising China: what elements of
588   urbanisation are most associated with diabetes? *Diabetologia* **55**, 3182-3192 (2012).
589

590   18.   Cheema, A., Adeloye, D., Sidhu, S., Sridhar, D. & Chan, K.Y. Urbanization and prevalence of type
591   2 diabetes in Southern Asia: A systematic analysis. *J Glob Health* **4**, 010404 (2014).
592

593   19.   Ruiz-Alejos, A. *et al.* Addressing the impact of urban exposure on the incidence of type 2 diabetes
594   mellitus: The PERU MIGRANT Study. *Scientific Reports* **8**, 5512 (2018).
595

596   20.   Gassasse, Z., Smith, D., Finer, S. & Gallo, V. Association between urbanisation and type 2 diabetes:
597   an ecological study. *BMJ Global Health* **2**, e000473 (2017).
598

599   21.   Roslund, M.I. *et al.* Long-term biodiversity intervention shapes health-associated commensal
600   microbiota among urban day-care children. *Environment International* **157**, 106811 (2021).
601

602   22.   Rothschild, D. *et al.* Environment dominates over host genetics in shaping human gut microbiota.
603   *Nature* **555**, 210 (2018).
604

605   23.   Anthony, M.A., Bender, S.F. & van der Heijden, M.G.A. Enumerating soil biodiversity.
606   *Proceedings of the National Academy of Sciences* **120**, e2304663120 (2023).
607

608   24.   Blakemore, R.J. Biodiversity restated: > 99.9% of global species in Soil Biota. *ZooKeys* **1224**, 283-
609   316 (2025).
610

611   25.   Zhou, G.-L. *et al.* Exposure to greenspaces sourced soils improves mice gut microbiota.
612   *Environmental Pollution* **362**, 124931 (2024).
613

614   26.   Liu, W. *et al.* Exposure to soil environments during earlier life stages is distinguishable in the gut
615   microbiome of adult mice. *Gut Microbes* **13**, 1-13 (2021).
616

617   27.   Liddicoat, C. *et al.* Naturally-diverse airborne environmental microbial exposures modulate the gut
618   microbiome and may provide anxiolytic benefits in mice. *Science of The Total Environment* **701**,
619   134684 (2020).
620

621   28.   Grieneisen, L.E. *et al.* Genes, geology and germs: gut microbiota across a primate hybrid zone are
622   explained by site soil properties, not host species. *Proceedings of the Royal Society B: Biological
623   Sciences* **286**, 20190431 (2019).
624

625   29.   Cao, T., Luo, Y., Shi, M., Tian, X. & Kuzyakov, Y. Microbial interactions for nutrient acquisition
626   in soil: Miners, scavengers, and carriers. *Soil Biology and Biochemistry* **188**, 109215 (2024).
627

628   30.   Adamczyk, B. How do terrestrial plants access high molecular mass organic nitrogen, and why does
629   it matter for soil organic matter stabilization? *Plant and Soil* (2021).
630

631 31. Qin, W., Song, P. & Zhang, S. Seasonal and Soil Microbiota Effects on the Adaptive Strategies of
632   Wild Goitered Gazelles Based on the Gut Microbiota. *Frontiers in Microbiology* **Volume 13 - 2022**
633   (2022).
634
635 32. Peng, H. *et al.* Longitudinal gut microbiota tracking reveals the dynamics of horizontal gene
636   transfer. *Nature Communications* **16**, 11543 (2025).
637
638 33. Bindels, L.B. *et al.* A blueprint for contemporary studies of microbiomes. *Microbiome* **13**, 95
639   (2025).
640
641 34. Finley, B.K. *et al.* Unraveling the influence of microbial necromass on subsurface microbiomes:
642   metabolite utilization and community dynamics. *ISME Communications* **5** (2025).
643
644 35. Roslund, M.I., Laitinen, O.H. & Sinkkonen, A. Scoping review on soil microbiome and gut
645   health—Are soil microorganisms missing from the planetary health plate? *People and Nature* **6**,
646   1078-1095 (2024).
647
648 36. Liddicoat, C. *et al.* Next generation restoration metrics: Using soil eDNA bacterial community data
649   to measure trajectories towards rehabilitation targets. *Journal of Environmental Management* **310**,
650   114748 (2022).
651
652 37. Delgado-Baquerizo, M. *et al.* Global homogenization of the structure and function in the soil
653   microbiome of urban greenspaces. *Science Advances* **7**, eabg5809 (2021).
654
655 38. Liddicoat, C. *et al.* Bioenergetic mapping of 'healthy microbiomes' via compound processing
656   potential imprinted in gut and soil metagenomes. *Science of The Total Environment* **940**, 173543
657   (2024).
658
659 39. Sun, X. *et al.* Unforeseen high continental-scale soil microbiome homogenization in urban
660   greenspaces. *Nature Cities* **2**, 759-769 (2025).
661
662 40. Van Hul, M. *et al.* What defines a healthy gut microbiome? *Gut* **73**, 1893 (2024).
663
664 41. Fierer, N., Wood, S.A. & Bueno de Mesquita, C.P. How microbes can, and cannot, be used to assess
665   soil health. *Soil Biology and Biochemistry* **153**, 108111 (2021).
666
667 42. Neveu, V., Nicolas, G., Amara, A., Salek, R.M. & Scalbert, A. The human microbial exposome:
668   expanding the Exposome-Explorer database with gut microbial metabolites. *Scientific Reports* **13**,
669   1946 (2023).
670
671 43. Mallick, H. *et al.* Predictive metabolomic profiling of microbial communities using amplicon or
672   metagenomic sequences. *Nature Communications* **10**, 3136 (2019).
673
674 44. Garza, D.R., van Verk, M.C., Huynen, M.A. & Dutilh, B.E. Towards predicting the environmental
675   metabolome from metagenomics with a mechanistic model. *Nature Microbiology* **3**, 456-460
676   (2018).
677
678 45. Sun, S. & Badgley, B.D. Changes in microbial functional genes within the soil metagenome during
679   forest ecosystem restoration. *Soil Biology and Biochemistry* **135**, 163-172 (2019).
680

681   46.   Karlsson, F.H. *et al.* Gut metagenome in European women with normal, impaired and diabetic
682   glucose control. *Nature* **498**, 99-103 (2013).
683
684   47.   Qin, J. *et al.* A metagenome-wide association study of gut microbiota in type 2 diabetes. *Nature*
685   **490**, 55-60 (2012).
686
687   48.   Bissett, A. *et al.* Introducing BASE: the Biomes of Australian Soil Environments soil microbial
688   diversity database. *GigaScience* **5**, 21 (2016).
689
690   49.   Rivas-Ubach, A. *et al.* Moving beyond the van Krevelen Diagram: A New Stoichiometric Approach
691   for Compound Classification in Organisms. *Analytical Chemistry* **90**, 6152-6160 (2018).
692
693   50.   Wu, X. *et al.* Microbial Interactions With Dissolved Organic Matter Drive Carbon Dynamics and
694   Community Succession. *Frontiers in Microbiology* **9** (2018).
695
696   51.   Fisher, B. *et al.* Can nature deliver on the sustainable development goals? *The Lancet Planetary*
697   *Health* **3**, e112-e113 (2019).
698
699   52.   Marselle, M.R. *et al.* Pathways linking biodiversity to human health: A conceptual framework.
700   *Environment International* **150**, 106420 (2021).
701
702   53.   Yoon, Y., Lee, H., Lee, S., Kim, S. & Choi, K.-H. Membrane fluidity-related adaptive response
703   mechanisms of foodborne bacterial pathogens under environmental stresses. *Food Research*
704   *International* **72**, 25-36 (2015).
705
706   54.   Gozdzik, P., Magkos, F., Sledzinski, T. & Mika, A. Monomethyl branched-chain fatty acids: Health
707   effects and biological mechanisms. *Progress in Lipid Research* **90**, 101226 (2023).
708
709   55.   Yang, S.-j., Yu, X.-k. & Zuo, Q. Branched- Chain Fatty Acids and Obesity: A Narrative Review.
710   *Nutrition Reviews* (2025).
711
712   56.   Mao, S. *et al.* Branched-Long-Chain Monomethyl Fatty Acids: Are They Hidden Gems? *Journal of*
713   *Agricultural and Food Chemistry* **71**, 18674-18684 (2023).
714
715   57.   Maheshwari, G. *et al.* Branched-Chain Fatty Acids as Mediators of the Activation of Hepatic
716   Peroxisome Proliferator-Activated Receptor Alpha by a Fungal Lipid Extract. *Biomolecules* **10**
717   (2020).
718
719   58.   Brown, E.M., Clardy, J. & Xavier, R.J. Gut microbiome lipid metabolism and its impact on host
720   physiology. *Cell Host & Microbe* **31**, 173-186 (2023).
721
722   59.   Zhu, M. *et al.* Monomethyl branched-chain fatty acid mediates amino acid sensing upstream of
723   mTORC1. *Developmental Cell* **56**, 2692-2702.e2695 (2021).
724
725   60.   Zhu, H., Shen, H., Sewell, A.K., Kniazeva, M. & Han, M. A novel sphingolipid-TORC1 pathway
726   critically promotes postembryonic development in Caenorhabditis elegans. *eLife* **2**, e00429 (2013).
727
728   61.   Takahara, T., Amemiya, Y., Sugiyama, R., Maki, M. & Shibata, H. Amino acid-dependent control
729   of mTORC1 signaling: a variety of regulatory modes. *Journal of Biomedical Science* **27**, 87 (2020).
730

731 62. Liu, L. *et al.* Human fetal intestinal epithelial cells metabolize and incorporate branched chain fatty
732   acids in a structure specific manner. *Prostaglandins, Leukotrienes and Essential Fatty Acids* **116**,
733   32-39 (2017).
734

735 63. Ezzine, C. *et al.* Fatty acids produced by the gut microbiota dampen host inflammatory responses by
736   modulating intestinal SUMOylation. *Gut Microbes* **14**, 2108280 (2022).
737

738 64. Neinast, M., Murashige, D. & Arany, Z. Branched Chain Amino Acids. *Annual Review of*
739   *Physiology* **81**, 139-164 (2019).
740

741 65. Sen, S. *et al.* Growth-Environment Dependent Modulation of Staphylococcus aureus Branched-
742   Chain to Straight-Chain Fatty Acid Ratio and Incorporation of Unsaturated Fatty Acids. *PLOS ONE*
743   **11**, e0165300 (2016).
744

745 66. Yao, J. & Rock, C.O. How Bacterial Pathogens Eat Host Lipids: Implications for the Development
746   of Fatty Acid Synthesis Therapeutics*. *Journal of Biological Chemistry* **290**, 5940-5946 (2015).
747

748 67. Booth, S.C., Smith, W.P.J. & Foster, K.R. The evolution of short- and long-range weapons for
749   bacterial competition. *Nature Ecology & Evolution* **7**, 2080-2091 (2023).
750

751 68. Stewart, C.J. *et al.* Temporal development of the gut microbiome in early childhood from the
752   TEDDY study. *Nature* **562**, 583-588 (2018).
753

754 69. Chen, S. *et al.* Plant diversity enhances productivity and soil carbon storage. *Proceedings of the*
755   *National Academy of Sciences* **115**, 4027 (2018).
756

757 70. Kooch, Y., Amani, M. & Abedi, M. The effect of shrublands degradation intensity on soil organic
758   matter-associated properties in a semi-arid ecosystem. *Science of The Total Environment* **853**,
759   158664 (2022).
760

761 71. Cunningham, S.C. *et al.* Reforestation with native mixed-species plantings in a temperate
762   continental climate effectively sequesters and stabilizes carbon within decades. *Global Change*
763   *Biology* **21**, 1552-1566 (2015).
764

765 72. Avera, B.N., Strahm, B.D., Burger, J.A. & Zipper, C.E. Development of ecosystem structure and
766   function on reforested surface-mined lands in the Central Appalachian Coal Basin of the United
767   States. *New Forests* **46**, 683-702 (2015).
768

769 73. Song, Y. *et al.* Soil metabolomics: Deciphering underground metabolic webs in terrestrial
770   ecosystems. *Eco-Environment & Health* **3**, 227-237 (2024).
771

772 74. Näsholm, T., Kielland, K. & Ganeteg, U. Uptake of organic nitrogen by plants. *New Phytologist*
773   **182**, 31-48 (2009).
774

775 75. Kielland, K., McFarland, J.W., Ruess, R.W. & Olson, K. Rapid Cycling of Organic Nitrogen in
776   Taiga Forest Ecosystems. *Ecosystems* **10**, 360-368 (2007).
777

778 76. Zhang, W.-P. *et al.* Plant litter strengthens positive biodiversity–ecosystem functioning relationships
779   over time. *Trends in Ecology & Evolution* **38**, 473-484 (2023).
780

781   77.   Gunina, A. & Kuzyakov, Y. Sugars in soil and sweets for microorganisms: Review of origin,
782   content, composition and fate. *Soil Biology and Biochemistry* **90**, 87-100 (2015).
783
784   78.   Nagar, B.R. Free Monosaccharides in Soil Organic Matter. *Nature* **194**, 896-897 (1962).
785
786   79.   Takeuchi, T. *et al.* Gut microbial carbohydrate metabolism contributes to insulin resistance. *Nature*
787   **621**, 389-395 (2023).
788
789   80.   Le Bouguénec, C. & Schouler, C. Sugar metabolism, an additional virulence factor in
790   enterobacteria. *International Journal of Medical Microbiology* **301**, 1-6 (2011).
791
792   81.   Liddicoat, C. *et al.* Can bacterial indicators of a grassy woodland restoration inform ecosystem
793   assessment and microbiota-mediated human health? *Environment International* **129**, 105-117
794   (2019).
795
796   82.   Bulsiewicz, W.J. The Importance of Dietary Fiber for Metabolic Health. *American Journal of*
797   *Lifestyle Medicine* **17**, 639-648 (2023).
798
799   83.   Cryan, J.F. *et al.* The Microbiota-Gut-Brain Axis. *Physiological Reviews* **99**, 1877-2013 (2019).
800
801   84.   Flinders_University. DeepThought (HPC). Retrieved from
802   https://doi.org/10.25957/FLINDERS.HPC.DEEPTHOUGHT. 2021.
803
804   85.   Pawsey_Supercomputing_Research_Centre. Setonix Supercomputer. Perth, Western Australia;
805   2023.
806
807   86.   Langmead, B. & Salzberg, S.L. Fast gapped-read alignment with Bowtie 2. *Nature Methods* **9**, 357-
808   359 (2012).
809
810   87.   Overbeek, R. *et al.* The SEED and the Rapid Annotation of microbial genomes using Subsystems
811   Technology (RAST). *Nucleic Acids Research* **42**, D206-D214 (2013).
812
813   88.   Silva, G.G.Z., Green, K.T., Dutilh, B.E. & Edwards, R.A. SUPER-FOCUS: a tool for agile
814   functional analysis of shotgun metagenomic data. *Bioinformatics* **32**, 354-361 (2015).
815
816   89.   Oksanen, J. *et al.* vegan: Community Ecology Package. R package version 2.5-7.; 2020.
817
818