Widely Used Variants of the Farquhar-von-Caemmerer-Berry Model Can Cause Errors in Parameter Estimation

Edward B. Lochocki[1] and Justin M. McGrath[1,2,3*]

[1]Carl R. Woese Institute for Genomic Biology, University of Illinois, Urbana–Champaign, Urbana, IL, USA

[2]Plant Biology Department, University of Illinois, Urbana–Champaign, Urbana, IL, USA

[3]USDA ARS Global Change and Photosynthesis Research Unit, Urbana, IL, USA

*Corresponding author's email address: justin.mcgrath@usda.gov

SUMMARY STATEMENT: The Farquhar-von-Caemmerer-Berry model of net CO_2 assimilation plays a key role in plant research, but many publications use variants of the model that differ from the original version and can potentially introduce errors in estimates of photosynthetic parameters.

ABSTRACT: The Farquhar-von-Caemmerer-Berry (FvCB) model is the most widely-used mechanistic model of C_3 net CO_2 assimilation, and it plays a significant role in plant physiology, ecology, climate science, and Earth system modeling. The model's equations are not written consistently across publications, and a common variant is to use the minimum of assimilation rates rather than the minimum of carboxylation rates. This variant, which we call the "min-*A* variant," makes different predictions than the original FvCB model, yet appears in approximately half of highly-cited publications and software tools that use the FvCB model. Another concern is that although leaf biochemistry restricts the range of CO_2 partial pressures where limitations due to triose phosphate utilization (TPU) can occur, this restriction is commonly omitted from the model's equations. Among other potential issues, these variations can introduce errors exceeding 20% when estimating photosynthetic parameter values from CO_2 response curves. It is therefore important to be aware of this source of error when fitting

25   the model, to avoid using the min-*A* variant, and to include the biochemically-derived CO_2

26   threshold for TPU limitations.

27   **KEYWORDS:** Fitting *A*-C_i curves, photosynthesis at low CO_2 concentrations, triose phosphate

28   utilization (TPU), ribulose 1,5-bisphosphate (RuBP) depletion

1. INTRODUCTION

The Farquhar-von-Caemmerer-Berry (FvCB) model for C_3 net CO_2 assimilation is a cornerstone

of modern plant biology, ecology, and climate science, having been highly successful in

explaining experimental measurements and making predictions at scales ranging from single

cells to the entire globe (Farquhar et al., 2001; Von Caemmerer, 2013). Although commonly

referred to by the names of the authors of a key 1980 publication (Farquhar et al., 1980), the

FvCB model is nonetheless built on decades of research that predates 1980, and it has been

improved by other researchers since its initial description (Yin et al., 2021). It is a mechanistic

model based on a simplified version of the light-dependent reactions, the Calvin-Benson-

Bassham (CBB) cycle, and the photorespiratory cycle, and it consists of a small set of equations

for predicting net CO_2 assimilation rates (Farquhar et al., 1980; Farquhar and von Caemmerer,

1982; Kirschbaum and Farquhar, 1984). In a plant physiology context, the many possible

applications of the FvCB model include assessing the impact of elevated atmospheric CO_2

concentrations on photosynthesis (Bernacchi et al., 2005), identifying limitations to

photosynthetic CO_2 assimilation (Busch and Sage, 2017), evaluating the genetic diversity of

ribulose-1,5-bisphosphate carboxylase-oxygenase (Rubisco) carboxylation efficiency within a

species (De Souza et al., 2020), and helping to guide strategies for developing sustainable and

high-performing crops in the face of climate change (He and Matthews, 2023; Matthews et al.,

2022; Wu et al., 2023; Yin and Struik, 2017).

The FvCB model works by separately calculating three potential rates of ribulose 1,5-

bisphosphate (RuBP) carboxylation as catalyzed by Rubisco, which are limited by either Rubisco

activity (W_c), RuBP regeneration (W_j), or triose phosphate utilization (TPU) (W_p). The slowest

potential rate is chosen as the actual RuBP carboxylation rate (V_c). The net leaf-level CO_2

assimilation rate (A_n) can then be determined by subtracting CO_2 losses due to RuBP

oxygenation and non-photorespiratory CO_2 release from the gains due to RuBP carboxylation.

Mathematically, this process can be expressed by five equations:

$$W_c = \frac{C \cdot V_{c,max}}{C + K_c \cdot \left(1 + \frac{O}{K_o}\right)} \tag{1a}$$

$$W_j = \frac{C \cdot J}{4 \cdot C + 8 \cdot \Gamma^*} \tag{1b}$$

$$W_p = \begin{cases} \infty, & C \leq \Gamma^* \cdot (1 + 3 \cdot \alpha) \\ \frac{3 \cdot C \cdot T_p}{C - \Gamma^* \cdot (1 + 3 \cdot \alpha)}, & C > \Gamma^* \cdot (1 + 3 \cdot \alpha) \end{cases} \tag{1c}$$

$$V_c = \min\{W_c, W_j, W_p\} \tag{1d}$$

$$A_n = V_c \cdot \left(1 - \frac{\Gamma^*}{C}\right) - R_L \tag{1e}$$

Here, C is the partial pressure of CO_2 in the vicinity of Rubisco. Definitions for all the symbols

used in these equations can be found in Table **1**. Although the earliest expressions of the FvCB

model did not include TPU, they did use the same reasoning when identifying the limiting

process, so we refer to Equation **1** as the "original FvCB model" to distinguish it from the other

variants discussed below.

The conditional form of Equation **1c** ensures that $W_p \geq 0$ (as is necessary for a carboxylation

rate) and that TPU cannot impose a limit on V_c when $C \leq \Gamma^* \cdot (1 + 3 \cdot \alpha)$. This lower C

threshold for TPU limitations will be discussed later in more detail. Note that in a more complex

version of the model, the TPU limitation separately includes glycolate carbon leaving the

photorespiratory pathway as glycine, serine, or 5,10-methylene-tetrahydrofolate (CH_2-THF),

and Γ^* is modified to reflect changes in photorespiratory CO_2 release due to glycine and CH_2-

THF synthesis (Busch, 2020; Busch et al., 2018). We use the simpler version for clarity because

the Busch *et al*. update does not affect the general conclusions presented here (Supplemental

Section **S1**).

Most applications of the FvCB model in scientific research require knowledge of the values of

key parameters such as $V_{c,max}$, J, and T_p; for example, parameter values may be used to

characterize groups of plants or as inputs to computational models. These parameter values

can be estimated from experimentally measured CO_2 response curves, which are obtained by

using a gas exchange system to record A_n and C_i as the CO_2 concentration around a leaf is

varied under otherwise constant conditions of incident light, humidity, and temperature. By

comparing the measured curve against predictions from the FvCB model made using $C = C_i$ or,

if possible, $C = C_c$, it is possible to find values of the model parameters that best reproduce the

measured assimilation rates. Many approaches to curve fitting can be found in the literature,

some of which include additional types of data that can be used to estimate C_c, such as

chlorophyll fluorescence (Bellasio et al., 2016; Duursma, 2015; Gu et al., 2010; Lochocki et al.,

2025; Moualeu-Ngangue et al., 2017; Sharkey et al., 2007; Stinziano et al., 2021; Wang et al.,

2017; Xiao et al., 2021).

A complication to curve fitting and other applications is that the FvCB model equations are not

always written consistently across scientific publications and computational tools. Many

variations are superficial (Supplemental Section **S1**), but one is not: Equations **1d** and **1e** are

often replaced by a single equation calculating A_n as the minimum of three potential net

assimilation rates (Figure **1a**). As described in detail below, this variation, which we refer to as

the "min-*A* variant," alters the model's predictions for A_n when $C < \Gamma^*$ such that they disagree

with measurements. Another issue is that although TPU limitations are understood to only

occur at high C (Harley and Sharkey, 1991; Sharkey, 2019; Sharkey et al., 2007), the explicit

lower C threshold for TPU included in Equation **1c** is often omitted, leading to situations where

the model incorrectly predicts TPU to be limiting. The prevalence of these model variations in

the literature and the potential consequences of using them in parameter estimates have not

been explored.

While it is difficult or impossible to determine how often researchers use a particular variant,

estimates can be made by surveying peer-reviewed publications (Supplemental Section **S2**). The

min-*A* variant first appears in the literature around 1990, with Collatz et al. (1990) and Collatz et

al. (1991) being the earliest examples found in the survey. Out of the highest-cited papers

published since 1990 that discuss the FvCB model equations, slightly more than half present the

min-*A* variant (Figure **1b**). The min-*A* variant was identified as an "incorrect form" of the FvCB

model in a 2010 publication (Gu et al., 2010), but its problems were not explained or

demonstrated, and thus highly-cited publications have continued to use it with roughly the

same frequency since 2013 (Figure **1b**). Likewise, the min-A variant is more likely to be used in

popular software tools for estimating FvCB model parameters from CO_2 response curves (Figure

1b). Among the publications identified in this survey, only one includes an explicit lower C

threshold for TPU limitations: Gu et al. (2010).

The survey of software tools also reveals the presence of two additional sub-variants, which we

refer to as the "forced Rubisco limitation" (FRL) and "fixed TPU threshold" (FTT) modifications.

The FRL modification applies to the min-A variant and enforces Rubisco limitations at low C

such that $A_n = A_c$ for $C \leq \Gamma^*$. The FTT modification applies to either variant and prevents TPU

limitations for C below a fixed threshold value, rather than the one in Equation **1c**, which is

determined from leaf biochemistry. These modifications address some, but not all, biological

and numerical issues related to variants of the FvCB model equations, as described below.

Figure 1: Description of the original FvCB model and the min-A variant, and an estimate of their relative prevalence in the

scientific literature. (a) Overview of the essential difference between the original FvCB model and the min-A variant. The

carboxylation rates W_c, W_j, and W_p that appear in the original FvCB model can be calculated using Equation 1, while the

assimilation rates A_c, A_j, and A_p that appear in the min-A variant can be calculated using Equation 2. (b) Fraction of highly cited

papers (or tools for estimating FvCB model parameter values from CO_2 response curves) that present (or use) the original FvCB

equations, the min-A variant, or other variations (Supplemental Section S2).

Although the literature survey presented here is limited (and does not include any software

tools used in climate or agricultural modeling), it strongly suggests that many researchers

working with biochemical models of C$_3$ net CO_2 assimilation have seen or used the min-A

variant rather than the original FvCB equations. It may even be possible that the min-*A* variant

is more prevalent overall. Many researchers may also be unaware of the lower C threshold for

TPU limitations. Given the central role of the FvCB model in several fields, it is necessary to

elucidate the differences between the original FvCB model and the min-*A* variant (with or

without the FRL and FTT modifications) and to quantify the errors resulting from use of the min-

A variant to estimate photosynthetic parameter values.

It might be argued that the differences between the original FvCB model and the min-*A* variant

are immaterial because both variants are incorrect when $C < \Gamma^*$ due to Rubisco deactivation

and RuBP depletion, processes that are not explicitly included in either variant. Based on this

reasoning, some researchers may intentionally avoid making gas exchange measurements or

performing curve fits at low C. Yet, the ability to fit CO_2 response curves for $C < \Gamma^*$ has not

been systematically tested. Thus, there is an opportunity to better understand net CO_2

assimilation at low C by first expanding both variants to include Rubisco deactivation and RuBP

depletion and then comparing each against measured assimilation rates for $C < \Gamma^*$.

The objectives of this study are to (1) describe the key differences between the original FvCB

model and its variants, (2) establish the biochemical basis of the lower C threshold for TPU, (3)

develop a simple way to include Rubisco deactivation and RuBP depletion in the FvCB

framework, and (4) test the validity of each variant by fitting CO_2 response curves that extend

to $C < \Gamma^*$. Throughout, we emphasize that the unrealistic predictions of the min-*A* variant

cannot be fixed through the FRL and FTT modifications, and that the min-*A* variant cannot be

extended to include Rubisco deactivation or RuBP depletion. Finally, the curve fits show that

the original FvCB model is able to represent C_3 net CO_2 assimilation at low C while the min-

A+FTT variant cannot, leading to underestimates of V_{cmax} and J by up to 23% and 12%,

respectively. Although errors are unlikely to be as severe in most situations, the min-*A* variant

does not provide better predictions in any situation.

2. DESCRIPTION

Symbol	Meaning	Units	Range	Typical value
A_n	Net CO_2 assimilation rate	$\mu mol\ m^{-2}\ s^{-1}$	-	-
A_c	Net CO_2 assimilation rate when RuBP carboxylation is limited by Rubisco activity	$\mu mol\ m^{-2}\ s^{-1}$	-	-
$A_c\|_{C=0}$	The value of A_c when C = 0	$\mu mol\ m^{-2}\ s^{-1}$	-	-
A_d	Net CO_2 assimilation rate when RuBP carboxylation is limited by substantial Rubisco deactivation or RuBP depletion at low C	$\mu mol\ m^{-2}\ s^{-1}$	-	-
A_j	Net CO_2 assimilation rate when RuBP carboxylation is limited by the rate of electron transport going to support RuBP regeneration[b]	$\mu mol\ m^{-2}\ s^{-1}$	-	-
$A_j\|_{C=0}$	The value of A_j when C = 0	$\mu mol\ m^{-2}\ s^{-1}$	-	-
A_p	Net CO_2 assimilation rate when RuBP carboxylation is limited by the rate of inorganic phosphate release from TPU	$\mu mol\ m^{-2}\ s^{-1}$	-	-
C	Partial pressure of CO_2 in the vicinity of Rubisco	μbar	$0 \leq C$	-
C_{cj}	The value of C where $W_c = W_j$	μbar	$0 \leq C_{cj}$	-
C_c	Chloroplastic partial pressure of CO_2	μbar	$0 \leq C_c$	-
C_i	Intercellular partial pressure of CO_2	μbar	$0 \leq C_i$	-
C_i^*	The value of C_i when $C_c = \Gamma^*$	μbar	$0 \leq C_i^*$	-
J	Potential rate of electron transport going to support RuBP regeneration at a given light intensity	$\mu mol\ m^{-2}\ s^{-1}$	$0 \leq J$	170[a]
J_{eq}	The value of J where $A_c\|_{C=0} = A_j\|_{C=0}$	$\mu mol\ m^{-2}\ s^{-1}$	$0 \leq J_{eq}$	-
J_d	The value of J where the denominator of Equation **A5** is zero	$\mu mol\ m^{-2}\ s^{-1}$	$0 \leq J_d$	-
J_n	The value of J where the numerator of Equation **A5** is zero	$\mu mol\ m^{-2}\ s^{-1}$	$0 \leq J_n$	-
K_c	Michaelis-Menten constant for CO_2	μbar	$0 < K_c$	259[a]
K_o	Michaelis-Menten constant for O_2	$mbar$	$0 < K_o$	179[a]
O	Partial pressure of O_2 in the vicinity of Rubisco	$mbar$	$0 \leq O$	210
Q_{in}	Incident photosynthetically active flux density	$\mu mol\ m^{-2}\ s^{-1}$	$0 \leq Q_{in}$	-

R_L	Rate of non-photorespiratory CO_2 release in the light	µmol m^{-2} s^{-1}	$0 \leq R_L$	1[a]
R_{pc}	Net rate of inorganic phosphate consumption in the chloroplast due to photosynthesis and photorespiration[c]	µmol m^{-2} s^{-1}	-	-
T_p	Potential rate of TPU	µmol m^{-2} s^{-1}	$0 < T_p$	11.8[a]
V_c	RuBP carboxylation rate	µmol m^{-2} s^{-1}	$0 \leq V_c$	-
$V_{c,max}$	Maximum rate of Rubisco activity	µmol m^{-2} s^{-1}	$0 \leq V_{c,max}$	100[a]
V_o	RuBP oxygenation rate	µmol m^{-2} s^{-1}	$0 \leq V_o$	-
W_c	RuBP carboxylation rate limited by Rubisco activity	µmol m^{-2} s^{-1}	$0 \leq W_c$	-
W_d	RuBP carboxylation rate limited by substantial Rubisco deactivation or RuBP depletion at low C	µmol m^{-2} s^{-1}	$0 \leq W_d$	0
W_j	RuBP carboxylation rate limited by the rate of electron transport going to support RuBP regeneration[b]	µmol m^{-2} s^{-1}	$0 \leq W_j$	-
W_p	RuBP carboxylation rate limited by the rate of inorganic phosphate release from TPU	µmol m^{-2} s^{-1}	$0 \leq W_p$	-
α	Fraction of remaining glycolate carbon not returned to the chloroplast after accounting for carbon released as CO_2[d]	dimensionless	$0 \leq \alpha \leq 1$	0[a]
Γ^*	CO_2 compensation point in the absence of non-photorespiratory CO_2 release	µbar	$0 < \Gamma^*$	38.6[a]
Δ_c	Net rate of carbon production during the reduction of PGA to unbound RuBP	µmol m^{-2} s^{-1}	-	-

Table 1: Description of symbols used in this publication. Values marked with a superscript (a) are from the caption of Figure 2.6
in von Caemmerer (2000). The typical value for O in the table is the atmospheric value at sea level rounded to two significant
figures. (b) By convention, the term "RuBP regeneration limitations" refers to the case where RuBP regeneration is limited
primarily by electron transport, and possibly co-limited by the activity of CBB cycle enzymes such as sedoheptulose-1,7
bisphosphatase; limitations to RuBP regeneration imposed by the supply of inorganic phosphate or carbon are treated
separately. (c) Consumption refers to the incorporation of inorganic phosphate into organic molecules. (d) Even when α is zero,
one quarter of glycolate carbon is released as CO_2 during glycine decarboxylation (Harley and Sharkey, 1991), so the overall
fraction of glycolate carbon not returned to the chloroplast is $(1 + 3 \cdot \alpha)/4$. Also note that in an updated version of the FvCB
model, α is replaced by three parameters – α_g, α_s, and α_t – that represent fractions of glycolate carbon leaving the
photorespiratory pathway as glycine, serine, or CH_2-THF, respectively (Busch, 2020; Busch et al., 2018).

2.1. The Original FvCB Model

The original FvCB model is given above in Equation **1**. Here we note that within the time scale of a typical gas exchange measurement, the main environmental influences on the potential carboxylation rates are C, Q_{in}, O, and temperature. Note that some of this influence is not immediately obvious; for example, Γ^* depends on O, and J depends on Q_{in} and other factors (von Caemmerer, 2000). For simplicity, we assume constant temperature and treat Γ^* and J as inputs to the model, especially since several different approaches are available for calculating J from Q_{in} (Walker et al., 2021). As C, J, and O vary, the slowest carboxylation rate may change. Changes in the rate-limiting process as C increases from zero will be discussed later in more detail (Section **3.2**).

2.2. The Min-A Variant

When RuBP carboxylation is Rubisco-limited, the corresponding net assimilation rate (called A_c rather than A_n to indicate the rate-limiting process) can be found by setting $V_c = W_c$ in Equation **1e** and replacing it with the expression in Equation **1a**:

$$A_c = W_c \cdot \left(1 - \frac{\Gamma^*}{C}\right) - R_L = \frac{(C-\Gamma^*) \cdot V_{c,max}}{C+K_c \cdot \left(1+\frac{O}{K_o}\right)} - R_L. \tag{2a}$$

Likewise, when RuBP carboxylation is limited by RuBP regeneration or TPU, the corresponding net assimilation rates are given by analogous equations:

$$A_j = W_j \cdot \left(1 - \frac{\Gamma^*}{C}\right) - R_L = \frac{(C-\Gamma^*) \cdot J}{4 \cdot C + 8 \cdot \Gamma^*} - R_L, \tag{2b}$$

$$A_p = W_p \cdot \left(1 - \frac{\Gamma^*}{C}\right) - R_L = \frac{3 \cdot (C-\Gamma^*) \cdot T_p}{C - \Gamma^* \cdot (1+3 \cdot \alpha)} - R_L. \tag{2c}$$

In the min-*A* variant, A_c, A_j, and A_p are considered to be potential net CO_2 assimilation rates and the actual net CO_2 assimilation rate is calculated as the smallest of the three, in analogy with Equations **1d** and **1e** from the original FvCB model:

$$A_n = \min\{A_c, A_j, A_p\}. \tag{2d}$$

Thus, the essential difference between Equations **1** and **2** is that the original FvCB model

chooses a minimal potential carboxylation rate, while the min-*A* variant chooses a minimal

potential net CO_2 assimilation rate (Figure **1a**). Another difference is that Equation **1c** restricts

TPU limitations to to $C > \Gamma^* \cdot (1 + 3 \cdot \alpha)$, while Equation **2c** makes no such restriction. We do

not consider this to be an essential difference between the variants because many sources omit

the restrictions from the original FvCB equations as well. Nevertheless, unreasonable results

such as negative carboxylation rates and divergent behavior can occur in either variant when

this restriction is not included; these issues have been noticed before and the FTT modification

has been used to address them, although some issues remain when using a fixed TPU threshold

(Section **3.3**).

3. RESULTS

3.1. Minimal Carboxylation Rates Are Not Equivalent to Minimal Assimilation Rates

There are two irreconcilable differences between the original FvCB model and the min-*A* variant

(Figure **2**).

1) When $C < \Gamma^*$, the two variants always disagree about the rate-limiting process and the

values of V_c and A_n. This is a consequence of the fact that $(1 - \Gamma^*/C)$ is negative for $C < \Gamma^*$.

For example, neglecting TPU, suppose that $W_c < W_j$ for $C < \Gamma^*$. In this case, $\min\{W_c, W_j\} = W_c$,

so the original FvCB model predicts Rubisco limitations with $V_c = W_c$ and $A_n = A_c$ (Figure **2a**).

In contrast, the min-*A* variant uses the comparison $W_c \cdot (1 - \Gamma^*/C) > W_j \cdot (1 - \Gamma^*/C)$, and

thus predicts RuBP regeneration limitations with $V_c = W_j$ and $A_n = A_j$ (Figure **2b**), leading to a

discrepancy where the min-*A* variant predicts a larger V_c and smaller A_n (Figure **2c**). From this

analysis, we can see that the relationship between A_c, A_j, and A_n at low C in the original FvCB

model is actually

$$A_n = \max\{A_c, A_j\} \text{ if } C < \Gamma^* \tag{3}$$

rather than Equation **2d**. Typical values of Γ^* are close to 40 µbar (approximately 40 µmol mol^{-1}

or 40 ppm at typical atmospheric pressure), so this discrepancy between the variants only

occurs at low values of C. Historically, such low values of C have been rarely accessed in gas

exchange measurements because the available equipment could not easily reach them, but

modern instruments can achieve lower values and researchers are now more likely to

encounter $C < \Gamma^*$. As will be shown below, this small difference prevents the min-A variant

from matching measured CO_2 response curves, and it can have a large influence on parameter

estimation (Section **3.5**).

Figure 2: Comparison of outputs from the original FvCB model and the min-A variant for identical input parameter values with α

*= 0. (**a,b**) A_n (thick solid lines) as determined from potential carboxylation or assimilation rates (dashed lines) using Equations **1***

*and **2**, respectively. Crossover points where the rate-limiting process changes are marked with white stars. Solid rectangles*

*indicate ranges of C where carboxylation or assimilation is limited by Rubisco, RuBP regeneration, or TPU. (**c**) Direct comparison*

of the variants across a range of C values. Inset: close-up showing the difference between the variants when $C < \Gamma^*$. All

*parameters were set to the values specified in Table **1**.*

Irrespective of any practical consequences, the min-A variant is not logically consistent because

choosing the smallest net assimilation rate can lead to paradoxical results. For example, if V_{cmax}

were to become zero when $C < \Gamma^*$ due to substantial Rubisco deactivation (Section **3.4**), the

min-A variant would nevertheless predict $V_c = W_j$ and $A_n = A_j$, as discussed above. Assuming a

nonzero J, this is a contradictory prediction that $V_c > 0$ when $V_{cmax} = 0$. In contrast, choosing a

minimal carboxylation rate always produces self-consistent results.

2) The two variants always disagree about the number of points where the rate-limiting process

changes; these special values of C are called crossover points. In the min-A variant, $A_c = A_j =$

$-R_L$ when $C = \Gamma^*$ (Equations **2a** and **2b**), so a crossover point always exists at $C = \Gamma^*$ (Figure

2b). However, in the original FvCB model, W_c and W_j are generally not equal for $C = \Gamma^*$

(Equations **1a** and **1b**), so a crossover point does not always exist at $C = \Gamma^*$ (Figure **2a**). In other

words, there is a spurious crossover point at $C = \Gamma^*$ in the min-*A* variant. This spurious

crossover point has been noticed before (Dubois et al., 2007; Gu et al., 2010) and the FRL

modification has been used to address this issue, although this modification does not fully

reconcile the variants in all situations (Section **3.2**).

3.2. Rubisco Activity Does Not Always Limit Carboxylation at Low CO₂ Concentrations

Some fitting tools that implement the min-*A* variant introduce a constraint in their code where

$A_n = A_c$ is enforced for $C \leq \Gamma^*$ (or possibly below another threshold value), even when

Equation **2d** would predict otherwise (Figure **1** and Supplemental Section **S2.4**). Although rarely

written explicitly as an equation, this is equivalent to replacing Equation **2d** with

246
$$A_n = \begin{cases} A_c, & C \leq \Gamma^* \\ \min\{A_c, A_j, A_p\}, & C > \Gamma^* \end{cases}$$

We refer to this as the "forced Rubisco limitation" (FRL) modification, and such variants as min-

A+FRL variants. These variants agree with the original FvCB model in situations where

carboxylation is Rubisco limited for $C \leq \Gamma^*$.

However, the FRL modification will never be truly successful at fixing the min-*A* variant because

there are no simple rules governing the progression of rate-limiting processes as C increases.

Carboxylation can be limited by Rubisco activity or RuBP regeneration at low C, and a crossover

between limitations is not guaranteed. Neglecting TPU, it is possible to analytically determine

which process is limiting carboxylation at $C = 0$ and whether a crossover occurs (Supplemental

Section **S3**). This process can be performed for many different values of $V_{c,max}$ and J, producing

a map that shows where each possible sequence of assimilation rates occurs in the original

FvCB model (Figure **3a**). When including TPU, it is more straightforward to simulate a CO₂

response curve and extract the observed sequence of limitations; again, this can be performed

for many values of $V_{c,max}$ and J, producing a map (Figure **3b**).

*Figure 3: Predicted sequences of limiting processes as C increases from zero. (**a-d**) Maps of $(V_{c,max}, J)$-space, where labels within each region indicate the sequences as calculated by the original FvCB model (neglecting or including TPU) (**a-b**), the min-A variant (**c**), or the min-A+FRL variant (**d**). The triangles to the left of (**a**) and (**c**) indicate an approximate relation between J and Q_{in}. Dashed and solid black lines in (**a**) show J_d and J_n (Equations **A6** and **A7** in Supplemental Section **S3**). All labels in (**b-d**) refer to observed sequences below a threshold of 2000 μbar. All parameters other than $V_{c,max}$ and J were set to the values specified in Table **1**.*

This analysis shows that when $V_{c,max}$ is relatively small compared to J, the original FvCB model predicts Rubisco limitations for all values of C (dark blue regions labeled "1" in Figure **3**). This has been observed in modified tobacco with reduced Rubisco content (von Caemmerer et al., 1994), and this situation could potentially occur in plants under nitrogen stress (leading to low $V_{c,max}$) or in leaves exposed to high incident light levels (leading to high J). Likewise, when J is relatively small compared to $V_{c,max}$, the model predicts RuBP regeneration limitations for all values of C (red regions labeled "5" in Figure **3**). This situation could potentially occur in leaves exposed to low incident light levels (leading to low J), although this may be difficult to observe in practice due to a concurrent reduction in $V_{c,max}$ that tends to occur at low Q_{in} (Sage et al.,

2002; Taylor et al., 2022). Actual values of Q_{in} corresponding to high or low J depend strongly

on details of leaf biochemistry and growth environment, but for many plants, $Q_{in} \leq 100$ μmol

m^{-2} s^{-1} would correspond to low J while $Q_{in} \geq 1800$ μmol m^{-2} s^{-1} would correspond to high J.

The maps also show that $A_c \rightarrow A_j \rightarrow A_p$ is not the only predicted sequence involving TPU

limitations. In fact, a transition directly from Rubisco limited to TPU limited carboxylation ($A_c \rightarrow$

A_p) can be observed in plants grown in low-light conditions (Sharkey, 2019). Note that there is

no region corresponding to $A_j \rightarrow A_c$, indicating that this transition is not predicted to occur for

typical parameter values (Supplemental Sections **S3** and **S4**).

Similar maps generated using the min-A and min-A+FRL variants show large differences relative

to the original FvCB model map (Figures **3c** and **3d**). Because of the spurious crossover point at

Γ^* in the min-A variant (Section **3.1**), each region of the min-A variant map has one extra step

compared to its original FvCB analog, several of which do not occur in the original FvCB map.

For example, the $A_c \rightarrow A_p$ region of the original FvCB map (dark tan region labeled "4" in Figure

3b) becomes an $A_j \rightarrow A_c \rightarrow A_p$ region in the min-A variant map (teal region labeled "8" in

Figure **3c**). Because the min-A+FRL variant enforces Rubisco-limited assimilation at low C, its

map lacks several sequences that are present in the original FvCB map. For example, the A_j

region of the original FvCB map becomes an $A_c \rightarrow A_j$ region in the min-A+FRL map. These

results demonstrate that the original FvCB model and the min-A variant do not predict the

same transitions between limiting processes, and that the FRL modification does not fully bring

the min-A variant into agreement with the original FvCB model.

3.3. TPU Can Only Limit Carboxylation When Photosynthesis and Photorespiration Are

Net Consumers of Free Inorganic Phosphate

Additional differences between the original FvCB model and the min-A variant can appear when

$\alpha > 0$. This parameter is related to TPU and appears in Equations **1c** and **2c**. In the original FvCB

model, nonzero α introduces a downward slope to W_p (Figure **4a**). The resulting decrease in A_p

with increasing C, termed "reverse sensitivity," is an identifying signature of TPU limitations,

and the inclusion of α is essential to fit CO$_2$ response curves exhibiting reverse sensitivity (Busch

et al., 2018; Harley and Sharkey, 1991). In the min-A variant, nonzero α similarly introduces a

downward slope to A_p at high values of C, but also produces divergent behavior at lower values

of C (Figure **4b**). Mathematically, this behavior occurs because the denominator of Equation **2c**

becomes zero when $C = \Gamma^* \cdot (1 + 3 \cdot \alpha)$, and A_p approaches $-\infty$ as C approaches $\Gamma^* \cdot$

$(1 + 3 \cdot \alpha)$ from below. This issue does not occur for W_p because Equation **1c** sets W_p to ∞ for

$C \leq \Gamma^* \cdot (1 + 3 \cdot \alpha)$; however, it will occur if this condition is omitted from Equation **1c**.

Some fitting tools introduce a constraint in their code where TPU cannot set the net CO_2

assimilation rate below a fixed threshold value of C, which is often chosen to be 400 µbar

(Supplemental Section **S2.4**). Although rarely written explicitly as an equation, when using the

min-*A* variant this is equivalent to replacing Equation **2c** with

313
$$A_p = \begin{cases} \infty, & C \leq 400 \text{ µbar} \\ \frac{3 \cdot (C - \Gamma^*) \cdot T_p}{C - \Gamma^* \cdot (1 + 3 \cdot \alpha)} - R_L, & C > 400 \text{ µbar} \end{cases}$$

314　or when using the original FvCB model, replacing Equation **1c** with

315
$$W_p = \begin{cases} \infty, & C \leq 400 \text{ µbar} \\ \frac{3 \cdot C \cdot T_p}{C - \Gamma^* \cdot (1 + 3 \cdot \alpha)}, & C > 400 \text{ µbar} \end{cases}$$

We refer to this as the "fixed TPU threshold" (FTT) modification, and such variants as the min-

A+FTT or original FvCB+FTT variants. (Note that the min-*A*+FTT and min-*A*+FRL variants can be

combined to form another variant with both types of modifications: the min-*A*+FRL+FTT

variant.) These FTT variants agree with the original FvCB model in situations where W_p is not

the smallest carboxylation rate below the threshold value of C. However, if the original FvCB

model predicts $A_n = A_p$ for $\Gamma^* \cdot (1 + 3 \cdot \alpha) < C < 400$ µbar, then the original FvCB+FTT and

min-*A*+FTT variants will predict $A_n \neq A_p$ in that range, and then suddenly drop down to $A_n =$

A_p above the threshold (Figure **4c**). TPU limitations below 400 µbar are rare, but have been

observed (Harley and Sharkey, 1991). Another potential issue with the FTT modification is that

the divergent behavior of A_p will persist if the fixed threshold value is smaller than $\Gamma^* \cdot$

$(1 + 3 \cdot \alpha)$.

*Figure 4: Comparison of original FvCB model, min-A variant, and min-A+FTT variant outputs for identical input parameter values with $\alpha > 0$. (**a,b**) A_n (thick solid lines) as determined from potential carboxylation or assimilation rates (dashed lines) using Equations **1** and **2**, respectively. Crossover points where the rate-limiting process changes are marked with white stars. Solid rectangles indicate ranges of C where carboxylation or assimilation is limited by Rubisco, RuBP regeneration, or TPU. (**c**) Direct comparison of the variants across a range of C values. The following parameter values were used for these calculations: $V_{c,max} = 120 \ \mu mol \ m^{-2} \ s^{-1}$, $T_p = 8 \ \mu mol \ m^{-2} \ s^{-1}$, and $\alpha = 0.5$. All other parameters were set to the values specified in Table **1**.*

The FTT modification will never be truly successful at fixing the issues with Equation **2c** because

it does not account for an important biochemical aspect of TPU limitations: TPU can only limit

carboxylation when the net effect of photosynthesis and photorespiration is to shrink the pool

of free inorganic phosphate (P_i) in the chloroplast by incorporating it into organic compounds, a

process referred to as P_i consumption. P_i is required to synthesize adenosine triphosphate (ATP)

via photophosphorylation, which in turn is used to regenerate RuBP. To achieve sustained ATP

synthesis and RuBP regeneration, the net rate of P_i consumption due to photosynthesis and

photorespiration (R_{pc}) must be balanced by the rate at which P_i is returned to the chloroplast

through the utilization of triose phosphate for sugar production (T_p); otherwise, the pool of P_i

would become depleted (Harley and Sharkey, 1991; Sharkey, 1985; von Caemmerer, 2000). In

other words,

345
$$T_p = R_{pc} \tag{4a}$$

when TPU limits carboxylation. This requirement can be used to derive Equation **1c**, where the

key step in the derivation is to relate R_{pc} to the RuBP carboxylation rate. Here we show that the

restriction $C > \Gamma^* \cdot (1 + 3 \cdot \alpha)$ is a natural consequence of this derivation.

By considering the stoichiometry of the carboxylation and oxygenation cycles and allowing for

an additional release of P_i during photorespiration due to glycolate carbon that remains in the

cytosol, it can be shown that

$$R_{pc} = \frac{V_c}{3} - \frac{V_o}{6} - \frac{\alpha \cdot V_o}{2},$$ (4b)

353   where V_o is the rate of RuBP oxygenation (Harley and Sharkey, 1991; von Caemmerer, 2000).

354   Next, it can also be shown that V_o is related to V_c via $V_o = 2 \cdot \Gamma^* \cdot V_c / C$ (von Caemmerer, 2000),

355   allowing us to eliminate V_o from Equation **4b**:

$$R_{pc} = \frac{V_c}{3 \cdot C} [C - \Gamma^* \cdot (1 + 3 \cdot \alpha)].$$ (4c)

Here it is essential to note that, first, when $C = \Gamma^* \cdot (1 + 3 \cdot \alpha)$, the net rate of P_i consumption

is zero. Second, for smaller values of C, R_{pc} is negative, indicating that photosynthesis and

photorespiration are actually releasing P_i. In both of these situations, it is not possible for TPU

to become a limiting factor because photosynthesis and photorespiration are not shrinking the

pool of P_i in the chloroplast, and hence the return of P_i via TPU is not necessary to prevent P_i

depletion.

Finally, when TPU limits the carboxylation rate, V_c is denoted by W_p; in this case, we can now

use Equations **4a** and **4c** to express the requirement for TPU limitation as

$$T_p = R_{pc} = \frac{W_p}{3 \cdot C} [C - \Gamma^* \cdot (1 + 3 \cdot \alpha)].$$ (4d)

Solving Equation **4d** for W_p and remembering that photosynthesis and photorespiration only

consume P_i when $C > \Gamma^* \cdot (1 + 3 \cdot \alpha)$ yields Equation **1c**. This lower C threshold for TPU

limitations is a consequence of leaf biochemistry and its inclusion is guaranteed to prevent

numerical errors such as negative or undefined W_p, unlike the fixed threshold used in the FTT

modification. Although this threshold could be included in Equation **2c** from the min-*A* variant

as a conditional statement like the one in Equation **1c**, none of the studies in the literature

survey (Supplemental Section **2**) take this approach. Furthermore, doing so would not address

the other issues with the min-*A* variant (Sections **3.2**, **3.3**, and **3.5**).

3.4. The Original FvCB Model Can Be Extended To Include Substantial Rubisco Deactivation and RuBP Depletion at Low CO_2 Concentrations but the Min-A Variant Cannot

Besides the processes explicitly included in Equations **1** and **2**, Rubisco deactivation and RuBP depletion may also influence net CO_2 assimilation rates. Rubisco deactivation can occur when a leaf is held under low CO_2 conditions or under low light, likely by increasing the fraction of decarbamylated Rubisco, and the result of this process can be modeled as a reduction in V_{cmax} (Lochocki et al., 2025; Taylor et al., 2022; Sage et al., 2002). The original FvCB model and the min-A variant make different predictions in the hypothetical extreme case of full Rubisco deactivation for $C < \Gamma^*$, where V_{cmax} would become zero. Assuming no other parameter values change, W_c would also become zero but W_j and W_p would remain positive, so the original FvCB model would predict Rubisco limitations with $A_n = A_c = -R_L$ (Equation **1**). Yet, A_j would be less than $A_c = -R_L$, so the min-A variant would predict a different limiting process and a smaller value of A_n (Equation **2**). In other words, the min-A variant does not allow substantial Rubisco deactivation to limit net CO_2 assimilation at low C, even when no Rubisco sites are active.

Similarly, the min-A variant does not predict a change in A_n due to substantial RuBP depletion, which can occur when $C < \Gamma^*$ because RuBP cannot be regenerated in this range. RuBP regeneration depends not only on electron transport and the availability of P_i, but also on the supply of 3-phosphoglyceric acid (PGA) from photosynthesis and photorespiration, which is reduced to unbound RuBP in a multi-step enzymatic pathway (von Caemmerer, 2000). When $C < \Gamma^*$, the supply of carbon as PGA is insufficient to regenerate enough RuBP, eventually depleting the pool of unbound RuBP. This can be understood by considering the carbon stoichiometry of the reduction pathway, where Rubisco uses RuBP at a rate of $V_c + V_o$ and PGA is produced at a rate of $2V_c + 1.5V_o$ (von Caemmerer, 2000). RuBP and PGA are five- and three-carbon molecules, respectively, so the difference between carbon supply and demand (Δ_c) is

$$\Delta_c = 3 \cdot (2V_c + 1.5V_o) - 5 \cdot (V_c + V_o) = V_c \cdot \left(1 - \frac{\Gamma^*}{C}\right), \tag{5}$$

where $V_o = 2 \cdot \Gamma^* \cdot V_c/C$ has been used to simplify the final expression. There is a carbon

surplus ($\Delta_c > 0$) for $C > \Gamma^*$, which is exported from the chloroplast as triose phosphates.

Carbon supply and demand are exactly equal ($\Delta_c = 0$) when $C = \Gamma^*$, and there is a deficit ($\Delta_c <$

0) for $C < \Gamma^*$. In the latter case, continued Rubisco activity draws from the chloroplastic pool of

unbound RuBP without replenishing it.

Modeling RuBP depletion caused by insufficient carbon supply is more complex than Rubisco

deactivation because it does not simply alter the value of a parameter such as V_{cmax} or J.

Although not traditionally considered as part of the original FvCB model, here we include it by

adding a fourth potential limitation to Equation **1d**:

$$V_c = \min\{W_c, W_j, W_p, W_d\} \tag{6a}$$

where W_d is the carboxylation rate limited by RuBP depletion. W_d is taken to be zero when C

has been held below Γ^* long enough for Rubisco activity to completely deplete the pool of

unbound RuBP, and is otherwise assumed to not limit carboxylation. This can be described

mathematically as

$$W_d = \begin{cases} 0, & C < \Gamma^* \text{ for a sufficiently long time} \\ \infty, & \text{otherwise} \end{cases} \tag{6b}$$

Equation **6b** is not precise and omits two key details – incomplete RuBP depletion can lead to a

small but nonzero W_d at low C, and W_d should depend continuously on C, RuBP pool size, and

time. Nevertheless, this simplistic equation is instructive for understanding long-term steady-

state net CO_2 assimilation at low C. When substantial RuBP depletion limits carboxylation, $V_c =$

W_d and the corresponding net CO_2 assimilation rate (A_d) can be found with Equations **1a** and

6b:

$$A_d = \begin{cases} -R_L, & C < \Gamma^* \text{ for a sufficiently long time} \\ \infty, & \text{otherwise} \end{cases} \tag{7a}$$

With this, RuBP depletion can also be incorporated into the min-*A* variant by adding a fourth

potential assimilation rate to Equation **2d**:

425
$$A_n = \min\{A_c, A_j, A_p, A_d\}. \tag{7b}$$

Equations **6** and **7** make contradictory predictions for A_n when C is below Γ^* for enough time to

ensure substantial RuBP depletion. Assuming otherwise typical model parameter values in this

scenario, W_c, W_j, and W_p would each be positive, so $W_d = 0$ would always be the smallest

carboxylation rate, and the original FvCB model would predict RuBP depletion limitations with

$A_n = A_d = -R_L$ (Equation **6**). Conversely, A_c and A_j would each be smaller than $-R_L$, so the

min-*A* variant would never predict $A_n = A_d$. In fact, Equation **7** would always predict the same

rate and limiting process as Equation **2**. In other words, the original FvCB model predicts a shift

of A_n to $A_d = -R_L$ when substantial RuBP depletion occurs at low CO_2 concentrations, while

the min-*A* variant does not allow substantial RuBP depletion to limit net CO_2 assimilation.

3.5. Comparing Both Variants to Measured CO_2 Response Curves

Since the original FvCB model and the min-*A* variant make different predictions for $C < \Gamma^*$,

even when considering substantial Rubisco deactivation and RuBP depletion, it is possible to

test them against measurements to determine which variant better represents reality. To do

this, thirty-six CO_2 response curves were measured from *Nicotiana tabacum* (tobacco) cv.

Samsun leaves at multiple incident light intensities using Licor LI-6800 portable gas exchange

systems (Supplemental Section **S5.1**). Reference CO_2 concentration set-points ranging from $10 -$

1800 μmol mol^{-1} were used, where the lowest values ensured several points in each curve with

$C < \Gamma^*$. Each curve was fit with the *PhotoGEA* R package (Lochocki, 2024; Lochocki et al., 2025)

on a C_i basis (i.e., by setting $C = C_i$) using tobacco temperature response parameters (Sharkey

et al., 2007) and allowing $V_{c,max}$, J, R_L, T_p, and α to vary. The fits were performed in four ways:

- *Original FvCB model (all C_i)*: The entire curve was fit twice using Equations **1** and **6**, and

the result with smaller root mean square error (RMSE) was chosen as the best fit. If

Equation **6** produced the best fit, the curve was considered to exhibit A_d limitations;

otherwise, the curve was not considered to exhibit A_d limitations.

- *min-A+FTT variant (all C_i)*: The entire curve was fit using Equation **2** with a fixed TPU

threshold of 400 μbar. There is no need for an additional fit using Equation **7** because

Equations **2** and **7** make identical predictions.

•   *Original FvCB model (C_i > 45 µbar)*: Points from the curve where C_i is above 45 µbar

were fit using Equation **1**. Requiring C_i > 45 µbar ensured that no points with $C < \Gamma^*$

were used for the fit, since tobacco Γ^* is approximately 39 µbar at the measurement

temperature of 27 °C.

•   min-*A+FTT variant (C_i > 45 µbar)*: Points from the curve where C_i is above 45 µbar were

fit using Equation **2** with a fixed TPU threshold of 400 µbar.

Note that the original FvCB model predicts $A_n = -R_L$ when either substantial Rubisco

deactivation or RuBP depletion limits net CO_2 assimilation for $C < \Gamma^*$ (Section **3.4**), so it is not

possible to unambiguously attribute an observed A_d limitation to either process. Because of

this, we consider W_d and A_d to represent both processes in the context of curve fitting. Also

note that the min-*A*+FTT variant is used for these comparisons because the divergent behavior

of the min-*A* variant with nonzero α severely interferes with the fitting process (Section **3.3**).

For a curve without such A_d limitations, an extrapolation of the fit made using the original FvCB

model for points where C_i is above 45 µbar agrees well with the measured points at lower C_i

(Figure **5a**), suggesting that a good fit across all measured points could be achieved using

Equation **1**. However, an extrapolation of the fit made using the min-*A*+FTT variant greatly

diverges from the measured assimilation rates for C_i below 45 µbar, even though the two fits

are identical for higher C_i (Figure **5b**). For a curve with A_d limitations, a similar extrapolation of

the original FvCB model fit made using points where C_i is above 45 µbar does not match the

measured points at lower C_i, but $A_d = -R_L$ does lie close to those points (Figure **5e**),

suggesting that a good fit across all measured points could be achieved using Equation **6**. An

extrapolation of the min-*A*+FTT fit is even further from the measured points where C_i is below

45 µbar (Figure **5f**), but the min-*A* variant never predicts $A_n = A_d$, so a good fit across all C_i is

unlikely.

As expected from these extrapolations, the original FvCB model is able to closely match the

measured points in each curve when fitting the entire range of C_i (Figures **5c** and **5g**), while the

min-*A*+FTT variant is not (Figures **5d** and **5h**). This is reflected in the RMSE values, where a

lower RMSE indicates a better fit. The RMSE values of the original FvCB fits barely increase

when adding the points with C_i < 45 μbar (0.140 vs. 0.160 for the curve without A_d limitations and 0.156 vs. 0.185 for the curve with A_d limitations), while there is a large increase in the min-A+FTT RMSE values (0.140 vs. 0.728 for the curve without A_d limitations and 0.156 vs. 1.312 for the curve with A_d limitations) (Figure **5**).

*Figure 5: Testing FvCB model variants against measured CO₂ response curves. Fits are shown for one curve without A_d limitations (**a-d**) and one with A_d limitations (**e-h**), both of which were measured with Q_{in} = 600 μmol m⁻² s⁻¹. Fits were made using the original FvCB model (**a,c,e,g**) or the min-A+FTT variant (**b,d,f,h**), either limited to C_i above 45 μbar (**a,b,e,f**) or using all measured points (**c,d,g,h**). In each panel, measured points that were included in or excluded from the fit are shown as open black and gray circles, respectively. RMSE values are calculated from only the points included in each fit. A zoomed-in portion of each curve is shown to highlight differences at low C_i; see Supplemental Section **S5** for expanded views showing the fits across the entire range of measured C_i.*

These results hold across the entire set of thirty-six curves, where eighteen were found to exhibit A_d limitations at low C_i (Figure **6**). When fitting points at all C_i, the min-A+FTT variant generally produces larger RMSE values than the original FvCB model, especially for curves that exhibit A_d limitations (Figure **6a**). Across all curves, the mean RMSE for the original FvCB fits is 0.32 μmol m⁻² s⁻¹, while the mean RMSE for the min-A+FTT fits is more than twice as large at 0.79 μmol m⁻² s⁻¹ (see also Supplemental Figure **S10**). Fits made with the original FvCB model have similar RMSE values when including or excluding the points where C_i is below 45 μbar, but fits made with the min-A+FTT variant are appreciably worse when including points where C_i is

below 45 μbar, even for curves that do not exhibit A_d limitations. Overall, this indicates that the

original FvCB model is able to represent leaf net CO_2 assimilation across all C_i values when

potential limitations due to Rubisco deactivation and RuBP depletion are considered for $C < \Gamma^*$

(Equation **6**), while the min-*A*+FTT variant cannot.

Because the min-*A*+FTT variant does not closely match the observed values of A_n for $C < \Gamma^*$,

parameter values estimated using this variant are greatly influenced by including points where

C_i is below 45 μbar (Figures **6b-c**). As compared to fits made with the original FvCB model, the

min-*A*+FTT variant tends to underestimate both V_{cmax} and J. Values of V_{cmax} from min-*A*+FTT

fits can be as much as 12 μmol m^{-2} s^{-1} lower (as much as 23% lower on a relative basis), while

values of J can be as much as 17 μmol m^{-2} s^{-1} lower (as much as 12% lower on a relative basis).

Figure 6: Comparing fit results from thirty-six CO_2 response curves. Differences in (**a**) RMSE, (**b**) V_{cmax}, and (**c**) J that occur when
including all points in the fit as compared to only using points where C_i is above 45 μbar, where $\Delta RMSE = RMSE^{All\ C_i} -$
$RMSE^{C_i>45}$, $\Delta V_{cmax} = V_{cmax}^{All\ C_i} - V_{cmax}^{C_i>45}$, $\Delta J = J^{All\ C_i} - J^{C_i>45}$, and superscripts indicate which points were fit. Values from
original FvCB model and min-A+FTT variant fits are shown as filled blue circles and open yellow circles, respectively. Vertical gray
lines connect the values from each variant for each curve. Results from curves that exhibit or do not exhibit A_d limitations are
shown in the left and right columns, respectively. Some parameter values could not be reliably estimated from some curves; such
parameters are marked with black stars. The two curves shown in Figure **5** are marked with black arrows.

4. DISCUSSION

The broad goal of this study was to identify variants of the original FvCB model and to compare them using mathematical and experimental approaches. First, we showed that the original FvCB model and the min-A variant disagree about the value of A_n and the rate-limiting process whenever $C < \Gamma^*$ (Section **3.1**). The FRL modification can address this discrepancy in some situations but does not fully reconcile the two variants (Section **3.2**). Second, we demonstrated that unreasonable predictions can occur when the biochemically-derived lower C threshold for TPU limitations, $\Gamma^* \cdot (1 + 3 \cdot \alpha)$, is not included in the equations, or when a fixed threshold is used instead (Section **3.3**). Third, we showed that substantial Rubisco deactivation and RuBP depletion can be included in the original FvCB model, but that the min-A variant cannot accommodate these processes for $C < \Gamma^*$ (Section **3.4**). Finally, we demonstrated that these issues prevent the min-A+FTT variant from closely fitting measured CO_2 response curves that include points where $C < \Gamma^*$, a potential source of error when estimating FvCB parameter values (Section **3.5**).

Half of the curves fit in this study exhibited signs of substantial Rubisco deactivation or RuBP depletion for $C < \Gamma^*$, requiring Equation **6** rather than Equation **1** to achieve a good fit. At present, it is not possible to predict whether a particular leaf will exhibit A_d limitations, making Equations **1** and **6** useful for describing curves but not necessarily for simulating assimilation rates when $C < \Gamma^*$. It is likely that the simplistic treatment of Rubisco deactivation and RuBP depletion in Equation **6** can be improved, enabling predictive abilities and producing even better agreement with measurements. Yet, because the min-A variant never allows these processes to limit assimilation at low C, it is unlikely to ever agree with the measured curves.

Parameter estimation from experimentally measured CO_2 response curves is a major application of the FvCB model. Whenever a curve fitting tool uses the min-A, min-A+FRL, or min-A+FTT variants rather than the original FvCB model, there is a potential for errors in the estimated parameter values. These errors can occur whenever any of the points along the curve are measured in conditions where the original FvCB model disagrees with the other variants. This is likely to happen in the following situations:

• If the curve extends to low CO_2 concentrations (C roughly below 40 µbar), some of the
measured points may satisfy $C < \Gamma^*$, and the min-A variant will assign the wrong
limiting process in this range (Section **3.1**). This issue persists even when considering
substantial Rubisco deactivation and RuBP depletion (Section **3.4**).
• If the curve extends to low CO_2 concentrations and was measured with low light levels
around Q_{in} = 100 µmol m^{-2} s^{-1} (such that assimilation is limited by RuBP regeneration
across the entire measured range), then the min-A variant will predict a sensitivity of
the fit to $V_{c,max}$, even though $V_{c,max}$ would have no influence on the fit using the original
FvCB model. This error will persist even when adding the FRL modification (Section **3.2**).
• If the curve exhibits strong TPU limitations with reverse sensitivity, a nonzero value of α
will be necessary. This will cause divergent behavior in the min-A variant at lower CO_2
concentrations, preventing a good fit (Section **3.3**).
• If the curve exhibits strong TPU limitations where $A_n = A_p$ for C below 400 µbar, the
min-A+FTT variant will not be able to represent this behavior (Section **3.3**).

Here we show that estimates of J and $V_{c,max}$ made using the min-A+FTT variant are typically
lower than those from the original FvCB model, with underestimates exceeding 20%. Because
the differences between the variants only appear for certain environmental conditions and
plant characteristics, the only way to assess errors introduced by the min-A variant for a
particular CO_2 response curve is to compare parameter estimates against those made using the
original FvCB model. Rather than attempting to quantify errors in a range of situations, it is
simpler to always use the original FvCB model.

Across the scientific literature, most CO_2 response curves are measured under high light and
may only contain one or two points at very low CO_2 concentrations. It is also rare for TPU to be
present for C below 400 µbar. Thus, min-A variant usage is not likely to have led to significant
errors in published work. It is also possible that some researchers intentionally avoid the
complexities of fitting curves with points where $C < \Gamma^*$, either by only measuring points at
higher CO_2 set-points, or by excluding potentially problematic points (e.g., where $C_i < \Gamma^*$ or
$A_n < 0$) when fitting. Yet, this practice is not documented, even in recent comprehensive

guides to photosynthetic gas exchange measurements such as Busch et al. (2024). Regardless, a

lack of errors in published work does not justify using the min-*A* variant.

Furthermore, as research trends change and new research areas are formed, the likelihood of

encountering some of these problematic situations may become larger. For example, a recently

developed method for estimating leaf cuticle conductance using gas exchange measurements

actually necessitates the measurement of CO_2 response curves under low irradiance (Márquez

et al., 2022), a situation where the min-*A* and min-*A*+FRL variants can produce incorrect

parameter estimates. The Laisk method for estimating C_i^* and R_L also requires low-light CO_2

response curves (Laisk, 1977; Walker and Ort, 2015). Efforts to improve crop yield and food

security by engineering plants with new Rubisco homologs may create situations where

Rubisco-related parameters like Γ^* and K_c take values significantly different from the ones in

Table **1** (Amaral et al., 2024; Carmo-Silva et al., 2015; Long et al., 2015; Parry et al., 2013). The

most extreme situations may occur in the field of astrobiology, where researchers study

photosynthesis in hypothetical extraterrestrial conditions with ambient gas environments and

biochemical parameters that may be very different from those found on Earth (Covone et al.,

2021; Lehmer et al., 2021; Lingam and Loeb, 2019; Mullan and Bais, 2018). It is therefore more

important than ever for plant scientists to clearly describe their curve fitting methods, including

the equations used (either original FvCB or min-*A*), whether any measured points satisfied $C <$

Γ^*, and whether any points at low C_i or A_n were excluded from fits.

Apart from practical issues, the original FvCB model is based on well-founded knowledge of the

system it describes, and teaching it is tantamount to describing what is known about leaf

biochemistry. In contrast, using and describing the min-*A* variant requires one to explain how it

deviates from what is already known about CO_2 assimilation, which clearly is a hindrance to

teaching and understanding.

Appearing in approximately half of the most cited scientific articles and curve fitting tools that

discuss or use the FvCB model, the min-*A* variant has the potential to prevent clear

communication between scientists (who may be unknowingly using different model equations)

and to interfere with the process of interpreting experimental measurements. The two model

variants may also provide a source of confusion for researchers and students who are new to

the field. Most important, the min-*A* variant is logically inconsistent and disagrees with gas

exchange measurements in situations where the original FvCB model works well. For these

reasons, the original FvCB model – including the biochemically-derived lower threshold for TPU

limitations, $\Gamma^* \cdot (1 + 3 \cdot \alpha)$ – should be used in place of the other variants whenever possible.

One way to help achieve this goal is to encourage greater use of open source software options

with clearly written equations that can easily be validated by other researchers (Duursma,

2015; Holzworth et al., 2014; Lochocki et al., 2022; Stinziano et al., 2021). Simply drawing

attention to this issue will allow the broader photosynthesis research community to move past

it, potentially resulting in faster and more meaningful improvements to CO_2 assimilation

models.

ACKNOWLEDGEMENTS

The authors thank Stephen P. Long, Carl J. Bernacchi, Coralie E. Salesse-Smith, Yi Xiao, and Scott

Rohde for helpful discussions.

This work was supported by Bill & Melinda Gates Agricultural Innovations grant investment ID

57248.

Any opinions, findings, and conclusions or recommendations expressed in this publication are

those of the authors and do not necessarily reflect the views of the US Department of

Agriculture. Mention of trade names or commercial products in this publication is solely for the

purpose of providing specific information and does not imply recommendation or endorsement

by the US Department of Agriculture. USDA is an equal opportunity provider and employer.

COMPETING INTERESTS

The authors declare no competing interests.

AUTHOR CONTRIBUTIONS

E.B.L. contributed to the conceptualization, modeling, data analysis and writing (original draft).

J.M.M. contributed to conceptualization and revising the manuscript.

DATA AVAILABILITY

631 All data and R scripts for reproducing all analysis contained in this work are available online at

632 https://github.com/ripeproject/FvCB-min-A.

SUPPORTING INFORMATION

634 The supporting information includes Section **S1** (*Superficial Variations in FvCB Model Equations*

635 *and Nomenclature*), Section **S2** (*Literature Survey*), Section **S3** (*CO$_2$ Response in the Original*

636 *FvCB Model*) Section **S4** (*Sequences of Limiting Processes for High Γ^* and/or Low K_c*), and

637 Section **S5** (*Fitting Experimentally Measured CO$_2$ Response Curves*).

Works Cited

Amaral, J., Lobo, A.K.M., Carmo-Silva, E., 2024. Regulation of Rubisco activity in crops. New Phytologist 241, 35–51. https://doi.org/10.1111/nph.19369

Bellasio, C., Beerling, D.J., Griffiths, H., 2016. An Excel tool for deriving key photosynthetic parameters from combined gas exchange and chlorophyll fluorescence: theory and practice. Plant, Cell & Environment 39, 1180–1197. https://doi.org/10.1111/pce.12560

Bernacchi, C.J., Morgan, P.B., Ort, D.R., Long, S.P., 2005. The growth of soybean under free air [CO2] enrichment (FACE) stimulates photosynthesis while decreasing in vivo Rubisco capacity. Planta 220, 434–446. https://doi.org/10.1007/s00425-004-1320-8

Busch, F.A., 2020. Photorespiration in the context of Rubisco biochemistry, CO2 diffusion and metabolism. The Plant Journal 101, 919–939. https://doi.org/10.1111/tpj.14674

Busch, F.A., Ainsworth, E.A., Amtmann, A., Cavanagh, A.P., Driever, S.M., Ferguson, J.N., Kromdijk, J., Lawson, T., Leakey, A.D.B., Matthews, J.S.A., Meacham-Hensold, K., Vath, R.L., Vialet-Chabrand, S., Walker, B.J., Papanatsiou, M., 2024. A guide to photosynthetic gas exchange measurements: Fundamental principles, best practice and potential pitfalls. Plant, Cell & Environment 47, 3344–3364. https://doi.org/10.1111/pce.14815

Busch, F.A., Sage, R.F., 2017. The sensitivity of photosynthesis to O2 and CO2 concentration identifies strong Rubisco control above the thermal optimum. New Phytologist 213, 1036–1051. https://doi.org/10.1111/nph.14258

Busch, F.A., Sage, R.F., Farquhar, G.D., 2018. Plants increase CO2 uptake by assimilating nitrogen via the photorespiratory pathway. Nature Plants 4, 46–54. https://doi.org/10.1038/s41477-017-0065-x

Carmo-Silva, E., Scales, J.C., Madgwick, P.J., Parry, M. a. J., 2015. Optimizing Rubisco and its regulation for greater resource use efficiency. Plant, Cell & Environment 38, 1817–1832. https://doi.org/10.1111/pce.12425

Collatz, G.J., Ball, J.T., Grivet, C., Berry, J.A., 1991. Physiological and environmental regulation of stomatal conductance, photosynthesis and transpiration: a model that includes a laminar boundary layer. Agricultural and Forest Meteorology 54, 107–136. https://doi.org/10.1016/0168-1923(91)90002-8

Collatz, G.J., Berry, J.A., Farquhar, G.D., Pierce, J., 1990. The relationship between the Rubisco reaction mechanism and models of photosynthesis*. Plant, Cell & Environment 13, 219–225. https://doi.org/10.1111/j.1365-3040.1990.tb01306.x

Covone, G., Ienco, R.M., Cacciapuoti, L., Inno, L., 2021. Efficiency of the oxygenic photosynthesis on Earth-like planets in the habitable zone. Monthly Notices of the Royal Astronomical Society 505, 3329–3335. https://doi.org/10.1093/mnras/stab1357

De Souza, A.P., Wang, Y., Orr, D.J., Carmo-Silva, E., Long, S.P., 2020. Photosynthesis across African cassava germplasm is limited by Rubisco and mesophyll conductance at steady state, but by stomatal conductance in fluctuating light. New Phytologist 225, 2498–2512. https://doi.org/10.1111/nph.16142

Dubois, J.-J.B., Fiscus, E.L., Booker, F.L., Flowers, M.D., Reid, C.D., 2007. Optimizing the statistical estimation of the parameters of the Farquhar–von Caemmerer–Berry model

of photosynthesis. New Phytologist 176, 402–414. https://doi.org/10.1111/j.1469-
8137.2007.02182.x
Duursma, R.A., 2015. Plantecophys - An R Package for Analysing and Modelling Leaf Gas
Exchange Data. PLOS ONE 10, e0143346. https://doi.org/10.1371/journal.pone.0143346
Farquhar, G.D., von Caemmerer, S., 1982. Modelling of Photosynthetic Response to
Environmental Conditions, in: Lange, O.L., Nobel, P.S., Osmond, C.B., Ziegler, H. (Eds.),
Physiological Plant Ecology II: Water Relations and Carbon Assimilation, Encyclopedia of
Plant Physiology. Springer, Berlin, Heidelberg, pp. 549–587.
https://doi.org/10.1007/978-3-642-68150-9_17
Farquhar, G.D., von Caemmerer, S., Berry, J.A., 2001. Models of Photosynthesis. Plant
Physiology 125, 42–45. https://doi.org/10.1104/pp.125.1.42
Farquhar, G.D., von Caemmerer, S., Berry, J.A., 1980. A biochemical model of photosynthetic
CO2 assimilation in leaves of C3 species. Planta 149, 78–90.
https://doi.org/10.1007/BF00386231
Gu, L., Pallardy, S.G., Tu, K., Law, B.E., Wullschleger, S.D., 2010. Reliable estimation of
biochemical parameters from C3 leaf photosynthesis–intercellular carbon dioxide
response curves. Plant, Cell & Environment 33, 1852–1874.
https://doi.org/10.1111/j.1365-3040.2010.02192.x
Harley, P.C., Sharkey, T.D., 1991. An improved model of C3 photosynthesis at high CO2:
Reversed O2 sensitivity explained by lack of glycerate reentry into the chloroplast.
Photosynth Res 27, 169–178. https://doi.org/10.1007/BF00035838
He, Y., Matthews, M.L., 2023. Seasonal climate conditions impact the effectiveness of
improving photosynthesis to increase soybean yield. Field Crops Research 296, 108907.
https://doi.org/10.1016/j.fcr.2023.108907
Holzworth, D.P., Huth, N.I., deVoil, P.G., Zurcher, E.J., Herrmann, N.I., McLean, G., Chenu, K.,
van Oosterom, E.J., Snow, V., Murphy, C., Moore, A.D., Brown, H., Whish, J.P.M., Verrall,
S., Fainges, J., Bell, L.W., Peake, A.S., Poulton, P.L., Hochman, Z., Thorburn, P.J., Gaydon,
D.S., Dalgliesh, N.P., Rodriguez, D., Cox, H., Chapman, S., Doherty, A., Teixeira, E., Sharp,
J., Cichota, R., Vogeler, I., Li, F.Y., Wang, E., Hammer, G.L., Robertson, M.J., Dimes, J.P.,
Whitbread, A.M., Hunt, J., van Rees, H., McClelland, T., Carberry, P.S., Hargreaves,
J.N.G., MacLeod, N., McDonald, C., Harsdorf, J., Wedgwood, S., Keating, B.A., 2014.
APSIM – Evolution towards a new generation of agricultural systems simulation.
Environmental Modelling & Software 62, 327–350.
https://doi.org/10.1016/j.envsoft.2014.07.009
Kirschbaum, M.U.F., Farquhar, G.D., 1984. Temperature Dependence of Whole-Leaf
Photosynthesis in Eucalyptus pauciflora Sieb. Ex Spreng. Australian Journal of Plant
Physiology 11, 519–538. https://doi.org/10.1071/pp9840519
Laisk, A.K., 1977. Kinetics of photosynthesis and photorespiration of C3 plants. Nauka.
Lehmer, O.R., Catling, D.C., Parenteau, M.N., Kiang, N.Y., Hoehler, T.M., 2021. The Peak
Absorbance Wavelength of Photosynthetic Pigments Around Other Stars From Spectral
Optimization. Frontiers in Astronomy and Space Sciences 8.
Lingam, M., Loeb, A., 2019. Photosynthesis on habitable planets around low-mass stars.
Monthly Notices of the Royal Astronomical Society 485, 5924–5928.
https://doi.org/10.1093/mnras/stz847

Lochocki, E.B., 2024. PhotoGEA: Photosynthetic Gas Exchange Analysis v1.1.0.

Lochocki, E.B., Rohde, S., Jaiswal, D., Matthews, M.L., Miguez, F., Long, S.P., McGrath, J.M.,
2022. BioCro II: a Software Package for Modular Crop Growth Simulations. in silico
Plants diac003. https://doi.org/10.1093/insilicoplants/diac003

Lochocki, E.B., Salesse-Smith, C.E., McGrath, J.M., 2025. PhotoGEA: An R Package for Closer
Fitting of Photosynthetic Gas Exchange Data With Non-Gaussian Confidence Interval
Estimation. Under review.

Long, S.P., Marshall-Colon, A., Zhu, X.-G., 2015. Meeting the Global Food Demand of the Future
by Engineering Crop Photosynthesis and Yield Potential. Cell 161, 56–66.
https://doi.org/10.1016/j.cell.2015.03.019

Márquez, D.A., Stuart-Williams, H., Farquhar, G.D., Busch, F.A., 2022. Cuticular conductance of
adaxial and abaxial leaf surfaces and its relation to minimum leaf surface conductance.
New Phytologist 233, 156–168. https://doi.org/10.1111/nph.17588

Matthews, M.L., Marshall-Colón, A., McGrath, J.M., Lochocki, E.B., Long, S.P., 2022. Soybean-
BioCro: a semi-mechanistic model of soybean growth. in silico Plants 4, diab032.
https://doi.org/10.1093/insilicoplants/diab032

Moualeu-Ngangue, D.P., Chen, T.-W., Stützel, H., 2017. A new method to estimate
photosynthetic parameters through net assimilation rate–intercellular space CO2
concentration (A–Ci) curve and chlorophyll fluorescence measurements. New
Phytologist 213, 1543–1554. https://doi.org/10.1111/nph.14260

Mullan, D.J., Bais, H.P., 2018. Photosynthesis on a Planet Orbiting an M Dwarf: Enhanced
Effectiveness during Flares. ApJ 865, 101. https://doi.org/10.3847/1538-4357/aadfd1

Parry, M.A.J., Andralojc, P.J., Scales, J.C., Salvucci, M.E., Carmo-Silva, A.E., Alonso, H., Whitney,
S.M., 2013. Rubisco activity and regulation as targets for crop improvement. Journal of
Experimental Botany 64, 717–730. https://doi.org/10.1093/jxb/ers336

Sage, R.F., Cen, Y.-P., Li, M., 2002. The activation state of Rubisco directly limits photosynthesis
at low CO2 and low O2 partial pressures. Photosynthesis Research 71, 241–250.
https://doi.org/10.1023/A:1015510005536

Sharkey, T.D., 2019. Is triose phosphate utilization important for understanding
photosynthesis? Journal of Experimental Botany 70, 5521–5525.
https://doi.org/10.1093/jxb/erz393

Sharkey, T.D., 1985. Photosynthesis in intact leaves of C3 plants: Physics, physiology and rate
limitations. Bot. Rev 51, 53–105. https://doi.org/10.1007/BF02861058

Sharkey, T.D., Bernacchi, C.J., Farquhar, G.D., Singsaas, E.L., 2007. Fitting photosynthetic carbon
dioxide response curves for C3 leaves. Plant, Cell & Environment 30, 1035–1040.
https://doi.org/10.1111/j.1365-3040.2007.01710.x

Stinziano, J.R., Roback, C., Sargent, D., Murphy, B.K., Hudson, P.J., Muir, C.D., 2021. Principles of
resilient coding for plant ecophysiologists. AoB PLANTS 13, plab059.
https://doi.org/10.1093/aobpla/plab059

Taylor, S.H., Gonzalez-Escobar, E., Page, R., Parry, M.A.J., Long, S.P., Carmo-Silva, E., 2022.
Faster than expected Rubisco deactivation in shade reduces cowpea photosynthetic
potential in variable light conditions. Nat. Plants 1–7. https://doi.org/10.1038/s41477-
021-01068-9

Von Caemmerer, S., 2013. Steady-state models of photosynthesis. Plant, Cell & Environment 36,
1617–1630. https://doi.org/10.1111/pce.12098
von Caemmerer, S., 2000. Biochemical Models of Leaf Photosynthesis. CSIRO Publishing.
https://doi.org/10.1071/9780643103405
von Caemmerer, S., Evans, J.R., Hudson, G.S., Andrews, T.J., 1994. The kinetics of ribulose-1,5-
bisphosphate carboxylase/oxygenase in vivo inferred from measurements of
photosynthesis in leaves of transgenic tobacco. Planta 195, 88–97.
https://doi.org/10.1007/BF00206296
Walker, A.P., Johnson, A.L., Rogers, A., Anderson, J., Bridges, R.A., Fisher, R.A., Lu, D., Ricciuto,
D.M., Serbin, S.P., Ye, M., 2021. Multi-hypothesis comparison of Farquhar and Collatz
photosynthesis models reveals the unexpected influence of empirical assumptions at
leaf and global scales. Global Change Biology 27, 804–822.
https://doi.org/10.1111/gcb.15366
Walker, B.J., Ort, D.R., 2015. Improved method for measuring the apparent CO2
photocompensation point resolves the impact of multiple internal conductances to CO2
to net gas exchange. Plant, Cell & Environment 38, 2462–2474.
https://doi.org/10.1111/pce.12562
Wang, Q., Chun, J.A., Fleisher, D., Reddy, V., Timlin, D., Resop, J., 2017. Parameter Estimation of
the Farquhar—von Caemmerer—Berry Biochemical Model from Photosynthetic Carbon
Dioxide Response Curves. Sustainability 9, 1288. https://doi.org/10.3390/su9071288
Wu, A., Brider, J., Busch, F.A., Chen, M., Chenu, K., Clarke, V.C., Collins, B., Ermakova, M., Evans,
J.R., Farquhar, G.D., Forster, B., Furbank, R.T., Groszmann, M., Hernandez-Prieto, M.A.,
Long, B.M., Mclean, G., Potgieter, A., Price, G.D., Sharwood, R.E., Stower, M., van
Oosterom, E., von Caemmerer, S., Whitney, S.M., Hammer, G.L., 2023. A cross-scale
analysis to understand and quantify the effects of photosynthetic enhancement on crop
growth and yield across environments. Plant, Cell & Environment 46, 23–44.
https://doi.org/10.1111/pce.14453
Xiao, Y., Sloan, J., Hepworth, C., Osborne, C.P., Fleming, A.J., Chen, X., Zhu, X.-G., 2021.
Estimating uncertainty: A Bayesian approach to modelling photosynthesis in C3 leaves.
Plant, Cell & Environment 44, 1436–1450. https://doi.org/10.1111/pce.13995
Yin, X., Busch, F.A., Struik, P.C., Sharkey, T.D., 2021. Evolution of a biochemical model of steady-
state photosynthesis. Plant, Cell & Environment 44, 2811–2837.
https://doi.org/10.1111/pce.14070
Yin, X., Struik, P.C., 2017. Can increased leaf photosynthesis be converted into higher crop mass
production? A simulation study for rice using the crop model GECROS. Journal of
Experimental Botany 68, 2345–2360. https://doi.org/10.1093/jxb/erx085
803