1 **Title:** Spontaneous eye movements reveal that premotor cortex is involved in
2 human thinking
3

4 **Authors:** Artur Pilacinski[1,2,$,*], Lia Wagner[1,$], Gabriel Besson[2], Maria Joao Matos[2], Eduardo Araujo[2],
5 Christian Klaes[1]
6
7 [$] Equal contribution
8
9 **Affiliations:** [1]Medical Faculty, Ruhr University Bochum, Bochum, Germany; [2]Faculty of Psychology
10 and Educational Sciences, University of Coimbra, Coimbra, Portugal
11
12 ***Corresponding author:** AP, art.pilacinski@gmail.com

13
Abstract:

Non-visual saccades (NVS) are spontaneous eye movements humans make while thinking. Their
role is puzzling and there is no framework to explain them in terms of their underlying neural
systems. Here we studied neural activity preceding NVS when subjects performed standardized
abstract thinking tasks. We used high-density EEG and focused on frontal and parietal channels
representing oculomotor cortical areas. We found that NVS are preceded by neural activity
changes in frontal and parietal EEG channels qualitatively different to that for voluntary saccades.
However, only the rate of signal change in frontal channels, but not in parietal channels,
surpassed that observed when comparing NVS to normal saccades. Frequency spectrum analyses
showed activity patterns reflecting complex top-down and bottom-up processing. A source
reconstruction analysis additionally revealed that presaccadic signals originated mainly in the
premotor cortex. We propose that this presaccadic activity while thinking represents evidence
accumulation and attentional shifts, as shown before in the premotor cortex during sensorimotor
tasks. Our findings suggest that human thinking appears to function similarly to human actions
and employs premotor areas responsible for evaluating and manipulating objects also when
evaluating and manipulating concepts.

Main Text:

Introduction

Take a moment to answer the following questions: What do apples and oranges have in common? What is similar between crime and punishment? Why do we pay taxes? You might have noticed that, before answering, you probably made a few eye movements as if in search of the right answer. It is a known but often ignored fact that such spontaneous, non-visual saccadic eye movements (NVS) accompany human thinking, that is mental task solving and mnemonic retrieval processes (Ehrlichman & Micic, 2012). Why they occur and what function they serve remains unknown. Here, we show that NVS are immediately preceded by substantial changes in frontal regions including premotor cortex, indicating the crucial role of that area in conceptual operations (thinking).

Non-visual saccades remain a very understudied phenomenon and the existing research does not allow for building a coherent picture of their nature. Contrary to some opinions (Doherty-Sneddon and Phelps, 2005) NVS do not subserve others' gaze aversion as they are also performed in darkness or without other people present (Ehrlichman & Barrett, 1983). Restriction of NVS does not seem to impact memory retrieval performance (Micic et al., 2010) suggesting they may be an epiphenomenon rather than a functional process (for an alternative view see: Scholz et al., 2016). They are, however, not fully suppressible and people move their eyes while thinking even when instructed to fixate gaze (Micic et al., 2010) indicating that NVS are automatically accompanying memory operations. On the other hand, non-visual saccades have been demonstrated to reflect mnemonic content organization such as spatial arrangement of items (Johansson et al., 2022; Damiano and Walther, 2019; Scholz et al., 2016) or the point in time (Hartmann et al., 2014; Martarelli et al., 2016). Although one could suspect these NVS to merely reflect visuospatial organization of memorized items, it comes as a surprise that substantially more saccades are produced when operating on verbal than on visual material (Weiner and Ehrlichmann, 1976). This indicates that NVS do not directly reflect reactivating the visual scene in memory but rather other processes such as activation of conceptual knowledge necessary for processing semantics.

Organization of conceptual knowledge has been long attributed to medial temporal lobe (MTL) function (Bellmund et al., 2018; Constantinescu et al., 2016). The main cortical centers controlling saccades span across posterior parietal (PPC) and premotor cortex (PM), comprising frontal eye fields (FEF), the main area of the brain responsible for programming voluntary eye movements (Munoz, 2002). While there are some interactions between the MTL and eye movements systems, there is no mechanism that could possibly explain NVS. For example, activity in MTL reflects integrating spatial information about objects (Yeung et al., 2017), hippocampal maps are modulated by gaze direction (Meister et al., 2018) and parietal activity reflects that of the hippocampus (Whitlock et al., 2008). It is unclear, however, how retrieving mnemonic content could possibly initiate saccades.

The core cortical regions for programming voluntary saccades are posterior parietal and prefrontal cortical areas (Pierrot-Deseilligny et al., 1995; Andersen & Buneo, 2002). Pre-saccadic planning activity was described earlier as predominantly visible in parietal EEG channels (Everling

et al., 1997; Richards, 2003; Becker et al., 1972; Balaban & Weinstein, 1985). Posterior parietal
cortex (PPC) has been proposed to integrate between memory and actions (Whitlock et al., 2008;
Whitlock, 2017), and parietal oculomotor areas (Schluppeck et al., 2005) overlap with those
involved in long term memory search (Berryhill et al., 2007; Vilberg and Rugg, 2008). PPC is also
directly connected to superior colliculus (Pare and Wurtz, 1997), which suggests parietal activity
may be triggering NVS as a byproduct of memory search processes. Whether it is the memory
retrieval activity in parietal oculomotor areas that underlies NVS is unknown.

Posterior parietal cortex typically projects oculomotor programming signals to the frontal eye
field, the main saccade-controlling area in the primate brain (Bruce et al., 1985). In humans, FEF is
contained within the dorsal premotor cortex (Darby et al., 1996), indicating shared function of the
two areas. Premotor cortex (PM) has been repeatedly demonstrated to be involved in a number
of non-motor tasks, including working memory (Curtis et al., 2004; Curtis, 2006; Lindner 2010;
Pilacinski et al., 2020), evaluating evidence during decision-making (Thura et al., 2012) and
internal shifts of attention (Peelen et al., 2004). The premotor theory of attention further
suggests that these internal shifts of attention reutilize saccade programming circuitry (Rizzolatti
et al., 1996). Premotor cortex has also been shown to be involved in memory search (Kapur et al.,
1995).

The organization of conceptual knowledge has long been suggested to reutilize sensorimotor
computations, e.g. through representing cognitive spaces in a similar way as physical space
(Tolman, 1948; Bellmund et al., 2018). In our view, non-visual saccades are a window into the
evolution of human thinking by providing a link between the memory and the action systems in
the human brain. The available evidence suggests that the frontal and parietal cortex may be
involved in generation of NVS during memory search, although the particular involvement of
each of these areas has not been explored. It is likewise unknown how neural mechanisms
underlying NVS compare to typical programming of goal-directed saccades. Lastly, it is unclear
whether NVS are a function or merely a byproduct of human thinking.

Here, we tackled these questions and investigated neural signals immediately preceding non-
visual saccades. We used high-density EEG to record brain activity while volunteers were solving
abstract reasoning and knowledge tasks, with restricted and unrestricted gaze and with open and
structured memory search conditions. We compared the presaccadic activity for NVS with
planning activity for voluntary saccades. We reasoned that if NVS are triggered by parietal cortex
activity during memory search, then we should observe a substantial change in signals
immediately preceding NVS in parietal channels, similar to presaccadic activity reported earlier.
Conversely, if NVS were a product of additional processing of conceptual information, like
internal shifts of attention, they should be preceded by a change in frontal channels' activity.
Finally, if non-visually saccades arise from distinct neural processing compared to voluntary
saccadic preparation, the neural activity preceding them should substantially differ from that
preceding voluntary saccades.

121
122
123

Methods

Subjects

A total of 57 individuals (mean age 27.2 ± 8.1 years, 32 male, 5 left-handed) took part in the study.
All participants had normal or corrected-to-normal vision with no history of neurological or
psychiatric disorders. Participants were informed about the study procedures, were naive to the
experimental hypothesis and received no financial compensation for their participation. Before
participation, all subjects gave written informed consent in accordance with the Declaration of
Helsinki and local ethical guidelines. The procedures were approved by the local Ethical
Committee at the Faculty of Psychology and Educational Sciences of the University of Coimbra. All
datasets were anonymized before data collection began. Twelve participants were excluded from
all analyses due to missing data or poor data quality as described in the supplement, resulting in
an overall sample size of 45 participants (mean age 27.2 ± 8.0, 23 male, 4 left-handed) for the
behavioral analysis. For the EEG analysis, further exclusion criteria based on EEG signal quality
were applied as described in the supplement, resulting in a final sample size of 29 participants
(mean age 26.9 ± 7.0 years, 11 male, none left-handed). Due to a technical problem with data
logging, there were just 12 subjects considered in analyses of the memory saccade control task
(See Supplementary Methods).

Task

In the main experimental task, participants were supposed to answer forty questions taken from
the Portuguese adaptation of Wechsler Adult Intelligence Scale-III (Wechsler, 1997). This task
allowed us to have normalized stimuli, arranged progressively according to their difficulty and
abstractness levels. The questions were presented verbally, through headphones (Fig. 1 A). The
questions were split into "structured" and "open" according to the type of memory search they
involved: 20 questions from the Similarities subtest were considered "structured"; 10 questions
from the Information subtest, and 10 from the Comprehension subtest were considered "open".
The Information subtest evaluates long-term memory and general knowledge, e.g., "At what
temperature does water boil under normal circumstances?" The Comprehension subtest
assesses practical judgments, e.g., "Why do we pay taxes?" These were considered "open"
questions due to the unrestricted memory search required. The Similarities subtest assesses
abstract verbal reasoning by finding similarities between paired concepts, e.g., "How are an
orange and a banana comparable?". These were considered "structured" questions. Further
examples can be found in the supplementary materials (see Supplementary Table 1). Collectively,
these subtests assess verbal conceptualization, vocabulary, abstract reasoning, categorical
thinking, logical abstraction, and social intelligence, reflecting a broad range of cognitive
functions and culturally acquired knowledge. The forty questions were divided into four blocks of
10 trials each, with randomized order within each block, and increasing abstraction across blocks.
In each trial, a question was played through headphones and subjects were instructed to answer
as best as they could. Correctness of answers was manually assessed according to the test
instructions and saved in a binary form (correct/incorrect). The only displayed item during this
task was a centrally-positioned bright fixation cross on a dark grey background. Subjects were

randomly assigned to two groups: "fixed gaze" in which they were told to keep looking at the
fixation cross and "free gaze" in which they were not instructed to do so.
A control task was used to assess goal-directed saccadic programming, as well as to differentiate
between visual and memory saccade programming components (Fig. 1 B). The task was split into
the following conditions: visual vs. memory saccades and pro- vs. antisaccades. In the visual
condition, participants were instructed to make saccades either toward (prosaccade) or away
from (antisaccade) a blinking dot located in the periphery. In the memory condition, participants
were asked to make saccades to the location where a target dot had been before it disappeared.
In both conditions, the saccade had to be executed after a delay (when the fixation cross
disappeared).

175
Figure 1. Experimental tasks timeline. A) Main task. The question was played through
headphones; next, the participant was given time to think of an answer; after they responded, the
next trial started. B) Control tasks: subjects were instructed to saccade to (green cue) or away
from (red cue) the cued location (white dot). The saccade should be executed either towards the
visible (VST) or memorized (MST) target. Blue arrows indicate saccade epoch. Objects not to
scale.

Eyetracking
Eye movements were recorded at 100 Hz using the HTC VIVE Pro Eye, which has an embedded
eye-tracker in a VR head-mounted display and allows for accurate recording of eye data in
immersive environments (Sipatchin et al., 2021). The VR environments were developed to serve
as experimental tasks (Fig. 1 A, B), and an eye tracking algorithm was implemented in Unity with
the use of SRanipal software development kit from HTC.
For the behavioral oculomotor analysis, saccade and blink frequencies, pupil diameter, saccade
amplitudes, peak velocities, and angles were extracted for each trial and participant, focusing on
the period after each question until the end of each trial (See Supplementary Methods).
Parameters were averaged across combinations of factors for each participant, specifically blocks
1-4, condition type (fixed/free), and question type (open/structured). To assess eye parameters
during the cognitive engagement, GLMM analyses were used as an alternative to the repeated
measures ANOVA, with the fixed factors block (1-4), condition type (fixed/free) and question type
(open/structured). Subjects were considered as the random effects grouping factor. GLMM

analyses were performed separately for the dependent variable answer duration, saccade
frequency, saccade amplitude, blink frequency and pupil average. Only 'correct' trials were
included in the statistical analysis to minimize confounds in an already complex statistical model
and ensure a clearer interpretation of cognitive engagement. The final trial count in the EEG
analyses for the main tasks was 2884 (mean 100 ± 52.26 per participant), and for the control tasks
was 2168 (mean 94 ± 11.95 per participant).
203
EEG Recordings
EEG data was recorded using either a 64-electrode dry or a 256-electrode wet EEG system (ANT
Neuro, Enschede, Netherlands). Initially, the 256-channel system was used for the first eleven
participants, but due to technical issues, the 64-electrode system was used for the remaining
eighteen participants. There were no statistically significant differences between both systems in
terms of signal quality that would influence our results (See Supplementary Methods). Electrodes
were arranged in an equidistant montage with the left and right mastoids as online ground and
reference for the 64-channel setup, and CPz and the left mastoid for the 256-channel setup.
Impedances were kept below 25 kΩ, and signals were recorded at 500 Hz. Preprocessing was
done with EEGLAB and MNE-Python, including checks for sampling rate stability, application of a
1 Hz high-pass filter and 40 Hz low-pass filter, and removal of 25 Hz environmental noise using a
notch filter. A 50 Hz notch filter removed remaining line noise. Flatline channels were removed
using EEGLAB's pop_clean_rawdata function, and noisy channels were rejected using
pop_rejchan based on kurtosis, probability, and spectra, followed by manual inspection.
Interpolated channels were re-referenced to the average of all electrodes. ICA was not used to
avoid removing saccade-planning data; instead, eye movements were controlled using eye-
tracking data. EEG events were processed separately for the main and control tasks, and epochs
were extracted from −200 to 100 ms relative to saccade onset. Saccades preceded by other
saccades or blinks within 200 ms were excluded. A baseline correction from -200 to -170 ms was
performed using single-trial normalization. We decided for this time window based on typical
inter-saccadic latencies (Rayner et al., 1983; c.f. Fig 2 C). This allowed us to scrutinize signal
changes immediately preceding NVS. Epochs were rejected if oculomotor-related spikes were
detected, and semiautomatic artifact rejection occurred for amplitudes exceeding 50 μV
between samples or 100 μV over 50 ms. Four regions of interest (ROIs), frontal left (FL), frontal
right (FR), parietal left (PL), and parietal right (PR), were defined, distributed around
corresponding electrodes F3, F4, P3 and P4 in the 10–20 electrode system (see Fig. 3 A and Fig. S 1
A and B). Since the FEF and PPC are the primary areas of interest in saccade research and are both
superficial and spatially distinct, this setup allowed us to effectively distinguish neural activity in
these regions.
233
ERP analysis
235 For the EEG analysis, ERP analysis was performed using MNE-Python. An interval of -150 to -20 ms
236 prior to saccade onset was used for statistical evaluation of presaccadic activity (Revankar et al.,
237 2020). The mean amplitude of all epochs within this interval was calculated for each condition
238 and participant. Potentials were then averaged across electrodes within each ROI, allowing for

the assessment of neural dynamics associated with saccade preparation across different ROIs. Grand average ERPs were calculated for ERP visualization. For topographic visualizations of grand average ERPs, all electrodes from both EEG systems were included.

Statistical analysis of ERP data was performed either on each ROI or on combined frontal and parietal ROIs. A Kolmogorov-Smirnov test was used on averaged amplitudes to confirm normality prior to ANOVA analyses. Averaged EEG amplitudes of conditions were statistically compared using 4 × 2 or 2 × 2 repeated measures ANOVAs with ROI (4 ROIs and 2 ROIs, respectively) and condition type as factors. In cases of violation of sphericity (as indicated by Mauchly's test), Huynh-Feldt corrected p-values were used. For post hoc analysis, paired t-tests were conducted with Holm correction for multiple testing. To investigate the temporal dynamics of signal divergence between main and control tasks, the same statistical approach was applied using a sliding window of 10 ms within the previously defined interval. Mean amplitudes were calculated within each window for each participant and condition, and paired t-tests with Holm correction were used to identify the time points at which significant differences emerged.

Frequency spectrum analysis

Spectral decomposition was conducted using EEGLAB to assess neural activity in specific frequency bands associated with saccade preparation. The analysis focused on three frequency bands: alpha (8–12 Hz), beta (13–30 Hz), and gamma (30–45 Hz). Power spectral density (PSD) was computed for the interval of -150 to -20 ms prior to saccade onset. For each condition, PSD was averaged over time and the selected ROIs. Statistical analysis was performed separately for each frequency band on combined frontal and parietal ROIs. Repeated measures ANOVAs were conducted with ROI and task type as factors to evaluate differences in PSD values. In cases of violation of sphericity (as indicated by Mauchly's test), Huynh-Feldt corrected p-values were used. For post hoc analysis, paired t-tests were conducted with Holm correction for multiple testing.

Source localization

The forward model and the ERP for each condition were used to estimate an inverse model using the standardized low-resolution brain electromagnetic tomography (sLORETA) method (Pascual-Marqui, 2002). We extracted source estimates for each condition and computed spatial adjacency in the source space to enable cluster-based permutation testing. We performed a non-parametric, two-tailed cluster-based permutation test, involving voxel-wise comparisons to zero and correcting for multiple testing. We assessed statistical significance using a p-value threshold of 0.05. We projected t-values into sLORETA images to visualize source activity, highlighting regions with significant neural dynamics. We obtained anatomical labels using the FreeSurfer parcellation scheme to map source activity to specific cortical regions, focusing on cortical involvement in saccadic movements. Subcortical areas were not captured due to parcellation limitations. We aligned source estimates with these labels to investigate neural activity in distinct areas. For each significant cluster identified, we extracted the maximum t-value for each label.

Results

Behavioral and oculomotor results

First, a decrease in percentage of known answers can be observed across blocks for structured questions while open questions show a significant drop in block 2 with no overall decrease across blocks. Fewer known answers are reported for structured questions compared to open questions and in the fixed gaze condition compared to the free gaze condition. Additionally, interactions between question type and condition type revealed fewer known answers for the fixed gaze condition for structured questions only. A significant main effect was found of the factor block ($\chi2=53.151, p<0.001$), indicating that performance dropped as abstractness increased. Additionally, significant interaction effects were found for block × question type ($\chi2=112.722, p<0.001$) and question type × condition type ($\chi2=7.987, p=0.005$), suggesting that the effect of block varied across different question types (Fig. 2 A). Post hoc Wilcoxon signed-rank tests revealed a significant difference for block 1 and block 2, block 1 and 4, and block 3 and block 4 ($p<0.001$). Additionally, a significant difference was found between block 1 and block 3 ($p=0.031$). Post hoc Wilcoxon rank-sum tests also showed a difference between fixed and free gaze condition ($p=0.029$). For interactions between question type and condition type, post hoc Wilcoxon rank-sum tests interestingly confirmed a significant difference between open compared to structured questions for the fixed gaze condition only ($p=0.026$), but not for the free gaze condition ($p=1.000$) (Fig 2 B).

Saccade frequency for condition type and question type remains approximately the same, but increases over blocks for both fixed and free gaze conditions (Fig 2 C). For saccade frequency, a significant main effect was found of the factor block ($\chi2=11.500, p=0.009$). Post hoc Wilcoxon signed-rank tests revealed a significant difference between block 1 and 3 ($p=0.017$) and block 2 and 3 ($p=0.043$). However, no significant difference was observed for block 4, including comparisons between block 1 and block 4 ($p=0.055$), showing that while we see an increase in saccade frequency in block 3, the frequency is slightly reduced again in block 4. The post hoc Wilcoxon rank-sum test for condition type showed no significant difference ($p=0.604$).

Saccadic amplitudes slightly decreased across blocks (Fig. 2 D). For different question types, saccadic amplitude remains approximately the same, whereas in the fixed condition, mean saccadic amplitudes are decreased compared to the free gaze condition. For saccadic amplitude, a significant main effect of the factor block was found ($\chi2=10.563, p=0.014$). Also, a significant interaction effect of block × question type × condition type ($\chi2=9.784, p=0.020$) suggests that the effect of blocks varied across different question types as well as condition types. Regarding condition type, the post hoc Wilcoxon rank-sum test revealed a significant difference between fixed and free gaze conditions ($p=0.006$).

An increased pupil diameter as an indicator of cognitive load can be identified across blocks and in the fixed gaze condition compared to the free gaze condition (Fig. 2 E). Additionally, a slight increase in pupil diameter can be noted for open questions compared to structured questions. For pupil diameter, a significant main effect of the factor block ($\chi2=24.720, p<0.001$), question type ($\chi2=16.897, p<0.001$), and condition type ($\chi2=6.750, p=0.009$) was found. Additionally, a significant interaction between block × question type was identified

$(\chi 2=21.756, p<0.001)$. Post hoc Wilcoxon signed-rank tests revealed a significant difference for
block 1 compared to all other blocks $(p<0.001)$. A significant difference was also found between
block 2 and 3 $(p=0.029)$. No significant differences were found between block 2 and 4
$(p=0.194)$ or between block 3 and 4 $(p=0.311)$. These results suggest that pupil diameter
increased across the initial blocks, indicating rising cognitive load or engagement levels, but
slightly decreased again in block 4.
Lastly, for answer duration (Fig 2F), a significant main effect was found of the factor block
$(\chi 2=58.375, p<0.001)$ and question type $(\chi 2=6.074, p=0.014)$. There was also a significant
interaction effect of block × question type $(\chi 2=24.804, p<0.001)$, suggesting that the effect of
blocks varied across different question types. Post hoc Wilcoxon signed-rank tests revealed a
significant difference between block 1 and 2 $(p=0.002)$, and all other blocks $(p<0.001)$, except
block 3 and 4, where no significant difference was found $(p=0.697)$. The post hoc Wilcoxon rank-
sum test for condition type showed no significant difference $(p=0.153)$.

334
Figure 2. Behavioral metrics for different gaze conditions across fixed effects variables. Box plots
show the IQR with medians as horizontal lines. Mean values with 95% confidence intervals are
marked as colored points, and outliers as individual points. Only known response trials were
included (n = 45). A) Normalized correct responses are shown across blocks and B) different
question types. C) Saccade frequency, D) saccade amplitude, E) pupil diameter and F) answer
durations are shown across blocks. Error bars denote upper and lower quartiles, boxes denote
interquartile ranges, solid horizontal bars denote medians;
342
343 Presaccadic potentials were analyzed for main versus control tasks, with the control task
344 including both perception and memory tasks. The analyzed interval (-200 to 50 ms) is referenced
345 within the overall signal (compare Fig. S5 A), as determined based on typical saccadic latency time
346 (Rayner et al., 1983), meaning that the reported effects reflect relative changes rather than
347 absolute increases or decreases in activity. In the specified interval, the main task is characterized

by a presaccadic positive frontal and negative parietal potential, while the control task shows a positive parietal potential and a smaller frontal potential in contrast to the main task. Potentials peak approximately 20 ms prior to saccade onset. At saccade onset, a negative frontal and positive parietal spike can be observed for both task conditions. Averaged presaccadic amplitudes are larger for main saccades in frontal ROIs and smaller for main saccades in parietal ROIs compared to control saccades. A larger polarization is also observed for frontal and parietal ROIs for main saccades compared to control saccades, as confirmed in the topographical visualization. Topographic maps (Fig. 3 E) illustrate the evolving spatial distribution of ERPs leading up to saccade onset. As already indicated by the ERPs, the main task shows a strong positive frontal and negative parietal potential developing over time, peaking just before the saccade. In contrast, the control task exhibits a prominent positive potential in the parietal regions, which extends into the posterior frontal regions. However, overall, the frontal electrodes show less pronounced, negative potential. A larger cluster of activity is visible over the parietal electrodes. This pattern is congruent with previously reported presaccadic EEG activity (Everling et al., 1997; Richards, 2003; Becker et al., 1972; Balaban & Weinstein, 1985). To quantify these activity changes, we used repeated measures ANOVA using ERP amplitudes averaged over the presaccadic time window, with "task" (main/control) and "ROI" (frontal/parietal) as factors. This analysis showed significantly higher activity change preceding saccades during the thinking task in both channel groups. A significant main effect of "task" $(F(1,28)=16.205, p<0.001, \omega 2=0.137)$ and "ROI" $(F(1,28)=5.528, p=0.026, \omega 2=0.114)$ was found with a large and medium to large effect size, respectively. Additionally, a significant interaction effect of "task" × "ROI" $(F(1,28)=25.496, p<0.001, \omega 2=0.411)$ with a large effect size was observed. Post hoc paired t-tests revealed a significant difference between main and control tasks for the frontal channels $(p<0.001)$, as well as for the parietal channels $(p=0.003)$. Significant differences were also identified for the main task between frontal and parietal ROI $(p<0.001)$, while no significant difference was found for the control task between frontal and parietal ROI $(p=0.130)$.

As previous analyses showed that the main and the control tasks had distinct neural components, we next analyzed the magnitude of signal changes preceding saccades without the confounding effects of signal polarity. We compared the absolute values of ERP amplitudes by conducting repeated measures ANOVA with "task" (main/control) and "ROI" (frontal/parietal) as factors. A significant main effect of "task" $(F(1,28)=10.087, p=0.004, \omega 2=0.145)$ was found with a large effect size. No significant main effect was found for "ROI" $(F(1,28)=1.887, p=0.180, \omega 2=0.009)$. Additionally, a significant interaction effect of "task" × "ROI" $(F(1,28)=6.279, p=0.018, \omega 2=0.047)$ with a small effect size was observed. Post hoc paired t-tests revealed a significant difference between main and control tasks for the frontal channels $(p<0.001)$, but no significant difference for the parietal channels $(p=0.265)$. Significant differences were also identified for the main task between frontal and parietal ROI $(p=0.035)$, while no significant difference was found for the control task between frontal and parietal ROI $(p=0.477)$. These analyses show that absolute signal amplitude changes in prefrontal channels in the main task were greater than those in parietal channels and when compared to the saccadic preparatory activity in the control tasks.

The sliding window analysis revealed the time points at which ERP signals significantly separated
between the main and control tasks in the frontal and parietal ROIs. In the frontal ROI, signal
separation was first observed at -140 ms prior to saccade onset $(p=0.038)$. In the parietal ROI,
signal separation occurred later, at -110 ms prior to saccade onset $(p=0.042)$. The results
demonstrate a 30 ms difference between the onset of significant signal change in the frontal and
parietal ROIs, with frontal signal preceding parietal signal. This temporal difference underscores
the distinct timing of neural processes involved in NVS across these regions.
In the next step, we analyzed power spectral density for alpha, beta, and low-gamma bands. For
each band separately, we conducted an ANOVA with factors "Task" and "ROI". These ANOVAs
showed a significant increase in power spectral density in the main task for beta
$(F(1,28)=7.41, p=0.011, \omega2=0.029)$ and gamma bands
$(F(1,28)=9.016, p=0.006, \omega2=0.058)$. There was also a nearly-significant effect of Task in
alpha band $(F(1,28)=4.14, p=0.051, \omega2=0.01)$. There was a significant ROI main effect in
alpha $(F(1,28)=16.205, p<0.001, \omega2=0.137)$ and beta bands
$(F(1,28)=16.205, p<0.001, \omega2=0.137)$. No Task x ROI interaction effect was significant
$p>0.2$.
Lastly, source localization revealed strong, focused neural activity primarily in the right
hemisphere. Figure 3F shows clusters in the frontal regions, with the peak activation in the right
superior and middle frontal cortex (peak t = 9.3). Activity extends posteriorly into the motor
cortex and precentral gyrus, and medially, involving orbitofrontal and medial areas. Notable
activation was also found bilaterally in the superior frontal cortex (t = 8.17), precentral and
postcentral regions (t = 8.21), the insula (t = 8.14), and rostral anterior cingulate (t = 7.98).
Additional activation, though less intense, occurred in the superior parietal cortex (t = 7.95),
medial orbitofrontal cortex (t = 7.88), and precuneus (t = 7.97). This pattern was substantially
different in control tasks (See Supplementary Results).

Figure 3. EEG results for main vs. control tasks. A) ROI channels selection and ERPs relative to

saccade onset for main and the control tasks; dashed lines indicate significant difference between

signals in main and control tasks; grey box denotes the baseline period; shaded areas denote 95%

confidence intervals; B) ERP amplitudes across tasks. C) Absolute values of ERP amplitudes across
conditions. D) Modulation of alpha, beta and gamma frequency across tasks. E) Topographic
maps of presaccadic activity across main and control tasks. F) Source localization. Error bars
denote upper and lower quartiles, boxes denote interquartile ranges, solid horizontal bars
denote medians.
424
425

Discussion

Non-visual saccades are a puzzling phenomenon accompanying conceptual operations. Here, we
show that the neural processes preceding NVS are qualitatively different from how voluntary
saccades are programmed, with the main difference lying in the engagement of the frontal cortex
directly preceding each saccade. NVS are preceded by a substantial signal change in both frontal
and posterior channels, qualitatively different from the activity observed during voluntary
saccade planning. However, only the rate of activity change in frontal channels before NVS
significantly differs from that preceding voluntary saccades. The source localization analysis
additionally indicates that the source of these pre-NVS signal changes is the premotor cortex. The
use of abstract conceptual tasks allowed us to determine that these signal changes preceding NVS
are not due to processing visuospatial content.
436

We believe that the gradual buildup of activity before NVS likely represents neural computations
required for evaluating conceptual information, similar to evidence accumulation during
sensorimotor decision-making (Cisek et al., 2009; Thura and Cisek, 2014). The subsequent
saccade could be due to the shifting of attention to another sub-element of the currently-
processed set of concepts. It has been shown that covert shifts of attention indeed result in
involuntary eye movements, likely due to direct connections between frontal eye fields and the
superior colliculus (Peterson et al., 2004). Complex conceptual operations require multiple such
shifts of attention between concepts or their elements, and these shifts are likely to underlie NVS
(Scholz et al., 2017) in a way similar to how saccades scan the visual space, identifying its
(semantic) components. The premotor theory of attention proposes that covert attentional shifts
and saccade planning processes are identical (Rizzolatti et al., 1987; Smith and Schenk, 2012). This
theory is at odds with observations showing that covert and overt visual attention are managed
by distinct neural populations (Smith and Schenk, 2012), however, both are located closely within
the premotor cortex. Our results suggest that non-visual saccades, putatively resulting from shifts
of attention while thinking, result from different neural processes than voluntary saccade
planning. Nonetheless, the fact that neural activity preceding NVS seems to originate in the
premotor cortex, a common neuroanatomical substrate between NVS and voluntary saccades,
may explain why mental operations on concepts trigger NVS. As frontal eye fields directly project
to the superior colliculus, it is likely that mental operations involving premotor computations may
collaterally initiate eye movements, resulting in NVS being an epiphenomenon of covert
attentional shifts during thinking. The fact that restricting gaze did not substantially worsen
mnemonic performance additionally supports the notion that NVS are an epiphenomenon rather

than a primary function. An alternative theory considering the attentional account proposes that retrieving information from long-term memory requires an attentional shift to memory, disengaging from the visual scene, which in turn results in NVS (Servais et al., 2023). While this view does not seem congruent with the fact that NVS are also executed without vision, it nonetheless emphasizes the vital importance attention has in mnemonic operations.

Whether concepts are represented in pictorial/visuospatial or verbal/abstract form has long been a subject of dispute in philosophy, psychology, and neuroscience (see e.g., Pylyshyn, 2003). Edward Tolman proposed that past experience and spatial cognition are arranged in the form of spatially organized maps (Tolman, 1948). This theory implied that the knowledge map builds on representations of space, and the two likely share neural components. Since then, it has been confirmed that spatial maps are organized by the hippocampal-entorhinal system (O'Keefe & Dostrovsky, 1971) using a grid-like code (Hafting et al., 2005), and that this system also organizes conceptual knowledge using similar principles (Constantinescu et al., 2016). However, just as spatial maps need to be translated into neural codes understood by the motor system, stored conceptual knowledge needs to be processed for daily operations. It has been proposed that the parietal lobe acts as an interface connecting the MTL memory system with areas responsible for actions (Whitlock, 2017). The main role of the PPC seems to lie in integrating representations of world-centered space into body-centered maps used for locomotion and actions (Whitlock et al., 2008). In the cognitive domain, there is also strong evidence showing the parietal lobe's involvement in long-term memory search but not in the recognition of remembered objects, indicating that the parietal cortex is not merely involved in storage but rather in the active exploration of the knowledge space (Cabeza et al., 2008). Motor (and oculomotor) areas in the posterior parietal cortex share strong reciprocal connections with the premotor cortex (Pandya & Kuypers, 1969), as both regions perform complementary roles in controlling actions (Curtis, 2006; Cisek, 2007; Westendorff et al., 2010; Pilacinski and Lindner, 2019). This tight fronto-parietal interrelation in processing actions seems to be retained when it comes to operating on memory. Both PM and PPC seem to process different stages of search and retrieval of mnemonic content, with PM controlling memory search and PPC successful retrieval of information (Kapur et al., 1995). Attentional shifts underlying memory search were demonstrated to increase functional connectivity between PM and the parietal cortex, suggesting a tight information exchange in the process of retrieving and manipulating mnemonic information, which relies critically on selecting relevant and suppressing irrelevant information (Heinen et al., 2017). But why would the premotor cortex be involved in thinking? The answer lies in how it is involved in actions. The primary function of the premotor cortex seems to be its ability to represent complex, coordinated multi-joint movements (Graziano and Aflalo, 2007) such as those needed for programming hand trajectories or gestures (Pilacinski and Lindner, 2019; Wong et al., 2019), or handwriting (Kadmon Harpaz et al., 2014). The premotor cortex appears to process both motor and semantic levels (i.e., higher-order goals) of actions (Rizzolatti et al., 1996) and is critically involved in executing relevant and suppressing irrelevant actions from among options represented by the parietal cortex (Cisek, 2007; Lindner et al., 2010; Pilacinski and Lindner, 2019).

Neurophysiological data further suggest that the key function of the premotor cortex in
programming movements is selecting from among competing action plans represented by the
posterior parietal cortex, using weighting input from more frontal areas (Cisek, 2007). The tight
connectivity between PM and PPC further suggests that this inter-areal exchange facilitates the
promotion of relevant and suppression of irrelevant information, e.g., actions. In conceptual
processing, such reciprocal exchange between PM and PPC may be utilized in a similar way, in
which frontal areas select from competing information retrieved from long-term memory by
parietal areas through top-down modulation. This idea seems to be supported by our data,
showing that immediately before each NVS, we observe a substantial increase in all three EEG
bands we measured, indicating a complex computational process required by complex top-down
and bottom-up modulation of cognitive content (Karch et al., 2016; Strube et al., 2021). These
distinctive patterns of activity may reflect the evaluation of conceptual information by PM in a
way similar to that during sensorimotor decision-making (Cisek et al., 2009; Thura and Cisek,
2014).

Limitations

Our study focuses on cortical processing responsible for thinking, considering the role of frontal
and parietal areas in conceptual processing triggering NVS. Human cognition involves numerous
subcortical structures such as the hippocampus or basal ganglia, which are critically involved in
information storage and filtering. Due to EEG limitations, we cannot address the role these
structures   play   in   processing   resulting   in   NVS.
Although posterior parietal cortex and premotor cortex can directly exchange information
through a one-synapse connection (Pandya & Kuypers, 1969), we believe there may be
substantial involvement of subcortical structures in how conceptual information is processed.
Due to EEG's poor spatial resolution, our reasoning about the premotor cortex is mainly based on
assumptions given its role in eye movements and sensorimotor processing rather than direct
functional localization. We acknowledge that conceptual operations involve a long cascade of
processes including emotional evaluation, evidence weighting, and everything that constitutes
individual human cognition. While this research focused on a specific byproduct of conceptual
processing (i.e., NVS) and its accompanying neural components, we recognize the complex and
dynamic interplay within the brain resulting in the human ability to operate on abstract concepts.
In our results we seemingly uncovered an interplay between the type of memory search (open vs.
structured) and the instruction to maintain fixation, which cannot be fully explained based on our
data. Restricting NVS did not substantially impair memory performance in the open questions
task, suggesting they are a byproduct of conceptual processing rather than a function.
Lastly, our control task results were clearly biased towards the visual saccade task and did not
allow us to fully assess the role of working memory processing and maintenance in prefrontal
activity preceding NVS. We attempted to do it using the memory saccade task, which was strongly
underpowered in our case. Despite this, the data we obtained suggest that brain activity

preceding NVS is qualitatively different from that in the memory saccade task and does not merely represent maintenance of mnemonic content. Further research is required to assess this complex processing network, and we believe NVS can be a good behavioral model for that. The interplay between NVS and content reproduction type also requires further investigation.

Conclusions

In recent years, multiple authors suggested that conceptual knowledge builds on sensorimotor circuitry in the brain, operating in low-dimensional spaces organized similarly to those representing the physical space. The available evidence has suggested that organizing, storing and manipulating conceptual knowledge is shared by temporal and parietal systems, with the involvement of other areas, such as the frontal cortex, being less understood. Through the use of non-visual saccades as a behavioural model, our study extends the traditional views on the organization of human conceptual knowledge by showing that the premotor cortex is involved in exploring and manipulating items within conceptual space.

References and Notes

Abdoun, Oussama. "NeuroMap: A Spline-Based Interactive Open-Source Software for Spatiotemporal Mapping of 2D and 3D MEA Data." *Frontiers in Neuroinformatics* 4 (2011): 119. https://doi.org/10.3389/fninf.2010.00119.

Abe, Takashi, Hiroshi Nittono, and Tadao Hori. "Current Sources of the Brain Potentials before Rapid Eye Movements in Human REM Sleep." *International Congress Series* 1270 (August 2004): 241–44. https://doi.org/10.1016/j.ics.2004.04.092.

Agresti, Alan. *An Introduction to Categorical Data Analysis*. Wiley, 2007. https://doi.org/10.1002/0470114754.

Andersen, Richard A., and Christopher A. Buneo. "Intentional Maps in Posterior Parietal Cortex." *Annual Review of Neuroscience* 25, no. 1 (March 2002): 189–220. https://doi.org/10.1146/annurev.neuro.25.112701.142922.

Anderson, Carolyn J., Jay Verkuilen, and Timothy R. Johnson. *Applied Generalized Linear Mixed Models: Continuous and Discrete Data - For the Social and Behavioral Sciences*. Springer, 2012.

Babapoor-Farrokhran, Sahand, Martin Vinck, Thilo Womelsdorf, and Stefan Everling. "Theta and Beta Synchrony Coordinate Frontal Eye Fields and Anterior Cingulate Cortex during Sensorimotor Mapping." *Nature Communications* 8, no. 1 (February 2017): 13967. https://doi.org/10.1038/ncomms13967.

Balaban, Carey D., and Joel M. Weinstein. "The Human Pre-Saccadic Spike Potential: Influences of a Visual Target, Saccade Direction, Electrode Laterality and Instructions to Perform Saccades." *Brain Research* 347, no. 1 (November 1985): 49–57. https://doi.org/10.1016/0006-8993(85)90888-1.

Becker, Wolfgang, Ottomar Hoehne, Katsuhiko Iwase, and Hans H. Kornhuber. "Bereitschaftspotential, Prämotorische Positivierung Und Andere Hirnpotentiale Bei Sakkadischen Augenbewegungen." *Vision Research* 12, no. 3 (March 1972): 421–36. https://doi.org/10.1016/0042-6989(72)90087-9.

Bellmund, Jacob L. S., Peter Gärdenfors, Edvard I. Moser, and Christian F. Doeller. "Navigating Cognition: Spatial Codes for Human Thinking." *Science* 362, no. 6415 (November 2018). https://doi.org/10.1126/science.aat6766.

Berryhill, Marian E., Lisa Phuong, Lauren Picasso, Roberto Cabeza, and Ingrid R. Olson. "Parietal Lobe and Episodic Memory: Bilateral Damage Causes Impaired Free Recall of Autobiographical Memory." *The Journal of Neuroscience* 27, no. 52 (December 2007): 14415–23. https://doi.org/10.1523/JNEUROSCI.4163-07.2007.

Bijvank, J. A. Nij, A. Petzold, L. J. Balk, H. S. Tan, B. M. J. Uitdehaag, M. Theodorou, and L. J. van Rijn. "A Standardized Protocol for Quantification of Saccadic Eye Movements: DEMoNS." *PLOS ONE* 13, no. 7 (July 2018): e0200695. https://doi.org/10.1371/journal.pone.0200695.

Brandmeyer, Tracy, and Arnaud Delorme. "Closed-Loop Frontal Midlineθ Neurofeedback: A Novel Approach for Training Focused-Attention Meditation." *Frontiers in Human Neuroscience* 14 (June 2020): 246. https://doi.org/10.3389/fnhum.2020.00246.

Bruce, C. J., M. E. Goldberg, M. C. Bushnell, and G. B. Stanton. "Primate Frontal Eye Fields. II. Physiological and Anatomical Correlates of Electrically Evoked Eye Movements." *Journal of Neurophysiology* 54, no. 3 (September 1985): 714–34. https://doi.org/10.1152/jn.1985.54.3.714.

Cabeza, Roberto, Elisa Ciaramelli, Ingrid R. Olson, and Morris Moscovitch. "The Parietal Cortex and Episodic Memory: An Attentional Account." *Nature Reviews Neuroscience* 9, no. 8 (August 2008): 613–25. https://doi.org/10.1038/nrn2459.

Casals, Martí, Montserrat Girabent-Farrés, and Josep L. Carrasco. "Methodological Quality and Reporting of Generalized Linear Mixed Models in Clinical Medicine (2000–2012): A

Systematic Review." *PLoS ONE* 9, no. 11 (November 2014): e112653.
https://doi.org/10.1371/journal.pone.0112653.

Cisek, Paul. "Cortical Mechanisms of Action Selection: The Affordance Competition Hypothesis." *Philosophical Transactions of the Royal Society B: Biological Sciences* 362, no. 1485 (September 2007): 1585–99. https://doi.org/10.1098/rstb.2007.2054.

Cisek, Paul, Geneviève Aude Puskas, and Stephany El-Murr. "Decisions in Changing Conditions: The Urgency-Gating Model." *The Journal of Neuroscience* 29, no. 37 (September 2009): 11560–71. https://doi.org/10.1523/JNEUROSCI.1844-09.2009.

Constantinescu, Alexandra O., Jill X. O'Reilly, and Timothy E. J. Behrens. "Organizing Conceptual Knowledge in Humans with a Gridlike Code." *Science* 352, no. 6292 (June 2016): 1464–68. https://doi.org/10.1126/science.aaf0941.

Cordeiro, Gauss M., and Alexandre B. Simas. "The Distribution of Pearson Residuals in Generalized Linear Models." *Computational Statistics & Data Analysis* 53, no. 9 (July 2009): 3397–3411. https://doi.org/10.1016/j.csda.2009.02.025.

Cothi, William de, and Hugo J. Spiers. "Spatial Cognition: Goal-Vector Cells in the Bat Hippocampus." *Current Biology* 27, no. 6 (March 2017): R239–41. https://doi.org/10.1016/j.cub.2017.01.061.

Curtis, C. E. "Prefrontal and Parietal Contributions to Spatial Working Memory." *Neuroscience* 139, no. 1 (April 2006): 173–80. https://doi.org/10.1016/j.neuroscience.2005.04.070.

Curtis, Clayton E., Vikas Y. Rao, and Mark D'Esposito. "Maintenance of Spatial and Motor Codes during Oculomotor Delayed Response Tasks." *The Journal of Neuroscience* 24, no. 16 (April 2004): 3944–52. https://doi.org/10.1523/JNEUROSCI.5640-03.2004.

Darby, David G., Anna C. Nobre, Venkatesan Thangaraj, Robert Edelman, M.-Marsel Mesulam, and Steven Warach. "Cortical Activation in the Human Brain during Lateral Saccades Using EPISTAR Functional Magnetic Resonance Imaging." *NeuroImage* 3, no. 1 (February 1996): 53–62. https://doi.org/10.1006/nimg.1996.0006.

Delorme, Arnaud. "EEG Is Better Left Alone." *Scientific Reports* 13, no. 1 (February 2023): 2372. https://doi.org/10.1038/s41598-023-27528-0.

Delorme, Arnaud, and Scott Makeig. "EEGLAB: An Open Source Toolbox for Analysis of Single-Trial EEG Dynamics Including Independent Component Analysis." *Journal of Neuroscience Methods* 134, no. 1 (March 2004): 9–21. https://doi.org/10.1016/j.jneumeth.2003.10.009.

Desikan, Rahul S., Florent Ségonne, Bruce Fischl, Brian T. Quinn, Bradford C. Dickerson, Deborah Blacker, Randy L. Buckner, et al. "An Automated Labeling System for

Subdividing the Human Cerebral Cortex on MRI Scans into Gyral Based Regions of Interest." *NeuroImage* 31, no. 3 (July 2006): 968–80. https://doi.org/10.1016/j.neuroimage.2006.01.021.

Doherty-Sneddon, G., and F. G. Phelps. "Gaze Aversion: A Response to Cognitive or Social Difficulty?" *Memory & Cognition* 33, no. 4 (June 2005): 727–33. https://doi.org/10.3758/BF03195338.

Dunn, Peter K., and Gordon K. Smyth. *Generalized Linear Models With Examples in R*. Springer New York, 2018. https://doi.org/10.1007/978-1-4419-0118-7.

Ehrlichman, Howard, and John Barrett. "'Random' Saccadic Eye Movements during Verbal-Linguistic and Visual-Imaginal Tasks." *Acta Psychologica* 53, no. 1 (April 1983): 9–26. https://doi.org/10.1016/0001-6918(83)90013-6.

Ehrlichman, Howard, and Dragana Micic. "Why Do People Move Their Eyes When They Think?" *Current Directions in Psychological Science* 21, no. 2 (April 2012): 96–100. https://doi.org/10.1177/0963721412436810.

Ehrlichman, Howard, Dragana Micic, Amber Sousa, and John Zhu. "Looking for Answers: Eye Movements in Non-Visual Cognitive Tasks." *Brain and Cognition* 64, no. 1 (June 2007): 7–20. https://doi.org/10.1016/j.bandc.2006.10.001.

Everling, S., P. Krappmann, and H. Flohr. "Cortical Potentials Preceding Pro- and Antisaccades in Man." *Electroencephalography and Clinical Neurophysiology* 102, no. 4 (April 1997): 356–62. https://doi.org/10.1016/S0013-4694(96)96569-4.

Freeden, Willi. "Spherical Spline Interpolation—Basic Theory and Computational Aspects." *Journal of Computational and Applied Mathematics* 11, no. 3 (December 1984): 367–75. https://doi.org/10.1016/0377-0427(84)90011-6.

Gramfort, Alexandre. "MEG and EEG Data Analysis with MNE-Python." *Frontiers in Neuroscience* 7 (2013): 267. https://doi.org/10.3389/fnins.2013.00267.

Graziano, Michael S. A., and Tyson N. Aflalo. "Rethinking Cortical Organization." *The Neuroscientist* 13, no. 2 (April 2007): 138–47. https://doi.org/10.1177/1073858406295918.

Gurtner, Lilla M., Matthias Hartmann, and Fred W. Mast. "Eye Movements during Visual Imagery and Perception Show Spatial Correspondence but Have Unique Temporal Signatures." *Cognition* 210 (May 2021): 104597. https://doi.org/10.1016/j.cognition.2021.104597.

Hafting, Torkel, Marianne Fyhn, Sturla Molden, May-Britt Moser, and Edvard I. Moser. "Microstructure of a Spatial Map in the Entorhinal Cortex." *Nature* 436, no. 7052 (August 2005): 801–6. https://doi.org/10.1038/nature03721.

Harpaz, Naama Kadmon, Tamar Flash, and Ilan Dinstein. "Scale-Invariant Movement Encoding in the Human Motor System." *Neuron* 81, no. 2 (January 2014): 452–62. https://doi.org/10.1016/j.neuron.2013.10.058.

Heinen, Klaartje, Eva Feredoes, Christian C. Ruff, and Jon Driver. "Functional Connectivity between Prefrontal and Parietal Cortex Drives Visuo-Spatial Attention Shifts." *Neuropsychologia* 99 (May 2017): 81–91. https://doi.org/10.1016/j.neuropsychologia.2017.02.024.

Hollander, John, and Stephanie Huette. "Extracting Blinks from Continuous Eye-Tracking Data in a Mind Wandering Paradigm." *Consciousness and Cognition* 100 (April 2022): 103303. https://doi.org/10.1016/j.concog.2022.103303.

Imaoka, Yu, Andri Flury, and Eling D. de Bruin. "Assessing Saccadic Eye Movements With Head-Mounted Display Virtual Reality Technology." *Frontiers in Psychiatry* 11 (September 2020): 572938. https://doi.org/10.3389/fpsyt.2020.572938.

Inc, The MathWorks. "MATLAB (v9.14.0 (R2023a))," 2023.

Johansson, Roger, Marcus Nyström, Richard Dewhurst, and Mikael Johansson. "Eye-Movement Replay Supports Episodic Remembering." Research Square Platform, February 2021. https://doi.org/10.21203/rs.3.rs-275969/v1.

Kang, Seung Suk, Timothy J. Lano, and Scott R. Sponheim. "Distortions in EEG Interregional Phase Synchrony by Spherical Spline Interpolation: Causes and Remedies." *Neuropsychiatric Electrophysiology* 1, no. 1 (December 2015): 9. https://doi.org/10.1186/s40810-015-0009-5.

Kapur, S., Craik, F. I., Jones, C., Brown, G. M., Houle, S., & Tulving, E. (1995). Functional role of the prefrontal cortex in retrieval of memories: a PET study. *Neuroreport*, *6*(14), 1880–1884. https://doi.org/10.1097/00001756-199510020-00014

Karch, Susanne, Fabian Loy, Daniela Krause, Sandra Schwarz, Jan Kiesewetter, Felix Segmiller, Agnieszka I. Chrobok, Daniel Keeser, and Oliver Pogarell. "Increased Event-Related Potentials and Alpha-, Beta-, and Gamma-Activity Associated with Intentional Actions." *Frontiers in Psychology* 7 (January 2016). https://doi.org/10.3389/fpsyg.2016.00007.

Koval, Michael J., R. Matthew Hutchison, Stephen G. Lomber, and Stefan Everling. "Effects of Unilateral Deactivations of Dorsolateral Prefrontal Cortex and Anterior Cingulate Cortex on Saccadic Eye Movements." *Journal of Neurophysiology* 111, no. 4 (February 2014): 787–803. https://doi.org/10.1152/jn.00626.2013.

Kovalenko, Lyudmyla Y., and Niko A. Busch. "Probing the Dynamics of Perisaccadic Vision with EEG." *Neuropsychologia* 85 (May 2016): 337–48.

https://doi.org/10.1016/j.neuropsychologia.2015.12.012.

Larson, Eric, Alexandre Gramfort, Denis A. Engemann, Jaakko Leppakangas, Christian
Brodbeck, Mainak Jas, Teon L. Brooks, Jona Sassenhagen, Daniel McCloy, and Martin
Luessi. "MNE-Python (v1.7.1)." Zenodo, 2024.

Larsson, Linnea, Marcus Nystrom, and Martin Stridh. "Detection of Saccades and
Postsaccadic Oscillations in the Presence of Smooth Pursuit." *IEEE Transactions on
Biomedical Engineering* 60, no. 9 (September 2013): 2484–93.
https://doi.org/10.1109/TBME.2013.2258918.

Lindner, Axel, Asha Iyer, Igor Kagan, and Richard A. Andersen. "Human Posterior Parietal
Cortex Plans Where to Reach and What to Avoid." *The Journal of Neuroscience* 30, no.
35 (September 2010): 11715–25. https://doi.org/10.1523/JNEUROSCI.2849-09.2010.

Liu, Bo, Qi-Chao Zhao, Yuan-Yuan Ren, Qing-Ju Wang, and Xue-Lian Zheng. "An Elaborate
Algorithm for Automatic Processing of Eye Movement Data and Identifying Fixations in
Eye-Tracking Experiments." *Advances in Mechanical Engineering* 10, no. 5 (May 2018):
1–15. https://doi.org/10.1177/1687814018773678.

Mack, David J., Sandro Belfanti, and Urs Schwarz. "The Effect of Sampling Rate and Lowpass
Filters on Saccades – A Modeling Approach." *Behavior Research Methods* 49, no. 6
(December 2017): 2146–62. https://doi.org/10.3758/s13428-016-0848-4.

Martarelli, Corinna S., Fred W. Mast, and Matthias Hartmann. "Time in the Eye of the
Beholder: Gaze Position Reveals Spatial-Temporal Associations during Encoding and
Memory Retrieval of Future and Past." *Memory & Cognition* 45, no. 1 (January 2017):
40–48. https://doi.org/10.3758/s13421-016-0639-2.

McDowell, Jennifer E., Kara A. Dyckman, Benjamin P. Austin, and Brett A. Clementz.
"Neurophysiology and Neuroanatomy of Reflexive and Volitional Saccades: Evidence
from Studies of Humans." *Brain and Cognition* 68, no. 3 (December 2008): 255–70.
https://doi.org/10.1016/j.bandc.2008.08.016.

McPeek, Robert M., Alexander A. Skavenski, and Ken Nakayama. "Concurrent Processing of
Saccades in Visual Search." *Vision Research* 40, no. 18 (August 2000): 2499–2516.
https://doi.org/10.1016/S0042-6989(00)00102-4.

Meghanathan, Radha Nila, Cees van Leeuwen, Marcello Giannini, and Andrey R. Nikolaev.
"Neural Correlates of Task-Related Refixation Behavior." *Vision Research* 175 (October
2020): 90–101. https://doi.org/10.1016/j.visres.2020.07.001.

Meister, Miriam L. R., and Elizabeth A. Buffalo. "Neurons in Primate Entorhinal Cortex
Represent Gaze Position in Multiple Spatial Reference Frames." *The Journal of
Neuroscience* 38, no. 10 (March 2018): 2430–41.

https://doi.org/10.1523/JNEUROSCI.2432-17.2018.

Micic, Dragana, Howard Ehrlichman, and Rebecca Chen. "Why Do We Move Our Eyes While Trying to Remember? The Relationship between Non-Visual Gaze Patterns and Memory." *Brain and Cognition* 74, no. 3 (December 2010): 210–24. https://doi.org/10.1016/j.bandc.2010.07.014.

Munoz, Douglas P. "Commentary: Saccadic Eye Movements: Overview of Neural Circuitry." In *Progress in Brain Research*, 89–96. Elsevier, 2002. https://doi.org/10.1016/S0079-6123(02)40044-1.

Nikolaev, Andrey R., Peter Jurica, Chie Nakatani, Gijs Plomp, and Cees van Leeuwen. "Visual Encoding and Fixation Target Selection in Free Viewing: Presaccadic Brain Potentials." *Frontiers in Systems Neuroscience* 7 (2013): 26. https://doi.org/10.3389/fnsys.2013.00026.

Nouira, Ibtihel, Asma Ben Abdallah, and Mohamed Hedi Bedoui. "Three-Dimensional Interpolation Methods to Spatiotemporal EEG Mapping during Various Behavioral States." *Signal, Image and Video Processing* 10, no. 5 (July 2016): 943–49. https://doi.org/10.1007/s11760-015-0844-7.

Nyström, Marcus, and Kenneth Holmqvist. "An Adaptive Algorithm for Fixation, Saccade, and Glissade Detection in Eyetracking Data." *Behavior Research Methods* 42, no. 1 (February 2010): 188–204. https://doi.org/10.3758/BRM.42.1.188.

O'Keefe, J., and J. Dostrovsky. "The Hippocampus as a Spatial Map. Preliminary Evidence from Unit Activity in the Freely-Moving Rat." *Brain Research* 34, no. 1 (November 1971): 171–75. https://doi.org/10.1016/0006-8993(71)90358-1.

Pandya, Deepak N., and Henricus G. J. M. Kuypers. "Cortico-Cortical Connections in the Rhesus Monkey." *Brain Research* 13, no. 1 (March 1969): 13–36. https://doi.org/10.1016/0006-8993(69)90141-3.

Paré, Martin, and Robert H. Wurtz. "Monkey Posterior Parietal Cortex Neurons Antidromically Activated From Superior Colliculus." *Journal of Neurophysiology* 78, no. 6 (December 1997): 3493–97. https://doi.org/10.1152/jn.1997.78.6.3493.

Pascual-Marqui, R. D. "Standardized Low-Resolution Brain Electromagnetic Tomography (sLORETA): Technical Details." *Methods and Findings in Experimental and Clinical Pharmacology* 24 Suppl D (2002): 5–12.

Peelen, Marius V., Dirk J. Heslenfeld, and Jan Theeuwes. "Endogenous and Exogenous Attention Shifts Are Mediated by the Same Large-Scale Neural Network." *NeuroImage* 22, no. 2 (June 2004): 822–30. https://doi.org/10.1016/j.neuroimage.2004.01.044.

Perrin, F., J. Pernier, O. Bertrand, and J. F. Echallier. "Spherical Splines for Scalp Potential

787 and Current Density Mapping." *Electroencephalography and Clinical Neurophysiology*
788 72, no. 2 (February 1989): 184–87. https://doi.org/10.1016/0013-4694(89)90180-6.

789 Peterson, Matthew S., Arthur F. Kramer, and David E. Irwin. "Covert Shifts of Attention
790 Precede Involuntary Eye Movements." *Perception & Psychophysics* 66, no. 3 (April
791 2004): 398–405. https://doi.org/10.3758/BF03194888.

792 Pierrot-Deseilligny, Charles, Sophie Rivaud, Bertrand Gaymard, René Müri, and Anne-
793 Isabelle Vermersch. "Cortical Control of Saccades." *Annals of Neurology* 37, no. 5 (May
794 1995): 557–67. https://doi.org/10.1002/ana.410370504.

795 Pilacinski, Artur, and Axel Lindner. "Distinct Contributions of Human Posterior Parietal and
796 Dorsal Premotor Cortex to Reach Trajectory Planning." *Scientific Reports* 9, no. 1
797 (February 2019): 1962. https://doi.org/10.1038/s41598-019-39188-0.

798 Pylyshyn, Zenon. "Return of the Mental Image: Are There Really Pictures in the Brain?"
799 *Trends in Cognitive Sciences* 7, no. 3 (March 2003): 113–18.
800 https://doi.org/10.1016/S1364-6613(03)00003-2.

801 Rayner, Keith, Maria L. Slowiaczek, Charles Clifton, and James H. Bertera. "Latency of
802 Sequential Eye Movements: Implications for Reading." *Journal of Experimental*
803 *Psychology: Human Perception and Performance* 9, no. 6 (1983): 912–22.
804 https://doi.org/10.1037/0096-1523.9.6.912.

805 Revankar, Gajanan S., Noriaki Hattori, Yuta Kajiyama, Tomohito Nakano, Masahito Mihara,
806 Etsuro Mori, and Hideki Mochizuki. "Ocular Fixations and Presaccadic Potentials to
807 Explain Pareidolias in Parkinson's Disease." *Brain Communications* 2, no. 1 (January
808 2020): 1–15. https://doi.org/10.1093/braincomms/fcaa073.

809 Richards, John E. "Cortical Sources of ERP in Prosaccade and Antisaccade Eye Movements
810 Using Realistic Source Models." *Frontiers in Systems Neuroscience* 7 (2013): 27.
811 https://doi.org/10.3389/fnsys.2013.00027.

812 Richards, John E. "Cortical Sources of Event-related Potentials in the Prosaccade and
813 Antisaccade Task." *Psychophysiology* 40, no. 6 (November 2003): 878–94.
814 https://doi.org/10.1111/1469-8986.00106.

815 Rizzolatti, Giacomo, Luciano Fadiga, Vittorio Gallese, and Leonardo Fogassi. "Premotor
816 Cortex and the Recognition of Motor Actions." *Cognitive Brain Research* 3, no. 2
817 (March 1996): 131–41. https://doi.org/10.1016/0926-6410(95)00038-0.

818 Rizzolatti, Giacomo, Lucia Riggio, Isabella Dascola, and Carlo Umiltá. "Reorienting Attention
819 across the Horizontal and Vertical Meridians: Evidence in Favor of a Premotor Theory
820 of Attention." *Neuropsychologia* 25, no. 1 (January 1987): 31–40.
821 https://doi.org/10.1016/0028-3932(87)90041-8.

Ruíz, Josafhat Salinas, Osval Antonio Montesinos López, Gabriela Hernández Ramírez, and
Jose Crossa Hiriart. "Elements of Generalized Linear Mixed Models." In *Generalized*
Linear Mixed Models with Applications in Agriculture and Biology, 22–35. Springer
International Publishing, 2023. https://doi.org/10.1007/978-3-031-32800-8.

Schielzeth, Holger, Niels J. Dingemanse, Shinichi Nakagawa, David F. Westneat, Hassen
Allegue, Céline Teplitsky, Denis Réale, Ned A. Dochtermann, László Zsolt Garamszegi,
and Yimen G. Araya-Ajoy. "Robustness of Linear Mixed-effects Models to Violations of
Distributional Assumptions." *Methods in Ecology and Evolution* 11, no. 9 (September
2020): 1141–52. https://doi.org/10.1111/2041-210X.13434.

Schluppeck, Denis, Paul Glimcher, and David J. Heeger. "Topographic Organization for
Delayed Saccades in Human Posterior Parietal Cortex." *Journal of Neurophysiology* 94,
no. 2 (August 2005): 1372–84. https://doi.org/10.1152/jn.01290.2004.

Scholz, Agnes, Anja Klichowicz, and Josef F. Krems. "Covert Shifts of Attention Can Account
for the Functional Role of 'Eye Movements to Nothing.'" *Memory & Cognition* 46, no. 2
(February 2018): 230–43. https://doi.org/10.3758/s13421-017-0760-x.

Servais, Anaïs, Christophe Hurter, and Emmanuel J. Barbeau. "Attentional Switch from the
Outside to the Inner World: An Early and Critical, but Unexplored Stage of
Autobiographical Memory Retrieval Processes." *Brain and Cognition* 170 (August
2023): 106031. https://doi.org/10.1016/j.bandc.2023.106031.

Sipatchin, Alexandra, Siegfried Wahl, and Katharina Rifai. "Eye-Tracking for Clinical
Ophthalmology with Virtual Reality (VR): A Case Study of the HTC Vive Pro Eye's
Usability." *Healthcare* 9, no. 2 (February 2021): 180.
https://doi.org/10.3390/healthcare9020180.

Smith, Daniel T., and Thomas Schenk. "The Premotor Theory of Attention: Time to Move
On?" *Neuropsychologia* 50, no. 6 (May 2012): 1104–14.
https://doi.org/10.1016/j.neuropsychologia.2012.01.025.

Strube, Andreas, Michael Rose, Sepideh Fazeli, and Christian Büchel. "Alpha-to-Beta- and
Gamma-Band Activity Reflect Predictive Coding in Affective Visual Processing."
Scientific Reports 11, no. 1 (December 2021): 23492. https://doi.org/10.1038/s41598-
021-02939-z.

Svantesson, Mats, Håkan Olausson, Anders Eklund, and Magnus Thordstein. "Virtual EEG-
Electrodes: Convolutional Neural Networks as a Method for Upsampling or Restoring
Channels." *Journal of Neuroscience Methods* 355 (May 2021): 109126.
https://doi.org/10.1016/j.jneumeth.2021.109126.

Team, JASP. "JASP (v0.18.3)," 2024.

Thura, David, and Paul Cisek. "Deliberation and Commitment in the Premotor and Primary Motor Cortex during Dynamic Decision Making." *Neuron* 81, no. 6 (March 2014): 1401–16. https://doi.org/10.1016/j.neuron.2014.01.031.

Tolman, Edward C. "Cognitive Maps in Rats and Men." *Psychological Review* 55, no. 4 (1948): 189–208. https://doi.org/10.1037/h0061626.

Wechsler, David. *Wechsler Adult Intelligence Scale*. 3rd ed. The Psychological Corporation, 1997.

Weiner, Susan L., and Howard Ehrlichman. "Ocular Motility and Cognitive Process." *Cognition* 4, no. 1 (January 1976): 31–43. https://doi.org/10.1016/0010-0277(76)90009-3.

Westendorff, Stephanie, Christian Klaes, and Alexander Gail. "The Cortical Timeline for Deciding on Reach Motor Goals." *The Journal of Neuroscience* 30, no. 15 (April 2010): 5426–36. https://doi.org/10.1523/JNEUROSCI.4628-09.2010.

Whitlock, Jonathan R. "Posterior Parietal Cortex." *Current Biology* 27, no. 14 (July 2017): R691–95. https://doi.org/10.1016/j.cub.2017.06.007.

Whitlock, Jonathan R., Robert J. Sutherland, Menno P. Witter, May-Britt Moser, and Edvard I. Moser. "Navigating from Hippocampus to Parietal Cortex." *Proceedings of the National Academy of Sciences* 105, no. 39 (September 2008): 14755–62. https://doi.org/10.1073/pnas.0804216105.

Xia, Yinglin, and Jun Sun. "Linear Mixed-Effects Models for Longitudinal Microbiome Data." In *Bioinformatic and Statistical Analysis of Microbiome Data*. Springer International Publishing, 2023. https://doi.org/10.1007/978-3-031-21391-5.

Yeung, Lok-Kin, Rosanna K. Olsen, Hannah E. P. Bild-Enkin, Maria C. D'Angelo, Arber Kacollja, Douglas A. McQuiggan, Anna Keshabyan, Jennifer D. Ryan, and Morgan D. Barense. "Anterolateral Entorhinal Cortex Volume Predicted by Altered Intra-Item Configural Processing." *The Journal of Neuroscience* 37, no. 22 (May 2017): 5527–38. https://doi.org/10.1523/JNEUROSCI.3664-16.2017.

Acknowledgments: The authors would like to acknowledge support from Bial Foundation (grant 260/22) and Portuguese Foundation for Science and Technology (2023.08332.CEECIND) to AP, and Deutsche Forschungsgemeinschaft grant (122679504—SFB 874) to CK. We want to thank Alexandra Sipatchin and Fernanda Ponce for their awesome help with eyetracking data analysis, Diogo Branco for his help with the task and the setup, Soraia Oliveira for help with data collection and Susanne Dyck, Jon Walbrin, Fredrik Bergstroem, Elizabetta Del Re and Pieter Medendorp for helpful input and discussions.