Title: Spontaneous saccades reveal that human premotor cortex is involved in conceptual processing

Authors: Artur Pilacinski[1,2,*], Lia Wagner[1,$], Gabriel Besson[2], Maria Joao Matos[2], Eduardo Araujo[2], Christian Klaes[1]

[$] Equal contribution

Affiliations: [1] Medical Faculty, Ruhr University Bochum, Bochum, Germany; [2] Faculty of Psychology and Educational Sciences, University of Coimbra, Coimbra, Portugal

*Corresponding author: AP, art.pilacinski@gmail.com

Abstract:

Non-visual saccades (NVS) are spontaneous eye movements humans make while thinking. Their role is puzzling and there is no framework to explain them in terms of underlying neural systems. Here we studied neural activity preceding NVS when subjects performed standardized abstract thinking tasks. We used high-density EEG and focused on frontal and parietal channels representing oculomotor cortical areas. We found that NVS are distinctively preceded by neural activity changes in frontal and parietal EEG channels qualitatively different to that for voluntary saccades. However, only the ERP amplitude changes in frontal channels, but not in parietal channels, were different from saccadic preparation. A source reconstruction analysis additionally revealed that presaccadic neural activity originated mainly in the premotor cortex. Frequency spectrum analyses showed increased synchronization in alpha to low gamma bands suggesting complex top-down and bottom-up processing. We propose that this presaccadic activity while thinking represents evidence accumulation and attentional shifts, as shown before in the premotor cortex during sensorimotor tasks. Our findings suggest that human thinking appears to function similarly to human actions and engages premotor areas responsible for evaluating and manipulating objects also when evaluating and manipulating concepts.

Main Text:

Introduction

What do apples and oranges have in common? What is similar between crime and punishment? Why do we pay taxes? Take a moment to answer these questions and you might notice that, before answering, you probably made a few eye movements as if in search of the right answer. It is a known but often ignored fact that such spontaneous, non-visual saccadic eye movements (NVS) accompany human thinking, that is mental task solving and mnemonic retrieval processes (Ehrlichman & Micic, 2012). Why they occur and what function they serve remains unknown. Here, we show that NVS are immediately preceded by specific activity changes in frontal regions including premotor cortex, indicating the crucial role of that area in conceptual operations (thinking).

Non-visual saccades remain a very understudied phenomenon and the existing research does not allow for building a coherent picture of their nature. Contrary to some opinions (Doherty-Sneddon and Phelps, 2005) NVS do not subserve others' gaze aversion as they are also performed in darkness or without other people present (Ehrlichman & Barrett, 1983). Restriction of NVS does not seem to impact memory retrieval performance (Micic et al., 2010) suggesting they may be an epiphenomenon rather than a functional process (for an alternative view see: Scholz et al., 2016). They are, however, not fully suppressible and people move their eyes while thinking even when instructed to fixate gaze (Micic et al., 2010) indicating that NVS are automatically accompanying memory operations. On the other hand, non-visual saccades have been demonstrated to reflect mnemonic content organization such as spatial arrangement of items (Johansson et al., 2022; Damiano and Walther, 2019; Scholz et al., 2016) or the point in time (Hartmann et al., 2014; Martarelli et al., 2016). Although one could suspect these NVS to merely reflect visuospatial organization of memorized items, it comes as a surprise that substantially more saccades are produced when operating on verbal than on visual material (Weiner and Ehrlichmann, 1976). This indicates that NVS do not directly reflect reactivating the visual scene in memory but rather other processes such as activation of conceptual knowledge necessary for processing semantics.

Organization of conceptual knowledge has been long described as the core function of medial temporal lobe (MTL) (Bellmund et al., 2018; Constantinescu et al., 2016). Meanwhile, the main cortical centers controlling saccades span across posterior parietal (PPC) and premotor cortex (PM), comprising frontal eye fields (FEF), the main area of the brain responsible for programming voluntary eye movements (Munoz, 2002). While some known interactions exist between the MTL and eye movements systems, the mechanism that could possibly explain NVS remains unclear. For example, activity in MTL reflects integrating spatial information when exploring visual scenes (Leszczynski et al., 2024), about objects (Yeung et al., 2017), hippocampal maps are modulated by gaze direction (Meister et al., 2018) and parietal activity reflects that of the hippocampus (Whitlock et al., 2008). It is unclear, however, how retrieving mnemonic content could possibly initiate saccades. One recently suggested possibility is that hippocampus allows binding elements of the visual scene through receiving the corollary discharge accompanying visual exploration, thereby showing tight links between oculomotor system and memory encoding (Katz et al., 2022; Leszczynski et al., 2024). It is conceivable that such corollary discharge could then be reinstated too during memory retrieval, for example to enable shifting the focus of attention between elements of the conceptual space. While hypothetical, this perspective highlights the existing physiological links between medial temporal lobe and oculomotor regions in the primate brain that may be important for understanding NVS.

The core cortical regions for programming voluntary saccades are posterior parietal and prefrontal cortical areas (Pierrot-Deseilligny et al., 1995; Andersen & Buneo, 2002). Pre-saccadic planning activity was described earlier as predominantly visible in parietal EEG channels (Everling et al., 1997; Richards, 2003; Becker et al., 1972; Balaban & Weinstein, 1985). Posterior parietal cortex (PPC) has been proposed to integrate between memory and actions (Whitlock et al., 2008; Whitlock, 2017), and parietal oculomotor areas (Schluppeck et al., 2005) overlap with those involved in long term memory search (Berryhill et al., 2007; Vilberg and Rugg, 2008). PPC is also directly connected to superior colliculus (Pare and Wurtz, 1997), which suggests parietal activity may be triggering NVS as a byproduct of memory search processes. Whether it is the memory retrieval activity in parietal oculomotor areas that underlies NVS is unknown.

Posterior parietal cortex typically projects oculomotor programming signals to the frontal eye

field, the main saccade-controlling area in the primate brain (Bruce et al., 1985). In humans, FEF

is contained within the dorsal premotor cortex (Darby et al., 1996), indicating shared function of

the two areas. Premotor cortex (PM) has been repeatedly demonstrated to be involved in a

number of non-motor tasks, including working memory (Curtis et al., 2004; Curtis, 2006; Lindner

2010; Pilacinski et al., 2020), evaluating evidence during decision-making (Thura et al., 2012) and

internal shifts of attention (Peelen et al., 2004). The premotor theory of attention further

suggests that these internal shifts of attention reutilize saccade programming circuitry (Rizzolatti

et al., 1996). Premotor cortex has also been shown to be involved in memory search (Kapur et

al., 1995).

The organization of conceptual knowledge has long been suggested to reutilize sensorimotor

computations, e.g. through representing cognitive spaces in a similar way as physical space

(Tolman, 1948; Bellmund et al., 2018). In our view, non-visual saccades are a window into the

evolution of human thinking by providing a link between the memory and the action systems in

the human brain. The available evidence suggests that the frontal and parietal cortex may be

involved in generation of NVS during memory search, although the particular involvement of

each of these areas has not been explored. It is likewise unknown how neural mechanisms

underlying NVS compare to typical programming of goal-directed saccades. Lastly, it is unclear

whether NVS are a function or merely a byproduct of human thinking.

Here, we tackled these questions using high-density EEG, by investigating the neural activity

immediately preceding non-visual saccades, while volunteers were solving abstract reasoning

and knowledge tasks, with restricted and unrestricted gaze and with open and structured

memory search conditions. In addition, we compared the presaccadic activity for NVS with

voluntary saccades thereby controlling for unspecific preparatory signals such as readiness

potential. Our central question was whether NVS are preceded by specific neural signals. If that

was the case, we further reasoned that if NVS are a product of memory search mediated by

posterior parietal cortex (as suggested by prior studies), then we should observe a neural activity

change immediately preceding NVS in parietal channels. Conversely, if NVS were a product of

attentional processing of conceptual information, they should be preceded by neural activity

changes in frontal channels. Finally, if non-visual saccades arise from neural processing distinct

to saccadic preparation, the neural activity changes preceding them should not be present before

voluntary saccades.

Methods

Subjects

A total of 57 individuals (mean age 27.2 ± 8.1 years, 32 male, 5 left-handed) took part in the study.

All participants had normal or corrected-to-normal vision with no history of neurological or

psychiatric disorders. Participants were informed about the study procedures, were naive to the

experimental hypothesis and received no financial compensation for their participation. Before

participation, all subjects gave written informed consent in accordance with the Declaration of

Helsinki and local ethical guidelines. The procedures were accepted by the local Ethics and

Deontology Committee for Research of the Faculty of Psychology and Educational Sciences of the

University of Coimbra. All datasets were anonymized before data collection began. Twelve

participants were excluded from all analyses due to missing data or poor data quality as detailed

in the Supplementary Materials (see Table S5), leaving an overall sample size of 45 participants

(mean age 27.2 ± 8.0, 23 male, 4 left-handed) for the behavioral analysis. For the EEG analysis,

further exclusion criteria based on EEG signal quality were applied, as described in the

supplement, resulting in a final sample size of 29 participants (mean age 26.9 ± 7.0 years, 11

male, none left-handed). Due to a technical problem with data logging, there were just 12

subjects considered in analyses of the memory saccade control task (see Supplementary

Materials, Table S5).

Task

In the main experimental task, participants were supposed to answer forty questions taken from

the Portuguese adaptation of Wechsler Adult Intelligence Scale-III (Wechsler, 1997). This task

allowed us to have normalized stimuli, arranged progressively according to their difficulty and

abstractness levels. The questions were presented verbally, through headphones (see Figure 1

A). The questions were split into "structured" and "open" according to the type of memory search

they involved: 20 questions from the Similarities subtest were considered "structured"; 10
questions from the Information subtest, and 10 from the Comprehension subtest were
considered "open". The Information subtest evaluates long-term memory and general
knowledge, e.g., "At what temperature does water boil under normal circumstances?" The
Comprehension subtest assesses practical judgments, e.g., "Why do we pay taxes?" These were
considered "open" questions due to the unrestricted memory search required. The Similarities
subtest assesses abstract verbal reasoning by finding similarities between paired concepts, e.g.,
"How are an orange and a banana comparable?". These were considered "structured" questions.
Further examples can be found in the supplementary materials (see Table S1). Collectively, these
subtests assess verbal conceptualization, vocabulary, abstract reasoning, categorical thinking,
logical abstraction, and social intelligence, reflecting a broad range of cognitive functions and
culturally acquired knowledge. The forty questions were divided into four blocks of 10 trials each,
with randomized order within each block across subjects, and increasing abstraction across
blocks. In each trial, a question was played through headphones and subjects were instructed to
answer as best as they could. Correctness of answers was manually assessed according to the
test instructions and saved in a binary form (correct/incorrect). The only displayed item during
this task was a centrally-positioned bright fixation cross on a dark grey background. Subjects were
randomly assigned to two groups: "fixed gaze" in which they were told to keep looking at the
fixation cross and "free gaze" in which they were not instructed to do so.
A control task was used to assess goal-directed saccadic programming, as well as to differentiate
between visual and memory saccade programming components (see Figure 1 B). The task was
split into the following conditions: visual vs. memory saccades and pro- vs. antisaccades. In the
visual condition, participants were instructed to make saccades either toward (prosaccade) or
away from (antisaccade) a blinking dot located in the periphery. In the memory condition,
participants were asked to make saccades to the location where a target dot had been before it
disappeared. In both conditions, the saccade had to be executed after a delay (when the fixation
cross disappeared).

Figure 1. Experimental tasks timeline. A) Main task. The question was played through

headphones; next, the participant was given time to think of an answer; after they responded,

the next trial started. B) Control tasks: subjects were instructed to saccade to (green cue) or away

from (red cue) the cued location (white dot). The saccade should be executed either towards the

visible (VST) or memorized (MST) target. Blue arrows indicate saccade epoch. Objects not to

scale.

Eye Tracking

Eye movements were recorded at 100 Hz using the HTC VIVE Pro Eye, which has an embedded

eye tracker in a VR head-mounted display and allows for accurate recording of eye data in

immersive environments (Sipatchin et al., 2021). The VR environments were developed to serve

as experimental tasks (see Figure 1 A, B), and an eye tracking algorithm was implemented in Unity

with the use of SRanipal software development kit from HTC. The use of such a head-mounted

display allows for reliable EEG recordings (Tauscher et al., 2019, including EEG systems similar to

the one used here (Dejbara et al., 2019).

For the behavioral oculomotor analysis, saccade and blink frequencies, pupil diameter, saccade

amplitudes, peak velocities, and angles were extracted for each trial and participant, focusing on

the period after each question until the end of each trial (see Supplementary Methods).

Parameters were averaged across combinations of factors for each participant, specifically blocks

1-4, condition type (fixed/free), and question type (open/structured). To assess eye parameters

during the cognitive engagement, GLMM analyses were used as an alternative to the repeated

measures ANOVA, with the fixed factors block (1-4), condition type (fixed/free) and question type

(open/structured). Subjects were considered as the random effects grouping factor. GLMM

analyses were performed separately for the dependent variable answer duration, saccade

frequency, saccade amplitude, blink frequency and pupil average. Only 'correct' trials were

included in the statistical analysis to minimize confounds in an already complex statistical model

and ensure a clearer interpretation of cognitive engagement. The final trial count in the EEG

analyses for the main tasks was 2884 (mean 100 ± 52.26 per participant), and for the control

tasks was 2168 (mean 94 ± 11.95 per participant).

EEG Recordings

EEG data was recorded using either a 64-electrode dry or a 256-electrode wet EEG system (eego

mylab, ANT Neuro, Enschede, Netherlands), and event triggers were sent via the Lab Streaming

Layer (LSL) protocol between the computers of the VR system and the EEG software (eego

software, ANT Neuro), ensuring precise synchronization of the stimulus and eye tracking events

within the EEG data. Initially, the 256-channel system was used for the first eleven participants,

but due to technical issues, the 64-electrode system was used for the remaining eighteen

participants. There were no statistically significant differences between both systems in terms of

signal quality that would influence our results (see Supplementary Methods). Electrodes were

arranged in an equidistant montage with the left and right mastoids as online ground and

reference for the 64-channel setup, and CPz and the left mastoid for the 256-channel setup.

Impedances were kept below 25 kΩ, and signals were recorded at 500 Hz. Preprocessing was

done with EEGLAB and MNE-Python, including checks for sampling rate stability, application of a

1 Hz high-pass filter and 40 Hz low-pass filter, and removal of 25 Hz environmental noise using a

notch filter. A 50 Hz notch filter removed remaining line noise. Flatline channels were removed

using EEGLAB's pop_clean_rawdata function, and noisy channels were rejected using

pop_rejchan based on kurtosis, probability, and spectra, followed by manual inspection.

Interpolated channels were re-referenced to the average of all electrodes. Eye movements were

controlled using eye tracking data. EEG events were processed separately for the main and

control tasks, and epochs were extracted from -200 to 100 ms relative to saccade onset. Saccades

preceded by other saccades or blinks within 200 ms were excluded. A baseline correction from -

200 to -170 ms was performed using single-trial normalization. We decided for this time window

based on typical inter-saccadic latencies (Rayner et al., 1983). This allowed us to scrutinize signal

changes immediately preceding NVS. Epochs were rejected if spikes were detected, and

semiautomatic artifact rejection occurred for amplitudes exceeding 50 µV between samples or

100 µV over 50 ms. As the epochs were thus free of eye movements, ICA was not used to clean

further the signal to avoid the risk of eliminating relevant data (e.g. saccade planning),

particularly given recent evidence that automated ICA rejection, such as of eye movements

(clearly present in their data), does not reliably increase the actual contrasts studied (Delorme,

2023). Four regions of interest (ROIs), frontal left (FL), frontal right (FR), parietal left (PL), and

parietal right (PR), were defined, distributed around corresponding electrodes F3, F4, P3 and P4

in the 10-20 electrode system (see Figure 3 A and Figure S1 A, B). Since the FEF and PPC are the

primary areas of interest in saccade research and are both superficial and spatially distinct, this

setup allowed us to effectively distinguish neural activity in these regions.

ERP analysis

For the EEG analysis, ERP analysis was performed using MNE-Python. An interval of -150 to -20

ms prior to saccade onset was used for statistical evaluation of presaccadic activity (Revankar et

al., 2020). The mean amplitude of all epochs within this interval was calculated for each condition

and participant. Potentials were then averaged across electrodes within each ROI, allowing for

the assessment of neural dynamics associated with saccade preparation across different ROIs.

Grand average ERPs were calculated for ERP visualization. For topographic visualizations of grand

average ERPs, all electrodes from both EEG systems were included.

Statistical analysis of ERP data was performed either on each ROI or on combined frontal and

parietal ROIs. A Kolmogorov-Smirnov test was used on averaged amplitudes to confirm normality

prior to ANOVA analyses. Averaged EEG amplitudes of conditions were statistically compared

using 4 × 2 or 2 × 2 repeated measures ANOVAs with ROI (4 ROIs and 2 ROIs, respectively) and

condition type as factors. In cases of violation of sphericity (as indicated by Mauchly's test),

Huynh-Feldt corrected p-values were used. For post hoc analysis, paired t-tests were conducted

with Holm correction for multiple testing. To investigate the temporal dynamics of signal

divergence between main and control tasks, the same statistical approach was applied using a

sliding window of 10 ms within the previously defined interval. Mean amplitudes were calculated

within each window for each participant and condition, and paired t-tests with Holm correction

were used to identify the time points at which significant differences emerged.

Frequency spectrum analysis

Spectral decomposition was conducted using EEGLAB to assess neural activity in specific

frequency bands associated with saccade preparation. The analysis focused on three frequency

bands: alpha (8–12 Hz), beta (13–30 Hz), and low-gamma (30–45 Hz). Power spectral density

(PSD) was computed for the interval of -150 to -20 ms prior to saccade onset. For each condition,

PSD was averaged over time and the selected ROIs. Statistical analysis was performed separately

for each frequency band on combined frontal and parietal ROIs. Repeated measures ANOVAs

were conducted with ROI and task type as factors to evaluate differences in PSD values. In cases

of violation of sphericity (as indicated by Mauchly's test), Huynh-Feldt corrected p-values were

used. For post hoc analysis, paired t-tests were conducted with Holm correction for multiple

testing.

Source localization

The forward model and the ERP for each condition were used to estimate an inverse model using

the standardized low-resolution brain electromagnetic tomography (sLORETA) method (Pascual-

Marqui, 2002). We extracted source estimates for each condition and computed spatial

adjacency in the source space to enable cluster-based permutation testing. We performed a non-

parametric, two-tailed cluster-based permutation test, involving voxel-wise comparisons to zero

and correcting for multiple testing. We assessed statistical significance using a p-value threshold

of 0.05. We projected t-values into sLORETA images to visualize source activity, highlighting

regions with significant neural dynamics. We obtained anatomical labels using the FreeSurfer

parcellation scheme to map source activity to specific cortical regions, focusing on cortical

involvement in saccadic movements. Subcortical areas were not captured due to parcellation

limitations. We aligned source estimates with these labels to investigate neural activity in distinct

areas. For each significant cluster identified, we extracted the maximum t-value for each label.

Results

Behavioral and oculomotor results

First, a decrease in percentage of known answers can be observed across blocks for structured

questions while open questions show a significant drop in block 2 with no overall decrease across

blocks (see Figure 2 A). Additionally, interactions between question type and condition type

revealed fewer known answers for the fixed gaze condition for structured questions only (see

Figure 2 B). A significant main effect was found of the factor block ($\chi^2 = 53.151, p < 0.001$),

indicating that performance dropped as abstractness increased. No significant main effect was

found for question type ($\chi^2 = 3.398, p = 0.065$) or condition type ($\chi^2 = 2.702, p = 1.000$),

indicating that the percentage of known answers did not reliably differ between structured and

open questions or between fixed and free gaze conditions. Significant interaction effects were

found for block × question type ($\chi^2 = 112.722, p < 0.001$) and question type × condition type

($\chi^2 = 7.987, p = 0.005$), suggesting that the effect of block varied across different question

types. Post hoc Wilcoxon signed-rank tests revealed a significant difference for block 1 and block

2, block 1 and 4, and block 3 and block 4 ($p < 0.001$). Additionally, a significant difference was

found between block 1 and block 3 ($p = 0.031$). Post hoc Wilcoxon rank-sum tests also showed

a difference between fixed and free gaze condition ($p = 0.029$). For interactions between

question type and condition type (Figure 2 B), post hoc Wilcoxon rank-sum tests confirmed a

significant difference between open compared to structured questions for the fixed gaze

condition only ($p = 0.026$), but not for the free gaze condition ($p = 1.000$).

Saccade frequency for condition type and question type remains approximately the same, but

increases over blocks for both fixed and free gaze conditions (see Figure 2 C). For saccade

frequency, a significant main effect was found of the factor block ($\chi^2 = 11.500, p = 0.009$). Post

hoc Wilcoxon signed-rank tests revealed a significant difference between block 1 and 3 ($p = 0.017$) and block 2 and 3 ($p = 0.043$). However, no significant difference was observed for block

0.017) and block 2 and 3 ($p = 0.043$). However, no significant difference was observed for block

4, including comparisons between block 1 and block 4 ($p = 0.055$), showing that while we see

an increase in saccade frequency in block 3, the frequency is slightly reduced again in block 4.

The post hoc Wilcoxon rank-sum test for gaze condition type showed no significant difference

($p = 0.604$).

Saccadic amplitudes slightly decreased across blocks (see Figure 2 D). For different question

types, saccadic amplitude remains approximately the same, whereas in the fixed condition, mean

saccadic amplitudes are decreased compared to the free gaze condition. For saccadic amplitude,

a significant main effect of the factor block was found ($\chi^2 = 10.563, p = 0.014$). Also, a

significant interaction effect of block × question type × condition type ($\chi^2 = 9.784, p = 0.020$)

suggests that the effect of blocks varied across different question types as well as condition types.

Regarding condition type, the post hoc Wilcoxon rank-sum test revealed a significant difference

between fixed and free gaze conditions ($p = 0.006$).

An increased pupil diameter as an indicator of cognitive load can be identified across blocks and

in the fixed gaze condition compared to the free gaze condition (see Figure 2 E). Additionally, a

slight increase in pupil diameter can be noted for open questions compared to structured

questions. For pupil diameter, a significant main effect of the factor block ($\chi^2 = 24.720, p <$

0.001), question type ($\chi^2 = 16.897, p < 0.001$), and condition type ($\chi^2 = 6.750, p = 0.009$) was

found. Additionally, a significant interaction between block × question type was identified ($\chi^2 =$

$21.756, p < 0.001$). Post hoc Wilcoxon signed-rank tests revealed a significant difference for

block 1 compared to all other blocks ($p < 0.001$). A significant difference was also found

between block 2 and 3 ($p = 0.029$). No significant differences were found between block 2 and

4 ($p = 0.194$) or between block 3 and 4 ($p = 0.311$). These results suggest that pupil diameter

increased across the initial blocks, indicating rising cognitive load or engagement levels, but

slightly decreased again in block 4.

Lastly, for answer duration (see Figure 2 F), a significant main effect was found of the factor block

($\chi^2 = 58.375, p < 0.001$) and question type ($\chi^2 = 6.074, p = 0.014$). There was also a

significant interaction effect of block × question type ($\chi^2 = 24.804, p < 0.001$), suggesting that

the effect of blocks varied across different question types. Post hoc Wilcoxon signed-rank tests

revealed a significant difference between block 1 and 2 ($p = 0.002$), and all other blocks ($p <$

0.001), except block 3 and 4, where no significant difference was found ($p = 0.697$). The post

hoc Wilcoxon rank-sum test for condition type showed no significant difference ($p = 0.153$). This

indicates that participants took longer to respond for more abstract questions.

Figure 2. Behavioral metrics for different gaze conditions across fixed effects variables (n = 45).

Box plots show the IQR with medians as horizontal lines. Mean values with 95% confidence

intervals are marked as colored points, and outliers as individual points. Only known response

trials were included. A) Normalized correct responses are shown across blocks and B) different

question types. C) Saccade frequency, D) saccade amplitude, E) pupil diameter and F) answer

durations are shown across blocks. Error bars denote upper and lower quartiles, boxes denote

interquartile ranges, solid horizontal bars denote medians.

Presaccadic potentials were analyzed for main versus control tasks, with the control task

including both visual and memory saccade tasks. The analyzed interval (-200 to 50 ms) is

referenced within the overall signal (c.f. Figure S6 A), as determined based on typical saccadic

latency time (Rayner et al., 1983) to avoid intervening saccades. Within-condition baseline

reference means that the reported effects reflect relative changes rather than absolute increases

or decreases in activity. In the specified interval, the main task is characterized by a presaccadic

positive frontal and negative parietal potential, while the control task shows a positive parietal

potential and a smaller frontal potential in contrast to the main task (see Figure 3 A). Potentials
peak at approximately 20 ms prior to saccade onset. At saccade onset, a negative frontal and
positive parietal spike can be observed for both task conditions. Averaged presaccadic
amplitudes are larger for the main task in frontal ROIs and smaller for the main task in parietal
ROIs compared to the control task. Supplementary analyses confirmed that the observed ERP
differences between main and control tasks are not driven by differences in saccade velocity or
amplitude (see Supplementary Results and Figure S5). A larger polarization is also observed for
frontal and parietal ROIs for the main task compared to the control task, as confirmed in the
topographical visualization. Topographic maps illustrate the evolving spatial distribution of ERPs
leading up to saccade onset (see Figure 3 E). As already indicated by the ERPs, the main task
shows a strong positive frontal and negative parietal potential developing over time, peaking just
before the saccade. In contrast, the control task exhibits a prominent positive potential in the
parietal regions, which extends into the posterior frontal regions. However, overall, the frontal
electrodes show less pronounced, negative potential. A larger cluster of activity is visible over
the parietal electrodes. This pattern is congruent with previously reported presaccadic EEG
activity (Everling et al., 1997; Richards, 2003; Becker et al., 1972; Balaban & Weinstein, 1985). To
quantify these activity changes, we used repeated measures ANOVA using ERP amplitudes
averaged over the presaccadic time window (-150 to -20 ms), with "task" (main/control) and
"ROI" (frontal/parietal) as factors. This analysis showed significantly higher activity change
preceding saccades during the thinking task in both channel groups (see Figure 3 B). A significant
main effect of "task" ($F(1,28) = 16.205, p < 0.001, \omega^2 = 0.137$) and "ROI"   ($F(1,28) =$
$5.528, p = 0.026, \omega^2 = 0.114$) was found with a large and medium to large effect size,
respectively. Additionally, a significant interaction effect of "task" × "ROI" ($F(1,28) =$
$25.496, p < 0.001, \omega^2 = 0.411$) with a large effect size was observed. Post hoc paired t-tests
revealed a significant difference between main and control tasks for the frontal channels ($p <$
0.001), as well as for the parietal channels ($p = 0.003$). Significant differences were also
identified for the main task between frontal and parietal ROI ($p < 0.001$), while no significant
difference was found for the control task between frontal and parietal ROI ($p = 0.130$).

As previous analyses showed that the main and the control tasks had distinct neural components,

we next analyzed the absolute magnitude of pre-saccadic EEG activity changes (-150 to -20 ms)

independent of potentially confounding effects of signal polarity (see Figure 3 C). We compared

the absolute values of ERP amplitudes by conducting repeated measures ANOVA with "task"

(main/control) and "ROI" (frontal/parietal) as factors. A significant main effect of "task"

($F(1,28) = 10.087, p = 0.004, \omega^2 = 0.145$) was found with a large effect size. No significant

main effect was found for "ROI" ($F(1,28) = 1.887, p = 0.180, \omega^2 = 0.009$). Additionally, a

significant interaction effect of "task" × "ROI" ($F(1,28) = 6.279, p = 0.018, \omega^2 = 0.047$) with a

small effect size was observed. Post hoc paired t-tests revealed a significant difference between

main and control tasks for the frontal channels ($p < 0.001$), but no significant difference for the

parietal channels ($p = 0.265$). Significant differences were also identified for the main task

between frontal and parietal ROI ($p = 0.035$), while no significant difference was found for the

control task between frontal and parietal ROI ($p = 0.477$). These analyses show that absolute

signal amplitude changes in prefrontal channels in the main task were greater than those in

parietal channels and when compared to the saccadic preparatory activity in the control tasks.

The sliding window analysis revealed the time points at which ERP signals significantly separated

between the main and control tasks in the frontal and parietal ROIs (see Figure 3 A). In the frontal

ROI, signal separation was first observed at -140 ms prior to saccade onset ($p = 0.038$). In the

parietal ROI, signal separation occurred later, at -110 ms prior to saccade onset ($p = 0.042$). The

results demonstrate a 30 ms difference between the onset of significant signal change in the

frontal vs. parietal ROIs, with the frontal onset preceding the parietal one. This temporal

difference underscores the distinct timing of neural processes involved in NVS across these

regions.

In the next step, we analyzed power spectral density for alpha, beta, and low-gamma bands (see

Figure 3 D). For each band separately, we conducted an ANOVA with factors "Task" and "ROI".

These ANOVAs showed a significant increase in power spectral density in the main task due to a

significant "Task" main effect for beta ($F(1,28) = 7.414, p = 0.011, \omega^2 = 0.029$) and gamma

bands ($F(1,28) = 9.016, p = 0.006, \omega^2 = 0.058$). There was also a nearly-significant main effect

of Task in alpha band ($F(1,28) = 4.14, p = 0.051, \omega^2 = 0.011$). There was a significant ROI main

effect in alpha ($F(1, 28) = 6.171, p = 0.019, \omega^2 = 0.036$) and a near significant ROI main effect

in beta bands ($F(1, 28) = 4.174, p = 0.051, \omega^2 = 0.020$). No Task x ROI interaction effect was

significant ($p > 0.5$).

Lastly, source localization revealed strong, focused neural activity primarily in the right

hemisphere. Figure 3 F shows clusters in the frontal regions, with the peak activation in the right

superior and middle frontal gyrus ($peak\ t = 9.3$). Activity extends posteriorly into the motor

cortex and precentral gyrus, and medially, involving orbitofrontal and medial areas. Notable

activation clusters was also found bilaterally in the superior frontal cortex ($t = 8.17$), precentral

and postcentral regions ($t = 8.21$), the insula ($t = 8.14$), and rostral anterior cingulate ($t = $

7.98). Additional activation clusters, though less intense, occurred in the superior parietal cortex

($t = 7.95$), medial orbitofrontal cortex ($t = 7.88$), and precuneus ($t = 7.97$). This pattern was

different in control tasks (see Supplementary Results).

Figure 3. EEG results for main vs. visual and memory control tasks. A) ROI channels selection and ERPs relative to saccade onset for main and the control tasks; dashed lines indicate significant amplitude difference between ERP signals in main and control tasks; grey box denotes the baseline period; shaded areas denote 95% confidence intervals; B) Mean ERP amplitudes across tasks. C) Absolute values of ERP amplitudes across conditions. D) Modulation of alpha, beta and gamma frequency across tasks. E) Topographic maps of presaccadic activity across main and control tasks. F) Source localization. Error bars denote upper and lower quartiles, boxes denote interquartile ranges, solid horizontal bars denote medians.

Discussion

Non-visual saccades are a not well understood phenomenon accompanying conceptual operations. Here, we show that NVS are directly preceded by neural activity changes in frontal EEG channels. The source localization analysis additionally indicated that the pre-NVS activity originates in the premotor cortex. These pre-NVS changes are qualitatively different from signals preceding voluntary saccades indicating that NVS are driven by other processes than oculomotor preparation. While pre-NVS activity changes are seen in both frontal and posterior channels, only the activity change in frontal channels significantly differ from that preceding voluntary saccades. As we used abstract conceptual tasks, we can determine that neural activity before NVS is not merely reflecting mnemonic processing or retrieval of visuospatial content.

It has long been a subject of dispute in philosophy, psychology, and neuroscience whether concepts are represented in pictorial/visuospatial or verbal/abstract form (see e.g., Pylyshyn, 2003). Edward Tolman proposed that past experience and spatial cognition are arranged in the form of spatially organized maps (Tolman, 1948). This theory implied that the knowledge map builds on representations of space, and the two likely share neural components. Since then, it has been confirmed that spatial maps are organized by the hippocampal-entorhinal system (O'Keefe & Dostrovsky, 1971) using a grid-like code (Hafting et al., 2005), and that this system also organizes conceptual knowledge using similar principles (Constantinescu et al., 2016). However, just as spatial maps need to be translated into neural codes understood by the motor system, stored conceptual knowledge needs to be processed for daily operations. One recently

suggested possibility is that the hippocampus plays a crucial role in binding elements of a visual

scene by receiving corollary discharge signals that accompany visual exploration. By receiving

these signals, the hippocampus may be able to temporally align incoming visual information with

the sequence of eye movements, thereby supporting the formation of coherent scene

representations in memory (Katz et al., 2022; Leszczynski et al., 2024). It is conceivable that these

same corollary discharge signals are reinstated during memory retrieval to recreate aspects of

the original exploratory state, thereby enabling attention to be internally shifted between

different components of the remembered scene or conceptual space. This possibility aligns with

models of memory that propose dynamic and reconstructive processes, where internally

generated signals guide the reactivation of stored representations. In this context, the MTL may

contribute not only to encoding but also to the active reconstruction of memory traces via

simulated sensory-motor interactions.

It has been proposed that the parietal lobe acts as an interface connecting the MTL memory

system with areas responsible for actions (Whitlock, 2017). The main role of the PPC seems to lie

in integrating representations of world-centered space into body-centered maps used for

locomotion and actions (Whitlock et al., 2008). In the cognitive domain, there is also strong

evidence showing the parietal lobe's involvement in long-term memory search but not in the

recognition of remembered objects, indicating that the parietal cortex is not merely involved in

storage but rather in the active exploration of the knowledge space (Cabeza et al., 2008). Motor

(and oculomotor) areas in the posterior parietal cortex share strong reciprocal connections with

the premotor cortex (Pandya & Kuypers, 1969), as both regions perform complementary roles in

controlling actions (Curtis, 2006; Cisek, 2007; Westendorff et al., 2010; Pilacinski and Lindner,

2019). This tight fronto-parietal interrelation in processing actions seems to be retained when it

comes to operating on memory. Both PM and PPC seem to process different stages of search and

retrieval of mnemonic content, with PM controlling memory search and PPC successful retrieval

of information (Kapur et al., 1995). Attentional shifts underlying memory search were

demonstrated to increase functional connectivity between PM and the parietal cortex,

suggesting a tight information exchange in the process of retrieving and manipulating mnemonic

information, which relies critically on selecting relevant and suppressing irrelevant information

(Heinen et al., 2017). But why would the premotor cortex be involved in thinking? The answer

lies in how it is involved in actions.

The primary function of the premotor cortex seems to be its ability to represent complex,

coordinated multi-joint movements (Graziano and Aflalo, 2007) such as those needed for

programming hand trajectories or gestures (Pilacinski and Lindner, 2019; Wong et al., 2019), or

handwriting (Kadmon Harpaz et al., 2014). The premotor cortex appears to process both motor

and semantic levels (i.e., higher-order goals) of actions (Rizzolatti et al., 1996) and is critically

involved in executing relevant and suppressing irrelevant actions from among options

represented by the parietal cortex (Cisek, 2007; Lindner et al., 2010; Pilacinski and Lindner, 2019).

Neurophysiological data further suggest that the key function of the premotor cortex in

programming movements is selecting from among competing action plans represented by the

posterior parietal cortex, using weighting input from more frontal areas (Cisek, 2007). The tight

connectivity between PM and PPC further suggests that this inter-areal exchange facilitates the

promotion of relevant and suppression of irrelevant information, e.g., actions. In conceptual

processing, such reciprocal exchange between PM and PPC may be utilized in a similar way, in

which frontal areas (mediated by PM) select between competing information retrieved from

long-term memory by parietal areas through top-down modulation, as suggested by ERP

dynamics. This idea seems to be further supported by spectral analysis data, showing that

immediately before each NVS, we observe a substantial increase in all three EEG bands we

measured, indicating a complex computational process required by top-down and bottom-up

modulation of cognitive content (Karch et al., 2016; Strube et al., 2021). These distinctive

patterns of activity may reflect the evaluation of conceptual information by PM in a way similar

to that during sensorimotor decision-making (Cisek et al., 2009; Thura and Cisek, 2014).

We believe that the gradual change of activity before NVS likely represents neural computations

required for evaluating conceptual information, similar to evidence accumulation during

sensorimotor decision-making (Cisek et al., 2009; Thura and Cisek, 2014). The subsequent

saccade could then occur due to the shifting of attention to another sub-element of the currently-

processed part of conceptual space. It has been shown that covert shifts of attention indeed

result in involuntary eye movements, likely due to direct connections between frontal eye fields

and the superior colliculus (Peterson et al., 2004). Complex conceptual operations require

multiple such shifts of attention between concepts or their elements, and these shifts are likely

to underlie NVS (Scholz et al., 2017) in a way similar to how saccades scan the visual space,

identifying its (semantic) components. The premotor theory of attention proposes that covert

attentional shifts and saccade planning are identical or closely-related processes involving mainly

premotor cortex (Rizzolatti et al., 1987; Smith and Schenk, 2012). Our results suggest that non-

visual saccades arise from specific processing within the premotor cortex, distinct from

mechanisms involved in voluntary saccade planning. This interpretation is further supported by

the fact that pre-NVS activity remained consistent even when subjects were instructed to

withhold eye movements, indicating that non-visual saccades are preceded by a dedicated neural

substrate. Nonetheless, the fact that neural activity preceding NVS seems to originate in the

premotor cortex, a common neuroanatomical substrate between NVS and voluntary saccades,

may explain why mental operations on concepts trigger NVS. As frontal eye fields directly project

to the superior colliculus, it is likely that mental operations involving premotor computations

while processing concepts may collaterally initiate eye movements as a byproduct of thinking.

The fact that restricting gaze did not substantially worsen mnemonic performance supports the

notion that NVS are an epiphenomenon rather than a functional component of conceptual

processing.

Limitations

Our study focuses on cortical processing responsible for thinking, considering the role of frontal

and parietal areas in conceptual processing triggering NVS. Yet, human thinking involves

numerous subcortical structures such as the hippocampus or basal ganglia, which are critically

involved in information storage and selection. Due to EEG limitations, we cannot address the role

these subcortical structures play in processing resulting in NVS. Although posterior parietal cortex

and premotor cortex can directly exchange information through a one-synapse connection

(Pandya & Kuypers, 1969), theoretically allowing for a direct exchange of information, we believe

there may be substantial involvement of subcortical structures in how conceptual information is

processed (Cisek et al., 2009). Due to EEG's poor spatial resolution, our reasoning about the

premotor cortex is mainly based on assumptions given its role in eye movements and
sensorimotor processing rather than direct functional localization. We do acknowledge,
however, that conceptual operations involve a long cascade of processes including emotional
evaluation, evidence weighting, and everything that constitutes individual human cognition.
Although this research focused on a specific byproduct of conceptual processing (i.e., NVS) and
its accompanying neural components, we recognize the complex and dynamic processing within
the brain resulting in the human ability to operate on abstract concepts.

In our results we seemingly uncovered an interplay between the type of memory search (open
vs. structured) and the instruction to maintain fixation, which cannot be fully explained based on
our data. Restricting NVS did not substantially impair memory performance in the open questions
task, suggesting they are a byproduct of conceptual processing rather than a function. Yet,
restricted gaze resulted in worse performance in structured retrieval task, meaning that either
gaze fixation may require additional cognitive demands, or NVS help in some types of conceptual
processing. This remains to be determined.

Lastly, our control task results were clearly biased towards the visual saccade task and did not
allow us to fully assess the role of working memory processing and maintenance in modulating
prefrontal activity preceding NVS. We attempted to do it using the memory saccade task, which
was strongly underpowered in our case. Despite this, the data we obtained suggest that brain
activity preceding NVS is qualitatively different from that in the memory saccade task and does
not merely represent maintenance of mnemonic content. Further research is required to assess
this complex processing network, and we believe NVS can be a good behavioral model for that.
The interplay between NVS and content reproduction type also requires further investigation.

Conclusions

In recent years, multiple authors have suggested that conceptual knowledge builds on
sensorimotor circuitry in the brain, operating in low-dimensional conceptual spaces organized
similarly to the representation of physical space. The available evidence suggests that organizing,
storing, and manipulating conceptual knowledge is shared by temporal and parietal systems, with
the involvement of other areas, such as the frontal cortex, being less well understood. Through

the use of non-visual saccades as a behavioural model, our study extends established views on

the organization of human conceptual knowledge by showing that the premotor cortex is

involved in exploring and manipulating items within conceptual space.

References and Notes

Abdoun, Oussama. "NeuroMap: A Spline-Based Interactive Open-Source Software for

Spatiotemporal Mapping of 2D and 3D MEA Data." *Frontiers in Neuroinformatics* 4

(2011): 119. https://doi.org/10.3389/fninf.2010.00119.

Abe, Takashi, Hiroshi Nittono, and Tadao Hori. "Current Sources of the Brain Potentials

before Rapid Eye Movements in Human REM Sleep." *International Congress Series*

1270 (August 2004): 241–44. https://doi.org/10.1016/j.ics.2004.04.092.

Agresti, Alan. *An Introduction to Categorical Data Analysis*. Wiley, 2007.

https://doi.org/10.1002/0470114754.

Andersen, Richard A., and Christopher A. Buneo. "Intentional Maps in Posterior Parietal

Cortex." *Annual Review of Neuroscience* 25, no. 1 (March 2002): 189–220.

https://doi.org/10.1146/annurev.neuro.25.112701.142922.

Anderson, Carolyn J., Jay Verkuilen, and Timothy R. Johnson. *Applied Generalized Linear*

Mixed Models: Continuous and Discrete Data - For the Social and Behavioral Sciences.

Springer, 2012.

Babapoor-Farrokhran, Sahand, Martin Vinck, Thilo Womelsdorf, and Stefan Everling. "Theta

and Beta Synchrony Coordinate Frontal Eye Fields and Anterior Cingulate Cortex

during Sensorimotor Mapping." *Nature Communications* 8, no. 1 (February 2017):

13967. https://doi.org/10.1038/ncomms13967.

Balaban, Carey D., and Joel M. Weinstein. "The Human Pre-Saccadic Spike Potential:

Influences of a Visual Target, Saccade Direction, Electrode Laterality and Instructions

to Perform Saccades." *Brain Research* 347, no. 1 (November 1985): 49–57.

https://doi.org/10.1016/0006-8993(85)90888-1.

Becker, Wolfgang, Ottomar Hoehne, Katsuhiko Iwase, and Hans H. Kornhuber.

"Bereitschaftspotential, Prämotorische Positivierung Und Andere Hirnpotentiale Bei

Sakkadischen Augenbewegungen." *Vision Research* 12, no. 3 (March 1972): 421–36.

https://doi.org/10.1016/0042-6989(72)90087-9.

Bellmund, Jacob L. S., Peter Gärdenfors, Edvard I. Moser, and Christian F. Doeller.

"Navigating Cognition: Spatial Codes for Human Thinking." *Science* 362, no. 6415

(November 2018). https://doi.org/10.1126/science.aat6766.

Berryhill, Marian E., Lisa Phuong, Lauren Picasso, Roberto Cabeza, and Ingrid R. Olson.

"Parietal Lobe and Episodic Memory: Bilateral Damage Causes Impaired Free Recall of

Autobiographical Memory." *The Journal of Neuroscience* 27, no. 52 (December 2007):

14415–23. https://doi.org/10.1523/JNEUROSCI.4163-07.2007.

Bijvank, J. A. Nij, A. Petzold, L. J. Balk, H. S. Tan, B. M. J. Uitdehaag, M. Theodorou, and L. J.

van Rijn. "A Standardized Protocol for Quantification of Saccadic Eye Movements:

DEMoNS." *PLOS ONE* 13, no. 7 (July 2018): e0200695.

https://doi.org/10.1371/journal.pone.0200695.

Brandmeyer, Tracy, and Arnaud Delorme. "Closed-Loop Frontal Midlineθ Neurofeedback: A

Novel Approach for Training Focused-Attention Meditation." *Frontiers in Human*

Neuroscience 14 (June 2020): 246. https://doi.org/10.3389/fnhum.2020.00246.

Bruce, C. J., M. E. Goldberg, M. C. Bushnell, and G. B. Stanton. "Primate Frontal Eye Fields.

II. Physiological and Anatomical Correlates of Electrically Evoked Eye Movements."

Journal of Neurophysiology 54, no. 3 (September 1985): 714–34.

https://doi.org/10.1152/jn.1985.54.3.714.

Cabeza, Roberto, Elisa Ciaramelli, Ingrid R. Olson, and Morris Moscovitch. "The Parietal

Cortex and Episodic Memory: An Attentional Account." *Nature Reviews Neuroscience*

9, no. 8 (August 2008): 613–25. https://doi.org/10.1038/nrn2459.

Casals, Martí, Montserrat Girabent-Farrés, and Josep L. Carrasco. "Methodological Quality

and Reporting of Generalized Linear Mixed Models in Clinical Medicine (2000–2012): A

Systematic Review." *PLoS ONE* 9, no. 11 (November 2014): e112653.

https://doi.org/10.1371/journal.pone.0112653.

Cisek, Paul. "Cortical Mechanisms of Action Selection: The Affordance Competition

Hypothesis." *Philosophical Transactions of the Royal Society B: Biological Sciences* 362,

no. 1485 (September 2007): 1585–99. https://doi.org/10.1098/rstb.2007.2054.

Cisek, Paul, Geneviève Aude Puskas, and Stephany El-Murr. "Decisions in Changing

Conditions: The Urgency-Gating Model." *The Journal of Neuroscience* 29, no. 37

(September 2009): 11560–71. https://doi.org/10.1523/JNEUROSCI.1844-09.2009.

Constantinescu, Alexandra O., Jill X. O'Reilly, and Timothy E. J. Behrens. "Organizing

Conceptual Knowledge in Humans with a Gridlike Code." *Science* 352, no. 6292 (June

2016): 1464–68. https://doi.org/10.1126/science.aaf0941.

Cordeiro, Gauss M., and Alexandre B. Simas. "The Distribution of Pearson Residuals in

Generalized Linear Models." *Computational Statistics & Data Analysis* 53, no. 9 (July

2009): 3397–3411. https://doi.org/10.1016/j.csda.2009.02.025.

Cothi, William de, and Hugo J. Spiers. "Spatial Cognition: Goal-Vector Cells in the Bat

Hippocampus." *Current Biology* 27, no. 6 (March 2017): R239–41.

https://doi.org/10.1016/j.cub.2017.01.061.

Curtis, C. E. "Prefrontal and Parietal Contributions to Spatial Working Memory."

Neuroscience 139, no. 1 (April 2006): 173–80.

https://doi.org/10.1016/j.neuroscience.2005.04.070.

Curtis, Clayton E., Vikas Y. Rao, and Mark D'Esposito. "Maintenance of Spatial and Motor

Codes during Oculomotor Delayed Response Tasks." *The Journal of Neuroscience* 24,

no. 16 (April 2004): 3944–52. https://doi.org/10.1523/JNEUROSCI.5640-03.2004.

Darby, David G., Anna C. Nobre, Venkatesan Thangaraj, Robert Edelman, M.-Marsel

Mesulam, and Steven Warach. "Cortical Activation in the Human Brain during Lateral

Saccades Using EPISTAR Functional Magnetic Resonance Imaging." *NeuroImage* 3, no.

1 (February 1996): 53–62. https://doi.org/10.1006/nimg.1996.0006.

Delorme, Arnaud. "EEG Is Better Left Alone." *Scientific Reports* 13, no. 1 (February 2023):

2372. https://doi.org/10.1038/s41598-023-27528-0.

Delorme, Arnaud, and Scott Makeig. "EEGLAB: An Open Source Toolbox for Analysis of

Single-Trial EEG Dynamics Including Independent Component Analysis." *Journal of*

Neuroscience Methods 134, no. 1 (March 2004): 9–21.

https://doi.org/10.1016/j.jneumeth.2003.10.009.

Desikan, Rahul S., Florent Ségonne, Bruce Fischl, Brian T. Quinn, Bradford C. Dickerson,

Deborah Blacker, Randy L. Buckner, et al. "An Automated Labeling System for

Subdividing the Human Cerebral Cortex on MRI Scans into Gyral Based Regions of

Interest." *NeuroImage* 31, no. 3 (July 2006): 968–80.

https://doi.org/10.1016/j.neuroimage.2006.01.021.

Djebbara, Z., Fich, L. B., Petrini, L., & Gramann, K. (2019). Sensorimotor brain dynamics

reflect architectural affordances. *Proceedings of the National Academy of Sciences*,

116(29), 14769-14778. https://doi.org/10.1073/pnas.1900648116

Doherty-Sneddon, G., and F. G. Phelps. "Gaze Aversion: A Response to Cognitive or Social

Difficulty?" *Memory & Cognition* 33, no. 4 (June 2005): 727–33.

https://doi.org/10.3758/BF03195338.

Dunn, Peter K., and Gordon K. Smyth. *Generalized Linear Models With Examples in R*.

Springer New York, 2018. https://doi.org/10.1007/978-1-4419-0118-7.

Ehrlichman, Howard, and John Barrett. "'Random' Saccadic Eye Movements during Verbal-

Linguistic and Visual-Imaginal Tasks." *Acta Psychologica* 53, no. 1 (April 1983): 9–26.

https://doi.org/10.1016/0001-6918(83)90013-6.

Ehrlichman, Howard, and Dragana Micic. "Why Do People Move Their Eyes When They

Think?" *Current Directions in Psychological Science* 21, no. 2 (April 2012): 96–100.

https://doi.org/10.1177/0963721412436810.

Ehrlichman, Howard, Dragana Micic, Amber Sousa, and John Zhu. "Looking for Answers:

Eye Movements in Non-Visual Cognitive Tasks." *Brain and Cognition* 64, no. 1 (June

2007): 7–20. https://doi.org/10.1016/j.bandc.2006.10.001.

Everling, S., P. Krappmann, and H. Flohr. "Cortical Potentials Preceding Pro- and

Antisaccades in Man." *Electroencephalography and Clinical Neurophysiology* 102, no. 4

(April 1997): 356–62. https://doi.org/10.1016/S0013-4694(96)96569-4.

Freeden, Willi. "Spherical Spline Interpolation—Basic Theory and Computational Aspects."

Journal of Computational and Applied Mathematics 11, no. 3 (December 1984): 367–

75. https://doi.org/10.1016/0377-0427(84)90011-6.

Gramfort, Alexandre. "MEG and EEG Data Analysis with MNE-Python." *Frontiers in*

Neuroscience 7 (2013): 267. https://doi.org/10.3389/fnins.2013.00267.

Graziano, Michael S. A., and Tyson N. Aflalo. "Rethinking Cortical Organization." *The*

Neuroscientist 13, no. 2 (April 2007): 138–47.

https://doi.org/10.1177/1073858406295918.

Gurtner, Lilla M., Matthias Hartmann, and Fred W. Mast. "Eye Movements during Visual

Imagery and Perception Show Spatial Correspondence but Have Unique Temporal

Signatures." *Cognition* 210 (May 2021): 104597.

https://doi.org/10.1016/j.cognition.2021.104597.

Hafting, Torkel, Marianne Fyhn, Sturla Molden, May-Britt Moser, and Edvard I. Moser.

"Microstructure of a Spatial Map in the Entorhinal Cortex." *Nature* 436, no. 7052

(August 2005): 801–6. https://doi.org/10.1038/nature03721.

Harpaz, Naama Kadmon, Tamar Flash, and Ilan Dinstein. "Scale-Invariant Movement

Encoding in the Human Motor System." *Neuron* 81, no. 2 (January 2014): 452–62.

https://doi.org/10.1016/j.neuron.2013.10.058.

Heinen, Klaartje, Eva Feredoes, Christian C. Ruff, and Jon Driver. "Functional Connectivity

between Prefrontal and Parietal Cortex Drives Visuo-Spatial Attention Shifts."

Neuropsychologia 99 (May 2017): 81–91.

https://doi.org/10.1016/j.neuropsychologia.2017.02.024.

Hollander, John, and Stephanie Huette. "Extracting Blinks from Continuous Eye-Tracking

Data in a Mind Wandering Paradigm." *Consciousness and Cognition* 100 (April 2022):

103303. https://doi.org/10.1016/j.concog.2022.103303.

Imaoka, Yu, Andri Flury, and Eling D. de Bruin. "Assessing Saccadic Eye Movements With

Head-Mounted Display Virtual Reality Technology." *Frontiers in Psychiatry* 11

(September 2020): 572938. https://doi.org/10.3389/fpsyt.2020.572938.

Inc, The MathWorks. "MATLAB (v9.14.0 (R2023a))," 2023.

Johansson, Roger, Marcus Nyström, Richard Dewhurst, and Mikael Johansson. "Eye-

Movement Replay Supports Episodic Remembering." Research Square Platform,

February 2021. https://doi.org/10.21203/rs.3.rs-275969/v1.

Kang, Seung Suk, Timothy J. Lano, and Scott R. Sponheim. "Distortions in EEG Interregional

Phase Synchrony by Spherical Spline Interpolation: Causes and Remedies."

Neuropsychiatric Electrophysiology 1, no. 1 (December 2015): 9.

https://doi.org/10.1186/s40810-015-0009-5.

Kapur, S., Craik, F. I., Jones, C., Brown, G. M., Houle, S., & Tulving, E. (1995). Functional role of the prefrontal cortex in retrieval of memories: a PET study. *Neuroreport*, *6*(14), 1880–1884. https://doi.org/10.1097/00001756-199510020-00014

Karch, Susanne, Fabian Loy, Daniela Krause, Sandra Schwarz, Jan Kiesewetter, Felix Segmiller, Agnieszka I. Chrobok, Daniel Keeser, and Oliver Pogarell. "Increased Event-Related Potentials and Alpha-, Beta-, and Gamma-Activity Associated with Intentional Actions." *Frontiers in Psychology* 7 (January 2016). https://doi.org/10.3389/fpsyg.2016.00007.

Koval, Michael J., R. Matthew Hutchison, Stephen G. Lomber, and Stefan Everling. "Effects of Unilateral Deactivations of Dorsolateral Prefrontal Cortex and Anterior Cingulate Cortex on Saccadic Eye Movements." *Journal of Neurophysiology* 111, no. 4 (February 2014): 787–803. https://doi.org/10.1152/jn.00626.2013.

Kovalenko, Lyudmyla Y., and Niko A. Busch. "Probing the Dynamics of Perisaccadic Vision with EEG." *Neuropsychologia* 85 (May 2016): 337–48. https://doi.org/10.1016/j.neuropsychologia.2015.12.012.

Larson, Eric, Alexandre Gramfort, Denis A. Engemann, Jaakko Leppakangas, Christian Brodbeck, Mainak Jas, Teon L. Brooks, Jona Sassenhagen, Daniel McCloy, and Martin Luessi. "MNE-Python (v1.7.1)." Zenodo, 2024.

Larsson, Linnea, Marcus Nystrom, and Martin Stridh. "Detection of Saccades and Postsaccadic Oscillations in the Presence of Smooth Pursuit." *IEEE Transactions on Biomedical Engineering* 60, no. 9 (September 2013): 2484–93. https://doi.org/10.1109/TBME.2013.2258918.

Leszczynski, M., Espinal, E., Smith, E., Schevon, C., Sheth, S., & Schroeder, C. E. (2024). Eye movements organize excitability state, information coding and network connectivity in the human hippocampus. BioRxiv, 2024-12.

Lindner, Axel, Iyer A., Kagan I., and Andersen, Richard A. "Human Posterior Parietal Cortex Plans Where to Reach and What to Avoid." *The Journal of Neuroscience* 30, no. 35 (September 2010): 11715–25. https://doi.org/10.1523/JNEUROSCI.2849-09.2010.

Liu, Bo, Qi-Chao Zhao, Yuan-Yuan Ren, Qing-Ju Wang, and Xue-Lian Zheng. "An Elaborate

Algorithm for Automatic Processing of Eye Movement Data and Identifying Fixations in Eye-Tracking Experiments." *Advances in Mechanical Engineering* 10, no. 5 (May 2018): 1–15. https://doi.org/10.1177/1687814018773678.

Mack, David J., Sandro Belfanti, and Urs Schwarz. "The Effect of Sampling Rate and Lowpass Filters on Saccades – A Modeling Approach." *Behavior Research Methods* 49, no. 6 (December 2017): 2146–62. https://doi.org/10.3758/s13428-016-0848-4.

Martarelli, Corinna S., Fred W. Mast, and Matthias Hartmann. "Time in the Eye of the Beholder: Gaze Position Reveals Spatial-Temporal Associations during Encoding and Memory Retrieval of Future and Past." *Memory & Cognition* 45, no. 1 (January 2017): 40–48. https://doi.org/10.3758/s13421-016-0639-2.

McDowell, Jennifer E., Kara A. Dyckman, Benjamin P. Austin, and Brett A. Clementz. "Neurophysiology and Neuroanatomy of Reflexive and Volitional Saccades: Evidence from Studies of Humans." *Brain and Cognition* 68, no. 3 (December 2008): 255–70. https://doi.org/10.1016/j.bandc.2008.08.016.

McPeek, Robert M., Alexander A. Skavenski, and Ken Nakayama. "Concurrent Processing of Saccades in Visual Search." *Vision Research* 40, no. 18 (August 2000): 2499–2516. https://doi.org/10.1016/S0042-6989(00)00102-4.

Meghanathan, Radha Nila, Cees van Leeuwen, Marcello Giannini, and Andrey R. Nikolaev. "Neural Correlates of Task-Related Refixation Behavior." *Vision Research* 175 (October 2020): 90–101. https://doi.org/10.1016/j.visres.2020.07.001.

Meister, Miriam L. R., and Elizabeth A. Buffalo. "Neurons in Primate Entorhinal Cortex Represent Gaze Position in Multiple Spatial Reference Frames." *The Journal of Neuroscience* 38, no. 10 (March 2018): 2430–41. https://doi.org/10.1523/JNEUROSCI.2432-17.2018.

Micic, Dragana, Howard Ehrlichman, and Rebecca Chen. "Why Do We Move Our Eyes While Trying to Remember? The Relationship between Non-Visual Gaze Patterns and Memory." *Brain and Cognition* 74, no. 3 (December 2010): 210–24. https://doi.org/10.1016/j.bandc.2010.07.014.

Munoz, Douglas P. "Commentary: Saccadic Eye Movements: Overview of Neural Circuitry."

In *Progress in Brain Research*, 89–96. Elsevier, 2002. https://doi.org/10.1016/S0079-
6123(02)40044-1.

Nikolaev, Andrey R., Peter Jurica, Chie Nakatani, Gijs Plomp, and Cees van Leeuwen. "Visual
Encoding and Fixation Target Selection in Free Viewing: Presaccadic Brain Potentials."
Frontiers in Systems Neuroscience 7 (2013): 26.
https://doi.org/10.3389/fnsys.2013.00026.

Nouira, Ibtihel, Asma Ben Abdallah, and Mohamed Hedi Bedoui. "Three-Dimensional
Interpolation Methods to Spatiotemporal EEG Mapping during Various Behavioral
States." *Signal, Image and Video Processing* 10, no. 5 (July 2016): 943–49.
https://doi.org/10.1007/s11760-015-0844-7.

Nyström, Marcus, and Kenneth Holmqvist. "An Adaptive Algorithm for Fixation, Saccade,
and Glissade Detection in Eyetracking Data." *Behavior Research Methods* 42, no. 1
(February 2010): 188–204. https://doi.org/10.3758/BRM.42.1.188.

O'Keefe, J., and J. Dostrovsky. "The Hippocampus as a Spatial Map. Preliminary Evidence
from Unit Activity in the Freely-Moving Rat." *Brain Research* 34, no. 1 (November
1971): 171–75. https://doi.org/10.1016/0006-8993(71)90358-1.

Pandya, Deepak N., and Henricus G. J. M. Kuypers. "Cortico-Cortical Connections in the
Rhesus Monkey." *Brain Research* 13, no. 1 (March 1969): 13–36.
https://doi.org/10.1016/0006-8993(69)90141-3.

Paré, Martin, and Robert H. Wurtz. "Monkey Posterior Parietal Cortex Neurons
Antidromically Activated From Superior Colliculus." *Journal of Neurophysiology* 78, no.
6 (December 1997): 3493–97. https://doi.org/10.1152/jn.1997.78.6.3493.

Pascual-Marqui, R. D. "Standardized Low-Resolution Brain Electromagnetic Tomography
(sLORETA): Technical Details." *Methods and Findings in Experimental and Clinical*
Pharmacology 24 Suppl D (2002): 5–12.

Peelen, Marius V., Dirk J. Heslenfeld, and Jan Theeuwes. "Endogenous and Exogenous
Attention Shifts Are Mediated by the Same Large-Scale Neural Network." *NeuroImage*
22, no. 2 (June 2004): 822–30. https://doi.org/10.1016/j.neuroimage.2004.01.044.

Perrin, F., J. Pernier, O. Bertrand, and J. F. Echallier. "Spherical Splines for Scalp Potential

and Current Density Mapping." *Electroencephalography and Clinical Neurophysiology*

72, no. 2 (February 1989): 184–87. https://doi.org/10.1016/0013-4694(89)90180-6.

Peterson, Matthew S., Arthur F. Kramer, and David E. Irwin. "Covert Shifts of Attention

Precede Involuntary Eye Movements." *Perception & Psychophysics* 66, no. 3 (April

2004): 398–405. https://doi.org/10.3758/BF03194888.

Pierrot-Deseilligny, Charles, Sophie Rivaud, Bertrand Gaymard, René Müri, and Anne-

Isabelle Vermersch. "Cortical Control of Saccades." *Annals of Neurology* 37, no. 5 (May

1995): 557–67. https://doi.org/10.1002/ana.410370504.

Pilacinski, Artur, and Axel Lindner. "Distinct Contributions of Human Posterior Parietal and

Dorsal Premotor Cortex to Reach Trajectory Planning." *Scientific Reports* 9, no. 1

(February 2019): 1962. https://doi.org/10.1038/s41598-019-39188-0.

Pylyshyn, Zenon. "Return of the Mental Image: Are There Really Pictures in the Brain?"

Trends in Cognitive Sciences 7, no. 3 (March 2003): 113–18.

https://doi.org/10.1016/S1364-6613(03)00003-2.

Rayner, Keith, Maria L. Slowiaczek, Charles Clifton, and James H. Bertera. "Latency of

Sequential Eye Movements: Implications for Reading." *Journal of Experimental*

Psychology: Human Perception and Performance 9, no. 6 (1983): 912–22.

https://doi.org/10.1037/0096-1523.9.6.912.

Revankar, Gajanan S., Noriaki Hattori, Yuta Kajiyama, Tomohito Nakano, Masahito Mihara,

Etsuro Mori, and Hideki Mochizuki. "Ocular Fixations and Presaccadic Potentials to

Explain Pareidolias in Parkinson's Disease." *Brain Communications* 2, no. 1 (January

2020): 1–15. https://doi.org/10.1093/braincomms/fcaa073.

Richards, John E. "Cortical Sources of ERP in Prosaccade and Antisaccade Eye Movements

Using Realistic Source Models." *Frontiers in Systems Neuroscience* 7 (2013): 27.

https://doi.org/10.3389/fnsys.2013.00027.

Richards, John E. "Cortical Sources of Event-related Potentials in the Prosaccade and

Antisaccade Task." *Psychophysiology* 40, no. 6 (November 2003): 878–94.

https://doi.org/10.1111/1469-8986.00106.

Rizzolatti, Giacomo, Luciano Fadiga, Vittorio Gallese, and Leonardo Fogassi. "Premotor

835   Cortex and the Recognition of Motor Actions." *Cognitive Brain Research* 3, no. 2

836   (March 1996): 131–41. https://doi.org/10.1016/0926-6410(95)00038-0.

837 Rizzolatti, Giacomo, Lucia Riggio, Isabella Dascola, and Carlo Umiltá. "Reorienting Attention

838   across the Horizontal and Vertical Meridians: Evidence in Favor of a Premotor Theory

839   of Attention." *Neuropsychologia* 25, no. 1 (January 1987): 31–40.

840   https://doi.org/10.1016/0028-3932(87)90041-8.

841 Ruíz, Josafhat Salinas, Osval Antonio Montesinos López, Gabriela Hernández Ramírez, and

842   Jose Crossa Hiriart. "Elements of Generalized Linear Mixed Models." In *Generalized*

843   *Linear Mixed Models with Applications in Agriculture and Biology*, 22–35. Springer

844   International Publishing, 2023. https://doi.org/10.1007/978-3-031-32800-8.

845 Schielzeth, Holger, Niels J. Dingemanse, Shinichi Nakagawa, David F. Westneat, Hassen

846   Allegue, Céline Teplitsky, Denis Réale, Ned A. Dochtermann, László Zsolt Garamszegi,

847   and Yimen G. Araya-Ajoy. "Robustness of Linear Mixed-effects Models to Violations of

848   Distributional Assumptions." *Methods in Ecology and Evolution* 11, no. 9 (September

849   2020): 1141–52. https://doi.org/10.1111/2041-210X.13434.

850 Schluppeck, Denis, Paul Glimcher, and David J. Heeger. "Topographic Organization for

851   Delayed Saccades in Human Posterior Parietal Cortex." *Journal of Neurophysiology* 94,

852   no. 2 (August 2005): 1372–84. https://doi.org/10.1152/jn.01290.2004.

853 Scholz, Agnes, Anja Klichowicz, and Josef F. Krems. "Covert Shifts of Attention Can Account

854   for the Functional Role of 'Eye Movements to Nothing.'" *Memory & Cognition* 46, no. 2

855   (February 2018): 230–43. https://doi.org/10.3758/s13421-017-0760-x.

856 Servais, Anaïs, Christophe Hurter, and Emmanuel J. Barbeau. "Attentional Switch from the

857   Outside to the Inner World: An Early and Critical, but Unexplored Stage of

858   Autobiographical Memory Retrieval Processes." *Brain and Cognition* 170 (August

859   2023): 106031. https://doi.org/10.1016/j.bandc.2023.106031.

860 Sipatchin, Alexandra, Siegfried Wahl, and Katharina Rifai. "Eye-Tracking for Clinical

861   Ophthalmology with Virtual Reality (VR): A Case Study of the HTC Vive Pro Eye's

862   Usability." *Healthcare* 9, no. 2 (February 2021): 180.

863   https://doi.org/10.3390/healthcare9020180.

Smith, Daniel T., and Thomas Schenk. "The Premotor Theory of Attention: Time to Move On?" *Neuropsychologia* 50, no. 6 (May 2012): 1104–14. https://doi.org/10.1016/j.neuropsychologia.2012.01.025.

Strube, Andreas, Michael Rose, Sepideh Fazeli, and Christian Büchel. "Alpha-to-Beta- and Gamma-Band Activity Reflect Predictive Coding in Affective Visual Processing." *Scientific Reports* 11, no. 1 (December 2021): 23492. https://doi.org/10.1038/s41598-021-02939-z.

Svantesson, Mats, Håkan Olausson, Anders Eklund, and Magnus Thordstein. "Virtual EEG-Electrodes: Convolutional Neural Networks as a Method for Upsampling or Restoring Channels." *Journal of Neuroscience Methods* 355 (May 2021): 109126. https://doi.org/10.1016/j.jneumeth.2021.109126.

J. Tauscher et al., "Immersive EEG: Evaluating Electroencephalography in Virtual Reality," 2019 IEEE Conference on Virtual Reality and 3D User Interfaces (VR), Osaka, Japan, 2019, pp. 1794-1800;

Team, JASP. "JASP (v0.18.3)," 2024.

Thura, David, and Paul Cisek. "Deliberation and Commitment in the Premotor and Primary Motor Cortex during Dynamic Decision Making." *Neuron* 81, no. 6 (March 2014): 1401–16. https://doi.org/10.1016/j.neuron.2014.01.031.

Tolman, Edward C. "Cognitive Maps in Rats and Men." *Psychological Review* 55, no. 4 (1948): 189–208. https://doi.org/10.1037/h0061626.

Wechsler, David. *Wechsler Adult Intelligence Scale*. 3rd ed. The Psychological Corporation, 1997.

Weiner, Susan L., and Howard Ehrlichman. "Ocular Motility and Cognitive Process." *Cognition* 4, no. 1 (January 1976): 31–43. https://doi.org/10.1016/0010-0277(76)90009-3.

Westendorff, Stephanie, Christian Klaes, and Alexander Gail. "The Cortical Timeline for Deciding on Reach Motor Goals." *The Journal of Neuroscience* 30, no. 15 (April 2010): 5426–36. https://doi.org/10.1523/JNEUROSCI.4628-09.2010.

Whitlock, Jonathan R. "Posterior Parietal Cortex." *Current Biology* 27, no. 14 (July 2017): R691–95. https://doi.org/10.1016/j.cub.2017.06.007.

Whitlock, Jonathan R., Robert J. Sutherland, Menno P. Witter, May-Britt Moser, and Edvard

I. Moser. "Navigating from Hippocampus to Parietal Cortex." *Proceedings of the*

National Academy of Sciences 105, no. 39 (September 2008): 14755–62.

https://doi.org/10.1073/pnas.0804216105.

Xia, Yinglin, and Jun Sun. "Linear Mixed-Effects Models for Longitudinal Microbiome Data."

In *Bioinformatic and Statistical Analysis of Microbiome Data*. Springer International

Publishing, 2023. https://doi.org/10.1007/978-3-031-21391-5.

Yeung, Lok-Kin, Rosanna K. Olsen, Hannah E. P. Bild-Enkin, Maria C. D'Angelo, Arber

Kacollja, Douglas A. McQuiggan, Anna Keshabyan, Jennifer D. Ryan, and Morgan D.

Barense. "Anterolateral Entorhinal Cortex Volume Predicted by Altered Intra-Item

Configural Processing." *The Journal of Neuroscience* 37, no. 22 (May 2017): 5527–38.

https://doi.org/10.1523/JNEUROSCI.3664-16.2017.

Acknowledgments: The authors would like to acknowledge support from Bial Foundation (grant

260/22) and Portuguese Foundation for Science and Technology (2023.08332.CEECIND) to AP,

and Deutsche Forschungsgemeinschaft grant (122679504—SFB 874) to CK. We want to thank

Alexandra Sipatchin and Fernanda Ponce for their help with eye tracking data analysis, Diogo

Branco for his help with the task and the setup, Soraia Oliveira for help with data collection and

Marcin Leszczynski, Susanne Dyck, Jon Walbrin, Fredrik Bergstroem, Elizabetta Del Re and Pieter

Medendorp for helpful input and discussions.

Author Contributions: AP conceptualized the experiment. AP, MJM, EA and GB designed the

experiment and collected the data. LW, GB, AP and MJM analyzed the data. AP, LW, GB and MJM

wrote the manuscript. All authors revised the manuscript.

Competing interests statement: The authors declare no competing interests.

Data availability statement: Data underlying all figures is available for download at:

https://osf.io/8q4wh/?view_only=730dcb4c2cab415694c3f7989241f31a

List of Supplementary Materials:

1) Supplementary Methods

2) Supplementary Results