Multi-omic landscape of human gliomas from diagnosis to treatment and recurrence

Hadeesha Piyadasa*[,1], Benjamin Oberlton*[,1,2], Mikaela Ribi[1,3], Jolene S. Ranek[1], Inna Averbukh[1], Ke Leow[1,4], Meelad Amouzgar[1,2], Candace C. Liu[1,2], Noah F. Greenwald[1,4], Erin F. McCaffrey[1], Rashmi Kumar[1], Selena Ferrian[1], Albert G. Tsai[1], Ferda Filiz[1], Christine Camacho Fullaway[1], Marc Bosse[1], Sricharan Reddy Varra[1], Alex Kong[1], Cameron Sowers[1], Melanie Hayden Gephart[5], Pablo Nuñez-Perez[5], EnJun Yang[6], Mike Travers[6], Michael J. Schachter[6], Samantha Liang[6], Maria R. Santi[8], Samantha Bucktrout[6], Pier Federico Gherardini[6,7], John Connolly[6], Kristina Cole[8], Michael E. Barish[6,9], Christine E. Brown[10], Derek A. Oldridge[6,11,12], Richard R. Drake[13], Joanna J. Phillips[14,15], Hideho Okada[6,15], Robert Prins[6, 16], Sean C. Bendall[1, 2, 4, 6,#], Michael Angelo[1, 2, 4, 6#]

1. Department of Pathology, Stanford University School of Medicine, Stanford, CA, USA
2. Immunology Program, Stanford University School of Medicine, Stanford, CA, USA
3. Department of Chemistry and Sarafan ChEM-H, Stanford University, Stanford, CA, USA
4. Cancer Biology Program, Stanford University School of Medicine, Stanford, CA, USA
5. Department of Neurosurgery, Stanford University School of Medicine, Stanford, CA, USA
6. Parker Institute for Cancer Immunotherapy, San Francisco, CA, USA
7. Department of Biology, University of Rome "Tor Vergata", Rome, Italy
8. Children's Hospital of Philadelphia, Perelman School of Medicine, University of Pennsylvania, PA, USA
9. Department of Stem Cell Biology and Regenerative Medicine, Beckman Research Institute of the City of Hope, Duarte, CA, USA
10. Departments of Hematology & Hematopoietic Cell Transplantation and Immuno-Oncology, Beckman Research Institute of the City of Hope, Duarte, CA, USA
11. Department of Pathology and Laboratory Medicine, Perelman School of Medicine at the University of Pennsylvania, Philadelphia, PA, USA
12. Center for Computational and Genomic Medicine, Children's Hospital of Philadelphia, PA, USA
13. Department of Pharmacology and Immunology, Medical University of South Carolina, Charleston, SC, USA
14. Department of Pathology, University of California San Francisco, San Francisco, CA, USA
15. Department of Neurological Surgery, University of California San Francisco, San Francisco, CA, USA
16. Department of Neurosurgery, UCLA, Los Angeles, CA, USA

* Equal first authors
# Correspondence: bendall@stanford.edu & mangelo0@stanford.edu

Abstract

Gliomas are among the most lethal cancers, with limited treatment options. To uncover hallmarks of therapeutic escape and tumor microenvironment (TME) evolution, we applied spatial proteomics, transcriptomics, and glycomics to 670 lesions from 310 adult and pediatric patients. Single-cell analysis shows high B7H3+ tumor cell prevalence in glioblastoma (GBM) and pleomorphic xanthoastrocytoma (PXA), while most gliomas, including pediatric cases, express targetable tumor antigens in less than 50% of tumor cells, potentially explaining trial failures. Longitudinal samples of isocitrate dehydrogenase (IDH)-mutant gliomas reveal recurrence driven by tumor-immune spatial reorganization, shifting from T-cell and vasculature-associated myeloid cell-enriched niches to microglia and CD206+ macrophage-dominated tumors. Multi-omic integration identified N-glycosylation as the best classifier of grade, while the immune transcriptome best predicted GBM survival. Provided as a community resource, this study opens new avenues for glioma targeting, classification, outcome prediction, and a baseline of TME composition across all stages.

Introduction

Diffuse glioma are primary brain tumors that are among the most lethal forms of cancer, with limited treatment options and poor prognosis[1,2]. The average survival of patients with glioblastoma (GBM) is only 9–18 months, while diffuse midline glioma (DMG) is the leading cause of brain tumor-related deaths in children[3]. Despite advances in immunotherapies that have shown promise in other cancers, outcomes for gliomas have remained stagnant for decades, and the mechanisms underlying this therapeutic resistance remain elusive[4].

Due to their location and invasive growth into the normal brain, complete surgical resection of diffuse gliomas is often not possible making them prone to recurrence[5]. Additionally, the glioma tumor microenvironment (TME) is myeloid rich and highly immunosuppressive, rendering them resistant to treatment[6–8]. Clinical therapies currently in trials are attempting to replicate previous successes in other malignancies that use antibodies or engineered T-cells to target tumor-enriched antigens[9]. Indeed, tumor antigen (TA) expression, such as B7H3 and EGFR, represents a promising avenue for directly targeting tumor cells through CAR-T cell therapies and monoclonal antibodies[10–12]. Still, these therapies have either been ineffective or have led to only modest improvements in patient survival[13]. Perhaps related to these setbacks, the expression patterns of these TAs in human lesions are largely unknown[14,15]. Much of what is known about TAs and glioma TME has been gleaned from bulk transcriptomic analysis, patient-derived xenograft mouse models, and tumor cell lines[16–20]. Taken together, these findings point to a significant gap in our understanding of how TAs, immune-tumor interactions, and the functional states induced by them mutually reinforce one another to maintain a treatment resistant milieu. Additionally, though post-translational molecular features, such as tumor glycosylation, have been shown in other malignancies to play a crucial role in promoting tumor-immune-suppression, the glycome in human gliomas remains largely unexplored[21–23].

To address these questions, we assembled a clinically annotated human glioma tissue cohort from multiple sites across the US. This cohort includes 677 samples from 310 adult and pediatric patients that are classified as high or low-grade tumors, and encompass treatment-naive, post-immunotherapy (anti-PD1, vaccine, anti-CD27, combinatorial), recurrent, and longitudinal cases, making it one of the largest and diverse glioma cohorts assembled[24–27]. To capture multiple layers of molecular regulation, we employed complementary spatial profiling technologies to quantify biomolecule abundance, cellular states and identities (i.e., transcriptome, proteome, and glycome). Multiplexed Ion Beam Imaging (MIBI-TOF) provided high-resolution spatial mapping of 1.2 million cells consisting of neurons, endothelial, tumor and all immune cells (with 18 annotated sub-populations), quantifying targetable tumor antigen expression (i.e., B7H3, EGFR) and immune regulatory receptors (i.e. PD-L1, TIM3) at the single-cell level[28]. Co-registered tissue sections linked these single-cell maps to spatially resolved transcriptomic profiles of over 11,000 genes from immune-cell-rich versus -poor regions using digital spatial profiling (DSP)[29]. Lastly, co-registered tissue sections were also combined with spatial maps of the *N*-glycome quantified by matrix-assisted laser desorption-ionization mass spectrometry imaging (MALDI-MSI) which discovered 70 differentially expressed glycans across gliomas that that could influence immune recognition and tumor behavior[30–32]. Combining these modalities, we established direct correlations between cell composition, protein expression, gene expression, and glycan profiles

within the same spatial context across our diverse glioma cohort and how it relates to clinical metadata.

Multi-modal data analysis across these lesions provided new insights into how the glioma TME is shaped by tumor classification and treatment, revealing potential explanations for the limited efficacy of TA-targeted immunotherapies. Furthermore, both tracing the evolving structure of gliomas from primary to recurrent disease and aligning molecular data with patient metrics uncovered factors linked to overall survival[33]. Towards this, we conducted a spatial proteomics-based, quantitative assessment of glioma TAs, marking the first time the prevalence of these antigens and their co-occurrence within single cells has been systematically profiled. Our results indicate that, aside from the high abundance of B7H3+ tumor cells in GBM and pleomorphic xanthoastrocytoma (PXA), most tumors only show less than 50% TA-positive cells. Yet, co-expression analyses suggest that dual-agent strategies could meaningfully increase the fraction of tumor cells potentially targeted. Consistent with the modest clinical improvements seen to date, comparing lesions from patients who received immunotherapy versus standard of care revealed only minor differences tumor architecture[2]. To better understand the apparent stalling of anti-tumor immune activation, we next characterized the glioma glycome, identifying sialylated and core-fucosylated N-glycans whose shifts in abundance correlated strongly with tumor grade, representing both a new classification and therapeutically targetable entity. Spatial transcriptomics further reinforced molecular pathways influencing glycosylation in glioma biology[6]. Still, examining patient outcomes in primary GBM, we found that immune cell, as opposed to tumor cell, programs played the most prominent role in predicting overall survival. All the underlying data from this diverse glial tumor cohort is fully accessible, without restriction, via an interactive online portal (www.bruce.parkerici.org), providing a multi-omic resource to the community that should prove invaluable for future clinical design, beyond the insights provided herein.

Results

Diverse and clinically detailed glioma cohort and applied spatial multi-omics

The goal of this study was to create a more comprehensive understanding of the glioma TME and how the multitude of regulatory features at play may integrate to reconcile disease diversity, including the prevalence and amounts of targetable TA. To achieve this, we collaborated with five clinical sites across the US to build a multi-omic atlas of the glioma TME, assembling a cohort of glioma patients with detailed clinical annotations, covering a wide age range from under 1 year to 88 years (**Figure 1A, Figure S1A, Table S1**). The cohort spans World Health Organization (WHO) grades 2, 3, and 4, and includes patients diagnosed with GBM, IDH-mutant astrocytoma, IDH-mutant, oligodendroglioma, IDH-mutant and 1p/19q-codeleted PXA, DMG and other pediatric high-grade gliomas (pHGG). We also included two cohorts of recurrent GBM patients treated with neoadjuvant anti-PD1 immune checkpoint blockade, as well as longitudinal samples from patients with primary and recurrent IDH mutant astrocytoma, oligodendroglioma, and PXA, some of whom received neoadjuvant immunotherapy before surgical resection (**Figure 1B, Figure S1A, Table S1**). Clinical data, including mutational profiles, such as IDH status (Isocitrate Dehydrogenase), MGMT methylation (O6-Methylguanine-DNA Methyltransferase), PTEN (Phosphatase and Tensin Homolog), p53 (Tumor Protein 53), and ATRX status (Alpha Thalassemia/Mental Retardation Syndrome X-linked) along with overall survival, corticosteroid

Figure 1. BRUCE (BRain tUmor heterogeneity deCiphEred by high dimensional multi-omic analysis) cohort and data overview. (A) BRUCE cohort summary (310 patients, 677 samples), including patient demographics, **(B)** sample distribution and **(C)** the spatial modalities (transcriptomics, proteomics, N-glycome) used for data acquisition from FFPE sections **(D)** Illustration of web resource.

use, and other key attributes, were harmonized across all centers to create a unified, clinically annotated dataset (**Figures 1A and 1B, Figure S1A, Table S1**). This integrated resource enabled us to perform in-depth analyses and explore the molecular intricacies within the TME.

Multiple regions for each lesion were analyzed using single cell spatial proteomics, spatial transcriptomics and spatial N-glycomics using adjacent serial sections to map cellular phenotypes, whole transcriptome in immune-rich versus -poor regions, and N-glycan abundances and distribution (**Figure 1C, Figure S1A**). In total, we imaged over 1.2 million cells from 677 patient samples quantifying the expression and cellular distribution of 52 proteins. The functional status

of these cells was characterized using 14 immune regulatory markers and 8 TA across more than
1,000 sampled tumor regions. Our whole transcriptome analysis revealed over 11,000 genes
enriched in immune cell-rich and -poor regions, while N-glycomics uncovered 70 unique N-
glycans across 10 broad classes, including sialylated structures implicated in immune suppression
within the TME of other cancers[34–36].

We have made all data from this study available on an intuitive, open-access portal,
www.bruce.parkerici.org, allowing researchers and clinicians to explore complex spatial and
multi-omic data alongside clinical metadata, providing meaningful insights that may guide future
research and clinical decisions (**Figure 1D**). The platform, accessible even to non-computational
experts, includes interactive tools such as Vitessce for single-cell spatial MIBI image
exploration[37]. Additionally, all raw data, including single-cell tables, glycan expression data,
spatial glycan TIFF files, and raw MIBI images, are freely available for download without
restriction.

A spatial atlas of glioma: Single-cell profiling

Understanding the composition, distribution, and antigen expression of immune and tumor cells
in human gliomas is crucial for understanding drivers of therapeutic failure and for developing
new treatments[20,27,38,39]. With this in mind, we used spatial proteomics to provide a basic level of
identity to all cells within the TME, including the quantification of tumor cells and 18 distinct
immune cell subsets (**Figures 2A and 2C, Figures S2A – S2N, Tables S2 and S3**). Tumor cells
were labeled based on histology by expert pathologists and multiplexed phenotype. For the latter,
cells were labeled tumor if they were positive for any combination of Olig2, GFAP, CD133, B7H3,
EGFR, GM2/GD2, HER2, VISTA, GPC2, NG2, EGFRvIII, H3K27M, or IDH R132H while being
negative for immune, endothelial, or neuronal markers (**Figures S2M**). This approach was taken
because glioma tumor cells are nondescript and difficult to identify reliably based on a single
protein[40]. However, tumors with K27M mutation of the histone 3 isoforms (H3K27M) or R132H
mutation of IDH are an exception because mutation-specific antibodies can be used to specifically
identify tumor cells[41,42]. We used this subset of lesions to assess the accuracy of multiplexed tumor
phenotyping, demonstrating strong concordance (**Figures S3A – S3D**). The immune compartment
(18% of total cells) was dominated by multiple subsets of microglia, macrophages, dendritic cells
(DCs), and monocytes that were delineated by coexpression of CD11b, HLA-DR, CD14, CD68,
CD163, CD206, CD209, TMEM119 and CD141 (**Figure 2B, Figures S2M and S2N**). Lymphoid
cells were sparse (<1% of total cells) consisting of mostly CD8+ and CD4+ T cells (**Figure 2B,
Figures S2M and S2N**).

Because this cohort spans across glioma type and grade, we were able to identify how cellular
composition varies with respect to these clinical covariates (**Figures 2D and 2E, Figure S3E**).
For instance, CD163 expression, linked to macrophage wound healing and anti-inflammatory
behavior increases with WHO grade and is negatively correlated with overall survival in GBM[43–
45]. However, it remains unclear which specific myeloid subsets in each glioma type express
CD163[46]. In our cohort, we identified three major myeloid subsets expressing CD163: CD14+,
CD68+, and microglia (**Figure 2D, orange arrow**). While CD14+ and CD68+ subsets were found
to increase with grade, CD163+ microglia decreased with grade (**Figure 2D, orange arrow,
Figures S2F – S2H**), with grade 2 and 3 astrocytoma having the highest relative abundance (4.5%

245
246
247 **Figure 2. Immune cell and targetable tumor antigen (TA) characterization across glioma subtypes using**
248 **spatial proteomics. (A)** Antibody marker panel used to identify various cell types including tumor, immune,

neuronal, endothelial, and regulatory cells, as well as segmentation markers. **(B)** Cellular composition of the tumor microenvironment, highlighting targetable antigen positive tumor cells and immune cells, with a detailed breakdown of myeloid and lymphoid subpopulations. **(C)** Representative MIBI images highlighting various markers. **(D)** Heatmap displaying immune cell populations across different glioma subtypes. Color represents row wise Z-score average abundance and circle size represents Log10 average abundance (relative to all immune cells). **(E)** Representative spatial maps showing immune and tumor cell distribution in glioma subtypes, including GBM, PXA, oligodendroglioma, and astrocytoma. **(F)** Heatmap displaying TA across glioma subtypes. Color represents row wise Z-score intensity and circle size represents Log10 average abundance (relative to all immune cells)

and 7.3% respectively) (**Figure 2D**). Within GBM (grade 4), the proportion of CD14+ and CD68+ subsets expressing CD163 was remarkably high (15% and 29%) (**Figure 2D**). Interestingly, IDH1-mutant grade 4 astrocytoma exhibited a lower abundance of CD163+ myeloid populations compared to GBM, despite containing relatively high levels of total microglia, CD14+, and CD68+ myeloid cells (**Figure 2D, green arrow**). These findings highlight the nuanced CD163+ immune subsets across gliomas and grades and offer new insights into their potential roles in the TME.

These differential immune compositions measured also extend to rare gliomas, such as PXA, an astrocytic tumor affecting children and young adults accounting for less than 1% of gliomas. Due to the rarity of PXAs, our understanding of their TME remains limited[47]. To address this, we characterized the immune cell composition and TA expression across 15 pediatric (Pediatric HGG [other]), 13 grade 2, and 37 grade 3 PXA samples (**Figure 2D and 2E**), which revealed two notable insights. First, PXAs had the lowest relative abundance of myeloid cells phenotypically consistent with MDSCs (CD11b+, HLA-DR-) (**Figure 2D, purple arrow**). Second, Grade 2 PXAs had the highest abundance of T cells across all gliomas. Together, these findings highlight how the immune landscape can remodel with glioma type and grade, implicating the likely need for more nuanced immune targeting strategies instead of a one-size-fits all in glioma.

Quantifying tumor antigen variability: Implications for targeted therapy in gliomas

Beyond immune cell modulation, targeting of TAs remains a significant strategy for gliomas, in general. Still, the efficacy of TA-targeting therapies is dependent, in large part, on abundant and high frequency expression across tumor cells[48]. However, quantitative profiles describing how these antigens are co-expressed at single cell level with respect to tumor structure, type, and grade in human gliomas are not known. To this end, we quantified the expression of eight different targetable TA (B7H3, EGFR, EGFRvIII, HER2, VISTA, GPC2, NG2, GM2/GD2) that are currently in pre-clinical or clinical stage of investigation in glial tumors[49–56]. EGFRvIII was excluded from analysis due to the limited number of confirmed patients in our cohort with this variant. Because lipids like GD2 are typically lost during processing of archival tissues, we used GD2 synthase (GM2/GD2) as a surrogate for this TA. Of the TAs measured here, B7H3 showed the highest abundance across all tumor types and grades (**Figure 2F**). However, B7H3 was not restricted to tumor cells, with >60% immune and endothelial cells expressing this as well (**Figure S3I**). EGFR was also prominent but only in adult tumors (**Figure 2F**). We further investigated the correlation between the expression of these antigens and their transcripts using spatial transcriptomic data from serial samples. B7H3 and EGFR proteins showed significant correlation with their respective transcripts, while other targetable tumor antigens showed no significant correlations (**Figure S3J**).

Figure 3. Comprehensive analysis of tumor antigen diversity, coverage, and coexpression in gliomas. (A)
Boxplot of Shannon diversity index showing TA heterogeneity across different tumor subtypes. **(b-c)** Line graph
showing % tumor cell coverage with increasing TA. Error bars show SD. **(D)** Top two tumor antigen expression
heatmap shows % patients for combinations of selected top 2 TA yielding maximal coverage of tumor cells for high
TA diversity glioma. **(E)** Venn diagrams showing % of tumor cells positive and overlap for B7H3, EGFR and HER2
out of total tumor cells in high TA diversity gliomas. **(F)** Representative tumor CPM images, illustrating the
distribution of tumor antigens for coverage and coexpression. **(G)** Violin plot of % tumor cells negative for the 7 TA
measured. Each dot represents an individual patient.

We next examined *in silico* how tumor targeting might be improved using multi-agent therapy[57,58].
In high-diversity tumors (GBM, astrocytoma, oligodendroglioma) that more frequently express
multiple TAs (**Figure 3A**), our analysis suggests that dual therapy could lead to a significant
improvement in tumor coverage–defined here as the percentage of tumor cells expressing at least
one of the TAs of interest (**Figure 3B**). However, this benefit quickly plateaus and does not
improve further with three or more agents. In contrast, because low-diversity tumors (**Figure 3A**)
mostly expressed B7H3, little improvement in tumor coverage was seen by targeting multiple TAs
(**Figure 3C**). Given that dual-TA coverage was most effective in high-diversity tumors, we
assessed TA pairings to determine the optimal combination for each patient (**Figure 3D, Figure
S3K**). Dual therapy with B7H3 and EGFR was predicted to be optimal, covering up to 100% of
tumor cells, for 47% of GBM and 44% of oligodendroglioma cases. In astrocytoma, no dominant
pair emerged, though 22% of patients had either B7H3 with EGFR or HER2 (**Figure 3D**).

Convoluting this, antigen expression by non-tumor cells, as we saw with B7H3, can lead to off-
target effects and adverse events when using monospecific therapies alone or with multi-agent
regimens like the ones described above[59]. To circumvent these issues, bispecific antibodies and
CAR-T cells that require co-expression of two antigens on the same cell are being examined as a
means of improving tumor specificity[57,58]. With this in mind, we quantified how often B7H3,
EGFR, and HER2 were co-expressed on the same cell in high-diversity tumors to understand how
tumor coverage with a bispecific therapy compared with monospecific dual therapy (**Figures 3E
and 3F**). Overall, we found the added specificity of a bispecific therapy is predicted to come at
the cost of reduced tumor coverage. For example, in GBM, 85% of tumor cells express either
B7H3 or EGFR, but only 32% expressed both. Lastly, we would like to note that nearly all lesions

331   contained tumor cells that were pan-negative for all TAs quantified here, with their prevalence
332   varying widely from 0-99% (**Figure 3G**). Taken together, these results provide new guidance on
333   how therapeutic regimens might be improved while also highlighting the challenges in designing
334   broadly reactive, specific therapies in glioma.
335
336   **Benchmarking tumor antigen expression dynamics for targeted therapy in gliomas**
337
Besides frequency, the ideal therapeutic target would be consistently expressed by all tumor cells,
across patients, minimizing cell-to-cell variability within each patient's lesion[57–59]. Given the
prevalence of pan-negative tumor cells (**Figure 3G**), we next examined this more broadly to
understand how the frequency of TA-positive cells could impact the potential efficacy of tumor
cell targeting therapies. To contextualize these findings, we first compared our data with a tumor-
targeted therapy that has shown durable survival benefits, CD19 CAR-T therapy in B-cell
malignancies. Previous studies have shown a cohort median of 98% antigen-positive tumor cells
with individual patients ranging from 72%-100%[60]. Since B7H3 and EGFR emerged as the top
TAs for GBM, oligodendroglioma, and astrocytoma, we compared the percentage of tumor-
positive cells for these antigens with the rate of CD19 positivity in B-cell malignancies. This
analysis revealed two insights. First, of the three tumor types, GBM had the highest fraction of
patients falling in this range for both B7H3 and EGFR (62% and 30%, **Figures 4A and B**,
respectively). Second, and perhaps encouragingly, TA-positive cells for lesions falling within this
range also expressed on average higher amounts of antigen than those outside of it (B7H3 = 1.6X
higher, *p = 0.001* and EGFR = 4.2X higher, *p = 0.007*). However, cell-to-cell variability in TA
expression (i.e., intra-patient variation) was also greatest in these lesions. Given these findings, we
then examined how TA expression varied between patients. We quantified the dynamic range of
TA expression (5th-95th percentile) at a single-cell level across all tumor cells for all patients
(**Figures 4C and 4D**). Similar to our findings when examining variation within a single lesion, we
found EGFR and B7H3 to exhibit the largest interpatient variation as well, spanning a dynamic
range of 3.49 and 2.59 Log2 FC from median, respectively.
359
TA expression is often not spatially uniform throughout and can be biased to the central tumor
core or peripheral edge near the resection margin where residual disease is likely to occur[19,61].
With this in mind, we further examined regional variability of TA expression in WHO grade 4
tumors, focusing on the gradient from core to infiltrating edge (**Figures S4A – S4H)**. Here, we
found that two of the TAs exhibited non-uniformity with respect to these clinically significant
histologic regions. GM2/GD2 tended to be higher at the resection margin while B7H3 was
typically higher in the tumor core (**Figures S4A – S4H**). Taken together, these findings underscore
the complexity of developing effective therapies for gliomas, as TA variability, both within and
between tumors, complicates consistent targeting, even for abundant antigens like B7H3 and
EGFR.
370
371   **Clinically significant changes in disease status are spatially encoded in the glioma TME**
372
373   In previous work examining carcinomas, we found clinically significant events like disease
374   recurrence and metastasis to be tightly linked to corresponding changes in TME structure that
375   revealed insights into drivers of tumor progression and therapeutic resistance[8,62–65]. To determine
376   if TME structure and disease progression are similarly linked in gliomas, we compared

Figure 4. Dynamic range of expression of targetable tumor antigens (TA) in gliomas. Scatter plot for (A) B7H3 and (B) EGFR showing mean intensity on Y axis, and % positive tumor cells on the x axis. Size of each dot indicates the intrapatient dynamic range of expression. Each dot represents an individual patient. Colors identify tumor type. Shaded region shows theoretical mean intensity and % positive tumor cells of CD19 in B cell malignancies. (C) Point plot showing the log2 fold change from the median intensity. 10,000 cells per TA sampled. Color scale represents Log2 fold change capped at -1.5 and 1.5 (D) Representative MIBI CPM of tumor antigen expression. Color scale shows the log2 fold change from median shown in panel a. Scale bar = 100 μm.

longitudinally matched primary and recurrent lesions from patients with LGG (Astrocytoma IDH-mutant, Oligodendroglioma IDH-mutant 1p19q co-deletion and PXA) that received standard of care (SOC) using QUantitative InterCellular nicHe Enrichment (QUICHE) (**Figures 5A and 5B, green line**)[63]. QUICHE combines graph neighborhood analysis with statistical modeling to identify cellular niches that are differentially enriched in one condition over another. This analysis revealed systematic shifts in how tumor and immune cells colocalize in recurrent LGG disease (**Figures 5C and 5D, Figure S5A**). For example, in primary tumors, there were complex interactions between CD4+ and CD8+ T cells with CD68+CD163+ macrophages, CD14+ myeloid cells, tumor cells and endothelial cells (indicated by interconnected lines while thickness of lines indicate the prevenance across patients) which were absent in recurrent tumors. Instead, microglia with or without CD163, CD206+ macrophages and CD11+HLADR- myeloid cells formed distinct spatial niches with tumor cells in recurrent tumors.

These changes suggest that tumor cells in primary tumors interact more with vasculature and immature myeloid cells in comparison to recurrent tumors where tumor cells interact more with brain resident microglia and more effector myeloid populations[66]. Notably, while the TME

Figure 5: Analysis of distinct spatial niches and therapy-induced changes within the TME of longitudinally sampled low-grade gliomas. (A) Schematic overview of the QUICHE analysis pipeline. **(B)** Overview of the longitudinal LGG cohort with paired primary and recurrent samples, including a subset of patients treated with immunotherapy. Spatial interaction graphs derived from differentially abundant niches for **(C)** primary and **(D)** recurrent tumors. Edge width and color corresponds to the number of unique patients with the interaction. **(E)** Differential abundance analysis of immune cell types between recurrent and primary tumors in the SOC cohort, showing log2 fold changes and statistical significance (circle size indicates -log10 p-value). **(F)** Volcano plot comparing log2 fold changes of immune cells with functional marker classification between SOC and immunotherapy-treated patients, highlighting (red) significant differences. **(G)** Representative images of multiplexed single-cell spatial TME landscapes, showing functional marker co-expression within immune cell subsets.

structure differed significantly in primary and recurrent lesions, the prevalence of tumor and immune cell subsets composing the TME did not (**Figure 5E,** p > 0.05 for all populations). We next evaluated whether the cells within these differential niches were phenotypically distinct from cells outside of them to understand how TME structure and function are related. TIM3, IDO1, and CD86-positive cell fractions were notably enriched in DE niche neighborhoods (**Figures S5B –**

S5D). For example, TIM3+ CD4 and CD8 T cells were overrepresented in primary tumor niches
but were entirely absent in recurrent tumor niches. Collectively, these findings demonstrate that
although the overall cellular makeup remains constant, glioma recurrence is encoded by the spatial
a reorganization of tumor–immune cell interactions.

In addition to patients receiving SOC, our LGG cohort included recurrent samples from phase 1
clinical trials in which patients with Astrocytoma IDH-mutant or Oligodendroglioma IDH-mutant
1p19q co-deletion were treated with either a vaccine alone or a vaccine in combination with an
anti-CD27 agonistic antibody[25,26]. Although small sample sizes and limited follow-up precluded a
definitive assessment of clinical benefit, immunotherapies in glioma generally have not improved
clinical outcome. Thus, we hypothesized that comparison between patients receiving
immunotherapy and SOC would not reveal significant spatial differences. In line with this,
QUICHE analysis did not reveal any treatment-specific niches. However, bulk frequencies of
IDO1+, CD86+, and TIM3+ myeloid cells was higher in SOC-treated patients (**Figures 5F and
5G, Figure S5E**). Taken together, these findings suggest that TME niche enrichment is linked to
glioma clinical status through a sensitive and specific encoding that cannot be reliably inferred in
a spatially agnostic manner based on cell composition alone.

Multi-omic profiling of the TME reveals unique glycan profiles across glioma grades and
novel cellular and gene module associations.

N-glycosylation is a post-translational modification occurring on most membrane-bound proteins
that regulates a broad range of cell-cell interactions involved in cancer progression and immune
regulation[21–23]. For example, tumor sialylated *N*-glycans inhibit innate and adaptive responses
through binding of Siglec receptors on immune cells[34–36,67]. With this in mind, we used MALDI-
IMS (Bruker TimsTOF MALDI2) to map the spatial composition of glioma TME glycans to
determine which structural classes are present and how they change with grade[32,68]. Searching a
possible library of N-glycans between 800 and 4000 m/z, we identified 70 unique *N*-glycans across
10 different structural classes (each unique glycan may belong to more than a single structural
class) in WHO grade 2-4 tumor samples (**Figure 6A**, **Figure S6A**). The abundance of these glycan
classes was grade-specific[69]. For example, grade 2 lesions where enriched in high mannose,
fucosylated, hybrid, agalactosylated, and tetrantennary structures while grade 4 gliomas had higher
levels of sialylated, polylacnac, bi- and triantennary glycans (**Figure 6B and 6C, Figure S6A**).
Notably, we saw that IDH mutant grade 4 tumors were distinctly different from IDH mutant grade
2 and 3 tumors showing lower fucosylation and higher sialylation similar to GBM (**Figure S6B**)
indicating that the glycocalyx may be independent of IDH status.

To understand how the glycocalyx of gliomas could be remodeled with grade, we then integrated
differential N-glycan compositions with tissue whole transcriptome profiling (Nanostring DSP
GeoMx) from matching regions from adjacent serial sections. Here we identified glyco-regulatory
enzymes whose expression correlated with these trends across grade (**Figure 6D**). Of the ten
glycan classes analyzed, fucosylated and sialylated glycans exhibited the greatest concordance
with the transcript levels of the enzymes predicted to synthesize them (**Figure 6D**, *red triangle*
and *purple diamond*, respectively; **Figures S6C – S6D**). Interestingly however, robust trends with
gene expression matching N-glycan modifications, like this, were not found for other glycan
classes despite19 genes captures by whole transcriptome profiling. This discordance suggests that

468
Figure 6. Multi-omic profiling of the TME. (A) Bar graph % of glycans identified in each N-glycan class across
glioma samples **(B)** Heatmap depicting the relative intensity (Z-score) of glycan classes across WHO grades 2, 3,
and 4 **(C)** Boxplots of relative intensity for selected glycans from different glycan classes across WHO grades, with
corresponding representative images of glycan staining illustrating low and high glycan expression in different
patient samples. Glycan structures are displayed beside the boxplots **(D)** Heatmap showing the Z-score expression of
both glycans and RNA for glycan processing enzymes that were differentially expressed between any WHO grades.
Glycans are identified by cyan, and RNA is identified by dark blue. **(E)** Network diagram showing cell types linked
to glycan classes (fucosylated, agalactosylated, sialylated, tri-antennary) and their associated biological processes,
derived from multi-omic analysis. Edge weights between cells and glycans indicate correlation coefficient while
edge weights between glycans and biological processes indicate gene ratio (genes identified with high correlation /
total genes in pathway) **(F)** Heatmap of the top 25 genes correlated with glycan expression, with rows representing
glycan classes and columns representing genes, colored by correlation strength. **(G)** Scatter plot illustrating the
relationship between NRXN1 expression and fucosylated glycans, with dot size representing the neuron-to-tumor
cell ratio and color indicating WHO grade. **(H)** Scatter plot showing the association of GRN expression with
sialylated glycans, with dot size reflecting CD209+ DC/Mac abundance and color corresponding to WHO grade.

the stark remodeling in tumor glycans classes with grade cannot be reliably inferred from upstream
molecular information. Consequently, the regulatory role of glycans in glioma biology could be a
promising, yet understudied, target for both tumor classification and therapeutic targeting.
487
In order to better approximate the functional role of N-glycans in gliomas we looked to further
integrate the differential signals we overserved with the broader set cellular features mapped for
the TME. Therefore, we examined associations with both cellular abundances and total gene
(>11000 genes) expression to glycan classes. We further identified all transcripts correlated ($R^2 >$
0.4) with each glycan class and performed Gene Ontology (GO) analysis (**Figure 6E**). We found
that fucosylated and agalactosylated glycans, associated with brain-specific functions, including
neuronal development processes, were positively correlated with neuronal and tumor cells and
negatively correlated with most immune cell subsets (**Figure 6E**, **Figure S6E**). In contrast,
sialylated and tri-antennary glycans, associated with immune-related gene-expression modules,
such as antigen presentation and leukocyte activation, showed strong positive correlations with
immune cell abundance (**Figure 6E**). These data suggests that distinct glycosylation patterns may
play a key role in tumor related biological processes and influence disease state.
500
Finally, to identify the main drivers of these biological processes, we extracted the top 25 genes
based on correlative magnitude with glycan expression (**Figure 6F**). Among these were NRXN1,
NRXN2, NTRK2, LAPTM5, and GRN, all recognized for their involvement in glial tumor
biology, highlighting their potential roles in the TME[70–74]. In particular, NRXN1 (neurexin 1), a
cell surface receptor that binds neuroligins at synapses in the central nervous system, is known to
promote tumor growth through excitatory neuronal activity. Its strong positive correlation with
fucosylated glycans along with its positive correlation with the neuron-to-tumor cell ratio in
patients, highlights a potential link of surface fucose moieties with neuronal activation (**Figure
6G**). In contrast, GRN (granulin) showed a strong correlation with sialylated glycans. GRN has
been indicated to play a role in glioma tumor cell proliferation with increasing expression seen
with tumor grade[73,74]. Interestingly, our multi-omic analysis revealed that GRN showed a higher
weighted correlation when considering the abundance of CD209+ DC/Mac (**Figure 6H**). This
correlation suggests a potential involvement of specific myeloid subtypes with increased
sialylation involved in tumor growth. Together, these findings reveal a cohesive picture of glycan
involvement within the TME that was reinforced by correlations with cellular abundance and GO-
enriched biological pathways. This consistency across three orthogonal modalities provides a
holistic view of how glioma glycosylation could impact clinical outcomes.
518
519 **Multi-omic integration identifies distinct grade- and survival-specific TME features in GBM**
520
Integrating cellular protein, tissue glycan, and lineage gene expression can provide a deeper
understanding of the tumor microenvironment by capturing the complex interplay between
molecular pathways that drive tumor progression and immune responses[8]. Therefore, to understand
more holistically which aspects of this regulatory hierarchy drive glioma biology between disease
grades we trained a random forest classifier on multi-omic data from primary lesions to predict
WHO grade (**Figure 7A**). The model incorporated spatial proteomics (cell abundances, ratios,
marker intensity, spatial organization, etc.), glycomics, and transcriptomics from immune-
enriched (CD45+) or immune-poor (CD45-) glioma tissue regions. After removing sparse and
highly correlated features to reduce noise, the model achieved an AUC of 0.80 across 10 random

530
531
Figure 7. Multi-omic classification of glioma grades and survival features. (A) Schematic of the study design for
classifying glioma patients by WHO grade (2, 3, 4) and distinguishing between short-term and long-term survivors
in GBM using multi-omic data (transcriptome, proteome, N-glycome) **(B)** Boxplot showing the Area Under the
Curve (AUC) for the WHO grade classifier with original and permuted labels. **(C)** Top 75 Log10 importance scores
for different feature types (glycans, CD45+ and CD45- cell RNA, and MIBI) used in the WHO grade classifier. **(D)**
Top 25 feature importance ranking. Glycan (Cyan), MIBI (red), CD45- cell RNA (dark blue) and CD45+ cell RNA
(light blue) **(E)** Boxplot of endothelial cell CD31 intensity across WHO grades, with high and low representative
image of CD31 expression. Whole image shows CPM with endothelial cells colored in red **(F)** Inset shows raw
CD31 marker expression **(G)** Boxplot showing VEGFA RNA expression levels across WHO grades for CD45- cell
populations. **(H)** Boxplot showing AUC values for the survival classifier with original and permuted labels **(I)** Top
75 Log10 importance scores for different feature types (glycans, CD45+ and CD45- cell RNA, and MIBI) used in
the WHO grade classifier **(J)** Principal Component Analysis (PCA) plot using top 75 features from multi omic
classifier. Each dot represents a patient. blue for long term survivors (>1.5 years) and red for short term survivors
(<1.5 years).

trials with a permuted baseline of 0.57 (**Figure 7B**). The top 75 features by feature class revealed
that glycan features had the highest median importance, followed by MIBI features, RNA from
immune-poor tumor regions and RNA from immune-rich regions (**Figure 7C**). Among the most
important individual features were CD31 intensity in endothelial cells enriched in grade 4 (**Figures**

7D – 7F); H6N3F1 and H5N5F2, two fucosylated glycans enriched in LGG (**Figure 7D, Figure 6D**); and Vascular endothelial growth factor A (VEGFA), linked to vascular proliferation also enriched in grade 4 (**Figure 7G**). Notably, only 10% of the top 75 features were transcripts from immune-rich regions (**Table S4**) suggesting that the transcriptome of immune cells is more stable across grade than tumor cells. Interestingly these patterns were largely maintained when considering IDH mutation status (**Figure S7A**) suggesting that grade 4 IDH mutant astrocytoma is distinct from grade 2 and grade 3 IDH mutant tumors. Overall, in line with the initial N-glycan analysis findings (Figure 6), it appears that grade-driven alterations in glycan composition are key determinants of glioma biology, implicating the tumor glycocalyx as a central orchestrator of the molecular and cellular interactions that drive disease biology.

While tumor grading across diverse cancers, including gliomas, breast, and lung tumors, is typically determined by morphological changes, mutational status, and vascular proliferation, the molecular determinants of patient survival often diverge from these grading criteria[75]. To that end, using a similar approach, we trained a separate model for classifying short-term vs long-term survival in treatment-naive primary GBM to understand what aspects of the glioma TME separate these groups (**Figure 7A**). This model exhibited similar performance to the one above with an AUC of 0.81 and a permuted baseline of 0.59 (**Figure 7H**). However, the top 75 features for each model markedly differed with only the ratio of B7H3+ to TOX+ tumor cells being selected in both (**Tables S4 and S5**). In contrast to the grade-specific model, none of the top 75 survival-associated features included glycans (**Figure 7I**). Similarly, while transcripts from immune-rich regions had the lowest median importance for grade they had the highest median importance for survival predictions. This suggests that within the most aggressive disease, GBM, tumor immune responses are the predominant driving factor of survival. Transcripts such as MCM2, KIF3A, THOC6, and TMED3, all known to play a major role in tumor progression were among the most important features from immune-rich regions (**Figure S7B, Table S5**). To further validate this association by restricting the search space further, we confirmed that the top 25 features (**Figure S7B**) from the classifier were truly distinguishing between short- and long-term survivors, we performed PCA. The results showed a clear separation of the two groups along the PC1 axis, which accounted for 27% of the variance (**Figure 7J**). Together, these results demonstrate that while tumor glycan and vascularization features are most relevant for distinguishing / classifying tumor grade, immune-related transcriptional programs play a more critical role in driving survival signals, underscoring the value of multi-omic profiling in categorizing disease state progression and understanding and possibly perturbing patient outcomes.

Discussion

Gliomas are among the most aggressive tumors with the lowest overall survival (OS) rates of all cancers[1]. While significant improvements in OS have been achieved with immune or tumor targeted-therapies in extracranial malignancies like melanoma or lung adenocarcinoma, comparable progress in gliomas remains elusive[33,76,77]. This lack of progress suggests that mechanisms of tumor persistence in gliomas differ fundamentally from those occurring outside the blood brain barrier. To gain insights into why these approaches have stalled and how future therapies could circumvent these issues, we assembled a multicenter glioma cohort from 310 patients, representing 677 samples, to provide a unified, comprehensive view of this unique TME **(Figure 1)**. Multiomic integration of spatial proteomics, transcriptomics, and glycomics revealed

systematic differences in the structure, composition, and phenotype of gliomas with respect to tumor type and grade, a rationale for why targeted therapies have failed to improve OS, and new insights into how the tumor glycocalyx could be driving severity in high grade lesions. To enable future work to build upon these findings, we have made the data, analysis, and clinical endpoints freely available, without restriction, via an interactive web portal at www.bruce.parkerici.org **(Figure 1)** where users can browse annotated single cell data and view spatial proteomics images. This resource is also hosted on Bioimage Archive and GitHub where it can be downloaded in its entirety.

In previous work, we found that TME organization and morphology can be used to construct accurate trajectories of immune recruitment, infer immune phenotype, and to predict OS in both invasive and pre-invasive breast cancer[65,78]. Likewise, work by others have found this concept of spatial rulesets generalizes to other malignancies like melanoma and head and neck cancer as well[61,79]. In contrast, gliomas are highly diffuse and unstructured, often lacking the type of consistent spatial organization that is well suited for automated, extensible, and accurate scoring needed for developing these spatial rulesets using conventional tools.

To address these challenges on a larger scale with the less-structured TME of glioma, we used QUICHE to uncover subtle structural trends that were differentially expressed between patient groups[63]. For example, comparing IDH-mutant LGG structure in patient-matched primary and recurrent lesions revealed consistent and significant changes in how tumor and immune cell subsets localized with one another **(Figure 5)**. We would like to highlight two points from this analysis. First, while the TME structure differed significantly in primary and recurrent lesions, the prevalence of tumor and immune cell subsets did not. This means that the changes in cell localization observed here are likely due to corresponding differences in TME signaling rather than shifts in cell prevalence that could drive more happenstance interactions. Second, cells located within these differential niches were phenotypically distinct from cells outside of them. For instance, TIM3 and IDO1 were elevated in niches differentially enriched in primary tumors but not in recurrences. Intriguingly, previous work found the spatial organization of cells expressing TIM3 and IDO1 to be predictive of outcome in ovarian cancer, suggesting that the spatial organization of these cells are a surrogate for shifts between productive and tolerogenic immune responses. In contrast, when comparing on-therapy samples from patients receiving IO with those who received SOC, no structural differences were found. These findings suggest that while IO therapy may alter the functional status of certain immune populations, the lack of change in spatial organization could indicate these effects are transient and insufficient for recruiting durable antitumor immunity.

Tumor-targeting therapies such as small molecule EGFR inhibitors, monoclonal antibodies targeting B7H3, or CAR-T therapies against B7H3 and EGFR are another class of interventions that are being explored[10,50,80]. Like immunotherapy, these approaches have been effective in some tumors outside the brain but thus far in early clinical trials have not improved OS in glioma[4]. To understand why this might be, we conducted the first quantitative analysis of TA for seven targets currently in clinical trials **(Figure 2 and 3)**. This analysis revealed three insights for understanding why this approach has not yet been successful and how drug regimens might be optimized in future studies. First, ~57% of gliomas had a high proportion of tumor cells (>30%) that were pan-negative for all TAs quantified here, suggesting that these patients may not benefit from these therapies.

Second, for lesions with higher proportions of TA-positive cells, B7H3 and EGFR were the most
frequently expressed across all gliomas. Tumor cells in ~69% of GBM patients expressed one of
these antigens in comparable proportions to CD19 in B-cell malignancies (>72% antigen positive
tumor cells). However, B7H3 and EGFR also exhibited the highest degree of cell-to-cell variation
in expression (6-12 fold) (**Figures 4**). Given that low antigen copy number has been shown
previously to be associated with therapeutic failure in solid tumors, the proportion of glioma cells
with weak antigen expression could be a determinant here as well[81]. Lastly, personalized therapy
targeting one or more antigens could provide a means of improving tumor targeting and mitigating
antigen escape[57]. For example, early phase clinical trials using bispecific CD19/20 CAR-T to treat
Non-Hodgkin Lymphoma have shown >2 times the PFS compared to using CD19 CAR-T alone[82].
Based on our results, B7H3 and EGFR should offer the most coverage and overlap in GBM,
Astrocytoma and oligodendroglioma. In contrast, adult PXA patients showed high tumor cell
coverage with just B7H3, potentially making them ideal candidates for monotherapy. Still, given
the broad diversity of antigen expression (**Figure 2–4**) and the significant influence of immune
microenvironment on patient survival (**Figure 7**), effective long-term GBM treatment will likely
depend on strategies that harness and activate endogenous adaptive immune responses.
659
Recent advancements in glioma molecular classification, underscore the need to explore novel
biological dimensions beyond the proteome and transcriptome to fully capture tumor heterogeneity
and progression[83]. To understand how glioma tumor *N*-glycans are involved in driving disease
progression, we used a selective PNGaseF catalyzed cleavage assay with MALDI-IMS for label-
free, *de novo* detection of these structurally diverse posttranslational modifications[68]. *N*-glycans
are present on most membrane proteins where they comprise a large portion of glycocalyx, a
viscous carbohydrate coating surrounding the cell membrane that is involved in regulating a broad
range of cell-cell interactions[84]. Aberrant tumor glycosylation is a cancer hallmark that was first
identified over 50 years ago[85]. Since that time, innovations in chemical biology and mass
spectrometry have revealed the startlingly central role it plays in nearly every facet of disease
progression[86]. For example, TME sialylated glycans drive immune evasion by binding Siglecs
present on both lymphoid and myeloid cells[34–36].
672
Importantly, healthy brain tissue has distinct N-glycosylation patterns critical for normal neuronal
function and immune regulation[87]. Various sialylated structures in different brain subregions
modulate neuronal activity and intercellular communication, reflecting the tight regulation of the
brain's structural and immune milieu[88]. Disruption of these glycan profiles can have far-reaching
consequences, underscoring how aberrant sialylation in gliomas may potentiate immune evasion
and malignant infiltration.
679
Recognizing this, we quantified N-glycan class enrichments across glioma grades and found that
sialylated glycans were significantly enriched in grade 4 gliomas, where immune cells represent a
larger proportion of the tumor mass (**Figure 6**). To understand how these changes in the tumor
glycocalyx impact glioma composition and function, we correlated these findings with co-
registered spatial proteomics and transcriptomics. Tumor sialylation correlated with preferential
increases in specific immune subsets, including CD209+ DC/Mac, CD4+ T cells, and Tregs
(**Figure S6**). In line with these findings, previous studies have demonstrated that DC uptake of
sialylated antigens induces tolerogenic states that preferentially polarize naïve CD4+ T cells
toward Tregs[89].

689
Expanding on this, we trained a classifier on all three spatial modalities to simultaneously
determine contrasting glioma features that were predictive of grade and clinical outcome (**Figure**
7). Tumor glycans and features relating to neovascularization (i.e. CD31 endothelial expression,
VEGFA transcript abundance) accounted for the top five most predictive of glioma grade. In
contrast, these same spatial glycomic signals provided almost no value for predicting OS in
patients with GBM. Conversely, while spatial transcriptomics of immune-enriched CD45-positive
regions were the least valuable for predicting grade, they were the most valuable for predicting
GBM outcomes. These results indicate the divergent properties, specific to high grade disease,
should not be assumed to drive clinical outcome as well.
699
Taken together, this study provides a comprehensive view of the multi-omic landscape of human
gliomas. We established baseline measurements of tumor antigen variability, immune cell
landscapes, and the effects of therapeutic perturbations on the TME. Additionally, for the first time
we uncovered dynamic relationships between glycosylation, cellular features, and transcriptomes
in humans, creating a unified dataset that serves as a valuable resource for understanding glioma
biology and advancing translational research for this disease.
706
Going forward, multiple directions have emerged that future studies can build upon to deepen our
understanding of glioma pathophysiology. First, potential off-target consequences of B7H3-
directed treatments warrant scrutiny, given its prominent expression on non-tumor cells within the
brain. Additionally, strategies to enhance treatment efficacy should account for the notable
variability of tumor antigens at the single-cell level. Longitudinal studies will help us understand
how the tumor glycocalyx and TME architecture change over time, which could explain why
current therapies have limited effects. Using these insights in new clinical trials, especially those
testing bispecific or multiantigen-targeted treatments that account for individual glycocalyx and
antigen profiles, may help overcome the barriers that have held back glioma immunotherapy and
targeted therapies.
717

718 **Limitations of Study**
719
Despite the broad and diverse cohort in this study, having a larger sample size for each clinical
factor would have further increased the confidence of our findings. While we included longitudinal
sampling for some IDH mutant LGG lesions, having patient-matched primary and recurrent high-
grade samples would have provided even deeper insights into tumor evolution. The nature of this
study was entirely retrospective, and thus our findings, however significant, are primarily
correlative. Our use of TMA, though guided by expert pathologists to capture areas rich in tumor
and immune cells, inevitably limited the tissue coverage per patient and may have missed
important regions. In terms of single-cell analysis, although we measured a relatively large panel
of markers, future work could refine this panel to include additional markers for cytokines for
better immune cell functional characterization, as well as more robust markers for tumor cells. Our
spatial whole transcriptome profiling offered a perspective on immune-rich and immune-poor
regions, but a single-cell RNA approach might have yielded complementary data with more
cellular granularity more suited for multi-omic integration with our single-cell protein data.
Finally, practical constraints such as multi-site coordination and patient confidentiality meant we
could not obtain a complete clinical history for individuals, indicating that, while our dataset is

comprehensive, there is still room for improvement in both coverage and annotation.
736

Methods
738
<u>Study design and sample collection</u>
This cohort was assembled by combining tissue samples from several clinical trials and
institutional archives spanning multiple disease types, ages, treatment conditions, and WHO
grades. Multiple regions were collected per patient based on location and CD45/CD3 IHC for
immune rich regions. All samples were chosen under the supervision of a pathologist. A total of
677 cores from 310 unique patients were included on the 16 tissue blocks.
745

Phase I trial for GBM Response to Pembrolizumab (NCT02852655) of surgically accessible
recurrent/progressive glioblastoma that received 200 mg pembrolizumab (MK3475) intravenous
infusions. All patients provided written informed consent; the study was approved by institutional
review boards at all sites (Dana-Farber Cancer Institute; Huntsman Cancer Institute; M.D.
Anderson Cancer Center; Massachusetts General Hospital; Memorial Sloan Kettering Cancer
Center; University of California, Los Angeles; University of California, San Francisco) and
conducted according to the Declaration of Helsinki.
753

The LGG vaccine trial was made up of two phase I trials (NCT02924038, NCT02549833) (n =
53). Trial 1 was a pilot, randomized, two arm neoadjuvant vaccine study in human leukocyte
antigen-A2 positive (HLA-A2+) adults with World Health Organization (WHO) grade II glioma,
for which surgical resection of the tumor is clinically indicated. 4.96mg IMA950 and 1.4mg poly-
ICLC administered as one formulation followed by/or without 3mg/kg Varlilumab infusion
(intravenously) ~3 weeks before the date of scheduled standard-of-care surgery. Patients continue
receiving IMA950/poly-ICLC subcutaneous injections every week leading up to surgery. Trial 2
was a pilot neoadjuvant vaccine study in adults with WHO grade II glioma, for which surgical
resection of the tumor is clinically indicated. GBM6-AD lysate protein and poly-ICLC
administered as one formulation every week leading up to standard-of-care surgery to grade II
glioma ~3 weeks or was given post-surgery. Both studies were approved by the IRB at UCSF and
were conducted according to the Declaration of Helsinki, all patients provided written informed
consent.
767

The trial combining IL13Rα2-CAR-T cells with checkpoint inhibition (NCT04003649) is a phase
1, single-center, randomized study, comparing CAR T cells plus nivolumab with (Arm 1) and
without (Arm 2) neoadjuvant treatment with nivolumab and ipilimumab seven days before surgery.
The primary objectives are to establish the safety and feasibility of administering 50 x10^6
IL13(EQ)BBζ/CD19t+ T cells (100 x 106 T cells total) for 4 week cycles (n = 6 research subjects
per Arm) via dual delivery (administered through both the ICT and ICV catheters) in combination
with IV administration of checkpoint inhibitors in participants with recurrent glioblastoma and
IDH-mutant astrocytoma (grade 4). Samples collected were prior to CAR-T administration.
776

Treatment naive GBM samples tissues were acquired from Stanford Health Care's tissue
repository (n= 86) in which of the tumor samples there are, grade 2 astrocytoma (n = 5), grade 2
oligodendroglioma (n =5), grade 3 astrocytoma (n = 9), grade 3 oligodendroglioma (n =3), grade
4 astrocytoma (n = 8), and grade 4 GBM (n= 32). A select set of patient samples were labeled as

either core (n = 21), core-to-infiltrating (n = 8), or infiltrating edge (n = 14) depending on the
location of the sample within the tumor.
783
Pediatric H27M+ diffuse midline gliomas (DMG) and other pediatric high-grade gliomas (pHGG)
were acquired from Children's Hospital of Philadelphia (CHOP) (n =156). Of these, TMAs
included tumors designated as pediatric astrocytoma (n = 24), pediatric midline glioma (n = 25),
pediatric GBM (n = 33), pediatric ganglioglioma (n = 2), pediatric glioma (n = 16), pediatric high
grade gliomas (n = 8), pediatric PXA (n = 8), and pediatric thalamic glioma (n = 2), as well as
breast carcinoma (n = 4) and colon carcinoma (n = 4) as control tissues. More specific diagnostic
categorization of the pediatric cases, including integration of molecular diagnostic information
when available, can be obtained on request from the Children's Brain Tumor Network (CBTN,
https://cbtn.org/).
793
<u>Control TMA construction</u>
We constructed a control tissue microarray (TMA) to assess slide-to-slide staining variability. The
TMA comprised five 1.5mm control tissue cores sourced from archival FFPE specimens provided
by the Stanford pathology department, including duplicate cores of tonsil, lymph node, and
placenta, plus an extra tonsil core for asymmetry. Serial recuts of the control TMA, as well as the
cohort TMAs, were placed on the same slides. Additionally, a larger control TMA was constructed
of 2 mm cores of normal brain, tonsil, spleen, placenta, high grade glioma, thymus, lymph node,
testis, kikuchi disease, colon carcinoma, cholangiocarcinoma, liver, low grade glioma, melanoma,
ependymoma, breast carcinoma, ovarian cancer, lung scc, DIPG.
803
<u>MIBI staining</u>
Panel construction
Most antibodies used in this study were previously validated for MIBI-TOF[65,90,91]. Newly selected
target antibodies were validated via immunohistochemistry to confirm appropriate staining in
control tissues. All antibodies were metal-labeled using the Ionpath conjugation kit (IonPath,
Menlo Park, USA) according to the manufacturer's protocol. Calprotectin and Mast Cell markers
(Chymase and Tryptase) were conjugated to Ga69 and Ga71 respectively using monomeric
maleimido-mono-amide NOTA. To prevent degradation and prolong shelf life, labeled antibodies
were lyophilized individually with 100 mM trehalose in 1 µg or 5 µg aliquots. Titer optimization
for new targets was performed using serial dilutions (1 µg/mL, 0.5 µg/mL, 0.25 µg/mL, and 0.125
µg/mL) or the recommended titer for previously validated MIBI-TOF antibodies.
815
Cohort staining
To minimize batch effects, all TMAs were stained using two mastermixes: one for the tumor panel
and one for the immune panel. Serial TMA slices were stained separately for each panel. Fresh
antibody aliquots were reconstituted and combined into the mastermixes. To reduce errors,
staining was performed in pairs, with one person reading and the other pipetting. Detailed
procedures are available in our methods publication, with a step-by-step guide outlined below[28].
822
Interactive protocols
Reagent preparation: https://www.protocols.io/view/mibi-and-ihc-solutions-261geo7wyl47/v1
IHC staining: https://www.protocols.io/view/ihc-staining-x54v9moxmg3e/v1
MIBI staining: https://www.protocols.io/view/mibi-staining-dm6gprk2dvzp/v5

Sequenza staining: https://www.protocols.io/view/staining-sequenza-6qpvrdeo2gmk/v1
Antibody lyophilization: https://www.protocols.io/view/antibody-lyophilization-
kxygxex5kv8j/v1
830
<u>MIBI data generation</u>
MIBI run setup
MIBI data was acquired using the MIBIScope instrument (IonPath, Menlo Park, USA). TMA
cores were labeled by row and column for easy mapping to metadata. Each core was acquired on
two slides, one for the tumor panel and one for the immune panel, with screen captures taken to
align regions between slides. Automated checks, with user confirmation, corrected naming errors
in row and column assignments. Core order was randomized before acquisition to minimize
batch effects from instrument drift[92].
839
Acquisition settings
Consistent settings were applied across all MIBI data acquired in this study. Field of view (FOV)
size was selected based on sample availability. When feasible, FOVs measuring $(800 \ \mu m)^2$ and
$(2048 \ pixels)^2$ were captured. In cases of limited tissue, an FOV size of $(400 \ \mu m)^2$ and $(1024$
$pixels)^2$ was used instead. A custom preset with a beam current of 8.5nA and dwell time of 0.58ms
was applied to optimize acquisition speed and image clarity. All default background correction
and noise removal settings were disabled.
847
Image compensation with Rosetta
We used our previously described Rosetta Matrix Compensation as a method for correcting
contamination and background across channels in MIBI data[64]. In summary we empirically
determined signal overlap for each source-output channel pair in order to create a matrix of all
channel overlaps and use this as a template to subtract known overlapping source markers from
their target counterpart, similar to flow-cytometry compensation.
854
Intensity normalization using median pulse height
We used our previously described Intensity normalization using median pulse height (MPH) as a
method for correcting for decreased sensitivity across a run in MIBI data[64]. In summary we created
an exponential curve with a PMMA (polymethyl methacrylate) sample, linking MPH to a
normalization coefficient. Each channel then has a polynomial function that is fit individually per
mass over the run that ignores outliers in order to produce a predicted MPH for that image. This
MPH is then mapped to the earlier described exponential curve to get a normalization coefficient
for each image. We then use this coefficient to normalize the data, correcting for drift and restoring
consistent signal levels throughout the run. This approach mitigated the observed decrease in
signal over time, preserving data accuracy.
865
866 <u>MIBI data QC</u>
867 To detect potential batch effects, we assessed slide-to-slide variation. For batch effects across
868 slides, control samples on each slide served as references, allowing us to calculate changes in the
869 mean of non-zero pixels for each channel in each control sample.
870

Cell segmentation
We segmented all images in the cohort using the previously described Mesmer model[93]. Mesmer is a pre-trained deep learning model that utilizes two input channels: a nuclear marker and a membrane marker. For the nuclear channel, we used the HH3/dsDNA channel, and for the membrane channel, we combined the CD14, CD45, CD8, and HLA1 channels.

Cell clustering
Pipeline overview
Pixie is an unsupervised pixel clustering algorithm that uses the subcellular localization of proteins to improve cell classification from multiplexed image data[94]. We used pixie to phenotype cells in a four-step process. First, pixels were over-clustered in each image by training a self-organizing map (10 × 20 grid or 200 clusters) on the phenotypic lineage marker pixels across all images for both panels[95]. For the immune panel we used Calprotectin, CD3, Chym/Tryp, CD68, CD4, CD8, CD208, CD141, CD209, CD11b, CD14, CD123, CD20, CD206, CD31, CD45, CD133, CD163, GFAP, GLUT1, FoxP3, HLADR, NeuN, Olig2, TMEM119. For the tumor panel we used CD4, FOXP3, CD31, NeuN, CD8, CD3, CD163, TMEM119, HLADR, CD14, CD45, ApoE, NG2, B7H3, VISTA, EGFR, HER2, EGFRvIII, EphA2, CD70, GPC2, H3K27M, GM2 GD2, CD133, GFAP, HLA1, Olig2, IDH1 R132H, CD47. These pixel clusters were then grouped into 20 pixel metaclusters using consensus hierarchical clustering[96]. Next, for each cell in the image, we counted the proportion of pixels in each pixel metacluster to generate a *cell × pixel metacluster* matrix. Following the same procedure, cells were then over-clustered by training a self-organizing map (10 × 20 grid or 200 clusters) on this pixel metacluster matrix, then grouped into 20 cell metaclusters using consensus hierarchical clustering. We validated these predictions through manual inspection of cell type-to-marker expression using Adobe Photoshop. Pixie was implemented using the ark-analysis v0.7.0 package in Python.

Cluster inspection and cleanup
To confirm accurate clustering, tiles of all images within each panel were constructed. Cell assignments were manually verified by looking at images through Adobe Photoshop. Assignments were viewed and confirmed by toggling on the appropriate image channels, which were then overlaid with the segmentation boundary and cluster assignments. Systematic errors in clustering were then addressed by reclustering based on marker specific thresholds.

Combining Immune and Tumor Panels
To integrate immune and tumor panels collected from separate serial slices, each with its own cell table, a comprehensive table encompassing all tumor and immune markers was constructed. For each field of view (FOV), the immune FOV and tumor FOV were spatially aligned to ensure accurate integration.

Each FOV was evaluated and categorized into one of three groups: (1) already aligned, (2) requiring manual alignment, or (3) unable to align. For FOVs requiring manual alignment, cells from both panels were plotted, and one FOV was shifted along the x and y axes to match the spatial layout of the other. Post-alignment, cells lying outside the aligned space were excluded. FOVs that could not be aligned, due to discrepancies in serial slices, variations in FOV sizes, or missing FOVs, were excluded from further analysis.

917   From the immune panel (288,224 cells), immune cells, endothelial cells, and neurons were
918   retained, while tumor cells were excluded. From the tumor panel (869,069 cells), only tumor cells
919   were retained, with immune cells, endothelial cells, and neurons removed. This resulted in a single
920   final combined panel (1,157,293 cells) prevented double counting and ensured the accurate
921   integration of immune and tumor cells.
922
Feature extraction pipeline
Pipeline overview
Feature extraction pipeline incorporates a diverse array of features to thoroughly characterize the
tumor-immune microenvironment. For each sample in our cohort, we computed all features, such
as cell composition, functional marker expression, cell density, cell ratios, transcriptome gene
counts, and glycome relative intensities. Each feature was calculated at the image level. To ensure
accuracy, we applied minimum cell count thresholds for many of these calculations, omitting any
features that did not meet these thresholds in a given image. Once generated, all features were
transformed into a standardized format and consolidated into a single data frame for downstream
analysis.   Details   available   at   https://github.com/angelolab/publications/tree/main/2024-
Piyadasa_Oberlton_etal_Glioma.
934
935   *Aggregating computed features*
936   Once the image features were calculated, we integrated them into a unified data structure for
937   downstream analysis. Each feature was assigned a descriptive name, along with metadata
938   specifying the image compartment where it was calculated, the cell types and/or markers involved,
939   and its overarching feature category.
940
Gene Set Enrichment Analysis (GSEA)
For the proportion of either GM2/GD2 or B7H3 out of all tumor cells, the median value was
calculated across all samples. Log2 fold changes were computed for each sample by dividing the
feature value by the median value, followed by sorting in descending order to generate a ranked
list of samples. Samples were then grouped into two categories based on tumor region: "Tumor
Core" and "Tumor Infiltrating." Each group was treated as a gene set for GSEA. The ranked list
of samples, based on log2FC values, was used as input for GSEA. The minimum and maximum
sizes of gene sets were automatically calculated based on the group sizes. Enrichment scores were
then calculated for both tumor regions using R package fgsea v1.30.0.
950
QUICHE Analysis
We performed Quantitative InterCellular Niche Enrichment (QUICHE) analysis to identify
differentially enriched niche neighborhoods across conditions (e.g. LGG primary vs recurrent,
tumor core vs tumor infiltrating edge, and GBM vs Grade 4 astrocytoma)[63]. In summary, for each
condition comparison, we identified cellular niches according to spatial proximity within each
image, performed distribution-focused downsampling to select a subset of niches per patient
sample, constructed a k-nearest neighbor (k-NN) graph for differential neighborhood enrichment
testing, and labeled niche neighborhoods according to the most abundant cell types (knn = 50,
radius = 200, k_sim = 100, looking at top 3 cell types per niche, downsample was determined to
be within 1 standard deviation below the mean, and adjusted depending on the spread of cells per
FOV). Niche neighborhoods were considered significant if they had a spatial false discovery rate

(FDR) < 0.10. QUICHE was implemented using the Python package at

QUICHE vs Bulk Functional Marker Analysis

To compare the functional marker abundance within QUICHE neighborhoods versus on the bulk scale we started by first calculating, for each functional maker, the percent positive cells in each cell type within a differentially expressed neighborhood. From there we then aggregated all cells of the same type across all differentially expressed niches and calculated the mean percent positivity for each functional marker. We separately calculated, for each functional marker, the bulk cell percent positivity for all cells in each condition of the comparison criteria. For each condition we calculated the Log2FC between the QUICHE percent positivity and the bulk percent positivity. If <0.05 of bulk cells were positive for a given functional marker the Log2FC was filtered out. In order to avoid division by zero a pseudo count of 1e-6 was added to all values. In order to keep from large Log2FC values a cap was created at -5 and 5.

Random Forest Model

Data Import and Preprocessing:

We filter the data based on criteria such as modality, immunotherapy status, disease type, patient age, and tumor region to ensure that only relevant variables are included in the analysis. Further data cleaning is performed by removing highly correlated features. A correlation matrix is calculated to identify highly correlated features. Features exceeding a specified correlation threshold (0.6 for who grade and 0.72 for survival status in order to maintain a similar number of features between the conditions) had the feature with the largest mean absolute correlation removed, retaining only those that contribute unique information to the analysis. , eliminating redundancy in the dataset. We also exclude features with a large proportion of >50% missing values and features in which all values are the same.

Feature Harmonization and Normalization:

Each feature is normalized (Z-scored) to facilitate comparisons. For datasets with multiple fields of view (FOVs), feature values are averaged across FOVs to ensure uniform representation.

Model Setup:

Using the processed data, a random forest classifier is trained to predict survival status (i.e. short vs long) or who grade (i.e. 2, 3, 4). We implement k-fold cross-validation (k = 3) to evaluate model performance. To evaluate performance, we compared the predictions to the ground truth using the area under the receiver operator curve (AUC). We performed 3-fold cross validation in each of 10 runs to obtain a distribution of predictions.

Model Evaluation and Feature Importance:

For each training iteration, AUC values are recorded to assess model performance. The median importance of each feature across multiple runs is calculated using the Gini importance metric, and the top features are identified based on this metric.

Randomized Control:

To validate model performance, labels are randomly shuffled outcome labels without replacement and the model is re-run to compare AUC values against the original (non-randomized) model,

1008 random forest is implemented in R using caret v.6.0.94. A t-test is implemented in R using stats
1009 v.4.4.1 to determine statistical significance between the original and randomized AUC values,
1010 providing a measure of model reliability.
1011
1012 *Principal Component Analysis (PCA):*
1013 Principal components analysis (PCA) was performed on the top 20 features for visualization in R
1014 using stats v4.4.1.
1015
1016 <u>RNA-Glycan co-occurrence analysis</u>
1017 In order to investigate the correlation between the gene expression of enzymes involved in glycan
1018 generation and the abundance of glycans across different GBM stages, we conducted an analysis
1019 on differentially expressed genes and glycans.
1020
1021 Initially, we normalized the expression data for these genes and glycans by Z-scoring each gene
1022 or glycan's expression across GBM stages. Thereafter, k-means clustering was applied using k=6,
1023 a value driven by minimal k to account for all potential trends across the three stages of GBM.
1024 This ensured sufficient classes to account for any trends observed within our dataset, while
1025 minimizing clustering complexity.
1026
1027 Using literature references, glycans and genes were each assigned a glycan type, such as bi-
1028 antennary or fucosylated. It's noteworthy that a gene or glycan could be assigned to multiple types.
1029 Following this assignment, we quantified, for each glycan type, instances where both a glycan and
1030 an enzyme of that type shared the same k cluster.
1031
1032 For establishing a comparative basis, we generated a null distribution by randomly assigning genes
1033 and glycans to types 1,000 times. The previously calculated quantity was then Z-scored against
1034 this null distribution. A high Z-score indicates a glycan type with a more correlated expression of
1035 genes and glycans than one would expect by chance. This method accounts for the overall
1036 expression trends observed in our dataset as given by the distribution between k-clusters.
1037
<u>N-glycan MALDI imaging mass spectrometry of FFPE tissue slide preparation</u>
Tissue TMA slides were prepared for N-glycan imaging mass spectrometry analysis using a
standardized tissue preparation workflow which has been previously published for FFPE tissues[97].
Briefly, tissue slides were dewaxed with xylene and rehydrated using a gradient of ethanol in a
linear stainer (Leica Biosciences) programmed to 3 dips per wash for 30 seconds each. Antigen
retrieval was performed in citraconic anhydride buffer at pH 3 for 30 min in a heating chamber at
95 °C. Slides were briefly dried under vacuum and subject to enzyme treatment using HTX M3+
Sprayer (HTX Technologies, Chapel Hill, NC).5 passes of PNGaseF PRIME enzyme (N-Zyme
Scientifics, Doylestown, PA) at 0.1 µg/µL was applied at a rate 25 µL/min with a velocity of 1200
mm/min and a 3 mm offset at 10 psi and 45 °C. Slides were incubated in prewarmed humidity
chambers for 2 h at 37 °C. After PNGaseF digestion, 7 mg/mL CHCA matrix in 50% ACN/0.1%
TFA was applied to the slides at a rate of 100 µL/min with a velocity of 1300 mm/min and a 2.5
mm offset at 10 psi and 79 °C using the same sprayer. Slides were stored under vacuum until
analysis[97].
1052
<u>MALDI-MSI Data Acquisition and Processing</u>

All samples were analyzed with a timsTOF fleX MALDI-2 mass spectrometer (Bruker
Corporation, Billerica, MA). The following parameters were used: mass range 800-4000 m/z,
positive ion mode, 10 μm field size and 200 shots per pixel. Raw imaging data was processed in
SCiLS Lab version 2024b Pro (Bruker Corporation, Billerica, MA). In SCiLs Lab, N-glycan
spectra were normalized by total ion count and converted to vendor-neutral imzML format. The
imzML files were parsed using pyimzML in Python, and the expression at each m/z peak was
extracted as single-channel TIF images.
1061
Nanostring DSP Sample Preparation
Formalin-fixed, paraffin-embedded (FFPE) samples were prepared manually according to
NanoString GeoMx RNA-NGS protocols. Tissue microarrays (TMAs) were arranged to fit the
largest dimensions allowable for imaging on the DSP system. Five-micrometer sections were
placed on positively charged slides, baked, deparaffinized, and then sequentially washed in ethanol
and PBS. Antigen retrieval was performed with Tris-EDTA (pH 9.0) in a pressure cooker for 10
minutes at 100°C, followed by a PBS rinse. The samples were then incubated with Proteinase K
(1 µg/mL) for 15 minutes at 37°C and washed again in PBS.
1070
1071 Afterward, the slides were hybridized overnight at 37°C with human WTA at the recommended
1072 concentration. HybriSlip Hybridization Covers (Grace BioLabs) were applied during this step.
1073 Coverslips were detached by soaking the slides in 2× SSC plus 0.1% Tween-20, and any unbound
1074 probes were removed by two 25-minute washes in 50% formamide/2× SSC at 37°C, followed by
1075 a brief wash in 2× SSC.
1076
1077 For staining with morphology markers, slides were first incubated in blocking buffer at room
1078 temperature for 30 minutes in a humid chamber. They were then exposed to 500 nM SYTO13 and
1079 fluorescently labeled antibodies (CD45, GFAP) for 1–2 hours. A final wash in 2× SSC was
1080 performed before loading the slides onto the GeoMx DSP instrument.
1081
Nanostring DSP acquisition
DSP experiments were conducted in alignment with the NanoString GeoMx-NGS DSP Instrument
manual and established protocols[98]. In brief, slides were imaged across four fluorescence channels
(FITC/525 nm, Cy3/568 nm, Texas Red/615 nm, and Cy5/666 nm) to visualize the selected
morphology markers. Regions of interest (AOIs) were defined and collected for all samples, with
segmentation driven by CD45 high and CD45 low expression using the DSP auto-segmentation
tool and manually adjusted parameters. The selected AOIs were illuminated, and the released tags
were deposited into 96-well plates following the established protocol.
1090
Sequencing and data cleanup
Following the NanoString GeoMx-NGS Readout Library Prep guidelines, the dried DSP aspirate
was reconstituted in 10 µL of DEPC-treated water, and 4 µL of this solution was used for PCR
with NanoString SeqCode primers. These primers both amplify the tags and incorporate Illumina
adaptor sequences along with sample-specific barcodes. PCR products were pooled in fixed
volumes and then purified twice using AMPure XP beads (Beckman Coulter). The final libraries
were sequenced exclusively on an Illumina NovaSeq 6000 using 27 × 27 paired end reads.
1098

FASTQ files were handled with the NanoString GeoMx NGS Pipeline v2.0 or v2.2. Low-quality bases and adaptor sequences were trimmed, and paired-end reads were stitched together. Barcode and UMI regions were identified, allowing up to one mismatch for barcode matching. Reads sharing the same barcode were deduplicated by UMI. The raw read count correlated strongly with unique UMIs across AOIs (Supplemental Figure S9), supporting a generally consistent library amplification. Nevertheless, all analyses presented here employed UMI-deduplicated counts to account for potential PCR bias.

Count data were processed and normalized using GeoMxTools R package v1.0 (https://bioconductor.org/packages/release/bioc/html/GeomxTools.html). Specific parameters and detailed used can be found in https://github.com/angelolab/publications/blob/main/2024-Piyadasa_Oberlton_etal_Glioma/Data_Analysis/2_NS_DSP_cleanup.R

Resource availability

Lead contact

Requests for further information and resources should be directed to and will be fulfilled by the lead contact Michael Angel (mangelo0@stanford.edu).

Materials availability

All unique/stable reagents generated in this study are available from the lead contact with a completed materials transfer agreement.

Data and code availability

To provide broad access to our multi-omic data set (including single cell MIBI image overlays) and clinically annotated metadata, we have created an interactive online portal, www.bruce.parkerici.org. This portal further facilitates direct download of all raw data tables and features calculated. Raw MIBI images and segmentation files are deposited at BioImage Archive (https://www.ebi.ac.uk/biostudies/bioimages/studies/S-BIAD1579) for easy access.

Code to generate all figures is available at https://github.com/angelolab/publications/tree/main/2024-Piyadasa_Oberlton_etal_Glioma. Low-level processing code can be found at: https://github.com/angelolab/toffy. Spatial analysis code can be found here: https://github.com/angelolab/ark

Acknowledgements and funding

We thank the patients and their families for participating in all the trials. We thank Parker Institute for Cancer Immunotherapy (PICI) that supported the multi-site collaboration, funding the project and the development of the web resource. We also thank Alliance for Cancer Gene Therapy for their funding support for additional samples, as well as their scientific oversight. We acknowledge resources provided by the UCSF Brain Tumor SPORE Biorepository and Pathology Core. Dr. Piyadasa was supported by the Canadian Institutes of Health Research (CIHR) postdoctoral fellowship (FRN: 176490). N.F.G. was supported by NCI CA246880, NCI CA264307, and the

Stanford Graduate Fellowship. Dr. Phillips was partially supported by the National Institutes of Health (R01 NS131474). NIH/NCI P50CA097257 for JJP. NIH/NCI U01 CA242096 for RRD. Dr. Oldridge was supported by NHLBI R38 HL143613, NCI T32 CA009140, and a bridge fellowship from the Parker Institute for Cancer Immunotherapy and V Foundation for Cancer Research.

Declaration of interests

M.A. and S.B. are named inventors on patent US20150287578A1, which covers the mass spectrometry approach utilized by MIBI-TOF to detect elemental reporters in tissue using secondary ion mass spectrometry. M.A. and S.B. are board members and shareholders in IonPath, which develops and manufactures the commercial MIBI-TOF platform.

Author contributions

H.P., B.O., S.B. and M.A. conceived the project.

R.P., H.O., J.P., D.O., C.B., M.B, K.C., F.G., P.P., F.F., A.T., S.F. and M.G.H collected and processed FFPE tissues, clinical metadata and provided expert guidance and feedback throughout the project.

E.Y., M.T., M.S., S.B. and S.L. facilitated the multi institutional collaboration and provided funding for the project.

H.P., E.Y., M.T., C.S. and S.V. constructed the web portal and deposited data.

H.P., B.O., I.A. C.C.F., M.Bo., R.K., and M.R. generated and processed MIBI, MALDI and NanoString DSP data.

H.P., B.O., J.R., I.A., K.L., M.A., C. L., N.F.G., E.M., A.K., C.S. and M.S. analyzed multi omic data.

H.P., and B.O. generated figures.

H.P., B.O., S.B. and M.A. wrote the manuscript.

S.B. and M.A. supervised the project.

R.D. provided expert feedback on MALDI analysis.

All authors provided feedback on the manuscript.

Declaration of generative AI and AI-assisted technologies

During the preparation of this work, the author(s) used ChatGPT to refine the text and code. After using this tool or service, the author(s) reviewed and edited the content as needed and take(s) full responsibility for the content of the publication.

References

1. Ostrom, Q.T., Cioffi, G., Waite, K., Kruchko, C., and Barnholtz-Sloan, J.S. (2021). CBTRUS Statistical Report: Primary Brain and Other Central Nervous System Tumors Diagnosed in the United States in 2014–2018. Neuro-Oncology *23*, iii1–iii105. https://doi.org/10.1093/neuonc/noab200.

2.   Mohammed, S., Dinesan, M., and Ajayakumar, T. (2022). Survival and quality of life analysis in
glioblastoma multiforme with adjuvant chemoradiotherapy: a retrospective study. Rep Pract Oncol
Radiother *27*, 1026–1036. https://doi.org/10.5603/RPOR.a2022.0113.

3.   Al Sharie, S., Abu Laban, D., and Al-Hussaini, M. (2023). Decoding Diffuse Midline Gliomas: A
Comprehensive Review of Pathogenesis, Diagnosis and Treatment. Cancers (Basel) *15*, 4869.
https://doi.org/10.3390/cancers15194869.

4.   Liu, Y., Zhou, F., Ali, H., Lathia, J.D., and Chen, P. (2024). Immunotherapy for glioblastoma:
current state, challenges, and future perspectives. Cell Mol Immunol *21*, 1354–1375.
https://doi.org/10.1038/s41423-024-01226-x.

5.   Hervey-Jumper, S.L., and Berger, M.S. (2014). Role of Surgical Resection in Low- and High-Grade
Gliomas. Curr Treat Options Neurol *16*, 284. https://doi.org/10.1007/s11940-014-0284-7.

6.   Grabowski, M.M., Sankey, E.W., Ryan, K.J., Chongsathidkiet, P., Lorrey, S.J., Wilkinson, D.S.,
and Fecci, P.E. (2021). Immune suppression in gliomas. J Neurooncol *151*, 3–12.
https://doi.org/10.1007/s11060-020-03483-y.

7.   Karimi, E., Yu, M.W., Maritan, S.M., Perus, L.J.M., Rezanejad, M., Sorin, M., Dankner, M.,
Fallah, P., Doré, S., Zuo, D., et al. (2023). Single-cell spatial immune landscapes of primary and
metastatic brain tumours. Nature *614*, 555–563. https://doi.org/10.1038/s41586-022-05680-3.

8.   Ravi, V.M., Will, P., Kueckelhaus, J., Sun, N., Joseph, K., Salié, H., Vollmer, L., Kuliesiute, U.,
Ehr, J. von, Benotmane, J.K., et al. (2022). Spatially resolved multi-omics deciphers bidirectional
tumor-host interdependence in glioblastoma. Cancer Cell *40*, 639-655.e13.
https://doi.org/10.1016/j.ccell.2022.05.009.

9.   Xu, S., Tang, L., Li, X., Fan, F., and Liu, Z. (2020). Immunotherapy for glioma: Current
management and future application. Cancer Letters *476*, 1–12.
https://doi.org/10.1016/j.canlet.2020.02.002.

10.   Vitanza, N.A., Ronsley, R., Choe, M., Seidel, K., Huang, W., Rawlings-Rhea, S.D., Beam, M.,
Steinmetzer, L., Wilson, A.L., Brown, C., et al. (2025). Intracerebroventricular B7-H3-targeting
CAR T cells for diffuse intrinsic pontine glioma: a phase 1 trial. Nat Med, 1–8.
https://doi.org/10.1038/s41591-024-03451-3.

11.   Oprita, A., Baloi, S.-C., Staicu, G.-A., Alexandru, O., Tache, D.E., Danoiu, S., Micu, E.S., and
Sevastre, A.-S. (2021). Updated Insights on EGFR Signaling Pathways in Glioma. International
Journal of Molecular Sciences *22*, 587. https://doi.org/10.3390/ijms22020587.

12.   Monje, M., Mahdi, J., Majzner, R., Yeom, K.W., Schultz, L.M., Richards, R.M., Barsan, V., Song,
K.-W., Kamens, J., Baggott, C., et al. (2024). Intravenous and intracranial GD2-CAR T cells for
H3K27M+ diffuse midline gliomas. Nature, 1–8. https://doi.org/10.1038/s41586-024-08171-9.

13.   Lin, Y.-J., Mashouf, L.A., and Lim, M. (2022). CAR T Cell Therapy in Primary Brain Tumors:
Current Investigations and the Future. Front Immunol *13*, 817296.
https://doi.org/10.3389/fimmu.2022.817296.

14.   Nejo, T., Yamamichi, A., Almeida, N.D., Goretsky, Y.E., and Okada, H. (2020). Tumor antigens
in glioma. Seminars in Immunology *47*, 101385. https://doi.org/10.1016/j.smim.2020.101385.

15.   Barish, M.E., Weng, L., Awabdeh, D., Zhai, Y., Starr, R., D'Apuzzo, M., Rockne, R.C., Li, H.,
Badie, B., Forman, S.J., et al. (2022). Spatial organization of heterogeneous immunotherapy target
antigen expression in high-grade glioma. Neoplasia *30*, 100801.
https://doi.org/10.1016/j.neo.2022.100801.

16.   Miao, H., Choi, B.D., Suryadevara, C.M., Sanchez-Perez, L., Yang, S., Leon, G.D., Sayour, E.J.,
McLendon, R., Ii, J.E.H., Healy, P., et al. (2014). EGFRvIII-Specific Chimeric Antigen Receptor
T Cells Migrate to and Kill Tumor Deposits Infiltrating the Brain Parenchyma in an Invasive
Xenograft Model of Glioblastoma. PLOS ONE *9*, e94281.
https://doi.org/10.1371/journal.pone.0094281.

17.   Giakoumettis, D., Kritis, A., and Foroglou, N. (2018). C6 cell line: the gold standard in glioma
research. Hippokratia *22*, 105–112.

18. Hoogstrate, Y., Draaisma, K., Ghisai, S.A., Hijfte, L. van, Barin, N., Heer, I. de, Coppieters, W., Bosch, T.P.P. van den, Bolleboom, A., Gao, Z., et al. (2023). Transcriptome analysis reveals tumor microenvironment changes in glioblastoma. Cancer Cell *41*, 678-692.e7. https://doi.org/10.1016/j.ccell.2023.02.019.

19. Lam, K.H.B., Leon, A.J., Hui, W., Lee, S.C.-E., Batruch, I., Faust, K., Klekner, A., Hutóczki, G., Koritzinsky, M., Richer, M., et al. (2022). Topographic mapping of the glioblastoma proteome reveals a triple-axis model of intra-tumoral heterogeneity. Nat Commun *13*, 116. https://doi.org/10.1038/s41467-021-27667-w.

20. Puchalski, R.B., Shah, N., Miller, J., Dalley, R., Nomura, S.R., Yoon, J.-G., Smith, K.A., Lankerovich, M., Bertagnolli, D., Bickley, K., et al. (2018). An anatomic transcriptional atlas of human glioblastoma. Science *360*, 660–663. https://doi.org/10.1126/science.aaf2666.

21. Wildburger, N.C., Zhou, S., Zacharias, L.G., Kroes, R.A., Moskal, J.R., Schmidt, M., Mirzaei, P., Gumin, J., Lang, F.F., Mechref, Y., et al. (2015). Integrated Transcriptomic and Glycomic Profiling of Glioma Stem Cell Xenografts. J. Proteome Res. *14*, 3932–3939. https://doi.org/10.1021/acs.jproteome.5b00549.

22. Pinho, S.S., and Reis, C.A. (2015). Glycosylation in cancer: mechanisms and clinical implications. Nat Rev Cancer *15*, 540–555. https://doi.org/10.1038/nrc3982.

23. Sethi, M.K., Downs, M., Shao, C., Hackett, W.E., Phillips, J.J., and Zaia, J. (2022). In-Depth Matrisome and Glycoproteomic Analysis of Human Brain Glioblastoma Versus Control Tissue. Mol Cell Proteomics *21*, 100216. https://doi.org/10.1016/j.mcpro.2022.100216.

24. McFaline-Figueroa, J.R., Sun, L., Youssef, G.C., Huang, R., Li, G., Kim, J., Lee, E.Q., Nayak, L., Chukwueke, U., Beroukhim, R., et al. (2024). Neoadjuvant anti-PD1 immunotherapy for surgically accessible recurrent glioblastoma: clinical and molecular outcomes of a stage 2 single-arm expansion cohort. Nat Commun *15*, 10757. https://doi.org/10.1038/s41467-024-54326-7.

25. Saijo, A., Ogino, H., Butowski, N.A., Tedesco, M.R., Gibson, D., Watchmaker, P.B., Okada, K., Wang, A.S., Shai, A., Salazar, A.M., et al. (2024). A combinatory vaccine with IMA950 plus varlilumab promotes effector memory T-cell differentiation in the peripheral blood of patients with low-grade gliomas. Neuro Oncol *26*, 335–347. https://doi.org/10.1093/neuonc/noad185.

26. Ogino, H., Taylor, J.W., Nejo, T., Gibson, D., Watchmaker, P.B., Okada, K., Saijo, A., Tedesco, M.R., Shai, A., Wong, C.M., et al. (2022). Randomized trial of neoadjuvant vaccination with tumor-cell lysate induces T cell response in low-grade gliomas. J Clin Invest *132*, e151239. https://doi.org/10.1172/JCI151239.

27. Sussman, J.H., Oldridge, D.A., Yu, W., Chen, C.-H., Zellmer, A.M., Rong, J., Parvaresh-Rizi, A., Thadi, A., Xu, J., Bandyopadhyay, S., et al. (2024). A longitudinal single-cell and spatial multiomic atlas of pediatric high-grade glioma. Preprint at bioRxiv, https://doi.org/10.1101/2024.03.06.583588 https://doi.org/10.1101/2024.03.06.583588.

28. Keren, L., Bosse, M., Thompson, S., Risom, T., Vijayaragavan, K., McCaffrey, E., Marquez, D., Angoshtari, R., Greenwald, N.F., Fienberg, H., et al. (2019). MIBI-TOF: A multiplexed imaging platform relates cellular phenotypes and tissue structure. Science Advances *5*, eaax5851. https://doi.org/10.1126/sciadv.aax5851.

29. Zhang, Q., Abdo, R., Iosef, C., Kaneko, T., Cecchini, M., Han, V.K., and Li, S.S.-C. (2022). The spatial transcriptomic landscape of non-small cell lung cancer brain metastasis. Nat Commun *13*, 5983. https://doi.org/10.1038/s41467-022-33365-y.

30. Aichler, M., and Walch, A. (2015). MALDI Imaging mass spectrometry: current frontiers and perspectives in pathology research and practice. Lab Invest *95*, 422–431. https://doi.org/10.1038/labinvest.2014.156.

31. McDowell, C.T., Lu, X., Mehta, A.S., Angel, P.M., and Drake, R.R. (2023). Applications and continued evolution of glycan imaging mass spectrometry. Mass Spectrom Rev *42*, 674–705. https://doi.org/10.1002/mas.21725.

32. Wallace, E.N., West, C.A., McDowell, C.T., Lu, X., Bruner, E., Mehta, A.S., Aoki-Kinoshita, K.F., Angel, P.M., and Drake, R.R. (2024). An N-glycome tissue atlas of 15 human normal and cancer

tissue types determined by MALDI-imaging mass spectrometry. Sci Rep *14*, 489.
https://doi.org/10.1038/s41598-023-50957-w.

33.   Lim, M., Xia, Y., Bettegowda, C., and Weller, M. (2018). Current state of immunotherapy for
glioblastoma. Nat Rev Clin Oncol *15*, 422–442. https://doi.org/10.1038/s41571-018-0003-5.

34.   Wieboldt, R., Sandholzer, M., Carlini, E., Lin, C., Börsch, A., Zingg, A., Lardinois, D., Herzig, P.,
Don, L., Zippelius, A., et al. (2024). Engagement of sialylated glycans with Siglec receptors on
suppressive myeloid cells inhibits anticancer immunity via CCL2. Cell Mol Immunol *21*, 495–509.
https://doi.org/10.1038/s41423-024-01142-0.

35.   Dobie, C., and Skropeta, D. (2021). Insights into the role of sialylation in cancer progression and
metastasis. Br J Cancer *124*, 76–90. https://doi.org/10.1038/s41416-020-01126-7.

36.   Edgar, L.J., Thompson, A.J., Vartabedian, V.F., Kikuchi, C., Woehl, J.L., Teijaro, J.R., and
Paulson, J.C. (2021). Sialic Acid Ligands of CD28 Suppress Costimulation of T Cells. ACS Cent.
Sci. *7*, 1508–1515. https://doi.org/10.1021/acscentsci.1c00525.

37.   Keller, M.S., Gold, I., McCallum, C., Manz, T., Kharchenko, P.V., and Gehlenborg, N. (2025).
Vitessce: integrative visualization of multimodal and spatially resolved single-cell data. Nat
Methods *22*, 63–67. https://doi.org/10.1038/s41592-024-02436-x.

38.   Rozenblatt-Rosen, O., Regev, A., Oberdoerffer, P., Nawy, T., Hupalowska, A., Rood, J.E.,
Ashenberg, O., Cerami, E., Coffey, R.J., Demir, E., et al. (2020). The Human Tumor Atlas
Network: Charting Tumor Transitions across Space and Time at Single-Cell Resolution. Cell *181*,
236–249. https://doi.org/10.1016/j.cell.2020.03.053.

39.   Wirsching, H.-G., Felsberg, J., Prummer, M., Moisoiu, V., Lourman, R., Hertler, C., Antonios, M.,
Cimino, P.J., Roth, P., Gorlia, T., et al. (2023). Spatial immune profiling of glioblastoma identifies
an inflammatory, perivascular phenotype associated with longer survival. Acta Neuropathol *146*,
647–649. https://doi.org/10.1007/s00401-023-02617-6.

40.   Ludwig, K., and Kornblum, H.I. (2017). Molecular Markers in Glioma. Journal of neuro-oncology
134, 505. https://doi.org/10.1007/s11060-017-2379-y.

41.   Saratsis, A.M., Knowles, T., Petrovic, A., and Nazarian, J. (2023). H3K27M mutant glioma:
Disease definition and biological underpinnings. Neuro Oncol *26*, S92–S100.
https://doi.org/10.1093/neuonc/noad164.

42.   Han, S., Liu, Y., Cai, S.J., Qian, M., Ding, J., Larion, M., Gilbert, M.R., and Yang, C. (2020). IDH
mutation in glioma: molecular mechanisms and potential therapeutic targets. Br J Cancer *122*,
1580–1589. https://doi.org/10.1038/s41416-020-0814-x.

43.   Prosniak, M., Harshyne, L.A., Andrews, D.W., Kenyon, L.C., Bedelbaeva, K., Apanasovich, T.V.,
Heber-Katz, E., Curtis, M.T., Cotzia, P., and Hooper, D.C. (2013). Glioma grade is associated with
the accumulation and activity of cells bearing M2 monocyte markers. Clinical Cancer Research *19*,
3776–3786. https://doi.org/10.1158/1078-0432.CCR-12-1940.

44.   Liu, S., Zhang, C., Maimela, N.R., Yang, L., Zhang, Z., Ping, Y., Huang, L., and Zhang, Y. (2019).
Molecular and clinical characterization of CD163 expression via large-scale analysis in glioma.
OncoImmunology *8*, e1601478. https://doi.org/10.1080/2162402X.2019.1601478.

45.   Ferreira, D.W., Ulecia-Morón, C., Alvarado-Vázquez, P.A., Cunnane, K., Moracho-Vilriales, C.,
Grosick, R.L., Cunha, T.M., and Romero-Sandoval, E.A. (2020). CD163 overexpression using a
macrophage-directed gene therapy approach improves wound healing in *ex vivo* and *in vivo* human
skin models. Immunobiology *225*, 151862. https://doi.org/10.1016/j.imbio.2019.10.011.

46.   Chen, T., Chen, J., Zhu, Y., Li, Y., Wang, Y., Chen, H., Wang, J., Li, X., Liu, Y., Li, B., et al.
(2019). CD163, a novel therapeutic target, regulates the proliferation and stemness of glioma cells
via casein kinase 2. Oncogene *38*, 1183–1199. https://doi.org/10.1038/s41388-018-0515-6.

47.   Kumar, A., Mohamed, E., Tong, S., Chen, K., Mukherjee, J., Lim, Y., Wong, C.M., Boosalis, Z.,
Shai, A., Pieper, R.O., et al. (2022). CXCL14 Promotes a Robust Brain Tumor-Associated Immune
Response in Glioma. Clinical Cancer Research *28*, 2898–2910. https://doi.org/10.1158/1078-
0432.CCR-21-2830.

48. Chen, N., Li, X., Chintala, N.K., Tano, Z.E., and Adusumilli, P.S. (2018). Driving CARs on the
uneven road of antigen heterogeneity in solid tumors. Current Opinion in Immunology *51*, 103–
110. https://doi.org/10.1016/j.coi.2018.03.002.

49. Zhang, C., Burger, M.C., Jennewein, L., Genßler, S., Schönfeld, K., Zeiner, P., Hattingen, E.,
Harter, P.N., Mittelbronn, M., Tonn, T., et al. (2016). ErbB2/HER2-Specific NK Cells for Targeted
Therapy of Glioblastoma. JNCI: Journal of the National Cancer Institute *108*, djv375.
https://doi.org/10.1093/jnci/djv375.

50. Kontos, F., Michelakos, T., Kurokawa, T., Sadagopan, A., Schwab, J.H., Ferrone, C.R., and
Ferrone, S. (2021). B7-H3: An Attractive Target for Antibody-based Immunotherapy. Clinical
Cancer Research *27*, 1227–1235. https://doi.org/10.1158/1078-0432.CCR-20-2584.

51. Ghouzlani, A., Lakhdar, A., Rafii, S., Karkouri, M., and Badou, A. (2021). The immune checkpoint
VISTA exhibits high expression levels in human gliomas and associates with a poor prognosis. Sci
Rep *11*, 21504. https://doi.org/10.1038/s41598-021-00835-0.

52. Mączyńska, J., Raes, F., Da Pieve, C., Turnock, S., Boult, J.K.R., Hoebart, J., Niedbala, M.,
Robinson, S.P., Harrington, K.J., Kaspera, W., et al. (2022). Triggering anti-GBM immune
response with EGFR-mediated photoimmunotherapy. BMC Med *20*, 16.
https://doi.org/10.1186/s12916-021-02213-z.

53. GPC2-Directed Chimeric Antigen Receptor T Cells for the Treatment of Patients with Relapsed or
Refractory Neuroblastoma - NCI (2016). https://www.cancer.gov/about-cancer/treatment/clinical-
trials/search/v?id=NCI-2023-06712.

54. Poli, A., Wang, J., Domingues, O., Planagumà, J., Yan, T., Rygh, C.B., Skaftnesmo, K.O., Thorsen,
F., McCormack, E., Hentges, F., et al. (2013). Targeting glioblastoma with NK cells and mAb
against NG2/CSPG4 prolongs animal survival. Oncotarget *4*, 1527–1546.

55. Prapa, M., Chiavelli, C., Golinelli, G., Grisendi, G., Bestagno, M., Di Tinco, R., Dall'Ora, M., Neri,
G., Candini, O., Spano, C., et al. (2021). GD2 CAR T cells against human glioblastoma. npj Precis.
Onc. *5*, 1–14. https://doi.org/10.1038/s41698-021-00233-9.

56. Dong, X., Ren, J., Amoozgar, Z., Lee, S., Datta, M., Roberge, S., Duquette, M., Fukumura, D., and
Jain, R.K. (2023). Anti-VEGF therapy improves EGFR-vIII-CAR-T cell delivery and efficacy in
syngeneic glioblastoma models in mice. J Immunother Cancer *11*, e005583.
https://doi.org/10.1136/jitc-2022-005583.

57. Choe, J.H., Watchmaker, P.B., Simic, M.S., Gilbert, R.D., Li, A.W., Krasnow, N.A., Downey,
K.M., Yu, W., Carrera, D.A., Celli, A., et al. (2021). SynNotch-CAR T cells overcome challenges
of specificity, heterogeneity, and persistence in treating glioblastoma. Science Translational
Medicine *13*, eabe7378. https://doi.org/10.1126/scitranslmed.abe7378.

58. Schmidts, A., Srivastava, A.A., Ramapriyan, R., Bailey, S.R., Bouffard, A.A., Cahill, D.P., Carter,
B.S., Curry, W.T., Dunn, G.P., Frigault, M.J., et al. (2023). Tandem chimeric antigen receptor
(CAR) T cells targeting EGFRvIII and IL-13Rα2 are effective against heterogeneous glioblastoma.
Neuro-Oncology Advances *5*, vdac185. https://doi.org/10.1093/noajnl/vdac185.

59. Flugel, C.L., Majzner, R.G., Krenciute, G., Dotti, G., Riddell, S.R., Wagner, D.L., and Abou-el-
Enein, M. (2023). Overcoming on-target, off-tumour toxicity of CAR T cell therapy for solid
tumours. Nat Rev Clin Oncol *20*, 49–62. https://doi.org/10.1038/s41571-022-00704-3.

60. Horna, P., Nowakowski, G., Endell, J., and Boxhammer, R. (2019). Comparative Assessment of
Surface CD19 and CD20 Expression on B-Cell Lymphomas from Clinical Biopsies: Implications
for Targeted Therapies. Blood *134*, 5345. https://doi.org/10.1182/blood-2019-129600.

61. Arora, R., Cao, C., Kumar, M., Sinha, S., Chanda, A., McNeil, R., Samuel, D., Arora, R.K.,
Matthews, T.W., Chandarana, S., et al. (2023). Spatial transcriptomics reveals distinct and
conserved tumor core and edge architectures that predict survival and targeted therapy response.
Nat Commun *14*, 5029. https://doi.org/10.1038/s41467-023-40271-4.

62. Bastiancich, C., Snacel-Fazy, E., Fernandez, S., Robert, S., Stacchini, R., Plantureux, L.,
Boissonneau, S., Testud, B., Guillet, B., Debarbieux, F., et al. (2024). Tailoring glioblastoma

treatment based on longitudinal analysis of post-surgical tumor microenvironment. J Exp Clin
Cancer Res *43*, 311. https://doi.org/10.1186/s13046-024-03231-4.

63. Ranek, J.S., Greenwald, N.F., Goldston, M., Fullaway, C.C., Sowers, C., Kong, A., Mouron, S.,
Quintela-Fandino, M., West, R.B., and Angelo, M. (2025). QUICHE reveals structural definitions
of anti-tumor responses in triple negative breast cancer. Preprint at bioRxiv,
https://doi.org/10.1101/2025.01.06.631548 https://doi.org/10.1101/2025.01.06.631548.

64. Greenwald, N.F., Nederlof, I., Sowers, C., Ding, D.Y., Park, S., Kong, A., Houlahan, K.E., Varra,
S.R., Graaf, M. de, Geurts, V., et al. (2025). Temporal and spatial composition of the tumor
microenvironment predicts response to immune checkpoint inhibition. Preprint at bioRxiv,
https://doi.org/10.1101/2025.01.26.634557 https://doi.org/10.1101/2025.01.26.634557.

65. Keren, L., Bosse, M., Marquez, D., Angoshtari, R., Jain, S., Varma, S., Yang, S.-R., Kurian, A.,
Valen, D.V., West, R., et al. (2018). A Structured Tumor-Immune Microenvironment in Triple
Negative Breast Cancer Revealed by Multiplexed Ion Beam Imaging. Cell *174*, 1373-1387.e19.
https://doi.org/10.1016/j.cell.2018.08.039.

66. Schol, P., Elsas, M.J. van, Middelburg, J., Twilhaar, M.K.N., Hall, T. van, Sluis, T.C. van der, and
Burg, S.H. van der (2024). Myeloid effector cells in cancer. Cancer Cell *42*, 1997–2014.
https://doi.org/10.1016/j.ccell.2024.11.002.

67. Smith, B.A.H., and Bertozzi, C.R. (2021). The clinical impact of glycobiology: targeting selectins,
Siglecs and mammalian glycans. Nat Rev Drug Discov *20*, 217–243.
https://doi.org/10.1038/s41573-020-00093-1.

68. Drake, R.R., McDowell, C., West, C., David, F., Powers, T.W., Nowling, T., Bruner, E., Mehta,
A.S., Angel, P.M., Marlow, L.A., et al. (2020). Defining the human kidney N-glycome in normal
and cancer tissues using MALDI imaging mass spectrometry. Journal of Mass Spectrometry *55*,
e4490. https://doi.org/10.1002/jms.4490.

69. Drake, R.R., West, C.A., Mehta, A.S., and Angel, P.M. (2018). MALDI Mass Spectrometry
Imaging of N-Linked Glycans in Tissues. Adv Exp Med Biol *1104*, 59–76.
https://doi.org/10.1007/978-981-13-2158-0_4.

70. Fanlo, L., Gómez-González, S., Rozalén, C., Pérez-Núñez, I., Sangrador, I., Tomás-Daza, L.,
Gautier, E.L., Usieto, S., Rebollo, E., Vila-Ubach, M., et al. (2023). Neural crest-related NXPH1/α-
NRXN signaling opposes neuroblastoma malignancy by inhibiting organotropic metastasis.
Oncogene *42*, 2218–2233. https://doi.org/10.1038/s41388-023-02742-2.

71. Sun, P.-P., Liao, S.-X., Sang, P., Liu, M.-M., and Yang, J.-B. (2024). Lysosomal transmembrane
protein 5: Impact on immune cell function and implications for immune-related deficiencies.
Heliyon *10*. https://doi.org/10.1016/j.heliyon.2024.e36705.

72. Pattwell, S.S., Arora, S., Cimino, P.J., Ozawa, T., Szulzewsky, F., Hoellerbauer, P., Bonifert, T.,
Hoffstrom, B.G., Boiani, N.E., Bolouri, H., et al. (2020). A kinase-deficient NTRK2 splice variant
predominates in glioma and amplifies several oncogenic signaling pathways. Nat Commun *11*,
2977. https://doi.org/10.1038/s41467-020-16786-5.

73. Wang, M., Li, G., Yin, J., Lin, T., and Zhang, J. (2012). Progranulin overexpression predicts overall
survival in patients with glioblastoma. Med Oncol *29*, 2423–2431. https://doi.org/10.1007/s12032-
011-0131-6.

74. Bandey, I., Chiou, S.-H., Huang, A.-P., Tsai, J.-C., and Tu, P. -h (2015). Progranulin promotes
Temozolomide resistance of glioblastoma by orchestrating DNA repair and tumor stemness.
Oncogene *34*, 1853–1864. https://doi.org/10.1038/onc.2014.92.

75. Fallahpour, S., Navaneelan, T., De, P., and Borgo, A. (2017). Breast cancer survival by molecular
subtype: a population-based analysis of cancer registry data. Canadian Medical Association Open
Access Journal *5*, E734–E739. https://doi.org/10.9778/cmajo.20170030.

76. Weiss, S.A., Wolchok, J.D., and Sznol, M. (2019). Immunotherapy of Melanoma: Facts and Hopes.
Clinical Cancer Research *25*, 5191–5201. https://doi.org/10.1158/1078-0432.CCR-18-1550.

77. Saito, M., Suzuki, H., Kono, K., Takenoshita, S., and Kohno, T. (2018). Treatment of lung
adenocarcinoma by molecular-targeted therapy and immunotherapy. Surg Today *48*, 1–8.
https://doi.org/10.1007/s00595-017-1497-7.
78. Risom, T., Glass, D.R., Averbukh, I., Liu, C.C., Baranski, A., Kagel, A., McCaffrey, E.F.,
Greenwald, N.F., Rivero-Gutiérrez, B., Strand, S.H., et al. (2022). Transition to invasive breast
cancer is associated with progressive changes in the structure and composition of tumor stroma.
Cell *185*, 299-310.e18. https://doi.org/10.1016/j.cell.2021.12.023.
79. Thrane, K., Eriksson, H., Maaskola, J., Hansson, J., and Lundeberg, J. (2018). Spatially Resolved
Transcriptomics Enables Dissection of Genetic Heterogeneity in Stage III Cutaneous Malignant
Melanoma. Cancer Research *78*, 5970–5979. https://doi.org/10.1158/0008-5472.CAN-18-0747.
80. Abourehab, M.A.S., Alqahtani, A.M., Youssif, B.G.M., and Gouda, A.M. (2021). Globally
Approved EGFR Inhibitors: Insights into Their Syntheses, Target Kinases, Biological Activities,
Receptor   Interactions,   and   Metabolism.   Molecules   *26*,   6677.
https://doi.org/10.3390/molecules26216677.
81. Wagner, J., Wickman, E., DeRenzo, C., and Gottschalk, S. (2020). CAR T Cell Therapy for Solid
Tumors: Bright   Future   or   Dark   Reality?   Mol   Ther   *28*,   2320–2339.
https://doi.org/10.1016/j.ymthe.2020.09.015.
82. Larson, S.M., Chen, Y.Y., Johnston, J., Roy, S., Patel, P., and Benjamin, J.E. (2023). A
CD19/CD20-Directed Bispecific CAR-T Cell Therapy in Relapsed or Refractory Aggressive B-
Cell Non-Hodgkin Lymphoma (NHL). Blood *142*, 6892. https://doi.org/10.1182/blood-2023-
187614.
83. Louis, D.N., Perry, A., Wesseling, P., Brat, D.J., Cree, I.A., Figarella-Branger, D., Hawkins, C.,
Ng, H.K., Pfister, S.M., Reifenberger, G., et al. (2021). The 2021 WHO Classification of Tumors
of   the   Central   Nervous   System: a   summary.   Neuro   Oncol   *23*,   1231–1251.
https://doi.org/10.1093/neuonc/noab106.
84. Foote, C.A., Soares, R.N., Ramirez-Perez, F.I., Ghiarone, T., Aroor, A., Manrique-Acevedo, C.,
Padilla, J., and Martinez-Lemus, L.A. (2022). Endothelial Glycocalyx. Compr Physiol *12*, 3781–
3811. https://doi.org/10.1002/cphy.c210029.
85. Munkley, J., and Elliott, D.J. (2016). Hallmarks of glycosylation in cancer. Oncotarget *7*, 35478–
35489. https://doi.org/10.18632/oncotarget.8155.
86. Munkley, J., Mills, I.G., and Elliott, D.J. (2016). The role of glycans in the development and
progression of prostate cancer. Nat Rev Urol *13*, 324–333. https://doi.org/10.1038/nrurol.2016.65.
87. Williams, S.E., Noel, M., Lehoux, S., Cetinbas, M., Xavier, R.J., Sadreyev, R.I., Scolnick, E.M.,
Smoller, J.W., Cummings, R.D., and Mealer, R.G. (2022). Mammalian brain glycoproteins exhibit
diminished   glycan   complexity   compared   to   other   tissues.   Nat   Commun   *13*, 275.
https://doi.org/10.1038/s41467-021-27781-9.
88. Lee, J., Ha, S., Kim, M., Kim, S.-W., Yun, J., Ozcan, S., Hwang, H., Ji, I.J., Yin, D., Webster, M.J.,
et al. (2020). Spatial and temporal diversity of glycome expression in mammalian brain. Proc Natl
Acad Sci U S A *117*, 28743–28753. https://doi.org/10.1073/pnas.2014207117.
89. Castenmiller, C., Keumatio-Doungtsop, B.-C., van Ree, R., de Jong, E.C., and van Kooyk, Y.
(2021). Tolerogenic Immunotherapy: Targeting DC Surface Receptors to Induce Antigen-Specific
Tolerance. Front. Immunol. *12*. https://doi.org/10.3389/fimmu.2021.643240.
90. Hartmann, F.J., Mrdjen, D., McCaffrey, E., Glass, D.R., Greenwald, N.F., Bharadwaj, A., Khair,
Z., Verberk, S.G.S., Baranski, A., Baskar, R., et al. (2021). Single-cell metabolic profiling of human
cytotoxic T cells. Nat Biotechnol *39*, 186–197. https://doi.org/10.1038/s41587-020-0651-8.
91. McCaffrey, E.F., Donato, M., Keren, L., Chen, Z., Delmastro, A., Fitzpatrick, M.B., Gupta, S.,
Greenwald, N.F., Baranski, A., Graf, W., et al. (2022). The immunoregulatory landscape of human
tuberculosis granulomas. Nat Immunol *23*, 318–329. https://doi.org/10.1038/s41590-021-01121-x.
92. Piyadasa, H., Oberlton, B., Kong, A., Camacho Fullaway, C., Reddy Varra, S., Sowers, C., and
Tsai, A.G. (2023). Rapid Setup of Tissue Microarray and Tiled Area Imaging on the Multiplexed

Ion Beam Imaging Microscope using the Tile/SED/Array Interface. J Vis Exp.
https://doi.org/10.3791/65615.

93.   Greenwald, N.F., Miller, G., Moen, E., Kong, A., Kagel, A., Dougherty, T., Fullaway, C.C.,
McIntosh, B.J., Leow, K.X., Schwartz, M.S., et al. (2022). Whole-cell segmentation of tissue
images with human-level performance using large-scale data annotation and deep learning. Nat
Biotechnol *40*, 555–565. https://doi.org/10.1038/s41587-021-01094-0.

94.   Liu, C.C., Greenwald, N.F., Kong, A., McCaffrey, E.F., Leow, K.X., Mrdjen, D., Cannon, B.J.,
Rumberger, J.L., Varra, S.R., and Angelo, M. (2023). Robust phenotyping of highly multiplexed
tissue imaging data using pixel-level clustering. Nat Commun *14*, 4618.
https://doi.org/10.1038/s41467-023-40068-5.

95.   Van Gassen, S., Callebaut, B., Van Helden, M.J., Lambrecht, B.N., Demeester, P., Dhaene, T., and
Saeys, Y. (2015). FlowSOM: Using self-organizing maps for visualization and interpretation of
cytometry data. Cytometry Part A *87*, 636–645. https://doi.org/10.1002/cyto.a.22625.

96.   Monti, S., Tamayo, P., Mesirov, J., and Golub, T. (2003). Consensus Clustering: A Resampling-
Based Method for Class Discovery and Visualization of Gene Expression Microarray Data.
Machine Learning *52*, 91–118. https://doi.org/10.1023/A:1023949509487.

97.   Drake, R.R., Powers, T.W., Norris-Caneda, K., Mehta, A.S., and Angel, P.M. (2018). In Situ
Imaging of N-Glycans by MALDI Imaging Mass Spectrometry of Fresh or Formalin-Fixed
Paraffin-Embedded Tissue. Current Protocols in Protein Science *94*, e68.
https://doi.org/10.1002/cpps.68.

98.   Document Library NanoString University. https://university.nanostring.com/page/document-
library.

1511