Thermal cycling-hyperthermia attenuates rotenone-induced cell injury in SH-SY5Y cells through heat-activated mechanisms

Yu-Yi Kuo[1,2], Guan-Bo Lin[1,2], You-Ming Chen[1,2], Hsu-Hsiang Liu[1,2], Fang-Tzu Hsu[1,2], Yi Kung[1,2], and Chih-Yu Chao[1,2,3*]

[1]Department of Physics, Lab for Medical Physics & Biomedical Engineering, National Taiwan University, 106319 Taipei, Taiwan,

[2]Biomedical & Molecular Imaging Center, National Taiwan University, 106319 Taipei, Taiwan,

[3]Graduate Institute of Applied Physics, Biophysics Division, National Taiwan University, 106319 Taipei, Taiwan

*Correspondence to: Chih-Yu Chao; **email:** cychao@phys.ntu.edu.tw

ABSTRACT

Parkinson's disease (PD) is the second most prevalent neurodegenerative disease. It is characterized by mitochondrial dysfunction, increased reactive oxygen species (ROS), α-synuclein (α-syn) and phosphorylated-tau protein (p-tau) aggregation, and dopaminergic neuron cell death. Current drug therapies only provide temporary symptomatic relief and fail to stop or reverse disease progression due to the severe side effects or the blood-brain barrier. This study aimed to investigate the neuroprotective effects of thermal cycling-hyperthermia (TC-HT) in an *in vitro* PD model using rotenone (ROT)-induced human neural SH-SY5Y cells. Our results revealed that TC-HT reduced ROT-induced mitochondrial apoptosis and ROS accumulation in SH-SY5Y cells. TC-HT also inhibited the expression of α-syn and p-tau through heat-activated pathways associated with sirtuin 1 (SIRT1) and heat-shock protein 70 (Hsp70), involved in protein chaperoning, and resulted in the phosphorylation of Akt and glycogen synthase kinase-3β (GSK-3β), which inhibit p-tau formation. These findings underscore the potential of TC-HT as an effective treatment for PD, which can be practically implemented via the focused ultrasound (FUS).

Keywords: rotenone, Parkinson's disease, hyperthermia, neuroprotection, antioxidation

INTRODUCTION

Parkinson's disease (PD) is the second most prevalent neurodegenerative disease (NDD) after Alzheimer's disease (AD) in elderly individuals, and affects more than 8.5 million patients worldwide [1, 2]. PD is primarily characterized by the degeneration of dopaminergic neurons in the substantia nigra pars compacta, along with the abnormal accumulation of Lewy bodies in the striatum, resulting in clinical motor symptoms such as tremors, rigidity, akinesia, and postural instability [3]. Besides aging, exposure to pesticides such as rotenone (ROT) can also induce PD-like symptoms. ROT, a natural compound derived from plants such as *Derris* and *Lonchocarpus*, has been utilized as a vegetable pest control agent for over a century [4]. It specifically inhibits mitochondrial complex I, which causes mitochondrial dysfunction and increases reactive oxygen species (ROS) production [5, 6]. Furthermore, ROS promotes the aggregation of α-synuclein (α-syn) and phosphorylated-tau protein (p-tau), ultimately leading to neuronal death [7]. In animal models exposed to ROT, motor symptoms associated with PD are observed along with dopaminergic neurodegeneration in the substantia nigra, with Lewy body accumulation occurring in these neurons [8]. Hence, ROT-exposed cell and animal models serve as valuable models for research on PD therapy.

The mechanisms that trigger PD are currently under investigation. Current therapeutic approaches primarily focus on regulating dopamine and acetylcholine levels in the dopaminergic system of the striatum to alleviate associative motor symptoms [9]. These strategies include the use of levodopa to augment dopamine levels, monoamine oxidase-B inhibitors to prevent dopamine breakdown, and anticholinergic agents to reduce acetylcholine levels [10]. However, the effectiveness of drug treatment is often limited by the selective permeability of the blood-brain barrier or possible adverse effects, especially in elderly patients [10, 11]. While these treatments provide temporary symptomatic relief, they do not address the underlying cause of PD, which could be attributed to the accumulation of misfolded proteins in neurons [12, 13]. As a result, these treatments slow but do not halt or reverse disease progression. Therefore, it is crucial to find alternative treatments that target aggregated proteins, to cure the disease.

Heat therapy has long been recognized as an effective alternative treatment, with hyperthermia (HT) resulting in therapeutic effects across various clinical fields, including ophthalmology, cardiology, and oncology [14]. Additionally, HT-induced moderate thermal stress can activate neuroprotective mechanisms that defend against oxidative stress and inhibit protein aggregation [15]. This is achieved by up-regulating enzymes involved in degradation [16], activating heat-shock protein 70 (Hsp70) for protein chaperoning [15, 17], and increasing phosphorylated Akt (p-Akt) levels [18], which inactivate glycogen synthase kinase-3β (GSK-3β), thereby suppressing p-tau formation [19, 20]. Moreover, the application of focused ultrasound (FUS) for heat delivery has additional advantages for NDD treatment, including the capability to penetrate the blood-brain barrier and precisely elevate the temperature in

targeted brain regions [21, 22], thereby mitigating the risks associated with invasive
surgical procedures.

While heat therapy has therapeutic potential, caution is essential during its
continuous application due to potential adverse effects. Studies have shown that
continuous exposure to high temperatures may impair neural cells and potentially
damage the central nervous system [23, 24]. The continuous application of
conventional HT treatment is associated with risks such as neurological and systemic
complications, leading to potential irreversible neural damage [25]. Therefore, it is
crucial to develop a safer and more effective method for heat application during NDD
treatment. Our previous research introduced a mild thermal treatment, thermal-cycling
hyperthermia (TC-HT), which enhances protective mechanisms in SH-SY5Y cells
against oxidative stressors such as H_2O_2 and β-amyloid (Aβ) while preserving cell
viability [26]. Moreover, TC-HT has been demonstrated to reverse Aβ-induced
cognitive and memory impairments in AD mouse models, while concurrently
activating neuroprotective mechanisms in the hippocampus of these mice [27]. These
findings suggest that TC-HT could be a potential therapeutic strategy for PD, as it
might activate protective mechanisms in neuron cells without inducing adverse effects.

In this study, we used an ultrasound (US) exposure system to administer TC-
HT treatment to SH-SY5Y cells, aiming to investigate its therapeutic potential in a
ROT-induced *in vitro* PD model. Our findings indicate that TC-HT effectively
protects SH-SY5Y cells from ROT-induced damage by significantly inhibiting the
increase in mitochondrial apoptosis and the accumulation of intracellular ROS.
Importantly, the neuroprotective effects of TC-HT significantly exceed those of
continuous HT. Furthermore, TC-HT significantly reduced ROT-elevated PD markers,
α-syn and p-tau, by potentially up-regulating proteins with multiple neuroprotective
roles, including Hsp70 and its upstream regulator sirtuin 1 (SIRT1) for chaperoning,
and promoting the inactivation of GSK-3β by p-Akt to reduce p-tau formation. These
findings highlight TC-HT as an alternative PD therapy and provide insights into its
underlying regulatory mechanisms against protein aggregation.

RESULTS

TC-HT restores the ROT-impaired viability of SH-SY5Y cells

As a pesticide, ROT induces the loss of dopaminergic neurons, promotes Lewy body
aggregation in the brain, and causes movement dysfunction *in vivo*, making it a
commonly used agent for creating PD models *in vitro* and *in vivo* [5]. In this study,
we first evaluated ROT cytotoxicity in SH-SY5Y cells in a dose-dependent manner
using the CCK-8 assay. As shown in Figure 1A, treatment with 5 μM ROT reduces
SH-SY5Y cell viability to 65.3% compared to untreated controls, and this
concentration was subsequently used for further experiments. To determine the
optimal temperature that could maximize heat-associated neuroprotective effects, SH-

SY5Y cells were subjected to TC-HT at varying US intensities. Figure 1B indicates
that a 9-cycle TC-HT exhibits no protective effect at US intensities below 1.0 W. At
an intensity of 1.0 W, where the high temperature reached 40.0 ± 0.5 °C, TC-HT
significantly restored ROT-impaired cell viability from 67.2% to 90.4%. However, at
an intensity of 1.5 W, recovery effect decreased to 77.3% with no statistical
significance. These results suggest that the optimal US intensity for rescuing ROT-
injured SH-SY5Y cells was 1.0 W, corresponding to a stabilized high temperature of
40.0 ± 0.5 °C. Additionally, our previous study demonstrated that the total duration of
heat application influences the neuroprotective efficacy of TC-HT [26]. Thus, TC-HT
at a US intensity of 1.0 W was applied to SH-SY5Y cells for 6 and 9 cycles, and the
results obtained under these conditions were compared to those obtained with
continuous HT for equal heating durations of 1 and 1.5 h to assess the neuroprotective
effects of TC-HT and HT. As shown in Figure 1C, 9-cycle TC-HT significantly
improved ROT-reduced SH-SY5Y cell viability from 68.5% to 85.0%, outperforming
the non-significant recovery observed with 6-cycle TC-HT. In contrast, both 1 and 1.5
h HT did not result in significant protective effects on ROT-reduced cell viability
(from 68.5% to 73.2% and 71.6%, respectively), indicating that continuous thermal
treatment may be detrimental to neural cells. Overall, our data show that TC-HT with
1.0 W US intensity and 9 cycles results in optimal neuroprotective efficacy for ROT-
injured SH-SY5Y cells, which were used in subsequent experiments.

Figure 1. Neuroprotective effects of TC-HT on SH-SY5Y against ROT
cytotoxicity. CCK-8 assay was conducted to determine the viabilities of SH-SY5Y
cells following (**A**) different ROT concentrations, (**B**) TC-HT treatments with
different US intensities, and (**C**) TC-HT treatments with different cycles, along with
corresponding HT treatments with equal heating durations. The viabilities were
measured 24 h after ROT treatment. Data were presented as the mean $\pm$ standard
deviation in triplicate. Significance levels between indicated groups are denoted as
**$P < 0.01$ and ***$P < 0.001$.

TC-HT attenuates ROT-induced mitochondrial apoptosis

Rotenone (ROT) is known to inhibit mitochondrial complex I function, leading to
mitochondrial dysfunction and neural cell apoptosis [5]. To evaluate the protective
effects of TC-HT, we employed flow cytometry using the fluorescence dyes Annexin
V-FITC and PI for apoptosis assessment, along with JC-1 fluorescence staining, to
assess the MMP. As illustrated in Figure 2A, TC-HT significantly reduced the
apoptotic rates in ROT-exposed neural cells from 30.7% to 14.2%, greatly
outperforming continuous HT with statistical significance, which only reduced the
apoptotic rate to 25.7%. Additionally, the results of JC-1 staining in Figure 2B
demonstrate that TC-HT significantly inhibited ROT-induced MMP loss, with the
MMP level (normalized fluorescence intensity ratio of PE/FL1 channel) recovering
from 0.65-fold to 0.85-fold of control group, indicating a protective effect of TC-HT

on mitochondrial integrity against ROT-induced damage. In contrast, continuous HT
resulted in a slight and non-significant increase in the fluorescence intensity ratio to
0.67-fold of control group. Figure 2C presents confocal microscopy images of SH-
SY5Y cells 24 h after each treatment, with red/green fluorescence intensities
consistent with the results shown in Figure 2B. Collectively, these findings highlight
that TC-HT effectively protects SH-SY5Y cells from ROT-induced damage by
inhibiting mitochondrial dysfunction and subsequent apoptosis, compared to HT.

Figure 2. TC-HT attenuates ROT-induced mitochondrial apoptosis. (**A**)
Apoptosis after treatment was analyzed using flow cytometry using Annexin V-
FITC/PI double staining, and the apoptotic percentage (LR+UR) was calculated. (**B**)
MMP level after treatment was analyzed via flow cytometry with JC-1 fluorescense
dye, and the MMP levels after each treatment were determined by the normalized
fluorescence intensity ratio of PE/FL1 channel relative to control. (**C**) The red and
green fluorescence signals of JC-1 after treatment were imaged by a confocal
microscope. Scale bar = 100 μm. Data were presented as the mean ± standard
deviation in triplicate. Significance levels between indicated groups are denoted as *P
< 0.05, **P < 0.01, and ***P < 0.001.

TC-HT reduces ROT-induced intracellular ROS and promotes antioxidative
protein expression

ROS is implicated in the progression of NDDs such as AD and PD, with excessive
accumulation originating from mitochondrial dysfunction [28]. To investigate the
inhibitory effect of TC-HT on intracellular ROS levels, we performed flow cytometry
to assess intracellular ROS levels in SH-SY5Y cells stained with the fluorescent dye
DHE. As shown in Figure 3A, TC-HT significantly down-regulated the ROT-induced
increase in intracellular ROS levels in SH-SY5Y cells, decreasing it from 1.32-fold to
1.04-fold compared to the control. In contrast, continuous HT only slightly reduced
ROS levels to 1.28-fold of the control, without any statistical significance. This
suggests that TC-HT is more effective in mitigating ROT-induced ROS accumulation
than continuous HT. Furthermore, as shown in Figure 3B, we examined the
expression of the antioxidative enzyme SOD2, which reduced intracellular ROS
levels by converting harmful $O_2^{\bullet-}$ into less toxic hydrogen peroxide. Thus, TC-HT
significantly restored ROT-reduced SOD2 expression from 0.81-fold to 1.18-fold of
the control, surpassing the 0.76-fold expression observed with continuous HT. Thus,
TC-HT is an effective antioxidative treatment for inhibiting ROT-induced
intracellular ROS accumulation in SH-SY5Y cells, providing robust neuroprotection
in vitro.

Figure 3. TC-HT reduces ROT-induced intracellular ROS via SOD2 promotion.
(**A**) Intracellular superoxide radical anion ($O_2^{\bullet-}$) levels of SH-SY5Y cells were

determined by flow cytometry using the fluorescent dye DHE, and the DHE mean
fluorescence intensities after each treatment were calculated. **(B)** Representative
western blots of the antioxidative enzyme SOD2 and its expression levels after each
treatment were quantified. GAPDH was used as loading control. Data were presented
as the mean ± standard deviation in triplicate. Significance levels between indicated
groups are denoted as **$P < 0.01$ and ***$P < 0.001$.

TC-HT down-regulates the expression of ROT-induced PD marker proteins

The accumulation of α-syn and p-tau, which aggregate as Lewy bodies and tau fibrils
in the degenerated dopaminergic neurons of the substantia nigra, are critical
pathological hallmarks of PD [12, 13, 29]. In this study, we used the AT8 antibody to
recognize tau protein phosphorylated at the serine 202 and threonine 205 residues.
Our findings show that ROT treatment significantly increased the expression levels of
α-syn and AT8 in SH-SY5Y cells to 1.25-fold and 1.47-fold of the control,
respectively (Figure 4). In the study, TC-HT effectively reduced the expression level
of α-syn to 1.07-fold and AT8 to 1.15-fold of the control, with both reductions
achieving statistical significance. In contrast, continuous HT did not show significant
effect on the levels of these marker proteins compared to the levels observed with
ROT treatment, exhibiting expression levels of 1.28-fold for α-syn and 1.40-fold for
AT8. These results suggest that TC-HT is more effective in reducing the levels of PD
marker proteins, thereby providing enhanced protection for SH-SY5Y cells against
ROT-induced impairments.

Figure 4. TC-HT down-regulates the expression levels of ROT-induced PD
marker proteins. Representative western blots and the quantifications of **(A)** α-syn
and **(B)** AT8 after each treatment. GAPDH was used as loading control. Data were
presented as the mean ± standard deviation in triplicate. Significance levels between
indicated groups are denoted as *$P < 0.05$ and **$P < 0.01$.

TC-HT attenuates ROT-induced PD *in vitro* via heat-activated mechanisms

Given that TC-HT effectively down-regulates ROT-induced PD markers in SH-SY5Y
cells, we further investigated the neuroprotective mechanisms activated by TC-HT,
particularly focusing on the inhibition of α-syn and p-tau accumulation. Chaperones
play an important role in regulating the accumulation of misfolded proteins. Hsp70, a
member of the heat-shock protein family, is a heat-activated chaperone that prevents
and reduces α-syn and p-tau accumulation [30, 31]. Our findings show that both
Hsp70 and its upstream transcription factor, SIRT1 [32], were significantly up-
regulated following TC-HT treatment, with Hsp70 levels increasing from 0.67-fold to
0.97-fold and SIRT1 from 0.67-fold to 0.83-fold of the control (Figures 5A, B),
surpassing the expression levels of 0.64-fold and 0.53-fold observed with continuous

HT. These results suggest that TC-HT enhances protective effects against ROT by
promoting greater expressions of SIRT1 and Hsp70 chaperones compared to HT,
thereby inhibiting α-syn and p-tau accumulation. In addition to chaperoning,
preventing protein aggregation is another important neuroprotective mechanism for
PD. GSK-3β, a protein responsible for tau hyperphosphorylation, can be inactivated
by p-Akt through phosphorylation at the serine 9 residue [19] The reduced expression
of p-GSK-3β (Ser9) has been linked to increased AT8 levels [20]. Therefore,
regulating the expression of p-GSK-3β (Ser9) is critical for reducing p-tau
aggregation. As shown in Figures 5C & D, ROT treatment down-regulates p-Akt and
p-GSK-3β (Ser9), while TC-HT effectively reverses this decline, significantly
improving p-Akt level from 0.75-fold to 1.25-fold and p-GSK-3β (Ser9) level from
0.66-fold to 0.88-fold of the control. These results surpass the expression levels of
0.88-fold for p-Akt and 0.60-fold for p-GSK-3β (Ser9) observed with continuous HT,
which do not show statistical significance compared to ROT-treated SH-SY5Y cells.
Corresponding with the increased AT8 levels in Figure 4, these findings suggest that
TC-HT may reduce AT8 expression by activating the p-Akt/p-GSK-3β pathway,
thereby inhibiting its formation. Overall, TC-HT effectively down-regulates the
accumulation of ROT-induced PD markers through mechanisms involving both
chaperoning and prevention of protein aggregation, thus mitigating the stress
associated with their accumulation.

Figure 5. TC-HT promotes the neuroprotective mechanisms against the
accumulation of PD marker proteins. Representative western blots and the
quantifications of **(A)** Hsp70, **(B)** SIRT1, **(C)** p-Akt, and **(D)** p-GSK-3β (Ser9) after
each treatment. GAPDH was used as loading control. Data were presented as the
mean ± standard deviation in triplicate. Significance levels between indicated groups
are denoted as **$P < 0.01$, ***$P < 0.001$.

270 DISCUSSION

As the second most common NDD, PD is characterized by the loss of
dopaminergic neurons in the substantia nigra pars compacta, accompanied by the
extensive accumulation of Lewy bodies and tau fibrils, which consist of aggregated α-
syn and p-tau proteins [12, 13, 29]. Current primary treatments for PD are
pharmacological, but they provide only temporary symptomatic relief [10] and do not
address the underlying issue of protein aggregation in dysfunctional dopaminergic
neurons, making it challenging to halt PD progression. Therefore, heat application has
emerged as a promising alternative treatment, as it can potentially enhance
neuroprotective effects and bypass the challenges associated with crossing the blood-
brain barrier, enabling the direct targeting of affected brain regions. In this study, we
employ a modified heat treatment method, TC-HT, which offers enhanced safety and

superior neuroprotective effects compared to continuous HT, and investigate its
therapeutic potential in a ROT-induced *in vitro* PD model.

The accumulation of ROT, a widely used pesticide, can occur in humans
through the food chain. Studies have shown that ROT induces dopaminergic neuron
degeneration and Lewy body accumulation in the substantia nigra pars compacta in
mice [5], leading to PD-like motor dysfunction. Thus, ROT has become a widely used
PD inducer in both *in vitro* and *in vivo* studies. In this study, ROT was employed to
impair the dopaminergic-like neuroblastoma cell line SH-SY5Y to investigate the
neuroprotective effects and mechanisms of TC-HT. The results show that TC-HT
preserves the viability of SH-SY5Y cells under ROT exposure to a greater extent than
continuous HT (Figure 1), highlighting the stronger neuroprotective effects of TC-HT
against ROT.

Apoptosis, closely related to mitochondrial function, is a key mechanism of
cell death. Mitochondria are crucial for providing cellular energy and maintaining cell
viability. Studies have shown that ROT exposure could induce mitochondrial
dysfunction, leading to cell death [5]. However, it was found that moderate thermal
exposure could restore the impaired MMP [26]. Our results also demonstrate that TC-
HT significantly reduces apoptosis in ROT-injured SH-SY5Y cells, with superior
efficacy compared to continuous HT (Figure 2A). Additionally, JC-1 staining results
in Figures 2B & C suggest that TC-HT restores MMP more effectively in ROT-
treated SH-SY5Y cells than continuous HT. These results collectively highlight the
superior ability of TC-HT to reduce ROT-induced mitochondrial apoptosis compared
to continuous HT application in SH-SY5Y cells.

Besides apoptosis, impaired mitochondrial function also results in excessive
ROS production, which is detrimental to neural cells in PD [28]. Moreover, dopamine
metabolism results in ROS production in the neurons of the substantia nigra pars
compacta, making these neurons particularly susceptible to oxidative stress [7]. Hence,
controlling ROS levels is important for neuronal survival in PD. Studies have shown
that heat and ultrasound stimulations can alleviate intracellular oxidative stress by
activating the antioxidative mechanisms [26, 33, 34]. As shown in Figure 3A, our
study demonstrates that TC-HT effectively down-regulates ROT-elevated ROS levels
in SH-SY5Y cells, resulting in significantly higher antioxidative effects compared to
continuous HT. Notably, this reduction is accompanied by a significant increase in the
expression of the antioxidative enzyme SOD2 (Figure 3B). Therefore, TC-HT can
regulate intracellular ROS by up-regulating SOD2 and thereby mitigating ROT-
induced injury in SH-SY5Y cells.

The accumulation of α-syn and p-tau proteins contributes to damaged
mitochondria and excessive ROS production, which results in mitochondrial
dysfunction and dopaminergic neuron loss [29, 35, 36]. ROS overproduction can
further promote the aggregation of α-syn and p-tau proteins, exacerbating PD
progression [29, 37, 38]. Thus, the regulation of α-syn and p-tau expression levels is

critical for protecting neural cells in PD. As shown in Figure 4, TC-HT significantly
reduces the expression levels of α-syn and AT8 in ROT-damaged SH-SY5Y cells.
This reduction is significantly larger than that observed with continuous HT, and
aligns with the observed inhibitory effects on mitochondrial apoptosis (Figure 2) and
intracellular ROS levels (Figure 3). Hence, TC-HT demonstrates considerable
potential as a therapeutic approach for PD.

The expression levels of aggregated α-syn and p-tau can be modulated through
various mechanisms, with chaperoning playing a key role in preventing their
accumulation. Hsp70, an important heat-activated chaperone protein, helps to manage
misfolded proteins such as α-syn and p-tau, protecting neuronal cells from PD marker
accumulation and facilitating their degradation [30, 31]. Our results have shown that
TC-HT significantly increases the expression levels of Hsp70 and its upstream
transcriptional factor SIRT1 (Figures 5A, B), inducing stronger neuroprotective
effects on ROT-impaired SH-SY5Y cells compared to continuous HT and alleviating
the stress resulting from ROT-induced α-syn and p-tau accumulations more
effectively. Additionally, SIRT1 also acts as an upstream regulator of SOD2 [39].
Accompanied by the results in Figure 2, these findings suggest that TC-HT results in
stronger antioxidative effects than continuous HT against ROT-induced oxidative
stress by up-regulating SIRT1 more effectively. Besides chaperoning, preventing the
formation of aggregated proteins is crucial for inhibiting their pathological effects.
GSK-3β, a protein involved in tau hyperphosphorylation, can be inactivated by heat-
activated protein p-Akt through the phosphorylation of its Ser9 residue, which
reduces p-tau formation. As shown in Figures 5C & D, TC-HT significantly activates
greater p-Akt levels compared to continuous HT, consequently leading to the
inactivation of more GSK-3β through the increased expression of p-GSK-3β (Ser9).
These results, along with the reduced AT8 levels in Figure 4B, suggest that TC-HT
reduces p-tau protein accumulation by activating p-Akt, which inhibits p-tau
formation by decreasing p-GSK-3β (Ser9) activation. Collectively, TC-HT reduces
the expression levels of ROT-induced α-syn and p-tau more effectively than
continuous HT by robustly enhancing the neuroprotective mechanisms associated
with protein chaperoning and aggregation inhibition.

Physical stimulation is increasingly recognized as a potent therapeutic
approach for NDDs. Instead of continuous HT application, intermittent thermal
implementation can preserve neuroprotective effects while minimizing side effects
[26, 27]. On the other hand, it has been reported that a 40 Hz repetition frequency in
both light and US treatments might help to treat AD *in vivo* by decreasing Aβ
accumulation and attenuating neuroinflammation [40, 41], with phase I clinical trials
already underway [42]. Inspired by these findings, we integrated the 40 Hz US pulse
as a repetition frequency into our TC-HT treatment for a ROT-induced *in vitro* PD
model. Our results show that TC-HT with additional 40 Hz repetition US pulses also
exhibits a protective effect on the viability of SH-SY5Y cells against ROT, which is,
however, only slightly higher than that of mono TC-HT, and the difference is not

statistically significant (Figure 6). This suggests that further investigation of
optimized pulse repetition frequencies in physical stimulations could enhance
therapeutic efficacy, potentially expanding their applicability for treating NDDs.

Figure 6. The neuroprotective effect of TC-HT integrated with additional 40 Hz
repetition US pulses against ROT cytotoxicity. CCK-8 assay was conducted to
determine the viabilities of SH-SY5Y cells following mono TC-HT or TC-HT with a
40 Hz repetition US pulses. The viabilities were measured 24 h after ROT treatment.
Data were presented as the mean ± standard deviation in triplicate. Significance levels
between indicated groups are denoted as ***$P < 0.001$.

Heat therapy can facilitate the rescue of impaired neural cells, but continuous
heat exposure may lead to adverse effects. Our study shows that, with the same total
heat exposure time, TC-HT achieves significantly greater neuroprotective effects than
continuous HT in the ROT-induced *in vitro* PD model. This indicates that
interspersing low-temperature intervals during heat stimulation periods is crucial for
maximizing the neuroprotective effect of TC-HT. Particularly, FUS could serve as a
practical modality for delivering TC-HT to PD patients, providing programmable
control and precise, localized heating in the human brain in a non-invasive manner
[21, 22]. Besides, the heating area and temperature can be modulated via multi-focal
FUS, by adjusting the scanned spot size and intensity. This precise thermal regulation
allows efficient heat dissipation and helps achieve the desired temperature for TC-HT,
thereby enhancing its therapeutic potential. Therefore, FUS-based TC-HT represents a
feasible approach for safe and effective PD treatment.

In summary, this study presents TC-HT as a safer and more effective
hyperthermic method for protecting SH-SY5Y cells in an *in vitro* ROT-induced PD
model. TC-HT demonstrates superior protective effects in preserving ROT-impaired
cell viabilities and reducing apoptotic rates compared to continuous HT. Additionally,
TC-HT significantly reduces ROT-induced MMP loss and intracellular ROS
accumulation, while up-regulating the antioxidative enzyme SOD2. In comparison,
continuous HT shows only minimal effects on ROT-induced MMP loss and an
increase in ROS levels. Furthermore, TC-HT significantly outperforms continuous
HT in down-regulating PD marker proteins such as α-syn and p-tau. This enhanced
efficacy could be attributed to the substantial increase in the levels of Hsp70 and its
transcriptional regulator SIRT1, which together prevent the further accumulation of α-
syn and p-tau through effective chaperoning. Moreover, TC-HT results in the stronger
activation of Akt, leading to more effective inhibition of the hyperphosphorylation of
GSK-3β and thus resulting in a greater reduction in ROT-induced p-tau expression.
Collectively, our findings demonstrate the superior therapeutic potential of TC-HT
over HT for PD *in vitro*, highlighting its efficacy and safety as a hyperthermic
approach. Practically, programmable FUS could serve as an optimal heating source to
implement TC-HT as a treatment strategy for PD.

MATERIALS AND METHODS

Cell culture

The human neuroblastoma cell line SH-SY5Y was purchased from American Type Culture Collection (Manassas, VA, USA) and was cultured in Dulbecco's Modified Eagle Medium (DMEM) (HyClone; Cytiva, Marlborough, MA, USA) supplemented with 10% fetal bovine serum (FBS) (HyClone; Cytiva), 100 units/mL penicillin and 100 μg/mL streptomycin (Gibco Life Technologies, Grand Island, NY, USA). The cells were maintained in a humidified incubator with 5% carbon dioxide (CO_2) at a temperature of 37 °C. Once the confluence reached 80%, cells were subcultured by 0.05% trypsin–0.5 mM ethylenediamine tetraacetic acid (EDTA) solution (Gibco Life Technologies), and subsequently seeded into 96-well or 35-mm-diameter culture dishes (Thermo Fisher Scientific, Inc., Waltham, MA, USA) for *in vitro* experiments after 24 h incubation in the humidified 5% CO_2 incubator at 37 °C.

Ultrasound (US) exposure

The US exposure system consisted of a function generator (SG382; Stanford Research Systems, Sunnyvale, CA, USA), a power amplifier (25a250a; Amplifier Research, Souderton, PA, USA), and a 0.5 MHz ceramic planar US transducer (A101S-RM; Olympus NDT Inc., Waltham, MA, USA), as shown in Figure 7. The function generator produced continuous pulses with the following parameters: -10 dBm amplitude, 1 ms pulse period, and 0.5 ms pulse width. The cell culture plate or dish was positioned on the US transducer, with glycerol filling between the gap to transmit US waves into the culture plate or dish. The ceramic planar US transducer facilitated the conversion of electrical signals into acoustic and thermal energy, thereby administering both TC-HT and HT treatments to the SH-SY5Y cells.

Figure 7. Experimental setups for TC-HT and HT treatments. The US exposure system was composed of a function generator, a power amplifier, and a 0.5 MHz ceramic planar US transducer. The culture plate or dish was placed on the ceramic transducer, with glycerol utilized as the acoustic coupling medium to fill the gap. The transducer delivered acoustic and thermal energy to the SH-SY5Y cells for the administration of TC-HT and HT treatments.

TC-HT treatment

As described in our previous study [26, 27], the TC-HT treatment is a repetitive procedure comprising multiple cycles, each consisting of a designated high temperature stage followed by a cooling stage to achieve a series of short periods of

heat exposure over a specific duration. In the present study, the actual temperatures
experienced by SH-SY5Y cells were monitored using a K-type thermocouple in
conjunction with a temperature/process controller, and a 9-cycle repeated TC-HT was
implemented. The high temperature stage was sustained for 10 minutes during which
the maximum temperature was at 40 ± 0.5 °C with the US intensity set at 1.0 W,
followed by a natural thermal dissipation period of 1 minute for cooling. For the HT
application at the same US intensity, the temperature was continuously maintained at
40 ± 0.5 °C for a duration of 90 minutes. The monitored actual temperatures during
TC-HT or HT applications were depicted in Figure 8. In order to conduct TC-HT
utilizing US pulses at a frequency of 40 Hz, the modulation mode of the function
generator was set to pulse mode, characterized by a period of 25 ms and a pulse duty
cycle of 50%. The treatment of SH-SY5Y cells was performed at the same
temperature, maintaining the output US intensity at 1 W, with the maximum
temperature recorded at 40 ± 0.5 °C.

Figure 8. Actual temperatures recorded throughout the TC-HT and HT
exposure periods. The actual temperatures in the SH-SY5Y cells were measured
using a K-type needle thermocouple at 10 sec intervals throughout the exposure
durations of TC-HT and HT treatment.

Rotenone treatment

The SH-SY5Y cells were pretreated with TC-HT or HT at room temperature, and
subsequently maintained in the humidified 5% CO_2 incubator at 37 °C. At 4 h after
the thermal treatment, the pretreated SH-SY5Y cells were exposed to 5 μM rotenone,
which was purchased from Sigma-Aldrich (St. Louis, MO, USA) and dissolved in
dimethyl sulfoxide (DMSO) (Echo Chemical Co. Ltd., Miaoli, Taiwan) at a stock
concentration of 100 mM.

Cell viability assessment

Cell viability of the treated SH-SY5Y cells was evaluated using the Cell counting kit-
8 (CCK-8) assay (Dojindo, Kumamoto, Japan). Specifically, 10 μL CCK-8 solution
was added to each well containing either treated or untreated SH-SY5Y cells at 24 h
after ROT treatment, facilitating the formation of formazan from live cells. After a 3 h
incubation in the 5% CO_2 incubator at 37 °C, the absorbance of each well was
quantified using a Multiskan GO microplate spectrophotometer (Thermo Fisher
Scientific). The amount of formazan was determined by measuring the absorbance at
450 nm, with a background subtraction at 600 nm. Cell viability was expressed as a
percentage, with the viability of untreated control cells designated as 100%.

Apoptotic analysis via flow cytometry

The apoptotic rates of SH-SY5Y cells were assessed using Annexin V-Fluorescein
isothiocyanate (Annexin V-FITC) and propidium iodide (PI) double detection kit (BD
Biosciences, San Jose, CA, USA). At 24 h after ROT treatment, SH-SY5Y cells were
harvested and rinsed twice with ice-cold phosphate-buffered saline (PBS) (Hyclone).
The rinsed cells were then resuspended in a binding buffer containing Annexin V-
FITC and PI, followed by a 15-minute incubation at room temperature in the dark.
Apoptotic signals were measured by FACSCanto II system (BD Biosciences) with the
PE and FL1 channels.

Detection of ROS levels via flow cytometry

ROS levels of superoxide anion ($O_2^{\bullet-}$) were measured using fluorescent dye
dihydroethidium (DHE) (Sigma-Aldrich). At 24 h after ROT treatment, SH-SY5Y
cells were harvested and rinsed twice with PBS. The rinsed cells were then
resuspended in PBS containing 5 μM DHE, and incubated in the dark for 30 minutes
at 37 °C. ROS signals were detected using the FACSCanto II system with the PE
channel.

Mitochondrial membrane potential (MMP) analysis via flow cytometry

The loss of MMP was evaluated using the mitochondrial fluorescent dye 5,5′,6,6′-
Tetrachloro-1,1′,3,3′-tetraethylbenzimidazolylcarbocyanine   chloride   (JC-1)
(MedChemExpress Ltd., Monmouth Junction, NJ, USA). At 24 h after ROT treatment,
SH-SY5Y cells were harvested and rinsed with PBS before staining. The rinsed cells
were then resuspended in PBS containing 5 μg/mL JC-1 and incubated in the dark for
30 minutes at 37 °C. The fluorescent signals from JC-1 were measured using the
FACSCanto II system with the PE channel and FL1 channels. The levels of MMP
were determined by calculating the normalized fluorescence intensity ratio of PE/FL1
channel relative to the untreated control cells.

JC-1 fluorescence detection by laser scanning confocal microscopy

In order to observe fluorescent signals through microscopy, SH-SY5Y cells were
subjected to staining with JC-1 at a concentration of 5 μg/mL and incubated in the
dark for 30 minutes at 37 °C. A Zeiss LSM 880 inverted laser scanning confocal
microscope was employed to excite the red and green fluorescence of JC-1 at
wavelengths of 565 nm and 488 nm, respectively, facilitating the capture of
fluorescent images.

Western blot analysis

The protein expression levels in SH-SY5Y cells were analyzed using western blot
analysis. Following a wash with PBS, the cells were lysed in a lysis buffer (Millipore,
Billerica, MA, USA) containing 50 mM Tris-HCl (pH 7.4) and 0.15 M NaCl, which
was supplemented with an active protease/phosphatase inhibitor cocktail (#5827S;
Cell Signaling Technology, Danvers, MA, USA). After centrifugation at 23,000 × g
for 30 minutes at 4 °C, the supernatants were collected, and protein concentrations

were quantified using the Bradford protein assay (Bioshop, Inc.). Subsequently, 20 µg
of protein from each sample was separated by 10% SDS-polyacrylamide gel
electrophoresis (SDS-PAGE) and transferred onto polyvinylidene fluoride (PVDF)
membranes (Millipore). To block non-specific antibody binding, the membranes were
incubated with 5% bovine serum albumin in TBST (20 mM Tris-base, pH 7.6, 0.15 M
NaCl, 0.1% Tween 20 (Bioshop, Inc.)) for 1 h at room temperature. Primary
antibodies targeting superoxide dismutase 2 (SOD2), Hsp70, SIRT1, p-Akt (Cell
Signaling), α-syn, AT8 (p-tau), phosphorylated GSK-3β (serine 9) (p-GSK-3β (Ser9))
(Abcam), and glyceraldehyde-3-phosphate dehydrogenase (GAPDH) (Gentex, Irvine,
CA, USA) in blocking solution were subsequently applied to the membranes for
probing and incubated overnight at 4 °C. Following three washes with TBST, the
membranes were exposed to horseradish peroxidase-conjugated goat anti-rabbit
secondary antibody (Jackson ImmunoResearch Laboratories, West Grove, PA, USA)
in blocking solution at room temperature for 1 h. Immunoreactivity signals were
amplified by an enhanced chemiluminescence (ECL) substrate (Advansta, San Jose,
CA, USA) and visualized using an imaging system (Amersham Imager 600, GE
Healthcare Life Sciences). Images were analyzed with Image Lab software (Bio-Rad
Laboratories, Inc., Hercules, CA, USA). GAPDH was utilized as a loading control to
normalize the relative expression levels of target proteins.

Statistical analysis

All experiments were conducted in triplicate to ensure validation, and results were
expressed as mean ± standard deviation. Statistical analyses were performed using
one-way analysis of variance (ANOVA) with OriginPro 2015 software (OriginLab),
and results were deemed statistically significant when P-values were less than 0.05.

Author Contributions: Conceptualization, C.Y.C.; Methodology, Y.Y.K., G.B.L. and C.Y.C.; Validation, Y.Y.K., G.B.L., Y.M.C., H.H.L., F.T.H., and Y.K.; Formal Analysis, Y.Y.K., G.B.L., Y.M.C., H.H.L., F.T.H., Y.K., and C.Y.C.; Investigation, Y.Y.K., G.B.L. and C.Y.C.; Resources, C.Y.C.; Data Curation, Y.Y.K., G.B.L., Y.M.C., H.H.L., F.T.H., and Y.K.; Writing – Original Draft Preparation, Y.Y.K. and C.Y.C.; Writing – Review & Editing, Y.Y.K. and C.Y.C.; Supervision, C.Y.C.; Funding Acquisition, C.Y.C. All authors contributed to the article and approved the final manuscript.

Funding: This work was supported by grants from Ministry of Science and Technology (NSTC 112-2112-M-002-033, MOST 110-2112-M-002-004, and MOST 109-2112-M-002-004 to C.Y.C.) of the Republic of China. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: The data presented in this study are available on request from the corresponding author.

Conflicts of interest: The authors declare no conflicts of interest related to this study.

Abbreviations

AD	Alzheimer's disease
PD	Parkinson's disease
NDD	neurodegenerative disease
ROT	rotenone
ROS	reactive oxygen species
α-syn	α-synuclein
p-tau	phosphorylated-tau protein
HT	hyperthermia
Hsp70	heat-shock protein 70
p-Akt	phosphorylated Akt
GSK-3β	glycogen synthase kinase-3β
FUS	focused ultrasound
US	ultrasound
TC-HT	thermal cycling-hyperthermia

580	SIRT1	sirtuin-1
581	CCK-8	cell counting kit-8
582	$O_2^{\bullet-}$	superoxide anion
583	DHE	dihydroethidium
584	MMP	mitochondrial membrane potential
585	JC-1	5,5´,6,6´-Tetrachloro-1,1´,3,3´-
586		tetraethylbenzimidazolylcarbocyanine chloride
587	SOD2	superoxide dismutase
588	p-GSK-3β (Ser9)	phosphorylated GSK-3β (serine 9)
589	GAPDH	glyceraldehyde-3-phosphate dehydrogenase
590	ANOVA	analysis of variance.

References

1.   Prajjwal, P.; Flores Sanga, H.S.; Acharya, K.; Tango, T.; John, J.; Rodriguez,
R.S.C.; Dheyaa Marsool Marsool, M.; Sulaimanov, M.; Ahmed, A.; Hussin, O.A.
Parkinson's disease updates: addressing the pathophysiology, risk factors,
genetics, diagnosis, along with the medical and surgical treatment. *Ann. Med.*
Surg. (Lond). **2023**, 85, 4887-4902.
https://doi.org/10.1097/MS9.0000000000001142.

2.   Soilemezi, D.; Palmar-Santos, A.; Navarta-Sánchez, M.V.; Roberts, H. C.;
Pedraz-Marcos, A.; Haahr, A.; Sørensen, D.; Bragstad, L.K.; Hjelle, E.G.;
Haavaag, S.B.; et al. Understanding support systems for Parkinson's disease
management in community settings: a cross□national qualitative study *Health.*
Expect. **2023**, 26, 670-682. https://doi.org/10.1111/hex.13691.

3.   Frank, C.; Pari, G.; Rossiter, J.P. Approach to diagnosis of Parkinson disease.
Can. Fam. Physician. **2006**, 52, 862-868.

4.   Lee, J.; Huang, M.S.; Yang, I.C.; Lai, T.C.; Wang, J.L.; Pang, V.F.; Hsiao, M.;
Kuo, M.Y. Essential roles of caspases and their upstream regulators in rotenone-
induced apoptosis. *Biochem. Biophys. Res. Commun.* **2008**, 371, 33-38.
https://doi.org/10.1016/j.bbrc.2008.03.149.

5.   Watabe, M.; Nakaki, T. Rotenone induces apoptosis via activation of bad in
human dopaminergic SH-SY5Y cells. *J. Pharmacol. Exp. Ther.* **2004**, 311, 948-
953. https://doi.org/10.1124/jpet.104.071381.

6.   Li, N.; Ragheb, K.; Lawler, G.; Sturgis, J.; Rajwa, B.; Melendez, J.A.; Robinson,
J.P. Mitochondrial complex I inhibitor rotenone induces apoptosis through
enhancing mitochondrial reactive oxygen species production. *J. Biol. Chem.*
2003, 278, 8516-8525. https://doi.org/10.1074/jbc.M210432200.

7. Han, X.; Han, B.; Zhao, Y.; Li, G.; Wang, T.; He, J.; Du, W.; Cao, X.; Gan, J.;
Wang, Z.; et al. Rosmarinic acid attenuates rotenone-induced neurotoxicity in
SH-SY5Y Parkinson's disease cell model through Abl inhibition. *Nutrients.* **2022**,
14, 3508. https://doi.org/10.3390/nu14173508.

8. Kamal, R.E.; Menze, E.; Albohy, A.; Ahmed, H.I.; Azab, S.S. Neuroprotective
repositioning and anti-tau effect of carvedilol on rotenone induced neurotoxicity
in rats: Insights from an insilico & in vivo anti-Parkinson's disease study. *Eur. J.*
Pharmacol. **2022**, 932, 175204. https://doi.org/10.1016/j.ejphar.2022.175204.

9. Kempster, P.; Ma, A. Parkinson's disease, dopaminergic drugs and the plant
world. *Front.   Pharmacol.   * **2022**,   13,   970714.
https://doi.org/10.3389/fphar.2022.970714.

10. Zahoor, I.; Shafi, A.; Haq, E. Pharmacological treatment of Parkinson's disease.
In *Parkinson's Disease: Pathogenesis and Clinical Aspects [Internet]*; Stoker,
T.B. Stoker, Greenland J.C., Eds.; Publisher: Brisbane, Australia, 2018, pp 129-
144, ISBN 978-0-9944381-6-4.

11. Cheng, G.; Liu, Y.; Ma, R.; Cheng, G.; Guan, Y.; Chen, X.; Wu, Z.; Chen, T.
Anti-parkinsonian therapy: strategies for crossing the blood-brain barrier and
nano-biological effects of nanomaterials. *Nanomicro. Lett.* **2022**, 14, 105.
https://doi.org/10.1007/s40820-022-00847-z.

12. Srinivasan, E.; Chandrasekhar, G.; Chandrasekar, P.; Anbarasu, K.; Vickram,
A.S.; Karunakaran, R.; Rajasekaran, R.; Srikumar, P.S. Alpha-synuclein
aggregation in Parkinson's disease. *Front. Med. (Lausanne).* **2021**, 8, 736978.
https://doi.org/10.3389/fmed.2021.736978.

13. Pan, L.; Meng, L.; He, M.; Zhang, Z. Tau in the pathophysiology of Parkinson's
disease. *J. Mol. Neurosci.* **2021**, 71, 2179-2191. https://doi.org/10.1007/s12031-
020-01776-5.

14. Habash, R.W.; Bansal, R.; Krewski, D.; Alhafid, H.T. Thermal therapy, part 1: an
introduction to thermal therapy. *Crit. Rev. Biomed. Eng.* **2006**, 34, 459-489.
https://doi.org/10.1615/critrevbiomedeng.v34.i6.20.

15. Banerjee Mustafi, S.; Chakraborty, P.K.; Dey, R.S.; Raha, S. Heat stress
upregulates chaperone heat shock protein 70 and antioxidant manganese
superoxide dismutase through reactive oxygen species (ROS), p38MAPK, and
Akt.   *Cell.   Stress.   Chaperones.*   **2009**,   14,   579-589.
https://doi.org/10.1007/s12192-009-0109-x.

16. Tundo, G.R.; Sbardella, D.; Ciaccio, C.; Bianculli, A.; Orlandi, A.; Desimio,
M.G.; Arcuri, G.; Coletta, M.; Marini, S. Insulin-degrading enzyme (IDE): a
novel heat shock-like protein. *J. Biol. Chem.* **2013**, 288, 2281-2289.
https://doi.org/10.1074/jbc.M112.393108.

17. Sato, K.; Saito, H.; Matsuki, N. HSP70 is essential to the neuroprotective effect
of heat-shock *Brain. Res.* **1996**, 740, 117-123. https://doi.org/10.1016/s0006-
8993(96)00846-3.

18. Cao, L.; Cao, D.X.; Su, X.W.; Chen, L.J.; Liu, A.L.; Jian, W.J.; Wu, Y.P.; Qiu,

P.X.; Yan, G.M. Activation of PI3-K/Akt pathway for thermal preconditioning to
protect cultured cerebellar granule neurons against low potassium-induced
apoptosis. *Acta. Pharmacol. Sin.* **2007**, 28, 173-179.
https://doi.org/10.1111/j.1745-7254.2007.00504.x.

19. Sousa, L.; Guarda, M.; Meneses, M.J.; Macedo, M.P.; Vicente Miranda, H.
Insulin-degrading enzyme: an ally against metabolic and neurodegenerative
diseases. *J. Pathol.* **2021**, 255, 346-361. https://doi.org/10.1002/path.5777.

20. Noel, A.; Barrier, L.; Ingrand, S. The Tyr216 phosphorylated form of GSK3β
contributes to tau phosphorylation at PHF-1 epitope in response to Aβ in the
nucleus of SH-SY5Y cells. *Life. Sci.* **2016**, 158, 14-21.
https://doi.org/10.1016/j.lfs.2016.06.020.

21. Partanen, A.; Tillander, M.; Yarmolenko, P.S.; Wood, B.J.; Dreher, M.R.; Kohler,
M.O. Reduction of peak acoustic pressure and shaping of heated region by use of
multifoci sonications in MR-guided high-intensity focused ultrasound mediated
mild hyperthermia. *Med. Phys.* **2013**, 40, 013301.
https://doi.org/10.1118/1.4769116.

22. Chaplin, V.; Caskey, C.F. Multi-focal HIFU reduces cavitation in mild-
hyperthermia. *J. Ther. Ultrasound.* **2017**, 5, 12. https://doi.org/10.1186/s40349-
017-0089-8.

23. Lee, J.S.; Choi, J.C.; Kang, S.Y.; Kang, J.H.; Park, J.K. Heat stroke: increased
signal intensity in the bilateral cerebellar dentate nuclei and splenium on
diffusion-weighted MR imaging. *AJNR. Am. J. Neuroradiol.* **2009**, 30, E58.
https://doi.org/10.3174/ajnr.A1432.

24. Tabassum, S.; Raza, N.; Shah, S.Z. Outcome of heat stroke patients referred to a
tertiary hospital in Pakistan: a retrospective study. *East. Mediterr. Health. J.*
2019, 25, 457-464. https://doi.org/10.26719/emhj.18.059.

25. Cremer, O.L.; Kalkman, C.J. Cerebral pathophysiology and clinical neurology of
hyperthermia in humans. *Prog. Brain. Res.* **2007**, 162, 153-169.
https://doi.org/10.1016/S0079-6123(06)62009-8.

26. Chen, W.T.; Kuo, Y.Y.; Lin, G.B.; Lu, C.H.; Hsu, H.P.; Sun, Y.K.; Chao, C.Y.
Thermal cycling protects SH-SY5Y cells against hydrogen peroxide and β-
amyloid-induced cell injury through stress response mechanisms involving Akt
pathway. *PLoS One.* **2020**, 15, e0240022.
https://doi.org/10.1371/journal.pone.0240022.

27. Kuo, Y.Y.; Chen, W.T.; Lin, G.B.; Chen, Y.M.; Liu, H.H.; Chao, C.Y. Thermal
cycling-hyperthermia ameliorates Aβ25-35-induced cognitive impairment in
C57BL/6 mice. *Neurosci. Lett.* **2023**, 810, 137337.
https://doi.org/10.1016/j.neulet.2023.137337.

28. Guo, C.; Sun, L.; Chen, X.; Zhang, D. Oxidative stress, mitochondrial damage
and neurodegenerative diseases. *Neural. Regen. Res.* **2013**, 8, 2003-2014.
https://doi.org/10.3969/j.issn.1673-5374.2013.21.009.

29. Haque, M.M.; Murale, D.P.; Kim, Y.K.; Lee, J.S. Crosstalk between oxidative

stress and tauopathy. *Int. J. Mol. Sci.* **2019**, 20, 1959.
https://doi.org/10.3390/ijms20081959.

30. Tao, J.; Berthet, A.; Citron, Y.R.; Tsiolaki, P.L.; Stanley, R.; Gestwicki, J.E.;
Agard, D.A.; McConlogue, L. Hsp70 chaperone blocks α-synuclein oligomer
formation via a novel engagement mechanism. *J. Biol. Chem.* **2021**, 296, 100613.
https://doi.org/10.1016/j.jbc.2021.100613.

31. Patterson, K.R.; Ward, S.M.; Combs, B.; Voss, K.; Kanaan, N.M.; Morfini, G.;
Brady, S.T.; Gamblin, T.C.; Binder, L.I. Heat shock protein 70 prevents both tau
aggregation and the inhibitory effects of preexisting tau aggregates on fast
axonal transport. *Biochemistry.* **2011**, 50, 10300-10310.
https://doi.org/10.1021/bi2009147.

32. Raynes, R.; Brunquell, J.; Westerheide, S.D. Stress inducibility of SIRT1 and its
role in cytoprotection and cancer. *Genes. Cancer.* **2013**, 4, 172–182.
https://doi.org/ 10.1177/1947601913484497.

33. Cao, Q.; Liu, L.; Hu, Y.; Cao, S.; Tan, T.; Huang, X.; Deng, Q.; Chen, J.; Guo, R.;
Zhou, Q. Low-intensity pulsed ultrasound of different intensities differently
affects myocardial ischemia/reperfusion injury by modulating cardiac oxidative
stress and inflammatory reaction. *Front. Immunol.* **2023**, 14, 1248056.
https://doi.org/10.3389/fimmu.2023.1248056.

34. Aimaijiang, M.; Liu, Y.; Zhang, Z.; Qin, Q.; Liu, M.; Abulikemu , P.; Liu,
L.; Zhou, Y. LIPUS as a potential strategy for periodontitis treatment: A review
of the mechanisms. *Front. Bioeng. Biotechnol.* **2023**, 11, 1018012.
https://doi.org/10.3389/fbioe.2023.1018012.

35. Riederer, P.; Nagatsu, T.; Youdim, M.B.H.; Wulf, M.; Dijkstra, J.M.; Sian-
Huelsmann, J. Lewy bodies, iron, inflammation and neuromelanin: pathological
aspects underlying Parkinson's disease. *J. Neural. Transm. (Vienna).* **2023**, 130,
627-646. https://doi.org/10.1007/s00702-023-02630-9.

36. Deas, E.; Cremades, N.; Angelova, P.R.; Ludtmann, M.H.; Yao, Z.; Chen, S.;
Horrocks, M.H.; Banushi, B.; Little, D.; Devine, M.J.; et al. Alpha-synuclein
oligomers interact with metal ions to induce oxidative stress and neuronal death
in Parkinson's disease. *Antioxid. Redox. Signal.* **2016**, 24, 376-391.
https://doi.org/10.1089/ars.2015.6343.

37. Puspita, L.; Chung, S.Y.; Shim, J.W. Oxidative stress and cellular pathologies in
Parkinson's disease. *Mol. Brain.* **2017**, 10, 53. https://doi.org/10.1186/s13041-
017-0340-9.

38. Dorszewska, J.; Kowalska, M.; Prendecki, M.; Piekut, T.; Kozłowska, J.;
Kozubski, W. Oxidative stress factors in Parkinson's disease. *Neural. Regen. Res.*
2021, 16, 1383-1391. https://doi.org/10.4103/1673-5374.300980.

39. Singh, C.K.; Chhabra, G.; Ndiaye, M.A.; Garcia-Peterson, L.M.; Mack, N.J.;
Ahmad, N. The role of sirtuins in antioxidant and redox signaling. *Antioxid.*
Redox. Signal. **2018**, 28, 643-661. https://doi.org/10.1089/ars.2017.7290.

40. Martorell, A.J.; Paulson, A.L.; Suk, H.J.; Abdurrob, F.; Drummond, G.T.; Guan,

W. Multi-sensory gamma stimulation ameliorates Alzheimer's-associated
pathology and improves cognition. *Cell.* **2019**, 177, 256-271.
https://doi.org/10.1016/j.cell.2019.02.014.

41. Park, S.S.; Park, H.S.; Kim, C.J.; Kang, H.S.; Kim, D.H.; Baek, S.S.; Kim, T.W.
Physical exercise during exposure to 40-Hz light flicker improves cognitive
functions in the 3xTg mouse model of Alzheimer's disease. *Alzheimers Res. Ther.*
2020, 12, 62. https://doi.org/10.1186/s13195-020-00631-4.

42. Chan, D.; Suk, H.J.; Jackson, B.L.; Milman, N.P.; Stark, D.; Klerman, E.B.;
Kitchener, E.; Fernandez Avalos, V.S.; de Weck, G.; Banerjee, A.; et al. Gamma
frequency sensory stimulation in mild probable Alzheimer's dementia patients:
Results of feasibility and pilot studies. *PLoS One.* **2022**, 17, e0278412.
https://doi.org/10.1371/journal.pone.0278412.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Figure 8