Persistent viral infections impact key biological traits in *Drosophila melanogaster*

Mauro Castelló-Sanjuán[1], Rubén González[1], Ottavia Romoli[1], Hervé Blanc[1], Jared Nigg[1, #],

Maria-Carla Saleh[1].

[1]Institut Pasteur, Université Paris Cité, Viruses and RNA Interference Unit, Paris, France.

[#]Current address: Chan Zuckerberg Biohub, San Francisco, California, USA.

Correspondence: jared.nigg@czbiohub.org, carla.saleh@pasteur.fr

Short title: Consequences of persistent viral infections

ABSTRACT

Persistent viral infections have been assumed to have minimal fitness costs for insects.

Using *Drosophila melanogaster*, we established persistent mono-infections with four

different enteric RNA viruses (Drosophila A virus, Drosophila C virus, Bloomfield virus,

and Nora virus) and observed that these infections significantly reduce fly survival, alter

the number of viable offspring per female, modulate microbiome composition, and

impact locomotor abilities. These results demonstrate significant impacts of persistent

viral infections on key biological traits and challenge the assumption of the innocuity of

persistent viral infections for the host. In addition, the four viruses displayed different

accumulation kinetics and elicited unique transcriptional profiles with no common core

responses. The transcriptional changes triggered by DCV infection persisted even after

viral clearance. Our findings reveal that persistent viral infections modulate critical

aspects of insect biology and can have negative impacts on host physiology and

behavior.

INTRODUCTION

The increasing availability of sequencing data has revealed tens of thousands of novel viruses in multiple host organisms, suggesting the widespread occurrence of previously undetected infections (Ho & Tzanetakis, 2014; Kapun *et al.,* 2020; Wallace *et al.,* 2021). In some cases, these are persistent infections that represent a metastable state in which the physiological impacts of viral infection are maintained below a lethal threshold without leading to viral clearance (Goic & Saleh 2012). In contrast to acute viral infections, which are characterized by rapid viral replication that is ultimately limited by the death of the host or clearance of the virus, persistent infections endure over extended periods within the individual host. During this time, the virus is still capable of being transmitted to other organisms or the host's offspring despite the inherent limitations imposed by the host's immune responses and cellular mortality (Goic & Saleh 2012).

Persistence is a frequent phenomenon in viral infections across all kingdoms, as it can provide advantages for both the virus and the host, acting as a modulator of the ecosystem. For example, livestock and crops often suffer from infections caused by harmful or deadly viruses (Goic & Saleh 2012). Growing evidence shows that persistent infections have costs to the host (Nigg et al. 2024, Wallace and Obbard, 2024). Nevertheless, research has mostly focused on the acute effects of viral infections. Consequently, the impacts of persistent infections on host physiology and behavior remain understudied.

In this study, we aimed to shed light on the consequences of persistent infections using the fruit fly, *Drosophila melanogaster*. This insect has an extensively characterized genome and short generation times, making it an ideal model organism for studying persistent viral infections throughout its lifespan. In a previous study, we demonstrated

that persistent infections with natural viruses are associated with decreased lifespan in

D. melanogaster (Nigg *et al.,* 2024). In the present study, we aimed to further examine the

effects of persistent mono-infections with four distinct enteric RNA viruses—Drosophila

A Virus (DAV), Drosophila C Virus (DCV), Bloomfield virus, and Nora virus —on a wide

range of physiological traits in *D. melanogaster*. Specifically, we focused on survival

rates, fertility (reproductive output, here defined as the number of viable offspring),

bacteriome composition, and locomotor abilities. These traits are essential for

understanding the host's overall fitness and its ecological interactions. Survival rates and

reproductive output serve as key indicators of an organism's ability to thrive and

contribute to the next generation (Crone, 2001). The composition of the microbiome has

been observed to influence various aspects of the host's physiology, such as nutrition

status, immune function, behavior, and overall health (Gould *et al.,* 2018; Jia *et al.,* 2021;

Shin *et al.,* 2011; Tafesh-Edwards & Eleftherianos, 2023). Locomotor abilities are vital for

behaviors such as foraging, mating, and avoiding predators, all of which are crucial for

survival and reproduction (Irschick and Garland, 2001). By examining these traits, we

aimed to elucidate the broader implications of persistent viral infections on insect

biology.

RESULTS

The dynamics of viral RNA levels over time are virus specific

Our laboratory has previously established persistent viral mono-infections in *D.*

melanogaster flies of the w^{1118} genetic background using different viruses: DAV, DCV,

Bloomfield virus, and Nora virus (Nigg *et al.,* 2022, 2024). All these viruses naturally infect

D. melanogaster and are transmitted primarily through orofecal contamination. In this

work, we measured key biological traits over time in the infected populations and an uninfected control population (Figure 1A). To maintain continuous exposure to contaminated food while preventing excessive viral accumulation, flies from the different persistently infected lines were transferred to new food every other day. Persistent viral infections significantly reduce the survival of infected flies, as previously described (Figure 1B; Nigg *et al.,* 2024). DAV infection has the most pronounced effects on survival, followed by DCV, Bloomfield virus, and Nora virus (Mantel-Cox test infected vs uninfected $P < 0.001$ for all the virus-infected populations). This is reflected by the time at which 50% of the population has perished, the median survival (Figure 1B): 25 days post-eclosion (dpe) for DAV, 28 dpe for DCV, 36 dpe for Bloomfield virus, and 52 dpe for Nora virus, compared to 63 dpe for uninfected flies. This information enabled us to establish different time points for further characterization of the infections: 1 dpe as an early time point, 12 dpe as an intermediate time point just before the onset of virus-induced mortality and the median survival as a late time point specific for each condition. Additionally, we added an additional time point for DCV at 6 dpe, as mortality starts earlier for this virus. In the present study, these time points served as the basis for characterizing the impact of persistent viral infections on the physiology and behavior of infected hosts (Figure 1A).

To characterize the dynamics of viral load within the infected populations, we examined viral RNA levels in individual flies at the aforementioned early, intermediate, and late stages of infection. Viral loads were quantified by RT-qPCR by measuring viral RNA levels relative to the Drosophila housekeeping gene *Rp49*. Flies were classified as infected if their Ct values for viral RNA were below 35 with melting temperature profiles indicating the presence of a single amplicon. We observed distinct patterns of viral

accumulation for each virus (Figure 1C, upper panels). DAV exhibited a rapid increase in viral RNA levels early post-eclosion, with a median increasing by ~4 and ~5 $\log_{10}$ orders of magnitude at 12 and 25 dpe, respectively. In contrast, DCV displayed a relatively stable pattern in viral RNA levels over time, with no significant changes in viral loads. Interestingly, we noted that DCV-infected individuals stably clustered into two distinct groups, displaying viral RNA levels several orders of magnitude apart. Bloomfield virus RNA levels rose steadily over the lifetime of infected individuals, increasing by 1 and 2 $\log_{10}$ orders of magnitude at 12 and 36 dpe, respectively. In contrast, Nora virus RNA levels remained consistent at the earlier timepoint but rose significantly by ~3 $\log_{10}$ orders of magnitude by the final timepoint (GLM, letters over the conditions indicate grouping for multiple comparisons).

Our examination of viral RNA levels allowed us to estimate infection prevalence in the different populations over time (Figure 1C, lower panels). A viral infection is considered prevalent in a population if individuals continue to be infected over time, even as time progresses. DAV and Nora virus were consistently detected in 100% of individuals, except for a slight reduction in Nora virus prevalence at the later stage (~91% of individuals infected at 52 dpe). Bloomfield virus was highly prevalent in the population at all time points, although not all individuals were infected at any given time, and infection prevalence varied from ~91% to ~72%. DCV was detected in all individuals at 1 dpe, and its prevalence in the population progressively decreased over time, with only ~36% of the individuals infected at the latest time point.

Persistent viral infections impact viable offspring production

Persistent viral infections can reduce individual survival rates, which may consequently affect the viability of the population over time. To understand the broader impact of these infections, we investigated whether populations with a prevalent viral infection affect the production of viable offspring, which we define as the number of progeny that successfully emerge as adults. This metric is a crucial aspect of reproductive success and population dynamics. Any impact on the production of viable offspring would indicate that persistent infections influence additional critical biological traits related to fitness beyond the previously observed reduction in survival.

To explore this, we selected 20 females and 20 males from each infected and uninfected stock on the day of eclosion (0 dpe). Flies were allowed to mate for 24 hours, after which the males were discarded. The females were then transferred into new vials with food each day, and the viable offspring were counted as they emerged. The final progeny number was counted and normalized based on the number of female flies laying eggs in each vial.

Flies collected from DAV, DCV, Bloomfield virus, or Nora virus-infected populations exhibited distinct patterns of offspring production compared to uninfected controls over time (Figure 2A). Uninfected flies exhibited a peak in viable offspring production at 1 dpe with 10.3 ± 2.02 (mean ± SD) adults emerging per female. This was followed by a gradual decrease in offspring production until 6 dpe, after which the decline became more pronounced, ultimately reaching zero by 16 dpe. Similarly, populations infected with DAV and Bloomfield viruses also peaked in viable offspring production at 1 dpe, with averages of 11.5 ± 2.07 and 11.6 ± 1.30, respectively. These infected populations also experienced a sharp decline in offspring production after 6 dpe. However, the pattern of viable

offspring per female daily indicates that, at the early time points, DAV and Bloomfield

virus-infected populations produced more viable offspring per female than the

uninfected control population (GLMM, DAV P = 0.0223, Bloomfield P = 0.0309). In a

different way, the DCV-exposed flies displayed a unique pattern in viable offspring

production, surpassing the uninfected group only at later time points (GLMM, P < 0.0001).

In contrast, Nora virus-infected flies consistently produced fewer viable offspring

(GLMM, P = 0.0098), peaking at 4 dpe with an average of 5.28 ± 2.15 viable offspring

produced.

Studying both daily and cumulative offspring production offers a comprehensive

perspective on reproductive dynamics. Daily offspring production highlights short-term

effects and variability, while cumulative offspring production indicates overall

reproductive success throughout the female's lifespan.

When examining the cumulative offspring per female, we see the total offspring

production from infected and uninfected stocks over time. We found that DCV-infected

flies produced more viable offspring (91.5 ± 32.1) than the uninfected group (67.7 ± 11.6)

(Figure 2B, GLMM, P = 0.0044). We also noted that Nora virus-infected flies produced

significantly fewer offspring per female (41.5 ± 15.5, GLMM, P = 0.0335). Overall, the

results illustrate the diverse and virus-specific impacts of persistent viral infections on

fertility.

Persistent viral infections alter locomotor function in a virus-dependent manner

Locomotor function is a key indicator of an organism's overall health and physiological

status. In flies, locomotion reflects both muscular and neurological capacity, making it a

valuable marker for assessing the general impact of viral infections on host physiology.

We assessed locomotor function using the reverse geotaxis assay, which measures the flies' upward climbing instinct over a specified period (Gargano *et al.,* 2005). The assay was performed on 20 mated females from different uninfected and infected stocks at 1, 6, 12, and 20 dpe. This final time point was chosen as a common late date for all fly stocks, occurring prior to the day of median survival for all of the infected populations. Flies were placed in an empty tube and collected at the bottom by gently tapping the tubes. After waiting for 3 seconds, a picture was taken, and the position of each fly was recorded manually. Our observations indicated that persistent viral infections had a mixed effect on locomotor function over time (Figure 3). DCV- and Nora virus-infected flies consistently showed impaired locomotion abilities compared to uninfected flies at almost all the tested time points (Figure 3A). In contrast, DAV- and Bloomfield-infected flies showed mixed locomotion patterns. DAV significantly reduced mobility at 1 day post-exposure (dpe) but significantly increased it at 6 and 20 dpe (GLMM, $P < 0.0001$, $P < 0.0001$, and $P = 0.0108$, respectively). Bloomfield virus significantly reduced mobility at 1 dpe and increased it at 12 dpe (GLMM, $P < 0.0001$ at both time points). When examining the fly distribution patterns along the tubes we obtained similar results, with a higher proportion of DCV- and Nora virus-infected flies found in the bottom of the tube at almost all time points (Figure 3B). DAV infection resulted in a higher proportion of flies climbing to the top of the tube at 6 dpe, but a higher proportion of flies at the bottom at 12 dpe (CLMM, $P = 0.0008$ and $P = 0.0013$). Bloomfield virus-infected flies were found more often on the bottom of the tube at 1 and 20 dpe, while they climbed more to the top at 12 dpe (CLMM, $P < 0.0001$, $P = 0.0028$, and $P = 0.0322$, respectively). Taken together, these results indicate that persistent viral infections affect the climbing abilities of infected flies, with a general trend of lower climbing height, particularly in flies harboring Nora

virus infections. These findings underscore the systemic impact of persistent viral

infections on host physiology and behavior.

Harboring persistent viral infections changes fly bacteriome composition

We next investigated whether persistent viral infections influence the fly bacteriome—

the bacteria associated with the host. All fly populations in this study shared the same

genetic background and were reared under identical conditions, except for the presence

of different persistent viral infections. This controlled setup allowed us to attribute any

observed differences in the associated bacteria to the presence of different viral

infections.

Flies were collected at 12 dpe, surface sterilized and their DNA was extracted.

Microbiota composition was assessed by amplification and sequencing of the V3-V4

region of the *16S* rRNA bacterial gene. Although the microbiota of our flies was generally

dominated by *Lactobacillus* species, we detected changes in the relative bacterial

composition of infected flies compared to the uninfected control population.

Specifically, DAV- and DCV-infected flies showed an increase in the abundance of

Lactobacillus, while Nora virus infection increased the relative abundance of

Sphingomonas and *Thepidiphilus* genera (Figure 4). We are currently conducting

additional replicates and including Bloomfield virus in our analysis to further elucidate

the bacteriome changes induced by persistent viral infections. These ongoing studies will

provide a more comprehensive understanding of how different viruses influence the gut

microbial community and the potential implications for host physiology and fitness.

Host transcriptional response is virus specific

We observed that persistent infections are associated with modulations of the life history traits discussed above. To gain insight into the host response to persistent viral infections, we studied the transcriptional response of flies at early, intermediate, and late stages of infection. For each time point and condition, we prepared four pools consisting of RNA extracted from three to five individuals. These pools were prepared using the same RNA samples used for the RT-qPCR analysis shown in Figure 1C. Informed by the RT-qPCR data, we excluded non-infected individuals and randomly selected RNA from infected individuals to prepare RNA pools for sequencing.

Observing the general transcriptomic profile of each condition, we found that each enteric virus elicited a distinct transcriptional response in the host. Nora virus infection was characterized by the most unique transcriptional response at 1 dpe (Supplementary Figure 1), with the differential regulation of several genes (Supplementary Figure 2, 3). On the contrary, at the same early time point other viruses induced little changes in transcriptional profiles of infected flies and samples clustered together with uninfected flies (Supplementary Figures 1-3). At a more advanced time point (12 dpe) the transcriptional profiles of DCV and Bloomfield virus-infected flies clustered together with uninfected flies, while Nora virus and DAV-infected flies were characterized by unique transcriptional responses (Supplementary Figure 1). At the days of median survival, all viruses induced significant changes in transcriptional profiles (Supplementary Figure 1). Among viral infections at the same time points, no specific differentially expressed gene (DEG) could be identified as a common factor across all infections (Supplementary Figure 4). Furthermore, during a persistent infection, we could not identify a consistent set of host genes that respond to the infection over time, thereby illustrating its

differential impacts as the infections progress (Supplementary Figure 4). Even when

comparing the transcriptomic profiles at the days of median survival, no common

transcriptional profile was shared among all infections (Supplementary Figure 4). We

also observed an increase in the number of DEGs as infection progressed for the flies

persistently infected with DAV, DCV, and Bloomfield virus (Supplementary Figures 3-4).

Many of the top DEGs identified within the diverse transcriptomic profiles at 1 dpe and 12

dpe are uncharacterized genes (Supplementary Figure 3).

Focusing on the genes known to participate in antiviral immune responses

(Supplementary File 1), Nora virus elicited the highest levels of upregulation and

downregulation at 1 dpe (Figure 5A). At 12 dpe, among the top twenty differentially

expressed genes (DEGs), DAV had the most genes upregulated. We found that DAV

infection resulted in the upregulation of genes associated with immune defense and

stress response pathways, indicating a potent antiviral response (Figure 5A). Conversely,

specific immune genes that appear upregulated during DAV infection are downregulated

during Nora virus infection. This applies to *PGRP-SC2*, the *Peptidoglycan Recognition*

Protein SC2 gene, a negative regulator of the IMD pathway, and *PGRP-SC1a*, involved in

the Toll pathway signaling, as well as to *PGRP-SA* and *GNBP-like3,* involved in the Toll

pathway. This suggests that, although the immune response to these viruses may involve

some shared genes, it is fundamentally different, if not opposing (Figure 5A). Compared

to the other viruses, persistent DCV infection triggered the differential expression of a

modest set of immune genes; notably, the initial downregulation of *PGRP-SC1a* at 1 dpe

and a modest upregulation of immune genes as the infection progressed. Bloomfield

virus-infected flies also displayed initial downregulation of *PGRP-SC1a* at 1 dpe, followed

by an upregulation of *pirk* (IMD pathway) and *PGRP-SA* (Toll) (Figure 5A).

Upon analyzing the transcriptomic profiles of genes described to influence

locomotion (Supplementary File 1), we found that these genes were mostly notably

differentially expressed in Nora virus-infected flies at 1 dpe. Notably, *Lsp2,* a gene

reported to be involved in motor neuron axon and synaptic target inhibition (Inaki *et al.,*

2007), was upregulated in Nora virus-infected samples (Figure 5B). As infection

progresses, we observed an increasing number of DEGs related to locomotion, with Nora

virus infection showing an opposite transcriptional profile compared to DAV infection

(Figure 5B). At 12 dpe, Nora virus infection significantly upregulated *CG3857*, *Ptth*, and

Rh5, genes described to play roles in gravitaxis regulation, negative phototaxis, and light

reception, respectively (Armstrong *et al.,* 2006; Yamanaka *et al.,* 2013; Sprecher and

Desplan, 2008). In the same samples, we also observed an upregulation of genes related

to neuronal function, such as *nompC* and *Pdf* (Figure 5B) (Renn *et al.,* 1999; Walker *et al.,*

2000). Furthermore, Nora virus infection induced the upregulation of *M6*, a gene involved

in several neurological processes, including positive phototaxis and the downregulation

of *fz*, related to neuronal function, and *Neto*, a gene reported to not only have

neuromuscular roles but also to modulate hatching behavior (Kim *et al.,* 2012).

Conversely, for the DAV-infected samples, we observed an upregulation of *Neto*, along

with other genes like *esg* (related to stem cell maintenance and nervous system

development ; Bahrampour *et al.,* 2017; Biteau *et al.,* 2008; Hartl *et al.,* 2011) and *Mlc2*

(involved in flight ; Miller *et al.,* 2011) (Figure 5B). For the DCV-, and Bloomfield virus-

infected samples, we detected modest differential expression of locomotion genes.

Taken together, these results indicate that the host response to persistent infection is

virus specific.

Viral infection induced transcriptional changes even after viral clearance

Persistent viral infections, even if prevalent in the population and persistent in most individuals, may still be cleared over time by the host (Figure 1B; Mondote *et al.,* 2018; Nigg *et al.,* 2022). To investigate the long-term effects of viral clearance on the host, we analyzed the transcriptome of flies that cleared the DCV infection (Figure 6). In fact, in the DCV-infected population all individuals tested at 1 dpe were infected (Figure 1C), but more than half of the population had cleared the infection by 28 dpe (Figure 1C). Therefore, we analyzed the transcriptional response of flies that had cleared DCV infection by 28 dpe and found that it remained significantly altered compared to the never-infected control population (Figure 6A). We further explored the differences in the transcriptomic profiles of DCV-infected and DCV-cleared flies. These results indicate that only a few genes are differentially expressed between the two sets of samples (Figure 6B). Most of these genes were mitochondrial transfer RNAs, or genes encoding for mitochondrial proteins, such as a subunit of the mitochondrial ATPase (*mt:ATPase8*), a protein belonging to the cytochrome c oxidase assembly (*l(3)neo43*), a protein involved in mitochondrial fatty acid beta-oxidation (*Mcad*), a mitochondrial single stranded DNA-binding protein (*mtSSB*), a protein part of respiratory chain complex (*ND-MWFE*), a carrier protein of the inner mitochondrial membrane (*sea*). The transcriptomic profiles of the two sample groups showed significant changes when compared to the never-infected control population. Specifically, genes related to the immune response continued to be upregulated (Figure 6D). These results suggest that the host transcriptome does not completely revert to its pre-infection state even after viral clearance, indicating lingering effects on host physiology and immune function.

DISCUSSION

Our study provides a comprehensive examination of the impacts of persistent viral infection on key biological traits in *D. melanogaster*. By studying mono-infections with four different enteric RNA viruses, we reveal a complex interplay between viruses and their hosts that affects survival, reproductive output (fertility), microbiota composition, locomotor abilities, and transcriptional profiles.

We observed a trade-off between survival and reproductive output. This trade-off suggests a compensatory mechanism in which, when faced with specific stressors such as viral infections, the host organism may prioritize reproduction over immunity to ensure population survival (Pontzer and McGrosky, 2022; Pagán *et al.,* 2008). Indeed, for DCV, DAV, and Bloomfield viruses, we noted reduced survival but increased offspring production in infected flies, indicating a redirection of resources from surviving the infection to reproduction. Conversely, for Nora virus infections, resources appear to be redirected primarily toward surviving the infection, as flies infected by this virus exhibit the smallest impact on survival but a significant reduction in offspring production. These diverse interactions correspond with the unique host transcriptional changes induced by each virus. Notably, Nora virus causes a more substantial disruption to the transcriptome, suggesting a higher immune investment in combating the infection. The differing impacts on offspring production observed across various viral infections are consistent with previous findings (Wallace and Obbard, 2024).

The viruses studied in this work are enteric viruses, meaning that they are transmitted via the fecal-oral route, infecting the organism through the gut. Our observations show that persistent enteric infections significantly shape the bacterial community present within the flies, which is presumed to be located mainly within the gut. This finding agrees

with research focused on acute infections, wherein alterations in microbiome composition have been observed across diverse hosts. The microbiome can modulate immune responses, nutrient absorption, and overall host physiology and thereby affect the outcome of viral infections (González and Elena, 2021; Schissel *et al.,* 2021). Hence, some of the effects observed could be partially explained by the microbiota changes induced by the viral infections. However, it has been observed that for infections with DAV (Nigg *et al.,* 2024) and Nora virus (Franchet *et al.,* 2025), the relative impact on fly survival is largely independent of the microbiota.

Despite the enteric nature of these infections, our observations indicate that their impacts extend beyond the gut, as seen in the locomotion results. The effects on locomotion reflect changes in metabolic and/or neurological status (Grotewiel *et al.,* 2005) and may have profound implications for fly behavior and ecology. Impaired locomotion can affect activities such as foraging, mate searching, and dispersal, consequently influencing the fly's ability to colonize new locations and disseminate associated microbes. This has important implications for disease vectoring and the spread of pathogens. The systemic impact of enteric infections in flies has also been observed for enteric bacteria (Deshpande *et al.,* 2022). One potential explanation for this is the virus spreading from its primary location in the gut to other tissues. Such spreading has been documented for DCV (Ferreira *et al.,* 2014), DAV (Nigg *et al.,* 2024), Nora virus (Franchet *et al.,* 2025). Further studies are required to understand the extent and mechanisms of this dissemination. Another potential explanation for the systemic effects of enteric infections is the dissemination of immune signaling molecules. Molecules released in response to localized gut infections can have far-reaching effects on distant organs and tissues. For instance, studies have shown that gut-derived

inflammatory signals can influence brain function and behavior, a phenomenon known

as the gut-brain axis (Cryan and Dinan, 2012; Mayer, 2011). This interorgan

communication can modulate various physiological processes, including metabolism,

immunity, and even cognitive function. Therefore, the systemic impacts of enteric viruses

may be mediated not only by direct viral spread but also by the immune signaling

molecules that circulate throughout the host.

Finally, we would like to acknowledge some limitations of our work. First, while all

individuals are infected at the time of eclosion, not all flies remain infected throughout

their adult lifespan, despite the virus remaining prevalent in the population. For DAV, Nora

virus, and Bloomfield virus, prevalence remained high, with approximately 100%, 92%,

and 72% of individuals remaining infected, respectively. In contrast, most individuals in

the DCV-infected population cleared the virus by the time of median survival.

Nevertheless, we observed lasting changes to the host transcriptome in flies that cleared

DCV, suggesting that DCV infection may cause long-lasting or permanent alterations in

host physiology and immune function. This is reminiscent of the long-lasting effects of

acute infections (Choutka *et al.,* 2022). Another limitation is that our study focused on

females of a single genetic background. Both sex and genetic background are known to

influence the effects of viral infections in Drosophila (Vale and Jardine, 2015; Klutzier *et*

al., 2023)

In conclusion, our study demonstrates that persistent viral infections significantly

impact key biological traits in *D. melanogaster*. This adds to the growing evidence that

persistent infections profoundly impact host fitness and physiology.

MATERIAL AND METHODS

Fly stock

For all experiments we used *Wolbachia*-free w^{1118} flies maintained on a standard cornmeal diet (Bloomington). All experiments were done at 25°C under a 12:12 light:dark cycle.

Viral infections

Persistently infected flies were established following the method described by Nigg and colleagues in 2022 and are the same stocks used in the work of Nigg and colleagues (2024). The stocks were maintained at 25°C under a 12:12 light:dark cycle.

Survival analysis

Survival analyses of the flies coming from persistently infected stocks were conducted using 20 female flies/replicate sorted at 1 dpe. Male and female flies were collected on the day of eclosion and left to mate for 24 hours. Subsequently, only female flies were sorted in new tubes containing 20 females each. Each tube was considered a biological replicate and a total of nine biological replicates (from three independent experiments) were analyzed. Survival was monitored daily by counting the number of dead flies in each vial and flies were transferred to new vials every two days. Fly survival was assessed from the day of adult eclosion until death.

Viral quantification

Individual female flies were collected in 300 µl of TRIzol reagent (Invitrogen, 15596026). The samples were homogenized using a pestle, and RNA was extracted following

manufacturer's instructions, supplemented with GlycoBlue (Invitrogen, AM9516) during

the precipitation process. RNA concentrations were measured using the Qubit RNA BR

Assay Kit (Invitrogen, Q10211). RT was performed using random primers and Maxima H

Minus Reverse Transcriptase (Thermo scientific, EP0751) following manufacturer's

instructions. The resulting cDNA was diluted 1:10 with MiliQ water. The qPCR was

performed using Luminaris Color HiGreen qPCR Master Mix (Thermo Scientific, K0374).

The reactions were done in triplicates of 10 μl each and using the corresponding set of

primers for each virus (Supplementary Table 1). The cycling conditions were as follows: 2

minutes at 50°C, 10 minutes at 95°C, followed by 40 cycles of 15 seconds at 95°C and 60

seconds at 60°C. A standard melt curve analysis was performed after the cycling

protocol. For the analysis of the results, the threshold for viral RNA detection was set to

35 Ct (i.e. if a sample had Ct higher than 35 it was considered non infected). The Ct values

of the virus for a sample were normalized against the Ct value of the housekeeping gene

Rp49 for that sample (ΔCt). The $2^{-\Delta Ct}$ values were calculated per sample and their $\log_{10}$

plotted.

Viable offspring production

To assess viable offspring production, 20 female and 20 male flies from infected or

uninfected stocks were left to mate in clean fly vials for 24 hours after eclosion. After this

period, males were discarded, and the mated females were transferred to new tubes

every day until no more progeny was produced (17 days). All tubes were kept at 25 °C

under a 12:12 light:dark cycle, and the number of adult flies (i.e. the viable progeny) in

each tube was counted after 14 days. The number of offspring was normalized to the

number of adult females placed in each tube daily to account for those that did not

survive throughout the test duration.

Climbing assays

Climbing assays were performed following the method described by Barone and

Bohmann (2013). Briefly, a 9 cm empty tube was used, and 20 female flies were placed

inside. With the cotton cap on, tubes had a maximum climbable height of 7.2 cm. The

tubes were tapped three times over the span of three seconds, and the climbing height

was assessed by taking a picture 3 seconds after the last tap. Flies' height was assessed

manually on the pictures using a metric guide on the back of the tubes. For categorical

classification, flies were considered "Top" if they surpassed 6.2 cm, "Bottom" if they were

between 0.0-1.0 cm, and "Middle" if they were between 1.0 cm and 6.2 cm.

Microbiota composition

Female flies from uninfected or persistently infected stocks were collected at 12 dpe.

External microbes were removed by surface sterilization using subsequent washes in 10

% bleach for 5 min, 70 % ethanol for 5 min and three rinsing steps in sterile water. At least

20 flies per condition were collected in a single experiment. Flies were placed in a lysis

buffer (10 mM Tris-HCl (pH 8), 26 mM EDTA, 0.5% SDS, 5 mg/mL lysozyme) and

homogenized with sterile pestles. After incubating the lysates for 1 h at 37 °C, total DNA

was extracted using the DNeasy Blood & Tissue Kit (Qiagen) following the manufacturer's

instruction. The V3-V4 hypervariable region of the bacterial *16S* rRNA gene was amplified

using the primers 341F (5'-CCTACGGGNGGCWGCAG-3') and 805R (5'-

GACTACHVGGGTATCTAATCC-3'). Samples were amplified and sequenced at Microsynth

(Balgach, Switzerland) using Illumina MiSeq on paired-end mode. At least 104'000 reads

per sample were obtained. Analyses were performed using qiime 2 (version 2019.10).

After quality trimming, chimeras detection and for and rev read merging using dada2, at

least 75'000 reads per sample were obtained. Taxonomy was attributed to amplicon

sequence variants (ASVs) using the Silva classifier in qiime2. The resulting ASV table was

analyzed in R studio (version 4.2.1) using the phyloseq package (version 1.42.0).

Transcriptome profiling by high-throughput sequencing

For each condition, four samples were sequenced. Each sample was made by pooling

RNA from five individuals (3 ng of total RNA per individual), except for DCV at 28 dpe,

where the four pools were made from three individuals each. RNA-seq libraries were

prepared from 15 ng of pooled RNA using an NEBNext Ultra II RNA Library Prep Kit for

Illumina (New England Biolabs, E7770L) with NEBNext Multiplex Oligos for Illumina (Dual

Index Primers Set 1) (New England Biolabs, E7600S). All sequencing was performed on

an Illumina NextSeq 500 instrument using a NextSeq 500/550 High Output Kit v2.5 (75

cycles) (Illumina, 20024906). Raw sequencing reads were mapped to the *D.*

melanogaster genome release dmel-r6.56 with STAR version 2.7.11b (Dobin et al 2013).

Feature counting was performed with HTSeq version 0.11.2. (Putri et al. 2022), and

differential gene expression analysis was performed with DESeq2 in R Studio (version

2023.03.1). Log2 fold changes were transformed into Z-scores using the "ashr" package,

and only adjusted p-values were considered for analysis.

Statistical analysis

All statistical analyses were conducted using R version 3.6.1 within the Rstudio development environment version 2024.04.2+764. The level of significance was set at $P < 0.05$ and symbols indicate the significance of differences between conditions: *** $P < 0.001$; ** $P < 0.01$; * $0.01 < P < 0.05$.

To account for the block effect, we employed linear mixed-effects or general linear mixed-effects models. Generalized linear models (GLM) were fitted to the data using the *glm* function from the "stats" package. Generalized linear mixed models (GLMM) were fitted to the data using the *lmer* function from the "lme4" package version 1.1-35.5 . Cumulative link mixed models (CLMM) using the *clmm* function from the "ordinal" package version 2023.12-4.1. Post hoc pairwise comparisons were performed using the *emmeans* function from the "emmeans" package version 1.10.5. Tukey's or FDR correction were used to adjust for multiple comparisons. The letter-based grouping for pair-wise comparisons was created using the the *cld* function from the "multcomp" package version 1.4-26. The specific statistical methods applied to each figure are detailed in the legend of the corresponding figure.

REFERENCES

Armstrong JD, Texada MJ, Munjaal R, Baker DA, Beckingham KM. Gravitaxis in

Drosophila melanogaster: a forward genetic screen. Genes Brain Behav. 2006;5: 222–

239. doi:10.1111/j.1601-183X.2005.00154.x

Bahrampour S, Gunnar E, Jonsson C, Ekman H, Thor S. Neural Lineage Progression

Controlled by a Temporal Proliferation Program. Dev Cell. 2017;43: 332-348.e4.

doi:10.1016/j.devcel.2017.10.004

Barone MC, Bohmann D. Assessing Neurodegenerative Phenotypes in *Drosophila*

Dopaminergic Neurons by Climbing Assays and Whole Brain Immunostaining. JoVE.

2013; 50339. doi:10.3791/50339

Biteau B, Hochmuth CE, Jasper H. JNK Activity in Somatic Stem Cells Causes Loss of

Tissue Homeostasis in the Aging *Drosophila* Gut. Cell Stem Cell. 2008;3: 442–455.

doi:10.1016/j.stem.2008.07.024

Choutka J, Jansari V, Hornig M, Iwasaki A. Unexplained post-acute infection

syndromes. Nat Med. 2022;28: 911–923. doi:10.1038/s41591-022-01810-6

Crone EE. Is survivorship a better fitness surrogate than fecundity? Evolution. 2001;55:

2611–2614. doi:10.1111/j.0014-3820.2001.tb00773.x

Cryan JF, Dinan TG. Mind-altering microorganisms: the impact of the gut microbiota on

brain and behaviour. Nat Rev Neurosci. 2012;13: 701–712. doi:10.1038/nrn3346

Deshpande R, Lee B, Grewal SS. Enteric bacterial infection in *Drosophila* induces

whole-body alterations in metabolic gene expression independently of the immune

deficiency signaling pathway. Tennessen J, editor. G3. 2022;12: jkac163.

doi:10.1093/g3journal/jkac163

Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, et al. STAR: ultrafast

universal RNA-seq aligner. Bioinform. 2013;29: 15–21.

doi:10.1093/bioinformatics/bts635

Ferreira ÁG, Naylor H, Esteves SS, Pais IS, Martins NE, Teixeira L. The Toll-Dorsal

Pathway Is Required for Resistance to Viral Oral Infection in *Drosophila*. Schneider DS,

editor. PLoS Pathog. 2014;10: e1004507. doi:10.1371/journal.ppat.1004507

Franchet A, Haller S, Yamba M, Barbier V, Thomaz-Vieira A, Leclerc V, et al. Nora virus

proliferates in dividing intestinal stem cells and sensitizes flies to intestinal infection and

oxidative stress. bioRxiv. 2025. doi:10.1101/2025.01.30.635658

Gargano J, Martin I, Bhandari P, Grotewiel M. Rapid iterative negative geotaxis (RING):

a new method for assessing age-related locomotor decline in *Drosophila*. Exp Gerontol.

2005;40: 386–395. doi:10.1016/j.exger.2005.02.005

Goic B, Saleh M-C. Living with the enemy: viral persistent infections from a friendly

viewpoint. Curr Opin Microbiol. 2012;15: 531–537. doi:10.1016/j.mib.2012.06.002

González R, Elena SF. The Interplay between the Host Microbiome and Pathogenic

Viral Infections. mBio. 2021;12: e02496-21. doi:10.1128/mBio.02496-21

Gould AL, Zhang V, Lamberti L, Jones EW, Obadia B, Korasidis N, et al. Microbiome

interactions shape host fitness. Proc Natl Acad Sci USA. 2018;115.

doi:10.1073/pnas.1809349115

Grotewiel MS, Martin I, Bhandari P, Cook-Wiens E. Functional senescence in

Drosophila melanogaster. Ageing Res Rev. 2005;4: 372–397.

doi:10.1016/j.arr.2005.04.001

Hartl M, Loschek LF, Stephan D, Siju KP, Knappmeyer C, Kadow ICG. A New Prospero

and microRNA-279 Pathway Restricts CO_2 Receptor Neuron Formation. J Neurosci.

2011;31: 15660–15673. doi:10.1523/JNEUROSCI.2592-11.2011

Ho T, Tzanetakis IE. Development of a virus detection and discovery pipeline using next

generation sequencing. Virology. 2014;471–473: 54–60. doi:10.1016/j.virol.2014.09.019

Inaki M, Yoshikawa S, Thomas JB, Aburatani H, Nose A. Wnt4 Is a Local Repulsive Cue

that Determines Synaptic Target Specificity. Curr Biol. 2007;17: 1574–1579.

doi:10.1016/j.cub.2007.08.013

Irschick DJ, Garland T. Integrating Function and Ecology in Studies of Adaptation:

Investigations of Locomotor Capacity as a Model System. Annu Rev Ecol Syst. 2001;32:

367–396. doi:10.1146/annurev.ecolsys.32.081501.114048

Jia Y, Jin S, Hu K, Geng L, Han C, Kang R, et al. Gut microbiome modulates *Drosophila*

aggression through octopamine signaling. Nat Commun. 2021;12: 2698.

doi:10.1038/s41467-021-23041-y

Kapun M, Barrón MG, Staubach F, Obbard DJ, Wiberg RAW, Vieira J, et al. Genomic

Analysis of European *Drosophila melanogaster* Populations Reveals Longitudinal

Structure, Continent-Wide Selection, and Previously Unknown DNA Viruses. Falush D,

editor. Mol Biol Evol. 2020;37: 2661–2678. doi:10.1093/molbev/msaa120

Kim Y-J, Bao H, Bonanno L, Zhang B, Serpe M. *Drosophila* Neto is essential for

clustering glutamate receptors at the neuromuscular junction. Genes Dev. 2012;26: 974–

987. doi:10.1101/gad.185165.111

Kutzer MAM, Gupta V, Neophytou K, Doublet V, Monteith KM, Vale PF. Intraspecific

genetic variation in host vigour, viral load and disease tolerance during Drosophila C virus

infection. Open Biol. 2023;13: 230025. doi:10.1098/rsob.230025

Mayer EA. Gut feelings: the emerging biology of gut–brain communication. Nat Rev

Neurosci. 2011;12: 453–466. doi:10.1038/nrn3071

Miller MS, Farman GP, Braddock JM, Soto-Adames FN, Irving TC, Vigoreaux JO, et al.

Regulatory Light Chain Phosphorylation and N-Terminal Extension Increase Cross-Bridge

Binding and Power Output in *Drosophila* at In Vivo Myofilament Lattice Spacing. Biophys

J. 2011;100: 1737–1746. doi:10.1016/j.bpj.2011.02.028

Mondotte JA, Gausson V, Frangeul L, Blanc H, Lambrechts L, Saleh M-C. Immune

priming and clearance of orally acquired RNA viruses in *Drosophila*. Nat Microbiol.

2018;3: 1394–1403. doi:10.1038/s41564-018-0265-9

Nigg JC, Castelló-Sanjuán M, Blanc H, Frangeul L, Mongelli V, Godron X, et al. Viral

infection disrupts intestinal homeostasis via Sting-dependent NF-κB signaling in

Drosophila. Curr Biol. 2024;34: 2785-2800.e7. doi:10.1016/j.cub.2024.05.009

Nigg JC, Mongelli V, Blanc H, Saleh M-C. Innovative Toolbox for the Quantification of

Drosophila C Virus, Drosophila A Virus, and Nora Virus. J Mol Biol. 2022;434: 167308.

doi:10.1016/j.jmb.2021.167308

Pontzer H, McGrosky A. Balancing growth, reproduction, maintenance, and activity in

evolved energy economies. Curr Biol. 2022;32: R709–R719.

doi:10.1016/j.cub.2022.05.018

Putri GH, Anders S, Pyl PT, Pimanda JE, Zanini F. Analysing high-throughput sequencing

data in Python with HTSeq 2.0. Boeva V, editor. Bioinform. 2022;38: 2943–2945.

doi:10.1093/bioinformatics/btac166

Renn SCP, Park JH, Rosbash M, Hall JC, Taghert PH. A pdf Neuropeptide Gene Mutation

and Ablation of PDF Neurons Each Cause Severe Abnormalities of Behavioral Circadian

Rhythms in *Drosophila*. Cell. 1999;99: 791–802. doi:10.1016/S0092-8674(00)81676-1

Schissel M, Best R, Liesemeyer S, Tan Y-D, Carlson DJ, Shaffer JJ, et al. Effect of Nora

virus infection on native gut bacterial communities of *Drosophila melanogaster*. AIMS

Microbiol. 2021;7: 216–237. doi:10.3934/microbiol.2021014

Shin SC, Kim S-H, You H, Kim B, Kim AC, Lee K-A, et al. *Drosophila* Microbiome

Modulates Host Developmental and Metabolic Homeostasis via Insulin Signaling.

Science. 2011;334: 670–674. doi:10.1126/science.1212782

Sprecher SG, Desplan C. Switch of rhodopsin expression in terminally differentiated

Drosophila sensory neurons. Nature. 2008;454: 533–537. doi:10.1038/nature07062

Tafesh-Edwards G, Eleftherianos I. The role of *Drosophila* microbiota in gut

homeostasis and immunity. Gut Microbes. 2023;15: 2208503.

doi:10.1080/19490976.2023.2208503

Vale PF, Jardine MD. Sex-specific behavioural symptoms of viral gut infection and

Wolbachia in *Drosophila melanogaster*. J Insect Physiol. 2015;82: 28–32.

doi:10.1016/j.jinsphys.2015.08.005

Walker RG, Willingham AT, Zuker CS. A *Drosophila* Mechanosensory Transduction

Channel. Science. 2000;287: 2229–2234. doi:10.1126/science.287.5461.2229

Wallace MA, Coffman KA, Gilbert C, Ravindran S, Albery GF, Abbott J, et al. The

discovery, distribution, and diversity of DNA viruses associated with *Drosophila*

melanogaster in Europe. Virus Evol. 2021;7: veab031. doi:10.1093/ve/veab031

Wallace MA, Obbard DJ. Naturally occurring viruses of *Drosophila* reduce offspring

number and lifespan. Proc R Soc B. 2024;291: 20240518. doi:10.1098/rspb.2024.0518

Yamanaka N, Romero NM, Martin FA, Rewitz KF, Sun M, O'Connor MB, et al.

Neuroendocrine Control of *Drosophila* Larval Light Preference. Science. 2013;341: 1113–

1116. doi:10.1126/science.1241210

infected fly population. *A) Experimental design. The large arrow indicates the days post-eclosion (dpe) during which specific experiments were conducted. The survival of persistently infected stocks has been monitored from eclosion until death across all stocks. The viability of offspring was assessed after a 24-hour mating period, until no more embryos were laid (17 dpe). Black dots indicate the specific days when the climbing assay and the sampling for the bacteriome studies were performed for all stocks. At the bottom of the graph, regarding the transcriptomics experiment, each colored dot signifies the date of sample collection for RNA sequencing corresponding to each different fly stock. B) Left panel shows the survival of w1118 flies uninfected (green) or persistently infected with DAV (red), DCV (yellow), Bloomfield virus (dark blue) or Nora virus (purple) Shaded regions indicate 95% confidence intervals. The data (n = 180 flies/condition in 3 independent experiments with 3 biological replcates/experiment) were obtained from Nigg et al. (2024). Statistical significance was assessed via the Mantel-Cox test. Right panel indicates the day post-eclosion at which 50% of the population is deceased. C) Viral infection dynamics. The upper panel shows viral RNA levels for DAV, DCV, Bloomfield virus, and Nora virus in infected populations. Viral RNA levels were measured in whole individuals at various times post-eclosion. Only data from infected individuals is presented. The Y-axis denotes the relative viral RNA level as $\log_{10} 2^{\Delta Ct}$ between the viral RNA and the housekeeping gene, Rp49. The red bars represent the median values along with the interquartile range.* For each virus, significance was calculated using a general linear model where the time point was the fixed factor; Tukey contrasts were used for post hoc analyses. Letters over the bars indicate letter-based grouping for multiple comparisons. Bottom panel: viral prevalence showing the proportion of flies resulting infected (black) or non-infected (grey) at the corresponding time-point. *Numbers below the pie chart represent the number of positives between the number of individuals sampled.*

Figure 2. Impact of persistent viral infection on offspring production. *A) Daily viable offspring per female. Number of viable offspring produced by female flies uninfected (green) or persistently infected with DAV (red), DCV (yellow), Bloomfield virus (dark blue) or Nora virus (purple) after one day of mating (20 females paired with 20 males).* After one day, the males were discarded, and the females were transferred to new vials with food each day. Mated flies were kept for subsequent days, and the viable offspring were counted. *The number of viable offspring was normalized by the number of females laying eggs in each tube. Day 0 represents the day of eclosion and mating. Experiments were conducted on 20 females paired with 20 males in 9 replicates. Significance was calculated using a general linear-mixed model where the time point was the fixed factor and experiments a random effect; Tukey contrasts were used for post hoc analyses. B) Cumulative viable offspring shown in (A). Total viable offspring produced by each female was measured cumulatively across the different fly populations throughout the duration of the experiment. The Y-axis shows the total viable offspring per female and the X-axis the population studied. Significance was calculated using a general linear-mixed model where the virus status was the fixed factor and experiments a random effect; Tukey contrasts were used for post hoc analyses.*

Figure 3. Impact of persistent viral infection on locomotion. *Climbing assay results of flies uninfected (green) or persistently infected with DAV (red), DCV (yellow), Bloomfield virus (dark blue) or Nora virus (purple).* ***A)*** *Quantitative height climbed by individual flies. Climbing assay results for days 1, 6, 12, and 20 post-eclosion. Results of 9 different biological replicates with 20 female flies. At each time point and replicate, six pictures were taken, and the height of each fly in the tube was recorded. Each symbol on the graph represents the recorded height of a fly for each picture. Boxplots are shown on the left side of each stock, indicating the median and interquartile range of the heights climbed by the flies. The dotted line represents the median height climbed by the uninfected population. The half-violin of each plot shows the density of the values. Significance was calculated using a general linear-mixed model where the virus status and the time point were the fixed factors and experiments a random effect; Tukey contrasts were used for post hoc analyses.* ***B)*** *Qualitative measurements of the climbing assay. Top (green) indicates the flies present in the top 1 cm of the climbing tubes (6.2-7.2 cm). Bottom (red) indicates the percentage of flies located in the bottom 1 cm of the tube (0-1 cm), and middle (blue) includes all flies that climbed between (1-6.2 cm). Significance was calculated using a cumulative link mixed model where the virus status and the time point were the fixed factors and experiments a random effect; false discovery rate contrasts were used for post hoc analyses.*

Figure 4. Bacteriome composition of the studied fly populations except for the one infected with Bloomfield virus. *Relative abundance in percentage of bacterial genera detected by sequencing of the V3-V4 region of the bacterial 16S rRNA gene. Each color corresponds to a specific bacterial genus sequenced in the microbiota. Bacteria belonging to different amplicon sequence variants (ASVs) but belonging to the same bacterial genus are displayed with the same color.*

Figure 5. Transcriptomic profiles of uninfected or persistently infected flies. *A) Heatmaps expressing the Z-scores of gene expression fold change of infected populations in comparison with uninfected ones. Genes are represented in the Y-axis and the infecting virus on the X-axis. For each virus, the results are calculated on the average of 4 samples. Top left: heatmap showing all the genes reported to be involved in antiviral immunity (subset of 191 immune antiviral genes, Supplementary File 1) and their differential expression in our samples at dpe1. Top right: heatmap showing the top 20 differentially expressed immune genes at dpe1. Bottom left: heatmap showing all the genes reported to be involved in the insect's immune response and their differential expression in our samples at dpe12. Bottom right: heatmap showing the top 20 differentially expressed immune genes at dpe12. On the right side of each heatmap, a colored bar indicates the pathway in which the gene is reported to participate. B) Heatmaps showing all the genes reported to be involved in the insect's locomotion (a subset of 264 locomotion genes, Supplementary File 1) and their differential expression in our samples at dpe1. Top left: heatmap showing the top 20 differentially expressed locomotion genes at dpe1. Top right: heatmap showing all the genes reported to be involved in the insect's locomotion and their differential expression in our samples at dpe12. Bottom left: heatmap showing the top 20 differentially expressed immune genes at dpe12. Bottom right: heatmap showing the top 20 DEGs at dpe12.*

Figure 6. Transcriptomic profile of flies at 28 dpe that were infected with DCV but cleared the viral infection. *A) Principal component analysis of uninfected samples, DCV Infected, and DCV Cleared. On the X-axis, we indicate the first principal component (PC1) and the percentage of variance explained by this component in brackets. On the Y-axis, we indicate the second principal component (PC2) and the percentage of variance explained by this component in brackets. B) Volcano plot indicating the differentially expressed genes between DCV Cleared samples and DCV Infected ones. Genes significantly up- (in red) or down- (in blue) regulated. In grey, we represent those genes with differential expression values lower than 1.5 log2 fold change and p-adjusted values lower than 0.05. These thresholds are indicated by dotted lines in the graphs. The names of the DEGs are shown. C) Heatmaps representing the differential expression patterns between DCV Infected and DCV Cleared samples when compared to the Uninfected of the same age. Heatmap on the right shows the top 200 differentially expressed genes in our samples in terms of absolute log2 fold change value at dpe 28. Heatmap on the left shows the top 20 DEGs at dpe 28; gene names are indicated on the left. D) Heatmaps representing the differential expression patterns for reported immune genes between DCV Infected and DCV Cleared samples when compared to the Uninfected of the same age. Heatmap on the left shows all the genes reported to be involved in the insect's immune response and their differential expression in our samples at dpe 28. Heatmap on the right shows the top 20 differentially expressed immune genes at dpe 28; gene names are indicated on the left. On the right side of each heatmap, a colored bar indicates the pathway in which the gene is reported to involved.*

SUPPLEMENTARY FIGURES

Supplementary Figure 1. *Principal Component Analysis of the different samples sequenced. On the Y axis of every graph, we indicate the second principal component (PC2) and the percentage of variance explained by this component between brackets. On the X axis, we indicate the first principal component (PC1) and the percentage of variance explained by this component between brackets. Black lines indicate the origin of each axis. The do represent our fly stocks throughout the different days post-eclosion, in green, we have the Uninfected stock, in burgundy, the Drosophila A virus persistent flies, in yellow, the Drosophila C virus persistently infected flies, in dark blue, we have the Bloomfield virus persistently infected stock, and in lavender the Nora virus persistently infected flies.*

Supplementary Figure 2. *Transcriptomic results indicate differentially expressed genes in our infected samples. Volcano plots showing the number of genes significantly up- (in red) or down- (in blue) regulated. In grey, we represent those genes displaying differential expression values lower than 1.5 log2 fold change and/or with p-adjusted values lower than 0.05. These thresholds are indicated by dotted lines in the graphs. Each row presents the volcano plots of a specific fly stock over time (dpe), in the following order: DAV, DCV, Bloomfield, and Nora. Each infected sample is analysed in comparison to an uninfected control of the same age.*

Supplementary Figure 3. *Persistent viral infection alters your transcriptomic profile in a virus-independent manner. For all heatmaps, on the X-axis, we indicate the infected stock analysed. **A)** From left to right: heatmap showing the top 200 differentially expressed genes in our samples in terms of absolute log$_2$ fold change value at dpe1. Heatmap showing the top 20 DEGs at dpe1. **B)** From left to right: heatmap showing the top 200 differentially expressed genes in our samples in terms of absolute log$_2$ fold change value at dpe12. Heatmap showing the top 20 DEGs at dpe12. For the heatmaps on the right, the top 20 gene names are indicated on the left of the graph.*

Supplementary Figure 4. *Common DEGs across infections and time. Venn diagrams of differentially expressed genes with differential expression values lower than 1.5 log2 fold change and/or with p-adjusted values lower than 0.05.* ***A)*** *DEGs for days 1, 12, and the day of median survival for each infected individual stock.* ***B)*** *DEGs on the day of median survival of all persistently infected stocks.* ***C)*** *DEGs for each persistently infected stock over time.*

SUPPLEMENTARY TABLES

621 **Supplementary Table 1.** Primers used in qPCR step of RT-qPCR for viral load

622 quantification.

Primer name	Sequence
Rp49_Forward	CGGATCGATATGCTAAGCTGT
Rp49_Reverse	GCGCTTGTTCGATCCGTA
DAV_Forward	GTTGGATCAGGCTAGTGTAGG
DAV_Reverse	TGCAACCGGACTCCAAGTTC
DCV_Forward	AGGCTGTGTTTGCGCGAAG
DCV_Reverse	AATGGCAAGCGCACACAATTA
Bloomfield_Forward	CCGCCGGCTACCTACTTTAG
Bloomfield_Reverse	ACCAGCAAGCGTCCGAAATA
Nora_Forward	GGTTTGCAGTTGATCGCAG
Nora_Reverse	TCAGCTCATCGTTCACGAA

SUPPLEMENTARY FILES

625 **Supplementary File 1.** Genes classified with antiviral immune or locomotion function.