1 **Persistent viral infections impact key biological traits in *Drosophila melanogaster***
2

Mauro Castelló-Sanjuán[1], Rubén González[1], Ottavia Romoli[1], Hervé Blanc[1], Jared C. Nigg[1, #],

Maria-Carla Saleh[1].

[1]Institut Pasteur, Université Paris Cité, Viruses and RNA Interference Unit, Paris, France.

[#]Current address: Chan Zuckerberg Biohub, San Francisco, California, USA.

Correspondence: jared.nigg@czbiohub.org, carla.saleh@pasteur.fr

ABSTRACT

Persistent viral infections have been assumed to impose minimal fitness costs for

insects. We established persistent mono-infections of *Drosophila melanogaster* with

four different enteric RNA viruses: Drosophila A virus (DAV), Drosophila C virus (DCV),

Bloomfield virus, and Nora virus. We observed that these infections significantly reduce

fly survival, alter the number of viable offspring per female, modulate microbiome

composition, impact locomotor abilities, and change activity patterns. These results

demonstrate the significant impact of persistent viral infections on key biological traits

and expand our understanding of the fitness costs of persistent viral infections for the

host. In addition, the four viruses displayed different accumulation kinetics and elicited

unique transcriptional profiles with no common core responses. The transcriptional

changes triggered by DCV infection persisted even after viral clearance. This

comprehensive comparative dataset represents a valuable resource for researchers

studying host-pathogen interactions, providing detailed transcriptional profiles, and

behavioral measurements across different viral infections and time points. Our findings

reveal that persistent viral infections modulate critical aspects of insect biology, affecting

host physiology and behavior.

INTRODUCTION

The increasing availability of sequencing data has revealed tens of thousands of novel viruses in multiple host organisms, suggesting the widespread occurrence of previously undetected infections (Ho and Tzanetakis, 2014; Kapun *et al.,* 2020; Wallace *et al.,* 2021). Recent studies have revealed extensive diversity in the RNA virosphere, identifying thousands of previously unknown viruses across diverse environments, including the most extreme habitats worldwide (Dominguez-Huerta et al., 2023; Hou et al., 2024; Shi et al., 2016). Among these newly discovered viruses are many that naturally infect *Drosophila* (Obbard et al., 2020; Wallace et al., 2021; Webster et al., 2015, 2016). In some cases, these are persistent infections that represent a metastable state in which the physiological impacts of viral infection are maintained below a lethal threshold without leading to viral clearance (Goic & Saleh, 2012; Lee et al., 2019). In contrast to acute viral infections, which are characterized by a rapid process that results in the quick death of the host or in the clearance of the virus, persistent infections endure over extended periods within the individual host. During this time, the virus is still capable of being transmitted to other organisms, including the host's offspring, despite the inherent limitations imposed by the host's immune responses and cellular mortality (Goic & Saleh, 2012; Hogenhout et al., 2008; Lee et al., 2019; Salas-Benito & De Nova-Ocampo, 2015; Williams et al., 2017).

Persistence is a frequent phenomenon in viral infections across all kingdoms, as it can provide advantages for both the virus and the host, acting as a modulator of the ecosystem. For instance, the cucumber mosaic virus (CMV) infects beet plants (*Beta vulgaris*), improving their drought tolerance and enhancing their freezing tolerance (Xu et al., 2008). On the other hand, persistent human papillomavirus (HPV) infection must

undermine host antiviral defense mechanisms, disrupt the balance of cellular proliferation and differentiation, and hijack DNA damage signaling and repair pathways to replicate viral DNA in stratified epithelium. Together, these modulations of host physiology carry detrimental consequences for the host, putting cells at high risk for carcinogenesis (Della Fera et al., 2021). Nevertheless, research has mostly focused on the acute effects of viral infections. Consequently, the impacts of persistent infections on host physiology and behavior remain understudied.

In this study, we aimed to elucidate whether persistent infections in the fruit fly, *Drosophila melanogaster* have consequences on fly fitness or key biological traits. The fruit fly has an extensively characterized genome and short generation times, making it an ideal model organism for studying persistent viral infections throughout lifespan. Growing evidence shows that persistent infections have costs to the host (Nigg et al. 2024, Wallace and Obbard, 2024). In the present study, we aimed to further examine the effects of persistent mono-infections with four distinct enteric RNA viruses—Drosophila A Virus (DAV), Drosophila C Virus (DCV), Bloomfield virus, and Nora virus—on a wide range of *D. melanogaster* physiological traits. Specifically, we studied survival rates, fertility (reproductive output, here defined as the number of viable offspring), bacteriome load and composition, locomotor abilities, and activity patterns. These biological traits are essential for understanding the host's overall fitness and its ecological interactions. Survival rates and reproductive output serve as key indicators of an organism's ability to thrive and contribute to the next generation (Crone, 2001). The microbiome has been observed to influence various aspects of the host's physiology, such as nutrition status, immune function, behavior, and overall health (Gould *et al.,* 2018; Jia *et al.,* 2021; Shin *et al.,* 2011; Tafesh-Edwards and Eleftherianos, 2023). Locomotor abilities and activity

patterns are vital for behaviors such as foraging, mating, and avoiding predators, all of

which are crucial for survival and reproduction (Irschick and Garland, 2001). By

examining these traits, we highlighted the broader implications of persistent viral infections

on insect biology.

RESULTS

The dynamics of viral RNA levels over time are virus-specific

Our laboratory has previously established persistent viral mono-infections in *D.*

melanogaster flies of the w^{1118} genetic background using different viruses: DAV, DCV,

Bloomfield virus, and Nora virus (Nigg *et al.,* 2022, 2024). All these positive sense single-

strand RNA viruses belong to different viral families and naturally infect *D. melanogaster*.

They are transmitted primarily through orofecal contamination. In this work, we

measured key biological traits over time in the infected and the uninfected control

population (Figure 1A). To maintain continuous exposure to contaminated food while

preventing excessive viral accumulation, flies from the different persistently infected

lines were transferred to fresh food every other day. Persistent viral infections

significantly reduce the survival of infected flies, as previously described (Figure 1B; Nigg

et al., 2024). DAV infection has the most pronounced effects on survival, followed by DCV,

Bloomfield virus, and Nora virus (Mantel-Cox test infected vs uninfected $p < 0.001$ for all

the virus-infected populations). This is reflected by the time at which 50% of the

population has perished, the median survival (Figure 1B): 25 days post-eclosion (dpe) for

DAV, 28 dpe for DCV, 36 dpe for Bloomfield virus, and 52 dpe for Nora virus, compared to

63 dpe for uninfected flies. This information enabled us to establish different time points

for further characterization of the infections: 1 dpe as an early time point, 12 dpe as an

intermediate time point just before the onset of virus-induced mortality, and the median

survival as a late time point specific for each condition. Additionally, we added an

additional time point for DCV at 6 dpe, as mortality starts earlier for this virus. In the

present study, these time points served as the basis for characterizing the impact of

persistent viral infections on the physiology and behavior of infected hosts (Figure 1A).

Figure 1. Experimental set-up, survival, offspring production, viral dynamics, and bacteriome analysis of persistently infected fly populations. *A) Experimental design. The large arrow indicates the days post-eclosion (dpe) during which specific experiments were conducted. The survival of persistently infected stocks has been monitored from eclosion until death across all stocks. The viability of offspring was assessed after a 24-hour mating period, until no more embryos were laid (17 dpe). Black dots indicate the specific days when the climbing assay and the sampling for the bacteriome studies were performed for all stocks. At the bottom of the graph, regarding the transcriptomics experiment, each colored dot signifies the date of sample collection for RNA sequencing corresponding to each*

*different fly stock. **B)** Left panel shows the survival of w1118 flies uninfected (green) or persistently infected with DAV (red), DCV (yellow), Bloomfield virus (dark blue) or Nora virus (purple) Shaded regions indicate 95% confidence intervals. The data (n = 180 flies/condition in 3 independent experiments with 3 biological replicates/experiment) were obtained from Nigg et al. (2024). Statistical significance was assessed via the Mantel-Cox test. Right panel indicates the day post-eclosion at which 50% of the population is deceased. **C)** Viral infection dynamics. The upper panel shows viral RNA levels for DAV, DCV, Bloomfield virus, and Nora virus in infected populations. Viral RNA levels were measured in whole individuals at various times post-eclosion. Only data from infected individuals is presented. The Y-axis denotes the relative viral RNA level as $\log_{10} 2^{\Delta Ct}$ between the viral RNA and the housekeeping gene, Rp49. The red bars represent the median values along with the interquartile range. For each virus, significance was calculated using a general linear model where the time point was the fixed factor; Tukey contrasts were used for post-hoc analyses. Letters over the bars indicate letter-based grouping for multiple comparisons. Bottom panel: viral prevalence showing the proportion of f infected (black) or non-infected (grey) flies at the corresponding time-point. Numbers below the pie charts represent the number of infected individuals over the number of individuals sampled. **D)** Cumulative viable offspring for each population, normalized over female number. Total viable offspring produced by each female was measured cumulatively across the different fly populations throughout the duration of the experiment, accounting for fly mortality. The Y-axis shows the total viable offspring per female and the X-axis the population studied. Significance was calculated using a general linear-mixed model where the virus status was the fixed factor and experiments a random effect; Tukey contrasts were used for post-hoc analyses. **E)** Relationship between the offspring and the survival of fly stocks. On the Y axis the total offspring per female. On the X axis, the median day of survival, in dpe, of each infected stock. Each dot on the graph represents a total offspring experiment. All infected samples from the same infected stock have the same median survival. For each virus-infected stock, the mean offspring is denoted as a black rhombus. Pearson's correlation coefficient r = -0.633 and p value < 0.001 are depicted in the graph. **F)** Bacterial composition of pools of at least 20 females from the studied fly populations except for the one infected with Bloomfield virus. Females coming from the uninfected or persistently infected stocks were collected at 12dpe, surface sterilized and their DNA was extracted. For each population, microbiome composition is expressed as the relative abundance in percentage of bacterial genera detected by sequencing of the full bacterial 16S rRNA gene. Each color corresponds to a specific bacterial genus sequenced in the microbiome. Bacteria belonging to different operational taxonomy units (OTUs) but belonging to the same bacterial genus are displayed with the same color. For all the figure, *** p < 0.001, ** p < 0.01, * p < 0.05. **G)** Total bacterial load of pools of 10-20 females coming from the uninfected or persistently infected stocks except for the one infected with Bloomfield virus. Females coming from the uninfected or persistently infected stocks were collected at 12dpe, surface sterilized and their DNA was extracted. The experiment was performed twice, with three pools per line analyzed in each experiment. Bacterial load was assessed via qPCR targeting the 16S bacterial ribosomal gene. The Y-axis denotes the relative bacterial DNA level as $\log_{10} 2^{\Delta Ct}$ between the bacterial 16S gene and the housekeeping gene, Rp49 (mean $\pm$SEM). Shape fill indicates the replicate. Statistical significance was determined with a generalized linear model using $\log_{10} 2^{\Delta Ct}$ values and condition as fixed factors ($F_{(4,24)}$ = 26.63, p < 0.001) and the replicate as a random factor. Tukey's adjustment was applied to post-hoc comparisons.*

To characterize the dynamics of viral load within the infected populations, we examined viral RNA levels in individual flies at the aforementioned early, intermediate, and late stages of infection. Viral loads were quantified by RT-qPCR by measuring viral RNA levels relative to the *Drosophila* housekeeping gene *Rp49*. Flies were classified as infected if their Ct values for viral RNA were below 35, with melting temperature profiles indicating the presence of a single amplicon. We observed distinct patterns of viral accumulation for each virus (Figure 1C, upper panels). DAV exhibited a rapid increase in viral RNA levels early post-eclosion, with a median increase of ~4 and ~5 $\log_{10}$ orders of magnitude at 12 and 25 dpe, respectively. In contrast, DCV displayed a relatively stable

pattern in viral RNA levels over time, with no significant changes in viral loads. Interestingly, we noticed that DCV-infected individuals stably clustered into two distinct groups, displaying viral RNA levels several orders of magnitude apart. Bloomfield virus RNA levels rose steadily over the lifetime of infected individuals, increasing by 1 and 2 $\log_{10}$ orders of magnitude at 12 and 36 dpe, respectively. In contrast, Nora virus RNA levels remained consistent at the earlier timepoint but rose significantly by ~3 $\log_{10}$ orders of magnitude by the final timepoint (General Linear Model -GLM- letters over the conditions indicate grouping for multiple comparisons).

Our examination of viral RNA levels allowed us to estimate infection prevalence in the different populations over time (Figure 1C, lower panels). A viral infection is considered prevalent in a population if individuals continue to be infected over time. DAV and Nora virus were consistently detected in 100% of individuals, except for a slight reduction in Nora virus prevalence at the later stage (~91% of individuals infected at 52 dpe). Bloomfield virus was highly prevalent in the population at all time points, although not all individuals were infected at any given time, and infection prevalence varied from ~91% to ~72%. DCV was detected in all individuals at 1 dpe, and its prevalence in the population progressively decreased over time, with only ~36% of the individuals infected at the latest time point.

Persistent viral infections impact viable offspring production

Persistent viral infections can reduce individual survival rates, which may consequently affect the viability of the population over time. To understand the broader impact of these infections, we investigated whether flies from populations with a prevalent viral infection produced altered numbers of viable offspring compared to flies from the uninfected

population. For this experiment, we measured viable offspring production as the number

of progeny that successfully emerged as adults. This metric is a crucial aspect of

reproductive success and population dynamics. Any impact on the production of viable

offspring would indicate that persistent infections influence additional critical biological

traits related to fitness beyond the previously observed reduction in survival.

To explore potential changes in viable offspring production in the infected populations,

20 males and 20 females from each infected population or from the uninfected

population were collected on the day of eclosion and placed together in fresh vials for 24

hours to allow mating. After 24 hours, males were discarded, and the 20 females were

transferred into new vials with fresh food each day. Initial female density across all

populations was uniformly established at 20 females per tube, as these conditions

provide optimal surface area and total tube space, thereby minimizing crowding effects

and promoting normal development. The viable offspring were counted daily as they

emerged and the viable offspring count was normalized based on the number of live

females remaining in each vial. This normalization allowed us to account for changes in

the density of females per tube due to mortality over the course of the experiment.

Flies collected from all four virus-infected populations exhibited distinct patterns of

offspring production compared to uninfected controls (Supplementary Figure 1).

Uninfected flies exhibited a peak in viable offspring production at 1 dpe (10.3 ± 2.02

adults emerging per female, mean ± SD), followed by a gradual decrease in offspring

production until 6 dpe and a pronounced decline thereafter, reaching zero by 16 dpe.

Similarly, populations infected with DAV and Bloomfield viruses also peaked in viable

offspring production at 1 dpe (11.5 ± 2.07 and 11.6 ± 1.30 offspring per female,

respectively) and experienced a sharp decline in offspring production after 6 dpe.

However, during early time points, DAV and Bloomfield virus-infected populations

produced more viable offspring per female than the uninfected control population

(GLMM, DAV P = 0.022, Bloomfield P = 0.031). DCV-infected flies displayed a distinct

pattern, producing significantly more viable offspring than the uninfected ones only at

later time points (GLMM, P < 0.001). In contrast, Nora virus-infected flies consistently

produced fewer viable offspring (GLMM, P = 0.010), with their offspring production

peaking at 4 dpe (5.28 ± 2.15 viable offspring per female).

Studying both daily and cumulative offspring production offers a comprehensive

perspective on reproductive fitness. While daily offspring production captures temporal

patterns and short-term reproductive dynamics, cumulative counts reveal the total

reproductive output across a female's lifespan.

Analysis of cumulative offspring per female reveals that DCV-infected flies produced

more viable offspring (91.5 ± 32.1) than the uninfected group (67.7 ± 11.6) (Figure 1D,

GLMM, P = 0.004). In contrast, Nora virus-infected flies produced significantly fewer

offspring per female (41.5 ± 15.5, GLMM, P = 0.034). Although DAV and Bloomfield virus-

infected flies showed higher early production rates, their lifetime reproductive output did

not differ significantly from uninfected controls. These results demonstrate that

persistent viral infections have virus-specific impacts on reproductive fitness, with some

viruses enhancing and others diminishing total offspring production.

Analysis of the relationship between total offspring production and female survival

reveals a negative correlation across virus-infected stocks (r = -0.633, P < 0.001 ; Figure

1E). DAV and DCV infections resulted in higher offspring production but shorter lifespans,

while Bloomfield and Nora virus infections showed the opposite pattern: lower

reproductive output but longer survival. This tradeoff suggests a potential reallocation of

resources from reproduction to defense mechanisms in response to viral infections.

Harboring persistent viral infections changes fly bacteriome

We next investigated whether persistent viral infections influence the fly bacteriome—

the bacteria associated with the host. All fly populations in this study shared the same

genetic background and were reared under identical conditions, except for the presence

of different persistent viral infections. This controlled setup allowed us to attribute any

observed differences in the associated bacteria to the presence of different viral

infections.

Pools of at least 20 flies in three replicates were collected at 12 dpe, surface sterilized

and their DNA was extracted. Microbiome composition was assessed by amplification

and sequencing of the full *16S* rRNA bacterial gene through Oxford Nanopore

sequencing. Despite good DNA yields for all samples, bacterial *16S* amplification was

not successful for Bloomfield virus infected flies, even when DNA was extracted from

pools of 40 flies. We therefore could not determine the bacterial composition of

Bloomfield virus infected flies.

An average of 82,053 reads was obtained per sample (min: 25,673; max: 139,361). As

previously described, we found a relatively low bacterial diversity in all the analyzed

populations, with only 7 to 21 operational taxonomy units (OTUs) identified in our

samples. Alpha diversity analyses did not reveal any significant difference in the OTU

diversity of uninfected or persistently infected flies (Supplementary Figure 3A, pairwise

Wilcoxon test on Chao1 and Shannon indexes, $P > 0.46$ for all comparisons). Similarly,

beta diversity analysis did not identify specific differences in the bacterial community

structure of the different fly lines (Supplementary Figure 3B, PERMANOVA on infection, F = 1.57, P = 0.24). After taxonomical assignation of reads, we confirmed that the microbiome of all our *Drosophila* populations was dominated by *Lactiplantibacillus* bacteria characterized by a *16S* sequence identity higher than 99% (Figure 1F). The failed amplification of the *16S* gene of Bloomfield virus-infected flies suggested a significant difference in the total microbiome load of our fly populations. We thus determined the total bacterial load in whole flies by *16S* qPCR. Nora virus-infected flies showed a significant increase in total bacterial loads compared to uninfected flies. Similarly, flies persistently infected with DAV or DCV were characterized by a non-significant increase in their total bacterial loads (Figure 1G). Although we successfully amplify bacterial *16S* from Bloomfield virus-infected flies (with amplification cycles earlier than no-template controls), their Ct values closely matched those of extraction negative controls – blank samples processed alongside fly samples. This confirms that flies persistently infected with Bloomfield virus harbor a markedly reduced microbiota load.

Taken together, these data indicate that persistent viral infection does not influence the composition of the microbiome but significantly impacts the total bacterial load colonizing the fly.

Persistent viral infections alter locomotor function in a virus-dependent manner

Locomotor function is a key indicator of an organism's overall health and physiological status. In flies, locomotion reflects both muscular and neurological capacity, making it a valuable marker for assessing the general impact of viral infections on host physiology.

We assessed locomotor function using the reverse geotaxis assay, which measures the flies' upward climbing instinct over a specified period (Gargano *et al.,* 2005). The

assay was performed on 20 mated females from the uninfected and the different infected

stocks at 1, 6, 12, and 20 dpe. This final time point was chosen as a common late date for

all fly stocks, occurring prior to the day of median survival for all of the infected

populations. Flies were placed in an empty tube and collected at the bottom by gently

tapping the tubes. After 3 seconds, a picture was taken and the position of each fly was

recorded manually. Our observations indicated that persistent viral infections had a

mixed effect on locomotor function over time (Supplementary Figure 2). DCV- and Nora

virus-infected flies consistently showed impaired locomotion abilities compared to

uninfected flies at almost all the tested time points (Supplementary Figure 2). In contrast,

DAV- and Bloomfield virus-infected flies showed mixed locomotion patterns. DAV

significantly reduced mobility at 1 dpe but significantly increased it at 6 and 20 dpe

(GLMM, $P < 0.001$, $P < 0.001$, and $P = 0.011$, respectively). Bloomfield virus significantly

reduced mobility at 1 dpe and increased it at 12 dpe (GLMM, $P < 0.001$ at both time

points). When examining the fly distribution patterns along the tubes we obtained similar

results, with a higher proportion of DCV- and Nora virus-infected flies found in the bottom

of the tube at almost all time points (Supplementary Figure 2). DAV infection resulted in

a higher proportion of flies climbing to the top of the tube at 6 dpe, but a higher proportion

of flies at the bottom at 12 dpe (CLMM, $P < 0.001$ and $P = 0.001$). Bloomfield virus-infected

flies were found more often on the bottom of the tube at 1 and 20 dpe, while they climbed

more to the top at 12 dpe (CLMM, $P < 0.001$, $P = 0.003$, and $P = 0.032$, respectively). Taken

together, these results indicate that persistent viral infections affect the climbing abilities

of infected flies, with a general trend of lower climbing height, particularly in flies

harboring Nora virus infections. These findings underscore the systemic impact of

persistent viral infections on host physiology and behavior.

Persistent viral infections alter the fly activity and sleep duration while preserving circadian rhythms

We continuously monitored fly movement over a 72-hour period under 12:12 h light:dark conditions to assess the impact of persistent viral infections on activity patterns and sleep behavior. The movement data revealed that all experimental groups maintained clear circadian rhythmicity, with peaks in activity occurring at light-dark transitions and similar overall temporal patterns of activity (Figure 2A).

Figure 2. Activity patterns and sleep behavior of persistently infected flies. *A) Mean locomotor activity over a 72-hour period under 12:12 light:dark conditions (T0-T72). Activity was quantified as infrared beam crossings per minute using Drosophila Activity Monitors. Colored lines represent different infection conditions (uninfected: green, Bloomfield virus: blue, DAV: red, DCV: yellow, and Nora virus: purple), with shaded areas indicating 95% confidence intervals. Gray shaded vertical bands indicate dark periods. Time is shown in hours, with T0 representing the beginning of the first light cycle after one night cycle of acclimation. B) Sleep time comparison during night and day phases. Sleep was defined as periods with no activity for at least 5 consecutive minutes. Relative sleep time is shown as proportion of total available time in each phase. Box plots indicate the median and interquartile range; each point represents an individual fly. Statistical significance compared to uninfected controls was determined using a linear mixed-effects model with experimental replicate as a random effect (*** P < 0.001, ** P < 0.01, * P < 0.05). C) Activity intensity during*

Analysis of sleep periods, defined as immobility bouts lasting at least 5 minutes (Cirelli

and Bushey, 2009), revealed significant differences between infected and uninfected

flies (Figure 2B). During the night phase, flies infected with Bloomfield, DAV, or Nora virus

spent significantly more time sleeping compared to uninfected controls (GLMM, all $P <$

0.001). DCV-infected flies showed no significant difference in night sleep compared to

uninfected controls. Day-phase sleep patterns displayed a similar profile, with all

infected flies exhibiting significantly increased sleep duration compared to uninfected

controls (GLMM, all $P < 0.001$).

We further analyzed the intensity of movement during active periods (when flies were

not asleep) to determine whether viral infections affected the quality of movement

(Figure 2C). Bloomfield-infected flies showed significantly reduced movement intensity

during their active periods compared to uninfected controls (GLMM, $P < 0.05$), suggesting

that even when Bloomfield virus-infected flies were active, they moved less vigorously

than uninfected flies. In contrast, flies infected with DAV, DCV, or Nora virus showed

movement intensities during active periods that were comparable to uninfected controls,

with no significant differences detected. Together, these results demonstrate that while

persistent viral infections do not disrupt fundamental circadian rhythmicity in

Drosophila, they significantly modify both sleep duration and movement intensity in a

virus-specific manner.

Host transcriptional response is virus specific

To better understand the alterations induced by persistent viral infections, we examined

the transcriptional response of flies at early (1 dpe), intermediate (12 dpe), and late

stages of infection (at median survival times). For each time point and condition, RNA

was extracted from individual flies for individual measurements of viral load via RT-qPCR.

Subsequently, aliquots of RNA from three to five infected individuals harboring similar

viral load levels (Figure 1C) were pooled together for the generation of bulk RNA-seq

libraries.

Principal component analysis revealed that each enteric virus elicited a distinct

transcriptional response in the host (Supplementary Figure 4). Nora virus infection was

characterized by the most unique transcriptional response at 1 dpe, with the differential

regulation of several genes (Supplementary Figure 5,6). In contrast, at the same early

time point, other viruses induced little changes in transcriptional profiles of infected flies

and samples clustered together with uninfected flies (Supplementary Figures 4-6). At a

more advanced time point (12 dpe) the transcriptional profiles of DCV and Bloomfield

virus-infected flies clustered together with uninfected flies, while Nora virus and DAV-

infected flies were characterized by unique transcriptional responses (Supplementary

Figure 4). By the days of median survival, all viruses had induced significant changes in

transcriptional profiles (Supplementary Figure 4).

No common differentially expressed gene (DEG) was identified across all viral

infections at any time point (Supplementary Figure 7). Furthermore, we could not identify

a consistent set of host genes responding to the infection over time, thereby illustrating

the differential impacts of viral infection on the host as the infection progresses

(Supplementary Figure 7). Even at median survival, no common transcriptional profile

was shared among all infections (Supplementary Figure 7). We also observed an increase

in the number of DEGs as infection progressed for the flies persistently infected with DAV,

DCV, and Bloomfield virus (Supplementary Figures 6-7). Many of the top DEGs identified

at early and intermediate time points are uncharacterized (Supplementary Figure 6).

Focusing on the genes known to participate in antiviral immune responses, Nora virus

elicited the highest levels of immune regulation at 1 dpe (Figure 3A). At 12 dpe, DAV had

the highest upregulation of genes associated with immune defense and stress response

pathways, indicating a potent antiviral response (Figure 3A). Interestingly, specific

immune genes upregulated during DAV infection were downregulated during Nora virus

infection, suggesting fundamentally different immune responses to these viruses (Figure

3A).

Figure 3. Transcriptomic profiles of uninfected or persistently infected flies. *A) Heatmaps expressing the Z-scores of gene expression fold change of infected populations in comparison with uninfected ones. Genes are represented in*

Analysis of genes involved in locomotion revealed significant differential expression in

Nora virus-infected flies at 1 dpe. Notably*, Lsp2,* a gene reported to be involved in motor

neuron axon and synaptic target inhibition (Inaki *et al.,* 2007), was upregulated in Nora

virus-infected samples (Figure 3B). As infection progresses, we observed an increasing

number of DEGs related to locomotion, with Nora virus infection showing an opposite

transcriptional profile compared to DAV infection (Figure 3B). At 12 dpe, Nora virus

infection upregulated genes involved in gravitaxis regulation (*CG3857*), negative

phototaxis (*Ptth*), light reception (*Rh5*), and neuronal function (*nompC*, *Pdf*) (Armstrong

et al., 2006; Yamanaka *et al.,* 2013; Sprecher and Desplan, 2008, Renn *et al.,* 1999;

Walker *et al.,* 2000). Furthermore, Nora virus infection induced the upregulation of *M6*

(involved in several neurological processes, including positive phototaxis) and the

downregulation of *fz* (neuronal function) and *Neto* (neuromuscular roles and hatching

behavior) (Kim *et al.,* 2012). Conversely, DAV-infected samples showed upregulation of

Neto, esg (related to stem cell maintenance and nervous system development) and *Mlc2*

(flight musculature) (Bahrampour *et al.,* 2017; Biteau *et al.,* 2008; Hartl *et al.,* 2011, Miller

et al., 2011). DCV and Bloomfield virus infections induced more modest differential

expression of locomotion-related genes. Taken together, these results indicate that the

host response to persistent infection is virus-specific.

Viral infection induced transcriptional changes even after viral clearance

While persistent viral infections remain prevalent at the population level, individual flies can clear the infection over time (Figure 1C; Mondote *et al.,* 2018; Nigg *et al.,* 2022). This was particularly evident for DCV, where all individuals were infected at 1 dpe, but more than half had cleared the virus by 28 dpe (Figure 1C). This provided an opportunity to examine whether viral clearance restores normal gene expression or leaves a lasting transcriptional signature.

Therefore, we analyzed the transcriptional response of flies that had cleared DCV infection by 28 dpe and found that it remained significantly altered compared to the never-infected control population (Figure 4A). Comparison of the transcriptional response of currently infected and cleared flies showed minimal differences between the two (Figure 4B), with most differentially expressed genes related to mitochondrial function. These included mitochondrial transfer RNAs and mitochondrial proteins such as *mt:ATPase8* (mitochondrial ATPase subunit), *l(3)neo43* (cytochrome c oxidase assembly), *Mcad* (mitochondrial fatty acid beta-oxidation), *mtSSB* (mitochondrial single-stranded DNA-binding protein), *ND-MWFE* (respiratory chain complex component), and *sea* (inner mitochondrial membrane carrier protein). There is a marked downregulation of mitochondrial genes in the cleared samples, which indicates that energy metabolism is upregulated during DCV infection.

Figure 4. Transcriptomic profile of flies at 28 dpe that were infected with DCV but cleared the viral infection. *A) Principal component analysis of uninfected samples, DCV Infected, and DCV Cleared. On the X-axis, we indicate the first principal component (PC1) and the percentage of variance explained by this component in brackets. On the Y-axis, we indicate the second principal component (PC2) and the percentage of variance explained by this component in brackets. B) Volcano plot indicating the differentially expressed genes between DCV Cleared samples and DCV Infected ones. Genes significantly up- (in red) or down- (in blue) regulated. In grey, we represent those genes with differential expression values lower than 1.5 log2 fold change and p-adjusted values lower than 0.05. These thresholds are indicated by dotted lines in the graphs. The names of the DEGs are shown. C) Heatmaps representing the differential expression patterns between DCV Infected and DCV Cleared samples when compared to the Uninfected of the same age. Heatmap on the right shows the top 200 differentially expressed genes in our samples in terms of absolute log2 fold change value at dpe 28. Heatmap on the left shows the top 20 DEGs at dpe 28; gene names are*

Both infected and virus-cleared flies maintained significant transcriptional differences

compared to never-infected controls, including sustained upregulation of immune

response genes (Figure 4D). These results indicate that viral infections induce long-

lasting or potentially permanent alterations in host gene expression, particularly affecting

mitochondrial and immune functions, that persist even after viral clearance.

DISCUSSION

Our study provides a comprehensive examination of how persistent viral infection

impacts key biological traits in *D. melanogaster*. By studying mono-infections with four

different enteric RNA viruses, we reveal a complex interplay between viruses and their

hosts that affects survival, reproductive output, bacteriome load, locomotor abilities,

and transcriptional profiles.

We observed a trade-off between survival and reproductive output. This pattern

suggests a compensatory mechanism in which, when faced with specific stressors, such

as viral infections, the host organism may prioritize reproduction over immunity (Pontzer

and McGrosky, 2022; Pagán *et al.,* 2008). Indeed, for DCV, DAV, and Bloomfield virus, we

noted reduced survival but increased offspring production in infected flies, indicating a

redirection of resources from surviving the infection to reproduction. Conversely, for Nora

virus infections, resources may be redirected primarily toward surviving the infection, as

flies infected by this virus exhibit the smallest impact on survival but a significant

reduction in offspring production. These diverse interactions correspond with the unique

host transcriptional changes induced by each virus. Notably, Nora virus causes a more

substantial disruption to the transcriptome, suggesting a higher immune investment in

combating the infection. This reduced survival appears to result from an acceleration of

biological aging, with our transcriptional dataset revealing that more pathogenic viruses

induce more rapid biological aging (González et al., 2025). The differing impacts on

offspring production observed across various viral infections align with previous findings

(Wallace and Obbard, 2024). For instance, it has been reported that DCV-infected flies

show increased fecundity compared to uninfected mothers (Siva-Jothy et al., 2018). Of

note, our protocol for measurement of offspring production involved transferring parental

females to fresh food daily. Whether the frequency of transfer to fresh food has an impact

on the dynamics of viral infection in the parental females is currently unknown and

therefore this caveat should be taken into account when considering our data regarding

offspring production.

The fitness costs we observed have important evolutionary implications for virus-host

dynamics. Persistent viral infections that reduce survival while altering reproductive

output could drive strong selection for resistance or tolerance mechanisms in *Drosophila*

populations (Obbard et al., 2006; Martins et al., 2014). Indeed, natural *Drosophila*

populations show considerable genetic variation in susceptibility to viral infections,

suggesting ongoing host-pathogen coevolution (Cao et al., 2017; Cogni et al., 2016;

Kutzer et al., 2023; Palmer et al., 2018). The virus-specific effects we documented may

explain why different antiviral mechanisms have evolved in insects, as each virus likely

imposes distinct selective pressures on host populations. Moreover, the persistence of

transcriptional changes even after viral clearance suggests that initial exposure events

could have lasting effects on host fitness. These findings highlight that persistent

infections may shape host evolution differently than acute infections, warranting further

investigation into the consequences of persistent viral infections on population genetics

in natural *Drosophila* communities.

The viruses studied in this work are enteric viruses, meaning that they are naturally

transmitted via the fecal-oral route, infecting the organism through the gut. Our

observations show that persistent enteric infections do not affect microbiome

composition but significantly alter the size of the bacterial community present within the

flies, which is mainly located in the gut. However, the bacteriome of laboratory-reared

flies does not fully recapitulate that of wild flies (Chandler et al., 2011). The microbiome

can modulate immune responses, nutrient absorption, and overall host physiology and

thereby affect the outcome of viral infections (Tafesh-Edwards & Eleftherianos, 2023,

González and Elena, 2021; Schissel *et al.,* 2021). Hence, some of the effects observed in

gene expression or locomotor activity could be partially explained by the microbiome

changes induced by the viral infections. However, it has been observed that for infections

with DAV (Nigg *et al.,* 2024) and Nora virus (Franchet *et al.,* 2025), the relative impact on

fly survival is largely independent of the microbiome.

The enteric viruses studied here primarily infect the gut. During DAV infection, the guts

of *D. melanogaster* exhibited overproliferation of intestinal stem cells, dysplasia, and a

loss of gut integrity. (Nigg *et al.,* 2024). However, our observations indicate that their

impacts extend beyond the gut, as seen in the locomotion results. The effects on

locomotion reflect changes in metabolic and/or neurological status (Grotewiel *et al.,*

2005) and may have profound implications for fly behavior and ecology. Impaired

locomotion can affect activities such as foraging, mate searching, and dispersal,

consequently influencing the fly's ability to colonize new locations and disseminate

associated microbes. This has important implications for disease vectoring and the

spread of pathogens. The systemic impact of enteric infections in flies has also been

observed for enteric bacteria (Deshpande *et al.,* 2022). One potential explanation for this

is the virus spreading from its primary location in the gut to other tissues. Such spreading

has been documented for DCV (Ferreira *et al.,* 2014), DAV (Nigg *et al.,* 2024), and Nora

virus (Franchet *et al.,* 2025). Further studies are required to understand the extent and

mechanisms of this viral dissemination. Another potential explanation for the systemic

effects of enteric infections is the dissemination of immune signaling molecules.

Molecules released in response to localized gut infections can have far-reaching effects

on distant organs and tissues. For instance, studies have shown that gut-derived

inflammatory signals can influence brain function and behavior, a phenomenon known

as the gut-brain axis (Cryan and Dinan, 2012; Mayer, 2011). This interorgan

communication can modulate various physiological processes, including metabolism,

immunity, and even cognitive function. Therefore, the systemic impacts of enteric viruses

may be mediated not only by direct viral spread but also by the immune signaling

molecules that circulate throughout the host.

Our analysis of activity patterns and sleep behavior further demonstrates the systemic

effects of persistent viral infections. While all viral infections preserved the flies'

fundamental circadian rhythmicity, they significantly altered sleep duration, with most

infected flies spending more time sleeping. This suggests that persistent infections affect

specific sleep regulatory pathways rather than disrupting the core circadian clock

mechanism (Allada and Chung, 2010). The increased sleep duration may represent a

host-driven response similar to the "sickness behavior" observed in vertebrates (Dantzer

and Kelley, 2007; Lopes et al., 2021), potentially serving as an energy conservation

strategy during infection. Interestingly, only Bloomfield virus-infected flies exhibited

reduced movement intensity during active periods, suggesting virus-specific effects on

neuromuscular and/or metabolic function. These findings align with previous studies

showing that immune challenges alter sleep patterns in *Drosophila* (Shirasu-Hiza et al.,

2007; Kuo et al., 2010; Ai et al., 2024) and highlight how different viral pathogens may

interact with distinct host pathways. The virus-specific alterations in sleep and activity

patterns likely reflect differences in how each virus affects energy metabolism, neural

signaling, or immune activation, further emphasizing the complex and multifaceted

nature of host-pathogen interactions. The lasting transcriptomic alterations we observed

in DCV-cleared flies provide molecular evidence for these persistent physiological

changes. The downregulation of mitochondrial genes in cleared flies compared to

currently infected flies suggests that mitochondrial reprogramming occurs during active

infection and persists after viral clearance. This shift could serve dual purposes: meeting

increased energetic demands during infection while simultaneously restricting cellular

resources available for viral replication. The importance of mitochondrial function in

infection outcomes is further supported by recent evidence that mitochondrial DNA

variants can confer broad protection against diverse pathogens, with protective

mitotypes showing upregulated mitochondrial respiration genes even in uninfected

states (Salminen et al., 2024).

Our study has certain limitations. Although infections were established at eclosion,

not all flies remained infected throughout their lifespan. While DAV, Nora virus, and

Bloomfield virus maintained high prevalence (approximately 100%, 92%, and 72% of

individuals, respectively), most individuals in the DCV-infected population cleared the

virus by the time of median survival. Nevertheless, flies that cleared DCV showed lasting

transcriptional changes, suggesting long-term effects reminiscent of post-acute

infection syndromes (Choutka *et al.,* 2022). Additionally, viral loads varied among

individuals within the same infected stock, driving variation in life history traits within

populations. However, viral load quantification requires tissue homogenization and fly

sacrifice, preventing longitudinal studies of the same individuals across behavioral and

fitness assays. Regarding measurements of offspring production, daily transfers could

theoretically reduce viral accumulation in the environment compared to a two-day

interval (as was employed for the survival curve and viral RNA accumulation

experiments), potentially influencing infection dynamics. However, our one-day protocol

was selected to maximize precision in offspring quantification with fine temporal

resolution while avoiding larval overcrowding and maintaining natural viral load

fluctuations. Finally, our study focused on females of a single genetic background,

whereas both sex and genetic background are known to influence the effects of viral

infections in *Drosophila* (Vale and Jardine, 2015; Kutzer *et al.,* 2023).

Our work presents a comprehensive multi-parameter assessment of how persistent

viral infections affect *Drosophila* biology, providing valuable reference datasets and

analytical frameworks for the research community. The primary contributions of our

study include: (1) detailed transcriptional profiles across different timepoints and

infection states that serve as reference resources for future investigations, (2) integrative

analyses connecting molecular signatures to physiological and behavioral phenotypes,

and (3) demonstrating virus-specific impacts on gut microbiome. Our study offers the

first holistic view of how different persistent infections affect interconnected biological

processes from gene expression to whole-organism fitness. The comparative datasets

we provide enable researchers to examine the relationships between transcriptional

responses, microbiome changes, and physiological outcomes across different viral

infections. These reference resources support future investigations into viral persistence

mechanisms, host-pathogen coevolution, and the broader ecological implications of

persistent infections in insect populations. In conclusion, our study demonstrates that

persistent viral infections significantly impact key biological traits in *D. melanogaster*.

Our work adds to the growing evidence that persistent infections profoundly shape host

fitness and physiology through complex and diverse mechanisms.

MATERIAL AND METHODS

Fly stocks

For all experiments we used *Wolbachia*-free w^{1118} flies maintained on a standard cornmeal diet. This diet was prepared by combining 440g inactive dry yeast, 440g corn meal, and 60g agar in 6L osmotic water. The mixture was autoclaved and after cooling, 150ml moldex solution (20% methylhydroxybenzoate) and 29ml propionic acid were added as preservatives. All experiments were done at 25°C under a 12:12 light:dark cycle. All fly lines were cleaned of possible persistent infections (viruses and Wolbachia) as described previously (Merkling & van Rij, 2015).

Viral infections

Persistently infected w^{1118} flies were established following the method described by Nigg and colleagues in 2022 and are the same stocks used in the work of Nigg and colleagues (2024). Persistent infections of four RNA viruses were established in w^{1118} *Drosophila melanogaster* using tailored methodologies. For DCV, a persistently infected bw1;st1 Ago3t3/TM6B;Tb+ line (Bloomington #28270) was used to environmentally transfer DCV to naïve w^{1118} flies. Infected donors were housed in a fresh vial for 3 days; after their removal, w^{1118} flies were exposed to the contaminated environment for 3 days. The F0 generation was transferred to a new vial, and their F1 progeny were confirmed as persistently infected. For DAV, the Australian isolate (DAV_{HD} from the van Rij lab) was injected into w^{1118} adults (50 nl/fly). Identical environmental transfer followed: injected flies contaminated a vial for 3 days before naïve w^{1118} exposure. Persistently infected F1 progeny were isolated and maintained. Bloomfield virus was identified in a contaminated Dipt-GFP stock (BDSC Cat#55709). Homogenate from these flies was filtered (0.22 µm)

and injected into w^{1118} adults. Injected F0 flies were removed after 3 days; F1 and F2

progeny underwent sequential generational passages (5–9 days/vial) to establish

persistence in F3 adults confirmed via RNA sequencing. The Nora virus-persistent line

originated from a w^{1118} line persistently infected with Nora virus that was gifted by Dr.

Stefan Ameres. Upon receipt, we repeatedly backcrossed the infected w^{1118} flies to the

w^{1118} line maintained in our laboratory. Subsequently, we confirmed that the backcrossed

flies retained persistent infection with Nora virus by RNA sequencing. The stocks were

maintained at 25°C under a 12:12 light:dark cycle.

The presence or absence of persistent infections was determined by RT-PCR with

specific primers for Nora virus, Drosophila A virus, DCV (Supplementary Table 2). In

addition, total RNA from persistently infected adults was extracted and sequenced. The

resulting reads were mapped to a database of all known *Drosophila* viruses

(https://obbard.bio.ed.ac.uk/data.html) confirming that only the desired virus was

present (Nigg et al., 2022, 2024).

Survival analysis

Survival analyses of the flies coming from persistently infected stocks were conducted

using 20 female flies/replicate sorted at 1 dpe. Male and female flies were collected on

the day of eclosion and left to mate for 24 hours. Subsequently, only female flies were

sorted in new tubes containing 20 females each. Each tube was considered a biological

replicate, and a total of nine biological replicates (from three independent experiments)

were analyzed. Survival was monitored daily by counting the number of dead flies in each

vial, and flies were transferred to new vials every two days. Fly survival was assessed from

the day of adult eclosion until death.

Viral quantification

Individual female flies were collected in 300 µl of TRIzol reagent (Invitrogen, 15596026).

The samples were homogenized using a pestle, and RNA was extracted following

manufacturer's instructions, supplemented with GlycoBlue (Invitrogen, AM9516) during

the precipitation process. RNA concentrations were measured using the Qubit RNA BR

Assay Kit (Invitrogen, Q10211). Following extraction, 4µl of each RNA sample were

treated with DNAse I (Roche, 04716728001), and 2µl of the final volume were used in the

RT reaction. RT was performed using random primers and Maxima H Minus Reverse

Transcriptase (Thermo scientific, EP0751) following manufacturer's instructions. The

resulting cDNA was diluted 1:10 with MilliQ water. The qPCR was performed using

Luminaris Color HiGreen qPCR Master Mix (Thermo Scientific, K0374) in a QuantStudio 7

flex machine (Thermo Scientific). The reactions were done in triplicates of 10 µl each and

using the corresponding set of primers targeting a replication protein for each virus

(Supplementary Table 1). The cycling conditions were as follows: 2 minutes at 50°C, 10

minutes at 95°C, followed by 40 cycles of 15 seconds at 95°C and 60 seconds at 60°C. A

standard melt curve analysis was performed after the cycling protocol. For the analysis

of the results, the threshold for viral RNA detection was set to 35 Ct (i.e. if a sample had

Ct higher than 35 it was considered non infected). The Ct values of the virus for a sample

were normalized against the Ct value of the housekeeping gene *Rp49* for that sample

(ΔCt). The $2^{-\Delta Ct}$ values were calculated per sample and their $\log_{10}$ plotted.

Viable offspring production

To assess viable offspring production, 20 female and 20 male flies from infected or

uninfected stocks were left to mate in clean fly vials for 24 hours after eclosion. After this

period, males were discarded, and the mated females were transferred to new tubes

daily until no more progeny was produced (17 days). All tubes were kept at 25 °C under a

12:12 light:dark cycle, and the number of adult flies (i.e. the viable progeny) in each tube

was counted after 14 days. The number of offspring was normalized to the number of

adult females placed in each tube daily to account for those that did not survive

throughout the test duration (Supplementary File 2).

Climbing assays

Climbing assays were performed following the method described by Barone and

Bohmann (2013). Briefly, a 9 cm empty tube was used, and 20 female flies were placed

inside. With the cotton cap on, tubes had a maximum climbable height of 7.2 cm. The

tubes were tapped three times over the span of three seconds, and the climbing height

was assessed by taking a picture 3 seconds after the last tap. Flies' height was assessed

manually on the pictures using a metric guide on the back of the tubes. For categorical

classification, flies were considered "Top" if they surpassed 6.2 cm, "Bottom" if they were

between 0.0-1.0 cm, and "Middle" if they were between 1.0 cm and 6.2 cm.

Behavioral analysis

We conducted behavioral analyses using DAM5H Drosophila Activity Monitors

(Trikinetics, Princeton, MA, USA). Using the DAM5H we tracked the locomotor activity of

individual flies housed in a 5mm-diameter, 80mm-length tube. The system records

activity when a fly crosses one of 15 infrared beams of each tube. The monitor quantifies

movement by registering infrared beam disruptions as activity "counts." A fly that moves

continuously through a beam zone or returns to the same beam multiple times generates

several counts, whereas a fly that simply enters a beam area and stops moving is

recorded as only a single count. The count data were registered per minute and were used

to evaluate the locomotion activity and to identify sleep periods, which were defined as

immobility bouts lasting at least 5 minutes. All behavioral experiments utilized 12-day-

old mated female flies. Flies were individually housed in DAM tubes containing standard

Bloomington cornmeal diet. All experiments were conducted at 25°C under a 12:12

light:dark cycle. To allow for acclimation, flies were placed in the monitoring tubes one

hour before the start of a dark cycle. Data collection began at the onset of the following

light cycle (designated as T0) and continued for three complete cycles (until T2). We

performed three independent experiments, each including 8-12 flies per experimental

condition.

Microbiome composition

Female flies from uninfected or persistently infected stocks were collected at 12 dpe.

External microbes were removed by surface sterilization using subsequent washes in 10

% bleach for 5 min, 70 % ethanol for 5 min and three rinsing steps in sterile water. Since

the fly microbiota is fairly stable in populations reared together, microbiota composition

was analyzed in three pools of 20 flies per condition. Flies were placed in a lysis buffer

(10 mM Tris-HCl (pH 8), 26 mM EDTA, 0.5% SDS, 5 mg/mL lysozyme) and homogenized

with sterile pestles. Blank samples consisting of lysis buffer were processed alongside

the flies and served as extraction negative controls. After incubating the lysates for 1 h at

37 °C, total DNA was extracted using the DNeasy Blood and Tissue Kit (Qiagen) following

the manufacturer's instruction. For each sample, 5 µL of undiluted genomic DNA was

used as template in a 20 µL final reaction volume for amplification of the 16S rRNA gene.

The primers containing locus specific sequence 8/27_Forward and 1492_Reverse were

employed (Supplementary Table 1). PCR amplification was carried out for a total of 35

cycles. No visible amplicons were obtained from extraction negative controls. The

resulting amplicons were used for library preparation with the Ligation Sequencing Kit

V14 (SQK-LSK114, Oxford Nanopore Technologies) following the manufacturer's

instructions. Subsequently the ONT DNA libraries were sequenced on a PromethION 2

Integrated apparatus using a FLO-PRO114M flowcell. The raw ONT sequencing data was

base-called (super-accurate algorithm), demultiplexed and sequencing adaptor

trimmed using the software dorado (version 7.3.11). The produced long-reads in fastq

format were quality checked using the software NanoPlot (version 1.42.0) and then

quality filtered using the software filtlong (version 0.2.1), discarding any long-reads with

an average quality score below Q17 and read lengths not in the range of 1200-2300bp.

The surviving long-reads were in-silico PCR amplified to verify the locus using the usearch

software (version v11.0.667), then clustered using the software isONclust (version

0.0.6.1). For each cluster a consensus sequence was generated to represent it as an

operational taxonomic unit (OTU). OTU sequences were compared to the reference

sequences   of   the   NCBI   targeted   loci   DB

(https://www.ncbi.nlm.nih.gov/refseq/targetedloci/ - downloaded 15. Feb. 2024). Taxa

were predicted and confidences were calculated using the SINTAX algorithm

implemented in the usearch software (version v11.0.667). DNA library construction,

sequencing, read processing and taxonomy identification described in these sections

were performed by Microsynth AG (Balgach, Switzerland). Alpha and beta diversity

measures were estimated using the Chao1 and Shannon indices (alpha diversity) and the

Bray-Curtis dissimilarity (beta diversity) in R using the phyloseq package (version 1.42.0).

Both alpha and beta diversities were calculated on microbiota composition at the OTU

level. Pairwise Wilcoxon tests were used to determine differences in alpha diversity

measures among the different fly lines, while PERMANOVA was used to determine the

effect of the infection status on Bray-Curtis dissimilarities. The barplot was produced in

Prism (version 10.4.2). As *Lactiplantibacillus* OTU sequences were characterized by a

sequence identity > 99%, we decided to display the bacterial genus without indicating

the bacterial species. Raw sequencing reads will be made available upon publication.

754   Bacteriome load

Female flies from uninfected or persistently infected stocks were collected at 12 dpe.

External microbes were removed as described above. 10 to 20 flies per condition were

collected in two independent experiments. Total DNA was extracted as described above.

Bacterial DNA was quantified via amplification of the *16S* rRNA bacterial gene and the fly

ribosomal gene *Rp49* via qPCR using the Luminaris Color HiGreen qPCR Master Mix

(Thermo Scientific, K0374) in a QuantStudio 7 flex machine (Thermo Scientific). Primer

sequences are listed in Supplementary Table 1. Ct values were retained only if they were

lower than those of both no-template and extraction blank controls. Total bacterial load

was calculated via the ΔCt method.

765   Transcriptome profiling by high-throughput sequencing

Raw sequencing reads can be found in the Sequence Read Archive under the BioProject

PRJNA1235228. For each condition, four samples were sequenced. Each sample was

made by pooling RNA from five individuals (3 ng of total RNA per individual), except for

DCV at 28 dpe, where the four pools were made from three individuals each. RNA-seq

libraries were prepared from 15 ng of pooled RNA using an NEBNext Ultra II RNA Library

Prep Kit for Illumina (New England Biolabs, E7770L) with NEBNext Multiplex Oligos for

Illumina (Dual Index Primers Set 1) (New England Biolabs, E7600S). All sequencing was

performed on an Illumina NextSeq 500 instrument using a NextSeq 500/550 High Output

Kit v2.5 (75 cycles) (Illumina, 20024906). Raw sequencing reads were mapped to the *D.*

melanogaster genome release dmel-r6.56 with STAR version 2.7.11b (Dobin et al 2013).

Feature counting was performed with HTSeq version 0.11.2. (Putri et al. 2022), and

differential gene expression analysis was performed with DESeq2 in R Studio (version

2023.03.1). Differential gene expression analysis results can be found in Supplementary

File 3. Log2 fold changes were transformed into Z-scores using the "ashr" package, and

only adjusted p-values were considered for analysis.

Statistical analysis

All statistical analyses were conducted using R version 3.6.1 within the Rstudio

development environment version 2024.04.2+764. The level of significance was set at *P*

< 0.05 and symbols indicate the significance of differences between conditions: *** *P* <

0.001; ** *P* < 0.01; * 0.01 < *P* < 0.05.

To account for the block effect, we employed linear mixed-effects or general linear

mixed-effects models. Generalized linear models (GLM) were fitted to the data using the

g*lm f*unction from the "stats" package. Generalized linear mixed models (GLMM) were

fitted to the data using the *lmer f*unction from the "lme4" package version 1.1-35.5.

Cumulative link mixed models (CLMM) using the *clmm* function from the "ordinal"

package version 2023.12-4.1. *Post-hoc* pairwise comparisons were performed using the

emmeans function from the "emmeans" package version 1.10.5. Tukey's or FDR

correction were used to adjust for multiple comparisons. The letter-based grouping for

pair-wise comparisons was created using the the *cld* function from the "multcomp"

package version 1.4-26. The specific statistical methods applied to each figure are

detailed in the legend of the corresponding figure.

FUNDING

This work was supported by funding from the French Government's Investissement

d'Avenir program, Laboratoire d'Excellence Integrative Biology of Emerging Infectious

Diseases (grant ANR-10-LABX-62-IBEID), the Agence Nationale de la Recherche (grant

ANR-23-CE15-0038-01, INFINITESIMAL), Fondation iXcore - iXlife - iXblue Pour La

Recherche and the Explore Donation, MIE project to M.-C.S. R.G. salary is supported by

a Pasteur-Roux-Cantarini fellowship of Institut Pasteur. This project has received funding

from the European Union's Horizon 2020 research and innovation program under the

Marie Skłodowska-Curie grant agreement No 101024099 to J.C.N. The funders had no

role in study design, data collection and analysis, decision to publish, or preparation of

the manuscript

ACKNOWLEDGEMENTS

We thank all members of the Saleh lab for their insightful discussions. We are especially

grateful to Vanesa Mongelli, whose early conversations sparked this project and who

guided Mauro through his first steps in the Drosophila field.

AUTHOR CONTRIBUTIONS

M.C.-S., R.G., O.R., J.C.N., and M.-C.S designed the experiments; M.C.-S., R.G. O.R., H.B.,
and J.C.N. performed the experiments; M.C.-S., R.G., O.R., J.C.N., and M.-C.S analyzed
the data; M.C.-S., and R.G., wrote the first draft with inputs from O.R., J.C.N., and M.-C.S.
All authors reviewed and approved the final version of the manuscript.

REFERENCES

Ai X, Deng H, Li X, Wei Z, Chen Y, Yin T, et al. cGAS-like receptors drive a systemic STING-dependent host response in Drosophila. Cell Rep. 2024; 43: 115081. doi:10.1016/j.celrep.2024.115081

Allada R, Chung BY. Circadian Organization of Behavior and Physiology in Drosophila. Annu Rev Physiol. 2010; 72: 605–624. doi:10.1146/annurev-physiol-021909-135815

Armstrong JD, Texada MJ, Munjaal R, Baker DA, Beckingham KM. Gravitaxis in *Drosophila melanogaster*: a forward genetic screen. Genes Brain Behav. 2006; 5: 222–239. doi:10.1111/j.1601-183X.2005.00154.x

Bahrampour S, Gunnar E, Jonsson C, Ekman H, Thor S. Neural Lineage Progression Controlled by a Temporal Proliferation Program. Dev Cell. 2017; 43: 332-348.e4. doi:10.1016/j.devcel.2017.10.004

Barone MC, Bohmann D. Assessing Neurodegenerative Phenotypes in *Drosophila* Dopaminergic Neurons by Climbing Assays and Whole Brain Immunostaining. JoVE. 2013; 50339. doi:10.3791/50339

Biteau B, Hochmuth CE, Jasper H. JNK Activity in Somatic Stem Cells Causes Loss of Tissue Homeostasis in the Aging *Drosophila* Gut. Cell Stem Cell. 2008; 3: 442–455. doi:10.1016/j.stem.2008.07.024

Cao C, Cogni R, Barbier V, Jiggins FM. Complex Coding and Regulatory Polymorphisms in a Restriction Factor Determine the Susceptibility of *Drosophila* to Viral Infection. Genetics. 2017; 206: 2159–2173. doi:10.1534/genetics.117.201970

Chandler, J. A., Morgan Lang, J., Bhatnagar, S., Eisen, J. A., & Kopp, A. Bacterial

Communities of Diverse Drosophila Species: Ecological Context of a Host–Microbe

Model System. PLoS Genet. 2011; 7(9), e1002272.

doi.org/10.1371/journal.pgen.1002272

Choutka J, Jansari V, Hornig M, Iwasaki A. Unexplained post-acute infection

syndromes. Nat Med. 2022; 28: 911–923. doi:10.1038/s41591-022-01810-6

Cogni, R., Cao, C., Day, J. P., Bridson, C., & Jiggins, F. M. The genetic architecture of

resistance to virus infection in Drosophila. Mol Ecol. 2016; 25(20), 5228–5241.

doi.org/10.1111/mec.13769

Crone EE. Is survivorship a better fitness surrogate than fecundity? Evolution. 2001;

55: 2611–2614. doi:10.1111/j.0014-3820.2001.tb00773.x

Cryan JF, Dinan TG. Mind-altering microorganisms: the impact of the gut microbiota on

brain and behaviour. Nat Rev Neurosci. 2012; 13: 701–712. doi:10.1038/nrn3346

Dantzer R, Kelley KW. Twenty years of research on cytokine-induced sickness

behavior. Brain Behav Immun. 2007; 21: 153–160. doi:10.1016/j.bbi.2006.09.006

Della Fera, A. N., Warburton, A., Coursey, T. L., Khurana, S., & McBride, A. A. Persistent

Human Papillomavirus Infection. Viruses. 2021; 13(2), 321. doi.org/10.3390/v13020321

Deshpande R, Lee B, Grewal SS. Enteric bacterial infection in *Drosophila* induces

whole-body alterations in metabolic gene expression independently of the immune

deficiency signaling pathway. G3. 2022; 12: jkac163. doi:10.1093/g3journal/jkac163

Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, et al. STAR: ultrafast

universal   RNA-seq   aligner.   Bioinform.   2013;   29:   15–21.

doi:10.1093/bioinformatics/bts635

Ferreira ÁG, Naylor H, Esteves SS, Pais IS, Martins NE, Teixeira L. The Toll-Dorsal

Pathway Is Required for Resistance to Viral Oral Infection in *Drosophila*. Schneider DS,

editor. PLoS Pathog. 2014; 10: e1004507. doi:10.1371/journal.ppat.1004507

Franchet A, Haller S, Yamba M, Barbier V, Thomaz-Vieira A, Leclerc V, et al. Nora virus

proliferates in dividing intestinal stem cells and sensitizes flies to intestinal infection and

oxidative stress. bioRxiv. 2025; doi:10.1101/2025.01.30.635658

Gargano J, Martin I, Bhandari P, Grotewiel M. Rapid iterative negative geotaxis (RING):

a new method for assessing age-related locomotor decline in *Drosophila*. Exp Gerontol.

2005; 40: 386–395. doi:10.1016/j.exger.2005.02.005

Goic B, Saleh M-C. Living with the enemy: viral persistent infections from a friendly

viewpoint. Curr Opin Microbiol. 2012; 15: 531–537. doi:10.1016/j.mib.2012.06.002

González R, Elena SF. The Interplay between the Host Microbiome and Pathogenic

Viral Infections. mBio. 2021;12: e02496-21. doi:10.1128/mBio.02496-21

González R, Castelló-Sanjuán M, Saleh M-C. Viral infections reduce *Drosophila*

lifespan through accelerated aging. bioRxiv. 2025; doi:10.1101/2025.03.13.643076

Gould AL, Zhang V, Lamberti L, Jones EW, Obadia B, Korasidis N, et al. Microbiome

interactions   shape   host   fitness.   Proc   Natl   Acad   Sci   USA.   2018;   115.

doi:10.1073/pnas.1809349115

Grotewiel MS, Martin I, Bhandari P, Cook-Wiens E. Functional senescence in
Drosophila melanogaster. Ageing Res Rev. 2005; 4: 372–397.
doi:10.1016/j.arr.2005.04.001

Hartl M, Loschek LF, Stephan D, Siju KP, Knappmeyer C, Kadow ICG. A New Prospero
and microRNA-279 Pathway Restricts CO_2 Receptor Neuron Formation. J Neurosci.
2011; 31: 15660–15673. doi:10.1523/JNEUROSCI.2592-11.2011

Ho T, Tzanetakis IE. Development of a virus detection and discovery pipeline using next
generation sequencing. Virology. 2014; 471–473: 54–60. doi:10.1016/j.virol.2014.09.019

Hogenhout, S. A., Ammar, E.-D., Whitfield, A. E., & Redinbaugh, M. G. (2008). Insect
Vector Interactions with Persistently Transmitted Viruses. Annual Review of
Phytopathology, 46(1), 327-359. https://doi.org/10.1146/annurev.phyto.022508.092135

Hou, X., He, Y., Fang, P., Mei, S.-Q., Xu, Z., Wu, W.-C., Tian, J.-H., Zhang, S., Zeng, Z.-Y.,
Gou, Q.-Y., Xin, G.-Y., Le, S.-J., Xia, Y.-Y., Zhou, Y.-L., Hui, F.-M., Pan, Y.-F., Eden, J.-S., Yang,
Z.-H., Han, C., … Shi, M. Using artificial intelligence to document the hidden RNA
virosphere. Cell. 2024; 187(24), 6929-6942.e16.
https://doi.org/10.1016/j.cell.2024.09.027

Inaki M, Yoshikawa S, Thomas JB, Aburatani H, Nose A. Wnt4 Is a Local Repulsive Cue
that Determines Synaptic Target Specificity. Curr Biol. 2007; 17: 1574–1579.
doi:10.1016/j.cub.2007.08.013

Irschick DJ, Garland T. Integrating Function and Ecology in Studies of Adaptation:
Investigations of Locomotor Capacity as a Model System. Annu Rev Ecol Syst. 2001; 32:
367–396. doi:10.1146/annurev.ecolsys.32.081501.114048

Jia Y, Jin S, Hu K, Geng L, Han C, Kang R, et al. Gut microbiome modulates *Drosophila*

aggression through octopamine signaling. Nat Commun. 2021; 12: 2698.

doi:10.1038/s41467-021-23041-y

Kapun M, Barrón MG, Staubach F, Obbard DJ, Wiberg RAW, Vieira J, et al. Genomic

Analysis of European *Drosophila melanogaster* Populations Reveals Longitudinal

Structure, Continent-Wide Selection, and Previously Unknown DNA Viruses. Falush D,

editor. Mol Biol Evol. 2020; 37: 2661–2678. doi:10.1093/molbev/msaa120

Kim Y-J, Bao H, Bonanno L, Zhang B, Serpe M. *Drosophila* Neto is essential for

clustering glutamate receptors at the neuromuscular junction. Genes Dev. 2012;26: 974–

987. doi:10.1101/gad.185165.111

Kuo T-H, Pike DH, Beizaeipour Z, Williams JA. Sleep triggered by an immune response

in Drosophila is regulated by the circadian clock and requires the NFκB Relish. BMC

Neurosci. 2010; 11: 17. doi:10.1186/1471-2202-11-17

Kutzer MAM, Gupta V, Neophytou K, Doublet V, Monteith KM, Vale PF. Intraspecific

genetic variation in host vigour, viral load and disease tolerance during Drosophila C virus

infection. Open Biol. 2023; 13: 230025. doi:10.1098/rsob.230025

Lee, W.-S., Webster, J. A., Madzokere, E. T., Stephenson, E. B., & Herrero, L. J. Mosquito

antiviral defense mechanisms: A delicate balance between innate immunity and

persistent viral infection. Parasit Vectors. 2019 ; 12(1), 165. doi.org/10.1186/s13071-019-

3433-8

Martins NE, Faria VG, Nolte V, Schlötterer C, Teixeira L, Sucena É, et al. Host adaptation

to viruses relies on few genes with different cross-resistance properties. Proc Natl Acad

Sci USA. 2014; 111: 5938–5943. doi:10.1073/pnas.1400378111

Mayer EA. Gut feelings: the emerging biology of gut–brain communication. Nat Rev

Neurosci. 2011; 12: 453–466. doi:10.1038/nrn3071

Miller MS, Farman GP, Braddock JM, Soto-Adames FN, Irving TC, Vigoreaux JO, et al.

Regulatory Light Chain Phosphorylation and N-Terminal Extension Increase Cross-Bridge

Binding and Power Output in *Drosophila* at In Vivo Myofilament Lattice Spacing. Biophys

J. 2011; 100: 1737–1746. doi:10.1016/j.bpj.2011.02.028

Mondotte JA, Gausson V, Frangeul L, Blanc H, Lambrechts L, Saleh M-C. Immune

priming and clearance of orally acquired RNA viruses in *Drosophila*. Nat Microbiol. 2018;

3: 1394–1403. doi:10.1038/s41564-018-0265-9

Nigg JC, Castelló-Sanjuán M, Blanc H, Frangeul L, Mongelli V, Godron X, et al. Viral

infection disrupts intestinal homeostasis via Sting-dependent NF-κB signaling in

Drosophila. Curr Biol. 2024; 34: 2785-2800.e7. doi:10.1016/j.cub.2024.05.009

Nigg JC, Mongelli V, Blanc H, Saleh M-C. Innovative Toolbox for the Quantification of

Drosophila C Virus, Drosophila A Virus, and Nora Virus. J Mol Biol. 2022; 434: 167308.

doi:10.1016/j.jmb.2021.167308

Lopes PC, French SS, Woodhams DC, Binning SA. Sickness behaviors across

vertebrate taxa: proximate and ultimate mechanisms. J Exp Biol. 2021; 224: jeb225847.

doi:10.1242/jeb.225847

Obbard DJ, Jiggins FM, Halligan DL, Little TJ. Natural Selection Drives Extremely Rapid

Evolution   in   Antiviral   RNAi   Genes.   Curr   Biol.   2006;   16:   580–585.

doi:10.1016/j.cub.2006.01.065

Obbard, D. J., Shi, M., Roberts, K. E., Longdon, B., & Dennis, A. B. A new lineage of

segmented   RNA   viruses   infecting   animals.   Virus   Evol.   2020;   6(1),   vez061.

doi:10.1093/ve/vez061

Pagán I, Alonso-Blanco C, García-Arenal F. Host Responses in Life-History Traits and

Tolerance to Virus Infection in *Arabidopsis thaliana*. Holmes EC, editor. PLoS Pathog.

2008; 4: e1000124. doi:10.1371/journal.ppat.1000124

Palmer W, Varghese F, Van Rij R. Natural Variation in Resistance to Virus Infection in

Dipteran Insects. Viruses. 2018; 10: 118. doi:10.3390/v10030118

Pontzer H, McGrosky A. Balancing growth, reproduction, maintenance, and activity in

evolved   energy   economies.   Curr   Biol.   2022;   32:   R709–R719.

doi:10.1016/j.cub.2022.05.018

Putri GH, Anders S, Pyl PT, Pimanda JE, Zanini F. Analysing high-throughput sequencing

data   in   Python   with   HTSeq   2.0.   Bioinform.   2022;   38:   2943–2945.

doi:10.1093/bioinformatics/btac166

Renn SCP, Park JH, Rosbash M, Hall JC, Taghert PH. A pdf Neuropeptide Gene Mutation

and Ablation of PDF Neurons Each Cause Severe Abnormalities of Behavioral Circadian

Rhythms in *Drosophila*. Cell. 1999; 99: 791–802. doi:10.1016/S0092-8674(00)81676-1

Salas-Benito, J. S., & De Nova-Ocampo, M. (2015). Viral Interference and Persistence

in Mosquito-Borne Flaviviruses. Journal of Immunology Research, 2015, 1-14.

https://doi.org/10.1155/2015/873404

Salminen TS, Vesala L, Basikhina Y, Kutzer M, Tuomela T, Lucas R, et al. A naturally

occurring mitochondrial genome variant confers broad protection from infection in

Drosophila. PLoS Genet. 2024; 20: e1011476. doi:10.1371/journal.pgen.1011476

Schissel M, Best R, Liesemeyer S, Tan Y-D, Carlson DJ, Shaffer JJ, et al. Effect of Nora

virus infection on native gut bacterial communities of *Drosophila melanogaster*. AIMS

Microbiol. 2021; 7: 216–237. doi:10.3934/microbiol.2021014

Shi, M., Lin, X.-D., Tian, J.-H., Chen, L.-J., Chen, X., Li, C.-X., Qin, X.-C., Li, J., Cao, J.-P.,

Eden, J.-S., Buchmann, J., Wang, W., Xu, J., Holmes, E. C., & Zhang, Y.-Z. Redefining the

invertebrate RNA virosphere. Nature. 2016; 540(7634), 539–543.

doi.org/10.1038/nature20167

Shin SC, Kim S-H, You H, Kim B, Kim AC, Lee K-A, et al. *Drosophila* Microbiome

Modulates Host Developmental and Metabolic Homeostasis via Insulin Signaling.

Science. 2011; 334: 670–674. doi:10.1126/science.1212782

Shirasu-Hiza MM, Dionne MS, Pham LN, Ayres JS, Schneider DS. Interactions between

circadian rhythm and immunity in *Drosophila melanogaster*. Curr Biol. 2007; 17: R353–

R355. doi:10.1016/j.cub.2007.03.049

Siva-Jothy, J. A., Monteith, K. M., & Vale, P. F. Navigating infection risk during oviposition

and cannibalistic foraging in a holometabolous insect. Behav Ecol. 2018; 29(6), 1426–

1435. doi.org/10.1093/beheco/ary106

Sprecher SG, Desplan C. Switch of rhodopsin expression in terminally differentiated

Drosophila sensory neurons. Nature. 2008; 454: 533–537. doi:10.1038/nature07062

Tafesh-Edwards G, Eleftherianos I. The role of *Drosophila* in gut homeostasis and

immunity. Gut Microbes. 2023; 15: 2208503. doi:10.1080/19490976.2023.2208503

Vale PF, Jardine MD. Sex-specific behavioural symptoms of viral gut infection and

Wolbachia in *Drosophila melanogaster*. J Insect Physiol. 2015; 82: 28–32.

doi:10.1016/j.jinsphys.2015.08.005

Walker RG, Willingham AT, Zuker CS. A *Drosophila* Mechanosensory Transduction

Channel. Science. 2000; 287: 2229–2234. doi:10.1126/science.287.5461.2229

Wallace MA, Coffman KA, Gilbert C, Ravindran S, Albery GF, Abbott J, et al. The

discovery, distribution, and diversity of DNA viruses associated with *Drosophila*

melanogaster in Europe. Virus Evol. 2021; 7: veab031. doi:10.1093/ve/veab031

Wallace MA, Obbard DJ. Naturally occurring viruses of *Drosophila* reduce offspring

number and lifespan. Proc R Soc B. 2024; 291: 20240518. doi:10.1098/rspb.2024.0518

Webster, C. L., Longdon, B., Lewis, S. H., & Obbard, D. J. Twenty-Five New Viruses

Associated with the Drosophilidae (Diptera). Evol Bioinform. 2016; 12s2, EBO.S39454.

doi: 10.4137/EBO.S39454

Webster, C. L., Waldron, F. M., Robertson, S., Crowson, D., Ferrari, G., Quintana, J. F.,

Brouqui, J.-M., Bayne, E. H., Longdon, B., Buck, A. H., Lazzaro, B. P., Akorli, J., Haddrill, P.

R., & Obbard, D. J. The Discovery, Distribution, and Evolution of Viruses Associated with

Drosophila melanogaster. PLoS Biol. 2015; 13(7), e1002210.

doi.org/10.1371/journal.pbio.1002210

Williams, T., Virto, C., Murillo, R., & Caballero, P. (2017). Covert Infection of Insects by

Baculoviruses.   Frontiers   in   Microbiology,   8,   1337.

https://doi.org/10.3389/fmicb.2017.01337

Xu, P., Chen, F., Mannas, J. P., Feldman, T., Sumner, L. W., & Roossinck, M. J. Virus

infection improves drought tolerance. New Phytol. 2008; *180*(4), 911–921.

doi.org/10.1111/j.1469-8137.2008.02627.x

Yamanaka N, Romero NM, Martin FA, Rewitz KF, Sun M, O'Connor MB, et al.

Neuroendocrine Control of *Drosophila* Larval Light Preference. Science. 2013; 341:

1113–1116. doi:10.1126/science.1241210

SUPPLEMENTARY FIGURES

Supplementary Figure 1. *Impact of persistent viral infection on offspring production. Daily viable offspring per female. Number of viable offspring produced by female flies uninfected (green) or persistently infected with DAV (red), DCV (yellow), Bloomfield virus (dark blue) or Nora virus (purple) after one day of mating (20 females paired with 20 males). After one day, the males were discarded, and the females were transferred to new vials with food each day. Mated flies were kept for subsequent days, and the viable offspring were counted. The number of viable offspring was normalized by the number of alive females eggs in each tube. Day 0 represents the day of eclosion and mating. Experiments were conducted on 20 females paired with 20 males in 9 replicates. Significance was calculated using a general linear-mixed model where the time point was the fixed factor and experiments a random effect; Tukey contrasts were used for post-hoc analyses. Significance values are *** P < 0.001, ** P < 0.01, * P < 0.05.*

1022

1023

1024

Supplementary Figure 2.*Impact of persistent viral infection on locomotion. Climbing assay results of uninfected flies (green) or flies persistently infected with DAV (red), DCV (yellow), Bloomfield virus (dark blue) or Nora virus (purple). A) Quantitative height climbed by individual flies. Climbing assay results for days 1, 6, 12, and 20 post-eclosion. Results of 9 different biological replicates with 20 female flies. At each time point and replicate, six pictures were taken, and the height of each fly in the tube was recorded. Each symbol on the graph represents the recorded height of a fly for each picture. Boxplots are shown on the left side of each stock, indicating the median and interquartile range of the heights climbed by the flies. The dotted line represents the median height climbed by the uninfected population. The half-violin of each plot shows the density of the values. Significance was calculated using a general linear-mixed model where the virus status and the time point were the fixed factors and experiments a random*

effect; Tukey contrasts were used for post-hoc analyses. **B)** *Qualitative measurements of the climbing assay. Top (green) indicates the flies present in the top 1 cm of the climbing tubes (6.2-7.2 cm). Bottom (red) indicates the percentage of flies located in the bottom 1 cm of the tube (0-1 cm), and middle (blue) includes all flies that climbed between (1-6.2 cm). Significance was calculated using a cumulative link mixed model where the virus status and the time point were the fixed factors and experiments a random effect; false discovery rate contrasts were used for post-hoc analyses. For all the figure, *** P < 0.001, ** P < 0.01, * P < 0.05.*

1025

1026

Supplementary Figure 3. Bacteriome d*iversity analyses of pools containing at least 20 female flies per condition (uninfected or mono-infected). **A)** Alpha diversity analyses on Chao1 and Shannon indexes reveal no significant differences in the species diversity of uninfected or persistently infected flies (pairwise Wilcoxon test on Chao1 and Shannon indexes, P > 0.46 for all comparisons). **B)** Principal component analysis of the Bray-Curtis dissimilarity matrix of the different studied fly populations at the OTU level. This analysis does not identify any specific differences in the bacterial community structure among the various fly lines (PERMANOVA on infection, F = 1.57, P = 0.24).*

Supplementary Figure 4. *Principal Component Analysis of the transcriptional responese of the different samples sequenced. These samples consist on pools of RNA coming from individual Drosophila females from the uninfecetd and infected stocks. Infection was confirmed through RT-qPCR. All samples in a pool (3-5) come from the same collection dpe and stock. On the Y axis of every graph, we indicate the second principal component (PC2) and the percentage of variance explained by this component between brackets. On the X axis, we indicate the first principal component (PC1) and the percentage of variance explained by this component between brackets. Black lines indicate the origin of each axis. The do represent our fly stocks throughout the different days post-eclosion, in green, we have the Uninfected stock, in burgundy, the Drosophila A virus persistent flies, in yellow, the Drosophila C virus*

persistently infected flies, in dark blue, we have the Bloomfield virus persistently infected stock, and in lavender the Nora virus persistently infected flies.

1028

1029

Supplementary Figure 5. *Volcano plots of the differentially expressed genes between the infected samples versus the uninfected flies. Transcriptomic results indicate differentially expressed genes in our infected samples. Volcano plots showing the number of genes significantly up- (in red) or down- (in blue) regulated. In grey, we represent those genes displaying differential expression values lower than 1.5 log2 fold change and/or with p-adjusted values lower than 0.05. Gene names are indicated fot the top 10 genes in absolute Log2FoldChange values. These thresholds are*

1030

Supplementary Figure 6. *Top differentially expressed genes in the infected samples over the uninfected samples. Persistent viral infection alters Drosophila melanogaster's transcriptomic profile in a virus-independent manner. For all heatmaps, on the X-axis, we indicate the infected stock analysed. **A)** From left to right: heatmap showing the top 200 differentially expressed genes in our samples in terms of absolute log2 fold change value at dpe1. Heatmap showing the top 20 DEGs at dpe1. **B)** From left to right: heatmap showing the top 200 differentially expressed genes in our samples in terms of absolute log2 fold change value at dpe12. Heatmap showing the top 20 DEGs at dpe12. For the heatmaps on the right, the top 20 gene names are indicated on the left of the graph.*

1031

Supplementary Figure 7. *Common differentially expressed genes (DEGs) across infections and time. DEGs for each infected samples over their uninfected controls. Venn diagrams of differentially expressed genes with differential expression values lower than 1.5 log2 fold change and/or with p-adjusted values lower than 0.05. **A)** DEGs for days 1, 12, and the day of median survival for each infected individual stock. **B)** DEGs on the day of median survival of all persistently infected stocks. **C)** DEGs for each persistently infected stock over time.*

1032

 SUPPLEMENTARY TABLES

Supplementary Table 1. Primers used in qPCR step of RT-qPCR for viral load quantification.

Primer name	Sequence
Rp49_Forward	CGGATCGATATGCTAAGCTGT
Rp49_Reverse	GCGCTTGTTCGATCCGTA
DAV_Forward	GTTGGATCAGGCTAGTGTAGG
DAV_Reverse	TGCAACCGGACTCCAAGTTC
DCV_Forward	AGGCTGTGTTTGCGCGAAG
DCV_Reverse	AATGGCAAGCGCACACAATTA
Bloomfield_Forward	CCGCCGGCTACCTACTTTAG
Bloomfield_Reverse	ACCAGCAAGCGTCCGAAATA
Nora_Forward	GGTTTGCAGTTGATCGCAG
Nora_Reverse	TCAGCTCATCGTTCACGAA
8/27_Forward	AGAGTTTGATCMTGGCTCAG
1492_Reverse	CGGTTACCTTGTTACGACTT
16S_qPCR_Forward	TCCTACGGGAGGCAGCAGT
16S_qPCR_Reverse	GGACTACCAGGGTATCTAATCCTGTT
Rp49_gDNA_Forward	GAAGTTCCTGGTGCACAACG
Rp49_gDNA_Reverse	CTTGCGCTTCTTGGAGGAGA

Supplementary Table 2. Primers used for RT-PCR to assess the presence of viruses and *Wolbachia*.

DAV_Forward	AGAGTGGCTGTGAGGCAGAT
DAV_Reverse	GCCATCTGACAACAGCTTGA
DCV_Forward	GTTGCCTTATCTGCTCTG
DCV_Reverse	CGCATAACCATGCTCTTCTG
Nora_Forward	ATGGCGCCAGTTAGTGCAGACCT
Nora_Reverse	CCTGTTGTTCCAGTTGGGTTCGA
Bloomfield_Forward_Seg4	CTATGGTTATCGATTGCATGGTCC
Bloomfield_Reverse_Seg4	GTAAACAAATCAAAACCATC
Bloomfield_Forward_Seg7	ATTTTTGGACTCAGATTGG
Bloomfield_Reverse_Seg7	GCCAAAATACTTGTTCCAG
Wsp_Forward	TGGTCCAATAAGTGATGAAGAAAC
Wsp_Reverse	AAAAATTAAACGCTACTCCA
WspB_Forward	TTTGCAAGTGAAACAGAAGG
WspB_Reverse	GCTTTGCTGGCAAAATGG

1039

SUPPLEMENTARY FILES

1040

Supplementary File 1. Genes associated with antiviral immune or locomotion function.

Supplementary File 2. Viable offspring production results.

Supplementary File 3. Differential gene expression analysis results. Table including all
the genes annotated for the differential dene expression analysis. All infected samples
are analyzed as in relationship to the uninfected control of the same day. All uninfected
samples are analyzed to the uninfected of day 1 post-eclosion.

1047

A Movement over time

B Sleep time

C Activity when awake

A — Antiviral genes

All genes / Top 20 DEGs

Immune pathway

1 dpe:
- CNT2
- kay
- Parp
- CG8046
- pirk
- PGRP-LB
- Tehao
- Spn42Dd
- spirit
- PGRP-SA
- psh
- GNBP-like3
- SPE
- grass
- GNBP3
- GNBP1
- modSP
- PGRP-SC1b
- PGRP-SC1a
- lncRNA:CR33942

12 dpe:
- cGlr2
- kay
- pirk
- PGRP-SD
- PGRP-LA
- PGRP-LC
- PGRP-LF
- PGRP-LB
- upd2
- Socs36E
- upd3
- PGRP-SA
- Hayan
- Toll-7
- GNBP-like3
- SPE
- PGRP-SC1b
- PGRP-SC2
- PGRP-SC1a
- lncRNA:CR33942

Columns: DAV, DCV, Bloomfield, Nora

Immune pathway legend:
- cGAS-STING
- cGAS-STING+IMD
- cGAS-STING+IMD+JAK-STAT
- IMD
- JAK-STAT
- JAK-STAT+IMD
- JAK-STAT+Toll
- Toll
- Toll+IMD

B — Locomotion genes

All genes / Top 20 DEGs

1 dpe:
- fz
- Neto
- Lsp2
- tim
- Pvf3
- ple
- Msp300
- rdgC
- Pu
- unc-5
- fz2
- Pvf2
- pyx
- nompC
- hh
- Rh5
- mt:ATPase6
- Fhos
- CG3857
- M6

12 dpe:
- CG3857
- Ptth
- M6
- nompC
- Rh5
- Pdf
- fz
- Neto
- santa-maria
- yuri
- bnl
- Trim9
- dpp
- NetA
- esg
- Pvr
- mt:scb
- Timp
- kay
- Mlc2

Columns: DAV, DCV, Bloomfield, Nora

A

Uninfected | DAV | DCV | Bloomfield | Nora

1 dpe: DAV ** | Bloomfield *** | Nora ***

6 dpe: DAV *** | DCV ** | Nora ***

12 dpe: DCV ** | Bloomfield *** | Nora ***

20 dpe: DAV * | Nora ***

B

Top | Middle | Bottom

Uninfected | DAV | DCV | Bloomfield | Nora

1 dpe: Bloomfield *** | Nora ***

6 dpe: DAV *** | DCV *** | Nora ***

12 dpe: DAV ** | DCV *** | Bloomfield ** | Nora ***

20 dpe: DCV ** | Bloomfield * | Nora ***

A Top 200 DEGs dpe1 | Top 20 DEGs dpe1

B Top 200 DEGs dpe12 | Top 20 DEGs dpe12