Title: pH-dependent beta-lactam resistance in *Klebsiella pneumoniae* is mediated by paralogous

class B PBPs and the class A PBP, PBP1b.

Authors

Sarah Beagle[1] and Petra Anne Levin[1*]

Affiliations

[1]Washington University in St. Louis, Department of Biology, St. Louis, Missouri, USA

*Correspondence: plevin@wustl.edu

Abstract

Klebsiella pneumoniae is a leading cause of global deaths due to antibiotic resistance. Of

particular concern is the rapid expansion of resistance to beta-lactam antibiotics within *K.*

pneumoniae lineages. The environmental factors that influence pathogen physiology and,

subsequently, antibiotic resistance remain poorly understood. Here we demonstrate that

physiologically-relevant reductions in pH increased *K. pneumoniae* beta-lactam resistance as

much as 64-fold, with the most dramatic increase observed for beta-lactams that specifically

inhibit cell division. We identified two genes that contribute to acid-dependent beta-lactam

resistance, the class A PBP, PBP1b, and the paralogous class B PBP, PBP3$_{PARA}$. Loss of either

PBP1b or PBP3$_{PARA}$ increases *K. pneumoniae* susceptibility to beta-lactams at low pH. Altogether

these data emphasize the importance of functional redundancy among cell wall synthesis

enzymes which allows for specialization and ensures robust cell wall synthesis across a range of

environmental conditions.

Importance

Beta-lactams are the most prescribed class of antibiotics, but their effectiveness is threatened by

a global rise in antimicrobial resistance. How the environment within a host or infection site shapes

pathogen response to antibiotics is frequently overlooked in assessments of antibiotic

effectiveness. We demonstrate that growth at physiologically-relevant low pH substantially

increases *Klebsiella pneumoniae* resistance to clinically important beta-lactams. An important

finding of this study is that during growth in acidic pH *K. pneumoniae* has a different repertoire of

cell wall synthesis genes available than during growth at neutral pH due to the presence of acid-

inducible paralogous copies of essential cell wall synthesis enzymes, PBP2 and PBP3. An

additional functionally-redundant enzyme, PBP1b, also contributes to acid-dependent beta-

lactam resistance. Together, these findings expand our understanding of how bacteria maintain

cell wall synthesis across diverse physiochemical environments and highlight potential new

therapeutic targets.

Introduction

Infections caused by the ESKAPE pathogen *K. pneumoniae* are increasingly difficult to treat due to

dramatic rises in antibiotic resistance. *K. pneumoniae* is a dominant member of two different Beta-

lactam-resistant groups of bacteria, Extended-spectrum-Beta-lactamase (ESBL)-producing

Enterobacteriaceae and Carbapenem-resistant *Enterobacteriaceae* (CRE) which are labeled

serious and urgent public health threats by the CDC[1]. In 2019, *K. pneumoniae* was the third leading

cause of global deaths where the death was directly due to or associated with antibiotic

resistance[2]. While ESBLs and related enzymes are major contributors to beta-lactam resistance in

Gram-negative bacteria, data from our group and others identifies physicochemical parameters

consistent with a host environment are under-appreciated drivers of resistance. Changes in pH,

nutrient availability, and accumulation of the alarmone, ppGpp, increase antibiotic resistance

across a range of bacterial pathogens, highlighting the need to better understand how

environmental factors impact pathogen physiology[3–6].

Despite the prevalence of resistance markers such as ESBLs, beta-lactams remain the most

prescribed antibiotic class[7]. Beta-lactams interfere with the synthesis of the cell wall or

peptidoglycan (PG), an essential structure that provides shape and structural rigidity. PG is

comprised of glycan strands of repeating linked N-acetylglucosamine (NAG) and N-acetylmuramic

acid (NAM) sugars that are crosslinked together by short peptide chains [8]. Synthesis and

maintenance of PG requires many enzymatic steps that occur across multiple cellular locations;

PG precursors are built in the cytoplasm, transported through the inner membrane, and fully

assembled into PG within the periplasmic space. Amongst the *Enterobacteriaceae,* periplasmic PG

assembly is highly enzymatically redundant, with an average of four enzymes capable of carrying

out a single periplasmic PG synthesis reaction[9] . This apparent functional redundancy is

particularly striking when compared to the earlier cytoplasmic PG synthesis reactions, which occur

in a nearly 1:1 enzyme-to-reaction ratio[10]. While precursors are assembled in the relatively buffered

cytoplasm, the cell wall itself is built in the periplasm and thus exposed to the physicochemical

environment.

Penicillin-binding proteins (PBPs) carry out essential transpeptidation reactions in the periplasm.

Class A PBPs are bi-functional enzymes that both link the NAG-NAM sugar moieties together

(transglycosylation, TG) as well as crosslink the peptide stems of adjacent glycan strands

(transpeptidation, TP). In *Escherichia coli,* the primary class A PBPs are PBP1a and PBP1b, which

are encoded by *mrcA* and *mrcB* respectively. PBP1a and PBP1b are individually dispensable for

growth, but loss of both is lethal[11–13]. A third class A PBP, PBP1c (*pbpC*), has an unknown role in PG

synthesis and maintenance and is dispensable under standard laboratory conditions[14]. Class B

PBPs are monofunctional transpeptidases; these PBPs work in conjunction with cognate

glycosyltransferases. PBP2 and PBP3 are the canonical class B PBPs in Gram-negative bacteria,

and they function within the elongation and division machinery respectively.

Beta-lactams interfere with PG synthesis by irreversibly binding to the transpeptidase domain of

PBPs and inhibiting transpeptidation[7]. Our previous work indicates that growth in acidic pH (~pH 5)

substantially increases *E. coli* resistance to PBP2- and PBP3-targeting Beta-lactams. Both growth

in pH values < 5, as well as acid-dependent Beta-lactam resistance required the Class A PBP,

PBP1b, while full fitness during growth in alkaline pH required PBP1a[3]. To determine if this

response is conserved amongst other *Enterobacteriaceae*, we evaluated the impact of

environmental pH on *K. pneumoniae* resistance to beta-lactams.

Here we report that *K. pneumoniae* beta-lactam resistance increased as much as 64-fold during

growth in acidic conditions. Genetic analysis identified two cell wall synthesis genes, *mrcB* and

ftsI2, that are critical for pH-dependent changes in antibiotic resistance. *mrcB* encodes PBP1b, a

bifunctional class A PBP previously identified as a contributor to pH-dependent antibiotic

resistance in *E. coli*[3]. Induced at low pH, *ftsI2* encodes $PBP3_{PARA,}$ a paralogous copy of the

canonical, division-associated class B PBP, PBP3. $PBP3_{PARA}$ is required for both acid-dependent

beta-lactam resistance and robust growth and division at low pH. Altogether our findings highlight

that functional redundancy in bacterial PBPs confers fitness advantages and contributes to

environmentally-driven antibiotic resistance.

Results

***Klebsiella pneumoniae* exhibits acid-dependent resistance to Beta-lactam antibiotics.**

To determine the impact of pH on *K. pneumoniae* beta-lactam susceptibility, we assessed the

response of a urinary tract infection (UTI) isolate, *K. pneumoniae* TOP52[15] to a panel of beta-

lactams during growth across a range of physiologically relevant pH conditions. The drug panel

included beta-lactams that preferentially target PBPs in the divisome (i.e. Cephalexin, Aztreonam,

and Piperacillin) or the elongasome (Meropenem, Doripenem), as well as non-specific inhibitors

(Ampicillin, Carbenicillin) that are reported to have similar affinities for all PBPs[16,17]. Relative

susceptibility or resistance to a given Beta-lactam was determined by comparing the minimal

inhibitory concentration of drug at neutral pH to that at other pH values.

K. pneumoniae exhibited a 2- to 64-fold increase in resistance to the entire spectrum of tested

beta-lactams when grown in LB at pH values < 5 relative to resistance at pH 7. (Fig 1A, Supp Table

4). We observed strong acid-dependent increases in resistance for drugs that preferentially target

PBP3, with Aztreonam (AZT) and Cephalexin (CEX) MICs increasing as much as 20- and 64-fold

respectively at pH 4.8. Resistance to PBP2-targeting Beta-lactams increased modestly at low pH,

with MICs to Doripenem (DOR) and Meropenem (MER) increasing 2- to 4-fold. In contrast to *E. coli*[3],

we also observed modest 2- to 4-fold increases in resistance to the non-specific Beta-lactams,

Ampicillin (AMP) and Carbenicillin (CAR) at low pH. At these pH values, increases in MICs during

growth at low pH are not due to significant antibiotic degradation[3].

Low pH not only enabled growth at concentrations of drug that are lethal at neutral pH but also

preserved cell morphology (Fig 1 C & D). Consistent with increased intrinsic resistance at low pH*,*

the characteristic rod-shaped morphology of *K. pneumoniae* was preserved during growth under

acidic conditions in concentrations of beta-lactams that cause either cell rounding (Meropenem)

or filamentation (Cephalexin) at neutral pH. The observed acid-dependent increased resistance to

beta-lactams is not strain specific behavior as *K. pneumoniae* strains NTUH-K2044 and ATCC

43816 also exhibit a 4-fold increase to Meropenem (MER) and an ~32-fold increase to Cephalexin

(CEX) when grown at pH 4.8 versus pH 6.8 (Fig. 1B).

To determine if this behavior also occurs under conditions that better reflect a native host

environment, we assayed how *K. pneumoniae* responded to beta-lactams during growth in an

artificial urine growth medium at low and neutral pH[18]. Similar to growth in LB, we observed 32- to

>64-fold increases in resistance to Ampicillin and Cephalexin in acidified artificial urine with and

without amino acid supplementation compared to neutral pH (Sfig 1, Supp table 4 & 5). While the

MIC for a given beta-lactam was generally higher during growth at low pH independent of media

composition, we observed that for certain antibiotics the media composition coupled with

acidification of the media significantly altered the susceptibility. The MIC for Ampicillin at low pH in

LB was significantly lower (mean MIC = 104 ug/mL ± 37) than during growth in AU (mean MIC =

2500 ug/mL ±0.0), indicating an interplay between nutrient composition and pH. We were unable to

determine the exact foldchange in MIC for Cephalexin, as *K. pneumoniae* grew in all tested

concentrations of Cephalexin in both AU and AU+AA (Sfig 1, table 4 & 5). Together our data suggest

that acid-dependent beta-lactam resistance occurs across a range of nutrient conditions and is

conserved amongst *K. pneumoniae* lineages.

Acid-dependent Beta-lactam resistance is independent of endogenous beta-lactamase

expression.

K. pneumoniae TOP52 encodes an endogenous class-A beta-lactamase, *bla$_{SHV-1}$*, which confers

increased resistance to the Penicillin subclass of Beta-lactams[7,15,19]. To determine if expression of

bla$_{SHV-1}$ is influencing the resistance profiles at low pH, we generated a deletion strain (Δ*bla$_{SHV-1}$*)

and repeated the MIC assays under both acidic and neutral pH conditions. We observed increased

sensitivity to the Penicillin sub-classes of Beta-lactams tested (AMP, CAR, PIP) in both pH

environments (Sup Table 5). Consistent with the specificity of *bla$_{SHV-1}$*, loss of *bla$_{SHV-1}$* did not alter

the MICs to non-Penicillin beta-lactams at either pH (Sup Table 5). However, the Δ*bla$_{SHV-1}$* mutant

still displayed increased resistance to all tested Beta-lactams at low pH relative to neutral pH (Fig 2

A-D, Supp table 5), demonstrating that acid-dependent resistance is independent of endogenous

beta-lactamase expression.

pH modulates expression of Class B PBP paralogs in *K. pneumoniae*.

To ascertain the impact of pH on expression of genes encoding PBPs or other cell wall synthesis

genes, we performed RNAseq in *K. pneumoniae* comparing cells cultured in pH 4.8 LB versus pH

6.8 LB (Supp table 7). Consistent with previous studies, pH does not impact expression of class A

PBPs, *mrcA* (PBP1a) or *mrcB* (PBP1b), in either *E. coli* or *K. pneumoniae*[20,21] (Table 1). Class A PBP

activity is regulated by cognate outer membrane activators, LpoA and LpoB[22–24]. We observed that

the PBP1b activator, encoded by *lpoB*, was down-regulated ~2-fold at low pH.PBP1c, a cryptic

class A PBP with no demonstrated role in cell wall synthesis under standard laboratory conditions,

exhibited an ~2-fold decrease in expression during growth in acidic pH (Table 1, Supp table 7). We

observed no significant changes expression of the class B PBPs between pH conditions, but there

was an order of magnitude increase in expression at low pH of genes we designated *mrdA2* and

ftsI2, which encode paralogous copies of class B PBPs, referred to here as PBP2$_{PARA}$ and PBP3$_{PARA}$

(Fig 3 A&B). In addition to the class B paralogs, several other cell wall synthesis genes displayed

pH-responsive expression profiles. Most notably, two carboxypeptidases, *dacA* (PBP5) and *dacD*

(PBP6b), exhibited modest increases of 2- to 3-fold in expression during growth at low pH. Both

dacA and *dacD* are implicated in maintaining proper cell morphology in *E. coli*, with *dacD* being

required for proper morphology specifically at low pH[25,26].

Loss of PBP3$_{PARA}$ results in filamentation in *K. pneumoniae* at low pH.

Duplications of the class B PBPs, PBP2 and PBP3, that have expanded the repertoire of cell wall

synthesis enzymes have been described in several pathogens[27–29]. Class B PBP paralogs in

Salmonella enterica serovar Typhimurium (*S.* Typhimurium) referred to as PBP2$_{SAL}$ and PBP3$_{SAL}$ are

thought to be important for maximal fitness during growth at low pH. Expression of both PBP2$_{SAL}$

and PBP3$_{SAL}$ is maximally induced at low pH (<5.8), and loss of PBP2$_{SAL}$ has been reported to impair

rod-shape morphology under acidic environments[28,29].

As with PBP2$_{SAL}$ and PBP3$_{SAL}$, PBP2$_{PARA}$ and PBP3$_{PARA}$ share significant sequence identity with their

canonical counterparts (59% for PBP2$_{PARA}$ and 61% for PBP3$_{PARA}$) and retain the sequence motifs

associated with catalysis, but their contributions to cell wall maintenance in *Klebsiella* are

unknown (Fig 3 C&D, Sfig 2). We generated individual gene deletion strains and examined the

mutants using cell size analysis and time-lapse microscopy to evaluate the role of the class B PBP

paralogs in cell size and division. We cultured cells at both acidic and neutral pH and sampled

cultures at early exponential phase (OD$_{600}$ = 0.1-0.2) for cell size analysis. While loss of PBP2$_{PARA}$ did

not impact cell size or morphology at either pH, we found that during growth at pH 4.8 the PBP3$_{PARA}$

deletion mutant exhibited heterogeneous cell lengths, with many cells displaying filamentous

behavior indicative of cell division defects (Fig4 A&B, Sfig 3). Using a definition of filamentation as a

cell length of >15 μM which represents ~3X the average length of WT cells at pH 4.8 (3.9 ± 0.2 μM,

Fig 4B), we determined that 21-58% of PBP3$_{PARA}$ deficient cells within a given population were

filamentous, with a smaller proportion of cells (3-17%) reaching lengths of >40 μM. The division

defects associated with loss of PBP3$_{PARA}$ were specific to growth in acidic conditions, as growth at

neutral pH resulted in an average cell length for the PBP3$_{PARA}$ deficient cells consistent with WT cell

lengths (Fig 3C).

Time-lapse microscopy of acid-grown Δ*ftsI2* cells spotted onto acidic LB agarose pads revealed a

population of cells with heterogeneous lengths that displayed clear division defects as cells

continued to elongate without successful division resulting in long filamentous cells. Whereas

spotting acid-grown Δ*ftsI2* cells onto LB agarose pads at neutral pH restored division to the

filamentous cells (Sfig 4). Complementation of *ftsI2* in trans from a medium-copy number plasmid

enabled division and restored cell lengths consistent with WT at low pH (Fig 4 A&B). Together these

data suggest that the activity of the canonical PBP3 is impaired under acidic conditions and that

PBP3$_{PARA}$ is the predominant transpeptidase supporting cell division in *K. pneumoniae* at low pH.

Loss of PBP1b or PBP3$_{PARA}$ differentially impacts Beta-lactam resistance at low pH in *Klebsiella*

The increased expression of genes encoding PBP2$_{PARA}$ and PBP3$_{PARA}$ and other cell wall

maintenance proteins at low pH raised the possibility that these proteins might contribute to acid-

dependent Beta-lactam resistance. To address this hypothesis, we generated strains individually

defective in a subset of *K. pneumoniae* genes implicated in cell wall or shape maintenance and

assayed their response to beta-lactam antibiotics during growth at acidic and neutral pH

conditions. In addition to genes that displayed pH-dependent expression, we also evaluated the

role of class A PBPs in *K. pneumoniae* response to beta-lactams across pH conditions, as these are

known to influence antibiotic response in *E. coli*. While most individual mutations did not

significantly impact acid-dependent-Beta-lactam resistance, loss of PBP1b or PBP3$_{PARA}$ altered

sensitivity to Beta-lactams at low pH (Fig 2A-D, Supp table 6).

Loss of PBP1b (Δ*mrcB*) most dramatically affected acid-dependent resistance to Cephalexin. A

Δ*mrcB* mutant displayed only a 4-fold increase in cephalexin MIC at low pH versus at neutral pH,

while the WT has a 64- to 128-fold increase in resistance under the same conditions. The Δ*mrcB*

mutant also no longer exhibited acid-dependent resistance to the generalist (AMP) and PBP2-

targeting- (DOR, MER) beta-lactams, (Fig 2 A-D, Sfig 6, Supp table 6).The increased beta-lactam

sensitivity of a PBP1b deletion mutant at low pH is likely due to loss of the enzyme activity under

this condition. Deletion of PBP1b's cognate outer membrane activator lpoB (Δ*lpoB*) results in

similar sensitization levels at low pH as a PBP1b mutant (Sfig 6, Supp Table 6). Defects in PBP1b

also led to a nearly 10-fold increase in susceptibility to Cefsulodin in both acidic and neutral pH

compared to WT (Sup Table 6). This pH-independent increase in sensitivity of the ΔPBP1b mutant

is consistent with data suggesting that PBP1a is the primary Class A target of Cefsulodin[30,31].

Loss of PBP3$_{PARA}$ differentially impacted resistance to beta-lactams. The PBP3$_{PARA}$ null mutant no

longer displayed acid-dependent resistance to Ampicillin and Cefsulodin at low pH (Fig 2A & B). In

fact, a PBP3$_{PARA}$ deletion mutant was more sensitive to Ampicillin at low pH than at neutral pH (Fig

2A). When challenged with PBP3-targeting Beta-lactams (CEX, AZT, PIP), the PBP3$_{PARA}$ deletion

mutant still retained mild acid-dependent resistance but was 2- to 8-fold more sensitive at low pH

than WT (Fig 4 D, 5 B). Surprisingly, the PBP3$_{PARA}$ null mutant exhibited increased resistance when

challenged with PBP2-targeting beta-lactams at low pH. The Meropenem and Doripenem MICs

obtained for the PBP3$_{PARA}$ null mutant were 12-16-fold higher at acidic pH versus neutral pH. For

comparison, WT cells only exhibited a 2-4-fold increase in MIC to these two drugs at acidic pH (Fig

4C, 5A). Despite the ~10-fold increases in expression under acidic conditions, loss of *mrdA2,*

encoding PBP2, had no significant impact on Beta-lactam susceptibility across pH values.

Increased acid-dependent resistance to PBP2-targeting compounds in the PBP3$_{PARA}$ null

mutant requires PBP2$_{PARA}$.

We reasoned that the unexpected increase to PBP2-inhibitors at low pH observed in the Δ*ftsI2*

mutant could be influenced by the presence of PBP2$_{PARA}$. To address this possibility, we generated

a double deletion strain lacking both class B paralogs (Δ*ftsI2* Δ*mrdA2*) and assayed its response to

beta-lactams with high specificity for PBP2 or PBP3 at both neutral and acidic pH. Deletion of both

class B PBP paralogs eliminated the dramatic increase in resistance at low pH (4.8), restoring WT-

level responses to both Doripenem and Meropenem. When challenged with PBP3-targeting Beta-

lactams, the double mutant class B paralog mutant phenocopied the Δ*ftsI2* mutant, displaying

similar reductions in resistance at low pH (Fig 5). Together these data suggest that both PBP2$_{PARA}$

and PBP3$_{PARA}$ are involved in mediating resistance to PBP2-inhibitors, whereas resistance to PBP3-

targeting Beta-lactams is primarily facilitated by PBP3$_{PARA}$.

Discussion

As observed in *E. coli,* beta-lactam resistance in *K. pneumoniae* increases substantially during

growth in acidified medium (Fig 1A). Importantly, we observed pH-dependent increases in beta-

lactam resistance in LB as well as in a more physiologically relevant medium, artificial urine (Fig 1,

Sfig 1). We find that the class A PBP, PBP1b, and the acid-responsive paralogous copy of the class

B PBP, PBP3$_{PARA}$, are major genetic contributors to acid-mediated beta-lactam resistance in *K.*

pneumoniae. Disruption of *mrcB*, the gene encoding PBP1b, eliminates acid-mediated resistance

to the class B PBP inhibitors, Meropenem, Doripenem, and Cephalexin. Loss of PBP3$_{PARA}$

differentially impacts beta-lactam resistance, with mutants showing sensitization towards

Ampicillin, Cefsulodin, and all PBP3-targeting beta-lactams but increased resistance towards

PBP2-targeting antibiotics.

While both *E. coli* and *K. pneumoniae* demonstrate acid-dependent beta-lactam resistance,

factors influencing this phenomenon differ between the two organisms. Acid-mediated beta-

lactam resistance in *E. coli* appears to be solely due to intrinsic properties of the PBP enzymes,

particularly PBP1b[3], while in *K. pneumoniae* the contribution of PBP1b is supplemented by pH-

dependent induction of genes encoding paralogous class B PBPs. Although we only observed

strong changes to antibiotic susceptibility in a PBP3$_{PARA}$ null mutant, loss of PBP2$_{PARA}$ impacted the

resistance to PBP2-targeting antibiotics in a PBP3$_{para}$ null background, as well as the morphological

response at low pH (Fig 5 C). This result suggests that PBP2$_{PARA}$ plays a role in maintaining PG and

cell shape at low pH. Together, our findings support a modular model of cell wall synthesis within

the *Enterobacteriaceae* family where apparent function redundancy amongst cell wall synthesis

enzymes is protective by enabling PG synthesis across a range of pH conditions.

The Class A PBP, PBP1b, is important for acid-dependent beta-lactam resistance in *K.*

pneumoniae

PBP1b is required for resistance to PBP2- and PBP3-targeting beta-lactams and for growth at low

pH (<5) in *E. coli*[3]. In contrast, deletion of PBP1b in *K. pneumoniae* did not significantly impair

growth in acidified culture medium (pH 4.8), suggesting that there are key differences in how PBP1b

influences growth in acidic environments between *E. coli* and *K. pneumoniae* (Sfig 5). While

expression of *mrcB,* encoding PBP1b, was not influenced by pH, loss of PBP1b impaired *K.*

pneumoniae's resistance to Ampicillin, Doripenem, Meropenem, and Cephalexin at low pH (Fig 2,

Supp Table 5). Consistent with MIC data, a PBP1b-deficient strain exhibits faster killing kinetics

than a WT strain in the presence of Cephalexin at low pH (Fig 2 E &F). While filamentation is a

characteristic morphology associated with division-targeting beta-lactams[32,33], at low pH a PBP1b-

deficient *K. pneumoniae* strain is sensitive to Cephalexin at concentrations that do not elicit

filamentation (Fig 4E). The time frame for lysis of the PBP1b mutant is rapid with more than 50% of

the population exhibiting lysis or loss of membrane integrity as denoted by propidium iodide

staining within 90 minutes of exposure to Cephalexin at low pH (Fig 2F).

Notably, at neutral pH loss of PBP1b or its cognate outer membrane activator, LpoB, mutant did not

significantly alter MICs across beta-lactam classes (Sup Table 5), suggesting that PBP1B specifically

mediates beta-lactam resistance at low pH in *K. pneumoniae*. This finding differs from what has

been observed in *E. coli* where loss of either PBP1b or LpoB increases sensitivity to multiple classes

of beta-lactams at neutral pH[34,35]. This result is also consistent with prior data suggesting that PBP1a

does not function optimally under acidic conditions and therefore cannot compensate for loss of

PBP1b activity at low pH[3]. Our data demonstrating that a PBP1b mutant is more sensitive to PBP2-

and PBP3-targeting beta-lactams is consistent with prior studies reporting interactions between

PBP1b and key division and elongasome proteins[36,37,29]. Based on these data, we propose that PBP1b

supports cell wall integrity at low pH through interactions supporting other cell division proteins.

Class B PBP paralogs differentially modulate pH-dependent changes in beta-lactam

resistance

Although, low pH induced expression of genes encoding both *K. pneumoniae's* PBP2 and PBP3

paralogs (Table 1), only loss of *ftsI2,* encoding PBP3$_{PARA}$, sensitized cells to beta-lactams during

growth in acidified medium. These data are consistent with observations in *S.* Typhimurium that

loss of its PBP3 paralog, PBP3$_{SAL}$sensitizes cells to beta-lactams targeting division[38].Defects in

mrdA2, encoding PBP2$_{PARA}$, did not significantly impact beta-lactam resistance at either acidic or

neutral pH. In *K. pneumoniae* loss of PBP3$_{PARA}$ at low pH sensitized cells to not only PBP3 inhibitors,

but also to a Cephalosporin targeting class A PBPs and the penicillin, Ampicillin (Figure 4 A&B).

While loss of PBP3$_{PARA}$ generally sensitized *K. pneumoniae* to beta-lactams under acidic conditions,

it increased resistance to compounds targeting PBP2 at pH 4.8 up to 12- to 16-fold (Figure 5A). This

increase in resistance in PBP3$_{PARA}$ deficient cells was eliminated in a double mutant lacking both

PBP3$_{PARA}$ and PBP2$_{PARA}$(Δ*ftsI2* Δ*mrdA2*). Under acidic conditions the double class B paralog mutant

exhibited WT-level MICs when challenged with Meropenem or Doripenem (Figure 5).

It is unclear why the loss of PBP3$_{PARA}$ renders cells resistant to PBP2-targeting compounds.

Morphological data of WT *K. pneumoniae* challenged with Meropenem at low pH reveals hybrid

morphologies, with cells displaying signatures of both divisome and elongasome inhibition. Cells

first undergo lengthening and then display large, rounded septal bulges (Fig 5C). These results

could reflect altered target binding by Meropenem at low pH, where PBP3/PBP3$_{PARA}$ are also

inhibited along with PBP2/PBP2$_{PARA}$. Alternatively, physical interactions between class B PBPs

within the elongasome and divisome have been reported in *E. coli*[39], and data suggests that the

protein components of the PG synthesis complexes in *Salmonella* vary depending on whether the

canonical or paralogous class B PBP is present in the complex[29,40]. Alterations to interactions

between class B PBP paralogs and other cell wall synthesis proteins due to beta-lactam binding or

in the class B PBP paralog mutant backgrounds could influence morphology and antibiotic

resistance. Additional work is needed to fully understand the complex role that the class B PBP

paralogs play in resistance to PBP2-targeting beta-lactams.

PBP3$_{PARA}$ is critical for *K. pneumoniae* division at low pH

Our data suggest that PBP3$_{PARA}$ is the major transpeptidase supporting cross wall synthesis at low

pH. Loss of PBP3$_{PARA}$ did not impact cell length at neutral pH but impaired division at low pH,

resulting in filamentous cells. However, these acid-cultured PBP3$_{PARA}$-deficient filamentous cells

were able to resume division and return to WT-lengths if they were transitioned to neutral pH

(Sfig3), suggesting that the canonical PBP3 is unable to fully compensate for the loss of PBP3$_{PARA}$ at

low pH. We attempted to disrupt the canonical PBP3 under acidic culture conditions where

PBP3$_{PARA}$ is likely the dominate division-specific transpeptidase; however, despite multiple

attempts, we were unsuccessful. Moreover, a small percentage of Δ*ftsI2* cells maintained WT-

lengths at low pH, suggesting that the canonical PBP3 retains limited, but critical activity at low pH

(Sfig 3). These data differ from *Salmonella* where the canonical PBP3 and PBP3 paralog PBP3$_{SAL}$ are

individually dispensable at low pH. The canonical PBP3 is capable of supporting growth at both low

and neutral pH, but PBP3$_{SAL}$ is unable to support division at neutral pH[28]. Our results support a

model in which *K. pneumoniae* PBP3 functions optimally at neutral pH while PBP3$_{PARA}$ functions

predominately at acidic pH, with both working together to ensure that division remains robust

across pH conditions.

Expansion and pH specialization of *K. pneumoniae* cell wall synthesis enzymes provide novel

therapeutic targets

Identification of paralogous class B PBPs has thus far only been described in pathogens [27–29,41], but

the role of these paralogous cell wall synthesis proteins in pathogen physiology remains largely

uncharacterized. In *Salmonella,* the acid-responsiveness of class B PBP paralogs raised

speculation that they could be critical to support bacterial survival during intracellular growth

within acidified host immune cells. However, loss of the class B PBP paralogs did not impair

replication within macrophages or attenuate virulence in a mouse infection model[28].

In *Klebsiella pneumoniae,* our data suggests that class B PBP paralogs are part of a general strategy of functional redundancy to preserve critical enzymatic activity across a wide range of environmental conditions. Colonization of a host exposes a bacterial pathogen to a wide range of pH conditions[42–44] . We demonstrate that the acidification of growth medium to physiologically-relevant levels is sufficient to increase resistance to beta-lactams. The class B PBP paralogs modulate that resistance in part, suggesting that these pathogen-specific proteins may provide novel therapeutic targets. We propose that future screens for new antimicrobial targets factor environmental pH as a variable to identify novel compounds that can bind to acid-responsive targets such as the Class B PBP paralogs. Altogether, our results highlight the need to evaluate pathogen physiology in diverse environments beyond the standard laboratory conditions.

Acknowledgements

We thank members of the Levin lab for their feedback and discussion regarding this study, with a special thanks to Kyra Raines and Dr. Sarah Anderson for critical reading of this manuscript. We are also grateful to David Rosen and his group for the gift of TOP52 and welcome advice on genetic manipulation of *K. pneumoniae*. This work has been funded by the NIH through an R35-GM127331 award to P.A.L.

Contributions

S.B.: Conceptualization, Investigation, Formal analysis, Methodology, Visualization, Writing-original draft, Writing – review and editing.

P.A.L.: Conceptualization, Funding acquisition, Resources, Supervision, Writing – review and editing.

Data availability

All raw and processed data required to evaluate the conclusions of this study are included in the text or supplemental data. The RNA sequencing data for gene expression studies at acidic and neutral pH have been deposited in the NCBI GEO repository: GSE291667.

Methods

Bacterial strains, growth conditions, and antibiotics.

Bacterial strains and plasmids used in this study are listed in Supplementary table 1 and

Supplementary table 2. Strains were routinely cultured in Lysogeny Broth (LB – 1% Tryptone, 0.5%

Yeast extract, and 1% NaCl) in large glass tubes at 37 °C with 200 RPM aeration. The pH of LB was

adjusted with 1:10 MMT buffer (1:2:2 molar ratio of D/L-malic acid, MES, and Tris base). Artificial

urine (AU) was prepared as described[18], and cells were cultured in either base AU or AU

supplemented with all twenty amino acids at a final concentration consistent with what is used in

EZ rich defined medium (https:www.genome.wisc.edu/resources/protocols/ezmedium.htm). The

pH of AU or AU+AA was adjusted with MMT buffer immediately prior to use. For ease and clarity,

media and amino acid composition is listed in Supp table 8.

Where indicated cultures were supplemented with Kanamycin (50 µg/mL), Spectinomycin (100

µg/mL), Chloramphenicol (50 µg/mL), or IPTG (0.1 mM) . The pRDC3 plasmid (p15A ori, SpecR,

Plac) was a gift from Fabrizio Arigoni. Plasmid pSDB8 was generated from pRDC3 using the IVA

cloning method to insert *ftsI2* under control of the Plac promoter[45]. Primers were acquired from

Integrated DNA Technologies (Coralville, IA) and sequences used are listed in Supp Table 3.

Chemically competent *E. coli* TOP10 (Invitrogen) were used to assemble vectors via IVA. Plasmid

sequence was confirmed via whole plasmid sequencing at Plasmidsaurus prior to transformation

into *K. pneumoniae* strains.

For growth studies, cultures were started in LB supplemented with MMT buffer to the indicated pH

and grown in large glass tubes at 37 °C with 200 RPM aeration. Cultures were harvested once they

reached mid-exponential phase (OD_{600} =0.2-0.6), concentrated to an OD_{600} =1.0, and serially

diluted 10-fold. 1.5 µL of the 10^{-5} dilution was used to inoculate wells of sterile 96-well plates

containing 150 µL of LB + MMT at indicated pH. The plate was covered with a Breathe-Easy strip

(Diversified Biosciences) and growth was monitored in a BioTek Synergy H1 plate reader with the

following settings: 37 °C, continuous double orbital shaking at 282 cpm, with OD_{600} readings taken

every 10 minutes.

For *ftsI2* cell size complementation studies, un-induced overnight cultures of strains carrying either

pRDC3 (Empty vector, EV) or pSDB8 were back-diluted 1:500 into 3 mL of fresh LB buffered to

either pH 4.8 or pH 6.8 containing Spectinomycin (100 µg/mL). Cultures were grown at 37 °C with

200 RPM aeration for 1.5 hrs. Expression of *ftsI2* was induced with the addition of 0.1 mM IPTG and

cultures were allowed to continue growing until early exponential phase (OD_{600} = 0.1-0.3) was

reached (~40-90 additional minutes). Cultures were back-diluted into fresh buffered media

containing Spectinomycin (100 µg/mL) and 0.1 mM IPTG to an OD_{600} ~= 0.005. Growth was

monitored every 30 minutes, and cells were harvested when cultures reached early exponential

phase (OD_{600} = 0.1-0.3). Cells were spotted onto media pads (LB + MMT (pH 4.8 or 6.8) +

Spectinomycin (100 µg/mL) + 0.1 mM IPTG + 1% agarose) and imaged live with phase microscopy.

The following antibiotics were used for antibiotic susceptibility screenings. Ampicillin sodium salt,

Aztreonam, Cephalexin hydrate, Cefsulodin sodium salt hydrate, Meropenem Trihydrate, and

Piperacillin sodium salt were obtained from Sigma Adrich (St. Louis, MO). Carbenicillin sodium salt

was from EMD Millipore (now Millipore Sigma, Germany). Doripenem was obtained from Cayman

Chemicals (Ann Arbor, MI).

Genetic manipulation: Gene disruptions were carried out in the *K. pneumoniae* TOP52

background using a Lambda Red vector modified to carry a Spectinomycin resistance cassette as

described previously [46,47]. Briefly, DreamTaq Polymerase (Thermo Scientific) was used to PCR

amplify the Kanamycin resistance cassette from the template plasmid pKD4 with primers

containing 40-60 bp of homology to the chromosomal targets. Primers were designed to leave the

start and stop codons of the targeted gene intact. In cases where the two gene sequences

overlapped, primers were designed to avoid disruption to any downstream genes. Linear PCR

products were purified using a Purelink PCR purification kit (Invitrogen), and 2,000-3,000 ng of

purified linear fragments were electroporated into *K. pneumoniae* cells grown in the presence of

0.7% L-arabinose to induce expression of the Lambda Red recombinase system from the pKD46S

plasmid. Electroporated cells were recovered in SOC (Lloyd's recipe – 2% Tryptone, 0.5% yeast

extract, 10 mM NaCl, 2.5 mM KCl, 10 mM $MgSO_4$, 20 mM glucose) for 4-8hrs at 30°C. Deletion

mutants were selected for by plating transformation culture onto LB agar plates containing

Kanamycin (50 µg/mL) and incubated at 37°C. Mutants were confirmed by colony PCR and

sequencing as necessary. For marker-less deletions, cells were transformed with pCP20 at 30°C in

LB + Chloramphenicol (50 µg/mL) to express the FLP to excise the FRT-flanked Kanamycin

cassette. After excision, cells were cultured at 42°C to promote loss of pCP20. Kanamycin and

Chloramphenicol sensitivity was confirmed for strains prior to use.

RNA sequencing: *K. pneumoniae* TOP52 or *E. coli* MG1655 was cultured overnight in LB buffered

to either pH 4.8 or 6.8 with MMT buffer. Overnight cultures were back diluted 1:100 into fresh media

and cultured at 37°C with 200 RPM aeration until cultures reached early exponential phase (OD_{600}

=~0.2). ~ 9.6×10^8 cells were transferred to 15 mL conical Falcon tubes and pelleted by

centrifugation at 4°C (4,000 RPM for 10 mins). The supernatant was discarded, cell pellets were

resuspended in 350 uL of RNAwiz, and the RNA was extracted following the protocol for the

Ribopure-Bacteria RNA extraction kit [Invitrogen]. RNA was eluted in 50 uL of elution solution.

Purified RNA was subjected to a DNase I treatment to remove any contaminating chromosomal

DNA. RNA sequencing and differential gene expression analysis was performed by MiGS (now

SeqCenter). Quality control and adapter trimming of reads was performed with bcl2fastq. Read

mapping was performed with HISAT2, and read quantification was performed using Subread's

featureCounts. Read counts were normalized in R using edgeR's Trimmed Mean of M values (TMM)

algorithm, and subsequent values were converted to counts per million (CPM). Differential

expression analysis was performed using edgeR's Quasi-Linear F-Test (qlfTest) functionality

against treatment groups, and DEGs were considered as those with |logFC| > 1 and p < .05. Three

biological replicates were submitted per pH condition.

Antibiotic susceptibility testing. Cells were cultured from a single colony in LB at the indicated

pH to exponential phase (OD_{600} =~0.2-0.6) at 37 °C with 200 RPM aeration. For MICs in AU or

AU+AA, overnight cultures were back-diluted 1:100 into fresh media and cultured at 37 °C with 200

RPM aeration at the indicated pH to exponential phase (OD_{600} =~0.2-0.6). Harvested cells were

concentrated to an OD_{600} =1 and serially diluted 10-fold. 1.5 µL of the 10^{-2} dilution was used to seed

wells containing 150 µL of LB broth + MMT (matching pH) in sterile 96-well microtiter plates

containing a range of two-fold dilutions of beta-lactam antibiotics. Plates were sealed with a

Breathe-Easy strip (Diversified Biotech) and incubated for 20 hours at 37°C with 200 RPM aeration.

The minimal inhibitory concentration (MIC) was determined by visual inspection and reflects the

lowest drug concentration that completely inhibited bacterial growth.

Terminal morphology Assessment. At the end of 20 hours for antibiotic susceptibility tests, cells

were taken from wells following growth at either neutral or acidic pH with the indicated antibiotic

concentrations. 5 µL of cells were spotted onto LB + 1% agarose and imaged by phase contrast

microscopy.

Microscopy, cell size analysis, and time-lapse Imaging. All phase contrast and fluorescence

images were acquired on either a Nikon Ti-E inverted microscope or a Ti2-E inverted microscope

(Nikon Instruments). The Ti-E is equipped with a 100X Plan Apo oil objective (N.A. = 1.45 Ph3

objective, X-Cite 120 LED light source (Lumen Dynamics) and an OrcaERG CCD camera

(Hammamatsu Photonics, Bridgewater, N.J.) Filter cubes were purchased from Chroma

Technology Corporation. The Ti2-E inverted microscope is equipped with a 100X Plan Apo oil

objective (N.A. = 1.45, Ph3 objective, Lumencor Sola LED light engine (Lumencore), and an orca-

Fusion sCMOS camera (Hammamatsu Photonics). Filter cubes were purchased from Chroma

Technology Corporation. Both microscopes have an enclosed temperature-controlled enclosure to

pre-heat the objective to 37 °C (OXO enclosure) prior to experiments. Nikon Elements software was

used for image capture, and analysis was performed using the MicrobeJ plug-in in ImageJ.

For cell size analysis, cells were grown from a single colony in LB with MMT buffer at indicated pH

at 37 °C and 200 RPM aeration to exponential phase (OD_{600} = 0.2-0.4) and back-diluted to an OD_{600}

= 0.005 into 7 mL of pre-warmed LB + MTT (same pH). Growth was monitored via optical density,

and when cultures reached early exponential phase (OD_{600} =0.1- 0.2) cells were harvested. 5 μL of

cells were spotted onto LB + MTT (same pH)+ 1% agarose pads and imaged by phase contrast

microscopy. Cell morphology and size dimensions were determined using the MicrobeJ[48] plug-in

for ImageJ.

For time-lapse imaging, cultures were grown from a single colony in LB with MMT buffer at

indicated pH at 37 °C and 200 RPM aeration to exponential phase (OD_{600} = 0.1-0.4). 5 μL of cells

were spotted onto LB + MTT (same pH)+ 1% agarose pads that contain antibiotics at indicated

concentrations and 1.5 μM propidium iodide (PI) for monitoring loss of membrane integrity. Phase

images were obtained every 2 minutes, and fluorescent images were collected every 10 minutes

(Chroma AT560/40X/ Chroma AT600 DC/ Chroma at635/60nm with 100 mS exposure/10%

intensity). For each experiment a total of 2-5 different XY coordinates were monitored throughout

the duration of the experiment. Quantification of PI+ cells was done using the MicrobeJ[48] plug-in for

ImageJ. Briefly the number of cells in the phase image and the number of corresponding

fluorescent cells were quantified. Data was reported as either percentage of PI+ cells (number of PI+

cells/total number of cells *100) or percent alive (number of cells-PI+ cells/total number of

cells*100).

Bibliography

1.   Centers for Disease Control and Prevention (U.S.). *Antibiotic Resistance Threats in the United*

States, 2019. https://stacks.cdc.gov/view/cdc/82532 (2019) doi:10.15620/cdc:82532.

2. Murray, C. J. *et al.* Global burden of bacterial antimicrobial resistance in 2019: a systematic

analysis. *The Lancet* **399**, 629–655 (2022).

3. Mueller, E. A., Egan, A. J. F., Breukink, E., Vollmer, W. & Levin, P. A. Plasticity of Escherichia coli

cell wall metabolism promotes fitness and antibiotic resistance across environmental

conditions. *eLife* **8**, (2019).

4. Vinella, D., Albrecht, C., Cashel, M. & D'Ari, R. Iron limitation induces SpoT-dependent

accumulation of ppGpp in Escherichia coli. *Mol Microbiol* **56**, 958–970 (2005).

5. Fanous, J. *et al.* Limited impact of Salmonella stress and persisters on antibiotic clearance.

Nature **639**, 181–189 (2025).

6. Honsa, E. S. *et al.* RelA Mutant Enterococcus faecium with Multiantibiotic Tolerance Arising in

an Immunocompromised Host. *mBio* **8**, (2017).

7. Bush, K. & Bradford, P. A. β-Lactams and β-Lactamase Inhibitors: An Overview. *Cold Spring*

Harbor Perspectives in Medicine **6**, a025247 (2016).

8. Egan, A. J. F., Errington, J. & Vollmer, W. Regulation of peptidoglycan synthesis and

remodelling. *Nature Reviews Microbiology 2020 18:8* **18**, 446–460 (2020).

9. Mueller, E. A. & Levin, P. A. Bacterial cell wall quality control during environmental stress. *mBio*

11, 1–15 (2020).

10. Pazos, M., Peters, K. & Vollmer, W. Robust peptidoglycan growth by dynamic and variable

multi-protein complexes. *Current Opinion in Microbiology* **36**, 55–61 (2017).

11. Kato, J., Suzuki, H. & Hirota, Y. Dispensability of either penicillin-binding protein -1a or -1b

involved in the essential process for cell elongation in Escherichia coli. *Molec. Gen. Genet.*

200, 272–277 (1985).

12. Suzuki, H., Nishimura, Y. & Hirota, Y. On the process of cellular division in Escherichia coli: a

series of mutants of E. coli altered in the penicillin-binding proteins. *Proceedings of the*

National Academy of Sciences **75**, 664–668 (1978).

13. Yousif, S. Y., Broome-Smith, J. K. & Spratt, B. G. Lysis of Escherichia coli by beta-lactam

antibiotics: deletion analysis of the role of penicillin-binding proteins 1A and 1B. *J Gen*

Microbiol **131**, 2839–2845 (1985).

14. Schiffer, G. & Höltje, J.-V. Cloning and Characterization of PBP 1C, a Third Member of the

Multimodular Class A Penicillin-binding Proteins of *Escherichia coli* *. *Journal of Biological*

Chemistry **274**, 32031–32039 (1999).

15. Johnson, J. G., Spurbeck, R. R., Sandhu, S. K. & Matson, J. S. Genome sequence of Klebsiella

pneumoniae urinary tract isolate Top52. *Genome Announcements* **2**, 668–682 (2014).

16. Sutaria, D. S. *et al.* First penicillin-binding protein occupancy patterns of - Lactams and -

lactamase inhibitors in klebsiella pneumoniae. *Antimicrobial Agents and Chemotherapy* **62**,

(2018).

17. Kocaoglu, O. & Carlson, E. E. Profiling of β-Lactam Selectivity for Penicillin-Binding Proteins in

Escherichia coli Strain DC2. *Antimicrobial Agents and Chemotherapy* **59**, 2785–2790 (2015).

18. Sarigul, N., Korkmaz, F. & Kurultak, İ. A New Artificial Urine Protocol to Better Imitate Human

Urine. doi:10.1038/s41598-019-56693-4.

19. Bush, K. & Jacoby, G. A. Updated functional classification of β-lactamases. *Antimicrobial*

Agents and Chemotherapy **54**, 969–976 (2010).

20. Kannan, G. *et al.* Rapid acid treatment of Escherichia coli: Transcriptomic response and

recovery. *BMC Microbiology* **8**, 37 (2008).

21. Maurer, L. M., Yohannes, E., Bondurant, S. S., Radmacher, M. & Slonczewski, J. L. pH Regulates

Genes for Flagellar Motility, Catabolism, and Oxidative Stress in Escherichia coli K-12. *Journal*

of Bacteriology **187**, 304–319 (2005).

22. Sardis, M. F., Bohrhunter, J. L., Greene, N. G. & Bernhardt, T. G. The LpoA activator is required

to stimulate the peptidoglycan polymerase activity of its cognate cell wall synthase PBP1a.

Proc Natl Acad Sci U S A **118**, e2108894118 (2021).

23. Typas, A. *et al.* Regulation of peptidoglycan synthesis by outer-membrane proteins. *Cell* **143**,

1097–1109 (2010).

24. Typas, A. *et al.* Regulation of peptidoglycan synthesis by outer-membrane proteins. *Cell* **143**,

1097–1109 (2010).

25. Choi, U., Park, S. H., Lee, H. B., Son, J. E. & Lee, C.-R. Coordinated and Distinct Roles of

Peptidoglycan Carboxypeptidases DacC and DacA in Cell Growth and Shape Maintenance

under Stress Conditions. *Microbiology Spectrum* **11**, e00014-23 (2023).

26. Peters, K. *et al.* The redundancy of peptidoglycan carboxypeptidases ensures robust cell shape

maintenance in Escherichia coli. *mBio* **7**, (2016).

27. Identification of a penicillin-binding protein 3 homolog, PBP3x, in Pseudomonas aeruginosa:

gene cloning and growth phase-dependent expression.

https://journals.asm.org/doi/epdf/10.1128/jb.179.5.1490-1496.1997

doi:10.1128/jb.179.5.1490-1496.1997.

28. Castanheira, S. *et al.* A Specialized Peptidoglycan Synthase Promotes Salmonella Cell Division

inside Host Cells. *mBio* **8**, 10.1128/mbio.01685-17 (2017).

29. Castanheira, S. & García-del Portillo, F. Evidence of two differentially regulated elongasomes in

Salmonella. *Commun Biol* **6**, 923 (2023).

30. Sub-Inhibitory Cefsulodin Sensitization of E. coli to β-lactams Is Mediated by PBP1b Inhibition |

PLOS One. https://journals.plos.org/plosone/article?id=10.1371/journal.pone.0048598.

31. Ramachandran, V. *et al.* Screen for Inhibitors of the Coupled Transglycosylase-Transpeptidase

of Peptidoglycan Biosynthesis in Escherichia coli. *Antimicrobial Agents and Chemotherapy* **50**,

1425–1432 (2006).

32. Curtis, N. A. C., Eisenstadt, R. L., Turner, K. A. & White, A. J. Inhibition of penicillin-binding

protein 3 of Escherichia coli K-12. Effects upon growth, viability and outer membrane barrier

function. *Journal of Antimicrobial Chemotherapy* **16**, 287–296 (1985).

33. Buijs, J., Dofferhoff, A. S. M., Mouton, J. W., Wagenvoort, J. H. T. & van der Meer, J. W. M.

Concentration-dependency of β-lactam-induced filament formation in Gram-negative

bacteria. *Clinical Microbiology and Infection* **14**, 344–349 (2008).

34. Paradis-Bleau, C. *et al.* Lipoprotein Cofactors Located in the Outer Membrane Activate

Bacterial Cell Wall Polymerases. *Cell* **143**, 1110–1120 (2010).

35. Nichols, R. J. *et al.* Phenotypic Landscape of a Bacterial Cell. *Cell* **144**, 143–156 (2011).

36. Boes, A., Olatunji, S., Breukink, E. & Terrak, M. Regulation of the Peptidoglycan Polymerase

Activity of PBP1b by Antagonist Actions of the Core Divisome Proteins FtsBLQ and FtsN. *mBio*

10, 10.1128/mbio.01912-18 (2019).

37. Bertsche, U. *et al.* Interaction between two murein (peptidoglycan) synthases, PBP3 and

PBP1B, in Escherichia coli. *Molecular microbiology* **61**, 675–690 (2006).

38. Castanheira, S. *et al.* An alternative penicillin-binding protein involved in *Salmonella* relapses

following ceftriaxone therapy. *eBioMedicine* **55**, 102771 (2020).

39. van der Ploeg, R. *et al.* Colocalization and interaction between elongasome and divisome

during a preparative cell division phase in scherichia coli. *Molecular Microbiology* **87**, 1074–

1087 (2013).

40. Castanheira, S., López-Escarpa, D., Paradela, A. & García-del Portillo, F. In Vivo Cross-Linking

Sheds Light on the Salmonella Divisome in Which PBP3 and PBP3 Compete for Occupancy.

Molecular Microbiology **n/a**,.

41. Hardy, C. D. Essential Paralogous Proteins as Potential Antibiotic Multitargets in Escherichia

coli. *Microbiology Spectrum* **10**, e02043-22 (2022).

42. Machen, T. E. & Paradiso, A. M. Regulation of intracellular pH in the stomach. *Annu Rev Physiol*

49, 19–33 (1987).

43. Fallingborg, J. Intraluminal pH of the human gastrointestinal tract. *Dan Med Bull* **46**, 183–196

(1999).

44. O'Hanlon, D. E., Come, R. A. & Moench, T. R. Vaginal pH measured in vivo: lactobacilli

determine pH and lactic acid concentration. *BMC Microbiol* **19**, 13 (2019).

45. García-nafría, J., Watson, J. F. & Greger, I. H. OPEN IVA cloning : A single-tube universal cloning

system exploiting bacterial In Vivo Assembly. *Nature Publishing Group* 1–12 (2016)

doi:10.1038/srep27459.

46. Bachman, M. A. *et al.* Genome-wide identification of Klebsiella pneumoniae fitness genes

during lung infection. *mBio* **6**, (2015).

47. Datsenko, K. A. & Wanner, B. L. One-step inactivation of chromosomal genes in Escherichia

coli K-12 using PCR products. *Proceedings of the National Academy of Sciences* **97**, 6640–6645

(2000).

48. Ducret, A., Quardokus, E. M. & Brun, Y. V. MicrobeJ, a tool for high throughput bacterial cell

detection and quantitative analysis. *Nature Microbiology* **1**, 1–7 (2016).

Figure Legends

Fig 1. *Klebsiella pneumoniae* exhibits acid-dependent resistance to Beta-lactams.

(A) Heatmap displaying the fold-change in minimal inhibitory change (MIC) for *K. pneumoniae*

TOP52 at indicated pH relative to neutral pH. Each cell represents the median fold-change value

from a minimum of three biological replicates performed in LB buffered to indicated pH with MMT

buffer. Specific Beta-lactam antibiotics along with their preferential PBP target are indicated on the

Y-axis. NS = non-specific (B) The fold-change in Meropenem (MER) or Cephalexin (CEX) MIC of *K.*

pnuemoniae strains ATCC 43816 and NHUT-K2044 grown at pH 4.8 compared to growth at pH 6.8.

Bars represent mean and error bars denote standard deviation. C-D) Representative micrographs

of terminal morphologies of *K. pneumoniae* TOP52 following growth for 20 hours in LB (pH 4.8 or

6.8) and treated with Meropenem (C) or Cephalexin (D) at indicated concentrations. Scale bar = 2

µm.

Fig 2. **Loss of either PBP1b or PBP3$_{PARA}$ results in differential impacts to beta-lactam**

resistance at low pH. (A-D) Fold change in MICs at pH 4.8 relative to pH 6.8 shown for generalist

(A), class A-targeting (B), PBP2-targeting (C), and PBP3-targeting (D) beta-lactams. Data are

graphed as median values with range. (E) Micrographs from time-lapse microscopy of WT (top) or

Δ*mrcB* (bottom) growing in the presence of CEX at low pH. Microscopy pads comprised of LB

buffered with MMT to pH 4.8 + 15.6 µg/mL CEX + 1.5 µM propidium iodide + 1% agarose. The CEX

concentration reflects the 1X MIC for the Δ*mrcB* mutant and 0.06X MIC for the WT strain.

Fig 3. *Klebsiella pneumoniae* possesses acid-responsive paralogous copies of the Class B

PBPs. (A) Genomic regions containing the canonical PBP2, encoded by *mrdA*, and the paralogous

gene *mrdA2,* which encodes PBP2$_{PARA}$ (B) Genomic regions containing the canonical PBP3,

encoded by *ftsI,* and the paralogous gene *ftsI2,* which encodes PBP3$_{PARA}$. (C & D) Domain

conservation between canonical and paralogous copies of PBP2 (C) and PBP3 (D). Domains are

denoted as follows: Transmembrane (TM), Lipoprotein lipid attachment site (LP, PS51257) PBP-

dimer (Pfam: PF03717), Transpeptidase domain (Pfam: PF00905). The location of motifs that are

critical for enzyme catalysis are highlighted in purple.

Fig 4. Loss of PBP3$_{PARA}$ impairs division at low pH. (A) Representative micrographs of indicated

strains following growth in LB buffered to pH 4.8 (top panels) or growth in LB buffered to pH 6.8

(bottom panels). For complementation studies with either empty vector (EV) or p*ftsI2,* the media

was supplemented with 0.1 mM IPTG. Scale bar = 10 uM. (B-C) Quantification of cell length in (B).

Each data point represents the mean of a biological replicate (>50 cells analyzed for each

replicate). The bar represents the mean of three biological replicates, and the error bars represent

the standard error of the mean. Statistical significance was determined by a one-way ANOVA with

Tukey's correction for multiple comparisons with asterisks denoting significance as follows:** =

$p<0.005$, * = $p<0.5$.

Fig 5. Loss of either PBP1b or PBP3$_{PARA}$ results in differential impacts to Beta-lactam

resistance at low pH. (A-D) Fold change in MICs at pH 4.8 relative to pH 6.8 shown for generalist

(A), class A-targeting (B),PBP2-targeting (C), and PBP3-targeting Beta-lactams. For comparison,

data for MER and CEX are re-graphed from Fig 4. Data are presented as median values with range.

(E) Micrographs from time-lapse microscopy of WT, Δ*ftsI2,* Δ*mrdA2,* and Δ*mrdA2* Δ*ftsI2* growing in

the presence of MER at low pH. Microscopy pads comprised of LB buffered with MMT to pH 4.8 +

0.2 µg/mL MER + 1.5 µM propidium iodide + 1% agarose. The MER concentration reflects 1X MIC for

WT, ~0.25X MIC for Δ*ftsI2,* ~2X MIC for Δ*mrdA2,* and ~0.5X MIC for Δ*mrdA2* Δ*ftsI2.*

Fig 2

A

B

C

D

E

F

Fig 3

Fig 4

Fig 5

Table 1. Transcriptional response to pH of key beta-lactam targets in *K. pneumoniae* and *E. coli*

	Gene	Enzyme Encoded	Description	log$_2$FC (pH 4.8/6.8)
K. pneumoniae				
D1637_RS05825	*mrcA*	PBP1a	bifunctional glycosyl transferase/transpeptidase	**0.60**
D1637_RS12385	*mrcB*	PBP1b	bifunctional glycosyl transferase/transpeptidase	**-0.39**
D1637_RS01040	*pbpC*	PBP1c	peptidoglycan glycosyltransferase	**-1.61**
D1637_RS15400	*mrdA*	PBP2	canonical peptidoglycan DD-transpeptidase	**0.84**
D1637_RS12000	*ftsI*	PBP3	canonical peptidoglycan DD-transpeptidase	**0.23**
D1637_RS21590	*mrdA2*	PBP2$_{para}$	paralogous peptidoglycan DD-transpeptidase	**3.41**
D1637_RS24275	*ftsI2*	PBP3$_{para}$	paralogous peptidoglycan DD-transpeptidase	**3.28**
D1637_RS15385	*dacA*	PBP5	D-Ala-D-Ala-carboxypeptidase	**1.20**
D1637_RS25010	*dacD*	PBP6b	D-Ala-D-Ala carboxypeptidase	**1.63**
D1637_RS17685	*lpoB*	LpoB	Outer membrane activator of PBP1b	**-1.12**
E. coli				
b3396	*mrcA*	PBP1a	bifunctional glycosyl transferase/transpeptidase	**0.36**
b0149	*mrcB*	PBP1b	bifunctional glycosyl transferase/transpeptidase	**-0.23**
b2519	*pbpC*	PBP1c	peptidoglycan glycosyltransferase	**-0.43**
b0635	*mrdA*	PBP2	canonical peptidoglycan DD-transpeptidase	**0.47**
b0084	*ftsI*	PBP3	canonical peptidoglycan DD-transpeptidase	**-0.64**
b0632	*dacA*	PBP5	D-alanyl-D-alanine carboxypeptidase	**0.95**
b2010	*dacD*	PBP6b	D-alanyl-D-alanine carboxypeptidase	**2.63**
B1105	*LpoB*	LpoB	Outer membrane activator of PBP1b	**-1.33**

*Shaded genes represent those that were differentially regulated by pH as determined by our cutoff of |logFC| > 1 and $p < .05$. Blue Log$_2$FC values indicate that the gene was down-regulated during growth at pH 4.8 versus pH 6.8. Magenta Log$_2$FC values indicate that the gene was up-regulated during growth at pH 4.8 versus pH 6.8.