Title: pH-dependent beta-lactam resistance in *Klebsiella pneumoniae* is mediated by paralogous

class B PBPs and the class A PBP, PBP1b.

Authors

Sarah Beagle[1] and Petra Anne Levin[1*]

Affiliations

[1]Washington University in St. Louis, Department of Biology, St. Louis, Missouri, USA

*Correspondence: plevin@wustl.edu

Abstract

Klebsiella pneumoniae is a leading cause of global deaths due to antibiotic resistance. Of

particular concern, is the rapid expansion of resistance to beta-lactam antibiotics within *K.*

pneumoniae lineages. The environmental factors that influence pathogen physiology and,

subsequently, antibiotic resistance remain poorly understood. Here we demonstrate that

physiologically-relevant reductions in pH increased antibiotic resistance as much as 64-fold, with

the most dramatic increases observed for beta-lactams that inhibit cell division. We identified two

genes that contribute to acid-dependent beta-lactam resistance, the class A PBP, PBP1b, and the

paralogous class B PBP, PBP3$_{PARA}$. Loss of either gene increases *K. pneumoniae* susceptibility to

beta-lactams at low pH. Our data supports previous work from our group and others indicating that

functional redundancy among cell wall synthesis enzymes allows for specialization and ensures

robust cell wall synthesis across a range of environmental conditions.

Introduction

The cell wall or peptidoglycan (PG) is an essential structure that provides shape and structural

rigidity to bacterial cells. PG is comprised of glycan strands of repeating linked N-

acetylglucosamine (NAG) and N-acetylmuramic acid (NAM) sugars that are crosslinked together by

short peptide chains [1]. Synthesis and maintenance of PG requires many enzymatic steps that

occur across multiple cellular locations; PG precursors are built in the cytoplasm, transported

through the inner membrane, and fully assembled into PG within the periplasmic space. Within

members of *Enterobacteriaceae* the process of periplasmic PG assembly is highly enzymatically

redundant, with approximately four enzymes capable of carrying out a single periplasmic PG

synthesis reaction. This apparent functional redundancy is particularly striking when compared to

the earlier cytoplasmic PG synthesis reactions, which occur in a nearly 1:1 enzyme-to-reaction

ratio[2]. While cell wall precursors are assembled in the relatively protected cytoplasm, the wall

itself is built in the periplasm and thus exposed to the physicochemical environment.

Penicillin-binding proteins (PBPs) carry out essential transpeptidation reactions in the periplasm.

Class A PBPs are bi-functional enzymes that both link the NAG-NAM sugar moieties together

(transglycosylation, TG) as well as crosslink the peptide stems of adjacent glycan strands

(transpeptidation, TP). In *Escherichia coli,* the primary class A PBPs are PBP1a and PBP1b, which

are encoded by *mrcA* and *mrcB* respectively. PBP1a and PBP1b are individually dispensable for

growth, but loss of both is lethal[3–5]. A third class A PBP, PBP1c (*pbpC*), has an unknown role in PG

synthesis and maintenance, as loss of this gene does not result in discernable defects under

standard laboratory conditions[6]. Class B PBPs are monofunctional transpeptidases; these PBPs

work in conjunction with cognate glycosyltransferases. PBP2 and PBP3 are the canonical class B

PBPs in Gram-negative bacteria, and they function within the elongation and division machinery

respectively.

The essentiality of the bacterial cell wall renders the enzymes that synthesize PG attractive drug

targets. Beta-lactams, the most prescribed cell-wall-targeting class of antibiotics, interfere with PG

synthesis by irreversibly binding to the transpeptidase domain of PBPs and inhibiting

transpeptidation[7]. Work done in *Escherichia coli* demonstrated that resistance to Beta-lactam

varies across physiochemical conditions. Specifically, *E. coli* exhibits increased resistance to

PBP2- and PBP3-targeting Beta-lactams when cells are grown in mildly acidic pH conditions (~pH

5). Growth at low pH as well as acid-dependent Beta-lactam resistance required the Class A PBP,

PBP1b, while full fitness during growth in alkaline pH required PBP1a[8]. These results indicate that

the apparent functional redundancy in PG synthesis enzymes allows for specialization of enzymes,

thereby ensuring that cell wall synthesis is robust across a wide range of environmental conditions.

To determine if this response is conserved in a pathogen, we assessed the impact of environmental

pH on Beta-lactam resistance in *Klebsiella pneumoniae,* a Gram-negative, encapsulated,

opportunistic human pathogen.

Infections caused by *K. pneumoniae,* a close relative of *E. coli,* are increasingly difficult to treat

due to dramatic rises in antibiotic resistance. *K. pneumoniae* is a dominant member of two

different Beta-lactam-resistant groups of bacteria, Extended-spectrum-Beta-lactamase (ESBL)-

producing *Enterobacteriaceae* and Carbapenem-resistant *Enterobacteriaceae* (CRE) that have

been labeled serious and urgent public health threats by the CDC[9]. In 2019, *K. pneumoniae* was the

third leading cause of global deaths where the death was directly due to or associated with

antibiotic resistance[10], highlighting the need to explore how environmental factors impact

pathogen physiology and subsequently, antibiotic resistance.

Here we evaluated the impact of pH on *K. pneumoniae* response to beta-lactams and found that

resistance increased as much as 64-fold during growth in mildly acidic conditions. Genetic

analysis identified two cell wall synthesis genes, *mrcB* and *ftsI2* that are critical for pH-dependent

changes in antibiotic resistance. *mrcB* encodes PBP1b, a bifunctional class A PBP we previously

identified as a contributor to pH-dependent antibiotic resistance in *E. coli*[8]. Induced at low pH,

ftsZ2 encodes PBP3$_{PARA,}$ a paralogous copy of the canonical, division-associated class B PBP,

PBP3. PBP3$_{PARA}$ is required for both acid-dependent beta-lactam resistance and robust growth and

division at low pH, highlighting the fitness advantage conferred by functional redundancy amongst

bacterial PBPs. **Results**

***Klebsiella pneumoniae* exhibits acid-dependent resistance to Beta-lactam antibiotics.**

To determine the impact of pH on *K. pneumoniae* beta-lactam susceptibility, we assessed the

response of a urinary tract infection (UTI) isolate, *K. pneumoniae* TOP52[11] to a panel of beta-

lactams during growth across a range of physiologically relevant pH conditions. The drug panel

included beta-lactams that preferentially target PBPs in the divisome (i.e. Cephalexin, Aztreonam,

and Piperacillin) or the elongasome (Meropenem, Doripenem), as well as non-specific inhibitors

(Ampicillin, Carbenicillin) that have similar affinities for all PBPs[12]. Relative susceptibility or

resistance to a given Beta-lactam was determined by comparing the minimal inhibitory

concentration of drug at neutral pH to that at other pH values.

K. pneumoniae exhibited a 2-64-fold increase in resistance to the entire spectrum of tested beta-

lactams at pH values < 5 relative to resistance at pH 7. (Fig 1A, Sup Table 3). We observed strong

acid-dependent increases in resistance for drugs that preferentially target PBP3, with Aztreonam

(AZT) and Cephalexin (CEX) MICs increasing as much as 20- and 64-fold respectively at pH 4.8.

Resistance to PBP2-targeting Beta-lactams increased modestly at low pH, with MICs to Doripenem

(DOR) and Meropenem (MER) increasing 2- to 4-fold. In contrast to *E. coli*[8], we also observed a

modest 2- to 4-fold increases in resistance to the non-specific Beta-lactams, Ampicillin (AMP) and

Carbenicillin (CAR) at low pH.

Low pH not only enabled growth at concentrations of drug that are lethal at neutral pH but also

preserved cell morphology (Fig 1 C & D). Consistent with increased intrinsic resistance at low pH,

the characteristic rod-shaped morphology of *K. pneumoniae* was preserved during growth under

acidic conditions in concentrations of Beta-lactams that cause either cell rounding (Meropenem)

or filamentation (Cephalexin) at neutral pH. The observed acid-dependent increased resistance to

beta-lactams is not strain specific behavior as *K. pneumoniae* strains NTUH-K2044 and ATCC

43816 also exhibit a 4-fold increase to Meropenem (MER) and an ~32-fold increase to Cephalexin

(CEX) when grown at pH 4.8 versus pH 6.8 (Fig. 1B). Together, these data suggest that acid-

dependent beta-lactam resistance is a conserved response within *K. pneumoniae.*

Acid-dependent Beta-lactam resistance is independent of endogenous beta-lactamase

expression.

K. pneumoniae encodes an endogenous class-A beta-lactamase, *bla$_{SHV-1}$,* which confers increased

resistance to the Penicillin subclass of Beta-lactams[7,11,13]. To determine if expression of *bla$_{SHV-1}$* is

influencing the resistance profiles at low pH, we generated a deletion strain (Δ*bla$_{SHV-1}$*) and

repeated the MIC assays under both acidic and neutral pH conditions. We observed increased

sensitivity to the Penicillin sub-classes of Beta-lactams tested (AMP, CAR, PIP) in both pH

environments (Sup Table 5). Consistent with the specificity of *bla$_{SHV-1}$,* loss of *bla$_{SHV-1}$* did not alter

the MICs to non-Penicillin beta-lactams at either pH (Sup Table 5). However, the Δ*bla$_{SHV-1}$* mutant

still displayed increased resistance to all tested Beta-lactams at low pH relative to neutral pH (Fig 2

A-D, Supp table 5), demonstrating that acid-dependent resistance is independent of endogenous

beta-lactamase expression.

pH modulates expression of Class B PBP paralogs in *K. pneumoniae*.

To ascertain the impact of pH on expression of genes encoding PBPs or other cell wall synthesis

genes, we performed RNAseq in *K. pneumoniae* comparing cells cultured in acidic (pH 4.8) versus

neutral pH media (pH 6.8). Consistent with previous studies, pH does not impact expression of

class A PBPs, *mrcA* (PBP1a) or *mrcB* (PBP1b), in either *E. coli* or *K. pneumoniae*[14,15] (Table 1) .

PBP1c, a cryptic class A PBP with no demonstrated role in cell wall synthesis under standard

laboratory conditions, exhibits an ~2-fold decrease in expression during growth in acidic pH. We

observed no significant changes expression of the class B PBPs between pH conditions, but there

was an order of magnitude increase in expression at low pH of genes we designated *mrdA2* and

ftsI2, which encode paralogous copies of class B PBPs, referred to here as PBP2$_{PARA}$ and PBP3$_{PARA}$

(Fig 3 A&B). In addition to the class B paralogs, several other cell wall synthesis genes displayed

pH-responsive expression profiles. Most notably, two carboxypeptidases, *dacA* (PBP5) and *dacD*

(PBP6b)*,* exhibited modest increases of 2- to 3-fold in expression during growth at low pH. Both

dacA and *dacD* are implicated in maintaining proper cell morphology in *E. coli*, with *dacD* being

required for proper morphology specifically at low pH[16,17].

Loss of PBP3$_{PARA}$ results in filamentation in *K. pneumoniae* at low pH.

Duplications of the class B PBPs, PBP2 and PBP3, have expanded the repertoire of cell wall

synthesis enzymes in several pathogens within the *Enterobacteriaceae* family[18–20]. Class B PBP

paralogs in *Salmonella enterica* serovar Typhimurium (*S.* Typhimurium) referred to as PBP2$_{SAL}$ and

PBP3$_{SAL}$ are thought to be important for maximal fitness during growth at low pH. Expression of both

PBP2$_{SAL}$ and PBP3$_{SAL}$ is maximally induced at low pH (<5.8), and loss of PBP2$_{SAL}$ has been reported to

impair rod-shape morphology under acidic environments[19,20].

As with PBP2$_{SAL}$ and PBP3$_{SAL}$, PBP2$_{PARA}$ and PBP3$_{PARA}$ share significant sequence identity with their

canonical counterparts (59% for PBP2$_{PARA}$ and 61% for PBP3$_{PARA}$) and retain the sequence motifs

associated with catalysis, but their contributions to cell wall maintenance in *Klebsiella* is unknown

[Fig 3 C&D, Sfig 2]. We generated individual gene deletion strains and examined the mutants using

cell size analysis and time-lapse microscopy to evaluate the role of the class B PBP paralogs in cell

size and division. We cultured cells at both acidic and neutral pH and sampled cultures at early

exponential phase (OD$_{600}$ = 0.1-0.2) for cell size analysis. While loss of PBP2$_{PARA}$ did not impact cell

size or morphology at either pH, we found that during growth at pH 4.8 the PBP3$_{PARA}$ deletion mutant

exhibited heterogeneous cell lengths, with many cells displaying filamentous behavior indicative of

cell division defects (Fig4 A&B, Sfig 2). Using a definition of filamentation as a cell length of >15 µM

which represents ~3X the average length of WT cells at pH 4.8 (3.9 ± 0.2 µM, Fig 4B), we determined

that 21-58% of PBP3$_{PARA}$ deficient cells within a given population were filamentous, with a smaller

proportion of cells (3-17%) reaching lengths of >40 µM. The division defects associated with loss of

PBP3$_{PARA}$ were specific to growth in acidic conditions, as growth at neutral pH resulted in an average

cell length for the PBP3$_{PARA}$ deficient cells that was consistent with WT cell lengths (Fig 3C).

Time-lapse microscopy of acid-grown Δ*ftsI2* cells spotted onto acidic LB agarose pads revealed a

population of cells with heterogeneous lengths that displayed clear division defects as cells

continued to elongate without successful division resulting in long filamentous cells. Whereas

spotting acid-grown Δ*ftsI2* cells onto LB agarose pads at neutral pH restored division to the

filamentous cells (Sfig 3). Complementation of *ftsI2* in trans from a medium-copy number plasmid

enabled division and restored cell lengths consistent with WT at low pH (Fig 4 A&B). Together these

data suggest that the activity of the canonical PBP3 is impaired under acidic conditions and that

PBP3$_{PARA}$ is the predominant transpeptidase supporting cell division in *K. pneumoniae* at low pH.

Loss of PBP1b or PBP3$_{PARA}$ differentially impacts Beta-lactam resistance at low pH in *Klebsiella*

The increased expression of genes encoding PBP2$_{PARA}$ and PBP3$_{PARA}$ and other cell wall

maintenance proteins at low pH raised the possibility that these proteins might contribute to acid-

dependent Beta-lactam resistance. To address this hypothesis, we generated strains individually

defective in a subset of *K. pneumoniae* genes implicated in cell wall or shape maintenance and

assayed their response to beta-lactam antibiotics during growth at acidic and neutral pH

conditions. In addition to genes that displayed pH-dependent expression, we also evaluated the

role of class A PBPs in *K. pneumoniae* response to beta-lactams across pH conditions. The class A

PBPs are known to influence antibiotic response in *E. coli*. While most individual mutations did not

significantly impact acid-dependent-Beta-lactam resistance, loss of PBP1b or PBP3$_{PARA}$ altered

sensitivities to Beta-lactams at low pH (Fig 2A-D).

Loss of PBP1b (Δ*mrcB*) most dramatically affected acid-dependent resistance to Cephalexin. A

Δ*mrcB* mutant displayed only a 4-fold increase in cephalexin MIC at low pH versus at neutral pH,

while the WT has a 64- to 128-fold increase in resistance under the same conditions. The Δ*mrcB*

mutant also no longer exhibited acid-dependent resistance to the generalist (AMP) and PBP2-

targeting- (DOR, MER) beta-lactams, (Fig 2 A-D). Defects in PBP1b also led to a nearly 10-fold

increase in susceptibility to Cefsulodin in both acidic and neutral pH compared to WT (Sup Table

5). The increase in sensitivity at all pHs of the ΔPBP1b mutant is consistent with data suggesting

that PBP1a is more sensitive to Cefsulodin[21,22].

Loss of PBP3$_{PARA}$ differentially impacted resistance to beta-lactams. The PBP3$_{PARA}$ null mutant no

longer displayed acid-dependent resistance to Ampicillin and Cefsulodin at low pH (Fig 2A & B). In

fact, a PBP3$_{PARA}$ deletion mutant was more sensitive to Ampicillin at low pH than at neutral pH (Fig

2A). When challenged with PBP3-targeting Beta-lactams (CEX, AZT, PIP), the PBP3$_{PARA}$ deletion

mutant still retained mild acid-dependent resistance but was 2- to 8-fold more sensitive at low pH

than WT (Fig 4 D, 5 B). Surprisingly, the PBP3$_{PARA}$ null mutant exhibited increased resistance when

challenged with PBP2-targeting beta-lactams at low pH. The Meropenem and Doripenem MICs

obtained for the PBP3$_{PARA}$ null mutant were 12-16-fold higher at acidic pH versus neutral pH. For

comparison, WT cells only exhibited a 2-4-fold increase in MIC to these two drugs at acidic pH (Fig

4C, 5A). Despite the ~10-fold increases in expression under acidic conditions, loss of *mrdA2,*

encoding PBP2, had no significant impact on Beta-lactam susceptibility across pH values.

187   Increased acid-dependent resistance to PBP2-targeting compounds in the PBP3$_{PARA}$ null

188   mutant requires PBP2$_{PARA}$

We reasoned that the unexpected increase to PBP2-inhibitors at low pH observed in the Δ*ftsI2*

mutant could be influenced by the presence of PBP2$_{PARA}$. To address this possibility, we generated

a double deletion strain lacking both class B paralogs (Δ*ftsI2* Δ*mrdA2*) and assayed its response to

beta-lactams with high specificity for PBP2 or PBP3 at both neutral and acidic pH. Deletion of both

class B PBP paralogs eliminated the dramatic increase in resistance at low pH (4.8), restoring WT-

level responses to both Doripenem and Meropenem. When challenged with PBP3-targeting Beta-

lactams, the double mutant class B paralog mutant phenocopied the Δ*ftsI2* mutant, displaying

similar reductions in resistance at low pH (Fig 5). Together these data suggest that both PBP2$_{PARA}$

and PBP3$_{PARA}$ are involved in mediating resistance to PBP2-inhibitors, whereas resistance to PBP3-

targeting Beta-lactams is primarily facilitated by PBP3$_{PARA}$.

199   Discussion

As observed in *E. coli,* beta-lactam resistance in *K. pneumoniae* increases substantially during

growth in acidified medium (Fig 1A).We find that the class A PBP, PBP1b, and the acid-responsive

paralogous copy of the class B PBP, PBP3$_{PARA}$, are major genetic contributors to acid-mediated

beta-lactam resistance in *K. pneumoniae*. Disruption of *mrcB*, the gene encoding PBP1b,

eliminates acid-mediated resistance to the class B PBP inhibitors, Meropenem, Doripenem, and

Cephalexin. Loss of PBP3$_{PARA}$ differentially impacts beta-lactam resistance, with mutants showing

sensitization towards Ampicillin, Cefsulodin, and all PBP3-targeting beta-lactams but increased

resistance towards PBP2-targeting antibiotics.

While both *E. coli* and *K. pneumoniae* demonstrate acid-dependent beta-lactam resistance,

factors influencing this phenomenon differ between the two organisms. Acid-mediated beta-

lactam resistance in *E. coli* appears to be solely due to intrinsic properties of the PBP enzymes,

particularly PBP1b[8], while in *K. pneumoniae* the contribution of PBP1b is supplemented by pH-

dependent induction of genes encoding paralogous class B PBPs. Although we only observed

changes to antibiotic susceptibility in a PBP3$_{PARA}$ null mutant, loss of PBP2$_{PARA}$ impacted the

morphological response to PBP2-targeting antibiotics at low pH (Fig 5 C). This result suggests that

PBP2$_{PARA}$ plays a role in maintaining PG and cell shape at low pH. Together, our findings support a

modular model of cell wall synthesis within the *Enterobacteriaceae* family where apparent function

redundancy amongst cell wall synthesis enzymes is protective by enabling PG synthesis across a

range of pH conditions.

The Class A PBP, PBP1b, is important for acid-dependent beta-lactam resistance in *K.*

pneumoniae

PBP1b is required for resistance to PBP2- and PBP3-targeting beta-lactams and for growth at low

pH (<5) in *E. coli*[8]. In contrast, deletion of PBP1b in *K. pneumoniae* did not significantly impair

growth in acidified culture medium (pH 4.8), suggesting that there are key differences in how PBP1b

influences growth in acidic environments between *E. coli* and *K. pneumoniae* (Sfig 4). While

expression of *mrcB,* encoding PBP1b, was not influenced by pH, loss of PBP1b impaired *K.*

pneumoniae's resistance to Ampicillin, Doripenem, Meropenem, and Cephalexin at low pH (Fig 2,

Supp Table 5). Consistent with MIC data, a PBP1b-deficient strain exhibits faster killing kinetics

than a WT strain in the presence of Cephalexin at low pH (Fig 2 E &F). While filamentation is a

characteristic morphology associated with division-targeting beta-lactams[23,24], at low pH a PBP1b-

deficient *K. pneumoniae* strain is sensitive to Cephalexin at concentrations that do not elicit

filamentation (Fig 4E). The time frame for lysis of the PBP1b mutant is rapid with more than 50% of

the population exhibiting signs of membrane integrity as denoted by propidium iodide staining

within 90 minutes of exposure to Cephalexin at low pH (Fig 2F).

Notably, at neutral pH a *ΔmrcB* mutant did not display significant changes in MIC across beta-

lactam classes (Sup Table 5), suggesting that PBP1B specifically mediates beta-lactam resistance

at low pH in *K. pneumoniae*. This finding differs from what has been observed in *E. coli* where loss

of either PBP1b or its cognate outer membrane activator, LpoB, increases sensitivity to multiple

classes of beta-lactams at neutral pH[25,26]. Our data demonstrating that a PBP1b mutant is more

sensitive to PBP2- and PBP3-targeting beta-lactams is consistent with prior studies reporting

interactions between PBP1b and key division and elongasome proteins[27,2820]. Based on these data,

we propose that PBP1b supports cell wall integrity at low pH through interactions with other cell

division proteins.

Class B PBP paralogs modulate pH-dependent changes in beta-lactam resistance

Although, low pH induced expression of genes encoding both *K. pneumoniae's* PBP2 and PBP3

paralogs (Table 1), only defects in *ftsI2,* encoding PBP3$_{PARA}$, sensitized cells to beta-lactams during

growth in acidified medium. Defects in *mrdA2*, encoding PBP2$_{PARA}$, did not significantly impact

beta-lactam resistance at either acidic or neutral pH. These data are consistent with what has

been observed in *S.* Typhimurium in which loss of its PBP3 paralog, PBP3$_{SAL}$sensitizes cells to beta-

lactams targeting division[29]. In *K. pneumoniae* loss of PBP3$_{PARA}$ at low pH sensitized cells to not only

PBP3 inhibitors, but also to a Cephalosporin targeting class A PBPs and the penicillin, Ampicillin

(Figure 4 A&B). While loss of PBP3$_{PARA}$ generally sensitized *K. pneumoniae* to beta-lactams under

acidic conditions, it increased resistance to compounds targeting PBP2 at pH 4.8 up to 12- to 16-

fold (Figure 5A). This increase in resistance in PBP3$_{PARA}$ deficient cells was eliminated in a double

mutant lacking both PBP3$_{PARA}$ and PBP2$_{PARA}$ (Δ*ftsI2* Δ*mrdA2*). Under acidic conditions the double

class B paralog mutant exhibited WT-level MICs when challenged with Meropenem or Doripenem

(Figure 5).

It is unclear why the loss of PBP3$_{PARA}$ renders cells resistant to PBP2-targeting compounds.

Morphological data of WT *K. pneumoniae* challenged with Meropenem at low pH reveals hybrid

morphologies, with cells displaying signatures of both divisome and elongasome inhibition. Cells

first undergo lengthening and then display large, rounded septal bulges (Fig 5C). These results

could reflect altered target binding by Meropenem at low pH, where PBP3/PBP3$_{PARA}$ are also

inhibited along with PBP2/PBP2$_{PARA}$. Alternatively, physical interactions between class B PBPs

within the elongasome and divisome have been reported in *E. coli*[30], and data suggests that the

protein components of the PG synthesis complexes in *Salmonella* vary depending on whether the

canonical or paralogous class B PBP is present in the complex[20,31]. Alterations to interactions

between class B PBP paralogs and other cell wall synthesis proteins due to beta-lactam binding or

in the class B PBP paralog mutant backgrounds could influence morphology and antibiotic

resistance. Additional work is needed to fully understand the complex role that the class B PBP

paralogs play in resistance to PBP2-targeting beta-lactams.

PBP3$_{PARA}$ is critical for *K. pneumoniae* division at low pH

Our data suggest that PBP3$_{PARA}$ is the major transpeptidase supporting cross wall synthesis at low

pH. Loss of PBP3$_{PARA}$ did not impact cell length at neutral pH but impaired division at low pH

resulting in filamentous cells. However, these acid-cultured PBP3$_{PARA}$-deficient filamentous cells

were able to resume division and return to WT-lengths if they were transitioned to neutral pH

(Sfig3), suggesting that the canonical PBP3 is unable to fully compensate for the loss of PBP3$_{PARA}$ at

low pH. We attempted to disrupt the canonical PBP3 under acidic culture conditions where

PBP3$_{PARA}$ is likely the dominate division-specific transpeptidase; however, despite multiple

attempts, we were unsuccessful. Moreover, a small percentage of Δ*ftsI2* cells maintained WT-

lengths at low pH, suggesting that the canonical PBP3 retains limited, but critical activity at low pH

(Sfig2). These data differ from *Salmonella* where the canonical PBP3 and PBP3 paralog PBP3$_{SAL}$ are

individually dispensable at low pH. The canonical PBP3 is capable of supporting growth at both low

and neutral pH, but PBP3$_{SAL}$ is unable to support division at neutral pH[19]. Our results support a

model in which *K. pneumoniae* PBP3 functions optimally at neutral pH while PBP3$_{PARA}$ functions

predominately at acidic pH, with both working together to ensure that division remains robust

across pH conditions.

Expansion and pH specialization of *K. pneumoniae* cell wall synthesis enzymes provide novel

therapeutic targets

Identification of paralogous class B PBPs has thus far only been described in pathogens [18–20,32], but

the role of these paralogous cell wall synthesis proteins in pathogen physiology remains largely

uncharacterized. In *Salmonella,* the acid-responsiveness of class B PBP paralogs raised

speculation that they could be critical to support bacterial survival during intracellular growth

within acidified host immune cells. However, loss of the class B PBP paralogs did not impair

replication within macrophages or attenuate virulence in a mouse infection model[19].

In *Klebsiella pneumoniae* our data suggests that class B PBP paralogs are part of a general strategy

of functional redundancy to preserve critical enzymatic activity across a wide range of

environmental conditions. Colonization of a host exposes a bacterial pathogen to a wide range of

pH conditions[33–35] . We demonstrate that the acidification of growth medium to physiologically-

relevant levels is sufficient to increase resistance to beta-lactams. The class B PBP paralogs

modulate that resistance in part, suggesting that these pathogen-specific proteins may provide

novel therapeutic targets. We propose that future screens for new antimicrobial targets factor

environmental pH as a variable in order to identify novel compounds that can bind to acid-

responsive targets such as the Class B PBP paralogs. Altogether, our results highlight the need to

evaluate pathogen physiology in diverse environments beyond the standard laboratory conditions.

Acknowledgements

We are indebted to members of the Levin lab for their feedback and discussion regarding this

study. We alsothank Kyra Raines and Dr. Sarah Anderson for critical reading of this manuscript.

This work has been funded by the NIH through an G35-GM127331 award to P.A.L.

Contributions

S.B.: Conceptualization, Investigation, Formal analysis, Methodology, Visualization, Writing-

original draft, Writing – review and editing.

P.A.L.: Conceptualization, Funding acquisition, Resources, Supervision, Writing – review and

editing.

Data availability

All raw and processed data required to evaluate the conclusions of this study are included in the

text or supplemental data. The RNA sequencing data for gene expression studies at acidic and

neutral pH have been deposited in the NCBI GEO repository: GSE291667.

Methods

Bacterial strains, growth conditions, and antibiotics.

Bacterial strains and plasmids used in this study are listed in Supplementary table 1 and

Supplementary table 2. Strains were routinely cultured in Lysogeny Broth (LB – 1% Tryptone, 0.5%

Yeast extract, and 1% NaCl) in large glass tubes at 37 °C with 200 RPM aeration. The pH of LB was

adjusted with 1:10 MMT buffer (1:2:2 molar ratio of D/L-malic acid, MES, and Tris base). Where

indicated cultures were supplemented with Kanamycin (50 µg/mL), Spectinomycin (100 µg/mL),

Chloramphenicol (50 µg/mL), or IPTG (0.1 mM) . The pRDC3 plasmid (p15A ori, SpecR, Plac) was a

gift from Fabrizio Arigoni. Plasmid pSDB8 was generated from pRDC3 using the IVA cloning method

to insert *ftsI2* under control of the Plac promoter[36]. Primers were acquired from Integrated DNA

Technologies (Coralville, IA) and sequences used are listed in Supp Table 3. Chemically competent

E. coli TOP10 were used to assemble vectors via IVA. Plasmid sequence was confirmed via whole

plasmid sequencing at Plasmidsaurus prior to transformation into *K. pneumoniae* strains.

For growth studies, cultures were started LB supplemented with MMT buffer to the indicated pH

and grown in large glass tubes at 37 °C with 200 RPM aeration. Cultures were harvested once they

reached mid-exponential phase (OD_{600} =0.2-0.6), concentrated to an OD_{600} =1.0, and serially

diluted 10-fold. 1.5 µL of the 10^{-5} dilution was used to inoculate wells of sterile 96-well plates

containing 150 µL of LB + MMT at indicated pH. The plate was covered with a Breathe-Easy strip

(Diversified Biosciences) and growth was monitored in a BioTek Synergy H1 plate reader with the

following settings: 37 °C, continuous double orbital shaking at 282 cpm, with OD_{600} readings taken

every 10 minutes.

For *ftsI2* cell size complementation studies, un-induced overnight cultures of strains carrying either

pRDC3 (Empty vector, EV) or pSDB8 were back-diluted 1:500 into 3 mL of fresh LB buffered to

either pH 4.8 or pH 6.8 containing Spectinomycin (100 µg/mL). Cultures were grown at 37 °C with

200 RPM aeration for 1.5 hrs. Expression of *ftsI2* was induced with the addition of 0.1 mM IPTG and

cultures were allowed to continue growing until early exponential phase (OD_{600} = 0.1-0.3) was

reached (~40-90 additional minutes). Cultures were back-diluted into fresh buffered media

containing Spectinomycin (100 µg/mL) and 0.1 mM IPTG to an OD_{600} ~= 0.005. Growth was

monitored every 30 minutes, and cells were harvested when cultures reached early exponential

phase (OD_{600} = 0.1-0.3). Cells were spotted onto media pads (LB + MMT (pH 4.8 or 6.8) +

Spectinomycin (100 µg/mL) + 0.1 mM IPTG + 1% agarose and imaged live with phase microscopy.

The following antibiotics used for antibiotic susceptibility screenings. Ampicillin sodium salt,

Aztreonam, Cephalexin hydrate, Cefsulodin sodium salt hydrate, Meropenem Trihydrate, and

Piperacillin sodium salt were obtained from Sigma Adrich (St. Louis, MO). Carbenicillin sodium salt

was from EMD Millipore (now Millipore Sigma, Germany). Doripenem was obtained from Cayman

Chemicals (Ann Arbor, MI).

Genetic manipulation: Gene disruptions were carried out in the *K. pneumoniae* TOP52

background using a Lambda Red vector modified to carry a Spectinomycin resistance cassette as

described previously [37,38]. Briefly, DreamTaq Polymerase (Thermo Scientific) was used to PCR

amplify the Kanamycin resistance cassette from the template plasmid pKD4 with primers

containing 40-60 bp of homology to the chromosomal targets. Primers were designed to leave the

start and stop codons of the targeted gene intact. Linear PCR products were purified using a

Purelink PCR purification kit (Invitrogen), and electroporated into *K. pneumoniae* cells expressing

the Lambda Red recombinase system from the pKD46S plasmid. Electroporated cells were

recovered in SOC (Lloyd's recipe) for 4-8hrs at 30°C. Deletion mutants were selected for by plating

transformation culture onto LB agar plates containing Kanamycin (50 µg/mL) and incubated at

37°C. Mutants were confirmed by colony PCR and sequencing as necessary. For marker-less

deletions, cells were transformed with pCP20 at 30°C in LB + Chloramphenicol (50 µg/mL) to

express the FLP to excise the FRT-flanked Kanamycin cassette. After excision, cells were cultured

at 42°C to promote loss of pCP20. Kanamycin and Chloramphenicol sensitivity was confirmed for

strains prior to use.

RNA sequencing: *K. pneumoniae* TOP52 or *E. coli* MG1655 was cultured overnight in LB buffered

to either pH 4.8 or 6.8 with MMT buffer. Overnight cultures were back diluted 1:100 into fresh media

and cultured at 37°C with 200 RPM aeration until cultures reached early exponential phase (OD_{600}

=~0.2). ~ 9.6×10^8 cells were transferred to 15 mL conical Falcon tubes and pelleted by

centrifugation at 4°C (4,000 RPM for 10 mins). The supernatant was discarded, cell pellets were

resuspended in 350 uL of RNAwiz, and the RNA was extracted following the protocol for the

Ribopure-Bacteria RNA extraction kit [Invitrogen]. RNA was eluted in 50 uL of elution solution.

Purified RNA was subjected to a DNase I treatment to remove any contaminating chromosomal

DNA. RNA sequencing and differential gene expression analysis was performed by MiGS. Quality

control and adapter trimming of reads was performed with bcl2fastq. Read mapping was

performed with HISAT2, and read quantification was performed using Subread's featureCounts.

Read counts were normalized in R using edgeR's Trimmed Mean of M values (TMM) algorithm, and

subsequent values were converted to counts per million (CPM). Differential expression analysis

was performed using edgeR's Quasi-Linear F-Test (qlfTest) functionality against treatment groups,

and DEGs were considered as those with |logFC| > 1 and p < .05. Three biological replicates were

submitted per pH condition.

Antibiotic susceptibility testing. Cells were cultured from a single colony in LB at the indicated

pH to exponential phase (OD_{600} =~0.2-0.6) at 37 °C with 200 RPM aeration. Harvested cells were

concentrated to an OD_{600} =1 and serially diluted 10-fold. 1.5 µL of the 10^{-2} dilution was used to seed

wells containing 150 µL of LB broth + MMT (matching pH) in sterile 96-well microtiter plates

containing a range of two-fold dilutions of beta-lactam antibiotics. Plates were sealed with a

Breathe-Easy strip (Diversified Biotech) and incubated for 20 hours at 37°C with 200 RPM aeration.

The minimal inhibitory concentration (MIC) was determined by visual inspection and reflects the

lowest drug concentration that completely inhibited bacterial growth.

Terminal morphology Assessment. At the end of 20 hours for antibiotic susceptibility tests, cells

were taken from wells following growth at either neutral or acidic pH with the indicated antibiotic

concentrations. 5 µL of cells were spotted onto LB + 1% agarose and imaged by phase contrast

microscopy.

Microscopy, cell size analysis, and time-lapse Imaging. All phase contrast and fluorescence

images were acquired on either a Nikon Ti-E inverted microscope or a Ti2-E inverted microscope

(Nikon Instruments). The Ti-E is equipped with a 100X Plan Apo oil objective (N.A. = 1.45 Ph3

objective, X-Cite 120 LED light source (Lumen Dynamics) and an OrcaERG CCD camera

(Hammamatsu Photonics, Bridgewater, N.J.) Filter cubes were purchased from Chroma

Technology Corporation. The Ti2-E inverted microscope is equipped with a 100X Plan Apo oil

objective (N.A. = 1.45, Ph3 objective, Lumencor Sola LED light engine (Lumencore), and an orca-

Fusion sCMOS camera (Hammamatsu Photonics). Filter cubes were purchased from Chroma

Technology Corporation. Both microscopes have an enclosed temperature-controlled enclosure to

pre-heat the objective to 37 °C (OXO enclosure) prior to experiments. Nikon Elements software was

used for image capture, and analysis was performed using the MicrobeJ plug-in in ImageJ.

For cell size analysis, cells were grown from a single colony in LB with MMT buffer at indicated pH

at 37 °C and 200 RPM aeration to exponential phase (OD_{600} = 0.2-0.4) and back-diluted to an OD_{600}

= 0.005 into 7 mL of pre-warmed LB + MTT (same pH). Growth was monitored via optical density,

and when cultures reached early exponential phase (OD_{600} =0.1- 0.2) cells were harvested. 5 µL of

cells were spotted onto LB + MTT (same pH)+ 1% agarose pads and imaged by phase contrast

microscopy. Cell morphology and size dimensions were determined using the MicrobeJ[39] plug-in

for ImageJ.

For time-lapse imaging, cultures were grown from a single colony in LB with MMT buffer at

indicated pH at 37 °C and 200 RPM aeration to exponential phase (OD_{600} = 0.1-0.4). 5 µL of cells

were spotted onto LB + MTT (same pH)+ 1% agarose pads that contain antibiotics at indicated

concentrations and 1.5 µM propidium iodide (PI) for monitoring loss of membrane integrity. Phase

images were obtained every 2 minutes, and fluorescent images were collected every 10 minutes

(100 mS exposure/10% intensity). For each experiment a total of 2-5 different XY coordinates were

monitored throughout the duration of the experiment. Quantification of PI$^+$ cells was done using

the MicrobeJ[39] plug-in for ImageJ. Briefly the number of cells in the phase image and the number of

corresponding fluorescent cells were quantified. Data was reported as either percentage of PI$^+$

cells (number of PI$^+$ cells/total number of cells *100) or percent alive (number of cells-PI$^+$ cells/total

number of cells*100).

Bibliography

1. Egan, A. J. F., Errington, J. & Vollmer, W. Regulation of peptidoglycan synthesis and

remodelling. *Nature Reviews Microbiology 2020 18:8* **18**, 446–460 (2020).

2. Pazos, M., Peters, K. & Vollmer, W. Robust peptidoglycan growth by dynamic and variable

multi-protein complexes. *Current Opinion in Microbiology* **36**, 55–61 (2017).

3. Kato, J., Suzuki, H. & Hirota, Y. Dispensability of either penicillin-binding protein -1a or -1b

involved in the essential process for cell elongation in Escherichia coli. *Molec. Gen. Genet.*

200, 272–277 (1985).

4. Suzuki, H., Nishimura, Y. & Hirota, Y. On the process of cellular division in Escherichia coli: a

series of mutants of E. coli altered in the penicillin-binding proteins. *Proceedings of the*

National Academy of Sciences **75**, 664–668 (1978).

5. Yousif, S. Y., Broome-Smith, J. K. & Spratt, B. G. Lysis of Escherichia coli by beta-lactam

antibiotics: deletion analysis of the role of penicillin-binding proteins 1A and 1B. *J Gen*

Microbiol **131**, 2839–2845 (1985).

6. Schiffer, G. & Höltje, J.-V. Cloning and Characterization of PBP 1C, a Third Member of the

Multimodular Class A Penicillin-binding Proteins of *Escherichia coli* *. *Journal of Biological*

Chemistry **274**, 32031–32039 (1999).

7. Bush, K. & Bradford, P. A. β-Lactams and β-Lactamase Inhibitors: An Overview. *Cold Spring*

Harbor Perspectives in Medicine **6**, a025247 (2016).

8.　Mueller, E. A., Egan, A. J. F., Breukink, E., Vollmer, W. & Levin, P. A. Plasticity of Escherichia coli

cell wall metabolism promotes fitness and antibiotic resistance across environmental

conditions. *eLife* **8**, (2019).

9.　Centers for Disease Control and Prevention (U.S.). *Antibiotic Resistance Threats in the United*

States, 2019. https://stacks.cdc.gov/view/cdc/82532 (2019) doi:10.15620/cdc:82532.

10.　Murray, C. J. *et al.* Global burden of bacterial antimicrobial resistance in 2019: a systematic

analysis. *The Lancet* **399**, 629–655 (2022).

11.　Johnson, J. G., Spurbeck, R. R., Sandhu, S. K. & Matson, J. S. Genome sequence of Klebsiella

pneumoniae urinary tract isolate Top52. *Genome Announcements* **2**, 668–682 (2014).

12.　Sutaria, D. S. *et al.* First penicillin-binding protein occupancy patterns of - Lactams and -

lactamase inhibitors in klebsiella pneumoniae. *Antimicrobial Agents and Chemotherapy* **62**,

(2018).

13.　Bush, K. & Jacoby, G. A. Updated functional classification of β-lactamases. *Antimicrobial*

Agents and Chemotherapy **54**, 969–976 (2010).

14.　Kannan, G. *et al.* Rapid acid treatment of Escherichia coli: Transcriptomic response and

recovery. *BMC Microbiology* **8**, 37 (2008).

15.　Maurer, L. M., Yohannes, E., Bondurant, S. S., Radmacher, M. & Slonczewski, J. L. pH Regulates

Genes for Flagellar Motility, Catabolism, and Oxidative Stress in Escherichia coli K-12. *Journal*

of Bacteriology **187**, 304–319 (2005).

16.　Choi, U., Park, S. H., Lee, H. B., Son, J. E. & Lee, C.-R. Coordinated and Distinct Roles of

Peptidoglycan Carboxypeptidases DacC and DacA in Cell Growth and Shape Maintenance

under Stress Conditions. *Microbiology Spectrum* **11**, e00014-23 (2023).

17.　Peters, K. *et al.* The redundancy of peptidoglycan carboxypeptidases ensures robust cell shape

maintenance in Escherichia coli. *mBio* **7**, (2016).

18. Identification of a penicillin-binding protein 3 homolog, PBP3x, in Pseudomonas aeruginosa:

gene cloning and growth phase-dependent expression.

https://journals.asm.org/doi/epdf/10.1128/jb.179.5.1490-1496.1997

doi:10.1128/jb.179.5.1490-1496.1997.

19. Castanheira, S. *et al.* A Specialized Peptidoglycan Synthase Promotes Salmonella Cell Division

inside Host Cells. *mBio* **8**, 10.1128/mbio.01685-17 (2017).

20. Castanheira, S. & García-del Portillo, F. Evidence of two differentially regulated elongasomes in

Salmonella. *Commun Biol* **6**, 923 (2023).

21. Sub-Inhibitory Cefsulodin Sensitization of E. coli to β-lactams Is Mediated by PBP1b Inhibition |

PLOS One. https://journals.plos.org/plosone/article?id=10.1371/journal.pone.0048598.

22. Ramachandran, V. *et al.* Screen for Inhibitors of the Coupled Transglycosylase-Transpeptidase

of Peptidoglycan Biosynthesis in Escherichia coli. *Antimicrobial Agents and Chemotherapy* **50**,

1425–1432 (2006).

23. Curtis, N. A. C., Eisenstadt, R. L., Turner, K. A. & White, A. J. Inhibition of penicillin-binding

protein 3 of Escherichia coli K-12. Effects upon growth, viability and outer membrane barrier

function. *Journal of Antimicrobial Chemotherapy* **16**, 287–296 (1985).

24. Buijs, J., Dofferhoff, A. S. M., Mouton, J. W., Wagenvoort, J. H. T. & van der Meer, J. W. M.

Concentration-dependency of β-lactam-induced filament formation in Gram-negative

bacteria. *Clinical Microbiology and Infection* **14**, 344–349 (2008).

25. Paradis-Bleau, C. *et al.* Lipoprotein Cofactors Located in the Outer Membrane Activate

Bacterial Cell Wall Polymerases. *Cell* **143**, 1110–1120 (2010).

26. Nichols, R. J. *et al.* Phenotypic Landscape of a Bacterial Cell. *Cell* **144**, 143–156 (2011).

27. Boes, A., Olatunji, S., Breukink, E. & Terrak, M. Regulation of the Peptidoglycan Polymerase

Activity of PBP1b by Antagonist Actions of the Core Divisome Proteins FtsBLQ and FtsN. *mBio*

10, 10.1128/mbio.01912-18 (2019).

28. Bertsche, U. *et al.* Interaction between two murein (peptidoglycan) synthases, PBP3 and

PBP1B, in Escherichia coli. *Molecular microbiology* **61**, 675–690 (2006).

29. Castanheira, S. *et al.* An alternative penicillin-binding protein involved in *Salmonella* relapses

following ceftriaxone therapy. *eBioMedicine* **55**, 102771 (2020).

30. van der Ploeg, R. *et al.* Colocalization and interaction between elongasome and divisome

during a preparative cell division phase in scherichia coli. *Molecular Microbiology* **87**, 1074–

1087 (2013).

31. Castanheira, S., López-Escarpa, D., Paradela, A. & García-del Portillo, F. In Vivo Cross-Linking

Sheds Light on the Salmonella Divisome in Which PBP3 and PBP3 Compete for Occupancy.

Molecular Microbiology **n/a**,.

32. Hardy, C. D. Essential Paralogous Proteins as Potential Antibiotic Multitargets in Escherichia

coli. *Microbiology Spectrum* **10**, e02043-22 (2022).

33. Machen, T. E. & Paradiso, A. M. Regulation of intracellular pH in the stomach. *Annu Rev Physiol*

49, 19–33 (1987).

34. Fallingborg, J. Intraluminal pH of the human gastrointestinal tract. *Dan Med Bull* **46**, 183–196

(1999).

35. O'Hanlon, D. E., Come, R. A. & Moench, T. R. Vaginal pH measured in vivo: lactobacilli

determine pH and lactic acid concentration. *BMC Microbiol* **19**, 13 (2019).

36. García-nafría, J., Watson, J. F. & Greger, I. H. OPEN IVA cloning : A single-tube universal cloning

system exploiting bacterial In Vivo Assembly. *Nature Publishing Group* 1–12 (2016)

doi:10.1038/srep27459.

37. Bachman, M. A. *et al.* Genome-wide identification of Klebsiella pneumoniae fitness genes

during lung infection. *mBio* **6,** (2015).

38. Datsenko, K. A. & Wanner, B. L. One-step inactivation of chromosomal genes in Escherichia

coli K-12 using PCR products. *Proceedings of the National Academy of Sciences* **97**, 6640–6645

(2000).

39. Ducret, A., Quardokus, E. M. & Brun, Y. V. MicrobeJ, a tool for high throughput bacterial cell

detection and quantitative analysis. *Nature Microbiology* **1**, 1–7 (2016).

Figure Legends

Fig 1. *Klebsiella pneumoniae* exhibits acid-dependent resistance to Beta-lactams that is
independent of Beta-lactamase expression.

(A) Heatmap displaying the fold-change in minimal inhibitory change (MIC) for *K. pneumoniae*
TOP52 at indicated pH relative to neutral pH. Each cell represents the median fold-change value
from a minimum of three biological replicates performed in LB buffered to indicated pH with MMT
buffer. Specific Beta-lactam antibiotics along with their preferential PBP target are indicated on the
Y-axis. NS = non-specific (B) The fold-change in Meropenem (MER) or Cephalexin (CEX) MIC of *K.*
pnuemoniae strains ATCC 43816 and NHUT-K2044 grown at pH 4.8 compared to growth at pH 6.8.
Bars represent mean and error bars denote standard deviation. C-D) Representative micrographs
of terminal morphologies of *K. pneumoniae* TOP52 following growth for 20 hours in LB (pH 4.8 or
6.8) and treated with Meropenem (C) or Cephalexin (D) at indicated concentrations. Scale bar = 2
µm.

Fig 2. **Loss of either PBP1b or PBP3$_{PARA}$ results in differential impacts to beta-lactam**
resistance at low pH. (A-D) Fold change in MICs at pH 4.8 relative to pH 6.8 shown for generalist
(A), class A-targeting (B), PBP2-targeting (C), and PBP3-targeting (D) beta-lactams. Data are
graphed as median values with range. (E) Micrographs from time-lapse microscopy of WT (top) or
Δ*mrcB* (bottom) growing in the presence of CEX at low pH. Microscopy pads comprised of LB
buffered with MMT to pH 4.8 + 15.6 µg/mL CEX + 1.5 µM propidium iodide + 1% agarose. The CEX
concentration reflects the 1X MIC for the Δ*mrcB* mutant and 0.06X MIC for the WT strain.

Fig 3. *Klebsiella pneumoniae* possesses acid-responsive paralogous copies of the Class B PBPs. (A) Genomic regions containing the canonical PBP2, encoded by *mrdA*, and the paralogous gene *mrdA2,* which encodes PBP2$_{PARA}$ (B) Genomic regions containing the canonical PBP3, encoded by *ftsI,* and the paralogous gene *ftsI2,* which encodes PBP3$_{PARA}$. (C & D) Domain conservation between canonical and paralogous copies of PBP2 (C) and PBP3 (D). Domains are denoted as follows: Transmembrane (TM), Lipoprotein lipid attachment site (LP, PS51257) PBP-dimer (Pfam: PF03717), Transpeptidase domain (Pfam: PF00905). Location of motifs that are critical for enzyme catalysis are highlighted in purple.

Fig 4. Loss of PBP3$_{PARA}$ impairs division at low pH. (A) Representative micrographs of indicated strains following growth in LB buffered to pH 4.8 (top panels) or growth in LB buffered to pH 6.8 (bottom panels). For complementation studies with either empty vector (EV) or p*ftsI2,* the media was supplemented with 0.1 mM IPTG. Scale bar = 10 uM. (B-C) Quantification of cell length in (B). Each data point represents the mean of a biological replicate (>50 cells analyzed for each replicate). The bar represents the mean of three biological replicates, and the error bars represent the standard error of the mean. Statistical significance was determined by a one-way ANOVA with Tukey's correction for multiple comparisons with asterisks denoting significance as follows:** = $p<0.005$, * = $p<0.5$.

Fig 5. Loss of either PBP1b or PBP3$_{PARA}$ results in differential impacts to Beta-lactam resistance at low pH. (A-D) Fold change in MICs at pH 4.8 relative to pH 6.8 shown for generalist (A), class A-targeting (B),PBP2-targeting (C), and PBP3-targeting Beta-lactams. For comparison, data for MER and CEX are re-graphed from Fig 4. Data are presented as median values with range. (E) Micrographs from time-lapse microscopy of WT, Δ*ftsI2,* Δ*mrdA2,* and Δ*mrdA2* Δ*ftsI2* growing in the presence of MER at low pH. Microscopy pads comprised of LB buffered with MMT to pH 4.8 + 0.2 µg/mL MER + 1.5 µM propidium iodide + 1% agarose. The MER concentration reflects 1X MIC for WT, ~0.25X MIC for Δ*ftsI2,* ~2X MIC for Δ*mrdA2,* and ~0.5X MIC for Δ*mrdA2* Δ*ftsI2.*

Fig 1

A

B

C

D

Fig 2

Fig 3

A

B

C

D

Fig 4

Fig 5

578 **Table 1. Transcriptional response to pH of key beta-lactam targets in *K. pneumoniae* and *E.***
579 ***coli***

	Gene	Enzyme Encoded	Description	log$_2$FC (pH 4.8/6.8)
K. pneumoniae				
D1637_RS05825	*mrcA*	PBP1a	bifunctional glycosyl transferase/transpeptidase	0.60
D1637_RS12385	*mrcB*	PBP1b	bifunctional glycosyl transferase/transpeptidase	-0.39
D1637_RS01040	*pbpC*	PBP1c	peptidoglycan glycosyltransferase	-1.61
D1637_RS15400	*mrdA*	PBP2	canonical peptidoglycan DD-transpeptidase	0.84
D1637_RS12000	*ftsI*	PBP3	canonical peptidoglycan DD-transpeptidase	0.23
D1637_RS21590	*mrdA2*	PBP2$_{para}$	paralogous peptidoglycan DD-transpeptidase	3.41
D1637_RS24275	*ftsI2*	PBP3$_{para}$	paralogous peptidoglycan DD-transpeptidase	3.28
D1637_RS15385	*dacA*	PBP5	D-Ala-D-Ala-carboxypeptidase	1.20
D1637_RS25010	*dacD*	PBP6b	D-Ala-D-Ala carboxypeptidase	1.63
E. coli				
b3396	*mrcA*	PBP1a	bifunctional glycosyl transferase/transpeptidase	0.36
b0149	*mrcB*	PBP1b	bifunctional glycosyl transferase/transpeptidase	-0.23
b2519	*pbpC*	PBP1c	peptidoglycan glycosyltransferase	-0.43
b0635	*mrdA*	PBP2	canonical peptidoglycan DD-transpeptidase	0.47
b0084	*ftsI*	PBP3	canonical peptidoglycan DD-transpeptidase	-0.64
b0632	*dacA*	PBP5	D-alanyl-D-alanine carboxypeptidase	0.95
b2010	*dacD*	PBP6b	D-alanyl-D-alanine carboxypeptidase	2.63

580 *Shaded genes represent those that were differentially regulated by pH as determined by our cutoff
581 of |logFC| > 1 and $p < .05$. Blue Log$_2$FC values indicate that the gene was down-regulated during
582 growth at pH 4.8 versus pH 6.8. Magenta Log$_2$FC values indicate that the gene was up-regulated
583 during growth at pH 4.8 versus pH 6.8.