ADAR1 regulates alternative splicing through an RNA editing-independent

mechanism.

Eduardo A. Sagredo[1,2*], Victor Karlström[2], Alejandro Blanco[3], Paloma Moraga[4,5],

Matias Vergara[3], Aino I. Jarvelin[6], Neus Visa[1], Katherine Marcelain[7,8], Alfredo

Castello[6,9], and Ricardo Armisén[3,*].

1: Department of Molecular Biosciences, The Wenner-Gren Institute, Stockholm University, Svante Arrhenius

20C, Stockholm, 106 91, Sweden.

2: Science for Life Laboratory, Department of Microbiology, Tumor and Cell Biology, Karolinska Institutet, 171 65,

Solna, Sweden.

3: Centro de Genética y Genómica, Instituto de Ciencias e Innovación en Medicina, Facultad de Medicina Clínica

Alemana Universidad del Desarrollo, Santiago, Chile.

4: Biomedical Neuroscience Institute, Faculty of Medicine, University of Chile, Santiago, Chile.

5: FONDAP Center for Geroscience, Brain Health and Metabolism (GERO), Santiago, Chile.

6: Department of Biochemistry, University of Oxford, South Parks Road, Oxford OX1 3QU, UK.

7: Departamento de Oncología Básico Clínica, Facultad de Medicina, Universidad de Chile, Santiago, Chile.

8: Centro para la Prevención y el Control del Cáncer, Universidad de Chile, Santiago, Chile.

9: MRC-University of Glasgow Centre for Virus Research, The University of Glasgow, Glasgow, United Kingdom.

* To whom correspondence should be addressed: Eduardo A. Sagredo (eduardo.sagredo.campos@ki.se) &

Ricardo Armisen (+56 2 25785579, rarmisen@udd.cl)

Abstract

Adenosine-to-inosine (A-to-I) RNA editing and alternative splicing occur co-

transcriptionally and can regulate and influence each other. The RNA editing

enzymes ADAR1 and ADAR2 catalyze A-to-I RNA editing, where it has been shown

that expression changes in both RNA editing enzymes exert alternative splicing.

ADAR1 manipulation has a significant impact on alternative splicing. While many of

those changes are related to changes in their A-to-I RNA editing activity, we

speculate that ADAR1 may also influence splicing in an editing-independent manner.

In this work, the protein-protein interactome of ADAR1 revealed that ADAR1

interacts with spliceosome co-factors and auxiliary splicing regulators. We confirmed

that ADAR1 does not only influence splicing through editing but also throughout

editing-independent functions with even greater penetrance. We show that ADAR1

modulates the splicing of transcripts encoding splicing factors, with many of these

splice sites overlapping with known splicing changes induced by the splicing factor

ACIN1, suggesting indirect effects upon ADAR1 expression. In summary, we show

that ADAR1 can regulate splicing in an editing-independent manner, which likely

occurs by widespread alteration of the splicing factor isoform landscape.

Introduction

The adenosine deaminase acting on RNA (ADAR) enzymes catalyze adenosine-

to-inosine (A-to-I) RNA editing, one of the most prevalent RNA modifications in

mammals [1–3]. Inosine is recognized as guanosine by the cellular machinery, which

can cause profound effects on RNA structure and translational fidelity[4–6]. The ADAR

protein family consists of three members (ADAR1-3), two of which are catalytically

active (ADAR1 and ADAR2)[3]. All ADAR enzymes contain two to three double-

stranded RNA binding domains and a deamination domain[7]. ADAR1 is ubiquitously

expressed and has two isoforms, ADAR1p110, which is primarily nuclear and

nucleolar, while ADAR1p150 is induced by interferon and can shuttle between the

nucleus and the cytoplasm[8,9]. Changes in ADAR1 expression and activity are

associated with different diseases, including amyotrophic lateral sclerosis[10], systemic

lupus erythematosus[11], and several types of cancers[12]. Recent data suggest a broad

role of ADAR proteins in cell fate, showing that they can modify mRNA splicing

sites[13] and alternative polyadenylation sites[14] and affect miRNA

processing/maturation[15,16] and RNA stability[17–19] by altering canonical RNA-RNA and

RNA-proteins interactions[20]. Several studies have established how RNA editing may

influence alternative splicing by modifying splicing regulatory elements or by altering

dsRNA structures. These alterations may affect the affinity of the splicing machinery

and trans-acting factors for those regions, leading to alterations in splicing

efficiency[13,21–24]. One of the most well-characterized RNA editing-dependent splicing

events involves the glutamate receptor subunit 2 (GRIA2) transcript[25], where a single

recoding site of the GRIA2 pre-mRNA, in the amino acid 607 (known as Q/R site), is

required for the excision of the intron 11, regulating the processing and proper

mRNA cytoplasmic localization[26–28]. Moreover, ADAR2 can edit its transcript,

generating a new splice site resulting in a transcript variant that encodes a truncated

non-functional protein[27,29]. As virtually all mRNAs are spliced and a vast majority

undergo RNA editing co-transcriptionally, it has been proposed that these two

processes can mutually influence each other. Indeed, transcriptomic analyses

revealed that ADAR1 and ADAR2 expression alterations significantly affect exon usage and alternative splicing in animal and cellular models [13,21,30,31]. In counterpart, it has been shown that the pharmacological inhibition of the spliceosome influences ADAR editing function in those sites that compromise an intron or other RNA structures that can enhance RNA editing due to secondary structures[21,31].

ADAR1 knock-out mice die at early embryonic stages due to widespread apoptosis[32–34]. This phenotype can be rescued to a varying extent by adding back catalytically inactive ADAR1 mutants, indicating that A-to-I editing is not the only role of ADAR1 in RNA metabolism[35,36]. In agreement, it has been shown that ADAR can interact with other RBPs and influence their activity, further reinforcing the idea of an editing-independent function[20]. In the early days of genomic studies, Solomon et al. (2013)[13] intersected alternative splicing and RNA editing analyses upon ADAR1 knockdown, showing that RNA editing can regulate mRNA variant generation. However, the extensive changes in alternative splicing observed upon ADAR1 knockdown could not be completely connected to the editing function, and, indeed, only a few editing sites were detected in the vicinity of near regulatory splicing sequences. Similarly, Licht et al. (2016)[23] demonstrated that ADAR2 RNA-binding could influence splicing efficiency in an editing-independent manner for specific ADAR2 target minigene constructs. In agreement, Kapoor et al. (2020)[22] recently showed that Adar –/– mouse-derived tissues display extensive editing-independent splicing changes.

In this work, we investigated the role of ADAR1 in alternative splicing in four transgenic non-related cell lines, focusing on the connections between RNA editing and alternative splicing, showing that most splicing events upon ADAR1 knockdown or overexpression are not related to RNA editing events. We also established the ADAR1p110 protein-protein interactome, showing interactions with several key spliceosome co-factors. Furthermore, using two ADAR1p110 inducible cell lines, we found that most ADAR1-related splicing events do not involve RNA editing near the affected splicing junctions. Finally, we show that overexpression of ADAR1p110 alters the exon usage of splicing factor isoforms within the cell, suggesting that ADAR1 does exert indirect effects on splicing. These results highlight a complex

99　regulatory network of splicing mediated by ADAR1 that includes direct RNA editing
100　and indirect effects through the regulation of the splicing machinery.

Methods

Cell Culture

103　　HeLa and HEK293 cells were cultured under standard conditions at 37 °C in a
104　humidified incubator containing 5% CO_2. HeLa ADAR1p110 Flp-In T-REx and GFP
105　Flp-In T-REx cells were generated, subcloning the ADAR1p110 cds into the
106　pCDNA5-FRT-cGFP plasmid to further integrate the construct using the Flp-In
107　System following manufacturer instructions (Thermo Fisher Scientific).

Immunoblotting and co-immunoprecipitation

HeLa ADAR1p110 Flp-In T-REx cells were lysed (one 10 cm dish at 80%
(800.000 cells) confluency per immunoprecipitation experiment) in 1mL lysis buffer
(10 mM Tris/Cl pH 7.5; 150 mM NaCl; 0.5 mM EDTA; 0.5% Triton X) supplemented
with protease inhibitors (Halt, Thermo Fisher Scientific) and 250 units of Benzonase
(Sigma Aldrich), and placed on ice for 30 mins. At the same time, extensively shred
using a G15 syringe. Cell lysates were centrifuged at 20.000 rpm for 10 min at 4°C,
and the supernatant was transferred to a low binding pre-cooled 1.5 mL tube. 40 µL
pre-washed control beads (Pierce) (washed three times in cold lysis buffer
completed with DTT 1mM and proteinase inhibitors) slurry were added to the lysate
and incubated for 30 min at 4°C under rotation. Further, the lysate was spun down at
2500 rpm for 2 min, and the supernatant transferred to a new low binding pre-cooled
1.5 mL and incubated with 40µL of pre-washed GFP-trap beads (washed three times
in cold lysis buffer completed with DTT (Thermo Fisher Scientific) 1mM and protease
inhibitors) during four h under gentle rotation at 4°C, the complexed beads were
spun down at 4°C at 2500 rpm for 2 min and the supernatant was discarded to
further wash them three times with lysis buffer (supplemented with protease
inhibitors). Finally, the complexed beads were resuspended 40 µL. For co-
immunoprecipitation experiments, 10 µL of the complexed beads were boiled with 4x
NuPAGE Buffer (Thermo Fisher Scientific) and three µL 1M DTT following
manufacturer recommendations. Western blotting was performed as stated
previously in Sagredo et al. (2017)[37], and for mass spectrometry, the beads were

eluted by pH elution approach as indicated in the manufacturer's protocol (GFP-

trap_A; Chromotek). Mass spectrometry was performed as previously described in

Avolio et al. (2018)[38].

Reporter assay and transfections

Reporter assays were performed according to Armisén et al. (2011)[39]. Briefly,

HeLa cells were transfected with one µg of the CMV-LUC2CP/ARE plasmid

(Addgene #62857), 0.5 µg of Renilla plasmid (pRL-Renilla, Promega), and one µg of

the different ADAR1 isoform plasmids. ADAR1p110 deaminase inactive (DeaD) and

the ADAR1p110 dsRBD mutant (EAA) were cloned into a pCDNA3.1(+) backbone

derived from ADAR1p110 cDNA. Dual-Luciferase measurements were performed

using the Dual-Glo Luciferase system kit (Promega), using Cytation 3 Multi-Mode

Reader (BioTek Instruments). Renilla signal was used to normalize prior luciferase

comparisons between the different samples. Plasmid and siRNA transfections were

carried out as described in Sagredo et al. (2020)[18]. Briefly, the different plasmids

used in this study were transfected using Lipofectamine 3000, according to the

manufacturer's instructions (Thermo Fisher Scientific), using six-well plates at 80%

confluency. For siRNA control (Cell Signaling, 6568S) or ADAR1 (Thermo Fisher

Scientific, 119580), siRNAs were transfected at a final concentration of 20 nM and

incubated for 48 hours for further experiments. For plasmid transfection, 2.5 µg of

the different plasmids were transfected, and 16 h after transfection, the cells were

grown with new fresh media to be harvested 48 h after the transfection procedure.

RNA-seq experiments and workflow

RNA integrity and RNA sequencing were performed as described previously in

Sagredo et al. (2020)[18]. Fastq files were aligned using STAR (two-step mode)

against hg19, and transcripts were counted with HT-seq software. Differential exon

usage (DEU) analysis for the different cell lines and datasets included in this study

was performed using the DEXSeq[40] tool v1.36.0 (DEXseq_count.py script) to

perform the differential DEU analysis further as recommended by the software

developers. Features with an FDR<0.01 and Log2 > fold one were included in the

MDA-MB-231 analysis, and FDR<0.05 and Log2 fold > 0.75 features were included

in the ZR-751 analysis. Differential splicing events were called and analyzed using

rMATS-turbo v4.1.2 software[41,42], comparing control samples vs. ADAR1
overexpressing cells. Splice junction definitions from the UCSC goldenPath
repository were utilized for further differential splicing analysis. All generated data
was processed in R (3.6.1) or the Unix command line. Datasets from ZR-751 and
MDA-MB-231 cells are analyzed from Sagredo et al. (2020)[18] (SRA accession
numbers SRP200635 and SRP200634).

RNA editing and Splicing intersection analysis

Similarly to Kapoor et al. (2020)[22], we intersected the differential exon usage
analyzed using the DEXseq tool. We scanned the called editing sites in 50 nt, 200 nt,
500 nt, and 1000 bp windows on both sides of the exon/intron boundary. In addition,
reference sites present in RADAR and DARNED databases were also included[43,44].
This analysis was performed for significant differential exons, included/excluded in
the ZR-751 and MDA-MD-231 cells. DARNED and RADAR events were annotated
and intersected with the spliceosome iCLIP from Briese et al. (2019)[45] E-MTAB-
8182, using bedtools intersect v2.26.0, and the same window scanning strategy was
used to intersect both datasets.

rMATS splicing events were called in default mode (--b1 {bams_controls} --b2
{bams_oe} --gtf hg19.ensGene.gtf -t paired --readLength 100 --nthread 12 --od
./output_files/ --tmp ./output_files/tmp --libType fr-firststrand). For further analysis,
the files counted only the reads that used cross-splice junctions. The splicing events
with less than 20 reads between the sum of counts with and without in each
replicate, including the exon/intron/splice event, were removed. The Remaining
events were filtered by FDR ≤ 0.05, and The remaining events were filtered by FDR
≤ 0.05, with a difference between control and OE groups of | IncLevelDifference | ≥
0.1. Where IncLevelDifference is the difference of PSI (Percent Spliced In) values of
Control vs OE treatment groups. The filtered set of splicing events was then
annotated with external_gene_name using the biomaRt package v2.54.0 from
Bioconductor. For those significant splicing changes, we selected a set of windows
involving the splicing event. The window considered 50 bp from the splice junction
into the body of the exons and 200 bp into the introns of the event, except for RI
events, where 50 bp was considered for both directions of the splice junctions. This
set of windows was created for each significant splicing event and intersected with

the set of edited sites. The edited sites were filtered by total depth ≥ 5 (considering
all samples, 6 per cell type) and kept those observed at least in 2 replicates or
observed in RADAR and/or DARNED databases[43,44]. Then, both sets, splicing
events and edited sites, were intersected with bedtools v2.26.0 using bedtools
intersect. Also, the splicing events were annotated with RADAR+DARNED, obtained
from REDIportal [46] using the same "windows set" strategy to account for known
editions intersecting the splice site windows. If any splicing event intersects with the
sample's editions or a known edition from RADAR+DARNED, the event is tagged as
Editing Dependent; otherwise, it is tagged as Editing-Independent. Finally, rMATS
significant changes were additionally annotated and intersected with the ADAR1
iCLIP data from Chen et al. (2015) (GSE63709) [16] and Bahn et al. (2015)[14]
(GSE55363) and the irCLASH data from Song et al. (2020)[47](GSE136327).

Quantitative RT-PCR

Total RNA was extracted using the Mammalian Total RNA Isolation Kit following
the manufacturer's instructions (SIGMA-Aldrich). This includes Turbo DNAse
(Thermo Fisher Scientific) treatment following manufacturer instructions. 500 ng of
RNA was reverse-transcribed using an AffinityScript qRT-PCR cDNA Synthesis Kit
(Agilent Technologies Inc.) and diluted 4 times. qRT-PCR was performed using
specific primers (for further details, please check Supplementary Material 1) and the
KAPA SYBR Green master mix (Roche). The reactions were performed using the
MIC qPCR machine using the following thermal profile: 95 °C for 15 seconds, 58 °C
for 15 seconds, and 72 °C for 15 seconds at 40 cycles, including no template
controls. Expression values were calculated using the ΔΔCt method and expressed
as the fold of change relative to control samples. ACTNB was used as a
housekeeping gene. In addition, RESS-qPCR for AZIN1 and MDM2 targets was
performed as described in Crews et al. (2015)[48]. Primers were used to validate DEU
data, targeting one specific exon and normalizing it against GAPDH. For rMATS
validations, two primers were used for splicing change measurements, amplifying the
affected exon and a constant region of the targeted transcript.

Site level A to G(I) editing comparisons in HeLa and HEK293 FLP T-REx cells

Variant calling files, annotations, and filtering were performed as described
previously in Sagredo et al. (2020)[18] and Widmark et al. (2022)[49]. Only addressable
A to G(I) variants were used, Fisher's exact test was used to compare the
treatments, and a confidence level of 0.05 was used as a cut-off for further analysis.
All the data generated was processed in R (3.6.1).

Gene ontology and pathway enrichment analysis

To determine the relationship between ADAR1 and splicing regulation, a gene
ontology analysis was carried out using Cytoscape[50] v3.7.1 software and ClueGO[51]
v2.5.4 plugin or ShinyGO[52] 0.76.3 web application. For ClueGO analysis, a gene list
was submitted on this software using the Biological Process database (v06/01/2022)
for further comparisons. Only statistically significant groups were displayed, using a
Bonferroni step-down multiple comparison post-hoc test. A corrected $p<0.05$ was
considered statistically significant. For the protein-protein enrichment analysis, the
STRING[53] database webpage was used, using the complete list (Uniprot IDs) of
significant targets (FDR<0.05, compared to GFP samples).

Results

Editing of splicing motifs does not correlate with alterations in exon usage
upon ADAR1 expression.

To investigate ADAR's involvement in splicing regulation, we performed a
differential exon usage analysis datasets generated upon knockdown of ADAR1 by
either shRNA (ZR-751 ADAR1 knockdown cells) or overexpression by adenoviral
transduction (MDA-MB-231 cells overexpressing ADAR1p110). Our analysis
revealed 960 significant exon usage changes in the ZR-751 ADAR1 knock-down
cells, compared to controls, and 1958 significant exon usage changes in the MDA-
MB-231 ADAR1p110 overexpressing cells, compared to GFP transduced cells, used
as control. These changes mapped to 300 and 805 transcripts at the gene level,
respectively (Figure 1.A). Those transcripts that exhibited differential exon usage
after the alteration of ADAR1 expression were enriched in pathways related to
apoptosis, cell mobility, and regulation of cell proliferation (Figure 1.B). Only 24
transcripts overlapped between the datasets, suggesting that ADAR1-dependent
splicing regulation strongly varies across cell lines. Interestingly, and despite the low

overlap at the gene level, both cell lines showed significant changes in exon usage in

transcripts involved in cancer progression, including DUSP6, AKT1, WNT9A, IL6ST,

and FGFR1 (Supplementary Table 1).

To elucidate how ADAR RNA editing is related to splicing regulation, we selected

all the differential exons present in both ADAR1 knockdown and ADAR1p110

overexpressing cells and determined if editing sites map to the vicinity of intron

boundaries in the compromised mRNAs. To do so, we used the unfiltered RNA

variant calling analysis to maximize the detection of any potential RNA editing site.

This data was intersected with the coordinates of compromised exons applying

windows ranging from 50 to 1000 base pairs (bp) in either direction. We also

considered all the editing sites annotated in the RADAR and DARNED databases to

include any previously described editing sites that may not have been identified in

the knockdown and overexpression cells (Figure 1.C). As expected, the overlap

between differentially used splice sites and edited bases increased as the window

size increased. Despite applying the bigger window (1000 bp), only ~30% of the

differentially used exons overlapped with editing sites. This lack of overlap was

observed when RADAR and DARNED editing sites were used (Figure 1.D). These

surprising results suggest that most alternative exon usage events are associated

with ADAR-independent editing, even when using very permissive parameters.

To reinforce these results, we compiled the spliceosome binding sites reported by

previous iCLIP-published datasets. We tested the binding footprints on the RNA

against RNA editing events collected above. The overlapping between the

spliceosome iCLIP and the editing databases RADAR and DANRED was only

3.33%, yielding 96896 editing sites from the 2909674 total sites in RADAR and

DANRED. However, the overlapping increased to 85.4% when a 1000 bp window

was used. Altogether, these complementary analyses revealed an apparent lack of

correlation between ADAR1-mediated RNA edits and alternative splicing events

(Figure 1.E).

The protein-protein interactome of ADAR1p110 includes many splicing factors.

Classical biochemistry assays have associated ADAR1 with RNA-binding

proteins involved in splicing, such as HNRNPL, NONO, and SFPQ[54]. To study all

the protein partners of ADAR1p110, we generated HeLa Flp-In Trex cells that express a GFP-tagged ADAR1p110 protein in a doxycycline-dependent manner (Supplementary Figure 1). ADARp110 was immunoprecipitated using GFP-trap beads under stringent conditions, and interactors were identified by label-free quantitative proteomics (Figure 2A). GFP-expressing control cells were used to exclude contaminants. The analysis revealed 432 significantly enriched in ADAR1p110-GFP. As immunoprecipitation was performed in native conditions, these proteins reflect direct and indirect protein-bridged interactors. Notably, 270 of the interactors are annotated as RBPs, 163 of which harbor classical RNA-binding domains (RBDs), and 107 are non-canonical[55,56] (Figure 2B and Supplementary Table 2). To focus on the most prominent (stoichiometric) interactors, we selected the proteins with a higher eluate signal (q50 quantile of the IBAQ signal). We performed a pathway enrichment analysis using Reactome and STRING databases. We found a significant enrichment of proteins involved in 'ribosome' (FDR: 3.41E-80), 'spliceosome' (FDR: 1.49E-38), 'RNA transport' (FDR: 0.0273), 'mRNA surveillance' (FDR: 4.21E-23) and 'mismatch repair' (FDR: 9.53E-17). (Table 2. C and Supplementary Table 3). We observed a striking enrichment of members of the Sm complex and U2 snRNP complex in ADARp110 interactome (SF3A3, SF3B2, SF3B3, SNRPB2, SNRPA1, SNRPD3, SNRPE and SNRPG)[57] (Supplementary Table 2). We also identified splicing regulators such as HNRNPL, DHX15, SFPQ, and STRBP[58]. We validated several of the observed interactions by ADAR1 immunoprecipitation followed by western blot showing that Sm antigen, the splicing factor SF3A3, and two different heteronuclear proteins, HNRNPL and HNRPA/B, reinforcing the significance of our results (Figure 2D). These results imply that ADAR1 may also regulate splicing through interaction with the spliceosome and several of its cofactors.

Deconvoluting the role of ADAR1p110 in splicing

Our results indicate an editing-independent function of ADAR1 in splicing, consistent with many splicing factors in ADAR's interactome. We induced the expression of ADAR1p110-GFP in HeLa and HEK293 Flp-in TREx cells and analyzed the exon and splicing changes after ADAR1p110 induction. Using DEXseq, we observed that 4207 and 4235 exons had decreased usage in HeLa and HEK293,

while 3751 and 2720 exons showed higher usage than GFP-induced control cells
(Fig 3A and Supplementary Table 4). To characterize the transcripts with changes in
exon usage, we compared the results in both lines at the transcript level, showing
1712 transcripts differentially spliced in an ADAR-dependent manner in both cell
lines. These genes are involved in 'RNA processing' (FDR: 1.4E-30), 'protein
localization' (Protein localization to organelle (FDR: 4.8E-24), and 'viral processing'
(FDR: 3.2E-22) GO terms, among others. These terms are highly enriched in RBPs
and splicing-related factors (Figure 3. B and Supplementary Table 5).

Further, we utilized rMATS to analyze splicing events, an event-based
software that circumvents the limitations of differential exon usage based on exonic
counts. Using this tool, we detected 1537 significant alternative splicing changes in
the HeLa cells and 2827 in the HEK293 cells upon ADAR1p110-GFP
overexpression, compared to GFP-induced cells (Figure 3C). Most of the changes
were related to exon skipping (SE), representing 65,7% and 61,4% in HeLa and
HEK293 cells, respectively, upon ADAR1p110 overexpression. The following most
abundant splicing events were mutually excluded exon changes (MXE) (13% and
16.1%, respectively), intron retention (RI) (7.9% and 8.9%, respectively), and
alternative 5' and 3' splice site usage (13.3% and 13.6%, respectively) (Figure 3D
and Supplementary Table 6 and Supplementary Table 7), indicating that
ADAR1p110 exerts alternative splicing changes on many transcripts across different
cell types. However, many of the observed transcripts that exhibited splicing
changes do not overlap between cell lines, suggesting that regulation of alternative
splicing by ADAR1-p110 depends on the cell in which it is expressed (Fig 3D).

GO analysis revealed that transcripts with intron retention changes in both
cell lines are enriched in RNA splicing and RNA processing (FDR: 0.042) and
Cytoplasmic translation (FDR:0.042). The transcripts that exhibited significant exon
skipping changes are enriched in 'protein lipidation,' 'organelle organization'
(Organelle assembly and microtubule cytoskeleton organization, FDR: 0.009), and
DNA damage repair related GO terms (i.e., 'mitotic cell cycle' FDR: 0.009).
Conversely, significant mutually exclusive exon changes were enriched in protein
catabolic-related processes ('pos. reg of cellular protein metabolic processes', FDR:
0.02) (Figure 3.E). Finally, we found that several splicing factors, such as THOC1,

ACIN1, TIA1, SRSF6, HNRNPD, DDX46, and U2AF1, exhibited significant alterations to their splicing patterns upon ADAR1p110 induction. Furthermore, genes with central roles in the cell cycle and DNA damage response, such as BRCA1, PAX6, MCM3, MKI67, and BCLAF1, also displayed differential splicing in both cell lines.

Our next goal was to validate the ADAR1p110-induced changes in alternative splicing, testing their dependence on RNA editing and the RNA-binding activity of ADAR1p110. To do so, we transfected HEK293 cells with an editing deficient ADAR1 mutant (ADAR1-DeaD) and an RNA-binding deficient mutant (ADAR1-EAA) (Supplementary Figure 2.A). We tested the alternative splicing of mRNAs altered in our dataset, including PTPR, HNRNPC, LIG1, NME, and SRSF1. Using splice-site-specific primers, we tested exon exclusion and inclusion. Our results revealed that ADAR1 WT and its mutants induce similar changes in splicing, which was in line with our sequencing analyses. These results suggest that ADAR1p110 can affect the splicing in an RNA-binding and activity-independent manner (Supplementary Figure 2).

Elucidating the editing dependence of ADAR1-mediated changes to differential splicing

To characterize the editing-independent role of ADAR1p110 in alternative splicing, we performed an A-to-G variant calling on the whole transcriptome of HeLa and HEK293 T-REx cells and intersecting those sites with the altered splicing junctions in our data. As expected, most of the detected editing sites (~90%) occurred in Alu-elements situated in non-coding genomic regions (Supplementary Table 4), and less than 0.3% of detected edited sites were found in coding regions (Figure 4A). In addition, we performed a GO analysis on the significant edited sites shared between HEK293 and HeLa cells at the gene level (1598 genes), showing enrichment for 'cell cycle' (FDR:0.0001), 'DNA replication' (FDR:0.002), 'Fanconi anemia pathway' (FDR: 0.001), 'mismatch repair' (FDR:0.04), and 'homologous recombination' (FDR:0.02) GO terms, among others (Figure 4. B and Supplementary Table 5).

We designed a workflow to intersect the differential splicing events with the RNA
editing variants in our data and those RNA editing sites included in different RNA
editing databases, considering the splice junction and the intronic region nearby of
any significant splicing change (Figure 4. C and Materials and Methods).
Interestingly, less than 5% of the splicing changes in our data intersected with RNA
editing variants called in the cell lines (Supplementary Figure 3. A and 3. B, right
panels), suggesting that most changes cannot directly relate to RNA editing. From
the different splicing changes, mutually exclusive exon (MXE) changes had the
highest number of edited sites in their sequence. In contrast, the 5' splice site, 3'
splice site, and intron retention changes had the lower frequency of RNA editing
sites in their coordinates (Figure 4D, Supplementary Table 6 and Supplementary
Table 7), where a little fraction intersected in RADAR, DARNED and/or Reditools
databases (Figure 4.D, Supplementary Table 7 and Supplementary Figure 3. A and
3.B). Further, we selected several transcripts that exhibited alternative splicing
changes in the absence of RNA editing for validation by qRT-PCR. Those targets are
related to DNA damage response and alternative splicing, including MCM3, RPS9,
MKI67, and SRSF3. Strikingly, we observed that overexpression of ADAR1-DeaD
mutant led to significant changes in the alternative splicing of these transcripts,
supporting that the alteration of their splicing is editing-independent (Figure 4E).

ADAR1p110 regulates and contributes to transcriptome variability affecting
mRNAs that encode for splicing cofactors.

Our results showed that many ADAR1-regulated alternative splicing appears to
be editing-independent significantly. To test this phenomenon, we co-transfected the
different ADAR1p110 mutants with a splicing reporter consisting of a luciferase split
with an intron. Active luciferase synthesis thus requires splicing. We observed a
notable decrease in luciferase signal upon ADAR1p110-DeaD and ADAR1p110-EAA
overexpression in HEK293 cells (Figure 5.A). These results agree with Kapoor et al.
(2020)[22] and suggest that non-catalytically active ADAR1p110 influences the splicing
of the reporter in both an editing-independent and RNA-binding-independent
manner. We had two hypotheses: 1) ADAR regulates splicing factor function through
direct protein-protein interactions by changing their RNA-binding properties or

stability; 2) the activity of ADAR on mRNAs induces changes in splicing that alter the
configuration of splicing factors, leading to differential effects.

One splicing co-factor affected by ADAR overexpression was ACIN1, a protein
part of the apoptosis and splicing-associated protein complex (ASAP) that interacts
with the exon junction complex (EJC)[59,60]. The transcript of ACIN1 consistently
showed editing-independent alternative splicing and expression changes in both our
HeLa and HEK293 ADAR1p110 overexpressing cells (Figure 5. C and
Supplementary Figure 2.B). Indeed, the overexpression of both DeaD and EAA
ADARp110 in HEK293 cells showed a significant change in ACIN1 isoform pattern,
both at mRNA and protein levels (Supplementary Figure 2.B and Supplementary
Figure 2.D), suggesting that ADAR1 could indirectly modify the splicing process by
affecting the splicing configuration, result that we expanded to other splicing factors
(Supplementary Figure 2.C). To evaluate if the second hypothesis is correct, we
intersected publicly available transcriptome data from studies investigating the effect
of alternative splicing after ACIN1 knockdown[60,61], with our data produced after
overexpression of ADAR1p110 (Figure 5.B). Interestingly, some targets were
shared, affecting the same mRNA splicing variants between the ACIN1 dataset and
our experiments. Further, we performed qRT-PCR in the same splicing regions
associated with ACIN1 using the DeaD ADAR1p110 variant. Notably, the DeaD
ADAR1p110 overexpression can partially reassemble the effects of ACIN1 (Figure
5.D). Together, these results reinforce the idea that ADAR1 can indirectly regulate
alternative splicing of specific transcripts through modulation of alternative splicing
on transcripts that code for splicing factors, revealing an intricate role of ADAR1 in
regulating alternative splicing and downstream phenotypes.

Discussion

In this work, we provided new evidence supporting the role of ADAR1p110 in
regulating alternative splicing. It has previously been suggested that ADAR1
interacts with the spliceosome and that ADAR1p110 can be detected in the same
fraction as U1 and U2 spliceosome complexes[62]. Previous reports have shown that

the editing function can be regulated by splicing co-factors such as SRSF9 and
DDX15[63,64] and that RNA immunoprecipitation experiments for HNRNP proteins
interact with Alu regions and other features strongly associated with ADAR1 binding
sites[54,65]. Moreover, some cofactors of the spliceosome, such as U2AF2, can be
detected near editing sites[66], suggesting a plausible idea that these factors could
interact with or displace ADAR1, modifying ADAR1 activity. Our ADAR1p110 protein-
protein interactome experiment showed that many ADAR1p110 interactors are
indeed RNA binding proteins involved in RNA metabolism and alternative splicing
processes, which agrees with other recent ADAR1 proteomic works[67,68],
strengthening the argument that ADAR1p110 could act as a splicing regulator.

We investigated the changes in alternative splicing transcriptome-wide following
manipulation of ADAR1 expression and found thousands of changes in each of the
four individual cell lines tested. We observed that few of the detected differential
splicing events are shared between the two cancer cell lines (ZR-751 and MDA-MB-
231), and a significant proportion of those changes present in the HEK293 and HeLa
ADARp110-GFP cell lines generated by us, where most of the observed splicing
events appear to be cell line specific. This suggests that the regulation imposed by
ADAR1 on alternative splicing depends on the particular subcellular environment,
confirming the data of previous reports[22]. Furthermore, our data indicates that more
than half of the observed differential exon usage in ZR-751 and MDA-MB-231
ADAR1 manipulated cells do not harbor an editing site nearby. Even when using a
screening window of 1000 bp and relaxed criteria for editing determination, we found
no correlation between splice sites and editing in agreement with previous work[69].

Upon ADAR1p110-DeaD overexpression, we noted significant changes in the
luciferase reporter plasmid, noting that the overexpression of ADAR1p110-EAA also
inhibited the luciferase reporter's splicing when compared to the WT protein. In
agreement with this, previous works[21,22] also investigated the action of an editing-
deficient and RNA-binding deficient ADAR1 enzyme and observed that expression of
an RNA-binding deficient *Adar1* protein exhibited a more substantial impact on the
splicing landscape when compared to the editing-deficient *Adar1* in mice, suggesting
RNA editing independent effects. In further support of the editing-independent
regulation of alternative splicing by ADAR1, we observed a minor overlap of known

edited sites when intersecting their positions with the published iCLIP data for the

spliceosome[45], suggesting an alternative way that the editing-independent functions

of ADAR1 are responsible for the majority of the observed differential splicing

following its dysregulation.

After overexpressing ADAR1p110-GFP in HEK293 and HeLa cells, we observed

that the differentially spliced transcripts were enriched for DNA repair, mRNA

surveillance pathways, Splicing/spliceosome, and other related GO terms.

Interestingly, many of the observed changes to differential splicing generated

transcripts annotated as coding for novel protein isoforms. This phenomenon was

previously demonstrated for the ADAR-regulated differential splicing of the CCD15

and RELL2 transcripts[24]. Although the extent of the ADAR1p110 mediated

perturbations to alternative splicing remains to be investigated, it may influence

mRNA decay or protein isoforms, favoring mRNA variants that are targeted for

degradation to mechanisms such as NMD or producing novel protein isoforms.

These results could also add an extra layer of complexity in the pathological contexts

linked to ADAR1, where this protein is significantly overexpressed and could

influence transcript stability and isoform generation, alterations implicated in many

disorders and types of cancers[69,70].

ADAR1p110-GFP overexpression affected the splicing and expression of ACIN1,

a core component of the ASAP complex involved in alternative splicing[59], nonsense-

mediated decay[61], apoptosis, and chromatin condensation[71]. After intersecting the

observed differential splicing data in our HEK293 and HeLa cells with differential

splicing data available for ACIN1 knockdown[61], we could determine that a certain

proportion of splicing events exhibited overlap between the different datasets,

showing the same type of differential splicing on sharing gene coordinates. This

suggests that the observed perturbations in splicing could be due to the effect that

ADAR1p110 exerts on ACIN1 and may not be due to any direct relationship between

ADAR1p110 on those transcripts, results that we further validated by qRT-PCR.

RNA editing occurs co-transcriptionally, and most sites are present in intronic

regions[30,72]; a critical challenge of these studies to correlate editing with alternative

splicing is enough read coverage to limit false positive events. As our RNA-seq data

is based on a poly-A enriched strategy, which limits intronic regions, we

supplemented our editing data with the extensive data contained in Reditools,
RADAR, and DARNED databases to increase the sensitivity with capturing splicing
events that could be associated with RNA editing. However, even with very relaxed
parameters (1000 bp) in each direction from the differentially used exons, less than
30% contained edited sites near splicing junctions. Seeing that ADAR1p110 also
affects the splicing of multiple other splicing factors and interacts with them at a
protein-protein level, the full extent of ADAR1's involvement in alternative splicing
should be investigated systematically to determine these effects in cis or trans.

Figure Legends.

Figure 1. ADAR1 manipulation produces significant splicing changes not
entirely associated with their A to I activity. A) Differential Exon Usage (DEU)
after ADAR1 manipulation in ZR-751 and MDA-MB-231 cells. DEU analysis
comparing MDA-MB-231 ADAR1p110 overexpressing versus GFP transduced cells
(upper panel). ZR-751 SHC versus ZR-751 ADAR1 knockdown DEU analysis (lower
panel). B) Gene Ontology (Biological process) enrichment analysis for those
transcripts with significant DEU in MDA-MB-231 cells (Upper panel) and ZR-751
cells (Lower panel). Showing the significant terms based on gene number and Fold
enrichment (FDR). C) Diagram to identify RNA editing events associated with the
DEU changes, using the variant calling files, RADAR and DANRED and positional
information from Briese et al (2019)[45], spliceosome CLIP data, using different
spanning windows (50 bp to 1 Kbp) to identify possible A to G (I) variants associated
to DEU changes found in the MDA-MB-231 and ZR-751 cells, after ADAR1
manipulation. D) Number of DEU with addressable editing sites in the compromised
exon position with an extended boundary of 100 bp, 400, 1000, or 2000 bp, using
their variant calling information (left panel), RADAR, and DANRED databases (right
panel) in ZR-751 cells (upper) and MDA-MB-231 (lower) cells. E) Number of editing
sites present in RADAR and DANRED databases and their association with the
spliceosome CLIP positions based on Briese et al. (2019)[45] and their intersection
using an extended boundary of 0 bp, 50 bp, 200 bp, 500 bp, or 1000 bp.

Figure 2. HR-MS reveals that ADAR1p110 interacts with splicing machinery.
A) Model of work for HR-MS to capture ADAR1-p110 protein interactors. B) Volcano
plots showing significant ADAR protein-protein interactors, highlighting significant

(pink) and known RNA binding proteins (RBP) (blue) interactors. Dashed lines depict
significant -Log10 p-value (FDR) (y-axis> 1.75) and Log2 Fold change over GFP (x-
axis> 1). C) Reactome/STRING pathway enrichment for Q1 and Q2 IBAQ ranked
ADAR1p110 interaction partners. D) co-IP validations in HeLa cells overexpressing
GFP or ADAR1p110, showing inputs and IP elutions for Sm Antigen, SF3A3,
HNRPL, and HNRPAB.

Figure 3. ADAR overexpression on HeLa and HEK293 models produces
significant splicing changes. A) Differential exon usage counts (DEU) on HEK293
(left) and HeLa ADAR1p110 overexpressing cells, showing those exons significantly
overrepresented (included) in red and those excluded after ADAR1p110
overexpression (in green). B) The Venn diagram shows the overlap -at transcript
level- between those significant DEU changes in HEK293 and HeLa cells (upper),
showing the significant biological processes associated with those shared transcripts
with significant DEU in both cell lines. C) Pie charts showing the significant splicing
change types after rMATS analysis in HEK293 ADAR1p110 (left) and HeLa
ADAR1p110 (right) overexpressing cells. D) Venn diagrams showing the overlap in
the different splicing event changes between HEK293 and HeLa ADAR1p110
overexpressing cells. E) Gene enrichment analysis (Biological process) for those
splicing events shared between HEK293 and HeLa ADAR1p110 cells.

Figure 4. ADAR1p110 overexpression on HeLa and HEK293 cells produces
significant splicing changes that are not entirely associated with their A to I
activity. A) RNA editing characterization in HeLa and HEK293 cells after performing
variant calling, showing the number of editing sites, their distribution on Alu regions
and across the mRNA features b) Differentially RNA editing analysis for those
shared targets significantly edited in HeLa ADAR1p110 and HEK293 ADAR1p110,
compared to GFP overexpressing cells, showing significant Reactome gene
ontologies. C) Workflow to identify and recall splicing variants that are affected by
editing events and those recalled as editing independent events, briefly rMATS
significant sites were used to intersect further the variant calling files and
RADAR+DANRED databases, for the intersections of two regions that compromise
250 pb were used (for further details, see material and methods). D) Identification of
RNA editing-dependent and RNA-independent events on HeLa and HEK293 cells,

showing the different splicing change types on the data and the number of events (in blue) that can be described as editing-dependent. E) qRT-PCR validation for rMATS splicing events recalled as editing independent events after overexpressing HEK293 cells with an empty vector, ADAR1p110, and the DeaD ADAR1p110 plasmid. In addition, sashimi plots are shown for those targets that qRT-PCR validated. Two-tailed student's T-test was used to calculate differences between samples in E (n=3) (*=P<0.05, **=P<0.01, ***=P<0.001, ****=P<0.0001).

Figure 5. ADAR manipulation produces essential changes in splicing activity and the alternative splicing of other splicing factors. A) Splicing reporter activity after ADAR1 mutant overexpression. Briefly, a split Luciferase plasmid was co-transfected with the different ADAR1p110 mutants, and data was normalized using a renilla plasmid. B) Intersection between splicing events produced by ACIN1, after ACIN1 knockdown and the significant splicing events found in either HeLa or HEK293 ADAR1p110 overexpressing cells, displaying the intersection for all the different splicing event types. C) ADAR overexpression leads to a significant change in the splicing pattern of ACIN1 independent of the editing activity. D) qRT-PCR measuring ACIN1 splicing events after ADAR1 DeaD overexpression, showing significant changes in the splicing pattern of previously validated ACIN1 targets. Two-tailed T-test was used to calculate significant differences between samples (*=P<0.05, **=P<0.01, ***=P<0.001, ****=P<0.0001) in A, C, and D (n=3).

Supplementary Figure 1. Characterization of the ADAR1p110 Flp-In TREx system. A) schema of Flp-In TREx ADAR1p110 GFP doxycycline-inducible system. B) GFP fluorescence intensity signal produced in HeLa ADAR1p110-GFP cells after doxycycline induction, recorded during 24h. C) mRNA levels of ADAR1 after doxycycline induction on HEK and HeLa cells. D) RESS-qPCR for RNA editing targets (AZIN1 and MDM2) on HeLa and HEK293 ADAR1p110 after and before doxycycline induction. D) Western blot showing ADAR1p110-GFP after and before doxycycline induction, probing against GFP (upper panel); and confocal imaging showing ADAR1p110 localization, showing nuclear staining (DAPI) and ADAR1p110-GFP (FITC) (63X oil magnification) Two-tailed student's T-test was used to calculate differences between samples in C (n=3) (**=P<0.01, ***=P<0.001, ns: non-significant).

Supplementary Figure 2. ADAR1 manipulation produces significant exon usage changes on transcripts. A) ADAR mutants used for the different transfections, showing the different mutations present across ADAR1p110 domains. B) Sashimi plots for ACIN1 in HeLa and HEK293 overexpressing ADAR1p110 and GFP, showing ACIN1(L) and ACIN1(S) variants, with the corresponding affected exons. C) qRT-PCR validations for differential exon usage not related to ADAR1 editing activity, showing significant changes after DeaD and EAA overexpression in HEK293 cells for PTPRs, HNRNPC, LIG1, NME, and SRSF1. D) Representative Western blot showing ACIN1 isoforms after ADAR1 knockdown (siRNA), ADAR1p110 overexpression, and ADAR1 DeaD overexpression (left) in HEK293 cells, with their respective quantification for those bands associated with the larger isoform (200 KDa, 150 KDa, and 90 KDa) (right). A two-tailed T-test was used to calculate significant differences between samples (ns= non-significant, *=P<0.05, **=P<0.01, ***=P<0.001) in B and C (n=3).

Supplementary Table 1. Gene enrichment for those common transcripts affected after ADAR1 manipulation in MDA-MB-231 and ZR-751 cells.

Supplementary Table 2. ADAR1p110-GFP protein-protein interactome hits, showing the different proteins associated with ADAR1p110 compared to GFP.

Supplementary Table 3. STRING and Reactome enrichment analysis for ADAR1p110-GFP protein-protein interactome.

Supplementary Table 4. Differential exon usage changes in HeLa and HEK293 cells after ADAR1p110 induction.

Supplementary Table 5. GO (BP) enrichment analysis for those transcripts with DEU changes in HeLa and HEK293 cells after ADAR1p110 induction.

Supplementary Table 6. rMATS detected changes in HeLa cells after ADAR1p110 induction, annotated using RADAR and DARNED, Variant calling, spliceosome iCLIP, and ADAR1 iCLIP available data.

Supplementary Table 7. rMATS detected changes in HEK293 cells after
ADAR1p110 induction, annotated using RADAR and DARNED, Variant calling,
spliceosome iCLIP, and ADAR1 iCLIP available data.

Data availability

The data associated with this article are available in its online supplementary
material. Raw data will be made available on request.

Acknowledgments

We thank Dr. Albin Widmark and Dr. Yanara A. Bernal for their feedback on the
project and critical reading of the article. We acknowledge support from the National
Genomics Infrastructure in Stockholm funded by Science for Life Laboratory, the
Knut and Alice Wallenberg Foundation and the Swedish Research Council, and the
Swedish National Infrastructure for Computing/Uppsala Multidisciplinary Center for
Advanced Computational Science for assistance with massively parallel sequencing
and access to the UPPMAX computational infrastructure.

Author contributions:

Conceptualization, E.A.S., R.A. and A.B.; investigation, E.A.S., V.K., P.M., M.V.,
A.B., A.I.J. and M.V., visualization, E.A.S., A.B., writing-review and editing, E.A.S.,
V.K., K.M., A.C. and R.A. supervision, E.A.S., N.V., R.A., project administration and
funding acquisition R.A., N.V. and E.A.S. All authors have read and agreed to the
published version of the manuscript.

Funding

Swedish Research Council 2019-03853 to N.V.; Carl Trygger's Foundation (to
N.V.); V.K. was supported by the Department of Molecular Biosciences, Wenner-
Gren Institute, Stockholm University; E.A.S. was supported by SFO funding from the
Faculty of Science at the Stockholm University, O.E. and Edla Johanssons
Foundation and Royal Physiographic Society in Lund. Fondecyt regular 1221436
and 1151446 to R.A, Anillo en Ciencia y Tecnología ACT210079 to R.A.

Conflict of interest statement.

RA declares honoraria for conferences, advisory boards, and educational activities from Roche, grants, and support for scientific research from Illumina, Pfizer, Roche & Thermo Fisher Scientific, and honoraria for conferences from Thermo Fisher Scientific, Janssen & Tecnofarma. The funders had no role in the study's design, in the collection, analyses, or interpretation of data, in the writing of the manuscript, or in the decision to publish the results. The other authors declare that they have no competing interests.

References

1. Nishikura, K. Functions and regulation of RNA editing by ADAR deaminases. *Annual Review of Biochemistry* **79**, 321–349 (2010).

2. Tan, M. H. *et al.* Dynamic landscape and regulation of RNA editing in mammals. *Nature* **550**, 249–254 (2017).

3. Bass, B. L. RNA editing by adenosine deaminases that act on RNA. *Annual Review of Biochemistry* **71**, 817–846 (2002).

4. Keegan, L. P., Gallo, A. & O'Connell, M. A. The many roles of an RNA editor. *Nature Reviews Genetics* **2**, 869–878 (2001).

5. Ramaswami, G. *et al.* Identifying RNA editing sites using RNA sequencing data alone. *Nature Methods* **10**, 128–132 (2013).

6. Peng, Z. *et al.* Comprehensive analysis of RNA-Seq data reveals extensive RNA editing in a human transcriptome. *Nature Biotechnology* **30**, 253–260 (2012).

7. Savva, Y. A., Rieder, L. E. & Reenan, R. A. The ADAR protein family. *Genome biology* vol. 13 252 at https://doi.org/10.1186/gb-2012-13-12-252 (2012).

8. Eckmann, C. R., Neunteufl, A., Pfaffstetter, L. & Jantsch, M. F. The human but not the Xenopus RNA-editing enzyme ADAR1 has an atypical nuclear localization signal and displays the characteristics of a shuttling protein. *Molecular Biology of the Cell* (2001) doi:10.1091/mbc.12.7.1911.

9. Poulsen, H., Nilsson, J., Damgaard, C. K., Egebjerg, J. & Kjems, J. CRM1 Mediates the Export of ADAR1 through a Nuclear Export Signal within the Z-DNA Binding Domain. *Molecular and Cellular Biology* (2001) doi:10.1128/mcb.21.22.7862-7871.2001.

10. Hideyama, T. *et al.* Profound downregulation of the RNA editing enzyme ADAR2 in ALS spinal motor neurons. *Neurobiology of Disease* (2012) doi:10.1016/j.nbd.2011.12.033.

11. Laxminarayana, D., O'Rourke, K. S., Maas, S. & Olorenshaw, I. Altered editing
in RNA editing adenosine deaminase ADAR2 gene transcripts of systemic
lupus erythematosus T lymphocytes. *Immunology* **121**, 359–369 (2007).

12. Han, L. *et al.* The Genomic Landscape and Clinical Relevance of A-to-I RNA
Editing in Human Cancers. *Cancer Cell* **28**, 515–528 (2015).

13. Solomon, O. *et al.* Global regulation of alternative splicing by adenosine
deaminase acting on RNA (ADAR). *Rna* **19**, 591–604 (2013).

14. Bahn, J. H. *et al.* Genomic analysis of ADAR1 binding and its involvement in
multiple RNA processing pathways. *Nature Communications* **6**, 6355 (2015).

15. Ota, H. *et al.* ADAR1 forms a complex with dicer to promote MicroRNA
processing and RNA-induced gene silencing. *Cell* **153**, 575–589 (2013).

16. Chen, T. *et al.* ADAR1 is required for differentiation and neural induction by
regulating microRNA processing in a catalytically independent manner. *Cell
Research* **25**, 459–476 (2015).

17. Wang, I. X. *et al.* ADAR Regulates RNA Editing, Transcript Stability, and Gene
Expression. *Cell Reports* **5**, 849–860 (2013).

18. Sagredo, E. A. *et al.* ADAR1 Transcriptome editing promotes breast cancer
progression through the regulation of cell cycle and DNA damage response.
Biochimica et Biophysica Acta - Molecular Cell Research **1867**, (2020).

19. Karlström, V. *et al.* ADAR3 modulates neuronal differentiation and regulates
mRNA stability and translation. *Nucleic Acids Research* gkae753 (2024)
doi:10.1093/nar/gkae753.

20. Licht, K. & Jantsch, M. F. The Other Face of an Editor: ADAR1 Functions in
Editing-Independent Ways. *BioEssays* vol. 39 at
https://doi.org/10.1002/bies.201700129 (2017).

21. Licht, K., Kapoor, U., Mayrhofer, E. & Jantsch, M. F. Adenosine to Inosine
editing frequency controlled by splicing efficiency. *Nucleic Acids Research* **44**,
6398–6408 (2016).

22. Kapoor, U. *et al.* ADAR-deficiency perturbs the global splicing landscape in
mouse tissues. *Genome Research* **30**, 1107–1118 (2020).

23. Licht, K. *et al.* A high resolution A-to-I editing map in the mouse identifies
editing events controlled by pre-mRNA splicing. *Genome Research* **29**, 1453–
1463 (2019).

24. Tang, S. J. *et al.* Cis- and trans-regulations of pre-mRNA splicing by RNA
editing enzymes influence cancer development. *Nature Communications* **11**,
(2020).

25. Egebjerg, J., Kukekov, V. & Heinemann, S. F. Intron sequence directs RNA
editing of the glutamate receptor subunit GluR2 coding sequence. *Proceedings
of the National Academy of Sciences of the United States of America* **91**,
10270–10274 (1994).

26. Schoft, V. K., Schopoff, S. & Jantsch, M. F. Regulation of glutamate receptor B

pre-mRNA splicing by RNA editing. *Nucleic Acids Research* **35**, 3723–3732
(2007).

27.   Kawahara, Y., Ito, K., Ito, M., Tsuji, S. & Kwak, S. Novel splice variants of
human ADAR2 mRNA: Skipping of the exon encoding the dsRNA-binding
domains, and multiple C-terminal splice sites. *Gene* (2005)
doi:10.1016/j.gene.2005.07.028.

28.   Rosenthal, J. J. C. & Seeburg, P. H. A-to-I RNA Editing: Effects on Proteins
Key to Neural Excitability. *Neuron* at
https://doi.org/10.1016/j.neuron.2012.04.010 (2012).

29.   Rueter, S. M., Dawson, T. R. & Emeson, R. B. Regulation of alternative
splicing by RNA editing. *Nature* **399**, 75–80 (1999).

30.   Hsiao, Y. H. E. *et al.* RNA editing in nascent RNA affects pre-mRNA splicing.
Genome Research **28**, 812–823 (2018).

31.   Nilsen, T. W. & Graveley, B. R. Expansion of the eukaryotic proteome by
alternative splicing. *Nature* vol. 463 457–463 at
https://doi.org/10.1038/nature08909 (2010).

32.   Hartner, J. C. *et al.* Liver Disintegration in the Mouse Embryo Caused by
Deficiency in the RNA-editing Enzyme ADAR1. *Journal of Biological Chemistry*
279, 4894–4902 (2004).

33.   Hartner, J. C., Walkley, C. R., Lu, J. & Orkin, S. H. ADAR1 is essential for the
maintenance of hematopoiesis and suppression of interferon signaling. *Nature
Immunology* **10**, 109–115 (2009).

34.   XuFeng, R. *et al.* ADAR1 is required for hematopoietic progenitor cell survival
via RNA editing. *Proceedings of the National Academy of Sciences of the
United States of America* **106**, 17763–17768 (2009).

35.   Bajad, P. *et al.* An internal deletion of ADAR rescued by MAVS deficiency
leads to a minute phenotype. *Nucleic Acids Research* **48**, 3286–3303 (2020).

36.   Heraud-Farlow, J. E. *et al.* Protein recoding by ADAR1-mediated RNA editing
is not essential for normal development and homeostasis. *Genome Biology* **18**,
(2017).

37.   Sagredo, A. I. *et al.* TRPM4 regulates Akt/GSK3-β activity and enhances β-
catenin signaling and cell proliferation in prostate cancer cells. *Molecular
Oncology* **12**, 151–165 (2018).

38.   Avolio, R. *et al.* Protein Syndesmos is a novel RNA-binding protein that
regulates primary cilia formation. *Nucleic Acids Research* **46**, 12067–12086
(2018).

39.   Armisén, R. *et al.* TRPM4 enhances cell proliferation through up-regulation of
the β-catenin signaling pathway. *Journal of Cellular Physiology* **226**, 103–109
(2011).

40.   Anders, S., Reyes, A. & Huber, W. Detecting differential usage of exons from
RNA-seq data. *Genome Research* **22**, 2008–2017 (2012).

41. Shen, S. *et al.* rMATS: Robust and flexible detection of differential alternative
splicing from replicate RNA-Seq data. *Proceedings of the National Academy of*
Sciences of the United States of America **111**, E5593–E5601 (2014).

42. Wang, Y. *et al.* rMATS-turbo: an efficient and flexible computational tool for
alternative splicing analysis of large-scale RNA-seq data. *Nature Protocols*
(2024) doi:10.1038/s41596-023-00944-2.

43. Ramaswami, G. & Li, J. B. RADAR: A rigorously annotated database of A-to-I
RNA editing. *Nucleic Acids Research* **42**, D109-13 (2014).

44. Kiran, A. & Baranov, P. V. DARNED: A DAtabase of RNa editing in humans.
Bioinformatics **26**, 1772–1776 (2010).

45. Briese, M. *et al.* A systems view of spliceosomal assembly and branchpoints
with iCLIP. *Nature Structural and Molecular Biology* **26**, 930–940 (2019).

46. Picardi, E. & Pesole, G. REDItools: High-throughput RNA editing detection
made easy. *Bioinformatics* (2013) doi:10.1093/bioinformatics/btt287.

47. Song, Y. *et al.* irCLASH reveals RNA substrates recognized by human ADARs.
Nature Structural and Molecular Biology **27**, 351–362 (2020).

48. Crews, L. A. *et al.* An RNA editing fingerprint of cancer stem cell
reprogramming. *Journal of Translational Medicine* **13**, 52 (2015).

49. Widmark, A. *et al.* ADAR1- and ADAR2-mediated regulation of maturation and
targeting of miR-376b to modulate GABA neurotransmitter catabolism. *Journal*
of Biological Chemistry **298**, (2022).

50. Shannon, P. *et al.* Cytoscape: A software Environment for integrated models of
biomolecular interaction networks. *Genome Research* **13**, 2498–2504 (2003).

51. Bindea, G. *et al.* ClueGO: A Cytoscape plug-in to decipher functionally
grouped gene ontology and pathway annotation networks. *Bioinformatics* **25**,
1091–1093 (2009).

52. Ge, S. X., Jung, D., Jung, D. & Yao, R. ShinyGO: A graphical gene-set
enrichment tool for animals and plants. *Bioinformatics* **36**, 2628–2629 (2020).

53. Szklarczyk, D. *et al.* The STRING database in 2021: Customizable protein-
protein networks, and functional characterization of user-uploaded
gene/measurement sets. *Nucleic Acids Research* **49**, D605–D612 (2021).

54. Quinones-Valdez, G. *et al.* Regulation of RNA editing by RNA-binding proteins
in human cells. *Communications Biology* **2**, (2019).

55. Castello, A. *et al.* Comprehensive Identification of RNA-Binding Domains in
Human Cells. *Molecular Cell* **63**, 696–710 (2016).

56. Beckmann, B. M. *et al.* The RNA-binding proteomes from yeast to man
harbour conserved enigmRBPs. *Nature Communications* (2015)
doi:10.1038/ncomms10127.

57. Zhan, X., Yan, C., Zhang, X., Lei, J. & Shi, Y. Structures of the human pre-
catalytic spliceosome and its precursor spliceosome. *Cell Research* **28**, 1129–
1140 (2018).

58. Cvitkovic, I. & Jurica, M. S. Spliceosome database: A tool for tracking
components of the spliceosome. *Nucleic Acids Research* **41**, (2013).

59. Murachelli, A. G., Ebert, J., Basquin, C., Le Hir, H. & Conti, E. The structure of
the ASAP core complex reveals the existence of a Pinin-containing PSAP
complex. *Nature Structural and Molecular Biology* **19**, 378–386 (2012).

60. Lin, Y. C., Lu, Y. H., Lee, Y. C., Hung, C. S. & Lin, J. C. Altered expressions
and splicing profiles of Acin1 transcripts differentially modulate brown
adipogenesis through an alternative splicing mechanism. *Biochimica et
Biophysica Acta - Gene Regulatory Mechanisms* **1863**, (2020).

61. Rodor, J., Pan, Q., Blencowe, B. J., Eyras, E. & Caceres, J. F. The RNA-
binding profile of Acinus, a peripheral component of the exon junction complex,
reveals its role in splicing regulation. *Rna* **22**, 1411–1426 (2016).

62. Raitskin, O., Cho, D. S. C., Sperling, J., Nishikura, K. & Sperling, R. RNA
editing activity is associated with splicing factors in lnRNP particles: The
nuclear pre-mRNA processing machinery. *Proceedings of the National
Academy of Sciences of the United States of America* (2001)
doi:10.1073/pnas.111153798.

63. Tariq, A. *et al.* RNA-interacting proteins act as site-specific repressors of
ADAR2-mediated RNA editing and fluctuate upon neuronal stimulation.
Nucleic Acids Research (2013) doi:10.1093/nar/gks1353.

64. Shanmugam, R. *et al.* SRSF9 selectively represses ADAR2-mediated editing
of brain-specific sites in primates. *Nucleic Acids Research* **46**, 7379–7395
(2018).

65. Hendrickson, D., Kelley, D. R., Tenen, D., Bernstein, B. & Rinn, J. L.
Widespread RNA binding by chromatin-associated proteins. *Genome Biology*
17, 28 (2016).

66. Freund, E. C. *et al.* Unbiased Identification of trans Regulators of ADAR and A-
to-I RNA Editing. *Cell Reports* **31**, (2020).

67. Orecchini, E. *et al.* ADAR1 restricts LINE-1 retrotransposition. *Nucleic Acids
Research* **45**, (2017).

68. Hong, H. Q. *et al.* Bidirectional regulation of adenosine-to-inosine (A-to-I) RNA
editing by DEAH box helicase 9 (DHX9) in cancer. *Nucleic Acids Research* **46**,
7953–7969 (2018).

69. Shen, H. *et al.* ADARs act as potent regulators of circular transcriptome in
cancer. *Nature Communications* (2022) doi:10.1038/s41467-022-29138-2.

70. Bonnal, S. C., López-Oreja, I. & Valcárcel, J. Roles and mechanisms of
alternative splicing in cancer — implications for care. *Nature Reviews Clinical
Oncology* vol. 17 457–474 at https://doi.org/10.1038/s41571-020-0350-x
(2020).

71. Sahara, S. *et al.* Acinus is a caspase-3-activated protein required for apoptotic
chromatin condensation. *Nature* (1999) doi:10.1038/43678.

72. Laurencikiene, J., Källman, A. M., Fong, N., Bentley, D. L. & Öhman, M. RNA

editing and alternative splicing: The importance of co-transcriptional
coordination. *EMBO Reports* (2006) doi:10.1038/sj.embor.7400621.

Figure 1

A

DEXseq in MDA-MB-231 ADAR1p110 OV cells

DEXseq in ZR-751 SHADAR cells

B

GO: Biological Process MDA-MB-231 ADAR1p110 OV cells

GO: Biological Process ZR-751 SHADAR cells

C

D

Intersections in ZR-751 SHADAR cells

Intersections in MDA-MB-231 ADAR1p110 OV cells

E

Figure 2

A

DOX+ 16hrs

Lysis
1% Triton X-100
Benzonase (+)

IP purification
and washes

Elution

Silver staining

Western blot
Co-IP

Mass spectrometry

ADAR1-GFP   Off targets   Agarose beads   GFP-Trap beads   GFP-Trap Coupled beads

B

- Log$_{10}$ (Pvalue)

Log$_2$ fold change

● Is an RBP   ● Not an RBP   ● Not significant

C

Pathway description	Observed counts	FDR
Ribosome	**68**	**3.41E-80**
Spliceosome	**42**	**1.49E-38**
RNA transport	19	1.28E-08
mRNA surveillance pathway	14	8.64E-08
Gap junction	8	0.00897
Systemic lupus erythematosus	8	0.0184
Protein export	4	0.0254
Mismatch repair	4	0.0273

Pathway description	Observed counts	FDR
Ribosome	25	4.21E-23
Spliceosome	20	9.53E-17

D

	Input		IP	
	GFP	ADAR	GFP	ADAR

30 KDa — Sm Antigen

50 KDa — SF3A3

45 KDa — HNRPL

40 KDa — HNRPAB

Figure 3

Figure 4

Figure 5.