Coupling codon and protein constraints decouples drivers of variant pathogenicity

Ruyi Chen[1], Nathan Palpant[2], Gabriel Foley[1], Mikael Bodén[1,*]

[1]School of Chemistry and Molecular Biosciences, The University of Queensland, Brisbane, Australia.
[2]Institute for Molecular Bioscience, The University of Queensland, Brisbane, Australia.
[*]To whom correspondence should be addressed: `m.boden@uq.edu.au`.

Abstract

Predicting the functional impact of genetic variants remains a fundamental challenge in genomics. While existing models excel at identifying protein-intrinsic defects, they often overlook the regulatory syntax embedded within coding sequences. Here, we employ single-sequence codon (CaLM) and protein (ESM-2) language models as orthogonal probes to dissect the drivers of variant pathogenicity. Benchmarking on ClinVar variants demonstrates that the two modalities contribute evenly to distinguish between pathogenic and benign changes. Crucially, evaluating the same models on ClinMAVE data elucidates a mechanistic dichotomy: loss-of-function variants are predominantly governed by residue-level features, whereas gain-of-function variants and dosage-sensitive genes tend to rely more heavily on information at the codon-level, e.g. regulating protein biogenesis. These results highlight that pathogenicity stems from both the "product" *and* the "process," and that a duality of language models offer means to decode the mechanisms underlying disease.

Keywords: Protein language models, Codon language models, Mutation effects, Bayesian optimization, Variant interpretation, Codon degeneracy

1 Introduction

Predicting the functional impact of genetic variants remains a fundamental challenge in genomics. To address this, deep learning models have been deployed to extract evolutionary constraints directly from biological sequences via "zero-shot learning" [1–3]. These models can score variants by estimating how much a mutation disrupts the probability of observing a token (e.g., an amino acid) in its native sequence context. This disruption can be quantified as a Log-Likelihood Ratio (LLR) between the mutant and wild-type (wt) state, and generates effect scores that correlate strongly with experimental fitness and clinical pathogenicity [1, 4].

Existing models are broadly categorized into three classes based on their training modalities: protein sequence [1, 5], structure [6, 7], and alignment-by-homology [2, 8]. While integrating evolutionary and structural signals improves predictive performance, these proteocentric models treat coding sequences primarily as generators of proteins, overlooking the regulatory syntax embedded in the genomic context [9–12]. Notably, Boshar and colleagues demonstrated that a joint genomic/proteomic model can outperform specialized protein-only models on specific tasks [13].

Capitalizing on this perspective, we consider the relationship between DNA and protein as analogous to distinct natural languages. For example, English and German may convey semantics that are largely similar yet subtly different in interpretation (Fig. 1A) [14]. The flow of genetic information from DNA to mRNA to proteins (where naively "translation" appears to be a matter of table look-up) illustrates how core meanings can transcend languages. We reasoned that large language models trained on sequences, expressed in distinct "languages," such as protein-coding DNA (cDNA) and their corresponding protein sequences, are likely to capture nuances derived from, and be specific to the cDNA *or* the protein super-sets, respectively (Fig. 1B). The integration of variant-level information from both cDNA and protein sequences is here hypothesised to provide *complementary* biological signals, thereby revealing pathogenic mechanisms that cannot be explained by protein defects alone.

Here, we employ a dual-probe framework to identify pathogenic drivers that escape proteocentric models but still shape clinical phenotypes. Specifically, we use a Codon Language Model (CLM; CaLM [15]) and a Protein Language Model (PLM; ESM-2 [3]) to independently calculate the LLRs of mutations at the codon and amino acid levels, and integrate these scores via Bayesian optimisation (Fig. 1E). Extensive validation on the *in vivo* ClinVar (Fig. 1C; [16]) and *in vitro* ClinMAVE (Fig. 1D; [17]) reveals a clear mechanistic dichotomy: Loss-of-Function (LoF) variants are predominantly governed by residue-level features, whereas Gain-of-Function (GoF) variants and dosage-sensitive genes tend to rely heavily on information at the codon-level. Together, these results highlight that pathogenicity emerges from both the "product" and the "process," effectively captured by a duality of large language models.

2 Results

2.1 Dual-modality model captures pathogenicity missed by single-modality models

To investigate whether and to what extent distinct biological constraints contribute to variant pathogenicity, we curated a comprehensive dataset of missense variants from ClinVar, comprising 137,350 variants (39,280 pathogenic and 98,070 benign) across 13,791 genes (Fig. 1C and Supplementary Table. S1).

We benchmarked individual CaLM and ESM-2 against their hybrid model to quantify the information gained by reconciling the signals in the two modalities. While single-modality models generate LLRs independently, we use Bayesian optimisation to determine weights to balance the influence the LLRs of each modality (see "Materials and methods – Missense effect scores"). Despite distinct training data and model architectures, we confirmed that CaLM and ESM-2 displayed compatible LLR profiles, with nearly identical standard deviations, both globally and across comparison groups (Supplementary Table S3). This suggests that both models capture concordant evolutionary landscapes, that can be combined directly as raw LLRs.

We employed 10-fold cross-validation to partitioning variants by gene, thereby controlling for ascertainment bias. To quantify the relative contribution of each modality, we employed Bayesian optimisation to determine the weights that best explain the clinical variants. This unbiased optimisation yielded a mean weight of 0.49 for CaLM across training folds (Fig. 2A), with fold-specific details provided in Supplementary Fig. S1 (see "Materials and methods - Model training and validation"). This near-even balance suggests, in general, that information from codons and amino acids jointly defines the pathogenicity landscape.

For each fold, we applied the weight optimized on the training set to generate predictions for the corresponding validation set. Aggregating these predictions yielded composite effect scores for all three models (Fig. 2B and C). Notably, while CaLM effectively differentiates benign from pathogenic variants, their discriminative power is marginally lower than that of the protein-based ESM-2. Integrating both modalities achieved an AUROC of 0.862, significantly outperforming both ESM-2 (0.831) and CaLM (0.822) (Fig. 2B; two-tailed paired t-test, $p < 0.001$). Furthermore, the hybrid model maintained this advantage in the Area Under the Precision-Recall Curve (AUPRC) (Supplementary Fig. S2; two-tailed paired t-test, $p < 0.001$), confirming that the weighted signal is robust to class imbalance and independent of the primary optimisation metric. The performance gain suggests that these biological signals are complementary rather than redundant.

2.2 Divergent contributions of codon and protein constraints to functional variant phenotypes

To further investigate the relative contributions of codon- and residue-level constraints to variant function prediction, we used the dual-modality model on the ClinMAVE dataset. Multiplexed Assays of Variant Effect (MAVE)-derived data decouples a variant's biophysical impact from its native genomic context. To interrogate whether

codon-level constraints influence the functional fitness landscapes captured at the protein level, we first stratified the ClinMAVE dataset into three distinct categories: 176,803 functionally normal, 74,038 LoF, and 2,237 GoF variants (Fig. 1D and Supplementary Table S2). We subsequently formulated two parallel comparative analyses: functionally normal vs. LoF, and functionally normal vs. GoF. For each comparison, LLRs were computed using CaLM and ESM-2 independently, followed by a search for the optimal weights of their hybrid.

Given the limited number of unique genes associated with GoF variants, strict gene-stratified 10-fold cross-validation limits statistical robustness. Therefore, unlike the ClinVar benchmark where we prioritised generalisation (using gene-wise split), here we employed a random-split strategy for both LoF and GoF comparisons. This approach allows us to uniformly characterise the intrinsic information contribution of each modality within the available data, while the gene-stratified results are provided in Supplementary Fig. S3 to demonstrate cross-gene patterns.

In the "functionally normal vs. LoF" comparison, protein features play the dominant role. We observed that the AUROC peaks at a minimal CaLM weight of 0.13 and declines as codon influence increases (Fig. 3A and Supplementary Fig. S3). This trend was mirrored in the cross-validated effect scores, where ESM-2 offered superior resolution for LoF variants, while CaLM-derived scores were largely indistinguishable from the functionally normal background (Fig. 3B). As a result, the hybrid model's output converges with the ESM-2 baseline, indicating that protein features are sufficient to capture LoF pathogenicity without significant contribution from codon sequences (Supplementary Fig. S3 and S4).

Conversely, the "functionally normal vs. GoF" comparison reveals a distinct shift in evolutionary signals. Model performance peaks at a CaLM weight of 0.56, indicating that codon-level signals contribute meaningfully to discriminating GoF variants (Fig. 3C and Supplementary Fig. S3). However, the extensive distributional overlap across all three models and the wide confidence intervals in GoF ROC curves reflect the limited sample size and gene heterogeneity inherent to this variant class (Fig. 3D and Supplementary Fig. S3). Accordingly, the stark contrast between genes dependent solely on CaLM ($w = 1$) or ESM-2 ($w = 0$) suggests that codon-level contributions are gene-specific rather than universal, and should be interpreted with caution pending validation in larger datasets (Supplementary Fig. S5).

2.3 Codon-level constraints reflect nonsense and synonymous variant effects

We next evaluated the utility of codon-level information using CaLM in characterising *nonsense* and *synonymous* mutations across both ClinVar and ClinMAVE datasets. Because PLMs operate solely on amino acid sequences, they are inherently unable to detect nucleotide-level variant effects that do not alter the protein sequence or that result in premature truncation.

As illustrated in Fig. 4, CaLM successfully captured the broad distinctions between mutation categories, assigning significantly lower LLR scores to nonsense variants compared to synonymous ones. However, distinguishing mutation impact within these categories proved more challenging. Specifically, the LLR distributions for pathogenic

and benign synonymous variants exhibited substantial overlap (Fig. 4A), and the separation between functionally normal and LoF/GoF phenotypes was less distinct (Fig. 4B).

This may be partially driven by the inherent evolutionary frequencies of these variants: because natural selection strictly penalises nonsense mutations while tolerating synonymous ones, the CaLM model learns a strong prior for the latter. This implicitly aligns with the clinical observation that nonsense variants are predominantly pathogenic and synonymous ones benign (Fig. 1C and D). Crucially, although the specific discriminative signal for synonymous variants is subtle, CaLM captures a unique dimension of nucleotide-level constraint (evident in the nonsense vs. synonymous split) that remains completely invisible to PLMs.

2.4 Divergence between codon and protein language models reveals the impact of codon degeneracy

To investigate the biological drivers of model discordance, we isolated "conflict zones" within ClinVar where CaLM and ESM-2 predictions were maximally divergent. We identified 2,748 mutations exhibiting extreme LLR discrepancies, comprising 1,281 variants in the *CLM better* group and 1,467 in the *PLM better* group (see "Materials and methods - Identification of missense variant with divergent model predictions" and Fig. 5A). To characterise the specific substitution biases of each model, we quantified the enrichment of wt to mutant substitutions in each subset (e.g., *CLM better*) relative to the background distribution of the full missense dataset and assessed statistical significance using two-tailed Fisher's exact tests (see "Materials and methods - Characterisation of amino acid substitution preferences").

We found that the discordance between CLM and PLM predictions correlates strongly with shifts in codon degeneracy. Significant discrepancies are particularly pronounced in regions where mutations induce a transition between low- and high-codon degeneracy states (e.g., from Tryptophan (W), encoded by one codon, to Serine (S), encoded by six, or vice versa; highlighted in blue boxes in Fig. 5B and C).

This suggests that distinct evolutionary signals are being prioritised: substitutions preserving degeneracy (synonymous-like flexibility) tend to yield convergent effect scores, whereas dramatic shifts in degeneracy state expose the orthogonality of the models. The CLM appears to weigh the information loss in the nucleotide landscape, while the PLM focuses on the physicochemical impact of the residue change.

This divergence captures complementary biological constraints rather than simple model error. For instance, the PLM significantly outperforms the CLM in predicting Glycine (G) to Glutamic acid (E) substitutions (as shown by the enrichment in Fig. 5C and depletion in Fig. 5B). In this context, the PLM likely captures the specific structural intolerance (or tolerance) of the residue change, whereas the CLM may be overly sensitive to the reduction in codon state space (from 4 codons to 2). Taken together, these findings indicate that variant pathogenicity implies a composite function of residue fitness and codon optimality, modulated by the genetic code's structure.

2.5 Distinct constraints dictate variant pathogenicity across gene classes

Moving beyond individual amino acid substitutions, we asked whether the different predictive strengths of CLM and PLM are visible at the gene level and associated with specific biological functions.

We computed the absolute difference in AUROC (ΔROC_g) for each gene to measure their capacity to discriminate pathogenic missense variants. Based on this metric, we filtered the dataset and identified 29 CaLM-superior genes (*CLM better*) and 66 ESM-2-superior genes (*PLM better*), which were used for subsequent biological characterisation (see "Materials and methods — Gene-level performance evaluation via bootstrapping" and Fig. 5D). Within the *CLM better* subset, *PCYT1A* exhibited the most pronounced performance advantage for CaLM, followed by *DNMT1* and *MEF2C* (Fig. 5E). Notably, many of these genes, including *EYA1* [18], *GATA1* [19], and *EZH2* [20], function as transcriptional regulators or chromatin modifiers. Conversely, the genes favouring ESM-2 were led by *SUMF1* and *KCNT1* (Fig. 5F). This group predominantly encodes components of multiprotein complexes (e.g., *HBA1* [21], *RAG1* [22], and *TP53* [23]), representing properties manifested at the amino acid level like structural stability and complex assembly.

To evaluate constraint profiles for these gene subsets, we used two gnomAD metrics: probability of loss-of-function intolerance (pLI) and loss-of-function observed/expected upper bound fraction (LOEUF) [24]. We observed that genes in the *CLM better* subset exhibited significantly higher pLI scores and correspondingly lower LOEUF scores compared to the *PLM better* group, with both comparisons demonstrating a moderate effect size (two-sided Mann-Whitney U test, $p < 0.05$; rank-biserial $r = -0.352$ for pLI, and $r = 0.300$ for LOEUF; Fig. 5G and Supplementary Fig. S6). The consistently near-maximal pLI scores (≈ 1.0) and highly depleted LOEUF values within the *CLM better* subset indicate a strong biological trend toward haploinsufficiency. Consequently, the proper function of these genes is highly sensitive to precise gene dosage – a quantitative trait heavily regulated by translational kinetics. Unlike the PLM, the CLM uniquely captures codon-level signals that contribute to maintaining critical expression thresholds.

To further clarify the biological basis of this stratification, we performed functional enrichment analysis using Gene Ontology Molecular Function (GO:MF) terms. The *CLM better* genes were preferentially enriched in transcriptional regulatory functions, particularly transcription coregulator binding and chromatin binding (Fig. 5H). This concordance with near-maximal pLI scores suggests that the CLM captures the sequence constraints underlying gene expression regulation – where maintaining precise protein levels (dosage sensitivity) is as important as preserving protein function. We further stratified *PLM better* genes by pLI scores, yielding 26 high-pLI and 34 low-pLI genes. In the high-pLI subset, PLM-associated enrichment was dominated by complex membrane signalling machinery, including NMDA glutamate receptor and calcium channel activities. In contrast, the low-pLI subset showed enrichment for more localised enzymatic and ligand-binding functions, notably biotin binding (Fig. 5I). Collectively, these findings highlight the PLM's capacity to model both the

complex structural dependencies of membrane proteins and highly specific, localised biochemical interactions.

To concretely illustrate this mechanistic difference in dosage-sensitive genes, we examined a pathogenic variant in *MEF2C* (p.Leu38Pro, c.113T>C) (Fig. 5J). Here, we found a conflict: while the PLM assigned a moderate score (LLR_{y_k} = -4.77) reflecting structural risk, the CLM flagged a strong signal (LLR_{c_k} = -14.22). This discrepancy suggests that for *MEF2C*, the primary pathogenic driver may not be solely structural destabilisation, but rather a critical failure in translational efficiency required to maintain gene dosage.

Collectively, these results indicate that CLM and PLM capture complementary layers of biological information: the CLM is better suited to detecting disruptions in translational regulation and dosage sensitivity, whereas ESM-2 primarily resolves the biophysical constraints governing protein structure and stability.

3 Discussion

Traditional variant predictors often reduce the coding sequence to a mere precursor for amino acids, blind to codon identity. Moving beyond this view, we propose that pathogenic potential emerges from the combination of distinct biological modalities spanning mRNA and protein levels. To interrogate pathogenicity through sequence constraints, we prioritised single-sequence models, specifically ESM-2 and CaLM, over MSA-dependent architectures like AlphaMissense [25]. This strategic choice circumvents the confounding influence of alignment-based evolutionary bias, allowing us to cleanly isolate and quantify the independent contributions of PLM and CLM. Thus, our framework serves not as a competitor to clinical diagnostic tools, but as an analytical approach for dissecting variant pathogenicity.

We used the differences between LoF and GoF to validate the biological fidelity of this framework. Given that LoF pathogenicity is predominantly mediated by structural disruption that compromises protein stability [26–28], we postulated that codon-level signals would contribute less than biophysical constraints. The model confirmed this hierarchy by adaptively attenuating CaLM's weight to a minimal 0.13 for LoF variants, largely rendering the codon signal redundant. In sharp contrast, the model elevated CaLM to a substantial weight (0.56) for characterising GoF pathogenicity. These GoF findings are constrained by the limited sample size and genetic diversity, necessitating further validation in more extensive cohorts. This shift is consistent with prior observations: while GoF mutations often exert much milder effects on protein structure compared to LoF variants [28], they frequently implicate process-dependent constraints, such as dosage regulation [29, 30] or the precise kinetic control required for adopting active conformations [31, 32]. Thus, rather than naively ensembling two scores, the hybrid framework modulates information sources to recapitulate the underlying biological context, underscoring that accurate pathogenicity assessment requires integrating information from both protein and mRNA levels.

Our findings imply that codon-level constraints remain clinically significant in missense contexts, extending beyond synonymous variation [33, 34]. We conceptualise drastic optimality shifts as "translational shock," a phenomenon where abrupt changes in translational kinetics reduce protein output [35–37]. Crucially, our pan-cell line analysis dissociates this phenomenon from intrinsic background transcript turnover. We observed no significant difference in mRNA half-lives between *CLM better* and *PLM better* genes across 20 diverse cell types (Supplementary Table S4). This indicates that the effect is not driven by gene instability, but rather by variant-specific disruptions in translational kinetics that compromise expression efficiency. Mechanistically, this supports a composite fitness model wherein missense mutations compromise fitness through two complementary mechanisms: structural impairment (captured by PLM) and dosage reduction (captured by CLM) (Fig. 6). This model is strongly supported by the enrichment of *CLM better* variants in dosage-sensitive high-pLI genes (Fig. 5G), such as *EZH2* [38] and *MEF2C* [39]. For these developmental regulators, even small reductions in protein abundance can be clinically significant (Fig. 5H) [24]. Thus, the CLM uniquely captures signals consistent with an often overlooked dimension: disruptions in expression regulation that lie outside the scope of standalone PLMs.

Synonymous variants have historically been difficult to classify due to the absence of protein alteration. Our results suggest that pure sequence-based modelling of synonymous effects remains challenging (Fig. 4). This limitation likely stems from dependencies on mRNA secondary structure [40] or splicing regulation [41]. While CaLM may potentially capture local splicing motifs embedded within the coding sequence (e.g., exonic splicing enhancers), accurate prediction likely requires the intronic context or global structural awareness that remains outside the model's scope. Nevertheless, CaLM establishes a foundational baseline where PLMs provide no signal. This capability opens a new avenue for codon-aware prediction in non-coding or silent variant spaces.

Ultimately, the integration of CLM and PLM establishes a proof-of-concept for combining complementary biological signals. Beyond clinical applications, we envision this compositional strategy as a paradigm for future computational biology, enabling the flexible assembly of foundation models to address multi-layered biological questions.

4 Materials and methods

4.1 ClinVar variant dataset

ClinVar [16] is a comprehensive database that aggregates genetic variants associated with disease, together with clinical phenotypes and supporting evidence. For this study, we downloaded the complete set of variant annotations from `variant_summary.txt` (release through October 8, 2024) available via the ClinVar FTP site (https://ftp.ncbi.nlm.nih.gov/pub/clinvar). We restricted the dataset to single-nucleotide substitutions classified as *missense*, *nonsense*, or *synonymous*. To ensure data reliability, the following filtering criteria were implemented:

1. restricted to variants annotated with a definitive clinical significance of *benign* (including likely benign) or *pathogenic* (including likely pathogenic);
2. required a ClinVar review status of at least one star;
3. excluded variants located in extremely large genes (e.g., *TTN*); and
4. for genes with annotations across multiple transcripts, only the transcript with the largest number of annotated variants was retained, establishing a robust one-to-one gene-transcript mapping.

The final ClinVar dataset comprised 137,350 missense variants across 13,791 genes, 46,386 nonsense variants across 3,291 genes, and 527,579 synonymous variants across 11,593 genes (Fig. 1C and Supplementary Table S1).

4.2 ClinMAVE variant dataset

ClinMAVE [17] is a curated database intended to support the clinical translation of high-throughput functional evidence generated by MAVEs. ClinMAVE implements a multi-layer functional annotation framework that applies reviewed score thresholds to MAVE-derived measurements, assigning each variant to one of three functional categories: functionally normal, LoF, or GoF. In this study, we compiled a dataset of *missense*, *nonsense*, and *synonymous* variants (December 15, 2025) and applied the following filters:

1. MAVE variants were generated by Deep Mutational Scanning (DMS); and
2. variants with coding sequence lengths greater than 10,000 were excluded.

The final ClinMAVE dataset comprised 253,078 missense variants across 429 genes, 11,868 nonsense variants across 418 genes, and 18,991 synonymous variants across 23 genes (Fig. 1D and Supplementary Table S2).

4.3 Sequences collection

For each single-point variant compiled from ClinVar and ClinMAVE, we retrieved the corresponding full GenBank records from the NCBI GenBank database. We then parsed these records to identify coding DNA sequence features and extracted the associated cDNA sequences and their translated protein products.

4.4 CaLM and ESM-2

To ensure a fair comparison across modalities, we employed both CaLM (86 million parameters; downloaded from https://github.com/oxpig/CaLM) [15] and ESM-2 (150 million parameters; downloaded from https://huggingface.co/facebook/esm2_t30_150M_UR50D) [3], two transformer-based language models. They were trained via masked language modelling objectives; however, they differ in tokenization strategies, training datasets, and output representations. Specifically:

1. Tokens: CaLM tokenizes cDNA sequences into codons, whereas ESM-2 tokenizes protein sequences into residues
2. Training data: CaLM is trained on a corpus of 9 million cDNA sequences sourced from the European Nucleotide Archive, whereas ESM-2 leverages 65 million protein sequences derived from the UniRef database
3. Output logits: CaLM generates position-specific effect scores for all 64 codons at each codon position, while ESM-2 produces effect scores for each of the 20 canonical amino acids at each position

Neither model enforces sequence length constraints; moreover, we did not specifically design our approach for handling long sequences.

4.5 Missense effect scores

Language models can estimate the probability of a mutation at a given position by leveraging only the surrounding sequence context, without the need for additional training. The wt cDNA sequence was input into CaLM, while its corresponding wt protein sequence was processed using ESM-2. A softmax function was then applied to generate an uncalibrated probability distribution over codons or residues.

Consider a cDNA sequence $\mathbf{x} = (x_1, x_2, \ldots, x_{3L+3})$, where each nucleotide $x_i \in \{A, T, C, G\}$ and L denotes the number of residues encoded by the cDNA sequence. It comprises a start codon, a coding region, and a stop codon, resulting in a total length of $3L + 3$ nucleotides. A codon $c_k = (x_{3k-2}, x_{3k-1}, x_{3k})$, for $k = 1, 2, \ldots, L + 1$, where $k = L + 1$ corresponds to the stop codon (e.g., $c_1 = (A, T, G)$). During translation, each codon c_k is mapped to a residue according to the genetic code. We thus define the protein sequence as $\mathbf{y} = (y_1, y_2, \ldots, y_L)$, where each y_i represents one of the 20 standard amino acids (typically denoted by their one-letter codes, e.g., $\{A, R, N, D, C, \ldots\}$). (Note that although the cDNA sequence contains $L + 1$ codons, the last codon c_{L+1} is a stop codon and does not encode a residue.) Only the first L triplet of three nucleotides are encoded into amino acids.

A missense mutation at position i in a cDNA sequence, denoted by $\tilde{x}_i$, altering the wt codon c_k^{wt} to a mutated form $\tilde{c}_k$, thereby replacing the residue y_k^{wt} with $\tilde{y}_k$ in the protein sequence. To quantify the effect of this mutation, we compute LLRs at both codon and residue levels from the model's per-position log-likelihoods over all amino acids or codons, including the wt amino acid and wt codon, following the methods in [4]:

$$\text{LLR}_{c_k} = \log \frac{\Pr(c_k = \tilde{c}_k)}{\Pr(c_k = c_k^{\text{wt}})} \tag{1}$$

$$\text{LLR}_{y_k} = \log \frac{\Pr(y_k = \tilde{y}_k)}{\Pr(y_k = y_k^{\text{wt}})} \tag{2}$$

To quantify the relative contribution of each modality rather than obscuring it within a complex non-linear head (e.g., a neural network), we define a missense effect score as a linear combination of the codon- and residue-level LLRs:

$$\text{LLR}_{\text{hybrid}} = w \cdot \text{LLR}_{c_k} + (1 - w) \cdot \text{LLR}_{y_k} \tag{3}$$

where $w \in [0, 1]$ is a weighting parameter that controls the relative contribution of the codon model

4.6 Model training and validation

To strictly evaluate model generalisation, we employed a gene-wise 10-fold cross-validation split for the global "benign vs. pathogenic" classification task (ClinVar) using `Scikit-Learn` (v1.6.1) [42]. In this strategy, all variants of a given gene are assigned exclusively to either the training or the test set within each fold, preventing data leakage arising from gene-specific sequence dependencies.

In contrast, for the mechanistic dissection of functional phenotypes (ClinMAVE), we prioritised the statistical characterisation of intrinsic information content over zero-shot generalisation. Given the limited number of unique genes available for GoF variants, which precludes stable gene-wise estimation (as evidenced by high variance in cross-gene performance), we adopted a random split strategy for *both* functionally normal vs. LoF" and functionally normal vs. GoF" comparisons. This framework ensures a fair assessment of the information contribution of codon versus residue constraints. For completeness, we also performed gene-wise cross-validation for these tasks to assess cross-gene heterogeneity, with results reported in the Supplementary Material.

Within each cross-validation fold, the weighting parameter w (search range from 0 to 1) was optimised solely on the training data via Bayesian optimisation to maximise the AUROC [43, 44]. The optimised w was subsequently evaluated on the corresponding held-out test fold. This optimisation procedure was implemented using the `BayesianOptimization` function from the `bayesian-optimization` package (v2.0.3), configured with 10 initial exploration steps followed by 20 optimisation iterations.

4.7 Metrics for variant classification

For the ClinVar dataset, *pathogenic* variants were defined as the positive class and *benign* variants as the negative class. In the ClinMAVE dataset, *functionally normal* variants served as the negative class, whereas *LoF* and *GoF* variants were assigned to the positive class for their respective classification tasks.

The AUROC metric quantifies the trade-off between the True Positive Rate (TPR) and the False Positive Rate (FPR), thereby providing a robust overall assessment of classifier performance. However, in datasets with significant class imbalance, AUROC

may not fully capture nuanced differences. To address this limitation, we employed the AUPRC metric to assess the precision-recall trade-off. For both ROCs and PRs, we calculated the AUROC and AUPRC for each validation fold, respectively, and assessed the statistical significance of the differences using a two-tailed paired t-test via `scipy.stats` (v1.13.1) [45].

4.8 Identification of missense variants with divergent model predictions

To quantify the modality-specific discrepancy in mutational likelihoods learned by the CLM and PLM, we formulated a differential metric based on LLRs, denoted as ΔL_k, for each missense variant k:

$$\Delta \mathrm{L}_k = \mathrm{LLR}_{y_k} - \mathrm{LLR}_{c_k} \tag{4}$$

We isolated variants exhibiting the most extreme discrepancies by selecting the top and bottom 1% of the ΔL distribution. This selection criterion resulted in a dataset of 2,748 variants (representing the 2% most divergent ΔL), which ensures a robust sample size for the characterisation of specific amino acid substitution preferences. We further stratified these variants into two subsets based on which model's prediction was more consistent with the ClinVar's label:

1. *PLM better* (S_P): Variants where the PLM provided a superior score. Specifically, for pathogenic mutations, the PLM assigned a lower LLR ($\mathrm{LLR}_{y_k} < \mathrm{LLR}_{c_k}$), whereas for benign mutations, it assigned a higher LLR ($\mathrm{LLR}_{y_k} > \mathrm{LLR}_{c_k}$).
2. *CLM better* (S_C): Variants where the CLM provided the superior score, satisfying the inverse conditions (i.e., $\mathrm{LLR}_{c_k} < \mathrm{LLR}_{y_k}$ for pathogenic and $\mathrm{LLR}_{c_k} > \mathrm{LLR}_{y_k}$ for benign variants).

Based on these criteria, we finally identified 1,281 missense variants as *CLM better* (S_C) and 1,467 missense variants as *PLM better* (S_P).

4.9 Characterisation of amino acid substitution preferences

To assess whether the PLM and CLM exhibit distinct preferences for specific types of missense variants, we compared the frequency distributions of wt to mutant amino acid substitutions in each subset (S_P and S_C) against their overall distribution in the complete missense dataset (A).

The frequency of a given amino acid substitution p (e.g., $W \rightarrow S$) in subset S_i ($i \in \{P, C\}$) is defined as:

$$f_{S_i}(p) = \frac{n_{S_i}(p)}{N_{S_i}} \tag{5}$$

where $n_{S_i}(p)$ denotes the count of substitution p in subset S_i, and N_{S_i} represents the total count of all missense variants in S_i

Similarly, the background frequency of the same substitution in the complete dataset A, denoted as $f_A(p)$, was calculated using the total counts from A. To quantify the relative enrichment or depletion of specific substitutions, we computed the

Bias Ratio (BR). For a substitution p in subset S_i, the BR is defined as:

$$\mathrm{BR}_{S_i}(p) = \frac{f_{S_i}(p)}{f_A(p)} \qquad (6)$$

A substitution was classified as *enriched* in subset S_i if $\mathrm{BR}_{S_i}(p) > 1$, indicating a higher prevalence relative to the overall dataset, whereas substitutions with $\mathrm{BR}_{S_i}(p) < 1$ were considered *depleted*.

To assess the statistical significance of these enrichment patterns, we employed two-tailed Fisher's exact tests. This method was chosen for its robustness in handling small sample sizes associated with specific rare amino acid substitutions. For each substitution p and subset S_i, we constructed a 2×2 contingency table comparing the counts in the subset (S_i) against the counts in the remainder of the dataset ($A \setminus S_i$) to ensure independence. Tests were performed using the `fisher_exact` function from `scipy.stats` (v1.13.1). To account for multiple hypothesis testing across all unique substitution types, we applied the Benjamini-Hochberg (BH) procedure using the `multipletests` function from `statsmodels.stats.multitest` (v0.14.6) to control the False Discovery Rate (FDR) at $\alpha = 0.01$. Substitutions with adjusted p-values below this threshold were considered statistically significant.

4.10 Gene-level performance evaluation via bootstrapping

To quantify performance at the gene level, we computed the ROC for CaLM and ESM-2 across genes (g), separately. We restricted the analysis to genes with at least 20 variants and imposed a maximum class imbalance ratio of 5:1 (i.e., the ratio of pathogenic to benign variants, or vice versa, did not exceed 5). We defined the performance difference, $\Delta\mathrm{ROC}_g$, as:

$$\Delta\mathrm{ROC}_g = \mathrm{ROC}_P(g) - \mathrm{ROC}_C(g) \qquad (7)$$

where $\mathrm{ROC}_P(g)$ and $\mathrm{ROC}_C(g)$ denote the ROC scores of the ESM-2 (PLM) and CaLM (CLM) models for gene g, respectively

To rigorously assess the statistical significance of $\Delta\mathrm{ROC}_g$, we employed a stratified bootstrap approach. For each gene, we performed 1,000 iterations of resampling with replacement, stratified by the ClinVar's labels to maintain the original class distribution. A 95% Confidence Interval (CI) for the performance difference was constructed using the percentile method. Genes were categorised as having a significant model preference only if the 95% CI did not overlap with zero. This filtering resulted in the identification of 29 *CLM better* genes and 66 *PLM better* genes.

4.11 Gene-level constraint analysis

Evolutionary constraints were assessed using two complementary metrics retrieved from the gnomAD database (v2.1.1) [24]: pLI and LOEUF. The pLI score quantifies the probability of intolerance to heterozygous LoF variation, serving as a proxy for haploinsufficiency and dosage sensitivity [46]. Specifically, genes are stratified into two distinct categories to reflect their mutational constraint: high pLI (pLI ≥ 0.9),

which identifies genes highly sensitive to dosage changes, and low pLI (pLI ≤ 0.1), which comprises genes generally more tolerant of LoF mutations. Conversely, LOEUF provides a continuous and robust measure of mutational constraint, where lower values signify stronger purifying selection against LoF variants [24].

To ensure robustness, we included only genes with available high-confidence constraint metrics, resulting in a finalised dataset of 28 (out of 29) *CLM better* and 64 (out of 66) *PLM better* genes. Within the *PLM better* group, 26 genes were classified as high pLI and 34 as low pLI. Differences in the score distributions between these two cohorts were evaluated using a two-sided Mann-Whitney U test implemented in `scipy.stats` (v1.13.1). Additionally, the magnitude of these differences was quantified by calculating the rank-biserial correlation coefficient (r) as a measure of effect size [47].

4.12 mRNA stability analysis

To rigorously assess whether the performance gain of the CLM is driven by transcript instability, we downloaded mRNA half-life data from the Transcriptome Turnover Database [48]. We used 20 diverse human cell lines, including epithelial, hepatic, immune, and neural progenitor cells, to exclude tissue-specific biases. For each cell line, multiple half-life measurements associated with a single gene (e.g., from technical replicates) were aggregated by calculating the median value to derive a robust gene-level estimate. Cell lines with sufficient coverage (more than 5 genes in both the *CLM better* and *PLM better* groups) were included to ensure statistical validity. We evaluated the difference in stability between *CLM better* and *PLM better* gene groups using a one-sided Mann-Whitney U test implemented in `scipy.stats` (v1.13.1).

4.13 Functional enrichment analysis

Functional enrichment analysis for GO:MF was performed in R (v4.2.3) using the `gprofiler2` package [49]. The `gost` function was utilized to identify significantly enriched terms for the input gene list against the annotated *Homo sapiens* background. The *p*-values were adjusted for multiple testing using the `g_SCS` correction method, with a significance threshold of 0.05. To focus on specific biological mechanisms, broad terms were manually excluded.

4.14 Data analysis and visualization

Data analysis and plot preparation were performed in Jupyter Notebook v6.5.5 using Python v3.11.11 and the following libraries: numpy v1.26.4, pandas v2.2.2, scipy.stats v1.13.1, matplotlib v3.10.0, seaborn v0.13.2, and ggplot2 v4.0.1.

Acknowledgements

Figure 1 and 6 were created with BioRender (https://www.biorender.com/).
Author contributions: R.C. and M.B. designed the study. R.C. implemented code and methods. M.B., G.F. and N.P. supervised the project. All authors contributed to the writing, reviewed, and approved the final manuscript.

Conflict of interest

None declared.

Funding

This work was supported by University of Queensland Research Training Scholarships to R.C. N.P. and M.B. received support from Medical Research Future Fund (2016033).

Data availability

All the data and code necessary to reproduce the analyses, figures, and results are available on our GitHub repository: https://github.com/Cassie818/Viral-mut.

Fig. 1: Linguistic duality reveals orthogonal evolutionary constraints across the central dogma.

(A) Semantic alignment across natural languages. English and German sentences convey identical core meanings yet differ in syntactic structure and expression. **(B)** An analogy between natural languages and biological language models. Just as different languages convey similar meanings with subtle interpretational differences, two language models (i.e., CLM and PLM) trained on the same coding region may develop nuanced variations in their outputs. **(C–D)** Dataset composition and label distribution. The distribution of ground-truth labels for synonymous, missense, and nonsense variants in the (C) ClinVar dataset and (D) ClinMAVE dataset. **(E)** Schematic of the hybrid model architecture. The framework employs two parallel inference streams: CaLM processes wt cDNA sequences to compute codon-level effect scores (LLR_{c_k}), while ESM-2 independently processes protein sequences to derive residue-level scores (LLR_{y_k}). The two scores are then unified through a Bayesian-optimized integration strategy to predict variant pathogenicity. This figure was created using BioRender.com.

Fig. 2: Codon and protein constraints define a composite landscape of variant pathogenicity.

(A) Hyperparameter tuning for the combination weight (w) of CaLM for the "pathogenic vs. benign" task. The vertical dashed line indicates the optimal weight ($w = 0.49$) determined by maximizing ROC on the training set using a gene-stratified split. **(B)** ROC curves benchmarking model performance on the held-out test set. Shaded areas represent the standard deviation across 10 folds. The statistical significance of the performance difference between the hybrid model and each single-modality baseline was evaluated using a two-tailed paired t-test (***$p < 0.001$). **(C)** Distributions of LLR for benign (blue shades: benign, likely Benign) and pathogenic (red shades: pathogenic, likely pathogenic) variants in ClinVar, as predicted by ESM-2, CaLM, and the hybrid model. The dashed lines represent the combined density estimates for pathogenic (red) and benign (blue) classes. The yellow shaded region indicates the overlap area between these densities.

Fig. 3: Decoupling evolutionary drivers of LoF and GoF mechanisms.

(A) Hyperparameter tuning for the weight (w) of CaLM for the "functionally normal vs. LoF" task. The vertical dashed line indicates the optimal weight ($w = 0.13$) determined by maximizing AUROC on the training set using a random split. **(B)** Split violin plots showing the distribution of LLR for functionally normal and LoF variants. Dashed lines indicate the quartiles. **(C-D)** Same as (A-B), but applied to the "functionally normal vs. GoF" task.

Fig. 4: Codon-level constraints reflect nonsense and synonymous variant effects.

(A) Comparison of LLR distributions between pathogenic and benign variants in the ClinVar dataset, grouped by nonsense and synonymous subtypes. **(B)** Comparison of LLR distributions across functional categories from ClinMAVE datasets. The left panel contrasts functionally normal variants with LoF variants, while the right panel contrasts functionally normal variants with GoF variants, stratified by nonsense and synonymous categories. Data are presented as split violin plots with internal box plots representing the median and inter-quartile ranges.

Fig. 5: Dissecting the biological complementarity of CLM and PLM.

(A) Distribution of the difference in LLRs (ΔL) between the two models for ClinVar missense variants. The dashed blue lines mark the 1th and 99th percentiles, isolating variants where one modality captures pathogenic features significantly better than the other. **(B-C)** Heatmaps depicting amino acid substitution biases associated with superior performance of (B) CaLM and (C) ESM-2. Axes represent wt (x-axis) and mutant (y-axis) residues, ordered by increasing codon degeneracy; coloured background blocks along the diagonal distinguish amino acid groups sharing the same codon degeneracy count. Circle size and colour intensity quantify the relative enrichment (red) or depletion (blue) of specific substitutions compared to background distribution of all missense variants. Statistically significant enrichment are denoted by stars using two-tailed Fisher's exact tests. **(D)** Gene-level performance comparison. Scatter plot of AUROC values for individual genes predicted by CaLM (y-axis) versus ESM-2 (x-axis). Genes where CLM performs substantially better are coloured in red, while genes favouring PLM are in blue. **(E-F)** The top 10 genes exhibiting the largest performance gap favouring (E) CLM and (F) PLM. **(G)** Violin plots showing the distribution of pLI scores for genes in the *CLM better* and *PLM better* groups. Black dots represent individual genes. Statistical significance was determined by a two-sided Mann-Whitney U test ($p = 0.007$), with the effect size quantified by the rank-biserial correlation coefficient ($r = -0.352$). **(H-I)** Enriched GO:MF terms for (H) *CLM better* genes and (I) *PLM better* genes. Bar lengths represent the number of genes (count) associated with each term. The color gradient indicates the significance level (adjusted p-value). For *PLM better* genes, the enrichment results are further stratified into high pLI (top) and low pLI (bottom) gene sets. **(J)** Case study of effect score divergence in *MEF2C* (p.Leu38Pro). Bar chart showing LLRs for that pathogenic variant.

Fig. 6: Schematic of the proposed mechanism.

Missense variants exert deleterious effects through parallel mechanisms: (1) structural impairment (left): amino acid changes causing protein misfolding or instability, a dimension leveraged by the PLM. (2) dosage attenuation (right): suboptimal codon usage causing "translational shock," which triggers ribosome stalling, leading to reduced protein levels. This kinetic dimension is captured by the CLM. This figure was created using BioRender.com.

References

[1] Meier, J., Rao, R., Verkuil, R., Liu, J., Sercu, T., Rives, A.: Language models enable zero-shot prediction of the effects of mutations on protein function. Advances in neural information processing systems **34**, 29287–29303 (2021)

[2] Frazer, J., Notin, P., Dias, M., Gomez, A., Min, J.K., Brock, K., Gal, Y., Marks, D.S.: Disease variant prediction with deep generative models of evolutionary data. Nature **599**(7883), 91–95 (2021)

[3] Lin, Z., Akin, H., Rao, R., Hie, B., Zhu, Z., Lu, W., Smetanin, N., Verkuil, R., Kabeli, O., Shmueli, Y., *et al.*: Evolutionary-scale prediction of atomic-level protein structure with a language model. Science **379**(6637), 1123–1130 (2023)

[4] Brandes, N., Goldman, G., Wang, C.H., Ye, C.J., Ntranos, V.: Genome-wide prediction of disease variant effects with a deep protein language model. Nature Genetics **55**(9), 1512–1522 (2023)

[5] Rives, A., Meier, J., Sercu, T., Goyal, S., Lin, Z., Liu, J., Guo, D., Ott, M., Zitnick, C.L., Ma, J., *et al.*: Biological structure and function emerge from scaling unsupervised learning to 250 million protein sequences. Proceedings of the National Academy of Sciences **118**(15), 2016239118 (2021)

[6] Hsu, C., Verkuil, R., Liu, J., Lin, Z., Hie, B., Sercu, T., Lerer, A., Rives, A.: Learning inverse folding from millions of predicted structures. In: International Conference on Machine Learning, pp. 8946–8970 (2022). PMLR

[7] Yang, K.K., Zanichelli, N., Yeh, H.: Masked inverse folding with sequence transfer for protein representation learning. Protein Engineering, Design and Selection **36**, 015 (2023)

[8] Rao, R.M., Liu, J., Verkuil, R., Meier, J., Canny, J., Abbeel, P., Sercu, T., Rives, A.: Msa transformer. In: International Conference on Machine Learning, pp. 8844–8856 (2021). PMLR

[9] Notin, P., Van Niekerk, L., Kollasch, A.W., Ritter, D., Gal, Y., Marks, D.S.: Trancepteve: Combining family-specific and family-agnostic models of protein sequences for improved fitness prediction. bioRxiv, 2022–12 (2022)

[10] Su, J., Han, C., Zhou, Y., Shan, J., Zhou, X., Yuan, F.: Saprot: Protein language modeling with structure-aware vocabulary. bioRxiv, 2023–10 (2023)

[11] Li, M., Tan, Y., Ma, X., Zhong, B., Yu, H., Zhou, Z., Ouyang, W., Zhou, B., Hong, L., Tan, P.: Prosst: Protein language modeling with quantized structure and disentangled attention. bioRxiv, 2024–04 (2024)

[12] Zvyagin, M., Brace, A., Hippe, K., Deng, Y., Zhang, B., Bohorquez, C.O., Clyde, A., Kale, B., Perez-Rivera, D., Ma, H., *et al.*: Genslms: Genome-scale language

models reveal sars-cov-2 evolutionary dynamics. The International Journal of High Performance Computing Applications **37**(6), 683–705 (2023)

[13] Boshar, S., Trop, E., Almeida, B.P., Copoiu, L., Pierrot, T.: Are genomic language models all you need? exploring genomic language models on protein downstream tasks. Bioinformatics **40**(9), 529 (2024)

[14] Laubitz, Z.: The gain and loss of information during translation (1983)

[15] Outeiral, C., Deane, C.M.: Codon language embeddings provide strong signals for use in protein engineering. Nature Machine Intelligence **6**(2), 170–179 (2024)

[16] Landrum, M.J., Lee, J.M., Benson, M., Brown, G., Chao, C., Chitipiralla, S., Gu, B., Hart, J., Hoffman, D., Hoover, J., *et al.*: Clinvar: public archive of interpretations of clinically relevant variants. Nucleic acids research **44**(D1), 862–868 (2016)

[17] Ma, C., Li, Z., Tang, X., Li, P., Li, L., Wang, S., Wu, J., Luo, L., Liu, Y., Zhang, Z., et al.: Clinmave: a curated database for clinical application of data from multiplexed assays of variant effect. Nucleic Acids Research, 1102 (2025)

[18] Xu, P.-X., Adams, J., Peters, H., Brown, M.C., Heaney, S., Maas, R.: Eya1-deficient mice lack ears and kidneys and show abnormal apoptosis of organ primordia. Nature genetics **23**(1), 113–117 (1999)

[19] Crispino, J.D., Horwitz, M.S.: Gata factor mutations in hematologic disease. Blood, The Journal of the American Society of Hematology **129**(15), 2103–2110 (2017)

[20] Viré, E., Brenner, C., Deplus, R., Blanchon, L., Fraga, M., Didelot, C., Morey, L., Van Eynde, A., Bernard, D., Vanderwinden, J.-M., *et al.*: The polycomb group protein ezh2 directly controls dna methylation. Nature **439**(7078), 871–874 (2006)

[21] Barrera-Reyes, P.K., Tejero, M.E.: Genetic variation influencing hemoglobin levels and risk for anemia across populations. Annals of the New York Academy of Sciences **1450**(1), 32–46 (2019)

[22] Kim, M.-S., Chuenchor, W., Chen, X., Cui, Y., Zhang, X., Zhou, Z.H., Gellert, M., Yang, W.: Cracking the dna code for v (d) j recombination. Molecular cell **70**(2), 358–370 (2018)

[23] Joerger, A.C., Fersht, A.R.: The tumor suppressor p53: from structures to drug discovery. Cold Spring Harbor perspectives in biology **2**(6), 000919 (2010)

[24] Karczewski, K.J., Francioli, L.C., Tiao, G., Cummings, B.B., Alföldi, J., Wang,

Q., Collins, R.L., Laricchia, K.M., Ganna, A., Birnbaum, D.P., *et al.*: The mutational constraint spectrum quantified from variation in 141,456 humans. Nature **581**(7809), 434–443 (2020)

[25] Cheng, J., Novati, G., Pan, J., Bycroft, C., Žemgulytė, A., Applebaum, T., Pritzel, A., Wong, L.H., Zielinski, M., Sargeant, T., *et al.*: Accurate proteome-wide missense variant effect prediction with alphamissense. Science **381**(6664), 7492 (2023)

[26] Matreyek, K.A., Starita, L.M., Stephany, J.J., Martin, B., Chiasson, M.A., Gray, V.E., Kircher, M., Khechaduri, A., Dines, J.N., Hause, R.J., *et al.*: Multiplex assessment of protein variant abundance by massively parallel sequencing. Nature genetics **50**(6), 874–882 (2018)

[27] Cagiada, M., Johansson, K.E., Valanciute, A., Nielsen, S.V., Hartmann-Petersen, R., Yang, J.J., Fowler, D.M., Stein, A., Lindorff-Larsen, K.: Understanding the origins of loss of protein function by analyzing the effects of thousands of variants on activity and abundance. Molecular Biology and Evolution **38**(8), 3235–3246 (2021)

[28] Gerasimavicius, L., Livesey, B.J., Marsh, J.A.: Loss-of-function, gain-of-function and dominant-negative mutations have profoundly different effects on protein structure. Nature communications **13**(1), 3895 (2022)

[29] Lampson, B.L., Pershing, N.L., Prinz, J.A., Lacsina, J.R., Marzluff, W.F., Nicchitta, C.V., MacAlpine, D.M., Counter, C.M.: Rare codons regulate kras oncogenesis. Current Biology **23**(1), 70–75 (2013)

[30] Supek, F., Miñana, B., Valcárcel, J., Gabaldón, T., Lehner, B.: Synonymous mutations frequently act as driver mutations in human cancers. Cell **156**(6), 1324–1335 (2014)

[31] Kimchi-Sarfaty, C., Oh, J.M., Kim, I.-W., Sauna, Z.E., Calcagno, A.M., Ambudkar, S.V., Gottesman, M.M.: A" silent" polymorphism in the mdr 1 gene changes substrate specificity. Science **315**(5811), 525–528 (2007)

[32] Buhr, F., Jha, S., Thommen, M., Mittelstaet, J., Kutz, F., Schwalbe, H., Rodnina, M.V., Komar, A.A.: Synonymous codons direct cotranslational folding toward different protein conformations. Molecular cell **61**(3), 341–351 (2016)

[33] Bonekamp, F., Dalbøge, H., Christensen, T., Jensen, K.F.: Translation rates of individual codons are not correlated with trna abundances or with frequencies of utilization in escherichia coli. Journal of bacteriology **171**(11), 5812–5816 (1989)

[34] Akashi, H.: Synonymous codon usage in drosophila melanogaster: natural selection and translational accuracy. Genetics **136**(3), 927–935 (1994)

[35] Yu, C.-H., Dang, Y., Zhou, Z., Wu, C., Zhao, F., Sachs, M.S., Liu, Y.: Codon usage influences the local rate of translation elongation to regulate co-translational protein folding. Molecular cell **59**(5), 744–754 (2015)

[36] Liu, Y., Yang, Q., Zhao, F.: Synonymous but not silent: the codon usage code for gene expression and protein folding. Annual review of biochemistry **90**(1), 375–401 (2021)

[37] Walsh, I.M., Bowman, M.A., Soto Santarriaga, I.F., Rodriguez, A., Clark, P.L.: Synonymous codon substitutions perturb cotranslational protein folding in vivo and impair cell fitness. Proceedings of the National Academy of Sciences **117**(7), 3528–3534 (2020)

[38] Suri, T., Dixit, A.: The phenotype of ezh2 haploinsufficiency—1.2-mb deletion at 7q36. 1 in a child with tall stature and intellectual disability. American Journal of Medical Genetics Part A **173**(10), 2731–2735 (2017)

[39] Rocha, H., Sampaio, M., Rocha, R., Fernandes, S., Leão, M.: Mef2c haploinsufficiency syndrome: Report of a new mef2c mutation and review. European journal of medical genetics **59**(9), 478–482 (2016)

[40] Duan, J., Wainwright, M.S., Comeron, J.M., Saitou, N., Sanders, A.R., Gelernter, J., Gejman, P.V.: Synonymous mutations in the human dopamine receptor d2 (drd2) affect mrna stability and synthesis of the receptor. Human molecular genetics **12**(3), 205–216 (2003)

[41] Sauna, Z.E., Kimchi-Sarfaty, C.: Understanding the contribution of synonymous mutations to human disease. Nature Reviews Genetics **12**(10), 683–691 (2011)

[42] Pedregosa, F., Varoquaux, G., Gramfort, A., Michel, V., Thirion, B., Grisel, O., Blondel, M., Prettenhofer, P., Weiss, R., Dubourg, V., *et al.*: Scikit-learn: Machine learning in python. the Journal of machine Learning research **12**, 2825–2830 (2011)

[43] Auer, P.: Using confidence bounds for exploitation-exploration trade-offs. Journal of Machine Learning Research **3**(Nov), 397–422 (2002)

[44] Hie, B., Zhong, E.D., Berger, B., Bryson, B.: Learning the language of viral evolution and escape. Science **371**(6526), 284–288 (2021)

[45] Virtanen, P., Gommers, R., Oliphant, T.E., Haberland, M., Reddy, T., Cournapeau, D., Burovski, E., Peterson, P., Weckesser, W., Bright, J., *et al.*: Scipy 1.0: fundamental algorithms for scientific computing in python. Nature methods **17**(3), 261–272 (2020)

[46] Lek, M., Karczewski, K.J., Minikel, E.V., Samocha, K.E., Banks, E., Fennell, T., O'Donnell-Luria, A.H., Ware, J.S., Hill, A.J., Cummings, B.B., *et al.*: Analysis

of protein-coding genetic variation in 60,706 humans. Nature **536**(7616), 285–291 (2016)

[47] Kerby, D.S.: The simple difference formula: An approach to teaching nonparametric correlation. Comprehensive Psychology **3**, 11 (2014)

[48] Jiang, H., Xu, Z., Li, T., Wang, J., Xie, Z.: Ttdb: a comprehensive transcriptome turnover database for exploring mrna stability. Nucleic Acids Research **54**(D1), 350–357 (2026)

[49] Kolberg, L., Raudvere, U., Kuzmin, I., Adler, P., Vilo, J., Peterson, H.: g: Profiler—interoperable web service for functional enrichment analysis and gene identifier mapping (2023 update). Nucleic acids research **51**(W1), 207–212 (2023)