Accessory subunits of PRC2 mimic H3K27me3 to restrict the spread of Polycomb domains

Samuel C Agius[1], Marion Boudes[1], Evan Healy[1], Yi-Cheng Chang[2], Melanie Murray[1], Jan Manent[2], Jeanette Rientjes[3], Zhihao Lai[1], Sophie Nixon[1], Vitalina Levina[1], Edwina McGlinn[2]#, Chen Davidovich[1]#.

1. Department of Biochemistry and Molecular Biology, Biomedicine Discovery Institute, Faculty of Medicine, Nursing and Health Sciences, Monash University, Clayton, VIC, Australia.
2. Australian Regenerative Medicine Institute, Monash University, Clayton, VIC, Australia
3. Monash Genome Modification Platform, Monash University, Clayton, VIC, Australia.

Correspondence to Edwina.Mcglinn@monash.edu and Chen.Davidovich@monash.edu

Abstract

The Polycomb repressive complex 2 (PRC2) is essential for normal development by maintaining developmental gene repression. PRC2 deposits the repressive chromatin mark H3 lysine 27 tri-methyl (H3K27me3) through a read-write loop that involves direct interactions between PRC2 and H3K27me3. According to current models, the PRC2-H3K27me3 read-write loop is initiated by the PRC2 subunits JARID2 and PALI1 that mimic H3K27me3. However, it is unknown what restricts the PRC2-H3K27me3 read-write loop from spreading H3K27me3 indefinitely. To answer this question, we generated mutant mice where PRC2 subunits cannot mimic H3K27me3. Unexpectedly, the mutations led to a homeotic transformation characteristic of a Polycomb gain-of-function in vivo and the spread of H3K27me3 beyond Polycomb domains in stem cells. Collectively, we show that H3K27me3 mimicry evolved to compete against the PRC2-H3K27me3 read-write loop in a process that restrains PRC2 and restricts the spread of Polycomb domains.

Keywords

PRC2, H3K27me3, H3K27me3 mimicry, Polycomb, PALI1, JARID2, Synergy, Allostery, Epigenetics

Highlights

- H3K27me3 mimicry antagonises Polycomb function in vivo.
- JARID2 and PALI1 synergise to allosterically regulate PRC2 during development.
- H3K27me3 mimicry by JARID2 and PALI1 antagonises PRC2 in stem cells.
- JARID2 and PALI1 mimic H3K27me3 to restrict the spread of Polycomb domains.

Introduction

When cell type specific genes are repressed, they are packed into facultative heterochromatin, comprised of Polycomb domains. The repressed state of Polycomb domains is maintained by Polycomb repressive complexes [1]. Polycomb domains are defined by the repressive H3 lysine 27 tri-methyl (H3K27me3) chromatin mark, which is deposited exclusively by the histone methyltransferase Polycomb repressive complex 2 (PRC2) [2,3]. By maintaining the repressed state of cell-type-specific genes, both PRC2 [4–6] and its repressive H3K27me3 mark [7] are essential for normal development [1,8].

Broad repressive chromatin marks, such as H3K27me3, H3K9me3 and H2AK119ub, are commonly deposited by a "read-write" mechanism: a process where a chromatin modifier binds to its own product (reviewed in [9,10]). In the case of PRC2, this read-write mechanism [11] is especially potent, as it is combined with an allosteric activation [12]. Specifically, PRC2 is an allosteric enzyme that is largely inactive in its basal state. PRC2 becomes catalytically active when an allosteric effector binds to the allosteric regulatory centre in its core subunit EED [12]. The best studied allosteric effector of PRC2 is its own product — the H3K27me3 mark [12]. Upon the deposition of the H3K27me3 mark to chromatin, H3K27me3-modified histone tails can directly bind to

the PRC2 subunit EED [11] to trigger an allosteric activation of PRC2 [12]. This

mechanism leads to a positive feedback loop that is proposed to maintain the

H3K27me3 mark at Polycomb domains when cells are proliferating [11,12]. Accordingly,

pharmacological targeting of the H3K27me3 binding pocket of EED [13] or its

disruption by mutagenesis [14] leads to a loss of H3K27me3 in cells. These

perturbation experiments were important to link the allosteric regulatory activity of

PRC2 to its function in H3K27me3 deposition in cells. Yet, these experiments perturb

the allosteric regulatory activity of PRC2 entirely and, therefore, cannot separate

H3K27me3 from the other allosteric effectors of PRC2.

Indeed, three non-histone allosteric effectors of PRC2 were identified, including

JARID2 [15] and the two paralogous PALI proteins—PALI1 and PALI2 [16]. JARID2 and

PALI1 are accessory subunits of the two distinct types of holo-PRC2 complexes:

PRC2.2 and PRC2.1, respectively [17]. PALI2 was proposed to be an accessory

subunit based on sequence homology to PALI1 [18]. Mechanistically, each of the non-

histone allosteric regulators of PRC2 has an effector lysine residue that mimics H3

lysine 27 [15,16]. These effector lysines—JARID2 K116, PALI1 K1241 and PALI2

K1558—are di- and tri-methylated by PRC2. Once di- and tri-methylated, the effector

lysines bind to the allosteric center in EED to switch PRC2 into its stimulated state

[15,16]. H3K27me3 mimicry is presumed to be of significance, as it has emerged

independently at least twice throughout evolution: H3K27me3 mimicry by JARID2

K116me3 is conserved from fly to human while H3K27me3 mimicry by PALI1

K1241me3 and PALI2 K1558me3 emerged independently in vertebrate species

through convergent evolution.

How do the allosteric effectors of PRC2 work together? Early works concluded

contradictory roles for JARID2 as either an inhibitory [19,20] or stimulatory [21] accessory

subunit of PRC2. However, later biochemical works found that JARID2 enhances the

chromatin binding activity of PRC2 [22] and allosterically activates catalysis [15].

Accordingly, the prevalent paradigm is that the allosteric effector subunits of PRC2,

JARID2 and PALI1/2, are methylated by PRC2 to mimic H3K27me3 and then "jump

start" PRC2 during *de novo* deposition of H3K27me3 (reviewed in [23–32]). Once

H3K27me3 is initiated on chromatin, it allosterically activates PRC2 in a self-

reinforcing positive feedback loop that maintains H3K27me3 at Polycomb domains.

Yet, this model leaves open an unresolved conundrum: what prevents the PRC2-

H3K27me3 positive feedback loop from spreading H3K27me3 indefinitely, far

beyond polycomb domains? Moreover, while regulation of PRC2 by the H3K27me3-

mimicry activity of PALI1 and JARID2 has been supported based on enzymatic

assays *in vitro* and knockout with rescue experiments in cells, this model was never

tested *in vivo*. Hence, it is unknown what is the biological relevance of previous

models for the regulation of PRC2 by its non-histone effectors through H3K27me3

mimicry.

We set out to determine the physiological and biological relevance of allosteric

regulation by the H3K27me3-mimicking subunits of PRC2. To this end, we knocked

in allosteric-defective mutations into the endogenous loci of all known PRC2 non-

histone effector genes in mouse animals, namely *Jarid2*, *Pali1* and *Pali2*. Herein, we

show that the allosteric effector activities of the PRC2 subunits JARID2 and PALI1,

but not PALI2, synergise to support normal embryonic development through

H3K27me3 mimicry. Yet, contrary to the prevalent paradigm [23–32], allosteric defective

mouse mutants exhibited skeletal homeotic transformation characteristic of a

Polycomb gain-of-function, indicating that H3K27me3 mimicry antagonises

Polycomb function. Accordingly, the simultaneous knock-in of allosteric-defective

mutations into the endogenous *Jarid2* and *Pali1* loci led to a global gain of

H3K27me3 and the spread of Polycomb domains into intergenic regions in mouse

embryonic stem cells (ESC). We find that the allosteric regulatory activity of JARID2

and PALI1 is required to restrict H3K27me3 and prevent the spread of Polycomb

domains. We propose that H3K27me3 mimicry has repeatedly emerged through

evolution to control the spread of Polycomb domains through competition against the

PRC2-H3K27me3 read-write loop.

Results

The allosteric effector activity of JARID2, but not that of PALI1 and PALI2, is
required for mouse development

We aimed to identify the functional consequences of the allosteric regulation of

PRC2 by its non-histone effector proteins. To do this, we set out to remove the

allosteric regulatory activity of its allosteric effector subunits while keeping these

proteins otherwise functional (Figure 1A). Such separation of function (SOF)

mutations were identified in JARID2 (K116R) and PALI1 (PALI1 K1241R) within

previous biochemical studies [15,16]. We extended the mutagenesis here also to PALI2

(PALI2 K1558R) based on sequence homology (Figure 1B). These separation of

function mutations are minimal lysine-to-arginine perturbations, which only prevent

the methylation of the effector lysine and its interactions with EED to prevent the

allosteric activation of PRC2 [15,16]. We generated three separate knock-in mouse

lines, in each of them a single separation of function mutation was introduced into

one endogenous locus of a non-histone effector, either *Jarid2*, *Pali1* or *Pali2* (Figure

1C). Importantly, unlike a previously studied allosteric mutation in EED, which

disrupts the allosteric activity of PRC2 entirely [14], in these mouse lines PRC2 can still

be subjected to allosteric activation by H3K27me3 (Figure 1A, bottom).

Mice heterozygous for the JARID2 allosteric defective SOF mutation, JARID2 K116R

(*Jarid2*$^{+/SOF}$), were viable. Yet, crossing the *Jarid2*$^{+/SOF}$ mice did not produce

homozygous mutants (*Jarid2*$^{SOF/SOF}$) that survived to weaning (Figure 1D). JARID2 is

essential for mouse development [33,34], and our data indicate that its allosteric effector

activity is required for this process. While PALI1 is also essential for mouse

development [18], the allosteric defective PALI1 K1241R homozygous SOF mutant

(*Pali1*$^{SOF/SOF}$) animals were viable. Similarly, PALI2 K1558R homozygous SOF

mutant mice and PALI1 and PALI2 double homozygous SOF mutant

(*Pali1*$^{SOF/SOF}$;*Pali2*$^{SOF/SOF}$) mice were also viable (Figure 1E). These results

collectively indicate that allosteric regulation by PALI1 and PALI2 is dispensable

during development if *Jarid2* is not perturbed (more below). However, the allosteric

regulation of PRC2 by its non-histone effector JARID2 is required for mouse

development.

Allosteric-defective JARID2 leads to a Polycomb gain-of-function phenotype *in*

vivo

Past studies using knockout or loss-of-function mutations in various Polycomb group

proteins indicate that a hallmark Polycomb loss-of-function phenotype is anterior-to-

posterior homeotic transformation (reviewed in [8]). Therefore, we next assayed for

homeotic transformations in the allosteric defective *Jarid2* mouse line. After crossing

the *Jarid2*$^{+/SOF}$ mouse line, we carried out skeletal imaging on E12.5 and E14.5

embryos using immunofluorescence light sheet microscopy (Figure 1F and Figure S1A) and using alcian blue staining (Figure S1B), respectively. E14.5 was selected, as we find that at this stage most of the embryos of all of the resulting genotypes are still alive. Unexpectedly, we found that approximately half of the $Jarid2^{SOF/SOF}$ embryos underwent transformation of the first lumbar vertebral element (L1) to a thoracic identity (Figure 1F,G, Figure S1, Supplementary Movies 1-3 and Supplementary Data Table 1). This phenotype was evidenced by the presence of a well formed rib on L1—a posterior-to-anterior transformation—which is characteristic of polycomb gain-of-function [8]. We did not observe any consistent homeotic transformation phenotype in any combination of $Pali1$ and $Pali2$ mutants (Figure S2 and Figure S3C), in agreement with what was previously reported for $Pali1$ knockout mice [18]. These data indicate that the allosteric regulatory activity of JARID2 is essential for mouse development but indirectly implies that this activity antagonises Polycomb function.

The allosteric regulatory activities of PALI1 and JARID2 synergise to sustain embryogenesis

We next wanted to test for genetic interactions between $Jarid2$ and $Pali1$ or $Pali2$. To test this, we crossed mice heterozygous for both the $Jarid2$ and either $Pali1$ or $Pali2$ SOF mutations (Figure 2A,B). As expected based on our analysis of the single mutants (Figure 1D), there were no surviving mice homozygous for the $Jarid2$ SOF allele ($Jarid2^{SOF/SOF}$; Figure 2A). Importantly, synthetic lethality occurred when the heterozygous $Jarid2$ SOF mutation was present in the background of the homozygous $Pali1$ SOF mutation (genotype $Jarid2^{+/SOF}$;$Pali1^{SOF/SOF}$). Accordingly, only one mouse of the $Jarid2^{+/SOF}$;$Pali1^{SOF/SOF}$ genotype survived to weaning, compared to the expected 20 (Figure 2A, marked with a red arrow). This type of genetic interaction did not occur between $Jarid2$ and $Pali2$ (Figure 2B).

For genotypes that were identified as lethal prior to weaning, we characterised the axial skeleton at embryonic day E18.5. This analysis revealed consistent posterior-to-anterior homeotic transformations (Figure 2C, Figure S3A and Supplementary Data Table 1) and an increase in the total number of vertebrae (Figure S3B). These posterior-to-anterior homeotic transformations are in agreement with the posterior-to-

anterior transformation identified in the $Jarid2^{SOF/SOF}$ embryos (Figure 1F,G and

Figure S1) and are in agreement with a polycomb gain-of-function phenotype [35–44].

While some of the transformations appeared only in certain genotypes and with a

low penetrance, the most prevalent homeotic transformation is the one we identified

in the $Jarid2^{SOF/SOF}$ embryos (Figure 1F,G and Figure S1): a posterior-to-anterior

transformation of L1 to T13 (Figure 2C). Crucially, this phenotype was also seen in

about half of the $Jarid2$ heterozygotes when that mutation was present on the

background of at least one mutant PALI1 allele (specifically, $Jarid2^{+/SOF}$;$Pali1^{+/SOF}$ or

$Jarid2^{+/SOF}$;$Pali1^{SOF/SOF}$ in Figure 2C). This is important, as it indicates a synergy

between the allosteric regulatory activities of JARID2 and PALI1 [45], with an apparent

antagonistic effect on Polycomb function in $vivo$.

Allosteric-defective mutations in the PRC2 accessory subunits increases H3K27me3 in cells

The unexpected observation that JARID2 K116R and PALI1 K1241R mutations

exhibited polycomb gain of function phenotypes in $vivo$ indirectly suggests a PRC2

gain of function. To test this directly, we knocked-in either the JARID2 K116R or the

PALI1 K1241R homozygous mutations into the endogenous $Jarid2$ or $Pali1$ loci in

mouse embryonic stem cell (mESC) lines using CRISPR/Cas9 mediated gene

editing ($Jarid2^{SOF/SOF}$ and $Pali1^{SOF/SOF}$ cell lines, respectively; in Figure 3A). We then

knocked-in the homozygous JARID2 K116R mutation into the endogenous $Jarid2$

locus within the $Pali1^{SOF/SOF}$ cell line, to generate the homozygous double mutant

$Jarid2^{SOF/SOF}$;$Pali1^{SOF/SOF}$ (Figure 3A, bottom).

The $Jarid2^{SOF/SOF}$;$Pali1^{SOF/SOF}$ double mutant cell line exhibited a large increase in

H3K27me3 globally, as detected by flow cytometry (Figure 3B, Figure S5C) and

immunoblotting (Figure 3C, Figure S5A,B). Accordingly, quantitative ChIP-seq with a

reference exogenous genome [46] revealed substantial increments of H3K27me3 at

transcription start sites (TSS) in the $Jarid2^{SOF/SOF}$;$Pali1^{SOF/SOF}$ cells with respect to

the wild type (Figure 3D and Figure S4). The increment in H3K27me3 was observed

in PRC2-target genes, defined by their association with SUZ12 peaks in the wild type

cells, but also at non-PRC2 target genes and intergenic regions (Figure 3E and

Figure S4). The increment in H3K27me3 was accompanied by a smaller but

reproducible increment of the core PRC2 subunit SUZ12, the PRC2.2 subunit

JARID2 and the PRC2.1 subunit MTF2 (Figure 3E and Figure S4). This observation

is in agreement with previous reports that histone methyltransferase-defective PRC2

mutants exhibit reduced chromatin occupancy in cells [47–49]. This phenotype is

different from the JARID2 knockout phenotypes, which did not lead to change in

global H3K27me3 [50] and on average slightly reduced H3K27me3 at PRC2 target

genes in mECS, either undifferentiated [51] or differentiated [48]. We also demonstrated

that such a large increment of global H3K27me3 is unlikely to occur merely because

of the clonal selection (Figure S6B-D). These observations indicate that the same

JARID2 and PALI1 allosteric defective mutants that lead to a Polycomb gain-of-

function phenotype *in vivo* (Figure 1 and 2) also increase the histone

methyltransferase activity of PRC2 in cells (Figure 3).

The allosteric regulatory activity of PRC2 accessory subunits restricts the

spread of polycomb domains

In order to gain mechanistic insights on how the allosteric regulatory subunits of

PRC2 reduces its histone methyltransferase activity, we analysed the spread of

H3K27me3 (Figure 4A-D). We found that in the wild type cells H3K27me3 is mostly

restricted to CpG islands, but in the $Jarid2^{SOF/SOF}$;$Pali1^{SOF/SOF}$ cells H3K27me3 has

spread tens of kilobase pairs further away from the nucleation sites in both directions

(Figure 4A). SUZ12, MTF2 and JARID2 are still localised to the same sites on

chromatin in both the wild type and mutant cell lines (Figure 4A; antibody for the

PRC2.1 subunit MTF2 was used as there is no suitable ChIP antibody for PALI1).

This indicates that in the mutant cells, H3K27me3 is spread away from Polycomb

domains in a process that does not involve a spread of PRC2 beyond its natural

nucleation sites. In the $Jarid2^{SOF/SOF}$;$Pali1^{SOF/SOF}$ cells, H3K27me3 was increased

both inside and outside of PRC2-target TSS (Figure 3E). Yet, differential analysis of

the ChIP-Rx coverage indicated that the gain of H3K27me3 was comparatively

higher around PRC2 binding sites than inside them (Figure 4B-C). Accordingly,

substantially less ChIP-Rx H3K27me3 reads from within the mutant cell lines, in

comparison to the wild type, were mapped to SUZ12 peaks (Figure S4B).

Conversely, differential analysis of SUZ12 ChIP-Rx indicated an increase at the

PRC2 target TSS, defined using the wild type cells, but not beyond TSS (Figure 4C,

in green, and Figure 4D). Collectively, these data indicate that JARID2 and PALI1

synergise to restrain PRC2 *in vivo* by restricting the spread of H3K27me3 beyond

polycomb domains.

Discussion

Allosteric regulation is a central tenet in chromatin biology, activating chromatin

modifiers in the presence of the right effector to enable context-specific regulation

[24,52]. However, little is known about how multiple allosteric effectors synergise to

regulate the activity of a given chromatin modifier *in vivo*. Previous works revealed

that the methylation of effector lysines in JARID2 and PALI1 stimulates PRC2 *in vitro*

[15,16]. Based on knockout with rescue experiments, these works concluded that

JARID2 K116me2/3 and PALI1 K1241me2/3 stimulate, or "jump start", PRC2 to

deposit H3K27me3 *de novo* during gene repression [15,16]. This "jump start" model [23] is

largely accepted [23–32] and possibly correct. Yet, using a set of knock-in mouse lines

and stem cells we find that the main biological role of the effector lysines in JARID2

and PALI1 is the opposite to what was assumed so far: they largely restrain PRC2 *in*

vivo, not enhancing its activity. Our new data, when contrasted with previous works

[11,12,15,16], allow piecing together a model that explains how multiple allosteric

effectors synergise to regulate PRC2 in cells:

During *de novo* H3K27me3 deposition, JARID2 K116me2/3 and PALI1 K1241me2/3

may allosterically activate PRC2, as previously proposed [15,16]. Once H3K27me3 has

been deposited to Polycomb domains, PRC2 molecules that bind there use their

regulatory subunit EED to interact with the pre-deposited H3K27me3 [11]. The

interaction between H3K27me3 and PRC2 triggers the allosteric activation of PRC2

[12] and, at the same time, increases the affinity of PRC2 to chromatin [11]. Now, the

allosterically activated PRC2 introduces more H3K27me3 to chromatin. Importantly,

this process can take place even if not all the JARID2 and PALI1 molecules in the

cell are methylated, as these proteins have a strong chromatin binding activity

disregarding their effector lysines [16,22,48,53]. Since the chromatin-bound PRC2 is now

allosterically activated, it can occasionally also methylate JARID2 K116 [15] or PALI1

K1241 [16]. The affinity of the effector lysines JARID2 K116me2/3 and PALI1

K1241me2/3 for EED is about 3- to 5-fold higher than that of the H3K27me3 histone

tails to EED [15,16]. Moreover, JARID2 and PALI1 are already part of the holo-PRC2

complex [15,16], which increases the effective concentration of JARID2 K116me2/3 and

PALI1 K1241me2/3 around EED. Hence, as soon as an effector lysine in JARID2 or

PALI1 becomes di- or tri-methylated, it blocks the allosteric regulatory site in EED

and prevents H3K27me3 tails from binding there. JARID2 K116me2/3 and PALI1

K1241me2/3 can still allosterically activate PRC2, although they do not directly

contribute to chromatin binding [15,16]. Conversely, the interactions between

H3K27me3 and EED contribute to both allosteric activation [12] and chromatin binding

[11], because H3K27me3 is part of the chromatin. This likely makes H3K27me3 a

stronger activator of PRC2 compared to the effector lysines in PALI1 and JARID2.

This model fits with the antagonistic activity of the JARID2 and PALI1 effector lysines

in cells (Figure 3-4) and *in vivo* (Figure 1-2) and also with their stimulatory activity *in*

vitro [15,16].

We propose that JARID2, PALI1 and H3K27me3 work together to maintain a healthy

level of H3K27me3 on chromatin. A high level of H3K27me3 would increase the

instances where PRC2 is allosterically activated. Sustained allosteric activation of

PRC2 would eventually lead to the di/tri-methylation of JARID2 and PALI1 and the

subsequent interference with the PRC2-H3K27me3 feedforward loop. This

mechanism is unlikely to take place in active genes, where the H3K4me3 and

H3K36me3 histone mark inhibit PRC2 [54] and H3K4me3 may compete against

JARID2 K116me2/3 for binding to EED there [55]. When JARID2 K116 and PALI1

K1241 are mutated, the PRC2-H3K27me3 feedforward loop cannot be restrained,

leading to increased H3K27me3 at CpG islands (Figure 3) and eventually to its

spread away from Polycomb domains (Figure 4).

Intriguingly, mutations in and around JARID2 K116 and PALI1 K1241, including

recurrent mutations, have been observed in cancer. Examples are JARID2 K116T [56]

and K116N [57] in carcinoma and PALI1 P1243S [58–60] and P1243A [59] in melanoma

[61,62]. If these cancer-associated mutations interfere with the binding of the JARID2 or

PALI1 effector lysines to EED, then they may confer a PRC2 gain-of-function, as

with JARID2 K116R and PALI1 K1241R (Figure 1-4). PRC2 gain-of-function is

commonly associated with poor prognosis in cancer [63] and in some cases can be

targeted by the pharmacological inhibition of PRC2 [64].

We show that the allosteric regulatory activity of the accessory subunits of PRC2 is

required for embryonic development and that allosteric-defective JARID2 and PALI1

lead to homeotic transformations (Figure 1-2), a hallmark Polycomb gain-of-function

phenotype. Anterior-to-posterior axial skeletal transformations have been identified

as a loss-of-function phenotype in mice for various Polycomb group proteins [35–44]. To

our knowledge, no homeotic transformation has been reported in studies where *Pali1*

[18] or *Jarid2* [34,65–67] were knocked-out in mice. Perinatal or embryonic lethality was

observed in the case of both PALI1 [18] and JARID2 [34] knock-out (reviewed in [33]), and

the latter also led to organ or skeletal development defects [34,65–67]. However, it was

impossible to link these previously-identified phenotypes directly to Polycomb

function, especially in the case of PALI1 that also binds to the H3K9me2

methyltransferase G9A [18]. Moreover, regions of JARID2 and PALI1, external to their

effector lysines, enhance the affinity of PRC2 for nucleosomes [16,22] and are required

for chromatin binding by PRC2 [48,53]. Therefore, knockout experiments cannot link a

specific biochemical function of PALI1 or JARID2 to the observed phenotype. Our

results indicate that the allosteric regulatory activities of JARID2 (Figure 1) and

PALI1 (Figure 2) are required for normal development and reveal a synergy between

them (Figure 2) in antagonising Polycomb function (Figures 2-4).

PALI1 is a vertebrate-specific protein [18] while JARID2, including K116, is conserved

from fly to human [15]. Their mechanistic resemblance *in vitro* led to the proposal that

PALI1 emerged through a convergent evolution to mimic JARID2 at the molecular

level [16]. Our discovery of genetic interactions between JARID2 K116 and PALI1

K1241 indicate that JARID2 and PALI1 synergise to allosterically regulate PRC2 *in*

vivo through H3K27me3 mimicry. We propose that H3K27me3 mimicry by JARID2

evolved to restrict the read-write PRC2-H3K27me3 feedforward loop of PRC2.2 in

metazoans while PALI1 emerged in vertebrate species to carry out this function in

the context of PRC2.1. Collectively, JARID2 and PALI1 synergize to buffer the

PRC2-H3K27me3 feedforward loop, allowing PRC2 to sustain H3K27me3 at

Polycomb domains without expanding beyond them.

Acknowledgements

We thank the support of the Monash FlowCore, Monash University Palaeontology Lab, and the MASSIVE HPC facility. The mutant mouse animals and cells were produced via CRISPR genome editing by the Monash Genome Modification Platform (MGMP), Monash University, as a node of Phenomics Australia. Phenomics Australia is supported by the Australian Government Department of Education through the National Collaborative Research Infrastructure Strategy, the Super Science Initiative and the Collaborative Research Infrastructure Scheme. S.C.A. was supported through an Australian Government Research Training Program (RTP) Scholarship. C.D. was an EMBL-Australia Group Leader and is a Sylvia and Charles Viertel Senior Medical Research Fellow and acknowledges support from the ARC (DP190103407) and the NHMRC (APP1162921, APP1184637, APP2011767 and APP2020900). This research was funded partially by the Victoria State Government through mRNA Victoria (C.D.).

Authors contributions

S.C.A. and C.D. conceptualised the project, S.C.A, M.B., Y.C, M.M., J.M., Z.L., S.N., E.M. carried out experiments and investigated, S.C.A., M.B., V.L. and C.D. managed mouse colonies and animal ethics, J.R. generated cell lines and mouse animal lines, Z.L. produced reagents, E.H. carried out bioinformatic analyses, Y.C. and E.M. provided training and input into experimental design, C.D. and E.M. supervised, S.C.A. and C.D. wrote the original draft and all authors reviewed and edited the manuscript.

Declaration of interests

The authors declare no conflict of interest.

Data and materials availability

Next generation sequencing data (ChIP-Rx) are available under GEO accession number GSE278724.

Figures

383 **Figure 1: Allosteric defective JARID2, but not PALI1 and PALI2, cause**

384 **Polycomb gain of function *in vivo*.** A. Schematic diagram of allosteric regulation of

385 PRC2. Top: PRC2 methylates an effector lysine—JARID2 K116, PALI1 K1241,

386 PALI2 K1558, or H3K27—that subsequently binds to the allosteric regulatory centre

in the PRC2 core to trigger an allosteric activation of methyltransferase. Bottom:

Mutating effector lysines in the accessory subunits of PRC2 prevents allosteric

activation *in vitro* (dashed red arrow) [15,16], but the biological consequences *in vivo*

are unknown. B. Alignment of portions of the human and mouse JARID2, PALI1 and

PALI2 protein sequences, where the effector lysines are highlighted in yellow. C.

Three genetically modified mouse animal lines were generated using CRISPR/Cas

knock in of separation-of-function (SOF) mutations into the endogenous *Jarid2*, *Pali1*

and *Pali2* genes, as indicated. Each of these mutations include a single lysine-to-

arginine mutation of the effector lysine (highlighted in yellow in (B)). D. The

percentage of observed (black) and expected (grey) pups that survived to 21 days

postnatal as the result of a *Jarid2$^{+/SOF}$* X *Jarid2$^{+/SOF}$* crossing. The number of pups

from each genotype is indicated. The expected number of pups (adjacent to the grey

bars) is rounded to the nearest integer. The total number of pups is indicated by n.

The p-value was calculated using a chi-squared test. E. *Pali1$^{+/SOF}$* and *Pali2$^{+/SOF}$*

mouse lines were crossed to yield the *Pali1$^{+/SOF}$;Pali2$^{+/SOF}$* line. Subsequently,

Pali1$^{+/SOF}$;Pali2$^{+/SOF}$ X *Pali1$^{+/SOF}$;Pali2$^{+/SOF}$* crossing took place and the observed and

expected percentage of pups are indicated, as in (D). F. Light sheet fluorescence

microscopy of E12.5 embryos immunostained for SOX9. For each genotype, the

following are shown: 3D reconstruction of a whole embryo, lateral view (first row),

and a separate scan of the same embryo limited to only the region surrounding the

thoracic vertebrae, raw data (second row) and 3D rendering (third row). In the

second and third rows, only the ribs on one side of the embryo are shown. A second

independent comparison is shown in Figure S1A. G. Analysis of homeotic

transformation in *JARID2 K116R* mutant embryos. The frequency of homeotic

transformations shown in (F) was determined in E14.5 embryos stained with alcian

blue. The number of embryos that were examined from each genotype are indicated

with n. A schematic diagram representing the mouse vertebral column is shown,

where the transformed vertebra is coloured red, with an arrow indicating the direction

of the transformation. The number of embryos in which the transformation was

observed is indicated beside the corresponding vertebra (in red, left number within

the parentheses) and the total number of embryos from the same genotype where

that vertebra was visible (in red, right number within the parentheses).

Figure 2: Synergy between allosteric defective JARID2 and PALI1 during mouse embryogenesis. A. Double mutant mice were created by crossing between PALI1 and JARID2 single mutant mice, as indicated. The resulting PALI1 and JARID2 double-heterozygous mutant animals were then crossed with animals of the same genotype. The bar plot represents the percentages of observed (black) and expected (grey) mice of each genotype that survived to 21 days postnatal resulting from these crosses. For each genotype, the observed and expected number of pups are indicated to the right of the corresponding bar (expected number of pups are rounded to the nearest integer). The total number of pups that survived to day 21 postnatal is indicated by n. The p-value was calculated using a chi-squared test. SOF refers to a separation of function mutant: JARID2 K116R or PALI1 K1241R. B. Same as (A), except that PALI2 and JARID2 double mutants were assayed. C.

Homeotic transformations in the vertebral column of E18.5 embryos resulting from the *Pali1+/SOF;Jarid2+/SOF x Pali1+/SOF;Jarid2+/SOF* crossing. Top: A schematic diagram representing the mouse vertebral column, including the occipital bone up to the 4th caudal vertebra (no transformations were observed in the caudal vertebrae). Vertebrae in which homeotic transformation or malformation were observed with a frequency of at least 50% are coloured red, or black if it was observed with a frequency of less than 50%. The direction of homeotic transformations are indicated by an arrow. The number of embryos that the transformation was observed in is given next to the corresponding vertebra. Bottom: Representative photos of the T12-L2 vertebrae showing the L1 to T13 transformation, indicated with arrows.

Figure 3: Allosteric defective JARID2 and PALI1 lead to increased H3K27me3 in mESC. A. Two genetically modified mESC lines were generated using CRISPR/Cas knock-in of the indicated separation-of-function (SOF) mutations into the endogenous *Jarid2* and *Pali1* loci. Subsequently, a double mutant cell line was generated by knocking in the JARID2 K116R mutation into the *Jarid2* locus within the PALI1 K1241R cells. All knock-ins were homozygous. B. Fluorescence-activated

cell sorting (FACS) on cells possessing mutations or Tazemetostat treatment as

specified. The bar charts show the mean of the median fluorescence signal and error

bars represent the standard error across three independent biological replicates that

were harvested on different days. Statistical significance was calculated using a one-

way ANOVA with a post-hoc Tukey's test to correct for multiple comparisons. Non-

significant comparisons are not shown. C. Western blots on whole cell lysate of cells

possessing the indicated mutations or Tazemetostat treatment using antibodies as

specified. See Figure S5A,B for two more replicates and a densitometry analysis. D.

Quantitative ChIP-seq (ChIP-Rx) was carried out using cell lines and antibodies as

indicated and heatmaps are centred on all transcription start sites. E. Average ChIP-

Rx signal profiles at all transcription start sites at PRC2-target genes, non-PRC2-

target genes, or intergenic regions as indicated. Colour keys are presented in the

bottom plots. For a second independent ChIP-Rx replicate that was carried out on a

different day see Figure S4A,B.

463
Figure 4: The allosteric regulatory activity of JARID2 and PALI1 is required to
restrict the uncontrolled spread of polycomb domains. A. Gene track over a
representative locus showing ChIP-Rx signals from cell lines and antibodies as
indicated. PRC2 nucleation sites are defined by the SUZ12 peaks in the wild-type
cells and are highlighted in green. B. Gene tracks presenting the ratio of H3K27me3
(top) or SUZ12 (bottom) in the PALI1 K1241R and JARID2 K116R double mutant

cells compared to the wild type. Colour code is as in (A). C. The average change in H3K27me3 or SUZ12 at all PRC2 target sites. Data is plotted as a log2 ratio of reads in the PALI1 K1241R and JARID2 K116R double mutant to wild type. Antibodies are indicated. Different ranges around TSS are presented in the top and bottom plots, as indicated. D. Left: A schematic diagram showing the change in H3K27me3 localisation in the mutant with respect to the wild type cells. Middle: The H3K27me3 ChIP-Rx read density inside and within 30 kbp outside of SUZ12 peaks (highlighted within the illustration on the left in green and yellow, respectively). The median is shown with a dashed line. Right: The change in the H3K27me3 ChIP-Rx read density inside and outside of peaks in the mutant relative to the wild type cells. Statistical significance was assessed by a Mann-Whitney U test and p values are indicated. E. Schematic diagram showing a model for allosteric regulation of PRC2 by JARID2 and PALI1. In wild type cells, PRC2 methylates JARID2 K116 and PALI1 K1241, which then allow them to compete against H3K27me3 for binding to the regulatory center in PRC2. This process breaks the PRC2-H3K27me3 read-write positive feedback loop to retain normal levels of H3K27me3 at CpG islands. In the allosteric defective mutant cell line (right), the mutant PALI1 and JARID2 can no longer compete with H3K27me3. As a result, the read-write loop is no longer controlled, leading to high levels of H3K27me3 at CpG islands and a spread of H3K27me3 beyond Polycomb domains.

Supplementary information

Supplementary Figure 1: *Jarid2$^{SOF/SOF}$* **causes posterior to anterior homeotic**

transformation in mouse. A. Light sheet fluorescence microscopy of the region

surrounding the thoracic vertebrae in E12.5 embryos stained for SOX9. A 3D

rendering from surface segmentation is shown in the second row. The homeotic

transformation L1 to T13 is indicated with an arrow. Presented is another

independent comparison from the same experiment as in Figure 1F. B.

Representative images of the E14.5 embryos corresponding to the analysis in Figure

1G. All embryos in this analysis were stained with alcian blue and imaged in a

dissecting microscope with white light. Representative embryos from each of the

genotypes are shown, where the homeotic transformation L1 to T13 is indicated with

an arrow.

Supplementary Figure 2: PALI1 K1241 and PALI2 K1558 are dispensable for

skeletal patterning. A. Sanger sequencing traces confirming the correct knock-in

heterozygous mouse lines that were generated in this study. The codon

corresponding to the separation of function mutation is annotated below. B. No

homeotic transformation or other defects were observed in embryonic day 18.5

(E18.5) skeletons resulting from *Pali1^{+/SOF}* x *Pali1^{+/SOF}* crossing. The number of

embryos that were examined from each genotype is indicated by n. A representative

photo of each genotype is shown. C. The same as (B), for *Pali1^{+/SOF};Pali2^{+/SOF}* x

Pali1^{+/SOF};Pali2^{+/SOF} crossings. Vertebrae in which homeotic transformation or

malformation were observed are coloured black, with the direction of homeotic

transformations indicated by an arrow. The number of embryos that the

transformation was observed in is given next to the corresponding vertebra.

Supplementary Figure 3: The synergy between allosteric defective JARID2 and
PALI1 during mouse embryogenesis. (A) The photos shown in this figure
correspond to the cropped photos shown in Figure 2C. An exception is the
Pali1+/SOF;Jarid2+/SOF genotype, which was photographed using a microscopic lens
limited only to the area of interest in Figure 2C, whereas a photograph of the
complete embryo that was recorded separately is shown here, positioned similarly.
Genotypes are indicated. B. Total vertebral number for the embryos that were part of

the experiment and are shown in Figure 2C. Each dot on the graph represents the

total vertebral number for one embryo. Only embryos for which all vertebrae could be

confidently counted were included on this plot. The median for each genotype is

shown by a red line. Genotypes are indicated. A Kruskal-Wallis test with Dunn's test

for multiple comparisons was used to test for significance between all combinations

of genotypes and p values are presented only where differences are statistically

significant. C. Same as (B), except that the embryos were part of the experiment

shown in Supplementary Figure 1C.

Supplementary Figure 4: Allosteric defective JARID2 and PALI1 lead to a genome wide increase of H3K27me3 in mESC. A. Second replicate of the ChIP-Rx shown in Figure 3D, where heatmaps are centred on all transcription start sites. B. Same data set as in (A), where average ChIP-Rx signal profiles are shown over all transcription start sites at PRC2-target genes, non-PRC2-target genes, or

541   intergenic regions as indicated. C. Bar chart representing the percentage of ChIP-

542   Rx-normalised H3K27me3 reads that aligned to genomic regions within SUZ12

543   peaks in the wild type and mutant cell lines within two independent ChIP-Rx

544   replicates.

Supplementary Figure 5: Allosteric defective JARID2 and PALI1 lead to increased global H3K27me3 in mESC. A. Two more western blot replicates corresponding to Figure 3C. B. Bar charts representing the mean densitometry values of all three western blot replicates shown in (A) and in Figure 3C. Shown are densitometry values that were normalised to the average densitometry values of all

cell lines in the experiment, excluding the Tazemetostat treated sample. C. Flow

cytometry data corresponding to Fig 3B. Indicated are individual fluorescence values

for each cell that participated in the experiments. Cell lines and Tazemetostat

treatments are indicated. The medians that are presented in Fig 3B were derived

from this analysis and are presented here with horizontal black lines. Each replicate

is plotted separately, and three independent replicates of each sample are grouped

together.

Supplementary Figure 6: Clonal selection of wild type mESCs is insufficient to cause a global gain of H3K27me3. A. Sanger sequencing traces confirming the correct knock-in homozygous mutant mESC lines generated in this study. The codon corresponding to the separation of function mutation is annotated below. B. A schematic diagram showing how the cells were treated to assess for a clonal effect: The parental wild type mESCs were seeded at a low density on a culture dish, and incubated until colonies grew from the individual cells. Ten colonies were picked and transferred into separate dishes. The resulting ten clonal wild type cell lines were all expanded and then assayed for global H3K27me3 by flow cytometry. C. Flow cytometry of the ten clonal wild type cells (see scheme in (B)) stained with anti-

H3K27me3 antibodies. The "parental" sample represents the same parental wild

type cell line used in other experiments and also for clonal selection here.

Tazemetostat was used as a negative control. The bar charts show the mean of the

median fluorescence signal and error bars represent standard error across three

independent biological replicates that were harvested on different days. Statistical

significance was calculated using a one-way ANOVA with a post-hoc Tukey's test to

correct for multiple comparisons. Only comparisons involving the uncloned cells are

shown. The p-values of non-significant comparisons are not shown. D. Scatter plots

of the H3K27me3 flow cytometry values measured from each cell within the cell lines

shown in (B). This data was used to calculate the median values (marked by a line)

that are presented in (C). X-axis labels are as in (C), except that three independent

replicates are presented side-by-side for each of the wild type clonal cell lines.

Supplementary Figure 7: Gating strategy used for flow cytometry analysis.
Representative scatter plots showing the gating strategy used for measuring global
H3K27me2/3 by flow cytometry. A gate was used to include only the single cells (A),
and another gate to include the intact mESCs (B). Only the events that fell within
both gates were considered when measuring the median H3K27me2 or H3K27me3
for that sample.

Supplementary Data Table 1: Embryonic skeleton phenotypes. A spreadsheet

showing the phenotype along the axial skeleton of each embryo included in the

experiments shown in Figures 1G and 2C and Figures S1B, S2B,C and S3. Scoring

of the skeletons resulting from unique crossings of genotypes are provided in

separate tabs. A key is shown below the table in each tab.

Supplementary Movie 1: Wild type E12.5 embryo immunostained for SOX9. A

movie showing the complete light sheet fluorescence microscopy 3D reconstruction

in Figure 1F. First, the complete signal is shown without removing any non-relevant

regions. Next, the non-relevant signal is removed, as in the second row of Figure 1F.

Next, a rendering of both of the pairs of ribs are imposed over the previous model,

and finally only the side used to generate the bottom row in Figure 1F is shown.

Supplementary Movie 2: *Jarid2+/SOF* E12.5 embryo immunostained for SOX9. As

Supplementary Movie 1, except that a *Jarid2+/SOF* E12.5 embryo was imaged.

Supplementary Movie 3: *Jarid2SOF/SOF* E12.5 embryo immunostained for SOX9.

As Supplementary Movie 1, except that a *Jarid2SOF/SOF* E12.5 embryo was imaged.

Methods

Amino acid numbering system

For compatibility with previous works, the allosteric effector lysines in mouse cells

and animals are named based on the human numbering system. Accordingly,

references to PALI1 K1241, PALI2 K1558 or JARID2 K116 (PALI1 K1393, PALI2

K1543 or JARID2 K116, respectively, in mouse) are made in this way, using the

human numbering system, regardless of if these amino acids are in the human or

mouse proteins. The corresponding sequences of the mouse and human proteins

are shown in Figure 1B, where the human and mouse sequences were aligned using

EMBOSS needle [68,69].

Mice genotyping and ethical approval

All mice used in this study were maintained on a C57BL/6J background. The genetic

background was confirmed by Transnetyx using miniMUGA. Mouse lines possessing

mutations in multiple genes were generated by crossing the relevant single mutant

lines. Genotyping was performed by Transnetyx using real-time PCR. All animal

breeding and experimentation was conducted following the Australian National

Health and Medical Research Council Code of Practice for the Care and Use of

Animals for Scientific Purposes guidelines for housing and care of laboratory animals

and performed in accordance with Institutional regulations after pertinent review and

approval by Monash University Animal Ethics Committee under project numbers

36532 and 26125. Mice were bred and housed in specific-pathogen-free (SPF)

conditions at the Monash Animal Research Platform (MARP) and Animal Research

Facility (ARL) at Monash University.

Knock-in of point mutations to mice

PALI1 K1241R, PALI2 K1558R and JARID2 K116R were knocked-in using

CRISPR/Cas with homology-directed repair, using sgRNAs that were designed to

target the following sequences: TCCTAGCAGGAATGGCTCCA,

AAGGAGAAGAGACTGCAAATT and GGCCCAGGCTGCAAGCACAA, respectively.

To avoid embryonic lethality resulting from homozygous knock-in of the mutation,

two repair templates were used for each genome modification process: one that

contains an MspA1I restriction site but without amino acid change (Repair Template

1), and one that contains MspA1I and HaeII restriction sites, and also the amino acid

change (Repair Template 2).

The appropriate sgRNA and Cas protein were incubated with the guide RNA to form

a ribonucleoprotein (RNP) complex. Cas9 used the PALI1 K1241R and JARID2

K116R knock-in and Cas12a used for the PALI2 K1558R knock-in. The RNP

complex and repair template were then electroporated or microinjected into

C57BL/6J zygotes at the pronuclei stage and transferred into the uterus of pseudo

pregnant F1 females. The following conditions were used for each gene:

PALI1 K1241 knock-in: Repair Template 1 sequence was

CCTTCCACCATGTACCCTAGTTCACTACAGGCAGAACGCTTGAAAAAACATTTA

AAGAAATTCCCCGGGGCCATTCCTGCTAGGAATAATTGGAAGACACAGAAGCTA

TGGGCTAAACTACGAGAGAATCCTGA. Repair Template 2 was

AGTTTCCTCCTTCCACCATGTACCCTAGTTCACTACAGGCAGAACGCTTGAAAA

AACATTTAAAGCGCTTCCCCGGGGCCATTCCTGCTAGGAATAATTGGAAGACAC

AGAAGCTATGGGCTAAACTACGAGAGAATCCTGA. Delivery was done using the

microinjection of Cas9 nuclease (IDT Alt-R S.p. Cas9 Nuclease V3; 10 ng/mL),

sgRNA (10 ng/mL) and repair templates (30 ng/mL).

PALI2 K1558R knock-in: Repair Template 1 was

TTTAAGAGTGGTGAGCCAACCATTTTATGTAGTATATGAAATTTCCCAGCTGGTT

TTGAAGGAGAGCTCACTGCAAATTTTTTAAAGTGCTTCCTGAAGGGGCTGGCAT

TAAATCCACAGCATTTCACCCCTAACTTAATGCCCT. Repair Template 2 was

TTTAAGAGTGGTGAGCCAACCATTTTATGTAGTATATGAAATTTCCCAGCTGGTT

TTGAAGGAGAGCTCACTGCAAAACGTTTAAAGTGCTTCCTGAAGGGGCTGGCA

TTAAATCCACAGCATTTCACCCCTAACTTAATGCCCT. Delivery was done using

the microinjection of Cas12a nuclease (Alt-R A.s. Cas12a (Cpf1) Ultra; 100 ng/mL),

crRNA (100 ng/mL) and repair templates (50 ng/mL).

JARID2 K116R knock-in: Repair template 1 was

CTCCAGTCTCTGCTTAATTTAGTTTTACATTTCTTTTGTTGTTGCAGGCCCAGGC

TGCAAGCACAGCGGAAGTTTGCTCAATCTCAGCCGAATAGTCCCAGCACAACT

CCAGTGAAGATAGTGGAGCCTTTG. Repair template 2 was

CTCCAGTCTCTGCTTAATTTAGTTTTACATTTCTTTTGTTGTTGCAGGCCCAGGC

TGCAAGCACAGCGGCGCTTTGCTCAATCTCAGCCGAATAGTCCCAGCACAACT

CCAGTGAAGATAGTGGAGCCTTTGCTACC. Delivery was done using the

microinjection of Cas9 nuclease (IDT Alt-R S.p. Cas9 Nuclease V3; 10 ng/mL),

sgRNA (10 ng/mL) and repair templates (30 ng/mL) or electroporation of Cas9

nuclease (3.8 mM), sgRNA (4.2 mM) and repair templates (10 mM). For all knock-

ins, the genotypes of the founder and F1 mice were confirmed by sanger sequencing

(Figure S2A).

Analysis of Mendelian inheritance

Mice possessing the desired genotypes were crossed. The resulting pups that

survived to 21 days postnatal were genotyped, and the frequency of each genotype

among them was tallied. Chi-square tests were conducted using Excel for the pups

resulting from the crossings to test for statistical significance between the observed

and expected ratios of genotypes.

Skeleton preparation and imaging

Mice heterozygous for each of the mutations of interest were time-mated, and embryos were harvested at embryonic day 14.5 or 18.5 (E14.5 or E18.5, respectively).

E18.5 embryos were prepared as described previously [70]. Specifically, the E18.5 embryos were skinned, then incubated in 95% (v/v) ethanol for 2 days, followed by 2 days in acetone. The embryos were then stained for 1-2 days in a solution containing 150 µg/mL alcian blue (Sigma 05500), 50 µg/mL alizarin red (Sigma A5533), 5% (v/v) glacial acetic acid and 60% (v/v) ethanol. The embryos were then cleared in 1% (w/v) KOH for 3-5 days, then transferred through a graded series of solutions in which a gradually increasing proportion of the KOH was replaced with glycerol (25%, 50%, 75% then 100% glycerol), with one day between each solution change. All incubations took place at room temperature with rocking.

E14.5 embryos were stained as previously described [71] with some minor changes. Specifically, the E14.5 embryos were incubated while rocking in 70% (v/v) ethanol at 4°C overnight, then at room temperature in 95% (v/v) ethanol for 1 hour, followed by acetone overnight. The embryos were stained in a solution containing 0.03% (w/v) alcian blue, 80% (v/v) ethanol and 20% (v/v) acetic acid, for 4 hours. The embryos were cleared over 2-3 days in 0.5% (w/v) KOH, then transferred through a graded series of solutions in which a gradually increasing proportion of the KOH was replaced with glycerol (25%, 50%, 75% then 100% glycerol), with one day between each solution change. All incubations took place at room temperature with rocking.

Images of E18.5 embryos were acquired using a Vision Dynamic BK Lab Imaging System equipped with a Canon 5D Mark II with a 100 mm macro lens, or 7D Mark II with Mitutoyo 5x microscope lens. Z-stacked images were captured at multiple focal planes through the embryo, then combined to create a composite photograph using Zerene Stacker. Images of E14.5 embryos were taken with a ZEISS Stemi 508 microscope equipped with a ZEISS Axiocam 208 color camera. The images were set to black and white, and brightness and colour levels for each image were adjusted in Photoshop separately.

3D light sheet immunofluorescence microscopy of whole embryos

To visualise the developing skeleton, *in toto* immunofluorescence was performed as described previously [72]. E12.5 embryos were dissected out of uterine horns in cold PBS and fixed overnight in PBS with 4% PFA at 4°C with rocking. The following day, embryos were washed in PBS then dehydrated by transferring through a series of PBS-methanol solutions, each containing a higher concentration of methanol (20%, 40%, 60%, 80%, 100% methanol) for one hour each, then frozen in 100% methanol until further processing. Prior to staining, embryos were bleached in methanol with 7.5% H2O2 under white light at 4°C with rocking, and rehydrated by transferring through a series of PBS-methanol solutions, each containing a lower concentration of methanol (80%, 60%, 40%, 20%, 0% methanol) for one hour each.

The rehydrated embryos were permeabilised in PBS with 0.2% Triton X-100, 20% DMSO and 2.3% glycine, then blocked in PBSGT (PBS with 0.2% gelatin and 0.5% Triton X-100). Permeabilisation and blocking were performed at 37°C with rocking for 48 hours.

The embryos were incubated with anti-SOX9 (Merck AB5535, 1:100) and anti-TUBB3 (BioLegend 801201, 1:500) primary antibodies diluted in PBSGT at 37°C in the dark with rocking for at least 5 days. After the primary antibodies were removed, the embryos were washed at least four times in PBSGT for one hour per wash. The embryos were then incubated with anti-rabbit (Thermo Scientific A-31572, 1:500) and anti-mouse (Thermo Scientific A-21241, 1:500) secondary antibodies for at least four days at 37°C in the dark with rocking. After the secondary antibodies were removed, the embryos were washed at least four times in PBSGT for one hour per wash.

Next, the embryos were dehydrated again by transferring through a series of PBS-methanol solutions, each containing a higher concentration of methanol (20%, 40%, 60%, 80%, 100% methanol) for one hour each at room temperature in the dark, rotating. This was followed by another wash in methanol for one hour, then again overnight. Clearing was performed with a 2:1 ratio of dichloromethane (DCM) to methanol for 3 hours in the dark, rotating, followed by 100% DCM for one hour, and finally submerged into dibenzyl ether (DBE) without further agitation to avoid excessive oxidation. Embryos were imaged within one week of clearing.

Images were acquired on an Ultramicroscope 2 (La Vision, Miltenyi Biotec) equipped

with a 2X objective and a 1.6X magnification in DBE. Single side illumination, with 3

convergent light sheets set at a width of 80% and a numerical aperture of 0.134

Nyquist was used to scan the sample. Dynamic focus was used across the X axis,

with the proprietary blend algorithm in Imspector Pro 5.0 (La Vision, Miltenyi Biotec).

Image processing was performed in Imaris 10.2 (Bitplane, Oxford Instruments). The

SOX9 signal for the rib cage and spine was isolated by manually subtracting signal

from other non-relevant SOX9-expressing tissues (limbs, heart, skin, gonadal ridges)

using the surface function in Imaris. 3D rendering of the ribs was generated using

the surface function in Imaris. Images were acquired using the Snapshot function of

Imaris, and the brightness was adjusted in ImageJ. Movies were generated using the

Animation menu of Imaris.

Cell culture

Bruce-4 mouse embryonic stem cells were cultured at 37 °C with 5% CO_2 in DMEM

growth medium, supplemented with 15% FBS (Scientifix FBSAU-2007A), 1%

penicillin-streptomycin (Thermo Scientific 15140122), MEM Non-Essential Amino

Acids Solution (Thermo Scientific 11140050), Glutamax™ Supplement (Thermo

Scientific 35050061), 0.1 mM 2-Mercaptoethanol (Merck M3148) and 1:1000

homemade LIF. The cells were grown on gelatinised plates, passed every 2-3 days,

and the media was replaced daily. Bruce-4 cells (derived from C57BL/6J-Thy1.1

strain, kindly gifted to Monash Genome Modification Platform by Colin Stewart) were

authenticated by ATCC using short tandem repeat (STR) analysis. Cells were tested

periodically for mycoplasma contamination using PCR [73].

Knock-in of point mutations to mESCs

PALI1 K1241R, PALI2 K1558R and JARID2 K116R were knocked-in using

CRISPR/Cas with homology-directed repair, using the sgRNAs that were designed

for the following targets: TCCTAGCAGGAATGGCTCCA,

AAGGAGAAGAGACTGCAAATT and GGCCCAGGCTGCAAGCACAA, respectively.

Lipofectamine CRISPRMAX was used to transfect mESCs with the appropriate

sgRNA, Cas9 protein fused with GFP and repair template with the same sequence

that was indicated above, for mouse knock-in. Clonal cells were selected using flow

cytometry to sort individual GFP-positive cells. In all cases, clonal cells were

expanded and screened for the desired genotype using PCR and restriction enzyme

digestion. Positive clones were validated by sanger sequencing (Figure S6A). Cell

lines were occasionally genotyped by Transnetyx using real-time PCR.

Detection of H3K27me2/3 using flow cytometry

For a negative control, wild type cells were grown in 10 µM Tazemetostat

(Selleckchem S7128) for 7 days. Other cell lines were cultured without

Tazemetostat. To collect cells, the culture media was removed, replaced with trypsin

(Thermo Scientific 25200-056) solution and cells were incubated in trypsin at room

temperature for <5 minutes. The detached cells were resuspended in growth media

and collected by centrifugation for 5 minutes at 300 g. The cell pellet was then

resuspended in 4% paraformaldehyde solution in PBS (Santa Cruz sc-281692) and

incubated for 15 minutes at room temperature to fix the cells. The cells were then

washed twice in PBSTX (PBS with 0.1% (v/v) Triton X-100 (Sigma X100)), and

resuspended in PBSTX to be stored for up to a month at 4 °C before further

processing.

For the antibody staining, 2×10^5 cells per sample were resuspended in blocking

buffer (PBS with 0.1% (v/v) Triton X-100, 1% (w/v) BSA (Sigma 10735078001) and

5% (v/v) FBS) and blocked for 15 minutes.

The cells were centrifuged and resuspended in blocking buffer with H3K27me2

(Abcam ab24684, 1:5000) or H3K27me3 antibody (Cell Signalling #9733S, 1:500).

After 1 hour of incubation, the cells were then washed with PBSTX, centrifuged at

1000 g for 2-5 minutes and the pellet resuspended in blocking buffer with secondary

antibody (Thermo Scientific A32795, 1:15000) and 1 µg/mL DAPI (Sigma, D9542-

1MG) for 45 minutes in the dark. The cells were then washed with PBSTX,

centrifuged at 1000 g for 2-5 minutes, and resuspended in PBS with 10% FBS and

615 µM EDTA, and ran through a cell strainer (Falcon #352235). The cells were then

analysed on a BD LSRFortessa™ X-20 Cell Analyzer, using the R670-A detector.

Three independent biological replicates were initiated on three different days and

were carried out as described above. The data was gated for intact single cells

(Figure S7) using FlowJo and plotted using GraphPad Prism.

Immunoblotting

To collect cells, the culture media was removed, replaced with trypsin and incubated in the trypsin solution for <5 minutes at room temperature. The detached cells were resuspended in growth media and collected by centrifugation for 5 minutes at 300 g. The cells were then washed 3 times in PBS and lysed in Laemmli buffer (1% (v/v) SDS, 12.5% (v/v) glycerol, 35 mM Tris pH 7.5 at 25°C, 0.01% (w/v) bromophenol blue, 5 mM MgCl2, 1% (v/v) 2-mercaptoethanol) and 25 U/mL Benzonase (Merck #70746), and then heated to 95 °C for 5-10 minutes before they were loaded on a gel.

The protein samples were run on a 8-16% gel (Thermo Scientific XP08160BOX), then transferred to a nitrocellulose membrane (GE Life Sciences #10600002). Membranes were incubated in blocking buffer (Thermo Scientific #37539), followed by the specified primary and then secondary antibodies. All incubations were done either at room temperature for 1 hour, or at 4 °C overnight. Signal was developed using SuperSignal™ West Pico PLUS Chemiluminescent Substrate (Thermo Scientific #34580) and images were taken on a ChemiDoc™ imager. Densitometry of the bands was measured using BioRad Image Lab.

The antibodies used for immunoblotting included: anti-GAPDH (Proteintech 10494-1-AP, 1:4000), anti-H3 (Abcam #Ab1791, 1:100000), anti-H3K27me1 (Active Motif 61015, 1:2000), anti-H3K27me2 (Abcam ab24684, 1:1000), anti-H3K27me3 (Active Motif 61017, 1:2500), anti-EZH2 (Cell Signalling 5246S, 1:1000), anti-SUZ12 (Cell Signalling 3737, 1:1000), anti-EED (Cell Signalling 85322, 1:1000) anti-mouse-HRP-conjugated (Jackson Immuno-Reasearch #715-035-150, 1:5000), and anti-rabbit HRP-conjugated (Santa Cruz Biotechnology #sc-2357, 1:5000).

Quantitative chromatin immunoprecipitation with whole genome sequencing (ChIP-Rx)

Cells were collected by replacing the culture media with trypsin for <5 minutes, then resuspending the detached cells in growth media and centrifuging for 5 minutes at 300 g. The cells were first washed once in PBS, then crosslinked for 10 minutes at

room temperature in PBS with 1% (v/v) formaldehyde. The formaldehyde was

quenched by adding glycine to a final concentration of 0.125 M, and was then

washed twice in PBS. Cell pellets containing an equal number of cells in each

sample were snap frozen with liquid nitrogen and stored at -80°C.

The cell pellets were thawed on ice and lysed in lysis buffer (100 mM NaCl, 0.5%

(v/v) SDS, 5 mM EDTA, 50 mM Tris-HCl pH 8 and 1X protease inhibitor cocktail

(Sigma #4693132001)) on ice, then centrifuged for 6 minutes at 300 g. The resulting

chromatin pellet was resuspended in ChIP buffer (a 2:1 ratio of lysis buffer and

dilution buffer (100 mM NaCl, 5% (v/v), Triton X-100, 5mM EDTA, 100mM Tris-HCl

pH 8.5 and 1X protease inhibitor cocktail (Sigma #4693132001))). For every 1×10^7

cells, chromatin from 5×10^6 cross-linked NTERA-2 cells were added. All samples

were then sonicated at 4°C in a Bioruptor set to high, with 30 cycles of 30 s on and

30 s off. The sonicated samples were diluted to 1×10^7 cells per mL, and 1mL per

antibody was used for ChIP, with 1% kept as input.

Antibodies were added to each sample as specified and incubated overnight at 4 °C

rotating at low speed. The antibodies used for ChIP included: anti-SUZ12 (Cell

Signalling 3737, 0.42 µg), anti-H3K27me3 (Cell Signalling 35861SF, 3 µg) anti-

Jarid2 (Cell Signalling 13594S, 1.115 µg), and anti-MTF2 (Proteintech 16208-1-AP,

1.435 µg). After the antibody incubation, the lysate was centrifuged at max speed for

30 minutes at 4 °C. The pellet was discarded and 40 µL of Dynabeads Protein G

(Thermo #10003D), washed 5 times in ChIP buffer, were added to the supernatant of

each sample and incubated for 2-4 hours at 4°C rotating at low speed. The beads

were then washed 3 times in mixed micelle buffer (150 mM NaCl, 20 mM Tris-HCl

pH 8, 5 mM EDTA, 5.2% (w/v) sucrose and 1% (v/v) Triton X-100), twice in buffer

500 (0.1% (w/v) sodium deoxycholate, 1 mM EDTA, 50 mM HEPES pH 7.5 and 1%

(v/v) Triton X-100), twice in LiCl/detergent wash buffer (0.5% (w/v) sodium

deoxycholate, 1 mM EDTA, 250 mM LiCl, 0.5% (v/v) NP-40, 10 mM Tris-HCl pH 8)

and once in TE (10 mM Tris-HCl pH 8 and 1 mM EDTA). The beads were then

resuspended in 100 µL elution buffer (1% (v/v) SDS and 100 mM NaHCO3) and

incubated at 65°C for 1 hour with shaking. The eluate was removed from the beads,

and 10 µL of the input was combined with 90 µL of elution buffer, then all samples

were reverse-crosslinked by shaking at 65 °C overnight.

1 µL of RNase A (Thermo #EN0531) was then added to all samples and incubated

shaking at 37 °C for 1 hour, followed by 1 µL of Proteinase K (NEB #P8107S) with 2

hours of shaking at 55 °C. DNA was purified from these samples using the Qiagen

MinElute PCR purification kit (Qiagen #28006) according to the instructions of the

manufacturer. The purified DNA was dried overnight at room temperature in a

biosafety hood before shipment to Azenta. Library preparation and Illumina

sequencing with 20 million paired-end reads of 150bp per library was performed by

Azenta for all samples.

Bioinformatic analysis of ChIP-Rx datasets

Chromosome names for the mouse genome were modified with the prefix 'mm10_'

and a metagenome was created by concatenating the mouse and human (spikein)

reference genomes (mm10 and hg38, respectively) before indexing with Bowtie2 [74]

as described previously [46]. Reads were aligned to the metagenome using Bowtie2

with default parameters [74]. Non-unique read alignments were filtered out using

SAMtools [75] to exclude those with an alignment quality of <2. The mm10_ prefix

appended to chromosome names was used to separate reads aligned to the

reference mouse or spike-in human genomes. SAMtools was used to convert SAM

files to BAM files and to remove duplicate aligned reads. Spike-in normalization

factors were calculated for each ChIP dataset using the formula for normalized

reference-adjusted reads per million (RRPM) as described [46] (1 per million spike-in

reads). BigWig files were generated using the bamCoverage tool from the deepTools

suite (version 3.3.0) [76] with a bin size of 10 and data were subsequently visualized

as ChIP-Rx normalized tracks using the IGV genome browser.

Transcription start sites (TSS) for the mm10 mouse genome build were defined and

annotated as all unique protein coding TSS +/- 5kb (n=19,888; downloaded from

ENSEMBL biomart). Peaks were called for SUZ12 from in the wild type cells using

MACS2 with paired-end function (BAMPE) and an FDR < 0.01 [77]. mm10 blacklisted

regions were removed from all peak sets prior to downstream analyses [78]. "PRC2-

targets" (n=4,445) were defined as TSS that overlapped with a SUZ12 peak using

bedTools intersect with -v and -wa flags [79]. The remaining TSS were defined as

"non-PRC2 targets" (n=16,118). "Intergenic regions" (n=22,050) included all non-

blacklist regions after excluding annotated genes and 5 kb upstream and

downstream from the start (TSS) and end (TES) of annotated genes.The genome-wide $\log_2$ fold-change of the ChIP-Rx signals for H3K27me3 and SUZ12 in the mutant cell line was calculated using the bigwigCompare tool from the deepTools suite [76]. "Outside SUZ12 peak" regions were defined as regions that reside within 30 kb upstream or downstream from the boundaries of all SUZ12 peaks and do not otherwise overlap with other SUZ12 peaks. Reads from both inside and outside SUZ12 peak regions were quantified using featureCounts [80], Rx-normalised, and visualised as violin plots using the ggpubr R package. Pairwise statistical tests were also performed using the ggpubr package. The proportion of H3K27me3 reads overlapping with H3K27me3 peaks was calculated using featureCounts. All heatmaps and average enrichment profiles were generated using the computeMatrix, plotHeatmap and plotProfile tools from the deepTools suite [76].

References

1. Kim, J.J., and Kingston, R.E. (2022). Context-specific Polycomb mechanisms in development. Nat Rev Genet 23, 680–695. https://doi.org/10.1038/s41576-022-00499-0.
2. Blackledge, N.P., and Klose, R.J. (2021). The molecular principles of gene regulation by Polycomb repressive complexes. Nature Reviews Molecular Cell Biology 22, 815–833. https://doi.org/10.1038/s41580-021-00398-y.
3. Piunti, A., and Shilatifard, A. (2021). The roles of Polycomb repressive complexes in mammalian development and cancer. Nature Reviews Molecular Cell Biology 22, 326–345. https://doi.org/10.1038/s41580-021-00341-1.
4. Faust, C., Schumacher, A., Holdener, B., and Magnuson, T. (1995). The eed mutation disrupts anterior mesoderm production in mice. Development 121, 273–285. https://doi.org/10.1242/dev.121.2.273.
5. O'Carroll, D., Erhardt, S., Pagani, M., Barton, S.C., Surani, M.A., and Jenuwein, T. (2001). The polycomb-group gene Ezh2 is required for early mouse development. Mol Cell Biol 21, 4330–4336. https://doi.org/10.1128/MCB.21.13.4330-4336.2001.
6. Pasini, D., Bracken, A.P., Jensen, M.R., Lazzerini Denchi, E., and Helin, K. (2004). Suz12 is essential for mouse development and for EZH2 histone methyltransferase activity. EMBO J 23, 4061–4071. https://doi.org/10.1038/sj.emboj.7600402.
7. Pengelly, A.R., Copur, Ö., Jäckle, H., Herzig, A., and Müller, J. (2013). A histone mutant reproduces the phenotype caused by loss of histone-modifying factor Polycomb. Science 339, 698–699. https://doi.org/10.1126/science.1231382.
8. Condemi, L., Mocavini, I., Aranda, S., and Di Croce, L. (2024). Polycomb function in early mouse development. Cell Death Differ, 1–10. https://doi.org/10.1038/s41418-024-01340-3.
9. Uckelmann, M., and Davidovich, C. (2021). Not just a writer: PRC2 as a chromatin reader. Biochemical Society Transactions 49, 1159–1170. https://doi.org/10.1042/bst20200728.
10. Grewal, S.I.S. (2023). The molecular basis of heterochromatin assembly and epigenetic inheritance. Mol Cell 83, 1767–1785. https://doi.org/10.1016/j.molcel.2023.04.020.
11. Hansen, K.H., Bracken, A.P., Pasini, D., Dietrich, N., Gehani, S.S., Monrad, A., Rappsilber, J., Lerdrup, M., and Helin, K. (2008). A model for transmission of the H3K27me3 epigenetic

mark. Nat Cell Biol *10*, 1291–1300. https://doi.org/10.1038/ncb1787.

12. Margueron, R., Justin, N., Ohno, K., Sharpe, M.L., Son, J., Drury Iii, W.J., Voigt, P., Martin,
S.R., Taylor, W.R., De Marco, V., et al. (2009). Role of the polycomb protein EED in the
propagation of repressive histone marks. Nature *461*, 762–767.
https://doi.org/10.1038/nature08398.

13. Qi, W., Zhao, K., Gu, J., Huang, Y., Wang, Y., Zhang, H., Zhang, M., Zhang, J., Yu, Z., Li, L., et
al. (2017). An allosteric PRC2 inhibitor targeting the H3K27me3 binding pocket of EED. Nat
Chem Biol *13*, 381–388. https://doi.org/10.1038/nchembio.2304.

14. Lee, C.H., Yu, J.R., Kumar, S., Jin, Y., LeRoy, G., Bhanu, N., Kaneko, S., Garcia, B.A.,
Hamilton, A.D., and Reinberg, D. (2018). Allosteric Activation Dictates PRC2 Activity
Independent of Its Recruitment to Chromatin. Mol Cell *70*, 422-434.e6.
https://doi.org/10.1016/j.molcel.2018.03.020.

15. Sanulli, S., Justin, N., Teissandier, A., Ancelin, K., Portoso, M., Caron, M., Michaud, A.,
Lombard, B., da Rocha, S.T., Offer, J., et al. (2015). Jarid2 Methylation via the PRC2
Complex Regulates H3K27me3 Deposition during Cell Differentiation. Mol Cell *57*, 769–
783. https://doi.org/10.1016/j.molcel.2014.12.020.

16. Zhang, Q., Agius, S.C., Flanigan, S.F., Uckelmann, M., Levina, V., Owen, B.M., and
Davidovich, C. (2021). PALI1 facilitates DNA and nucleosome binding by PRC2 and triggers
an allosteric activation of catalysis. Nature Communications *12*, 4592.
https://doi.org/10.1038/s41467-021-24866-3.

17. Hauri, S., Comoglio, F., Seimiya, M., Gerstung, M., Glatter, T., Hansen, K., Aebersold, R.,
Paro, R., Gstaiger, M., and Beisel, C. (2016). A High-Density Map for Navigating the Human
Polycomb Complexome. Cell Rep *17*, 583–595.
https://doi.org/10.1016/j.celrep.2016.08.096.

18. Conway, E., Jerman, E., Healy, E., Ito, S., Holoch, D., Oliviero, G., Deevy, O., Glancy, E.,
Fitzpatrick, D.J., Mucha, M., et al. (2018). A Family of Vertebrate-Specific Polycombs
Encoded by the LCOR/LCORL Genes Balance PRC2 Subtype Activities. Mol Cell *70*, 408-
421.e8. https://doi.org/10.1016/j.molcel.2018.03.005.

19. Peng, J.C., Valouev, A., Swigut, T., Zhang, J., Zhao, Y., Sidow, A., and Wysocka, J. (2009).
Jarid2/Jumonji Coordinates Control of PRC2 Enzymatic Activity and Target Gene
Occupancy in Pluripotent Cells. Cell *139*, 1290–1302.
https://doi.org/10.1016/j.cell.2009.12.002.

20. Shen, X., Kim, W., Fujiwara, Y., Simon, M.D., Liu, Y., Mysliwiec, M.R., Yuan, G.C., Lee, Y.,
and Orkin, S.H. (2009). Jumonji modulates polycomb activity and self-renewal versus
differentiation of stem cells. Cell *139*, 1303–1314.
https://doi.org/10.1016/j.cell.2009.12.003.

21. Li, G., Margueron, R., Ku, M., Chambon, P., Bernstein, B.E., and Reinberg, D. (2010). Jarid2
and PRC2, partners in regulating gene expression. Genes Dev *24*, 368–380.
https://doi.org/10.1101/gad.1886410.

22. Kalb, R., Latwiel, S., Baymaz, H.I., Jansen, P.W.T.C., Müller, C.W., Vermeulen, M., and
Müller, J. (2014). Histone H2A monoubiquitination promotes histone H3 methylation in
Polycomb repression. Nat Struct Mol Biol *21*, 569–571. https://doi.org/10.1038/nsmb.2833.

23. Yu, J.R., Lee, C.H., Oksuz, O., Stafford, J.M., and Reinberg, D. (2019). PRC2 is high
maintenance. Genes Dev *33*, 903–935. https://doi.org/10.1101/gad.325050.119.

24. Davidovich, C., and Zhang, Q. (2021). Allosteric regulation of histone lysine
methyltransferases: from context-specific regulation to selective drugs. Biochemical
Society Transactions *49*, 591–607. https://doi.org/10.1042/BST20200238.

25. Yang, Y., and Li, G. (2020). Post-translational modifications of PRC2: signals directing its
activity. Epigenetics & Chromatin *13*, 47. https://doi.org/10.1186/s13072-020-00369-1.

26. Park, S.H., Fong, K.-W., Mong, E., Martin, M.C., Schiltz, G.E., and Yu, J. (2021). Going
beyond Polycomb: EZH2 functions in prostate cancer. Oncogene *40*, 5788–5798.

https://doi.org/10.1038/s41388-021-01982-4.

27. Liu, X. (2021). A Structural Perspective on Gene Repression by Polycomb Repressive
Complex 2. Subcell Biochem *96*, 519–562. https://doi.org/10.1007/978-3-030-58971-4_17.

28. Hananya, N., Koren, S., and Muir, T.W. (2024). Interrogating epigenetic mechanisms with
chemically customized chromatin. Nat Rev Genet *25*, 255–271.
https://doi.org/10.1038/s41576-023-00664-z.

29. Chittock, E.C., Latwiel, S., Miller, T.C.R., and Müller, C.W. (2017). Molecular architecture of
polycomb repressive complexes. Biochem Soc Trans *45*, 193–205.
https://doi.org/10.1042/BST20160173.

30. Kadoch, C., Copeland, R.A., and Keilhack, H. (2016). PRC2 and SWI/SNF Chromatin
Remodeling Complexes in Health and Disease. Biochemistry *55*, 1600–1614.
https://doi.org/10.1021/acs.biochem.5b01191.

31. van Kruijsbergen, I., Hontelez, S., and Veenstra, G.J.C. (2015). Recruiting polycomb to
chromatin. Int J Biochem Cell Biol *67*, 177–187.
https://doi.org/10.1016/j.biocel.2015.05.006.

32. Holoch, D., and Margueron, R. (2017). Mechanisms Regulating PRC2 Recruitment and
Enzymatic Activity. Trends Biochem Sci *42*, 531–542.
https://doi.org/10.1016/j.tibs.2017.04.003.

33. Laugesen, A., and Helin, K. (2014). Chromatin Repressive Complexes in Stem Cells,
Development, and Cancer. Cell Stem Cell *14*, 735–751.
https://doi.org/10.1016/j.stem.2014.05.006.

34. Takeuchi, T., Yamazaki, Y., Katoh-Fukui, Y., Tsuchiya, R., Kondo, S., Motoyama, J., and
Higashinakagawa, T. (1995). Gene trap capture of a novel mouse gene, jumonji, required for
neural tube formation. Genes Dev *9*, 1211–1222. https://doi.org/10.1101/gad.9.10.1211.

35. Blackledge, N.P., Farcas, A.M., Kondo, T., King, H.W., McGouran, J.F., Hanssen, L.L.P., Ito,
S., Cooper, S., Kondo, K., Koseki, Y., et al. (2014). Variant PRC1 complex-dependent H2A
ubiquitylation drives PRC2 recruitment and polycomb domain formation. Cell *157*, 1445–
1459. https://doi.org/10.1016/j.cell.2014.05.004.

36. Li, X., Isono, K., Yamada, D., Endo, T.A., Endoh, M., Shinga, J., Mizutani-Koseki, Y., Otte,
A.P., Casanova, M., Kitamura, H., et al. (2011). Mammalian polycomb-like Pcl2/Mtf2 is a
novel regulatory component of PRC2 that can differentially modulate polycomb activity
both at the Hox gene cluster and at Cdkn2a genes. Mol Cell Biol *31*, 351–364.
https://doi.org/10.1128/mcb.00259-10.

37. Isono, K., Fujimura, Y., Shinga, J., Yamaki, M., O. Wang J, Takihara, Y., Murahashi, Y.,
Takada, Y., Mizutani-Koseki, Y., and Koseki, H. (2005). Mammalian polyhomeotic
homologues Phc2 and Phc1 act in synergy to mediate polycomb repression of Hox genes.
Mol Cell Biol *25*, 6694–6706. https://doi.org/10.1128/mcb.25.15.6694-6706.2005.

38. Suzuki, M., Mizutani-Koseki, Y., Fujimura, Y.-I., Miyagishima, H., Kaneko, T., Takada, Y.,
Akasaka, T., Tanzawa, H., Takihara, Y., Nakano, M., et al. (2002). Involvement of the
Polycomb-group gene Ring1B in the specification of the anterior-posterior axis in mice.
Development *129*, 4171–4183. https://doi.org/10.1242/dev.129.18.4171.

39. Akasaka, T., Kanno, M., Balling, R., Mieza, M.A., Taniguchi, M., and Koseki, H. (1996). A role
for mel-18, a Polycomb group-related vertebrate gene, during the anteroposterior
specification of the axial skeleton. Development *122*, 1513–1522.
https://doi.org/10.1242/dev.122.5.1513.

40. van der Lugt, N.M., Domen, J., Linders, K., van Roon, M., Robanus-Maandag, E., te Riele, H.,
van der Valk, M., Deschamps, J., Sofroniew, M., and van Lohuizen, M. (1994). Posterior
transformation, neurological abnormalities, and severe hematopoietic defects in mice with
a targeted deletion of the bmi-1 proto-oncogene. Genes Dev *8*, 757–769.
https://doi.org/10.1101/gad.8.7.757.

41. Wang, S., He, F., Xiong, W., Gu, S., Liu, H., Zhang, T., Yu, X., and Chen, Y. (2007).

1049  Polycomblike-2-deficient mice exhibit normal left–right asymmetry. Developmental
1050  Dynamics *236*, 853–861. https://doi.org/10.1002/dvdy.21070.

1051 42. Vizán, P., Gutiérrez, A., Espejo, I., García-Montolio, M., Lange, M., Carretero, A., Lafzi, A., de
1052  Andrés-Aguayo, L., Blanco, E., Thambyrajah, R., et al. (2020). The Polycomb-associated
1053  factor PHF19 controls hematopoietic stem cell state and differentiation. Science Advances
1054  *6*, eabb2745. https://doi.org/10.1126/sciadv.abb2745.

1055 43. Takihara, Y., Tomotsune, D., Shirai, M., Katoh-Fukui, Y., Nishii, K., Motaleb, Md.A., Nomura,
1056  M., Tsuchiya, R., Fujita, Y., Shibata, Y., et al. (1997). Targeted disruption of the mouse
1057  homologue of the Drosophila polyhomeotic gene leads to altered anteroposterior
1058  patterning and neural crest defects. Development *124*, 3673–3682.
1059  https://doi.org/10.1242/dev.124.19.3673.

1060 44. Schumacher, A., Faust, C., and Magnuson, T. (1996). Positional cloning of a global regulator
1061  of anterior–posterior patterning in mice. Nature *383*, 250–253.
1062  https://doi.org/10.1038/383250a0.

1063 45. Pérez-Pérez, J.M., Candela, H., and Micol, J.L. (2009). Understanding synergy in genetic
1064  interactions. Trends Genet *25*, 368–376. https://doi.org/10.1016/j.tig.2009.06.004.

1065 46. Orlando, D.A., Chen, M.W., Brown, V.E., Solanki, S., Choi, Y.J., Olson, E.R., Fritz, C.C.,
1066  Bradner, J.E., and Guenther, M.G. (2014). Quantitative ChIP-Seq Normalization Reveals
1067  Global Modulation of the Epigenome. Cell Reports *9*, 1163–1170.
1068  https://doi.org/10.1016/j.celrep.2014.10.018.

1069 47. Lavarone, E., Barbieri, C.M., and Pasini, D. (2019). Dissecting the role of H3K27 acetylation
1070  and methylation in PRC2 mediated control of cellular identity. Nat Commun *10*, 1679.
1071  https://doi.org/10.1038/s41467-019-09624-w.

1072 48. Glancy, E., Wang, C., Tuck, E., Healy, E., Amato, S., Neikes, H.K., Mariani, A., Mucha, M.,
1073  Vermeulen, M., Pasini, D., et al. (2023). PRC2.1- and PRC2.2-specific accessory proteins
1074  drive recruitment of different forms of canonical PRC1. Mol Cell *83*, 1393-1411.e7.
1075  https://doi.org/10.1016/j.molcel.2023.03.018.

1076 49. Gail, E.H., Healy, E., Flanigan, S.F., Jones, N., Ng, X.H., Uckelmann, M., Levina, V., Zhang,
1077  Q., and Davidovich, C. (2024). Inseparable RNA binding and chromatin modification
1078  activities of a nucleosome-interacting surface in EZH2. Nat Genet *56*, 1193–1202.
1079  https://doi.org/10.1038/s41588-024-01740-8.

1080 50. Højfeldt, J.W., Hedehus, L., Laugesen, A., Tatar, T., Wiehle, L., and Helin, K. (2019). Non-
1081  core Subunits of the PRC2 Complex Are Collectively Required for Its Target-Site Specificity.
1082  Mol Cell *76*, 423-436.e3. https://doi.org/10.1016/j.molcel.2019.07.031.

1083 51. Healy, E., Mucha, M., Glancy, E., Fitzpatrick, D.J., Conway, E., Neikes, H.K., Monger, C., Van
1084  Mierlo, G., Baltissen, M.P., Koseki, Y., et al. (2019). PRC2.1 and PRC2.2 Synergize to
1085  Coordinate H3K27 Trimethylation. Mol Cell *76*, 437-452.e6.
1086  https://doi.org/10.1016/j.molcel.2019.08.012.

1087 52. Kim, J.-A., Kwon, M., and Kim, J. (2019). Allosteric Regulation of Chromatin-Modifying
1088  Enzymes. Biochemistry *58*, 15–23. https://doi.org/10.1021/acs.biochem.8b00894.

1089 53. Fong, K., Zhao, J.C., Lu, X., Kim, J., Piunti, A., Shilatifard, A., and Yu, J. (2022). PALI1
1090  promotes tumor growth through competitive recruitment of PRC2 to G9A-target chromatin
1091  for dual epigenetic silencing. Molecular Cell *82*, 4611-4626.e7.
1092  https://doi.org/10.1016/j.molcel.2022.11.010.

1093 54. Schmitges, F.W., Prusty, A.B., Faty, M., Stützer, A., Lingaraju, G.M., Aiwazian, J., Sack, R.,
1094  Hess, D., Li, L., Zhou, S., et al. (2011). Histone methylation by PRC2 is inhibited by active
1095  chromatin marks. Mol Cell *42*, 330–341. https://doi.org/10.1016/j.molcel.2011.03.025.

1096 55. Cookis, T., Lydecker, A., Sauer, P., Kasinath, V., and Nogales, E. (2024). Structural basis for
1097  the inhibition of PRC2 by active transcription histone posttranslational modifications.
1098  Preprint, https://doi.org/10.1101/2024.02.09.579730
1099  https://doi.org/10.1101/2024.02.09.579730.

56. van de Wetering, M., Francies, H.E., Francis, J.M., Bounova, G., Iorio, F., Pronk, A., van
Houdt, W., van Gorp, J., Taylor-Weiner, A., Kester, L., et al. (2015). Prospective derivation of
a Living Organoid Biobank of colorectal cancer patients. Cell *161*, 933–945.
https://doi.org/10.1016/j.cell.2015.03.053.
57. Arai, E., Sakamoto, H., Ichikawa, H., Totsuka, H., Chiku, S., Gotoh, M., Mori, T., Nakatani, T.,
Ohnami, S., Nakagawa, T., et al. (2014). Multilayer-omics analysis of renal cell carcinoma,
including the whole exome, methylome and transcriptome. Int J Cancer *135*, 1330–1342.
https://doi.org/10.1002/ijc.28768.
58. Bonilla, X., Parmentier, L., King, B., Bezrukov, F., Kaya, G., Zoete, V., Seplyarskiy, V.B.,
Sharpe, H.J., McKee, T., Letourneau, A., et al. (2016). Genomic analysis identifies new
drivers and progression pathways in skin basal cell carcinoma. Nat Genet *48*, 398–406.
https://doi.org/10.1038/ng.3525.
59. Hayward, N.K., Wilmott, J.S., Waddell, N., Johansson, P.A., Field, M.A., Nones, K., Patch,
A.-M., Kakavand, H., Alexandrov, L.B., Burke, H., et al. (2017). Whole-genome landscapes
of major melanoma subtypes. Nature *545*, 175–180. https://doi.org/10.1038/nature22071.
60. Shain, A.H., Garrido, M., Botton, T., Talevich, E., Yeh, I., Sanborn, J.Z., Chung, J., Wang, N.J.,
Kakavand, H., Mann, G.J., et al. (2015). Exome sequencing of desmoplastic melanoma
identifies recurrent NFKBIE promoter mutations and diverse activating mutations in the
MAPK pathway. Nat Genet *47*, 1194–1199. https://doi.org/10.1038/ng.3382.
61. Cosmic COSMIC - Catalogue of Somatic Mutations in Cancer.
https://cancer.sanger.ac.uk/cosmic.
62. Tate, J.G., Bamford, S., Jubb, H.C., Sondka, Z., Beare, D.M., Bindal, N., Boutselakis, H.,
Cole, C.G., Creatore, C., Dawson, E., et al. (2019). COSMIC: the Catalogue Of Somatic
Mutations In Cancer. Nucleic Acids Research *47*, D941–D947.
https://doi.org/10.1093/nar/gky1015.
63. Chammas, P., Mocavini, I., and Di Croce, L. (2020). Engaging chromatin: PRC2 structure
meets function. Br J Cancer *122*, 315–328. https://doi.org/10.1038/s41416-019-0615-2.
64. Morschhauser, F., Tilly, H., Chaidos, A., McKay, P., Phillips, T., Assouline, S., Batlevi, C.L.,
Campbell, P., Ribrag, V., Damaj, G.L., et al. (2020). Tazemetostat for patients with relapsed
or refractory follicular lymphoma: an open-label, single-arm, multicentre, phase 2 trial.
Lancet Oncol *21*, 1433–1442. https://doi.org/10.1016/S1470-2045(20)30441-1.
65. Motoyama, J., Kitajima, K., Kojima, M., Kondo, S., and Takeuchi, T. (1997). Organogenesis of
the liver, thymus and spleen is affected in jumonji mutant mice. Mech Dev *66*, 27–37.
https://doi.org/10.1016/s0925-4773(97)00082-8.
66. Takeuchi, T., Kojima, M., Nakajima, K., and Kondo, S. (1999). jumonji gene is essential for
the neurulation and cardiac development of mouse embryos with a C3H/He background.
Mech Dev *86*, 29–38. https://doi.org/10.1016/s0925-4773(99)00100-8.
67. Lee, Y., Song, A.J., Baker, R., Micales, B., Conway, S.J., and Lyons, G.E. (2000). Jumonji, a
nuclear protein that is necessary for normal heart development. Circ Res *86*, 932–938.
https://doi.org/10.1161/01.res.86.9.932.
68. Rice, P., Longden, I., and Bleasby, A. (2000). EMBOSS: the European Molecular Biology
Open Software Suite. Trends Genet *16*, 276–277. https://doi.org/10.1016/s0168-
9525(00)02024-2.
69. EMBOSS http://emboss.open-bio.org/.
70. Chang, Y.-C., Manent, J., Schroeder, J., Wong, S.F.L., Hauswirth, G.M., Shylo, N.A., Moore,
E.L., Achilleos, A., Garside, V., Polo, J.M., et al. (2022). Nr6a1 controls Hox expression
dynamics and is a master regulator of vertebrate trunk development. Nature
Communications *13*, 7766. https://doi.org/10.1038/s41467-022-35303-4.
71. Rigueur, D., and Lyons, K.M. (2014). Whole-mount skeletal staining. Methods Mol Biol *1130*,
113–121. https://doi.org/10.1007/978-1-62703-989-5_9.
72. Blain, R., Couly, G., Shotar, E., Blévinal, J., Toupin, M., Favre, A., Abjaghou, A., Inoue, M.,

Hernández-Garzón, E., Clarençon, F., et al. (2023). A tridimensional atlas of the developing
human head. Cell *186*, 5910. https://doi.org/10.1016/j.cell.2023.11.013.

73. Young, L., Sung, J., Stacey, G., and Masters, J.R. (2010). Detection of Mycoplasma in cell
cultures. Nat Protoc *5*, 929–934. https://doi.org/10.1038/nprot.2010.43.

74. Langmead, B., and Salzberg, S.L. (2012). Fast gapped-read alignment with Bowtie 2. Nat
Methods *9*, 357–359. https://doi.org/10.1038/nmeth.1923.

75. Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G., Abecasis, G.,
Durbin, R., and 1000 Genome Project Data Processing Subgroup (2009). The Sequence
Alignment/Map format and SAMtools. Bioinformatics *25*, 2078–2079.
https://doi.org/10.1093/bioinformatics/btp352.

76. Ramírez, F., Ryan, D.P., Grüning, B., Bhardwaj, V., Kilpert, F., Richter, A.S., Heyne, S.,
Dündar, F., and Manke, T. (2016). deepTools2: a next generation web server for deep-
sequencing data analysis. Nucleic Acids Res *44*, W160-165.
https://doi.org/10.1093/nar/gkw257.

77. Zhang, Y., Liu, T., Meyer, C.A., Eeckhoute, J., Johnson, D.S., Bernstein, B.E., Nusbaum, C.,
Myers, R.M., Brown, M., Li, W., et al. (2008). Model-based analysis of ChIP-Seq (MACS).
Genome Biol 9, R137. https://doi.org/10.1186/gb-2008-9-9-r137.

78. Amemiya, H.M., Kundaje, A., and Boyle, A.P. (2019). The ENCODE Blacklist: Identification of
Problematic Regions of the Genome. Sci Rep *9*, 9354. https://doi.org/10.1038/s41598-019-
45839-z.

79. Quinlan, A.R., and Hall, I.M. (2010). BEDTools: a flexible suite of utilities for comparing
genomic features. Bioinformatics *26*, 841–842.
https://doi.org/10.1093/bioinformatics/btq033.

80. Liao, Y., Smyth, G.K., and Shi, W. (2014). featureCounts: an efficient general purpose
program for assigning sequence reads to genomic features. Bioinformatics *30*, 923–930.
https://doi.org/10.1093/bioinformatics/btt656.

1177