Chronic lysosome damage boosts interferon responses to Palbociclib through a mitochondrial signalling axis

Mihaela Bozic[1*], Tian En Lim[1*], Alain J. Kemp[1], Katie Winnington-Ingram[1], Laura Murphy[2], Ashish

Dhir[2,3], Ann Wheeler[2], Natalia Jimenez-Moreno[1], Simon Wilkinson[1,4]

[1]Cancer Research UK Scotland Centre, Institute of Genetics and Cancer, University of Edinburgh, Edinburgh EH4 2XR, UK

[2]MRC Human Genetics Unit, Institute of Genetics and Cancer, University of Edinburgh, Edinburgh EH4 2XU, UK

[3]Current address: 4BASEBIO DISCOVERY LIMITED, 25 Norman Way, Over, Cambridge, Cambridgeshire, England, CB24 5QE, UK

[4]Corresponding author, s.wilkinson@ed.ac.uk

*these authors contributed equally to this work

Abstract

Acute lysosome damage triggers the endolysosome damage response (ELDR) in order to co-ordinate vesicle repair or removal by autophagy (lysophagy). However, it is unclear whether persistent damage, as occurs after chronic challenge to lysosome integrity, triggers wider cellular responses. Here, we show that longitudinal treatment with a lysosomotropic cancer therapeutic, the CDK4/6 inhibitor Palbociclib, invokes chronic lysosome damage in breast and lung cancer cells. Autophagy ameliorates but does not avert this phenotype, which persists over days. Damaged lysosomes form contacts with mitochondria, invoking mitochondrial stress and cytosolic efflux of immunostimulatory mitochondrial nucleic acids. Importantly, mitochondrial nucleic acid release is necessary for the anti-cancer interferon response to Palbociclib. In conclusion, chronic lysosome damage rewires cellular signalling responses through the mitochondrion and this effect should be considered when assessing the cellular actions of cancer therapeutics.

Introduction

Lysosomes are single membrane-bound hydrolytic organelles that act at the termini of endocytic and autophagic pathways (Zoncu and Perera, 2022). Although rupture of pathogen-containing phagolysosomes is a feature of some infections (Thurston et al., 2012), it is also well established that, in a sterile context, widespread loss of integrity of lysosomes can result in toxic leakage of hydrolytic proteases to the cytoplasm and subsequent cell death (Wang et al., 2018). However, acute, sublethal loss of lysosome membrane integrity typically induces the endolysosome damage response (ELDR), a collective term for pathways facilitating membrane stabilisation and repair (Radulovic et al., 2018; Skowyra et al., 2018; Tan and Finkel, 2022), new lysosome biogenesis (Jia et al., 2020b) or lysophagy (Papadopoulos et al., 2020), the latter being the process of selective degradation of the damaged lysosome via autophagy (Maejima et al., 2013). Lysophagy requires evolutionarily-conserved core autophagy proteins that facilitate generation of nascent autophagosome vesicles and the concurrent sequestration of damaged lysosome cargo therein (Dikic and Elazar, 2018; Vargas et al., 2023). Mature autophagosomes fuse with healthy lysosomes to facilitate degradation of sequestered cargo. Damaged lysosomes are initially recognised for degradation via "eat-me" signals (Papadopoulos et al., 2020; Hoyer et al., 2022). These include recruitment of galectin proteins, for instance Galectin 8 (GAL8) (Thurston et al., 2012; Jia et al., 2018), to glycosylated lysosomal membrane proteins, and conjugation of polyubiquitin chains to lysosomal proteins (Kravic et al., 2022; Gahlot et al., 2024).

Mechanistically, acute damage to lysosomes leads to galectin- and polyubiquitin-mediated recruitment of molecules that mediate lysophagy (Thurston et al., 2012; Chauhan et al., 2016; Papadopoulos et al., 2017; Jia et al., 2018; Jia et al., 2020a; Jia et al., 2020b; Eapen et al., 2021; Gallagher and Holzbaur, 2023; Park et al., 2025), in particular adaptors that bind directly to ATG8 (LC3/GABARAP) proteins conjugated to nascent autophagosome membranes (Johansen and Lamark, 2020) and which also recruit additional factors upstream of ATG8 conjugation (Lazarou et al., 2015;

Ravenhill et al., 2019; Turco et al., 2019; Vargas et al., 2019). Indeed, deletion of *ATG5,* which prevents ATG8 conjugation to autophagosomal (and lysosomal) membranes, is effective in stabilising damaged lysosomes by ablating lysophagy and also by compromising ATG8-mediated recruitment of repair factors (Bhattacharya et al., 2023; Cross et al., 2023). Importantly, the removal and repair responses described above are engaged acutely, within minutes lysosomal damage.

Notably, while extensive lysosome damage provokes cell death, linked to neurodegeneration (Wang et al., 2018; Zoncu and Perera, 2022) and disorders in other tissues such as kidney (Maejima et al., 2013), the broader cellular consequences of chronic - yet sublethal - damage are unclear. This is particularly true in the context where the acute restorative ELDR functions, such as those dependent upon *ATG5*, are overwhelmed. Recent data indicate that programmed signalling responses to lysosome damage may exist (Jia et al., 2018). For instance, lysosome rupture triggers RNA stress granule formation via PKR (Protein Kinase R) activation, which may maintain cell survival, over at least the first four hours of lysosomal challenge (Jia et al., 2022; Duran et al., 2024). However, on the whole, mechanistic studies have focussed heavily on acute damage after treatment with tool compounds, such as the membrane lytic peptide L-Leucyl-L-Leucine methyl ester (LLOMe) (Maejima et al., 2013). Importantly, numerous cancer therapeutics accumulate within lysosomes in either the cancer cells or other cell types. Generally, this phenomenon is viewed as detracting from therapeutic output by sequestering molecules from their primary targets. Nonetheless, it is possible that this may be consequential for the mechanism of the drugs. For instance, exposure of primary myeloid cells with lysosomotropic tyrosine kinase inhibitors may trigger lysosome damage and inflammasome activation, potentially contributing to treatment efficacy (Neuwirt et al., 2023).

Palbociclib (Pb) is an important anti-cancer drug that is sequestered from its enzymatic target, the cell-cycle progression kinase CDK4/6, due to lysosomal trapping (Llanos et al., 2019). Strikingly, in

addition to cell-cycle inhibition, Palbociclib also engages an anti-tumoural interferon response (Goel et al., 2017; Fan et al., 2023; Wu et al., 2024). Signalling occurs via detection of immunostimulatory moieties, purportedly from nuclear-derived double-stranded (ds) RNA (Roulois et al., 2015), by cytosolic pattern recognition receptors (PRRs) (Akira et al., 2006), including RIG-I (retinoic-acid inducible gene 1) working in tandem with MAVS (mitochondrial antiviral signaling protein) (Yoneyama et al., 2004; Seth et al., 2005; Sun et al., 2013; Rehwinkel and Gack, 2020). This action of Palbociclib has been shown to drive anti-tumour immunity (Goel et al., 2017). Indeed, discovering triggers and mechanisms of sterile interferon responses via PRRs is an important research goal in cancer therapy (Zitvogel et al., 2015; Crow and Stetson, 2022). Other potential triggers could include exogenous DNA, nuclear DNA, and mitochondrial DNA or dsRNA (Ishikawa and Barber, 2008; Mackenzie et al., 2017; Dhir et al., 2018; Tigano et al., 2021).

Here, we reveal a mechanistic link between chronic lysosome damage and interferon signalling, as part of a new pathway for engagement of sterile interferon signalling. We show that chronic lysosome damage can be provoked by Palbociclib, independently of CDK4/6, as well as the tool compound LLOMe. Chronic lysosome damage results in persistent interorganellar contacts between ruptured lysosomes and mitochondria. Lysosome damage triggers stress responses within mitochondria and ultimately results in accumulation of immunostimulatory mitochondrial nucleic acid in the cytosol, which we show is a requisite event for productive interferon responses. However, in the case of Palbociclib the interferon responses are synergistically facilitated by concomitant CDK4/6 inhibition. Overall, our data show that mitochondria act as priming hubs in triggering of sterile interferon responses by chronic lysosome damage. This molecular mechanism has implications for understanding of - and for amelioration or enhancement of - the cellular response to various chronic insults, including cancer therapeutics.

Results

Palbociclib induces chronic lysosome damage and ELDR

Given that Palbociclib (Pb) accumulates in lysosomes and causes swelling (Llanos et al., 2019), we hypothesised that it might trigger lysophagy, indicative of lysosome damage. Importantly, cancer treatments are applied chronically, over days and weeks, rather than on the acute minute-to-hour scales typically employed in lysophagy studies. We thus examined engagement of lysophagy flux by 24 h treatment with Pb, alongside the established lysosome-damaging compound L-Leucyl-L-Leucine methyl ester (LLOMe); these assays were performed using a new, sensitive reporter consisting of GAL8 fused to the C-terminus of SRAI (signal-retaining autophagy indicator, Fig. 1A) (Katayama et al., 2020; Jimenez-Moreno et al., 2023). Similar to LLOMe, Pb treatment was discovered to trigger lysophagy flux in A549 lung and MCF7 breast cancer cells (Fig. 1B-C, Supp. Fig. 1). Corroborating this, we also tested GAL8 fused to the conventional mRFP-GFP autophagy flux probe, and enumerated discrete individual autolysosomes containing damaged lysosome remnants (mRFP$^+$ GFP$^-$ galectolysosomes, wherein the acidic lysosome environment quenches the GFP signal, Fig. 1D-E). Again, these were induced by Pb treatment; additionally, consistent with elevated lysophagy flux, they were dependent upon *ATG5* (non-targeted wild-type, NT, compared with Δ*ATG5* cells, Fig. 1D-E).

Next, we examined the steady-state abundance of damaged lysosomes over both acute and chronic treatment regimens. YFP-GAL8 foci, indicative of undegraded and unrepaired damaged lysosomes, were observed in response to chronic 48 h Pb treatment in A549 cells (Fig. 1F), in particular within the autophagy-deficient Δ*ATG5* cells. These YFP-GAL8 foci co-stained with the lysosomal maker LAMP2, as expected (Fig. 1G). Quantifying these foci at multiple time points of treatment, it was observed that treatment with Pb provoked a biphasic response in wild-type A549 (NT) cells; a single dose invoked YFP-GAL8 foci accumulation within 1 h, peaking at 24 h, but then resolving by 48 h (Fig. 1H), whereas

in $\Delta ATG5$ cells damaged lysosomes persisted at 48 h. A similar trend was observed for single dose LLOMe treatment, although the peak abundance of damaged lysosomes was observed earlier and, while still evident at 8 h, was completely resolved by 24 h in wild-type (NT) cells (Fig. 1I). Broadly similar responses to Pb and LLOMe were seen in wild-type MCF7 cells, although Palbociclib-induced lysosome damage was still evident at 48 h here, suggesting a lower capacity than A549 cells for *ATG5*-dependent damage resolution (Fig. 1J). Finally, the lysosome damaging effect of Pb was observed to be independent of its nucleocytoplasmic targets, CDK4 and CDK6 (Fig. 1L-M); A549 $\Delta ATG5$ cells were stably knocked out for CDK4 ($\Delta CDK4$) and then CDK6 was acutely knocked out (g*CDK6*), immediately prior to Pb treatment (in order to avoid confounding senescence engaged by double knockout), yet this did not alter YFP-GAL8 focus induction by Pb (Fig. 1M).

Considering the above data together, we propose a new role for Pb in inducing chronic lysosome damage and lysophagy, independently of its primary target, CDK4/6. Moreover, the stabilisation of damaged lysosomes in A549 $\Delta ATG5$ cells, after single treatment with Pb, provides a facile experimental setting for unpicking the events downstream of chronic lysosome damage.

Damaged lysosomes associate with mitochondria

Under basal conditions, unperturbed lysosomes form transient interorganellar contacts with mitochondria (Wong et al., 2018). Given reports that Pb may also perturb mitochondrial physiology (Uzhachenko et al., 2021), we hypothesised that damaged lysosomes could associate with mitochondria. To explore this, we first performed transmission electron microscopy (TEM) of Pb-treated A549 cells (Fig. 2A). Here, particularly in *ATG5*-deficient cells, the expected presence of swollen, electron-lucent lysosomes was apparent when compared to untreated controls (Fig. 2B), and interorganellar contacts between lysosomes and the mitochondrial network were increased (Fig. 2C).

Next, using super-resolution immunofluorescence microscopy, we determined that YFP-GAL8 lysosome foci induced by Pb were associated with - and enwrapped - the mitochondrial network (demarcated by TOM20, Fig. 2D, Supp. Movie 1). The non-random nature of this association was quantified by analysing the distances of YFP-GAL8 foci from the outer surface of the mitochondrial network and comparing to those observed after YFP-GAL8 foci were computationally shuffled to random positions within the cytoplasmic space (Fig. 2E). Finally, foci dynamics were observed by super-resolution radial fluctuation (SRRF) time-lapse microscopy (Fig. 2F, Supp. Movies 2 and 3). YFP-GAL8 structures persistently resided along the mitochondrial network, with no or few dissociation events observed over recording times of 10 minutes (Fig. 2F, Supp. Movies 2 and 3). Taking the above data together, we conclude that chronic damage by Pb results in ruptured lysosomes - or fragments thereof - becoming stably associated with the mitochondrial network.

Lysosome damage triggers stereotypical markers of mitochondrial nucleic acid stress

Recently, nucleic acid efflux from the mitochondrion has been described as a signal that triggers interferon responses. This phenomenon frequently correlates with evidence of mitochondrial nucleic acid stress, as visualised by the aggregation of mitochondrial genome nucleoids (West et al., 2015; Kim et al., 2019; Torres-Odio et al., 2021). We hypothesised that contact between damaged lysosomes and mitochondria might induce such mitochondrial nucleic acid stress. Indeed, we observed the formation of large TFAM (mitochondrial transcription factor A)- and DNA-positive foci in the mitochondrial network in response to chronic Pb treatment, indicative of aggregated mitochondrial nucleoids (Fig. 3A-D). This was greatest in $\Delta ATG5$ cells, underscoring the relationship between sustained presence of damaged lysosomes and the mitochondrial response to Pb. Corroborating these data, similar results were obtained when aggregated nucleoids were alternately imaged by staining for FASTKD2, which marks nucleoid-associated RNA processing granules (Antonicka et al., 2013) (Fig. 3E-F). Last, further consolidating the lysosomal origin of the mitochondrial stress response, chronic

treatment with LLOMe also engendered nucleoid aggregation (Fig. 3G-H). Finally, robust

mitochondrial nucleic acid stress responses were also detected in wild-type MCF7 cells treated with

Pb or LLOMe (Fig. 3I-J). Taken together, the above data demonstrate that chronic lysosome damage

triggers mitochondrial stress responses.

Palbociclib treatment triggers mitochondrial nucleic acid release into the cytosol and interferon

responses

Given the detection of mitochondrial nucleic acid stress upon Pb-mediated lysosome damage, we next

sought to directly detect immunostimulatory nucleic acid release from mitochondria. We prepared

mitochondrion-free cytosolic extracts (Fig. 4A) from cell clones deleted for $\Delta ATG5$ or rescued by

reintroduction of *GFP-ATG5* (Newman et al., 2017); use of these paired lines diminished potential

clonal variation that might have obscured subtle mitochondrial nucleic acid fluctuations. Cytosolic

mitochondrial DNA and RNA were found to be elevated in the cytosol in response to chronic Pb; again,

this was more prominent in $\Delta ATG5$ cells where lysosome damage is more poorly resolved (Fig. 4B-C).

Given that Pb is known to induce interferon responses, we set out to interrogate whether lysosome

damage played a role (and potentially thus cytoplasmic accumulation of mitochondrial nucleic acids).

Firstly, we first established that chronic Pb triggers interferon signalling, analysing hallmarks such as

phosphorylation of STAT1 and transcriptional activation of type III interferons, i.e., *IFNL1*, and

interferon response genes (IRGs), i.e., *ISG15*, *IFIT1* and *CXCL10*. Pb indeed induces phosphorylation of

STAT1 (Fig. 5A-B, Supp. Fig. 2) and transcriptional activation (Fig. 5C) in A549 cells over 48 h, most

prominently in $\Delta ATG5$ cells, correlating with the persistent lysosome damage known to occur here.

Furthermore, phosphorylation of STAT1 was seen when testing lower doses of Pb akin to steady-state

plasma concentrations achieved during clinical treatment, although with greater latency (one week)

(Fig. 5D). Also corroborating the link with lysosome damage, chronic LLOMe treatment also induced

interferon responses over 48 h (Figs. 5E-F). Finally, wild-type MCF7 cells also exhibited interferon

responses to chronic treatment with either Pb or LLOMe (Fig 5G-J). Taken together, the above data indicate that chronic lysosome damage is a contributor to induction of interferon responses.

Mitochondrial nucleic acids are required for the interferon response to chronic lysosome damage

To test whether the observed release of mitochondrial nucleic acid triggers the interferon signalling consequent from Pb-induced lysosome damage, we next depleted mitochondrial genomes from A549 control (Rescue) and $\Delta ATG5$ cells, via chronic dideoxy-cytidine treatment, generating "Rho0" cultures (Supp. Fig. 3A). Importantly, mitochondrial nucleic acid depletion inhibited engagement of downstream markers of interferon production by Pb, including phosphorylation of STAT1 (Fig. 6A-B), and transcriptional upregulation of *IFNL1* and *ISG15* (Fig. 6C). Two modes of nucleic acid efflux from mitochondria have been postulated in the prior literature. The first is associated with stereotypical markers of mitochondrial nucleic acid stress, as observed in Fig. 3, and can be blocked using the VDAC channel inhibitor VBIT-4 (West et al., 2015; Kim et al., 2019; Torres-Odio et al., 2021). The second is characterised by a requirement for mitochondrial BAK and BAX pore-forming proteins (McArthur et al., 2018; Riley et al., 2018). We found that interferon responses to Pb were indeed abrogated by VBIT-4 treatment (Figs. 6D-E) but not *BAK* and *BAK* double knockout (Supp. Fig. 3B). This observation further corroborates the link between the mitochondrial nucleic acid stress induced by lysosome damage and interferon signalling. In line with this, chronic LLOMe treatment was also unable provoke phosphorylation of STAT1 in Rho0 cells (Fig. 6F-G). Taken together with our prior findings, the above data demonstrate that lysosome damage triggers accumulation of mitochondrial nucleic acid in the cytosol, and that this drives interferon responses. However, Pb also reportedly triggers interferon responses by suppressing CDK4/6-mediated silencing of nuclear genome-derived dsRNA (Goel et al., 2017). Given that we observe that CDK4/6-independent lysosome damage drives mitochondrial nucleic acid release, we hypothesised co-operation between on-target (kinase activity) and off-target (lysosomal-mitochondrial) effects of Pb. Indeed, *CDK4/6* deletion, while insufficient to engage robust

interferon responses alone, synergised with Pb treatment to induce a maximal response (Fig. 6H-I).

Taking the above data together, we conclude that sustained suppression of CDK4/6 kinase activity has

the potential to engage interferon signalling via non-mitochondrial routes (Goel et al., 2017), but that

the lysosome damage-mitochondrion axis must co-operate to fully realise the interferon effect, at

least in the context of Pb treatment.

Discussion

Our findings support the following model (Fig. 7). In response to chronic exposure to the

lysosomotropic cancer therapeutic Palbociclib (Pb), persistent lysosome rupture is invoked. Acute

lysosome damage typically leads to repair, removal and/or replacement of the damaged organelle.

These responses are at the core of the ELDR (the endolysosome damage response). We propose that

in cells where chronic lysosome damage is insufficient to trigger cell death, propagation of stress

signalling via damaged lysosome-mitochondrion communication leads to accumulation of cytosolic,

self-derived immunostimulatory nucleic acids, triggering interferon signalling. Indeed, other recent

data have hinted that stress signals emanating from acutely damaged lysosomes trigger wider changes

in cell physiology (Jia et al., 2022; Neuwirt et al., 2023; Duran et al., 2024). Thus, it is possible that the

lysosomal-mitochondrial signalling axis described here represents only one of several outward-facing

stress relays engaged by sublethal lysosomal perturbation (i.e., not necessarily impacting directly on

lysosomal repair, removal or replacement as part of the core ELDR). In the case of Pb treatment, the

mitochondrial response to lysosome damage is required for interferon signalling but maximal

stimulation occurs in co-operation with the known CDK4/6-inhibitory activity of the drug, explaining

fully this important anti-cancer facet of Pb action. Instructively, mitochondrial nucleic acids may

similarly be required to license interferon responses after *Salmonella* infection (Xu et al., 2022). This

is suggestive when one considers that bacterial phagosome damage by infectious agents can also

engage selective autophagy mechanisms with molecular parallels to lysophagy (Thurston et al., 2012).

Furthermore, it has been recently suggested that damage to lysosomes invoked by pathogens effects broad changes in mitochondrial proteome, secondary to vesicular rupture (Bussi et al., 2022). Taking these observations together, we speculate that the mitochondrial responses to lysosome damage may have initially evolved to consolidate infection responses by priming interferon responses, but not in itself triggering maximal signalling, i.e., co-operating with additional stimuli such as nucleic acids or other patterns physically derived from the infectious agent. However, these ideas require future exploration.

Our data show that the lysosome damage-induced interferon response depends on chronic perturbation. For instance, A549 cells deleted for Δ*ATG5*, which are deficient in the ELDR, show a more robust response to Pb and LLOMe, which correlates with more sustained lysosome damage. Similarly, wild-type MCF7 cells show evidence of resolution of lysosome damage but more slowly than wild-type A549 cells, and this again correlates with a more readily detectable interferon response. However, it is possible that other functions of *ATG5* in autophagy pathways, for instance degradation of PRRs, could also contribute to the observed *ATG5*-dependency in A549 cells (Nakahira et al., 2011; Mathew et al., 2014; Prabakaran et al., 2018; He et al., 2019). Our data nonetheless still establish a pathway between lysosome damage and engagement of interferon responses via mitochondrial release of nucleic acid.

Finally, our results highlight an important consideration when investigating small-molecule therapeutics for cancer. The lysosome is sometimes considered a passive problem for therapy, in that it sequesters basophilic drugs from their site of action (Llanos et al., 2019). Although the mitochondrion is implicated here in response to Pb, it is unlikely that it will ultimately prove to be the only consequential signalling hub responding to chronic lysosome damage. Indeed, we would suggest that off-target signalling effects of cancer therapeutic drugs consequent from chronic accumulation in

the lysosome should be factored into cell-based drug action studies in future. This will facilitate holistic understanding of therapeutic compound action and thus, ultimately, promote refinement of clinical applications.

Figures

Figure 1   Palbociclib induces lysophagy and chronic lysosome damage

(*Legend overleaf*)

Figure 1 Palbociclib induces lysophagy and chronic lysosome damage

A) Schematic of SRAI (signal-retaining autophagy indicator)-Galectin 8 (GAL8) reporter for lysophagy. In the cytosol, YPet emission is low because TOLLES acts as a FRET donor. Conversely, upon completion of lysophagy, YPet is quenched by low pH and TOLLES fluorescence is apparent.

B-C) A549- or MCF7-SRAI-GAL8 cells were treated with DMSO, 0.7 mM LLOMe or 10 μM Palbociclib (Pb) for 24 h then imaged by widefield fluorescence. **B)** Representative A549 images (also see **Supp. Fig. 1**). **C)** Quantifications of TOLLES:Ypet (lysophagy flux) indices (n = 3, ± S.E.M., 50-100 cells/replicate, *** = p < 0.001, **** = p < 0.0001, 1-way ANOVA , Holm-Šídák comparisons vs. DMSO).

D-E) A549 NT (non-targeted) or $\Delta ATG5$ cells expressing mRFP-GFP-GAL8 were treated with DMSO or 10 μM Pb for 24 h prior to imaging by widefield fluorescence. **D)** Representative images, dotted lines demarcate zoom areas. **E)** Quantifications of GFP$^-$ mRFP$^+$ galectolysosomes (n = 3, ± S.E.M., > 90 cells/condition, ** = p < 0.01, * = p < 0.05, 2-way ANOVA, Holm-Šídák comparisons).

F-G) A549 NT or $\Delta ATG5$ cells expressing YFP-GAL8 were treated with DMSO or 10 μM Pb for 48h. **F)** Representative widefield images, dotted lines demarcate zoom areas, arrows indicate foci. **G)** Representative confocal fluorescence images of LAMP2 co-staining, arrows indicate co-localisation foci in zoom area, dotted line delineates the intensity trace (right).

H-K) A549 NT, A549 $\Delta ATG5$ cells and MCF7 cells expressing YFP-GAL8 were treated for indicated time points with 10 μM Pb or 0.7 mM LLOMe, prior to imaging by widefield fluorescence. YFP-GAL8 foci quantifications are presented in **H, I** and **K** (n = 3, ± S.E.M., > 100 cells/condition, **** = p < 0.0001, *** = p < 0.001, ** = p < 0.01, * = p < 0.05, coloured asterisks indicate significance vs. 0 h for a given treatment, black asterisks indicate significance between different treatments, mixed-effects analysis, Holm-Šídák comparisons). Representative images of MCF7 cells at 24h shown in **J**.

L-M) A549 $\Delta ATG5$ cells or $\Delta ATG5$ $\Delta CDK4$ derivatives, stably expressing Cas9 and YFP-GAL8, were transduced with non-targeting sgRNA (gNT) or g*CDK6*. 72 h post gRNA transduction, cells were treated with DMSO (-) or 10 μM Pb (+), for 24h and either **L)** immunoblotted or **M)** fixed for widefield fluorescence microscopy quantification of YFP-GAL8 foci, comparing *CDK4/6* intact controls ($\Delta ATG5$ gNT) with *CDK4/6* double knockout cultures ($\Delta ATG5$/g$\Delta CDK4$ gCDK6) (n = 3, ± S.E.M., > 110 cells/condition, ** = p < 0.01, * = p < 0.05, 2-way ANOVA, Holm-Šídák comparisons).

Scale bars = 20 µm.

Figure 2   Damaged lysosomes are associated with the mitochondrial network

(*Legend overleaf)*

Figure 2   Damaged lysosomes are associated with the mitochondrial network

A - C) A549 NT (non-targeted) or $\Delta ATG5$ cells were treated with DMSO or 10 μM Palbociclib (Pb) for 48 h then analysed by TEM. Representative images shown in **A)**, dotted lines delineate zoom area showing swollen lysosomes contacting mitochondria (arrowheads delimit a contact site, defined as continuous membrane opposition of < 30 nm). **B)** The proportional cross-sectional area of lysosomes within the cytoplasm. **C)** The mean length of lysosome-mitochondria (LM) contact sites per unit area of cytoplasm (n = 20 images, ± S.D., * = p < 0.05, **= p < 0.01, *** = p < 0.001, 2-way ANOVA, Holm-Šídák comparisons). Scale bars, 0.5 μm.

D - E) A549 $\Delta ATG5$ cells expressing YFP-GAL8 were treated with 10 μM Pb for 24 h, then stained for TOM20 (mitochondrion) and imaged by Super Resolution Optical Photon Reassignment Microscopy (SoRA). **D)** Representative raw images in upper panels, dotted lines (1-3) delineate the zoom areas in the row below and the cognate mitochondrial surface rendered images in the bottom row. **E)** Relative frequency distribution of distances between YFP-GAL8 foci and rendered mitochondria (Actual), compared with *in silico* shuffling of these foci within the cytoplasm (Random) (n = 3, 400 measurements from 5 cells, Kolmogorov-Smirnov test, p < 0.0001, representative of 3 independent experiments). See also **Supp. Movie 1**. Scale bars, 5 μm (merge), 1 μm (zooms).

F) A549 $\Delta ATG5$ cells expressing YFP-GAL8 were treated with 10 μM Pb for 6 h, then MitoTracker-Red added 30 mins prior to super-resolution radial fluctuation (SRRF) processing-facilitated time-lapse microscopy. White arrows: YFP-GAL8 positive membranes associated with the mitochondrial network. See also: **Suppl. Movies 2 and 3**. Scale bars, 2 μm. Representative of 4 experiments.

Figure 3   Lysosome damage triggers stereotypical mitochondrial nucleic acid stress markers

A-F) A549 NT (non-targeting wild-type) or ΔATG5 cells were treated with DMSO or 10 μM Palbociclib (Pb) for 48 h. Cells were stained for mitochondria (TOM20) and mitochondrial nucleoids using **A,C)** anti-TFAM; **B,D)** anti-DNA; **E,F)** anti-FASTKD2 antibodies. **A,B)** Representative SoRA and **E)** confocal microscopy images. **C-D,F)** Wide-field fluorescence microscopy was used to quantify the aggregated nucleoids (TFAM foci with > 0.7 μm^2 cross-sectional area or prominent DNA/FASTKD2 foci) (n =3, ± S.E.M., ns = non-significant, > 250 cells/condition, * = $p < 0.05$, **= $p < 0.01$, **** = $p < 0.0001$, 2-way ANOVA, Holm-Šídák comparisons DMSO vs. Pb). Dotted lines delineate zoom areas.

G-H) A549 NT or ΔATG5 cells were treated with DMSO or 700 μM LLOMe for 24 h. Cells were analysed for aggregated mitochondrial nucleoids as in **A-F**, quantifications presented here (n =3, ± S.E.M, > 220 cells/condition, ns = non-significant, * = $p < 0.05$, ** = $p < 0.01$, **** = $p < 0.0001$, 2-way ANOVA, Holm-Šídák comparisons DMSO vs. LLOMe).

I-J) MCF7 cells were treated with DMSO, 700 μM LLOMe or 10 μM Pb, for 24 h, then analysed for aggregated TFAM-positive nucleoids as in **A** and **C. I)** Representative widefield fluorescence images. **J)** Quantifications (n = 3, ± S.E.M, > 150 cells/condition, * = $p < 0.05$, t-tests vs. DMSO).

Scale bars, 10 μm.

Figure 4   Mitochondrial nucleic acids are released into the cytosol by Palbociclib

A) A549 Rescue (ΔATG5 + GFP-*ATG5*) and ΔATG5 cells were treated with DMSO (-) or 10 μM Palbociclib, Pb (+), for 48 h. Pure cytosolic fraction (Cytosol) was separated from remaining cellular material including mitochondria (Insoluble) using digitonin extraction and immunoblotted to confirm purity (mitochondrial markers: TOM20; Cytochrome C; ATP5A, ER marker: Calnexin)

 B-C) Nucleic acids (**B**, DNA; **C**, RNA) were isolated from the cytosolic fraction and qPCR performed for mitochondrial genes *ND1*, *ND6*, *CYTB* and *COII*, normalised to *ACTB* from whole cell DNA (n >= 4 after outliers removed by Grubb's test, ± S.E.M., normalisation to Rescue + DMSO, ** = p < 0.01, *** = p < 0.001, ratio paired t-tests).

Figure 5   Chronic lysosome damage triggers interferon responses

(*Legend overleaf*)

Figure 5   Chronic lysosome damage triggers interferon responses

A-B) A549 NT (non-targeted wild-type) or Δ*ATG5* cells were treated with 10 μM Palbociclib (Pb) for the indicated time and then immunoblotted for interferon pathway activation (phospho-Y701 STAT1, P-STAT1, and ISG15). **A)** Representative blot. **B)** Quantifications (n = 3, ± S.E.M., outlier removed by Grubb's test, *** = p < 0.001, 2-tail t-tests vs. 0 h, Holm-Šídák correction).

C) A549 NT or Δ*ATG5* cells treated as in **A** and subjected to qRT-PCR for indicated genes (n = 3, ± S.E.M., relative to *18S*, normalised to NT + DMSO, ns = non-significant, ** = p < 0.01, *** = p < 0.001, 1-sample t-test for NT, 2-tailed t-test for Δ*ATG5*).

D) A549 Δ*ATG5* cells were treated with DMSO (-) or 500 nM Pb (+), for the indicated number of days (d), then immunoblotted for P-STAT1 (n = 3, representative replicate shown).

E-F) A549 NT (non-targeted wild-type) or Δ*ATG5* cells were treated with 700 μM LLOMe for the indicated times then immunoblotted for interferon pathway activation (P-STAT1, ISG15). **E)** Representative blot. **F)** Quantifications (n = 3, ± S.E.M., * = p < 0.05, 2-tail t-tests vs. 0 h, Holm-Šídák correction).

G-J) MCF7 cells were treated with 10 μM Palbociclib (Pb) (**G-H**) or 700nM LLOMe (**I-J**) for the indicated times then immunoblotted as shown. **G,I)** Representative blots. **H,J)** Quantifications (n = 3, ± S.E.M., * = p < 0.05, ** = p < 0.01, 1-sample t-tests).

Figure 6   Mitochondrial nucleic acids are required for interferon response to chronic lysosome damage

(*Legend overleaf*)

Figure 6   Mitochondrial nucleic acids are required for interferon responses to chronic lysosome damage

A-C) A549 Rescue (ΔATG5 + GFP-ATG5) and ΔATG5 cells were treated to generate control (Rho+, +) or mitochondrial genome depleted (Rho0, 0) cells. Cells were treated with DMSO or 10 μM Palbociclib (Pb) for 48 h and immunoblotted for interferon activation (phospho-Y701-STAT1, P-STAT1, ISG15). **A)** Representative blot. **B)** Quantification for Pb-treated ΔATG5 cells (n = 3, ± S.E.M., normalised to Rho+, ** = p < 0.01, ns = non-significant, 1-sample t-test).

C) A549 ΔATG5 cells (Rho+ or Rho0) were treated with 10 μM Pb for 48 h and qRT-PCR performed for *IFNL1* and *ISG15* (n = 3, ± S.E.M., relative to *18S*, normalised to Rho+, ** = p < 0.01, **** = p < 0.0001, 1-sample t-tests).

D-E) A549 Rescue or ΔATG5 cells were treated with DMSO or 10 μM Pb +/- 15 μM VBIT-4, then immunoblotted as shown. **D)** Representative blot. **E)** Quantification for Pb-treated ΔATG5 cells (n = 3, ± S.E.M, ns = non-significant, **** = p < 0.0001, 2-way ANOVA, Holm-Šídák comparisons DMSO vs. VBIT-4).

F-G) A549 ΔATG5 cells, Rho+ (+) or Rho0 (0), were treated with DMSO or 700 μM LLOMe for 48 h, then immunoblotted for P-STAT1. **F)** Representative blot. **G)** Quantification for LLOMe treated cells, normalised to Rho+ (n = 3, ± S.E.M., * = p < 0.05, 1-sample t-test).

H-I) A549 ΔATG5 cells or ΔATG5 ΔCDK4 derivatives, stably expressing Cas9, were transduced with non-targeting sgRNA (gNT) or g*CDK6*. 72 h post transduction, cells were treated with DMSO (-) or 10 μM Pb (+) for 24h, then immunoblotted as shown. **H)** Representative blot. **I)** Quantifications comparing CDK4/6 intact controls (ΔATG5 gNT) with CDK4/6 double knockout cultures (ΔATG5/ΔCDK4 g*CDK6*) (n = 3, ± S.E.M., ** = p < 0.01, * = p < 0.05, 2-way ANOVA, Holm-Šídák comparisons).

Figure 7   Model for response to chronic lysosome damage invoked by Palbociclib

See text for details. GAL8 = galectin 8. IFN = interferon. Blue dots depict mitochondrial nucleoids.

Acknowledgements

Work in the SW lab is funded by a Cancer Research UK (CRUK) Senior Fellowship (C20685/A29576). MB is funded by a CRUK studentship. TEL is funded by a CRUK TRACC clinical studentship. AW and LM are funded by the Medical Research Council. We thank Richard Youle, Viktor Korolchuk, Noor Gammoh, Felix Randow for reagents. We thank Noor Gammoh and Alex von Kriegsheim for feedback. We thank Steven Mitchell for assistance with electron microscopy. For the purpose of open access, the author has applied a CC-BY public copyright licence to any Author Accepted Manuscript version arising from this submission.

Author contributions

Study conception and funding acquisition by SW. Experiments designed by MB and SW. Experiments performed by MB, with assistance from AJK, TEL, KWI and NJM. Refinement of mitochondrial nucleic acid analysis and mitochondrial genome depletion experiments by AD. FIJI macro for GAL8-mitochondrion image analysis written by LM. Time-lapse video microscopy by AW. All data analysis by MB. Manuscript drafted by MB and SW. Editing performed by MB, NJM and SW.

Methods

Cell culture and cell lines

Cells were cultured in DMEM supplemented with 10% FBS, 2 mM L-glutamine and 10 U/ml 10 μg/ml penicillin/streptomycin, at 37 °C under 5% CO_2. At monthly intervals all cells in culture were tested for the presence of mycoplasma. A549 NT, $\Delta ATG5$ and Rescue ($\Delta ATG5$ + GFP-ATG5) lines were described previously by the authors and are a derivative of the A549:EcoR parental cell line, which is A549 human lung adenocarcinoma cells derivatised to express Ecotropic receptor to enable transduction with retrovirus with murine tropism (Newman et al., 2012). A549:EcoR and MCF7 breast carcinoma lines were from the authors' laboratory and were verified by microsatellite genotyping. YFP-GAL8 and YFP-GAL8 R69H R232H expressing retroviruses (see **Plasmids**) were used to generate stable A549 line derivatives as described previously (Newman et al., 2012); pools of transductants were selected with 10 μg/ml blasticidin for 10 days. A549-EcoR cells stably expressing mRFP-GFP-GAL8 were made by lentivirus transduction (see **Plasmids**) and a pool of transductants selected with 10 μg/ml blasticidin. A549-EcoR and MCF7 cells stably expressing SRAI-GAL8 were created by lentiviral transduction (see **Plasmids**) and selection with 10 μg/ml blasticidin. Then, cell populations expressing high YPet and TOLLES fluorescent levels were enriched using a FACSAria II (BD Bioscience, San Jose, CA). To enable both pooled and single clone CRISPR/Cas9 knockouts in A549-EcoR cell lines, stable expression of Cas9 was achieved by transducing lentivirus made from LentiCas9-Blast template and selection with 10 μg/ml blasticidin. CRISPR/Cas9-mediated knockout in the resulting lines expressing Cas9 was performed using lentiviral delivery of gRNA. Non-targeting/targeting sgRNA was cloned into lentiGuide-puro for lentivirus production, and transductants selected for with 2.5 μg/ml puromycin. gRNA sequences are tabulated in Methods Table 1. Where single cell clones were used, this was achieved via limiting dilution. All lentivirus was packaged in HEK293T cells according to standard protocols; briefly, viral vector was co-transfected with pMD2.G and psPAX2 packaging plasmids using Lipofectamine 2000.

Target name	Target
GFP (NT)	GTGAACCGCATCGAGCTGAA
Luciferase (NT-2)	AACGGCGGCGGGAAGTTCAC
MAVS	TCTTCAATACCCTTCAGCGG
CDK4	TGTAGTGATGCGCGACGACA
CDK6	CTATTACTGAGCTCGATGTG
BAX	GGCGGTGATGGACGGGTCCG
BAK	GGAACTCTGAGTCATAGCGT

Methods Table 1 gRNA target sequences for Lentiguide-mediated targeting in A549

Drug treatments

Seeding density was normalised per experiment treatment requirements. Cells were plated to be approaching confluency, but not yet confluent, at the point of analysis. Typically, one third of the number of cells to be treated with higher doses (e.g. 10 μM) Pb (Pb-HCL, Cayman #16273, dissolved in DMSO) were seeded relative to the cognate DMSO vehicle treatments for 48 h treatments, and one half for 24 h treatments, to compensate the cytostatic effect of Pb. For 10 μM Pb treatment, treatment was applied once for the entire duration. For longer term treatments using 500 nM Pb, cells were plated on day 0, treatment was initiated on day 1, and medium was refreshed with new drug or vehicle at day 4. LLOMe (LLOMe-HCL, Cayman #16008, dissolved in DMSO) was dosed at 700 μM. VBIT-4 (Selleckchem #S3544) was dissolved in DMSO and used at 10 μM.

Mitochondrial genome depleted cultures

To generate paired Rho+ and Rho0 cultures, cells were cultured in standard medium supplemented with 200 μM uridine (Sigma #U3750) and, for Rho0 culture, with an additional 10 μM ddC (2',3'-dideoxycytidine, Sigma #D5782) for 14 consecutive days (Tigano et al., 2021). Cells were then maintained in culture for a maximum of another week in medium still supplemented with ddC.

Plasmids

See Methods Table 2 for full list. Where Gateway cloning was performed this was as per standard protocols using BP clonase II and LR clonase II (Invitrogen). sgRNA cloning into lentiGuide-puro vector was performed using restriction digestion as per the standard protocols available from Addgene. Detailed sequence maps are available from authors upon request.

Name	Insert	Vector	ID and/or Source
pSpCas9(BB)-2A-Puro (PX459) V2.0	-	-	Addgene #62988. A gift from Feng Zhang (Ran et al., 2013).
gRNA *ATG5*	GAGATATGGTTTGAATATGA targeting sequence	gRNA cloning vector	(Newman et al., 2012)
pLX304 SRAI-DEST-C1-BLAST	-	-	(Jimenez-Moreno et al., 2023)
pLX304 SRAI-GAL8 BLAST	GAL8	pLX304 SRAI-DEST-C1-BLAST	This study; Gateway cloning from pDONR GAL8
pDONR GAL8	GAL8	pDONR223	This study; Gateway cloning from pANT7_cGST-LGALS8 (DNASU collection HsCD00079457), this study.
pLenti CMV DEST puro	-	-	Addgene #17452, A gift from Eric Campeau & Paul Kaufman (Campeau et al., 2009).
pLX304-DEST	-	-	pLX304 was a gift from David Root (Addgene plasmid # 25890) (Yang et al., 2011).
pLX304-mRFP-GFP-DEST-C1-BLAST	-	-	This study; permits C-terminal fusion of ORF to mRFP-GFP by Gateway cloning; map available on request from authors
pLX304 - mRFP-GFP-Gal8	mRFP-GFP-Gal8	pLX304-mRFP-GFP-DEST-C1-BLAST	This study; Gateway cloning from pDONR GAL8
pMSCV-YFP-Gal8	YFP-Gal8	pMSCV-Blasticidin	(Thurston et al., 2012)
lentiCas9-Blast	-	-	Addgene #52962, A gift from Feng Zhang.
lentiGuide-puro	-	-	Addgene #52963, A gift from Feng Zhang.
pMD2.G	-	-	Addgene #12259, A gift from Didier Trono
psPAX2	-	-	Addgene #12260, A gift from Didier Trono

Methods Table 2   Plasmids used in this study and their origins

Immunoblotting and antibodies

Cells were lysed in 4% SDS lysis buffer (4% SDS, 50mM Tris-HCl pH 7.5 and 150mM NaCl), followed by pulse sonication. Protein concentration was measured by BCA protein assay, using the Pierce™ BCA protein assay kit (Thermo Fisher, #23225). Normalised protein samples were diluted in Laemmli buffer (4% SDS, 20% glycerol, 10% 2-mercaptoethanol, 0.125 M Tris-HCl pH 6.8, bromophenol blue) and briefly heated to 95 °C. SDS-PAGE was carried out using either 4-12% NuPage™ Bis-Tris Precast mini, or 4-12% Novex™ Tris-Glycine precast midi gels (Thermo Fisher), in 1x NuPage™ MOPS SDS running buffer (Thermo Fisher) or Pierce™ Tris-Glycine SDS running buffer (Thermo Fisher), respectively. Wet transfer was used to transfer protein from mini gels, performed onto 0.2 μm nitrocellulose membrane (GE Life Sciences), and semi-wet transfer system (Trans-blot turbo, Bio-Rad) was used to transfer proteins from midi gels. Membranes were blocked in 5 % BSA (Sigma) in PBS for 1 h at room temperature (RT). All washes were performed three times in TBS - 0.1 % Tween 20 (TBST), for 10 min each at RT. Primary antibody dilution was in 0.1 % BSA in PBS with 0.05% sodium azide, typically at 1:1000. Membranes were incubated in primary antibody solution overnight at 4 °C. The following antibodies were used: anti-ATG5 (Sigma, #A0731), anti-Calnexin (CST, #2679), anti-CDK4 (CST, #12790), anti-CDK6 (CST, nat.no. 3136), anti-ISG15 (CST, #2743), anti-MAVS (CST, #3993), anti-phosho-Y701-STAT1 (CST, #9167), anti-STAT1 (CST #9176), anti-TOM20 (Santa Cruz, #sc-17764), anti-Cytochrome C apoptosis antibody cocktail (Abcam, #ab110415) and anti-α-Tubulin (Sigma, #T9026). Secondary antibody solution was 0.1 % powdered milk in TBST in 1:3000 dilution. Membranes were incubated with cognate HRP-linked secondary antibody (CST) for 1 h at RT. For detection, membranes were incubated in ECL reagent (ECL prime, Western blotting reagent, GE Life Sciences) for 5 min at RT, or ECL reagent (1M Tris-HCl pH 8.5, 250 mM luminol, 90mM coumaric acid and 0.002% H_2O_2) for 1 min at RT, followed by film exposure and development (AGFA automated developer), or chemiluminescent imaging (Amersham ImageQuant 800, software v1.2.0). Relative protein band intensities were quantified using FIJI (v2.1.0). For P-STAT1, this is given as a ratio to tubulin, as STAT1 itself is an interferon stimulated gene.

Immunostaining

For immunofluorescence experiments, cells were typically seeded onto round 16 mm cover glass (Fisher Scientific) except for super-resolution microscopy where cells were seeded on high-precision cover glass (Zeiss™, Fisher Scientific). Cells were fixed in 4% paraformaldehyde (PFA) / PBS for 10 min at RT and washed once in PBS. Permeabilization was performed for 10 min at RT using 0.25% Triton X-100 (Sigma) / PBS, followed by another PBS wash. Primary and secondary antibody solution was diluted in PBS / 1% BSA, at 1:200 (except p62 used at 1:1000). Secondary antibody solution typically contained DAPI at 1:2500. Cells were incubated with primary antibody solution for 1 h, at RT, followed by 3 x 5 min PBS washes. The following antibodies were used: anti-DNA (EMD Milipore, #CBL186), anti-FASTKD2 (Proteintech, #17464-1-AP), anti-LAMP2 (Abcam, #ab25631), anti-TFAM (CST, #8076), anti-TOM20 (CST, #42406) (used when co-staining with mouse antibodies), anti-TOM20 (Santa Cruz, #sc-17764) (used when co-staining with rabbit antibodies and YFP-GAL8) and anti-DNA (Merck-Millipore, #CBL186). Secondary antibody was cognate fluorophore linked secondary antibody (Thermo Fisher, Alexa Fluor). Secondary antibody solution was applied to cells for 45 min, at RT, followed by 3 x 5 min PBS washes. Regular cover glass was mounted onto a microscope slide using DAKO fluorescence mounting medium. High-precision cover glass super-resolution microscopy was mounted using Fluoromount™ aqueous mounting medium (Sigma).

Fluorescence Microscopy

For widefield immunofluorescence microscopy, images were taken with Olympus BX51 fluorescent microscope (Cell^F software). For confocal microscopy, images were taken using a Nikon A1R laser-scanning confocal microscope. Images are presented as maximum projection Z-stacks. Super-resolution images were acquired using Super Resolution Optical Photon Reassignment Microscopy on a Nikon Eclipse Ti2 inverted microscope stand and a SR Apo TIRF 100x 1.49NA oil lens (Nikon Instruments). The CMOS cameras used for acquisition were Teledyne Photometrics Prime 95B

(Teledyne Photometrics 3440 E.Britannia Drive, Tucson AZ) and 405, 488, 561, 640nm laser lines. Acquisition of images and deconvolution was carried out using Nikon NIS-Elements Advanced Research software (v5.21-03). For imaging SRAI-GAL8, cells were plated on glass bottom 96 well plates (P96-1.5H-N, Cellvis) at 70-90% confluency. Cells were fixed with 4% paraformaldehyde in PBS, pH 7.2 at room temperature for 10 mins. Cells were then washed twice with PBS and kept at 4 $^{\circ}$C. Cells were imaged 1-3 days after fixation while immersed in PBS. Images were captured with a Nikon Eclipse Ti-E inverted microscope (Nikon Instruments UK Ltd, Kingston Upon Thames, UK) using a 40X objective. Widefield illumination for this was provided by a CoolLED PE-4000 LED light source (CoolLED Ltd, Andover, UK) combined with LED-CFP/YFP/mCherry-3X-A (Pinkel) filter sets (IDEX Health & Science, LLC, Center of Excellence, 1180 John Street, Rochester, New York 14586, United States). Images were acquired with a Photometrics Prime BSI sCMOS camera (Teledyne Photometrics, 3440 E. Britannia Drive, Suite 100, Tucson, AZ 85706). Conditional imaging was setup and controlled using the Nikon Nis-Elements JOBS module. The JOB (script) incorporates automated two wavelength capture of 3D images (Z stacks for CFP and YFP filters) from an array of imaging sites (XY positions) using a motorised XY stage within user selected wells (a total of 7 fields per well were imaged). SRAI-GAL8 images are shown as z-projections from 5 z-steps.

SRRF Time-lapse videomicroscopy

Time-lapse videomicroscopy was performed by exchanging full medium for phenol-red free medium containing 0.132 µg/ml MitoTracker Red and fresh Pb 30 mins prior to imaging. Images were acquired on the multimodal Imaging Platform Dragonfly (Andor technologies, Belfast UK) equipped with 405, 445, 488, 514, 561, 640 and 680nm lasers built on a Nikon Eclipse Ti-E inverted microscope body with Perfect focus system (Nikon Instruments, Japan). Super Resolution Radial Fluctuations (SRRF) data were acquired using the iXon 888 EMCCD camera and a 100x SR Apo TIRF 1.49NA objective lens. Environmental control of the cultured cells was maintained during imaging with an Okolab bold line

stage top incubation chamber incorporating temperature and humidified CO2 control (Okolab S.R.L, Ottaviano, NA, Italy).

Transmission electron microscopy (TEM)

Fixative solution was prepared by mixing fresh aqueous 16% paraformaldehyde (from methanol-free powder, Sigma) and fresh 25% EM grade glutaraldehyde (Merck) to final concentration 2% paraformaldehyde, 2.5% glutaraldehyde in 0.1M PHEM buffer (120 mM PIPES, 50 mM HEPES, 4 mM $MgCl_2$ and 20 mM EGTA, pH 6.9). Two 15 cm dishes per condition of cells were used and cells pelleted by centrifugation 200g for 3 min in a 15 mL tube. Cells were resuspended in the fixative solution and incubated for 20 min at room temperature, on a slow-motion roller bank. Cells were then centrifuged, supernatant removed, and fresh fixative solution added. The second fixative solution was incubated with cells for 4 h at room temperature, on a slow-motion roller bank. The cells were washed 3 times with 0.1M PHEM buffer and stored in 0.1M PHEM until embedding in EPON resin at 4 °C (2-3 days). Sections were co-stained with uranyl acetate and lead citrate to provide membrane contrast. TEM images were taken by JEOL JEM-1400 series, 120 kV transmission electron microscope.

Image analyses

From SRAI-GAL8 images, a TOLLES:YPet index was obtained using an ImageJ macro that calculates the percent area of pixels with TOLLES:YPet ratio above a given ratio over the total area that the fluorophore occupies within the field (Jimenez-Moreno et al., 2023). A ratio of 1.75 was used. Substructures in cells (foci, lysosomes, nuclei) were generally counted manually using ImageJ cell counter plug-in. In the specific instance of TFAM nucleoid aggregates, these were counted using the ImageJ Particle Analysis tool (automated particle count with a minimum detected particle area threshold of 0.7 μm^2. Parameters from TEM images (cytosol area, lysosome area, mitochondria-

lysosome contact length) were measured using the ImageJ ROI manager. The number of galectolysosomes formed in mRFP-GFP-Galectin-8 cells were determined by counting GFP quenched, mRFP positive foci using the ImageJ MitoQC plug-in (Montava-Garriga et al., 2020).

Images obtained by SoRa were 3D deconvoluted using NIS elements imaging software (Nikon, v5.21-03) and further analysed using Imaris software (v9.9.1). YFP-GAL8 foci were modelled as centroids. Similarly, the mitochondria network surface masking was applied according to the intensity of TOM20 fluorescent signal. The same modelling settings were used on all images acquired. Standard Imaris measurements were extracted showing the distance from the YFP-GAL8 foci to the rendered mitochondrial network, measured from the centre of modelled YFP-GAL8 centroids to the outer surface of the modelled mitochondrial network, with the radius of the centroid then subtracted. To assess association of YFP-GAL8 foci with the mitochondrial network, they were compared with a shuffled control. In these controls, an ImageJ plug-in was created to randomly shuffle the YFP-GAL8 foci within the cytosolic area on a per cell basis (cytosolic area as outlined by the reach of the mitochondrial network).

Cytosol fractionation for nucleic acid quantification

Methods were adapted from previously published protocols (Dhir et al., 2018; Bryant et al., 2022). Cells were trypsinised and fresh medium added to inactivate trypsin, then centrifuged at 200 g at 4 °C. The supernatant was then aspirated, and cells washed by gentle pipetting in 1 ml of ice-cold PBS. The centrifugation and PBS wash was repeated once more. The pellet was then resuspended in 500µl (per 60mm dish) of ice-cold fractionation buffer (50 mM HEPES pH 7.4, 150 mM NaCl, 25 µg/ml fresh digitonin (Sigma, #11024-24-1) and 1 X Complete Protease Inhibitor Cocktail with EDTA, Roche). When RNA was to be isolated from the cytosolic fractions 4 U RNase inhibitor (Promega, #N2111) was added

to the fractionation buffer. The suspensions were incubated end-over-end for 10 mins at 4 °C to allow selective plasma membrane permeabilization, then centrifuged at 1000 g for 3 min to pellet cells (insoluble fraction). The cytosolic supernatant was transferred to a fresh tube, pelleted at 1000 g for 3 mins, and the resulting cleared supernatant transferred to a fresh tube. This cytosolic supernatant was finally centrifuged at 17000 g for 10 min to pellet any remaining non-cytosolic cellular material or debris. The cytosolic fraction was transferred to a fresh tube. The insoluble fraction was solubilised in in 4% SDS lysis buffer (4% SDS, 50mM Tris-HCl pH 7.5 and 150mM NaCl), 300 μL of lysis buffer per 60 mm dish, and pulse sonicated.

In those instances where the cytosolic fraction was to be used for DNA abundance quantification, an additional step was introduced into the above procedure where, after initial cell trypsinisation, one-third of the cell suspension was separately pelleted and lysed by addition of 400μl of 50 μM NaOH, followed by heating at 95 °C for 30 min to solubilise DNA, and neutralised by adding 50 μl 1 M Tris-HCL (pH 8.0). These extracts served as normalisation controls (*ACTB* qPCR, see below)

Nucleic acid purification

For total cell RNA isolation, cells were lysed directly in RLT lysis buffer with added β-mercaptoethanol as per manufacturer's protocol (Qiagen, RNeasy kit), and RNA was isolated using by spin-column RNA extraction kit (Qiagen, RNeasy kit). For RNA isolation from cytosolic fractions (above), Trizol / chloroform extraction was performed. Briefly, cytosolic fractions were mixed with Trizol (Thermo Fisher, #15596026) in a 1:1 ratio and incubated for 5 min. 0.2 mL of chloroform per 1 mL of Trizol was added to the mixture and vortexed vigorously for 15 seconds, followed by incubation at room temperature for 2 min. The mixture was centrifuged at 12,000 g for 15 mins at 4 °C. Following centrifugation, the upper aqueous phase was transferred carefully into a fresh RNase-free tube. 25 μg

of GlycoBlue coprecipitant (Thermo Fisher, #AM9516) was added and RNA precipitated by mixing with 0.5 mL of isopropanol per 1 ml of original Trizol reagent used, then incubation at room temperature for 10 min, followed by centrifugation at 12,000g for 10 min at 2 °C. The supernatant was removed completely and discarded. The RNA pellet was washed once with 1 ml of 75% ethanol per 1 ml of original Trizol reagent used, and subsequent centrifugation at 7,500g for 5 min at 2 °C. The ethanol wash was then fully removed, and the cleaned RNA pellet air-dried for 10 min, then re-dissolved in nuclease free water. To remove trace DNA, samples were treated with DNase-I using the RNAqueous DNase kit (Thermo Fisher), according to the manufacturer's protocol. For DNA isolation from cytosolic fractions (above), a spin-column DNA isolation kit (QIAamp, Qiagen) was used, according to the manufacturer's protocol. DNA was eluted in nuclease-free water. Nucleic acid concentration and purity was measured using a NanoDrop 2000c spectrophotometer (Thermo Scientific).

Quantitative PCR (qPCR)

Reverse transcription from RNA isolated from total cell lysates was done using QScript cDNA SuperMix (QuantaBio #95048), according to the manufacturer's protocol. Reverse transcription from RNA isolated from cytosolic fractions (above) was performed using the SuperScript III reverse transcriptase (Thermo Fisher, cat.no 18080093). In both instances, random DNA hexamers were used for priming. qPCR was performed using SYBR Select Master Mix (Applied Biosystems, #4472908), according to the manufacturer's protocol, on a StepOne Real-Time PCR System instrument (ThermoFisher). For cDNA from total cell RNA samples, transcripts were quantified using the $\Delta\Delta$Ct method, while standard curve methodology was used to quantify cDNA/DNA obtained from cytosolic fractions. Each biological replicate data point is typically the mean of 3 technical replicates. Primers used are listed in Methods Table 3.

Target	Forward sequence	Reverse sequence
12S	AAGATTACACATGCAAGCA	TTGATCGTGGTGATTTAGA
18S	GTAACCCGTTGAACCCCATT	CCATCCAATCGGTAGTAGCG
COII	CGTCTGAACTATCCTGCCCG	TGGTAGGGAGGGATCGTTG
CXCL10	TCTGAGACATTCCTCAATTGCT	AGAGGTACTCCTTGAATGCCA
CYTB	AGACAGTCCCACCCTCACAC	GGTGATTCCTAGGGGGTTGT
IFIT1	CCTCCTTGGGTTCGTCTACA	GGCTGATATCTGGGTGCC TA
IFNL1	GAAGCAGTTGCGATTTAGCC	TTCAGCTTGAGTGACTCTTCCA
ISG15	CATCTTTGCCAGTACAGGAG	CCGATCTTCTGGGTGATCTG
ND1	CACCCAAGAACAGGGTTTGT	AGGAATGCCATTGCGATTAG
ND6	ACCCCACAAACCCCATTACT	TTGGGGTCATTGGTGTTCTT
ACTB (**β**-actin)	GAGCACAGAGCCTCGCCTTT	ACATGCCGGAGCCGTTGTC

Methods Table 3   qPCR oligonucleotide primers used in this study

Statistical analysis

All statistical analysis was carried out using GraphPad (Prism) (version 9.5.1). All independent biological replicates are represented with standard error of the mean (S.E.M.). For electron microscopy image analyses, individual cells are represented as data points and variance described as standard deviation (S.D.). Tests are detailed in Figure Legends. In general, an assumption of normality was made and all tests were two-sided. Thus, where pairwise comparisons between two samples are performed, this is performed with an unpaired Student's t-test or a one-sample t-test (latter instance when comparing against the normalisation control). Where a putatively non-normal distribution was clearly observed, in Fig. 2J, a one-sample Wilcoxon test was instead used for the pairwise comparison. When multiple comparisons were performed in instances where there was only one independent variable, we generally used one-way ANOVA with Holm-Šídák testing. Where multiple comparisons were performed in instances of two independent variables in data set (e.g. cell line identity and/or drug application) we generally performed two-way ANOVA with Holm-Šídák testing. In the case of cytosolic RNA/DNA analysis, ratio paired t-tests were performed. Replicates containing outliers as detected by the Grubb's test were removed prior to the ratio paired t-test. Comparison of pairs of

frequency distributions were performed by the Kolmogorov-Smirnov test for each of three replicates, and a p-value for each replicate generated individually.

Code availability

The custom FIJI (ImageJ) plug-in macro to shuffle galectin-8 foci within cell space is reproduced below:

```
1. # get current image, a image with labelled objects
2. plus=WindowManager.getCurrentImage()
3. # wrap ImagePlus into 3D suite image format
4. img=ImageInt.wrap(plus)
5. # create a population of 3D objects
6. pop=Objects3DPopulation(img)
7. # create a mask where to shuffle objects
8. # here same as input image
9. IJ.selectWindow("mask");
10. plus2=WindowManager.getCurrentImage()
11. maskimg=ImageInt.wrap(plus2)
12. obj=Object3DVoxels(maskimg)
13. pop.setMask(obj)
14. # get the shuffled objects and build a population from them
15. listObjects=pop.shuffle()
16. pop2=Objects3DPopulation(listObjects)
17. # draw the results
18. res = ImageShort("res", img.sizeX, img.sizeY, img.sizeZ)
19. pop2.draw(res)
20. res.show("shuffle")
```

Data availability

Raw data not included in this manuscript are available from the corresponding author upon reasonable request.

References

Akira, S., Uematsu, S., and Takeuchi, O. (2006). Pathogen recognition and innate immunity. Cell *124*, 783-801.

Antonicka, H., Sasarman, F., Nishimura, T., Paupe, V., and Shoubridge, E.A. (2013). The mitochondrial RNA-binding protein GRSF1 localizes to RNA granules and is required for posttranscriptional mitochondrial gene expression. Cell Metab *17*, 386-398.

Bhattacharya, A., Mukherjee, R., Kuncha, S.K., Brunstein, M.E., Rathore, R., Junek, S., Munch, C., and Dikic, I. (2023). A lysosome membrane regeneration pathway depends on TBC1D15 and autophagic lysosomal reformation proteins. Nat Cell Biol *25*, 685-698.

Bryant, J.D., Lei, Y., VanPortfliet, J.J., Winters, A.D., and West, A.P. (2022). Assessing Mitochondrial DNA Release into the Cytosol and Subsequent Activation of Innate Immune-related Pathways in Mammalian Cells. Curr Protoc *2*, e372.

Bussi, C., Heunis, T., Pellegrino, E., Bernard, E.M., Bah, N., Dos Santos, M.S., Santucci, P., Aylan, B., Rodgers, A., Fearns, A.*, et al.* (2022). Lysosomal damage drives mitochondrial proteome remodelling and reprograms macrophage immunometabolism. Nat Commun *13*, 7338.

Campeau, E., Ruhl, V.E., Rodier, F., Smith, C.L., Rahmberg, B.L., Fuss, J.O., Campisi, J., Yaswen, P., Cooper, P.K., and Kaufman, P.D. (2009). A versatile viral system for expression and depletion of proteins in mammalian cells. PLoS One *4*, e6529.

Chauhan, S., Kumar, S., Jain, A., Ponpuak, M., Mudd, M.H., Kimura, T., Choi, S.W., Peters, R., Mandell, M., Bruun, J.A.*, et al.* (2016). TRIMs and Galectins Globally Cooperate and TRIM16 and Galectin-3 Co-direct Autophagy in Endomembrane Damage Homeostasis. Dev Cell *39*, 13-27.

Cross, J., Durgan, J., McEwan, D.G., Tayler, M., Ryan, K.M., and Florey, O. (2023). Lysosome damage triggers direct ATG8 conjugation and ATG2 engagement via non-canonical autophagy. J Cell Biol *222*.

Crow, Y.J., and Stetson, D.B. (2022). The type I interferonopathies: 10 years on. Nat Rev Immunol *22*, 471-483.

Dhir, A., Dhir, S., Borowski, L.S., Jimenez, L., Teitell, M., Rotig, A., Crow, Y.J., Rice, G.I., Duffy, D., Tamby, C.*, et al.* (2018). Mitochondrial double-stranded RNA triggers antiviral signalling in humans. Nature *560*, 238-242.

Dikic, I., and Elazar, Z. (2018). Mechanism and medical implications of mammalian autophagy. Nat Rev Mol Cell Biol *19*, 349-364.

Duran, J., Salinas, J.E., Wheaton, R.P., Poolsup, S., Allers, L., Rosas-Lemus, M., Chen, L., Cheng, Q., Pu, J., Salemi, M.*, et al.* (2024). Calcium signaling from damaged lysosomes induces cytoprotective stress granules. EMBO J *43*, 6410-6443.

Eapen, V.V., Swarup, S., Hoyer, M.J., Paulo, J.A., and Harper, J.W. (2021). Quantitative proteomics reveals the selectivity of ubiquitin-binding autophagy receptors in the turnover of damaged lysosomes by lysophagy. Elife *10*, e72328.

Fan, H., Liu, W., Zeng, Y., Zhou, Y., Gao, M., Yang, L., Liu, H., Shi, Y., Li, L., Ma, J.*, et al.* (2023). DNA damage induced by CDK4 and CDK6 blockade triggers anti-tumor immune responses through cGAS-STING pathway. Commun Biol *6*, 1041.

Gahlot, P., Kravic, B., Rota, G., van den Boom, J., Levantovsky, S., Schulze, N., Maspero, E., Polo, S., Behrends, C., and Meyer, H. (2024). Lysosomal damage sensing and lysophagy initiation by SPG20-ITCH. Mol Cell *84*, 1556-1569 e1510.

Gallagher, E.R., and Holzbaur, E.L.F. (2023). The selective autophagy adaptor p62/SQSTM1 forms phase condensates regulated by HSP27 that facilitate the clearance of damaged lysosomes via lysophagy. Cell Rep *42*, 112037.

Goel, S., DeCristo, M.J., Watt, A.C., BrinJones, H., Sceneay, J., Li, B.B., Khan, N., Ubellacker, J.M., Xie, S., Metzger-Filho, O.*, et al.* (2017). CDK4/6 inhibition triggers anti-tumour immunity. Nature *548*, 471-475.

He, X., Zhu, Y., Zhang, Y., Geng, Y., Gong, J., Geng, J., Zhang, P., Zhang, X., Liu, N., Peng, Y.*, et al.* (2019). RNF34 functions in immunity and selective mitophagy by targeting MAVS for autophagic degradation. EMBO J *38*, e100978.

Hoyer, M.J., Swarup, S., and Harper, J.W. (2022). Mechanisms Controlling Selective Elimination of Damaged Lysosomes. Curr Opin Physiol *29*, 100590.

Ishikawa, H., and Barber, G.N. (2008). STING is an endoplasmic reticulum adaptor that facilitates innate immune signalling. Nature *455*, 674-678.

Jia, J., Abudu, Y.P., Claude-Taupin, A., Gu, Y., Kumar, S., Choi, S.W., Peters, R., Mudd, M.H., Allers, L., Salemi, M.*, et al.* (2018). Galectins Control mTOR in Response to Endomembrane Damage. Mol Cell *70*, 120-135 e128.

Jia, J., Bissa, B., Brecht, L., Allers, L., Choi, S.W., Gu, Y., Zbinden, M., Burge, M.R., Timmins, G., Hallows, K.*, et al.* (2020a). AMPK, a Regulator of Metabolism and Autophagy, Is Activated by Lysosomal Damage via a Novel Galectin-Directed Ubiquitin Signal Transduction System. Mol Cell *77*, 951-969 e959.

Jia, J., Claude-Taupin, A., Gu, Y., Choi, S.W., Peters, R., Bissa, B., Mudd, M.H., Allers, L., Pallikkuth, S., Lidke, K.A.*, et al.* (2020b). Galectin-3 Coordinates a Cellular System for Lysosomal Repair and Removal. Dev Cell *52*, 69-87 e68.

Jia, J., Wang, F., Bhujabal, Z., Peters, R., Mudd, M., Duque, T., Allers, L., Javed, R., Salemi, M., Behrends, C.*, et al.* (2022). Stress granules and mTOR are regulated by membrane atg8ylation during lysosomal damage. J Cell Biol *221*.

Jimenez-Moreno, N., Salomo-Coll, C., Murphy, L.C., and Wilkinson, S. (2023). Signal-Retaining Autophagy Indicator as a Quantitative Imaging Method for ER-Phagy. Cells *12*.

Johansen, T., and Lamark, T. (2020). Selective Autophagy: ATG8 Family Proteins, LIR Motifs and Cargo Receptors. J Mol Biol *432*, 80-103.

Katayama, H., Hama, H., Nagasawa, K., Kurokawa, H., Sugiyama, M., Ando, R., Funata, M., Yoshida, N., Homma, M., Nishimura, T.*, et al.* (2020). Visualizing and Modulating Mitophagy for Therapeutic Studies of Neurodegeneration. Cell *181*, 1176-1187 e1116.

Kim, J., Gupta, R., Blanco, L.P., Yang, S., Shteinfer-Kuzmine, A., Wang, K., Zhu, J., Yoon, H.E., Wang, X., Kerkhofs, M.*, et al.* (2019). VDAC oligomers form mitochondrial pores to release mtDNA fragments and promote lupus-like disease. Science *366*, 1531-1536.

Kravic, B., Bionda, T., Siebert, A., Gahlot, P., Levantovsky, S., Behrends, C., and Meyer, H. (2022). Ubiquitin profiling of lysophagy identifies actin stabilizer CNN2 as a target of VCP/p97 and uncovers a link to HSPB1. Mol Cell *82*, 2633-2649 e2637.

Lazarou, M., Sliter, D.A., Kane, L.A., Sarraf, S.A., Wang, C., Burman, J.L., Sideris, D.P., Fogel, A.I., and Youle, R.J. (2015). The ubiquitin kinase PINK1 recruits autophagy receptors to induce mitophagy. Nature *524*, 309-314.

Llanos, S., Megias, D., Blanco-Aparicio, C., Hernandez-Encinas, E., Rovira, M., Pietrocola, F., and Serrano, M. (2019). Lysosomal trapping of palbociclib and its functional implications. Oncogene *38*, 3886-3902.

Mackenzie, K.J., Carroll, P., Martin, C.A., Murina, O., Fluteau, A., Simpson, D.J., Olova, N., Sutcliffe, H., Rainger, J.K., Leitch, A.*, et al.* (2017). cGAS surveillance of micronuclei links genome instability to innate immunity. Nature *548*, 461-465.

Maejima, I., Takahashi, A., Omori, H., Kimura, T., Takabatake, Y., Saitoh, T., Yamamoto, A., Hamasaki, M., Noda, T., Isaka, Y., and Yoshimori, T. (2013). Autophagy sequesters damaged lysosomes to control lysosomal biogenesis and kidney injury. EMBO J *32*, 2336-2347.

Mathew, R., Khor, S., Hackett, S.R., Rabinowitz, J.D., Perlman, D.H., and White, E. (2014). Functional role of autophagy-mediated proteome remodeling in cell survival signaling and innate immunity. Mol Cell *55*, 916-930.

McArthur, K., Whitehead, L.W., Heddleston, J.M., Li, L., Padman, B.S., Oorschot, V., Geoghegan, N.D., Chappaz, S., Davidson, S., San Chin, H.*, et al.* (2018). BAK/BAX macropores facilitate mitochondrial herniation and mtDNA efflux during apoptosis. Science *359*.

Montava-Garriga, L., Singh, F., Ball, G., and Ganley, I.G. (2020). Semi-automated quantitation of mitophagy in cells and tissues. Mech Ageing Dev *185*, 111196.

Nakahira, K., Haspel, J.A., Rathinam, V.A., Lee, S.J., Dolinay, T., Lam, H.C., Englert, J.A., Rabinovitch, M., Cernadas, M., Kim, H.P.*, et al.* (2011). Autophagy proteins regulate innate immune responses by inhibiting the release of mitochondrial DNA mediated by the NALP3 inflammasome. Nat Immunol *12*, 222-230.

Neuwirt, E., Magnani, G., Cikovic, T., Wohrle, S., Fischer, L., Kostina, A., Flemming, S., Fischenich, N.J., Saller, B.S., Gorka, O.*, et al.* (2023). Tyrosine kinase inhibitors can activate the NLRP3 inflammasome in myeloid cells through lysosomal damage and cell lysis. Sci Signal *16*, eabh1083.

Newman, A.C., Kemp, A.J., Drabsch, Y., Behrends, C., and Wilkinson, S. (2017). Autophagy acts through TRAF3 and RELB to regulate gene expression via antagonism of SMAD proteins. Nat Commun *8*, 1537.

Newman, A.C., Scholefield, C.L., Kemp, A.J., Newman, M., McIver, E.G., Kamal, A., and Wilkinson, S. (2012). TBK1 kinase addiction in lung cancer cells is mediated via autophagy of Tax1bp1/Ndp52 and non-canonical NF-kappaB signalling. PLoS One *7*, e50672.

Papadopoulos, C., Kirchner, P., Bug, M., Grum, D., Koerver, L., Schulze, N., Poehler, R., Dressler, A., Fengler, S., Arhzaouy, K.*, et al.* (2017). VCP/p97 cooperates with YOD1, UBXD1 and PLAA to drive clearance of ruptured lysosomes by autophagy. EMBO J *36*, 135-150.

Papadopoulos, C., Kravic, B., and Meyer, H. (2020). Repair or Lysophagy: Dealing with Damaged Lysosomes. J Mol Biol *432*, 231-239.

Park, N.Y., Jo, D.S., Yang, J.Y., Bae, J.E., Kim, J.B., Kim, Y.H., Kim, S.H., Kim, P., Lee, D.S., Yoshimori, T.*, et al.* (2025). Activation of lysophagy by a TBK1-SCF(FBXO3)-TMEM192-TAX1BP1 axis in response to lysosomal damage. Nat Commun *16*, 1109.

Prabakaran, T., Bodda, C., Krapp, C., Zhang, B.C., Christensen, M.H., Sun, C., Reinert, L., Cai, Y., Jensen, S.B., Skouboe, M.K.*, et al.* (2018). Attenuation of cGAS-STING signaling is mediated by a p62/SQSTM1-dependent autophagy pathway activated by TBK1. EMBO J *37*.

Radulovic, M., Schink, K.O., Wenzel, E.M., Nahse, V., Bongiovanni, A., Lafont, F., and Stenmark, H. (2018). ESCRT-mediated lysosome repair precedes lysophagy and promotes cell survival. EMBO J *37*, e99753.

Ran, F.A., Hsu, P.D., Wright, J., Agarwala, V., Scott, D.A., and Zhang, F. (2013). Genome engineering using the CRISPR-Cas9 system. Nat Protoc *8*, 2281-2308.

Ravenhill, B.J., Boyle, K.B., von Muhlinen, N., Ellison, C.J., Masson, G.R., Otten, E.G., Foeglein, A., Williams, R., and Randow, F. (2019). The Cargo Receptor NDP52 Initiates Selective Autophagy by Recruiting the ULK Complex to Cytosol-Invading Bacteria. Mol Cell *74*, 320-329 e326.

Rehwinkel, J., and Gack, M.U. (2020). RIG-I-like receptors: their regulation and roles in RNA sensing. Nat Rev Immunol *20*, 537-551.

Riley, J.S., Quarato, G., Cloix, C., Lopez, J., O'Prey, J., Pearson, M., Chapman, J., Sesaki, H., Carlin, L.M., Passos, J.F.*, et al.* (2018). Mitochondrial inner membrane permeabilisation enables mtDNA release during apoptosis. EMBO J *37*.

Roulois, D., Loo Yau, H., Singhania, R., Wang, Y., Danesh, A., Shen, S.Y., Han, H., Liang, G., Jones, P.A., Pugh, T.J.*, et al.* (2015). DNA-Demethylating Agents Target Colorectal Cancer Cells by Inducing Viral Mimicry by Endogenous Transcripts. Cell *162*, 961-973.

Seth, R.B., Sun, L., Ea, C.K., and Chen, Z.J. (2005). Identification and characterization of MAVS, a mitochondrial antiviral signaling protein that activates NF-kappaB and IRF 3. Cell *122*, 669-682.

Skowyra, M.L., Schlesinger, P.H., Naismith, T.V., and Hanson, P.I. (2018). Triggered recruitment of ESCRT machinery promotes endolysosomal repair. Science *360*.

Sun, L., Wu, J., Du, F., Chen, X., and Chen, Z.J. (2013). Cyclic GMP-AMP synthase is a cytosolic DNA sensor that activates the type I interferon pathway. Science *339*, 786-791.

Tan, J.X., and Finkel, T. (2022). A phosphoinositide signalling pathway mediates rapid lysosomal repair. Nature *609*, 815-821.

Thurston, T.L., Wandel, M.P., von Muhlinen, N., Foeglein, A., and Randow, F. (2012). Galectin 8 targets damaged vesicles for autophagy to defend cells against bacterial invasion. Nature *482*, 414-418.

Tigano, M., Vargas, D.C., Tremblay-Belzile, S., Fu, Y., and Sfeir, A. (2021). Nuclear sensing of breaks in mitochondrial DNA enhances immune surveillance. Nature *591*, 477-481.

Torres-Odio, S., Lei, Y., Gispert, S., Maletzko, A., Key, J., Menissy, S.S., Wittig, I., Auburger, G., and West, A.P. (2021). Loss of Mitochondrial Protease CLPP Activates Type I IFN Responses through the Mitochondrial DNA-cGAS-STING Signaling Axis. J Immunol *206*, 1890-1900.

Turco, E., Witt, M., Abert, C., Bock-Bierbaum, T., Su, M.Y., Trapannone, R., Sztacho, M., Danieli, A., Shi, X., Zaffagnini, G.*, et al.* (2019). FIP200 Claw Domain Binding to p62 Promotes Autophagosome Formation at Ubiquitin Condensates. Mol Cell *74*, 330-346 e311.

Uzhachenko, R.V., Bharti, V., Ouyang, Z., Blevins, A., Mont, S., Saleh, N., Lawrence, H.A., Shen, C., Chen, S.C., Ayers, G.D.*, et al.* (2021). Metabolic modulation by CDK4/6 inhibitor promotes chemokine-mediated recruitment of T cells into mammary tumors. Cell Rep *35*, 109271.

Vargas, J.N.S., Hamasaki, M., Kawabata, T., Youle, R.J., and Yoshimori, T. (2023). The mechanisms and roles of selective autophagy in mammals. Nat Rev Mol Cell Biol *24*, 167-185.

Vargas, J.N.S., Wang, C., Bunker, E., Hao, L., Maric, D., Schiavo, G., Randow, F., and Youle, R.J. (2019). Spatiotemporal Control of ULK1 Activation by NDP52 and TBK1 during Selective Autophagy. Mol Cell *74*, 347-362 e346.

Wang, F., Gomez-Sintes, R., and Boya, P. (2018). Lysosomal membrane permeabilization and cell death. Traffic *19*, 918-931.

West, A.P., Khoury-Hanold, W., Staron, M., Tal, M.C., Pineda, C.M., Lang, S.M., Bestwick, M., Duguay, B.A., Raimundo, N., MacDuff, D.A.*, et al.* (2015). Mitochondrial DNA stress primes the antiviral innate immune response. Nature *520*, 553-557.

Wong, Y.C., Ysselstein, D., and Krainc, D. (2018). Mitochondria-lysosome contacts regulate mitochondrial fission via RAB7 GTP hydrolysis. Nature *554*, 382-386.

Wu, J., Wang, J., O'Connor, T.N., Tzetzo, S.L., Gurova, K.V., Knudsen, E.S., and Witkiewicz, A.K. (2024). Separable Cell Cycle Arrest and Immune Response Elicited through Pharmacological CDK4/6 and MEK Inhibition in RASmut Disease Models. Mol Cancer Ther *23*, 1801-1814.

Xu, L., Li, M., Yang, Y., Zhang, C., Xie, Z., Tang, J., Shi, Z., Chen, S., Li, G., Gu, Y.*, et al.* (2022). Salmonella Induces the cGAS-STING-Dependent Type I Interferon Response in Murine Macrophages by Triggering mtDNA Release. mBio *13*, e0363221.

Yang, X., Boehm, J.S., Yang, X., Salehi-Ashtiani, K., Hao, T., Shen, Y., Lubonja, R., Thomas, S.R., Alkan, O., Bhimdi, T.*, et al.* (2011). A public genome-scale lentiviral expression library of human ORFs. Nat Methods *8*, 659-661.

Yoneyama, M., Kikuchi, M., Natsukawa, T., Shinobu, N., Imaizumi, T., Miyagishi, M., Taira, K., Akira, S., and Fujita, T. (2004). The RNA helicase RIG-I has an essential function in double-stranded RNA-induced innate antiviral responses. Nat Immunol *5*, 730-737.

Zitvogel, L., Galluzzi, L., Kepp, O., Smyth, M.J., and Kroemer, G. (2015). Type I interferons in anticancer immunity. Nat Rev Immunol *15*, 405-414.

Zoncu, R., and Perera, R.M. (2022). Built to last: lysosome remodeling and repair in health and disease. Trends Cell Biol *32*, 597-610.

Supplementary Figures and Movie Legends

SRAI-GAL8

Supplementary Figure 1

A549- or MCF7-SRAI-GAL8 cells were treated for 24 h with DMSO, 0.7 mM LLOMe (+/- 0.2 µM of the lysosome blocker Bafilomycin, BafA1), or 10 µM Palbociclib, Pb, then imaged by widefield fluorescence microscopy. TOLLES-only foci are generated in response to both drugs, but this is blocked by BafA1.

Supplementary Figure 2

A549 NT (non-targeted wild-type), ΔATG5 or Rescue cells (ΔATG5 + GFP-ATG5) were treated with DMSO (-) or 10 µM Palbociclib, Pb (+), for 48 h, then immunoblotted to assess interferon pathway activation (phospho-Y701 STAT1, P-STAT1) (n = 3, representative replicate shown).

Supplementary Figure 3

A) Rho+ and Rho0 cultures of A549 Rescue (ΔATG5 + GFP-ATG5) and ΔATG5 cells generated as described in Fig. 6A-C were subjected to total cellular DNA isolation and qPCR for mitochondrial DNA (mtDNA) sequences *COII* and *ND1* (normalised to *ACTB*, nuclear DNA, nDNA).

B) A549 ΔATG5 cells stably-expressing Cas9 were transduced with non-targeting sgRNA (gNT or gNT-2) or sequentially transduced with gRNAs against *BAX* and *BAK* in denoted order. Cells were treated with DMSO or 10 µM Pb for 48 h and immunoblotted for phospho-Y701-STAT1 (P-STAT1) and Bax/Bak knockout controls.

Supplementary Movie 1

180° rotation around y-axis of 3D rendered cell shown in **Fig. 2D** (equivalent to lower zoom panel 1). YFP-GAL8 in green, TOM20 in red. Mitochondrial volume has been rendered to model the space occupied by the mitochondrial network.

Supplementary Movie 2

Supplementary Movie 2 corresponds to still frames in **Fig. 2F** row 1. YFP-GAL8 in green, MitoTracker in red.

Supplementary Movie 3

Supplementary Movie 2 corresponds to still frames in **Fig. 2F** row 2. YFP-GAL8 in green, MitoTracker in red.